Endangered Przewalski's horse, *Equus pzrewalskii,* cloned from historically

cryopreserved cells

Ben J. Novak[1], Oliver A. Ryder[2], Marlys L. Houck[2], Andrea S. Putnam[2], Kelcey Walker[3], Lexie Russell[3],

Blake Russell[3], Shawn Walker[3], Sanaz Sadeghieh Arenivas[3], Lauren Aston[3], Gregg Veneklasen[4], Jamie A.

Ivy[5], Klaus-Peter Koepfli[6,7], Anna Rusnak[8], Jaroslav Simek[9], Anna Zhuk[8,10], Ryan Phelan[1]

1. Revive & Restore, 1505 Bridgeway #203, Sausalito, CA 94965, USA

2. Beckman Center for Conservation Research, San Diego Zoo Wildlife Alliance, Escondido, CA,

92027, USA

3. ViaGen Pets & Equine, 715 Discovery Blvd #410, Cedar Park, TX, 78613, USA

4. Timber Creek Veterinary Hospital, 19302 Farm to Market Rd 1541, Canyon, TX, 79015, USA

5. Species Conservation Toolkit Initiative, Chicago Zoological Society, Brookfield, Illinois, USA

6. Smithsonian-Mason School of Conservation, George Mason University, 1500 Remount Road,

Front Royal, VA 22630, USA

7. Center for Species Survival, Smithsonian's National Zoo and Conservation Biology Institute,

1500 Remount Road, Front Royal, VA 22630, USA

8. Institute of Applied Computer Science, ITMO University, 197101 St. Petersburg, Russia

9. Prague Zoo, U Trojskeho zamku 120/3 Prague 7, Czech Republic

10. Laboratory of Amyloid Biology, St. Petersburg State University, 199034 St. Petersburg, Russia

Abstract

Two endangered Przewalski's horse stallions were cloned from fibroblast cells cultured and

cryopreserved in 1980. These stallions are clones of a male that lived from 1975-1998 that pedigree

analyses identified as a genetically valuable male for present-day conservation breeding. This is the

first time that multiple healthy clones have been produced for an endangered species.

Main Text

We report the first clones of the endangered Przewalski's horse, *Equus przewalskii*, or Takhi as it is

called by Mongolians. The clones were produced via cross-species somatic cell nuclear transfer from

donor cells cryopreserved in 1980 and using domestic horses (*Equus caballus*) as oocyte donors and surrogate mothers. The identity of the clones was verified by a microsatellite panel, and whole genome sequencing of one clone verified a match to the original donor animal (Figure 1A) with minor somatic mutations, consistent with whole genome comparisons of clones of other species[1,2]. Given the clones were produced via cross-species somatic cell nuclear transfer, it was expected that their mitochondrial haplotypes should be a mismatch to the donor, which was verified by whole mitochondrial genome analysis (Figure 1B).

While several dozen domestic and wild species have been successfully cloned[3], according to the International Union for the Conservation of Nature (IUCN) assessment and classifications of endangerment (Near Threatened, Vulnerable, Endangered, Critically Endangered, Extinct in the Wild), this is only the fourth endangered species to be successfully cloned and the first time that multiple viable clones of an endangered species have survived past the neonatal period (Table 1; note several amphibian species[4-8] and one mammal, the cynomalgus monkey *Macaca fascicularis*[9], that are currently assessed as endangered were not considered endnagered at the time they were first cloned; and two mammals, the sand cat, *Felis margarita*[10], and the European mouflon, *Ovis aries musimon*[11], were erroneously considered endangered at the time of cloning but have since been reclassified).

Table 1. History of Cloning Endangered Species

Common Name	Species Name	IUCN Status	Year(s) Cloned	Number of Live Births	Number surviving neonatal period	Reproductive Capability	References
Gaur	*Bos gaurus*	Vulnerable	2001, 2012	1, 1	0	-	12, 13
Banteng	*Bos javanicus*	Endangered	2003	2	1	Infertile (cryptorchidism)	14, 12
Esfahan Mouflon	*Ovis gmelini isphahanica*	Near Threatened	2010, 2015	2, NA	0, NA	Not reported.	15, 16
Przewalksi's Horse	*Equus przewalskii*	Endangered	2020, 2023	2	2	Not yet sexually mature.	Reported here.

47
Figure 1. A) Euler diagram of biallelic private and shared single nucleotide variants (SNVs) for five Przewalski's horses of
multiple founder breeding lines, revealing nearly identical overlap of Kurt (clone) and Kuporovitch (donor). B) Maximum
likelihood phylogenetic tree of mitochondrial genomes of domestic horses (black colored labels) and Przewalski's horses
(blue colored labels) based on an alignment of 16,130 bp. The tree is rooted with the onager (*Equus hemionus*). NCBI
Genbank accession numbers are associated with domestic and Przewalski's horses sequenced in previous studies. Asterisks
indicate nodes with >90 bootstrap support values, out of 500 replicates. Przewalski's horse individuals that were newly
sequenced for this study are indicated by colored circles corresponding to A, revealing the expected mismatch between
Kurt and Kuporovitch. Scale bar at bottom refers to the number of substitutions per site along branches. C) Kurt, the first
clone born August 6 2020, at 25 months of age. D) Ollie, the second clone, born February 17, 2023, at 7 months of age.
Photographs courtesy of San Diego Zoo Wildlife Alliance. E) Still-frame from archival footage of SB615, Kuporovitch, 8 years
of age.
59

The two males produced so far, named Kurt and Ollie (Figure 1C, 1D), are clones of

International Studbook Number 615 (SB615), Kuporovitch (Figure 1E), born 1975 and died 1998.

Recruitment of this cell line was deemed genetically beneficial, as analyses suggested the cell line to be

underrepresented within multiple regionally managed conservation breeding programs (Methods,

Table 2). This is only the second species in which decades-old biobanked materials have been used to

produce clones for conservation management (the first time being a Banteng, *Bos javanicus*, cloned

from 25-year-old cells[14,12], Table 1) and the first viable opportunity for clones of an endangered species

to fulfill their purpose: to reproduce and rescue valuable genetic diversity for conservation breeding[17-

19]. While neither of the clones has yet reached sexual maturity during a breeding season event

(Przewalski's horses reach sexual maturity at 2.5-3 years of age[20]), the testicular development of each

clone is so far normal.

The Przewalksi's horse was extirpated from its wild range across the central Eurasian grasslands

with the last commonly cited wild sighting in the Dzungarian Gobi region in 1969[21], at which time,

according to the Przewalski's Horse International Studbook, 174 individuals survived scattered across

several European and North American zoos. Those animals were descended from several founder lines

that had been maintained for display. Fortunately, by the 1950s, zoo managers recognized that the

future of the species relied on the horses surviving in ex situ herds. In 1959, the first International

Symposium on the Preservation of the Przewalski's Horse was organized by the Prague Zoo, initiating

the first efforts to establish a plan to not only save the species from extinction but to organize a

coordinated breeding program to produce enough horses to begin restoring herds to the wild. The plan

was formalized at the 5th International Symposium in 1990[22]. The breeding program had been rather

successful and following the 5th Symposium, reintroductions began in the early 1990s. Today, several

hundred horses live in free-roaming and semi-free-roaming herds in Mongolia, China, Russia,

Kazakhstan, and Hungary. The world's ~3,000 living Przewalski's horses are the result of over 8,700

horses bred over nearly 20 generations from just 12 wild founders captured from 1898-1947.

While the history of the Przewalski's horse is similar to many other species saved from

extinction through conservation breeding and reintroductions, such as the California Condor,

Whooping Crane, or black-footed ferret, its breeding management history is complex due to its origins

preceding the modern era of zoo conservation. To illuminate by contrast, the founders of the

endangered black-footed ferret breeding program were captured over a short period of time, brought

into captivity together, and bred in a manner to attempt to retain as much genetic diversity as

possible. The result is that all living black-footed ferrets today trace their ancestry to all founders in

roughly equal proportions[23]. The Przewalski's horse founders were captured over a long period of time,

beginning in the 1890s, and scattered across several zoos in which they were bred as separate lineages

for many decades before the coordinated efforts for conservation were initiated. The founders were

not captured and bred with the intent to save the species nor necessarily to maximize the retention of

surviving genetic diversity. Thus, when coordinated efforts for restoration began, the population was

represented by five distinct lineages, two of which whose founders were interbred with domestic

horses (*Equus caballus*)[24].

Even when concerted restoration breeding efforts began, there were significant differences of

opinion as to how best to interbreed the five founder lineages. A desire for "genetic purity" led to

continued and problematic segregation. Combined with breeding policies aimed at purging domestic

horse ancestry within introgressed lines, these management practices led to significant losses of wild

Przewalski's horse genetic diversity and high inbreeding coefficients. Experts at the time disagreed

about the phenotypic impacts of the introgression of the Mongolian domestic mare, but to our

knowledge, a comprehensive morphological analysis has never been reported. Fortunately, amidst this

complex breeding history, initially uninformed by currently accepted practices of conservation

management, biopsies of a large number of historic Przewalski's horses were collected and

cryopreserved, establishing a biobank today of nearly 600 individuals (298 individuals represented with

cell lines and 277 represented by tissues preserved in DMSO from which cell lines could potentially be

derived post-thaw), with several individuals just 1-5 generations separated from the wild founders.

Many individuals with domestic horse ancestry were intentionally and detrimentally

underutilized in the program, including SB615, the donor sire of the clones reported here. Previous

pedigree analyses concluded that significant historic allelic diversity would be lost without increasing

representation of this lineage[24], yet despite those findings SB615 was only bred during four out of 21

potential breeding seasons in his lifetime. When selecting individuals to clone for ongoing genetic

management, the genetic values of available cell lines were evaluated in the context of current

conservation breeding programs. The cell line from SB615 was chosen to clone because this line had

the lowest mean kinship of all males, indicating a genetic benefit for recruitment (Methods, Table 2).

The reproducible success of cloning this decades-long cryopreserved cell line demonstrates the

potential to clone other cryopreserved cell lines for ongoing genetic management. Thanks to the

extensive biobanking efforts of cell lines and germplasm (gametes), cloning more historic individuals or

"resurrecting" stallions through artificial insemination[25] offers the chance to undo some of the genetic

erosion caused by decades of managing the small zoo populations. In practice, the extensive biobank

of Przewalski's horse cell lines and gametes can now be viewed as an active reproductive resource for

future breeding decisions. The use of advanced reproductive technologies is still rare in conservation,

owing partially to the nascency of reproductive knowledge and techniques for most species. While

biobanking of viable cells and gametes for endangered species has been established in multiple

centers, its potential impacts are to this point limited by the small number of taxa cryopreserved and

low degree of these collections to capture extant variation[26], despite well-established methods for

many species, particularly for mammals[27]. The Przewalski's horse recovery program could be the first

to demonstrate the ideal model system for biobanking for conservation impact[19].

Methods

<u>Pedigree Analysis</u>

For the 2020 clone, biobanked cell lines were analyzed to determine which would confer the

greatest genetic benefit to the cooperatively managed Association of Zoos and Aquarium (AZA)

Przewalski's horse population. Cooperative management for this population occurs through a strategy

of minimizing pedigree-based kinships[28,29]. The mean kinship of an individual is the average kinship

between that individual and all other living individuals in the population including itself. Reproduction

from breeding pairs with low mean kinships and inbreeding coefficients below the population's

average mean kinship are prioritized. Twelve cell lines housed at the San Diego Zoo Wildlife Alliance

Biodiversity Bank's Frozen Zoo® were each independently analyzed to compare their mean kinships,

should they be added to the population, with the average mean kinship of the managed population

living at the time of analyses. The population included both AZA and non-AZA facilities that were

participating in cooperative management. Data from the AZA Regional Asian Wild Horse (*Equus*

przewalskii) Studbook (current to December 31, 2019[30]) was exported from Poplink 2.5.2[31] to PMx

v.1.6.20190628[32] to calculate pedigree-based kinships. Sterilized males and females over the age of 25

were excluded from kinship analyses. Each cell line, independently of the others, was added to the

genetically managed AZA population to estimate the cell line's mean kinship within the population. The

mean kinship of each cell line was then ranked from lowest mean kinship to highest mean kinship

(Table 2).

Table 2: The mean kinship of each analyzed cell line (CL) relative to each population. Population size (N) and average mean
kinship (MK). Cell line 615, chosen for both the 2020 and 2023 clones, is in bold. *Mean kinship of cell line 615 after siring 7
hypothetical offspring. Identity of other cell line studbook numbers is not disclosed, as they are proprietary to multiple
institutions.
157

Cell Line ID	Cell Line Sex	N. America 2020 N = 121 MK = 0.2175	AZA 2023 N = 74 MK = 0.2046	EAZA N = 211 MK= 0.1659	EARAZA N = 106 MK =0.1946	Reintroduced N = 1161 MK = 0.1658
CL#615	**M**	**0.0930**	**0.1137 / 0.1407***	**0.1309**	**0.0913**	**0.1432**
CL#1	M	-	0.1848	0.1555	0.1563	0.136
CL#2	F	0.2322	0.2192	0.192	0.2047	0.1644
CL#3	F	-	0.1027	0.1346	0.0925	0.1466
CL#4	F	0.0842	0.0993	0.135	0.0927	0.1475
CL#5	M	0.1338	0.139	0.1561	0.1716	0.1589
CL#6	M	0.2211	0.2096	0.1567	0.1367	0.1316
CL#7	F	0.2091	0.1956	0.1581	0.1528	0.1308
CL#8	M	0.2437	0.2253	0.1788	0.1853	0.146
CL#9	F	-	0.097	0.1307	0.0928	0.146
CL#10	M	-	0.2243	0.1724	0.1777	0.1409
CL#11	M	0.2350	0.2184	0.1746	0.1801	0.1451
CL#12	M	-	0.196	0.1906	0.1899	0.1574
CL#13	M	0.2077	0.1949	0.1696	0.178	0.1395
CL#14	M	-	0.1962	0.1695	0.1852	0.1398
CL#15	F	-	0.2073	0.1847	0.1939	0.1531
CL#16	F	-	0.133	0.1382	0.1119	0.1382
CL#17	F	0.1335	0.138	0.1441	0.1127	0.1448
CL#18	F	0.2252	0.2111	0.1658	0.1613	0.1332
CL#19	M	-	0.1922	0.1838	0.186	0.1502
CL#20	M	-	0.1888	0.1817	0.1808	0.1509
CL#21	M	0.1945	0.1874	0.1562	0.1431	0.1404
CL#22	F	-	0.1253	0.1487	0.1176	0.1427
CL#23	M	-	0.1355	0.1241	0.1786	0.1578
CL#24	M	-	0.1287	0.1572	0.1475	0.1626

The kinship analysis to prioritize cell lines for the 2023 clone were performed using the same

methods as the 2020 clone, with the following modifications. Thirteen additional cell lines were added

to the genetic analyses, for a total of 25 cell lines examined. Data from the International Studbook for

Przewalski's horse was exported from the ZIMS Studbook (current to December 2021[33]) to PMx

v.1.7.0.20210915 to calculate pedigree-based kinships. In addition to comparing the mean kinship of

the cell lines in the context of the AZA *ex situ* population that included the young 2020 clone, mean

kinships of the cell lines within three other regional populations were examined: the European

Association of Zoos and Aquariums (EAZA) *ex situ* population, the Eurasian Regional Association of Zoos

and Aquariums (EARAZA) *ex situ* population, and the combined reintroduced free-roaming and semi-

free populations of Orenburg Nature Reserve (Russia), Gobi National Park (Mongolia), Hortobagy

National Park (Hungary), and Hustai Nuruu National Park (Mongolia). To account for the possible

genetic representation from future reproduction of the 2020 clone within the AZA population, 8

hypothetical offspring were added to the AZA population where the 2020 clone was the sire and 7

reproductive females residing at San Diego Zoo Safari Park were the dams. The mean kinship of each

cell line was then ranked from lowest mean kinship to highest mean kinship within each region (Table

2).

<u>Somatic Cell Nuclear Transfer</u>

Table 3. Cross-species cloning results for donor cell line SB615. *Gestation includes 7 days *in vitro*

Trial Year	Embryos Transferred	Surrogates	Pregnancies	Live Births (% total embryos)	Gestation Length*
2020	4	4	3 (75%)	1 (25%)	322 days
2023	7	7	4 (57%)	1 (14%)	337 days
Total	11	11	7 (63%)	2 (18%)	-

To clone SB615, a vial of the historically cryopreserved fibroblast cells was shipped from the San

Diego Zoo Wildlife Alliance Frozen Zoo   to the commercial cloning company ViaGen Pets & Equine's

Texas based facility where the cells were thawed and donor cells were fused with enucleated domestic

horse oocytes to produce a total of 11 reconstructed embryos. The reconstructed embryos were

transferred to 11 domestic horse surrogate mares (1 embryo per mare). Two ensuing pregnancies

resulted in live births (Table 3, Figure 1).

<u>Microsatellite Genotyping</u>

Blood samples were collected from the clone horses and shipped to the University of California

Davis Veterinary Medicine Veterinary Genetics Laboratory for nuclear microsatellite genotyping along

with a cell pellet sample of the SB615 donor cells. A panel of 15 nuclear microsatellite loci were

analyzed for each clone, which include the mandatory loci for assessing equine parentage

recommended by the International Society for Animal Genetics (https://www.isag.us/): *AHT4*, *AHT5*,

AME, *ASB2*, *ASB17*, *ASB23*, *HMS2*, *HMS3*, *HMS6*, *HMS7*, *HTG10*, *HTG4*, *LEX3*, *LEX33*, *VHL20*[34]. The two

clones were a match to the donor cells across all loci.

<u>Mitochondrial and Whole Genome Comparisons</u>

Given the close evolutionary relationship and introgression between domestic and Przewalski's

horses, previous mtDNA studies have found shared haplotypes between the species[35,36], and therefore

a single mtDNA locus PCR may not provide adequate resolution for identity comparisons. These

factors may also obscure microsatellite analyses. Therefore whole genome sequencing was performed

to compare nuclear/mitochondrial DNA identities of one of the two clones to the donor cells alongside a

selection of additional unrelated Przewalski's horses as a control data set. Cell pellets from cultured

fibroblasts of each horse were shipped to the commercial service lab Psomagen, Inc. (Rockville,

Maryland, USA) for short-read resequencing. Specifically, genomic DNA was extracted from cell pellets

using the Mag-Bind Blood and Tissue Kit (Omega Bio-Tek Inc., Norcross, GA). DNA concentrations

were assessed using Picogreen and Victor x2 fluorometry (Life Technologies, Carlsbad, CA) and an

Agilent 4200 Tapestation (Agilent Technologies, Santa Clara, CA), and quality was checked with 1%

TBE gel electrophoresis. Genomic DNAs were then sheared into 350 bp fragments with a Covaris S220

Ultrasonicator (Covaris, Woburn, MA). DNA fragments were used to generate genomic libraries using

the TruSeq DNA PCR-free library kit (Illumina, San Diego, CA), which were quality checked on an

Agilent 4200 Tapestation and quantitated using a Lightcycler qPCR assay (Roche Life Science, St.

Louis, MO). Libraries were paired-end sequenced (2 x 150 bp) to a depth of 20x on an Illumina

NovaSeq 6000 instrument, which generated ~56 Gb for each sample.

Paired-end reads were explored with FastQC to calculate and visualize sequence quality

metrics[37]. The reads were aligned to the reference genome assembly of domestic horse

(GCF_002863925.1) using the BWA-MEM- (v0.7.17) algorithm with default parameters[38], and the

reference was indexed, and a dictionary created using SAMtools (v.1.17)[39] and Picard (v2.27.5)[40]. The

mapped reads were sorted with SAMtools (v1.17), and duplicated reads were marked with Picard

(v2.27.5) using MarkDuplicates. Finally, alignment statistics were evaluated with Picard and SAMtools.

SNVs were called using GATK HaplotypeCaller (v4.2.6.1)[41], followed by the use of GATK

SelectVariants to obtain only biallelic SNVs. VariantFiltration was applied to filter the vcf files based on

the following parameters: "QD < 2.0" -filter-name "QD2", "QUAL < 30.0" -filter-name "QUAL30", "SOR >

3.0" -filter-name "SOR3", "FS > 60.0" -filter-name "FS60", "MQ < 40.0" -filter-name "MQ40", -filter

"MQRankSum < -12.5 ", -filter "ReadPosRankSum < -8.0". The variant call set was further filtered by

excluding multiallelic variants and setting a read depth threshold of DP < 5 using BCFtools (v1.17)[39].

<u>Private Alleles Analysis</u>

To illustrate the proportions of private SNVs and those that are shared between individuals, a

Euler diagram was created based on biallelic SNVs alone. VCF files were parsed and for each individual

we prepared a list of values containing SNVs and its benchmarks. Using the Eulerr R package[42] and

these lists, we then prepared a Euler diagram. The diagram showed a huge intersection across the

clone (Kurt) and the donor (Kuporovitch, SB 615); from a genetic point of view the two individuals are

equal. Number of alleles shared across Kurt and Kuporovitch (668,467) is comparable to numbers of

private alleles in other individuals which vary from 436,446 in Vintel SB1376 to 1,118,521 in Rosa

SB162 (representing the latest and earliest generations of the pedigree in our sample set, respectively).

Mitochondrial genome assembly and analysis

The mitochondrial genomes of the five sequenced Przewalski's horses were *de novo* assembled

using GetOrganelle version 1.7.7.0[43]. The full read set for each individual was used as input and the

complete mitochondrial genome of the domestic horse was employed as the seed, with the -F

animal_mt parameter and all remaining parameters set to their default values. The mean coverage of

the assemblies ranged from 342X – 1,619X, as estimated with Mosdepth[44]. We obtained mitochondrial

genome assemblies ranging in length from 16,576 – 16,593 bp, with the length differences

corresponding to the number of tandem repeats in the control region. The mitogenomes were

annotated in the MITOS2 webserver[45].

To assess the phylogenetic position of the five Przewalski's horses with respect to their

maternal ancestry, particularly the cloned individual Kurt, we downloaded mitochondrial genomes of

available Przewalski's horses (n = 16) as well as those from a diversity of domestic horse breeds (n =

32) from NCBI's Genbank, which were published in previous studies[46-49]. We used the mitochondrial

genome phylogeny reported by Der Sarkissian et al.[49] to select domestic horse breeds that provided

the maximal representation of mitochondrial lineage diversity. We also downloaded the mitogenome

of an onager, *Equus hemionus* (HM118851[50]), which was used as the outgroup to root the phylogenetic

tree. All mitogenomes were imported into Geneious Prime version 2023.2

(https://www.geneious.com), after which a 16,748 bp multiple sequence alignment was generated

using the MAFFT version 7.490 plugin with the following settings: AUTO algorithm, 200PAM / k=2

scoring matrix, 1.53 gap open penalty, and an offset value of 0.123[51]. Due to poor alignment and

missing sequence data for some individuals in the control region, we trimmed 618 bp from this region,

resulting in a final alignment of 16,130 bp. This alignment was used to construct a maximum likelihood

phylogeny using the Geneious Prime plugin of RAxML version 8.2.11[52]. The nucleotide substitution

model was set as GTRGAMMA and the search algorithm was set as "Rapid bootstrapping and search

for beast-scoring ML tree", with number of bootstrap replicates = 500, parsimony random seed = 3,

and starting tree = random. The resulting tree was visualized in Geneious Prime with subsequent

modifications done in Microsoft PowerPoint.

Phylogenetic analysis of mitochondrial genomes confirmed that Kurt's maternal ancestry is

derived from a domestic horse, as his mitogenome is nested within a clade of domestic horses

representing Westphalian, Holstein, Akhal-Teke, Black Forest, and Shetland breeds, with >90%

bootstrap support across most nodes within the clade. In contrast, the four other individuals with an

expected Przewalski's horse maternal ancestry, Bars, Kuporovitch, Rosa and Vintel, are each grouped

within one of the three clades containing other Przewalski's horse mitogenomes, which are supported

by >90% bootstrap values. Notably, the new mitogenomes of Bars and Kuporovitch fall within the same

haplogroup that includes the previous mitogenomes sequenced from these two individuals, KT368744

and KT368747, respectively.

Funding Statement

The work of OAR, MLH, ASP, genome sequencing, and the cloning and early life care of cloned Przewalski's horses conducted by KW, LR, BR, SW, SSA, LA, and GV was funded by San Diego Zoo Wildlife Alliance. The work of JAI, KK, AR, and JS was supported by their respective affiliations. ASZ was funded by Saint-Petersburg State University project #95444727. The work of BN and RP was supported by the generous donors of Revive & Restore.

References

1. Kim, HM., Cho, Y., Kim, H. *et al.* Whole genome comparison of donor and cloned dogs. *Sci Rep* 3, 2998 (2013).

2. Jivanji, S., Harland, C., Cole, S. *et al.* The genomes of precision edited cloned calves show no evidence for off-target events or increased *de novo* mutagenesis. *BMC Genomics* **22**, 457 (2021).

3. Klinger, B., & Schnieke, A. 25th ANNIVERSARY OF CLONING BY SOMATIC-CELL NUCLEAR TRANSFER Twenty-five years after Dolly: how far have we come? *Reproduction*, **162**(1), F1-F10 (2021).

4. Tung TC, Wu SC, Tung YYF, Yan SY, Tu M, Lu TY . Nuclear transplantation in fish. *Science bulletin, Academia Sinica (in Chinese)* **7**:60–1 (1963).

5. Sambuichi, H. The roles of the nucleus and the cytoplasm in development I. An interspecific hybrid frog, developed from a combination of *Rana nigro-maculata nigromaculata* cytoplasm with a diploid nucleus of *Rana nigromaculata brevipoda*. *Journal of Science, Hiroshima University Ser. B. Div.* **17**:33–41 (1957).

6. Signoret, J. Transplantations nucléaires et différenciation embryonnaire. *Arch. Biol.* **71**:591–606 (1965).

7. Aimar, C. Analyse par la greffe nucléaire des propriétés morphogéné- tiques des noyaux embryonnaires chez Pleurodeles waltlii (Amphibien Urodèle). Application à l'étude de la gémellarité expérimentale. *Ann. Embryol. Morphogen.* **5**:5–42 (1972).

8. Nishioka, M. Reciprocal nucleo-cytoplasmic hybrids between Rana brevipoda and Rana plancyi
chosenica. *Scientific Report of the Laboratory for Amphibian Biology, Hiroshima University*.
(1972).

9. Liu, Z., Y. Cai, Y. Wang, Y. Nie, C. Zhang, Y. Zu, X. Zhang, Y. Lu, Z. Wang, M. Poo, and Q. Sun.
Cloning of Macaque Monkeys by Somatic Cell Nuclear Transfer. *Cell* **172**:881–887 (2018).

10. Gómez, M. C., C. E. Pope, R. H. Kutner, D. M. Ricks, L. A. Lyons, M. Ruhe, C. Dumas, J. Lyons, M.
López, B. L. Dresser, and J. Reiser. Nuclear transfer of sand cat cells into enucleated domestic
cat oocytes is affected by cryopreservation of donor cells. *Cloning and Stem Cells* **10**:469–483
(2008).

11. Loi, P., Ptak, G., Barboni, B., Fulka, J., Cappai, P., & Clinton, M. Genetic rescue of an endangered
mammal by cross-species nuclear transfer using post-mortem somatic cells. *Nat Biotechnol* **19**,
962–964 (2001).

12. Friese, C. Cloning Wildlife: Zoos, Captivity, and the Future of Endangered Animals. *New York:*
New York University Press. (2013).

13. Srirattana, K., Imsoonthornruksa, S., Laowtammathron, C., Sangmalee, A., Tunwattana, W.,
Thongprapai, T., Chaimongkol, C., Ketudat-Cairns, M., & Parnpai, R.Full-term development of
gaur-bovine interspecies somatic cell nuclear transfer embryos: Effect of Trichostatin A
treatment. *Cellular Reprogramming* **14**, 248–257 (2012).

14. Janssen, D. L., M. L. Edwards, J. A. Koster, R. P. Lanza, & O. A. Ryder. 206 Postnatal management
of cryptorchid banteng calves cloned by nuclear transfer utilizing frozen fibroblast cultures and
enculeated cow ova. *Reproduction, Fertility and Development* **16**, 224 (2004).

15. Hajian, M., Hosseini, S. M., Forouzanfar, M. , Abedi, P., Ostadhosseini, S., Hosseini, L., Moulavi,
F., Gourabi, H., Shahverdi, A.H., Taghi Dizaj, A. V., Kalantari, S. A., Fotouhi, Z., Iranpour, R.,
Mahyar, H., Amiri-Yekta, A., & Nasr-Esfahani, M. H. "Conservation cloning" of vulnerable
Esfahan mouflon (Ovis orientalis isphahanica): In vitro and in vivo studies. *European Journal of*
Wildlife Research **57**, 959–969 (2011).

16. Dehghan, S. K. Scientists in Iran clone endangered mouflon – born to domestic sheep. *The*
Guardian. https://www.theguardian.com/science/2015/aug/05/iran-scientists-clone-
endangered-mouflon-domestic-sheep. (2015).

329 17. Ryder, O. A. and K. Benirschke. The potential use of "cloning" in the conservation effort. *Zoo*
330   *Biology* **16,** 295–300 (1997).

331 18. Ryder, O.A. Cloning Advances and Challenges for Conservation. *TRENDS in Biotechnology* **20**,
332   231-232 (2002).

333 19. Ballou, J.D., Lacy, R.C., Traylor-Holzer, K., Bauman, K., Ivy, J.A. and Asa, C. Strategies for
334   establishing and using genome resource banks to protect genetic diversity in conservation
335   breeding programs. *Zoo Biology* **42,** 175–184 (2023).

336 20. Monfort, S. L., Arthur, N. P., & Wildt, D. E. Reproduction in the Przewalski's Horse. In:
337   Przewalski's Horse: The History and Biology of an Endangered Species, ed. Boyd L., & Houpt,
338   K.A. *State University of New York Press* (1994).

339 21. Bouman D.T., Bouman J.G. The history of Przewalski's Horse. In: Boyd L., Houpt D.A., editors.
340   Przewalski's Horse: The History and Biology of an Endangered Species. State University of New
341   York Press; Albany, NY, USA: 1994. pp. 5–38.

342 22. Siefert, S. Proceedings of the Fifth International Symposium on the Preservation of the
343   Przewalski Horse. Leipzig Zoological Garden, Leipzig, Germany. (1992).

344 23. Marinari, P. & Lynch, C. AZA Population Analysis & Breeding and Transfer Plan: Black-footed
345   ferret (Mustela nigripes) AZA Species Survival Plan ® Yellow Program. (2021).

346 24. Geyer, C.J., Thompson, E.A. and Ryder, O.A. Gene survival in the Asian wild horse (Equus
347   przewalskii): II. Gene survival in the whole population, in subgroups, and through history. *Zoo*
348   *Biol.*, **8**, 313-329 (1989). https://doi.org/10.1002/zoo.1430080402

349 25. Pukazhenthi, B.S., Johnson, A., Guthrie, H.D., Songsasen, N., Padilla, L.R., Wolfe, B.A., da Silva,
350   M.C., Alvarenga, M.A. & Wildt, D.E. Improved sperm cryosurvival in diluents containing amides
351   versus glycerol in the Przewalski's horse (*Equus ferus przewalskii*). *Cryobiology* **68**(2), 205-214.
352   (2014).

353 26. Mooney, A., Ryder, O. A., Houck, M. L., Staerk, J., Conde, D. A., & Buckley, Y. M. Maximizing the
354   potential for living cell banks to contribute to global conservation priorities. *Zoo Biology*, 1– 12.
355   (2023).

27. Houck M.L., Lear T.L. and Charter S.J. Animal Cytogenetics, Chapter 24. In: The AGT

Cytogenetics Laboratory Manual, 4th ed. Arsham M., Barch M. and Lawce H., (eds), pp. 1055-

1102. John Wiley & Sons, Inc., Hoboken, NJ, USA (2017).

28. Ballou JD,& Lacy RC. Identifying genetically important individuals for management of genetic

diversity in pedigreed populations. In: Ballou JD, Gilpin M, Foose TJ, editors. Population

management for survival and recovery. New York: Columbia Press. P. 76–111 (1995).

29. Fernandez, J., & Toro, M. A. The use of mathematical programming to control inbreeding in

selection schemes. *Journal of Animal Breeding and Genetics* **116**, 447–466 (1999).

30. Falino, A. Asian Wild Horse (*Equus przewalskii*) AZA Regional Studbook, Association of Zoos and

Aquariums, Silver Springs, MD. (2021).

31. Faust, L.J., Bergstrom, Y.M., Thompson, S.D., & Bier, L. PopLink Version 2.5.2. Chicago, IL:

Lincoln Park Zoo. (2018).

32. Lacy RC, Ballou JD, Pollak JP. (2012). PMx: Software package for demographic and genetic

analysis and management of pedigreed populations. Methods in Ecology and Evolution, 3(2),

433–437.

33. ZIMS for Studbooks for Przewalski's Horse. (Simek, J., Prague Zoo, Current to January 2022).

Species360 Zoological Information Management System. Retrieved from

http://zims.Species360.org

34. Bowling, A. T., Eggleston-Stott, M. L., Byrns, G., Clark, R. S., Dileanis, S., & Wictum, E. Validation

of microsatellite markers for routine horse parentage testing. *Animal Genetics* **28**(4), 247–252

(1997).

35. Goto, H., Ryder, O.A., Fisher, A.R., Schultz, B., Kosakovsky Pond, S.L., Nekrutenko, & A., Makova,

K.D. A massively parallel sequencing approach uncovers ancient origins and high genetic

variability of endangered Przewalski's horses. *Genome Biol Evol.* **3,** 1096-106 (2011).

36. Vilstrup, J.T., Seguin-Orlando, A., Stiller, M., Ginolhac, A., Raghavan, M., Nielsen, S.C.A, et al.

Mitochondrial Phylogenomics of Modern and Ancient Equids. *PLoS ONE* **8**(2), e55950 (2013).

37. Andrews, S., Krueger, F., Segonds-Pichon, A., Biggins, L., Krueger, C., & Wingett, S. FastQC. A

quality control tool for high throughput sequence data, 370 (2010).

38. Li, H. and R. Durbin. Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics **25**(14): 1754-1760 (2009).

39. Danecek, P., J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan, M. O. Pollard, A. Whitwham, T. Keane,
S. A. McCarthy, R. M. Davies and H. Li. Twelve years of SAMtools and BCFtools. *GigaScience*
10(2) giab008 (2021).

40. Broad, I. Broad Institute. Picard Tools. Broad Institute, GitHub Repository. 2021.
http://broadinstitute.github.io/picard/. (2021).

41. Poplin, R., V. Ruano-Rubio, M. A. DePristo, T. J. Fennell, M. O. Carneiro, G. A. Van der Auwera,
D. E. Kling, L. D. Gauthier, A. Levy-Moonshine, D. Roazen, K. Shakir, J. Thibault, S. Chandran, C.
Whelan, M. Lek, S. Gabriel, M. J. Daly, B. Neale, D. G. MacArthur and E. Banks. Scaling accurate
genetic variant discovery to tens of thousands of samples. *bioRxiv* 201178 (2018).

42. Gibson, T. Eulerr: Area-Proportional Euler and Venn Diagrams with Circles or Ellipses. R package
version 6.1.1. (2021). Retrieved from https://cran.r-
project.org/web/packages/eulerr/vignettes/introduction.html

43. Jin, JJ., Yu, WB., Yang, JB. et al. GetOrganelle: a fast and versatile toolkit for accurate de novo
assembly of organelle genomes. *Genome Biology* **21**, 241 (2020).

44. Pedersen, B. S., & Quinlan, A. R. Mosdepth: quick coverage calculation for genomes and
exomes. *Bioinformatics* **34**(5), 867–868 (2018).

45. Donath, A., Jühling, F., Al-Arab, M., Bernhart, S. H., Reinhardt, F., Stadler, P. F., Middendorf, M.,
& Bernt, M. Improved annotation of protein-coding genes boundaries in metazoan
mitochondrial genomes. *Nucleic Acids Research* **47**(20), 10543–10552 (2019).

46. Xu, S., Luosang, J., Hua, S., He, J., Ciren, A., Wang, W., Tong, X., Liang, Y., Wang, J., & Zheng, X.
High altitude adaptation and phylogenetic analysis of Tibetan horse based on the mitochondrial
genome. *Journal of Genetics and Genomics* **34**(8), 720–729 (2007).

47. Lippold, S., Matzke, N. J., Reissmann, M., & Hofreiter, M. (2011). Whole mitochondrial genome
sequencing of domestic horses reveals incorporation of extensive wild horse diversity during
domestication. *BMC Evolutionary Biology* **11**, 328. (2011).

48. Achilli, A., Olivieri, A., Soares, P., Lancioni, H., Hooshiar Kashani, B., Perego, U. A., Nergadze, S.
G., Carossa, V., Santagostino, M., Capomaccio, S., Felicetti, M., Al-Achkar, W., Penedo, M. C.,

413  Verini-Supplizi, A., Houshmand, M., Woodward, S. R., Semino, O., Silvestrelli, M., Giulotto, E.,

414  Pereira, L., Bandelt, H-J., Torroni, A. Mitochondrial genomes from modern horses reveal the

415  major haplogroups that underwent domestication. *Proceedings of the National Academy of*

416  *Sciences of the United States of America* **109**(7), 2449–2454 (2012).

417 49. Der Sarkissian, C., Ermini, L., Schubert, M., Yang, M. A., Librado, P., Fumagalli, M., Jónsson, H.,

418  Bar-Gal, G. K., Albrechtsen, A., Vieira, F. G., Petersen, B., Ginolhac, A., Seguin-Orlando, A.,

419  Magnussen, K., Fages, A., Gamba, C., Lorente-Galdos, B., Polani, S., Steiner, C., Neuditschko, M.,

420  Jagannathan, V., Feh, C., Greenblatt, C. L., Ludwig, A., Abramson, N. I., Zimmermann, W.,

421  Schafberg, R., Tikhonov, A., Sicheritz-Ponten, T., Willerslev, E., Marques-Bonet, T., Ryder, O. A.,

422  McCue, M., Rieder, S., Leeb, T., Slatkin, M., Orlando, L. Evolutionary Genomics and Conservation

423  of the Endangered Przewalski's Horse. *Current Biology* **25**(19), 2577–2583 (2015).

424 50. Luo, Y., Chen, Y., Liu, F., Jiang, C., & Gao, Y. Mitochondrial genome sequence of the Tibetan wild

425  ass (Equus kiang). Mitochondrial DNA, **22**(1-2), 6–8 (2011).

426 51. Katoh, K., & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:

427  Improvements in Performance and Usability. *Molecular Biology and Evolution* **30**(4), 772–780

428  (2013).

429 52. Stamatakis, A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large

430  phylogenies. *Bioinformatics* **30**(9), 1312–1313 (2014).