Single-cell transcriptomic profiling of the neonatal oviduct and uterus reveals

new insights into upper Müllerian duct regionalization

Shuai Jia[1], Fei Zhao[1]

[1]Department of Comparative Biosciences, School of Veterinary Medicine, University of

Wisconsin-Madison, Madison, WI 53706, USA

*Corresponding author:

Fei Zhao, Ph.D., Department of Comparative Biosciences, School of Veterinary Medicine,

University of Wisconsin-Madison, Madison, WI 53706, USA, Tel: 608-890-2610.

ORCID: 0000-0003-3107-1576.

E-mail address: fei.zhao@wisc.edu

Abstract

The upper Müllerian duct (MD) is patterned and specified into two

morphologically and functionally distinct organs, the oviduct and uterus. It is known that

this regionalization process is instructed by inductive signals from the adjacent

mesenchyme. However, the interaction landscape between epithelium and

mesenchyme during upper MD development remains largely unknown. Here, we

performed single-cell transcriptomic profiling of mouse neonatal oviducts and uteri at the

initiation of MD epithelial differentiation (postnatal day 3). We identified major cell types

including epithelium, mesenchyme, pericytes, mesothelium, endothelium, and immune

cells in both organs with established markers. Moreover, we uncovered region-specific

epithelial and mesenchymal subpopulations and then deduced region-specific ligand-

receptor pairs mediating mesenchymal-epithelial interactions along the craniocaudal

axis. Unexpectedly, we discovered a mesenchymal subpopulation marked by

neurofilaments with specific localizations at the mesometrial pole of both the neonatal

oviduct and uterus. Lastly, we analyzed and revealed organ-specific signature genes of

pericytes and mesothelial cells. Taken together, our study enriches our knowledge of

upper Müllerian duct development, and provides a manageable list of potential genes,

pathways, and region-specific cell subtypes for future functional studies.

Keywords: Müllerian duct; Oviduct; Uterus; Regionalization; Craniocaudal patterning

Introduction

A fundamental question in developmental biology is how a primitive epithelial tube differentiates into morphologically and functionally distinct tubular organs along the craniocaudal axis. During development, the Müllerian duct (MD), the primordium of female reproductive tract organs, undergoes craniocaudal patterning from a simple tube into four distinct organs: the oviduct, uterus, cervix, and vagina (1). During postnatal differentiation, epithelial cells in these organs acquire unique phenotypes and functions. In the upper MD, epithelial cells are committed to become oviductal ciliated and secretory cells (2); in the middle MD, they differentiate into uterine luminal and glandular epithelial cell types (3); in the lower MD, they become basal and stratified squamous epithelium in the ectocervix and vagina (4). Impairment of MD differentiation by genetic and environmental factors can potentially lead to Müllerian anomalies, a major cause of female infertility in humans (5, 6).

MD regionalization is largely dependent on instructive signals from its surrounding mesenchyme. MD mesenchyme expresses region-specific Homeobox genes for controlling MD craniocaudal patterning: *Hoxa9* in the oviduct, *Hoxa10* in the uterus, *Hoxa11* in the posterior uterus and cervix; and *Hoxa13* in the cervix and upper vagina (7, 8). Genetic ablations of these *Hox* genes expressed in the mesenchyme lead to homeotic transformation in mice, providing critical evidence that the mesenchyme governs MD epithelial fate and differentiation. For example, *Hoxa10* knockout caused partial morphological alteration of the uterus to the oviductal appearance(9); when the uterine *Hox* gene *Hoxa11* was swapped with *Hoxa13* that is normally expressed in the mesenchyme of the cervix and vagina, the uterus underwent homeotic transformation

and became similar to the cervix and vagina (10). Classic tissue recombination studies

also demonstrate the instructive function of the mesenchyme in MD regionalization.

Heterotypic tissue recombinants of uterine mesenchyme and vaginal epithelium

developed into the uterus while recombinants of vaginal mesenchyme and uterine

epithelium underwent vaginal morphogenesis (11, 12). In tissue recombinants,

developmental plasticity of epithelial cells is gradually lost approximately by 10 days

postpartum, after which epithelial cells cannot be reprogrammed (11, 12). Therefore,

region-specific mesenchymal signals at early postnatal development govern MD

craniocaudal patterning.

The paracrine signals from the mesenchyme for determining the fate and

differentiation of the middle and lower MD have been discovered. WNT4 and WNT5A in

the mesenchyme regulates the differentiation of uterine glandular epithelium

(Kelleher.et al. 2019). Activin A, BMP4 and FGF7/10 from the lower MD mesenchyme

are independently required for inducing vaginal cell fate (Terakawa. et al. 2020).

However, it is unknown what mesenchymal factor(s) determines the fate of upper MD

epithelium to become oviductal epithelia. In an effort to address this knowledge gap, we

performed single-cell mRNA profiling of the neonatal oviduct and uterus to reveal

region-specific interaction landscape between epithelium and mesenchyme at postnatal

day 3 (PND3), a critical developmental timepoint for establishing regional patterns (11-

13).

Results

Overview and characterization of major cell populations in the neonatal oviduct and uterus.

Our scRNA-seq profiling of PND3 oviducts and uteri generated a dataset with 22150 individual cells after filtering with average 3006 genes/cell (**Supplementary file 1**). Based on their transcriptomic similarities, those individual cells were classified into 8 cell populations with distinct cell-type specific markers: oviductal mesenchymal cells (*Hoxc8,* 7438 cells without expression of proliferation markers, 38%), uterine mesenchymal cells (*Hoxa10* (14)*,* 4898 cells without expression of proliferation markers, 25%), proliferating mesenchymal cells (*Top2a* (15), 2840, 14%), epithelial cells (*Epcam1(16),* 1150, 6%), endothelial cells (*Pecam1(17),* 1058, 5%), pericytes (*Rgs5(18),* 778, 4%), mesothelial cells (*Msln(19), 1139, 6%*), and myeloid/immune cells (*Lzy2(20),* 424, 2%) (**Fig. 1A-1C**). Through differential expression analysis, we identified the top 5 significantly enriched genes in each cluster **(Fig. 1D)**, some of which could serve as new cell type markers at this developmental stage. For example, *Enpp2* for oviductal mesenchymal cells, *Tcf21* for uterine mesenchymal cells (validated in the following figures), and *Wfdc2* for epithelial cells **(Fig. 1D & Supplementary file 2)**.

When UMAP was annotated based on organ origins of each individual cells, both uterine and oviductal cells were overlapped in the clusters of the epithelia, mesothelia, endothelia, pericytes and myeloid cells but separated in two distinct mesenchymal clusters (**Fig. 1B**). These results demonstrate that both neonatal oviducts and uteri are composed of the same cell types but with the largest transcriptomic difference observed in their mesenchymal cells. This observation is consistent with the notion that region-specific mesenchymal signaling plays inductive roles in MD regionalization (11, 12).

Dissecting mesenchymal cells of the neonatal oviduct and uterus.

Given the importance of the mesenchyme in directing epithelial differentiation and morphogenesis (21), we further dissected heterogeneity of the mesenchymal population, which represented the largest population (63% of cells) in the neonatal oviduct and uterus. We isolated and re-classified mesenchymal population into 7 clusters demarcated by distinct marker gene expression and organ origins (**Fig. S1, 2A & 2B; Supplementary file 3**): 2 smooth muscle subpopulations distinguished by the organ-specific expression of *Tcf21* (in the uterus) and *Nrp2* (in the oviduct); 2 mesenchymal subpopulations (*Pkib*+ and *Pcdh10*+) in the uterus; 2 mesenchymal subpopulations (*Wfdc1*+ and *Angptl7*+) in the oviduct; 1 shared population (*Nefm*+) observed in both the uterus and oviduct.

The uterine and oviductal smooth muscle clusters both expressed *Cnn1*, an established smooth muscle marker (22) (**Fig. 2B**). This observation was consistent with the previous report that smooth muscle differentiation has initiated in the prospective circular myometrial layer at PND3 in both organs (22). However, our scRNA-seq analysis uncovered previously unrecognized gene expression differences between the two smooth muscle populations. *Tcf21* is a bHLH transcription factor and plays critical roles in organogenesis (23). Its expression was enriched specifically in the uterine smooth muscle cells (**Fig. 2B & 2C**). On the other hand, the oviducal smooth muscle cells expressed *Nrp2* which encodes a transmembrane protein Neuropilin 2 (24). Based on the online scRNA-seq dataset of adult murine oviducts (25), *Tcf21* expression was still not detected but *Nrp2* continued to be expressed in the oviductal smooth muscle cells.

In addition to smooth muscle cells, there were two mesenchymal subpopulations

in either oviducts or uteri. The two mesenchymal subpopulations in oviducts were

defined by the expression of *Wfdc1* and *Angptl7*, respectively (**Fig. 2A & 2B**). We

validated their expression in the mesenchyme of cranial and caudal oviduct by

RNAscope (**Fig. 2C**). At this developmental stage, epithelial cells in the cranial oviduct

(future infundibulum and ampulla) undergo longitudinal folding while those in the cauda

region (isthmus) remain unfolded (2) (**Fig. 2C**). Both cranial and caudal regions of the

oviduct possessed *Wfdc1+* and *Anpgtl7+* mesenchymal cells, suggesting that these two

populations were not region-specific. However, it seemed that *Wfdc1+* cells were more

enriched in the mesenchyme adjacent to the epithelium of the caudal oviduct, while

Angptl7 were more enriched in the mesenchyme of the cranial oviduct (**Fig. 2C**). Using

the online scRNA-seq dataset of adult oviducts (25), we found that *Wfdc1* was still

expressed specifically in the mesenchyme-derived fibroblasts in both cranial

(infundibulum and ampulla) and cauda (isthmus and uterotubal junction) regions while

Angptl7 expression became undetected. These results demonstrate that *Wfdc1* can

serve as a specific marker for oviductal mesenchyme at both neonatal and adult stages

while *Angptl7* is a transient marker for oviductal mesenchyme at the neonatal stage.

Contrary to the oviductal mesenchyme, the uterine mesenchyme was marked by

the expression of *Tcf21* and classified into *Pkib+* and *Pcdh10+* subpopulations (**Fig. 2B**

& 2C). Previous studies have identified two closely related mesenchymal

subpopulations as inner and outer stroma depending on their proximity to the epithelial

lumen in rat uteri at PND6 (26). Markers genes for inner stroma (*Plac8* and *Bmp7)* and

outer stroma (*Apoe*) were preferentially expressed in *Pkib+* and *Pcdh10+* mesenchymal

subpopulations, respectively (**Fig.S2**). This observation suggests that two uterine

mesenchymal subpopulations at PND3 in our study may correspond to inner and outer

stromal populations at a later developmental stage. In the adult uterine mesenchyme,

Kirkwood et al., identified three stromal subpopulations, inner stroma, outer stroma, and

the third one localized in the subepithelial space (27). One of the markers (*Col6a4*) for

the subepithelial stromal subpopulation in the adult uterus was overrepresented in the

Pkib+ mesenchyme (potentially inner stroma) in our dataset (**Fig. S2**).

Finally, we identified a new mesenchymal subpopulation demarcated by unique

expression of *Nefm* and *Nefl* (**Fig. 2B & Fig. S2**). We found that *Nefm* expression was

predominantly localized at the mesometrial pole. In the oviduct, *Nefm* was found in the

mesosalpinx ligament and mesenchymal cells at the mesometrial pole. Likewise, in the

uterus, it was expressed in the mesometrium as well as mesenchymal cells at the

mesometrial pole (**Fig. 2C**). Since *Nefm* (encode neurofilament medium polypeptide)

and *Nefl* (encode neurofilament light polypeptide) are neurofilament genes (28), we

initially thought that this mesenchymal subpopulation might represent neuronal cells.

However, these neurofilament proteins are also expressed in non-neuron cells (29); and

Rbfox3 (encode NeuN (30), a neuronal nuclei marker protein*)* was not expressed in this

population at all (**Fig. S2**). Therefore, we conclude that the *Nefm*+ subpopulation

localized at the mesometrial pole is a unique mesenchymal cell subtype expressing

neurofilaments as a part of its cytoskeleton network.

Dissecting epithelial cells of the neonatal oviduct and uterus

The mature epithelium of the oviduct and uterus consists of four epithelial types:

secretory and ciliated cells in the oviduct (31, 32); and luminal and glandular epithelial

cells in the uterus (3). Establishment of these cell types from MD nascent epithelium

does not occur until PND4 (2, 33-35). When comparing epithelial cells from the uterus

and oviduct (**Fig. 3A**), we identified 390 differentially expressed genes (DEGs) and the

top marker *Msx1* for uterine epithelium and *Lrpap1* for oviductal epithelium (**Fig. 3B and**

Supplementary file 4). We confirmed the specific expression of MSX1 in the uterine

epithelium but not in the oviductal epithelium on PND3 (**Fig. 3E**). It has been reported

that MSX1 is present exclusively in lower parts of the female reproductive tract (the

uterus, cervix and vagina) (36). *Msx1* expression continues to be absent in the oviduct

at adulthood based on the published scRNA-seq dataset (25). Therefore, we concluded

that *Msx1* represents a specific gene signature of postnatal uterine epithelium.

In the oviduct, morphological differences (epithelial folding vs non-folding) in the

cranial and caudal regions were noticed at PND3 (**Fig. 2C**). In addition, it is known that

the predominant cell types lining the epithelium of the future cranial and caudal regions

are distinct, ciliated epithelial cells enriched in the cranial region while secretory

epithelial cells enriched in the caudal region (31). Therefore, we suspect there could be

at least two epithelial subtypes in the oviduct at PND3. By increasing the clustering

resolution (0.1), we were able to identify *Gsap+* and *Gsap-* epithelial subtypes at PND3

(**Fig. 3C & 3D**). By performing RNAscope, we discovered that *Gsap* expression was

restricted to the cranial oviduct epithelium and barely detected in the caudal oviduct or

uterus (**Fig.3E**). Expression of a previously-characterized marker *Wt1* for the cranial

oviduct at PND4 (34) was higher in the *Gsap+* population compared to *Gsap-* cluster

(**Supplementary file 5**). These results demonstrate the spatial locations of three

epithelial subpopulations along the craniocaudal axis at PND3: *Gsap+* epithelia in the

cranial oviduct; *Gsap-* epithelia at the caudal oviduct, and *Msx1+* uterine epithelia.

In the UMAP clustering, we noticed an outlier distant from the major oviductal

epithelial population (**Fig. S3**). When comparing the transcriptome between this outlier

and the rest of oviductal cells, we found enrichment of markers for ciliated cells (*Foxj1*

and *Fam183b* (37)) and other well-established transcription factors (*Ruvbl1 (38)* and

Myb (39)) critical for multiciliogenesis, suggesting that this outlier cluster represented

ciliated epithelial cells (**Fig. S3**). Therefore, multiciliated cell differentiation has occurred

in some of oviductal epithelium at PND3, earlier than previous reports (2, 34). In the list

of genes overrepresented in this multiciliated cell cluter (**Fig. S3 and Supplementary**

file 6), we uncovered two new transcription factors *Pbx4* and *Aes* (also known as *Tle5*),

which might potentially play crucial roles in multiciliated cell differentiation.

Interrogate ligand-receptor mediated crosstalk between epithelial and

mesenchymal cells.

After uncovering mesenchymal heterogeneity and epithelial regionalization, we

next explored the cell-cell interactions between them by leveraging Cellchat, a

bioinformatic tool that quantitatively infers and analyzes ligand-receptor mediated

intercellular communication networks (40). The cell-cell interactions between three

spatially distinct epithelial subpopulations (*Gsap+* cranial oviductal epithelium, *Gsap-*

caudal oviductal epithelium, and *Msx1+* uterine epithelium) and their corresponding

mesenchymal populations were analyzed (**Fig. S4 and Supplementary files 7-9**).

First, we inferred the signals sending from the mesenchyme to the epithelium.

We identified 157 ligand-receptor pairs in the communication from oviductal

mesenchyme to *Gsap*+ cranial oviductal epithelium, 58 from oviductal mesenchyme to

Gsap- caudal oviductal epithelium and 47 from uterine mesenchyme to uterine

epithelium (**Fig. S4 and Supplementary files 10**). In the predicted cell-cell interaction

lists, a ligand was paired with multiple receptors/coreceptors (**Supplementary file 7-9**).

To concisely visualize the potentially functional ligand-receptor pairs, we focused on

mesenchyme-derived ligands for the following criteria: 1) the knockout mice had

reproductive phenotypes and/or embryonic lethality based on Mouse Genome

Informatics (MGI) (41); 2) Ligands are enriched only in either the oviductal or uterine

mesenchyme (**Fig. S4 and Supplementary files 7-9, 11**). Therefore, we depicted a

simplified version of mesenchymal-epithelial crosstalk represented by potentially

functional region-specific ligands (**Fig. 4**).

In the ligand-mediated communication from oviductal mesenchyme to epithelium,

Cxcl12, *Edn2*, *Inhba*, *Kitl*, *Postn*, *Tgfb2* ligands were specific to *Gsap*+ cranial oviduct

while *Bmp5* and *Nrg1* from the mesenchyme can act on both *Gsap*+ and *Gsap*-

oviductal epithelium (**Fig. 4**). *Gdf7*, a ligand in the TGF-β pathway, specifically mediated

the communication from uterine mesenchyme to epithelium (**Fig. 4**). Although *Gdf7*

expression was detected in the uterine stroma (42), its functional significance in uterine

differentiation remains to be covered.

The crosstalk between the epithelium and mesenchyme is never a one-way

street (12). We therefore inferred the signals sending from the epithelium to the

mesenchyme. We identified 144 ligand-receptor pairs from *Gsap*+ oviductal epithelium

(cranial oviduct) to oviductal mesenchyme, 70 pairs from *Gsap-* oviductal epithelium

(cauda oviduct) to oviductal mesenchyme and 54 ligand-receptor pairs from uterine

epithelium to uterine mesenchyme (**Fig. S4, Supplementary file 12**). Likewise, we

consolidated and focused on region-specific ligands that were potentially functional

(**Supplementary files 7-9,11**). The region-specific ligands deriving from the epithelium

to act on adjacent mesenchyme were *Nrg1* and *Nampt* in the cranial oviduct; *Wnt5a* in

the caudal oviduct; *Vegfa* in the uterus (**Fig. 4**). *Nrg1* (Neuregulin 1) belongs to the

epidermal growth factor and plays essential roles in the heart and nervous system (43).

Nampt (nicotinamide phosphoribosyltransferase) exists in both intracellular and

extracellular forms. The intracellular NAMPT regulates NAD biosynthesis while

extracellular NAMPT is proposed to act as an insulin-mimetic cytokine (44, 45). Global

mouse knockout of either *Nrg1* or *Nampt* leads to embryonic lethality (43, 44). The

availability of *Nrg1-flox* (46) and *Nampt-flox* (47) alleles makes it possible and as a

promising direction to investigate their potential roles in MD development. Our finding of

Wnt5a-mediated communication from the epithelium to the mesenchyme in the caudal

oviduct was consistent with the report of enriched *Wnt5a* expression in the epithelium of

caudal oviducts at PND28 (33). *Vegfa* have been implicated in embryonic development

(48) and adult uterine function (49); however, its specific role in female reproductive

tract development remains to be investigated. Taken together, our data provides a list of

region-specific ligands/their activated pathways that potentially mediate epithelial-

mesenchymal communication during MD regionalization.

Inquire gene expression differences between oviductal and uterine pericytes and mesothelial cells.

Lastly, we turned our attentions to two supporting cells, pericytes (*Rgs5*+ cluster in **Fig. 1**) and mesothelial cells (*Msln*+ cluster in **Fig. 1**), which are rarely studied despite their crucial roles in organ development (50, 51). Pericytes exhibited inter-organ molecular differences (number of DEGs=186, **Supplementary file 13**). The top enriched genes in the oviductal pericytes (*Cebpb*, *Rgs16*, *Nr2f1*, *Hoxc8*, *Cryab*) and uterine pericytes (*Amer1*, *Plac8*, *Hoxd11*, *Lsp1*, *Hoxa10*) were also overrepresented in mesenchymal populations (**Fig. 5A & 5B**), suggesting that pericytes in the oviduct and the uterus might originate from organ-specific mesenchymal cells and/or acquire the organ signature after specification from other cell types (52).

Mesothelium is an extensive monolayer of squamous-like epithelial cells lining the surface of the oviduct and uterus (53, 54). Both mesothelial and epithelial cells expressed cytokeratins *Krt8*, *Krt18*, and *Krt19* (**Fig. 6A**). However, mesothelial cells did not expression luminal epithelium marker *Epcam* and *Mecom* but instead exhibited the expression of *Msln* (19) and *Upk3b* (55) in both the oviduct and uterus (**Fig. 1B & Fig. 6A**). Transcriptomic analysis between oviductal and uterine mesothelial cells identified 260 DEGs (**Supplementary file 14**), including the top genes enriched in the oviductal mesothelium (*Lgals7*, *Igf1*, *Mt1*, *Ccl2*, and *Nrgn)* and uterine mesothelium (*Alcam*, *Rspo3*, *Fxyd3*, *Tcf21*, and *Hoxd11)* (**Fig. 6B**). It has been reported that *Igf1* is enriched in the cranial MD (future oviduct) (56). Our *Tcf21* RNAscope staining confirmed the enriched expression of *Tcf21* in the mesothelial layer of the uterus but not in the oviduct

(**Fig. 2C**). These results validated organ-specific signature genes for mesothelial cells in

the neonatal oviduct and uterus.

Discussion

Our study presented a single-cell atlas of the murine oviduct and uterus at PND3,

a critical developmental timepoint for establishing regional differentiation of MD

epithelium (11-13). Our major findings on upper MD regionalization were summarized in

Fig. 7.

Region-specific transcriptional signatures of the epithelium and mesenchyme in

the upper Müllerian duct.

Uncovering three epithelial subpopulations (*Gsap+* oviduct, *Gsap-* oviduct, *Msx+*

uterine epithelium) along the craniocaudal axis at PND3 is an important finding in our

study. Our results were consistent with the notion that the cranial and caudal oviducts

were composed of two distinct lineages, respectively (57). MD epithelium at this

developmental stage (PND3) is still uncommitted and possesses developmental

plasticity till PND10. When recombined with mesenchyme, MD epithelium can be

reprogrammed into epithelial cell types corresponding to the origin of recombined

mesenchyme. For example, heterotypic recombinant tissues of cranial oviductal

epithelium and caudal oviductal mesenchyme from PND3 neonates formed the typical

cell composition pattern of the caudal oviduct (the majority is the secretory cell and few

is the ciliated cell) (13). These observations suggest that while developmental fate of

MD epithelium remains plastic, regional transcriptional differences have already arisen,

which might play critical roles in MD regional patterning. Our identification of *Gsap* as

the cranial oviductal epithelium marker at PND3 is noteworthy. The predominant

number of epithelium in the cranial oviduct differentiate into multiciliate cells, which

requires NOTCH signaling inhibition (58). GSAP functions as gamma-secretase

activating protein that can switch the form of gamma-secretase away from Notch

cleavages (59).

We identified *Msx1* as an uterine signature gene which encodes a homeobox

protein functioning as a transcriptional repressor during embryogenesis (60). Previous

studies have shown that MSX1 expression is detected not only in uterine epithelium but

also vaginal epithelium (36). However, MSX1 expression in the vagina occurred only

within the first week of postnatal development (36). Because vaginal epithelium can be

elicited by uterine mesenchyme to become uterine epithelium only during the first week

of postnatal development, *Msx1* was proposed to play a role in maintaining

developmental responsiveness to become uterine epithelium (36). In the mature uterus,

Msx1 and Msx2 function redundantly in establishing epithelial polarity and receptivity for

embryo implantation (61, 62). Therefore, these previous results along with our

observations cement the notion that *Msx1* is a master regulator of various aspects of

uterine epithelial development and function.

Although our dataset was powerful for comparing oviductal and uterine

epithelium, it contained a relatively low number of uterine epithelia, which prevented us

from further probing uterine epithelial heterogeneity at PND3. Single-cell mRNA profiling

of neonatal mouse uterine epithelium at a relatively larger scale have been performed

and revealed a varied number of epithelial subtypes depending on the developmental

stage (63-65). These datasets together provide a wealth of information on intrinsic

regulatory and signaling factors for regulating upper MD epithelial regionalization and

postnatal uterine differentiation.

Our dataset also revealed distinct gene signatures for oviductal and uterine

mesenchymes and their derived smooth muscles. We used region-specific Hox genes

to distinguish oviductal and uterine mesenchymes at PND3 (7, 8). Since region-specific

Hox genes are established before birth (66), these gene differences observed at PND3

might result from transcriptional regulation of region-specific *Hox* genes. Smooth muscle

can be a mechanical sculptor for governing epithelial morphogenesis (67, 68). Those

differential gene expression between oviductal and uterine smooth muscles might be

implicated in regulating epithelial morphogenesis of these two organs: oviductal

epithelia undergoes folding in the luminal side along with tubal looping while uterine

epithelia are about developing epithelial budding into the surrounding stroma for forming

epithelial glands. A transcription factor *Tcf21* is specifically expressed in the uterine

smooth muscle. It has been reported that *Tcf21* in the mesenchyme is crucial for kidney

morphogenesis (69) thus it might be interesting to determine its potential role in uterine

development.

Decoding organ-specific mesenchymal factors that potentially determine the fate

of oviductal and uterine epithelia.

The specific mesenchyme-derived factor(s) for directing fate decisions of

oviductal or uterine epithelia *in vivo* has yet been identified. A previous study proposed

that a BMP antagonist, follistatin-like-1 (*Fstl1*), was the diffusible mesenchymal factor

for transforming nascent epithelial cells to oviductal ciliated cells in vitro mouse

oviductal cell lines (70). However, *Fstl1* was not restricted to the oviduct in our dataset,

consistent with its reported expression in the uterine region (70). In addition, *Fstl1*

knockout mice died from defective lung development shortly after birth without any

reports of reproductive tract abnormalities (71). The recent development of *Fstl1-flox*

allele has made it feasible to determine its organ-specific role in oviductal epithelial

differentiation in vivo (71, 72).

Our identifications of *Inhba* and *Gdf7* as oviduct and uterus specific

mesenchymal secreted factors are intriguing. Both ligands belong to TGF-β superfamily.

Inhba encodes inhibin βA, an essential subunit of activins and inhibin A (antagonist of

activins) (73). *Inhba* is expressed in the mesonephric mesenchyme before sexual

differentiation and exhibits craniocaudal expression gradients in early mesonephros

(74). *Gdf7* encodes growth differentiation factor 7 and its expression is enriched in

uterine stromal cells (75). In males, *Inhba* and *Gdf7* are critical for epithelial

morphogenesis and differentiation of cranial and caudal Wolffian duct (primordium of

the male reproductive tract), respectively (76, 77). Therefore, their significant roles in

cranial and caudal MD development in females are worthy of future investigations.

A novel mesenchymal subpopulation at the mesometrial pole.

An advantage of single cell RNA-seq profiling of multicellular tissues is its

capability of identifying rare cell types or subtypes that have been overlooked in the bulk

tissue studies. Here, we discovered a neurofilament-positive (*Nefm*+ and *Nefl*+)

mesenchymal subpopulation. As major cytoskeletal components in neuron cells,

neurofilaments have been detected only in two non-neuron cells: the embryonic renal

mesenchyme (78) and adrenal progenitor cells (29). Therefore, the expression of

neurofilaments in one specific mesenchymal subpopulation of the upper MD is quite

unexpected. In addition, *Nefm+* subpopulation resided in a unique localization, the

mesometrial pole of both the oviduct and uterus. It has been noticed that the neural

crest cells invade into the mesometrial pole of the Müllerian duct (79). Also, during

sexual differentiation, Wolffian ducts degenerate in females however the mesenchyme

surrounding Wolffian ducts is maintained and contributes to mesenchymal cells at the

mesometrial side of the uterus (56). These observations suggest that *Nefm+*

mesenchymal subpopulation might derive from neural crest cells and/or Wolffian duct

mesenchyme.

Extend our understanding of the less-studied mesothelial cells.

Mesothelial cells have been implicated in female reproductive tract diseases,

such as ectopic pregnancy and endometriosis (80, 81). In contrast to the MD epithelium

lining the inside lumen, mesothelial cells form the single epithelial layer outside the

female reproductive tract. Despite the opposite lining positions of mesothelial cells and

MD epithelium, they have a common ancestor, coelomic epithelium (1). Coelomic

epithelium is the primitive mesothelial cell lining the outside surface of the

mesonephros. During early development, coelomic epithelium undergoes specification

and invagination to establish the cranial portion of MD, which then extend caudally to

form the entire MD epithelial tube (1, 82). We observed gene expression similarity and

differences between epithelium and mesothelium, consistent with the notion that they

are two lineages derived from the same progenitor. In human fallopian tubes (the

oviduct), mesothelial cells were distinguished with two markers *CALB2* and *LRRN4*

(83), which are different from the mesothelium marker *Msln* used in our mouse study.

MSLN was instead enriched in human oviductal secretory cells (83-85), indicating

species differences in mesothelium gene signatures. We also observed gene

expression differences between uterine and oviductal mesothelial cells. During

postnatal development, the oviduct becomes elongated and coiled while the uterus

remains a straight tube with increased diameter. As the cellular layer lining the organ

surface, mesothelial cells presumably need to undergo differential gene programs and

morphological changes. For example, oviductal mesothelium had higher expression of

Igf1 which has been reported to enhance growth of oviductal organoids (86).

In summary, using scRNA-seq, we have dissected region-specific gene

signatures of the epithelium and mesenchyme and deduced ligand-receptor mediated

mesenchymal-epithelial interactions in the upper Müllerian duct at a critical

developmental stage. We have discovered a unique mesenchymal subpopulation

expressing neurofilaments and provided further understanding of pericyte and

mesothelium in the oviduct and uterus. However, there are several limitations of this

study. Single cell transcriptomic profiling of early or later developmental stages is

absent for generating trajectory of epithelial and mesenchymal lineage progression. The

role of specific subpopulations, signature genes or paracrine factors needs further

confirmation by cell or gene deletions. Therefore, future studies would be necessary for

comprehensively elucidating functional gene regulatory network and cell lineage

differentiation in Müllerian duct development.

Methods.

Mice. CD1 (ICR, Strain Code: 022) mice were purchased from Charles River. The day

of birth was determined by pup delivery and designated as postnatal day 0 (PND0). All

mouse procedures were approved by the University of Wisconsin-Madison (UW-

Madison) Animal Care and Use Committees and were in compliance with UW-Madison

approved animal study proposals and public laws.

Tissue dissociation and preparation of single-cell suspension. The upper female

reproductive tracts from 7 female neonates were collected freshly on PND3, when the

junction between the uterus and oviduct, known as the uterotubal junction, is formed (2,

87). The connective tissues were carefully removed using a 27G needle in a petri dish

with cool 1x PBS. Tracts from 3 and 4 neonatal females were pooled as two biological

replicates (N=2), respectively. Oviducts and uteri were then separated at the uterotubal

junction (Fig. 1A), generating four groups: oviducts (A1 and A2) and uteri (B1 and B2).

Oviducts or uteri in each group were fragmented into small pieces with a 27 G needle

and then transferred into a 1.5 mL tube with 250 µL dissociation media [0.04% BSA

(DOT Scientific, DSA30075-25), 1.2 U/mL Dispase II (Sigma, D4693-1G), 1 mg/mL

Collagenase B (Sigma, 11088807001), 5 U/mL DNase I (Sigma, DN25-100MG) in 1 mL

of 1x PBS (Gibco, 14040-133)]. Tissues were dissociated at 37 ºC for 20 min on an

orbital shaker (500 rpm). All four samples were pipetted up and down every 10 min with

a 200 µL pipette until there were no observable pieces of tissue by naked eyes. The

dissociation enzyme reactions were quenched by adding 1 µL 0.5 M EDTA. Media was

removed by centrifugation at 500 g for 5 min at 4 ºC, the supernatant discarded, and

pellet resuspend in 500 µL 1x PBS+0.04% BSA. Samples were transferred to 5 mL

falcon tube while passing through a 35 µm cell strainer (Corning, 352235). Suspended

cells were maintained on ice and sent to the Biotechnology Center at the UW-Madison

for quality control (cell viability 76% (A1), 83% (A2), 77% (B1), 81% (B2)) and

proceeded to 10x Genomics Single Cell mRNA Sequencing protocols.

Single cell RNA-seq (scRNA-seq) library preparation, sequencing and alignments.

Single cell sequencing libraries were prepared using Chromium Single Cell 3' v3.1

Reagents Kits (10X Genomics) according to manufacturer's instructions and sequenced

on a NovaSeq 6000 platform using a S1 2 x 50bp flowcell. Raw data fastq files from the

Illumina NovaSeq 6000 platform were aligned using default 10x Cell Ranger

parameters.

Single-cell data analysis using Seurat R package (V4.2.3)(88). Cells with 200 <

nFeature_RNA (genes per cell) < 8000, percent.mt (mitochondrial genes) < 20, and

percent.Hba & Hbb (blood cell genes) < 0.5 were used for downstream analyses. Genes

expressed in fewer than 3 cells were excluded from downstream analyses. The data

was then normalized using a global-scaling normalization method "LogNormalize" with

default setting. Highly variable features were determined by employing

FindVariableFeatures function to return 2,000 features per dataset. The top 2000

highest variable features were used for the principal component analysis (PCA) and the

optimal number (10 PCs) of PCA components was determined by the Elbow procedure.

Single cells were clustered by the K-nearest neighbor (KNN) graph algorithm in PCA

space; the dimension was reduced by the Uniform Manifold Approximation and

Projection (UMAP) to visualize cell clusters. Marker genes for each cluster were

identified by FindAllMarkers function (genes were detected in a minimum of 25% cells,

and logFC threshold was set at 0.25). The cell type identity for each cluster was

manually assigned based on established cell markers. Differentially expressed genes

between clusters/subclusters were identified using the Wilcoxon rank-sum test under

the Findmarkers function and then filtered by setting the adjusted p value < 0.05 and the

absolute log2FC > 0.3.

The epithelial and mesenchymal clusters were isolated by the Subset function

and then prefiltered, renormalized and reclustered with the following cutoffs and

parameters: mesenchymal cluster: 200 < nFeature_RNA < 7000, percent.mt < 10,

percent.Hba < 0.2, percent.Hbb < 0.2, UMAP resolution= 0.3; epithelial cluster: 200 <

nFeature_RNA < 8000, percent.mt < 20, percent.Hba < 0.4, percent.Hbb < 0.4, UMAP

resolution= 0.1. *Top2a*+ proliferating cells were excluded for eliminating the effects of

different cell cycle states on reclustering and analysis of differentially expressed gene

(89, 90). CellChat R package(40) was used to deduce ligand-receptor interactions

between mesenchymal and epithelium cells. Average gene expression per cell cluster

was calculated by using "computeCommuProb" function with the "truncatedMean"

method by setting the trim value as 0.05 (5%) for Communication probabilities inferring.

RNA-scope. The manufacturer's protocol was followed as previously described (56).

Briefly, paraffin sections were treated with antigen retrieval buffer and proteinase.

Sections were hybridized with the following individual probes (*Angptl7,* Cat #: 552821;

Wfdc1, Cat #: 471331; *Tcf21,* Cat #: 508661; *Nefm,* Cat #: 315611) and subjected to

rinsing and signaling detection at designated temperatures.

Immunofluorescence. Paraffin sections were subjected to antigen retrieval (VECTOR,

H-3300) using a microwave oven. After washing with PBST (1x PBS with 0.1% Triton X-

100), sections were incubated in a blocking buffer (5% normal donkey serum in PBST)

for 1 hour and then incubated with a primary antibody in blocking buffer overnight at

4 °C. On the next day, sections were washed three times with PBST and incubated with

secondary antibodies for 1 hour at room temperature and counterstained with DAPI

(Thermo Scientific, 62248,1:1000). The following primary and secondary antibodies

were used for immunofluorescence staining: Goat anti-human MSX1 Met1 Thr165

(R&D, AF5045, 1:200), Donkey anti-Goat IgG (H+L) Cross-Adsorbed Secondary

Antibody, Alexa Fluor™ 488 (Invitrogen, A11055, 1:200).

Acknowledgments

We are thankful to Animal Care Staff at the School of Veterinary Medicine for

taking care of our mice. The author(s) utilized the University of Wisconsin – Madison

Biotechnology Center's Gene Expression Center Core Facility (Research Resource

Identifier - RRID:SCR_017757) for scRNA library preparation and the DNA Sequencing

Facility (RRID:SCR_017759) for sequencing. We appreciate Dr. Jayshree Samanta's

suggestions on neuron cell and Dr. Shuo Xiao at Rutgers University for his help with our

initial data analysis.

Funding Sources

National Institute of Child Health and Development R00HD096051 and R01HD111425

(FZ).

Author Contributions

SJ and FZ conceived the study; FZ supervised the project; SJ performed all

experiments and analyzed the data. SJ and FZ wrote the manuscript. All authors read

and approved the final manuscript.

Competing interests:

Authors declare that they have no competing interests.

Data and materials availability:

All data are available in the main text or the supplementary materials upon reasonable

request. scRNA-seq data have been deposited in the GEO database under the

accession code GEO XXXXXX.

References:

1.   Mullen, R. D., and Behringer, R. R. (2014) Molecular genetics of Mullerian duct formation,
regression and differentiation. *Sex Dev* **8**, 281-296
2.   Stewart, C. A., and Behringer, R. R. (2012) Mouse oviduct development. *Results Probl Cell Differ*
55, 247-262
3.   Kelleher, A. M., DeMayo, F. J., and Spencer, T. E. (2019) Uterine Glands: Developmental Biology
and Functional Roles in Pregnancy. *Endocr Rev* **40**, 1424-1445
4.   Kurita, T. (2010) Developmental origin of vaginal epithelium. *Differentiation* **80**, 99-105
5.   Robboy, S. J., Szyfelbein, W. M., Goellner, J. R., Kaufman, R. H., Taft, P. D., Richard, R. M., Gaffey,
T. A., Prat, J., Virata, R., Hatab, P. A., McGorray, S. P., Noller, K. L., Townsend, D., Labarthe, D.,
and Barnes, A. B. (1981) Dysplasia and cytologic findings in 4,589 young women enrolled in
diethylstilbestrol-adenosis (DESAD) project. *Am J Obstet Gynecol* **140**, 579-586
6.   Connell, M., Owen, C., and Segars, J. (2013) Genetic Syndromes and Genes Involved in the
Development of the Female Reproductive Tract: A Possible Role for Gene Therapy. *J Genet Syndr*
Gene Ther **4**
7.   Taylor, H. S., VandenHeuvel, G. B., and Igarashi, P. (1997) A conserved Hox axis in the mouse and
human female reproductive system: Late establishment and persistent adult expression of the
Hoxa cluster genes. *Biol Reprod* **57**, 1338-1345
8.   Ma, L., Benson, G. V., Lim, H., Dey, S. K., and Maas, R. L. (1998) Abdominal B (AbdB) Hoxa genes:
regulation in adult uterus by estrogen and progesterone and repression in mullerian duct by the
synthetic estrogen diethylstilbestrol (DES). *Dev Biol* **197**, 141-154
9.   Benson, G. V., Lim, H., Paria, B. C., Satokata, I., Dey, S. K., and Maas, R. L. (1996) Mechanisms of
reduced fertility in Hoxa-10 mutant mice: uterine homeosis and loss of maternal Hoxa-10
expression. *Development* **122**, 2687-2696
10.   Zhao, Y., and Potter, S. S. (2001) Functional specificity of the Hoxa13 homeobox. *Development*
128, 3197-3207
11.   Cunha, G. R. (1976) Stromal induction and specification of morphogenesis and
cytodifferentiation of the epithelia of the Mullerian ducts and urogenital sinus during
development of the uterus and vagina in mice. *J Exp Zool* **196**, 361-370
12.   Kurita, T., Cooke, P. S., and Cunha, G. R. (2001) Epithelial-stromal tissue interaction in
paramesonephric (Mullerian) epithelial differentiation. *Dev Biol* **240**, 194-211
13.   Yamanouchi, H., Umezu, T., and Tomooka, Y. (2010) Reconstruction of oviduct and
demonstration of epithelial fate determination in mice. *Biol Reprod* **82**, 528-533
14.   Taylor, H. S., Vanden Heuvel, G. B., and Igarashi, P. (1997) A conserved Hox axis in the mouse
and human female reproductive system: late establishment and persistent adult expression of
the Hoxa cluster genes. *Biol Reprod* **57**, 1338-1345
15.   Whitfield, M. L., George, L. K., Grant, G. D., and Perou, C. M. (2006) Common markers of
proliferation. *Nat Rev Cancer* **6**, 99-106
16.   Huang, L., Yang, Y., Yang, F., Liu, S., Zhu, Z., Lei, Z., and Guo, J. (2018) Functions of EpCAM in
physiological processes and diseases (Review). *Int J Mol Med* **42**, 1771-1785
17.   Privratsky, J. R., and Newman, P. J. (2014) PECAM-1: regulator of endothelial junctional integrity.
Cell Tissue Res **355**, 607-619
18.   Cho, H., Kozasa, T., Bondjers, C., Betsholtz, C., and Kehrl, J. H. (2003) Pericyte-specific expression
of Rgs5: implications for PDGF and EDG receptor signaling during vascular maturation. *FASEB J*
17, 440-442
19.   Rinkevich, Y., Mori, T., Sahoo, D., Xu, P. X., Bermingham, J. R., Jr., and Weissman, I. L. (2012)
Identification and prospective isolation of a mesothelial precursor lineage giving rise to smooth

596   muscle cells and fibroblasts for mammalian internal organs, and their vasculature. *Nat Cell Biol*
597   **14**, 1251-1260

598 20.   Menon, M. B., Yakovleva, T., Ronkina, N., Suwandi, A., Odak, I., Dhamija, S., Sandrock, I.,
599   Hansmann, F., Baumgartner, W., Forster, R., Kotlyarov, A., and Gaestel, M. (2021) Lyz2-Cre-
600   Mediated Genetic Deletion of Septin7 Reveals a Role of Septins in Macrophage Cytokinesis and
601   Kras-Driven Tumorigenesis. *Front Cell Dev Biol* **9**, 795798

602 21.   Cunha, G. R. (2008) Mesenchymal-epithelial interactions: past, present, and future.
603   *Differentiation* **76**, 578-586

604 22.   Gao, Y., Bayless, K. J., and Li, Q. (2014) TGFBR1 is required for mouse myometrial development.
605   *Mol Endocrinol* **28**, 380-394

606 23.   Acharya, A., Baek, S. T., Huang, G., Eskiocak, B., Goetsch, S., Sung, C. Y., Banfi, S., Sauer, M. F.,
607   Olsen, G. S., Duffield, J. S., Olson, E. N., and Tallquist, M. D. (2012) The bHLH transcription factor
608   Tcf21 is required for lineage-specific EMT of cardiac fibroblast progenitors. *Development* **139**,
609   2139-2149

610 24.   Bielenberg, D. R., Seth, A., Shimizu, A., Pelton, K., Cristofaro, V., Ramachandran, A., Zwaans, B.
611   M., Chen, C., Krishnan, R., Seth, M., Huang, L., Takashima, S., Klagsbrun, M., Sullivan, M. P., and
612   Adam, R. M. (2012) Increased smooth muscle contractility in mice deficient for neuropilin 2. *Am*
613   *J Pathol* **181**, 548-559

614 25.   McGlade, E. A., Herrera, G. G., Stephens, K. K., Olsen, S. L. W., Winuthayanon, S., Guner, J.,
615   Hewitt, S. C., Korach, K. S., DeMayo, F. J., Lydon, J. P., Monsivais, D., and Winuthayanon, W.
616   (2021) Cell-type specific analysis of physiological action of estrogen in mouse oviducts. *FASEB J*
617   **35**, e21563

618 26.   Saatcioglu, H. D., Kano, M., Horn, H., Zhang, L., Samore, W., Nagykery, N., Meinsohn, M. C.,
619   Hyun, M., Suliman, R., Poulo, J., Hsu, J., Sacha, C., Wang, D., Gao, G., Lage, K., Oliva, E., Morris
620   Sabatini, M. E., Donahoe, P. K., and Pepin, D. (2019) Single-cell sequencing of neonatal uterus
621   reveals an Misr2+ endometrial progenitor indispensable for fertility. *Elife* **8**

622 27.   Kirkwood, P. M., Gibson, D. A., Smith, J. R., Wilson-Kanamori, J. R., Kelepouri, O., Esnal-Zufiaurre,
623   A., Dobie, R., Henderson, N. C., and Saunders, P. T. K. (2021) Single-cell RNA sequencing
624   redefines the mesenchymal cell landscape of mouse endometrium. *FASEB J* **35**, e21285

625 28.   Yuan, A., Rao, M. V., Veeranna, and Nixon, R. A. (2017) Neurofilaments and Neurofilament
626   Proteins in Health and Disease. *Cold Spring Harb Perspect Biol* **9**

627 29.   Neirijnck, Y., Sararols, P., Kuhne, F., Mayere, C., Weerasinghe Arachchige, L. C., Regard, V., Nef,
628   S., and Schedl, A. (2023) Single-cell transcriptomic profiling redefines the origin and specification
629   of early adrenogonadal progenitors. *Cell Rep* **42**, 112191

630 30.   Gusel'nikova, V. V., and Korzhevskiy, D. E. (2015) NeuN As a Neuronal Nuclear Antigen and
631   Neuron Differentiation Marker. *Acta Naturae* **7**, 42-47

632 31.   Li, S., and Winuthayanon, W. (2017) Oviduct: roles in fertilization and early embryo
633   development. *J Endocrinol* **232**, R1-R26

634 32.   Spencer, T. E., Dunlap, K. A., and Filant, J. (2012) Comparative developmental biology of the
635   uterus: insights into mechanisms and developmental disruption. *Mol Cell Endocrinol* **354**, 34-53

636 33.   Ghosh, A., Syed, S. M., and Tanwar, P. S. (2017) In vivo genetic cell lineage tracing reveals that
637   oviductal secretory cells self-renew and give rise to ciliated cells. *Development* **144**, 3031-3041

638 34.   Ford, M. J., Harwalkar, K., Pacis, A. S., Maunsell, H., Wang, Y. C., Badescu, D., Teng, K.,
639   Yamanaka, N., Bouchard, M., Ragoussis, J., and Yamanaka, Y. (2021) Oviduct epithelial cells
640   constitute two developmentally distinct lineages that are spatially separated along the distal-
641   proximal axis. *Cell Rep* **36**, 109677

642 35.   Komatsu, M., and Fujita, H. (1978) Electron-microscopic studies on the development and aging
643   of the oviduct epithelium of mice. *Anat Embryol (Berl)* **152**, 243-259

36. Pavlova, A., Boutin, E., Cunha, G., and Sassoon, D. (1994) Msx1 (Hox-7.1) in the adult mouse
uterus: cellular interactions underlying regulation of expression. *Development* **120**, 335-345
37. Beckers, A., Ott, T., Schuster-Gossler, K., Boldt, K., Alten, L., Ueffing, M., Blum, M., and Gossler,
A. (2018) The evolutionary conserved FOXJ1 target gene Fam183b is essential for motile cilia in
Xenopus but dispensable for ciliary function in mice. *Sci Rep* **8**, 14678
38. Dafinger, C., Rinschen, M. M., Borgal, L., Ehrenberg, C., Basten, S. G., Franke, M., Hohne, M.,
Rauh, M., Gobel, H., Bloch, W., Wunderlich, F. T., Peters, D. J. M., Tasche, D., Mishra, T., Habbig,
S., Dotsch, J., Muller, R. U., Bruning, J. C., Persigehl, T., Giles, R. H., Benzing, T., Schermer, B., and
Liebau, M. C. (2018) Targeted deletion of the AAA-ATPase Ruvbl1 in mice disrupts ciliary
integrity and causes renal disease and hydrocephalus. *Exp Mol Med* **50**, 1-17
39. Tan, F. E., Vladar, E. K., Ma, L., Fuentealba, L. C., Hoh, R., Espinoza, F. H., Axelrod, J. D., Alvarez-
Buylla, A., Stearns, T., Kintner, C., and Krasnow, M. A. (2013) Myb promotes centriole
amplification and later steps of the multiciliogenesis program. *Development* **140**, 4277-4286
40. Jin, S., Guerrero-Juarez, C. F., Zhang, L., Chang, I., Ramos, R., Kuan, C. H., Myung, P., Plikus, M. V.,
and Nie, Q. (2021) Inference and analysis of cell-cell communication using CellChat. *Nat*
Commun **12**, 1088
41. Smith, C. L., and Eppig, J. T. (2009) The mammalian phenotype ontology: enabling robust
annotation and comparative analysis. *Wiley Interdiscip Rev Syst Biol Med* **1**, 390-399
42. Rekker, K., Saare, M., Eriste, E., Tasa, T., Kukuskina, V., Roost, A. M., Anderson, K., Samuel, K.,
Karro, H., Salumets, A., and Peters, M. (2017) High-throughput mRNA sequencing of stromal
cells from endometriomas and endometrium. *Reproduction* **154**, 93-100
43. Meyer, D., and Birchmeier, C. (1995) Multiple essential functions of neuregulin in development.
Nature **378**, 386-390
44. Revollo, J. R., Korner, A., Mills, K. F., Satoh, A., Wang, T., Garten, A., Dasgupta, B., Sasaki, Y.,
Wolberger, C., Townsend, R. R., Milbrandt, J., Kiess, W., and Imai, S. (2007) Nampt/PBEF/Visfatin
regulates insulin secretion in beta cells as a systemic NAD biosynthetic enzyme. *Cell Metab* **6**,
363-375
45. Audrito, V., Messana, V. G., and Deaglio, S. (2020) NAMPT and NAPRT: Two Metabolic Enzymes
With Key Roles in Inflammation. *Front Oncol* **10**, 358
46. Yang, X., Arber, S., William, C., Li, L., Tanabe, Y., Jessell, T. M., Birchmeier, C., and Burden, S. J.
(2001) Patterning of muscle acetylcholine receptor gene expression in the absence of motor
innervation. *Neuron* **30**, 399-410
47. Rongvaux, A., Galli, M., Denanglaire, S., Van Gool, F., Dreze, P. L., Szpirer, C., Bureau, F., Andris,
F., and Leo, O. (2008) Nicotinamide phosphoribosyl transferase/pre-B cell colony-enhancing
factor/visfatin is required for lymphocyte development and cellular resistance to genotoxic
stress. *J Immunol* **181**, 4685-4695
48. Ferrara, N., Carver-Moore, K., Chen, H., Dowd, M., Lu, L., O'Shea, K. S., Powell-Braxton, L., Hillan,
K. J., and Moore, M. W. (1996) Heterozygous embryonic lethality induced by targeted
inactivation of the VEGF gene. *Nature* **380**, 439-442
49. Kim, M., Park, H. J., Seol, J. W., Jang, J. Y., Cho, Y. S., Kim, K. R., Choi, Y., Lydon, J. P., Demayo, F.
J., Shibuya, M., Ferrara, N., Sung, H. K., Nagy, A., Alitalo, K., and Koh, G. Y. (2013) VEGF-A
regulated by progesterone governs uterine angiogenesis and vascular remodelling during
pregnancy. *EMBO Mol Med* **5**, 1415-1430
50. Li, S. Y., Bhandary, B., Gu, X., and DeFalco, T. (2022) Perivascular cells support folliculogenesis in
the developing ovary. *Proc Natl Acad Sci U S A* **119**, e2213026119
51. Koopmans, T., and Rinkevich, Y. (2018) Mesothelial to mesenchyme transition as a major
developmental and pathological player in trunk organs and their cavities. *Commun Biol* **1**, 170

52. Yamazaki, T., and Mukouyama, Y. S. (2018) Tissue Specific Origin, Development, and
Pathological Perspectives of Pericytes. *Front Cardiovasc Med* **5**, 78
53. Mutsaers, S. E. (2004) The mesothelial cell. *Int J Biochem Cell Biol* **36**, 9-16
54. Nicosia, S. V., and Johnson, J. H. (1984) Surface morphology of ovarian mesothelium (surface
epithelium) and of other pelvic and extrapelvic mesothelial sites in the rabbit. *Int J Gynecol*
Pathol **3**, 249-260
55. Kanamori-Katayama, M., Kaiho, A., Ishizu, Y., Okamura-Oho, Y., Hino, O., Abe, M., Kishimoto, T.,
Sekihara, H., Nakamura, Y., Suzuki, H., Forrest, A. R., and Hayashizaki, Y. (2011) LRRN4 and
UPK3B are markers of primary mesothelial cells. *PLoS One* **6**, e25391
56. Zhao, F., Grimm, S. A., Jia, S., and Yao, H. H. (2022) Contribution of the Wolffian duct
mesenchyme to the formation of the female reproductive tract. *PNAS Nexus* **1**, pgac182
57. Ford, M. J., Harwalkar, K., Pacis, A. S., Maunsell, H., Wang, Y. C., Badescu, D., Teng, K. T.,
Yamanaka, N., Bouchard, M., Ragoussis, J., and Yamanaka, Y. (2021) Oviduct epithelial cells
constitute two developmentally distinct lineages that are spatially separated along the distal-
proximal axis. *Cell Reports* **36**
58. Brooks, E. R., and Wallingford, J. B. (2014) Multiciliated cells. *Curr Biol* **24**, R973-982
59. Wong, E., Liao, G. P., Chang, J. C., Xu, P., Li, Y. M., and Greengard, P. (2019) GSAP modulates
gamma-secretase specificity by inducing conformational change in PS1. *Proc Natl Acad Sci U S A*
116, 6385-6390
60. Wang, J., and Abate-Shen, C. (2012) Transcriptional repression by the Msx1 homeoprotein is
associated with global redistribution of the H3K27me3 repressive mark to the nuclear periphery.
Nucleus **3**, 155-161
61. Daikoku, T., Cha, J., Sun, X., Tranguch, S., Xie, H., Fujita, T., Hirota, Y., Lydon, J., DeMayo, F.,
Maxson, R., and Dey, S. K. (2011) Conditional deletion of Msx homeobox genes in the uterus
inhibits blastocyst implantation by altering uterine receptivity. *Dev Cell* **21**, 1014-1025
62. Nallasamy, S., Li, Q., Bagchi, M. K., and Bagchi, I. C. (2012) Msx homeobox genes critically
regulate embryo implantation by controlling paracrine signaling between uterine stroma and
epithelium. *PLoS Genet* **8**, e1002500
63. Wu, B., An, C., Li, Y., Yin, Z., Gong, L., Li, Z., Liu, Y., Heng, B. C., Zhang, D., Ouyang, H., and Zou, X.
(2017) Reconstructing Lineage Hierarchies of Mouse Uterus Epithelial Development Using
Single-Cell Analysis. *Stem Cell Reports* **9**, 381-396
64. Mucenski, M. L., Mahoney, R., Adam, M., Potter, A. S., and Potter, S. S. (2019) Single cell RNA-
seq study of wild type and Hox9,10,11 mutant developing uterus. *Sci Rep* **9**, 4557
65. Spencer, T. E., Lowke, M. T., Davenport, K. M., Dhakal, P., and Kelleher, A. M. (2023) Single-cell
insights into epithelial morphogenesis in the neonatal mouse uterus. *Proc Natl Acad Sci U S A*
120, e2316410120
66. Nakajima, T., Iguchi, T., and Sato, T. (2016) Retinoic acid signaling determines the fate of uterine
stroma in the mouse Mullerian duct. *Proc Natl Acad Sci U S A* **113**, 14354-14359
67. Jaslove, J. M., and Nelson, C. M. (2018) Smooth muscle: a stiff sculptor of epithelial shapes.
Philos Trans R Soc Lond B Biol Sci **373**
68. Koyama, H., Shi, D., and Fujimori, T. (2019) Biophysics in oviduct: Planar cell polarity, cilia,
epithelial fold and tube morphogenesis, egg dynamics. *Biophys Physicobiol* **16**, 89-107
69. Ide, S., Finer, G., Maezawa, Y., Onay, T., Souma, T., Scott, R., Ide, K., Akimoto, Y., Li, C., Ye, M.,
Zhao, X., Baba, Y., Minamizuka, T., Jin, J., Takemoto, M., Yokote, K., and Quaggin, S. E. (2018)
Transcription Factor 21 Is Required for Branching Morphogenesis and Regulates the Gdnf-Axis in
Kidney Development. *J Am Soc Nephrol* **29**, 2795-2808

70.   Umezu, T., Yamanouchi, H., Iida, Y., Miura, M., and Tomooka, Y. (2010) Follistatin-like-1, a
diffusible mesenchymal factor determines the fate of epithelium. *Proc Natl Acad Sci U S A* **107**,
4601-4606

71.   Sylva, M., Li, V. S., Buffing, A. A., van Es, J. H., van den Born, M., van der Velden, S., Gunst, Q.,
Koolstra, J. H., Moorman, A. F., Clevers, H., and van den Hoff, M. J. (2011) The BMP antagonist
follistatin-like 1 is required for skeletal and lung organogenesis. *PLoS One* **6**, e22616

72.   Jiang, H., Zhang, L., Liu, X., Sun, W., Kato, K., Chen, C., Li, X., Li, T., Sun, Z., Han, W., Zhang, F.,
Xiao, Q., Yang, Z., Hu, J., Qin, Z., Adams, R. H., Gao, X., and He, Y. (2020) Angiocrine FSTL1
(Follistatin-Like Protein 1) Insufficiency Leads to Atrial and Venous Wall Fibrosis via SMAD3
Activation. *Arterioscler Thromb Vasc Biol* **40**, 958-972

73.   Namwanje, M., and Brown, C. W. (2016) Activins and Inhibins: Roles in Development,
Physiology, and Disease. *Cold Spring Harb Perspect Biol* **8**

74.   Joseph, A., Yao, H., and Hinton, B. T. (2009) Development and morphogenesis of the
Wolffian/epididymal duct, more twists and turns. *Dev Biol* **325**, 6-14

75.   Marečková, M., Garcia-Alonso, L., Moullet, M., Lorenzi, V., Petryszak, R., Sancho-Serra, C.,
Oszlanczi, A., Mazzeo, C. I., Hoffman, S., Krassowski, M., Garbutt, K., Kelava, I., Gaitskell, K.,
Yancheva, S., Woon, E. V., Male, V., Granne, I., Hellner, K., Mahbubani, K. T., Saeb-Parsy, K.,
Lotfollahi, M., Prigmore, E., Southcombe, J., Dragovic, R. A., Becker, C. M., Zondervan, K. T., and
Vento-Tormo, R. (2023) An integrated single-cell reference atlas of the human endometrium.
bioRxiv, 2023.2011.2003.564728

76.   Settle, S., Marker, P., Gurley, K., Sinha, A., Thacker, A., Wang, Y., Higgins, K., Cunha, G., and
Kingsley, D. M. (2001) The BMP family member Gdf7 is required for seminal vesicle growth,
branching morphogenesis, and cytodifferentiation. *Dev Biol* **234**, 138-150

77.   Tomaszewski, J., Joseph, A., Archambeault, D., and Yao, H. H. (2007) Essential roles of inhibin
beta A in mouse epididymal coiling. *Proc Natl Acad Sci U S A* **104**, 11322-11327

78.   Sainio, K., Nonclercq, D., Saarma, M., Palgi, J., Saxen, L., and Sariola, H. (1994) Neuronal
characteristics in embryonic renal stroma. *Int J Dev Biol* **38**, 77-84

79.   McKey, J., Bunce, C., Batchvarov, I. S., Ornitz, D. M., and Capel, B. (2019) Neural crest-derived
neurons invade the ovary but not the testis during mouse gonad development. *Proc Natl Acad*
Sci U S A **116**, 5570-5575

80.   Yan, D., Liu, X., Xu, H., and Guo, S. W. (2020) Mesothelial Cells Participate in Endometriosis
Fibrogenesis Through Platelet-Induced Mesothelial-Mesenchymal Transition. *J Clin Endocrinol*
Metab **105**

81.   Zalecka, J., Pankiewicz, K., Issat, T., and Laudanski, P. (2022) Molecular Mechanisms Underlying
the Association between Endometriosis and Ectopic Pregnancy. *Int J Mol Sci* **23**

82.   Santana Gonzalez, L., Rota, I. A., Artibani, M., Morotti, M., Hu, Z., Wietek, N., Alsaadi, A.,
Albukhari, A., Sauka-Spengler, T., and Ahmed, A. A. (2021) Mechanistic Drivers of Mullerian Duct
Development and Differentiation Into the Oviduct. *Front Cell Dev Biol* **9**, 605301

83.   Lengyel, E., Li, Y., Weigert, M., Zhu, L., Eckart, H., Javellana, M., Ackroyd, S., Xiao, J., Olalekan, S.,
Glass, D., Iyer, S., Krishnan, R., Bilecz, A. J., Lastra, R., Chen, M., and Basu, A. (2022) A molecular
atlas of the human postmenopausal fallopian tube and ovary from single-cell RNA and ATAC
sequencing. *Cell Rep* **41**, 111838

84.   Ulrich, N. D., Shen, Y. C., Ma, Q., Yang, K., Hannum, D. F., Jones, A., Machlin, J., Randolph, J. F.,
Jr., Smith, Y. R., Schon, S. B., Shikanov, A., Marsh, E. E., Lieberman, R., Gurczynski, S. J., Moore, B.
B., Li, J. Z., and Hammoud, S. (2022) Cellular heterogeneity of human fallopian tubes in normal
and hydrosalpinx disease states identified using scRNA-seq. *Dev Cell* **57**, 914-929 e917

85.   Dinh, H. Q., Lin, X., Abbasi, F., Nameki, R., Haro, M., Olingy, C. E., Chang, H., Hernandez, L.,
Gayther, S. A., Wright, K. N., Aspuria, P. J., Karlan, B. Y., Corona, R. I., Li, A., Rimel, B. J., Siedhoff,

M. T., Medeiros, F., and Lawrenson, K. (2021) Single-cell transcriptomics identifies gene
expression networks driving differentiation and tumorigenesis in the human fallopian tube. *Cell*
Rep **35**, 108978
86.	Qin, G., Park, E. S., Chen, X., Han, S., Xiang, D., Ren, F., Liu, G., Chen, H., Yuan, G. C., and Li, Z.
(2023) Distinct niche structures and intrinsic programs of fallopian tube and ovarian surface
epithelial cells. *iScience* **26**, 105861
87.	Harwalkar, K., Ford, M. J., Teng, K., Yamanaka, N., Yang, B., Burtscher, I., Lickert, H., and
Yamanaka, Y. (2021) Anatomical and cellular heterogeneity in the mouse oviduct-its potential
roles in reproduction and preimplantation developmentdagger. *Biol Reprod* **104**, 1249-1261
88.	Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W. M., 3rd, Zheng, S., Butler, A., Lee, M. J., Wilk, A.
J., Darby, C., Zager, M., Hoffman, P., Stoeckius, M., Papalexi, E., Mimitou, E. P., Jain, J.,
Srivastava, A., Stuart, T., Fleming, L. M., Yeung, B., Rogers, A. J., McElrath, J. M., Blish, C. A.,
Gottardo, R., Smibert, P., and Satija, R. (2021) Integrated analysis of multimodal single-cell data.
Cell **184**, 3573-3587 e3529
89.	Heumos, L., Schaar, A. C., Lance, C., Litinetskaya, A., Drost, F., Zappia, L., Lucken, M. D., Strobl, D.
C., Henao, J., Curion, F., Single-cell Best Practices, C., Schiller, H. B., and Theis, F. J. (2023) Best
practices for single-cell analysis across modalities. *Nat Rev Genet* **24**, 550-572
90.	Khozyainova, A. A., Valyaeva, A. A., Arbatsky, M. S., Isaev, S. V., Iamshchikov, P. S., Volchkov, E.
V., Sabirov, M. S., Zainullina, V. R., Chechekhin, V. I., Vorobev, R. S., Menyailo, M. E., Tyurin-
Kuzmin, P. A., and Denisov, E. V. (2023) Complex Analysis of Single-Cell RNA Sequencing Data.
Biochemistry (Mosc) **88**, 231-252

Figure Legends

Figure 1. Single-cell transcriptional atlas of PND3 oviduct and uterus. (A) Tissue sample surveyed in this study. The dotted line indicated where the oviduct and the uterus were separated. Scale bar=2.5 mm. (B) Uniform manifold approximation and projection (UMAP) plot featuring general cell clusters from both the oviduct and uterus. In the lower right, the same UMAP was annotated according to cell's organ origin. (C) UMAPs featuring cell type markers, *Hoxc8* for oviductal mesenchyme, *Hoxa10* for uterine mesenchyme, *Top2a* for proliferating cells, *Epcam* for epithelium, *Msln* for mesothelium, *Pecam1* for endothelium, *Rgs5* for pericytes, and *Lyz2* for myeloid cells. (C) Heatmap of top five marker genes for each cluster.

Figure 2. Characterization of mesenchymal subpopulations in PND3 oviduct and uterus. (A) UMAP featuring the mesenchymal subpopulation after re-clustering with higher resolution. In the lower right, the same UMAP was annotated according to cell's organ origin. (B) UMAPs feature marker genes for cell subtypes: *Cnn1* for smooth muscle cells in both the oviduct and uterus, *Tcf21* for uterine smooth muscle, *Nrp2* for oviductal smooth muscle, *Pkib* & *Pcdh10* for two uterine mesenchymal subtypes, *Wfdc1* & *Angptl7* for two oviductal mesenchymal subtypes, and *Nefm* for a mesenchymal subtype found in both the oviduct and uterus. (C) mRNA expression of four markers for mesenchymal subpopulations, *Wfdc1*, *Angptl7*, *Nefm* and *Tcf21,* by RNAscope. N=3. Scale bar=50 μm.

Figure 3. Region-specific epithelial subtypes in PND3 oviduct and uterus. (A)

UMAP featuring oviductal and uterine epithelial cells extracted from the overall dataset

in Fig 1B. (B) UMAPs featuring marker genes *Msx1* and *Lrpap1* for uterine and

oviductal epithelium, respectively. (C) UMAP featuring oviductal epithelial cells

extracted from Fig 3A and then re-clustered. (D) Violin plot showing *Gsap* expression in

two subclusters in Fig 3C. (E) Immunofluorescent staining of MSX1 and RNAscope

staining of *Gsap* in the cranial oviduct, caudal oviduct and uterus at PND3. N=3. Scale

bar: 25 and 50 µm for immunofluorescence and RNAscope, respectively.

Figure 4. Mesenchymal-epithelial crosstalk represented by region-specific ligands

that are potentially functional. Only region-specific ligands whose knockout mice

display embryonic lethality and/or reproductive phenotype were included. Please see all

ligand-pair pairs in the Fig S4 and supplementary files 7-12.

Figure 5. Gene expression comparison between oviductal and uterine pericytes.

(A) Violin plots and UMAPs of higher expressed genes (*Cebpb*, *Rgs16*, *Nr2f1*, *Hoxc8*

and *Cryab*) in oviductal pericytes. (B) Violin plots and UMAPs of higher expressed

genes (*Amer1*, *Plac8*, *Hoxd11*, *Lsp1*, and *Hoxa10*) in uterine pericytes

Figure 6. Gene expression comparison between oviductal and uterine

mesothelium. (A) UMAPs featuring expression of epithelium-specific (*Epcam*, *Mecom*),

mesothelium-specific (*Upk3b*), and commonly shared (*Krt8*, *Krt18* and *Krt19*) marker

genes. (B) Violin plots of higher expressed genes in the oviductal mesothelium (up row:

Lgals7, *Igf1*, *Mt1*, *Ccl2*, and *Nrgn)* and uterine mesothelium (bottom row: *Alcam*, *Rspo3*,

Fxyd3, *Tcf21*, and *Hoxd11)*.

Figure 7. Graphical summary

Figure S1. Composition of mesenchymal subpopulations in PND3 oviduct and

uterus. (A) Pie chart representing mesenchymal compositions and percentages of each

mesenchymal subpopulation. (B) Heatmap of top five marker genes for each cluster.

The color bar for each cluster corresponds to the colors in the pie chart of Fig S1A.

Figure S2 Expression of representative genes in the UMAP of the mesenchymal

population.

Figure S3. Expression of multiciliogenesis related genes in the UMAP of oviductal

epithelia at PND3.

Figure S4. Identification of ligand-receptor pairs mediating mesenchyme-

epithelium interactions in the cranial oviduct, the caudal oviduct, and the uterus

at PND3. First, the number of ligand-receptor pairs mediating the crosstalk between

region-specific epithelium and organ-specific mesenchymal subpopulations was shown.

Then, venn diagrams showed the overlap between the lists of ligand-receptor pairs sent

from two mesenchymal subpopulations to epithelium or sent from the epithelium to the

two mesenchymal subpopulations. All the ligands were consolidated and selected for

generating Fig 4 based on the following criteria: the knockout mice had reproductive

phenotypes and/or embryonic lethality based on Mouse Genome Informatics (MGI), and

ligands are enriched only in either the oviduct or uterus. Please see all ligand-pair pairs

in the supplementary files 7-12.

A

Uterus mesenchymal cells (*Pcdh10*+)

Uterus mesenchymal cells (*Pkib*+)

Shared cluster (*Nefm*+)

Oviduct mesenchymal cells (*Wfdc1*+)

Oviduct smooth muscle cells (*Nrp2*+)

Uterus smooth muscle cells (*Tcf21*+)

Oviduct mesenchymal cells (*Angptl7*+)

UMAP2

UMAP1

oviduct
uterus

B

Cnn1

Tcf21

Nrp2

Pkib

Pcdh10

Angptl7

Wfdc1

Nefm

C

	Enriched in oviductal mesenchyme		Found in both the uterus and oviduct	Enriched in uterine mesenchyme
	Wfdc1	*Angptl7*	*Nefm*	*Tcf21*
Cranial Oviduct				
Caudal Oviduct				
Uterus				

50 μm

A Higher expressed genes in oviductal pericytes

B Higher expressed genes in uterine pericytes

A

Epithelium-specific

Mesothelium-specific

Expressed in both epithelium and mesothelium

B

Fig. 7 Graphical summary

Cranial

Cranial oviduct

Caudal oviduct

Uterus

Caudal

Anti-mesometrial ← → Mesometrial

Cranial oviduct epithelial cells (*Gsap+*)

Caudal oviduct epithelial cells (*Gsap-*)

Uterine epithelial cells (*Msx1+*)

Mesothelial cells

Oviductal mesenchymal cells (*Wfdc1+*, *Angptl7+*)

Uterine mesenchymal cells (*Pcdh10+*, *Pkib+*)

Nefm+ cells

Oviductal smooth muscle cells (*Nrp2+*)

Uterine smooth muscle cells (*Tcf21+*)