RESEARCH PAPER

The radiation of nodulated *Chamaecrista* species from the rainforest

into more diverse habitats has been accompanied by a reduction in

growth form and a shift from fixation threads to symbiosomes

Patricia Alves Casaes[1] | José Miguel Ferreira dos Santos[2] | Verônica Cordeiro Silva[1] |

Mariana Ferreira Kruschewsky Rhem[1] | Matheus Martins Teixeira Cota[3] | Sergio Miana de

Faria[4] | Juliana Gastaldello Rando[5] | Euan K. James[6] | Eduardo Gross[7]

[1]Programa de Pós-Graduação em Biologia e Biotecnologia de Microrganismos, Universidade

Estadual de Santa Cruz, Bahia, Brazil

[2]Faculdade Pitágoras de Medicina e Faculdade Espírito Santo, Bahia, Brazil

[3]Programa de Pós-Graduação em Botânica, Universidade Estadual de Feira de Santana, Bahia,

Brazil

[4]Embrapa-Agrobiologia, 465 km 07, Seropédica, Rio de Janeiro, Brazil

[5]Programa de Pós-Graduação em Ciências Ambientais, Universidade Federal do Oeste da

Bahia, Bahia, Brazil

[6]The James Hutton Institute, Dundee, United Kingdom.

[7]Departamento de Ciências Agrárias e Ambientais, Universidade Estadual de Santa Cruz, Bahia,

Brazil

Correspondence

Eduardo Gross

Departamento de Ciências Agrárias e Ambientais

UESC – Universidade Estadual de Santa Cruz

Rod. Jorge Amado, Km 16

Ilhéus-BA, 45662-900, Brazil

E-mail: egross@uesc.br

Euan K. James

The James Hutton Institute,

Invergowrie,

Dundee DD2 5DA,

UK.

E-mail: euan.james@hutton.ac.uk

Abstract

All non-mimosoid nodulated genera in the legume subfamily Caesalpinioideae confine their rhizobial symbionts within cell wall-bound "fixation threads" (FTs). The exception is the large genus *Chamaecrista* in which shrubs and subshrubs house their rhizobial bacteroids more intimately within symbiosomes, whereas large trees have FTs. This study aimed to unravel the evolutionary relationships between *Chamaecrista* growth habit, habitat, nodule bacteroid type, and rhizobial genotype. The growth habit, bacteroid anatomy, and rhizobial symbionts of 30 nodulated *Chamaecrista* species native to different biomes in the Brazilian state of Bahia, a major centre of diversity for the genus, was plotted onto an ITS-*TrnL-F*-derived phylogeny of *Chamaecrista*. The bacteroids from most of the *Chamaecrista* species examined were enclosed in symbiosomes (SYM-type nodules), but those in arborescent species in the section *Apoucouita*, at the base of the genus, were enclosed in cell wall material containing homogalacturonan (HG) and cellulose (FT-type nodules). Most symbionts were *Bradyrhizobium* genotypes grouped according to the growth habits of their hosts, but the tree, *C. eitenorum*, was nodulated by *Paraburkholderia*. *Chamaecrista* has a range of growth habits that allow it to occupy several different biomes and to co-evolve with a wide range of (mainly) bradyrhizobial symbionts. FTs represent a less intimate symbiosis linked with nodulation losses, so the evolution of SYM-type nodules by most *Chamaecrista* species may have (a) aided the genus-wide retention of nodulation, and (b) assisted in its rapid speciation and radiation out of the rainforest into more diverse and challenging habitats.

Keywords: *Chamaecrista*, nodulation, homogalacturonan (HG), cellulose, fixation thread (FT), Caesalpinioideae, *Bradyrhizobium, Paraburkholderia, nodC, nifH.*

Introduction

The monophyletic legume genus *Chamaecrista* (L.) Moench (Leguminosae - Caesalpinioideae) has its centre of diversification in South America (Conceição et al., 2009). Most species occur in Brazil, where 268 of approximately 366 recognized species are distributed in various biomes and vegetation types. About 223 species are considered as Brazilian endemics (LPWG 2020; Rando et al., 2020) particularly in the states of Bahia (BA) and Minas Gerais (MG) which harbour 94 species distributed in such diverse environments as savannah (Cerrado), Campo rupestre (upland rocky fields), semiarid ecosystems (Caatinga), and tropical rain forest (Coutinho et al., 2016; Rando et al., 2016, 2020).

Chamaecrista is the ninth largest genus in the Leguminosae (Fabaceae) and the third largest in the Caesalpinioideae subfamily (after *Acacia* and *Mimosa*); it contains a wide variety of plant growth habits and sizes, ranging from trees through to shrubs, and subshrubs/woody herbaceous perennials (Lewis, 2005; Coutinho et al., 2016; Mendes et al. 2017; LPWG 2021). Furthermore, the distribution of *Chamaecrista* is unique in being the only nodulated caesalpinioid genus which has species that have colonized temperate regions (Sprent et al., 2013). All *Chamaecrista* species so far examined form symbiotic root nodules with nitrogen-fixing bacteria collectively known as rhizobia (Gyaneshwar et al., 2011; Peix et al., 2015; Sprent et al., 2017), while related genera comprising the Cassia clade (LPWG 2017), p. ex. *Cassia* L. and *Senna* Mill. do not nodulate (Sprent, 2001, 2009). The independent rise of nodulation and its variation in *Chamaecrista* (Delaux et al., 2015; Naisbitt et al., 1992) suggests that the genus has a pivotal position in the evolution of nodulation (Sprent et al., 2013), and could be used as a model for detailed studies of interactions between plants and nitrogen-fixing bacteria (Singer et al., 2009). Indeed, it is for this reason that Sprent et al. (2013) suggested studying the nitrogen-fixing nodules across *Chamaecrista* species in more depth with a focus on their structure and rhizobial symbionts.

There is a paucity of information about nodule anatomy and development in the paraphyletic grade comprising the Caesalpinioideae (*i.e.,* excluding the Mimosoid clade), which includes *Chamaecrista*. What we do know is that all nodules so far studied from the nine known nodulating non-Mimosoid Caesalpinioideae genera (*Campsiandra*, *Chamaecrista*, *Dimorphandra*, *Dinizia*, *Erythrophleum*, *Jacqueshuberia*, *Melanoxylon*, *Moldenhawera* and *Tachigali*) are indeterminate, retaining meristematic activity (Sprent, 2009; Fonseca et al., 2012; Sprent et al., 2013, 2017; Faria et al. 2022). Moreover, while the nodules in most papilionoid and all mimosoid species so far examined, have their

symbiotic rhizobia (bacteroids) released into membrane-bound vesicles called
symbiosomes (Sprent, 2009; Sprent et al., 2013, 2017), some papilionoid, and all
nodulated (non-mimosoid) caesalpinioid trees so far studied have their bacteroids
confined within cell wall-bound modified infection threads termed ''persistent infection
threads'' (PITs) or ''fixation threads'' (FTs) (Faria et al., 1987, 2022; Naisbitt et al., 1992;
Sprent, 2009; Fonseca et al., 2012; Sprent et al., 2013, 2017). In *Chamaecrista*, however,
nodules are anatomically diverse with their ultrastructure being apparently related to plant
growth habit i.e. the rhizobial bacteroids are enclosed in FTs in tree species, but within
membrane-bound symbiosomes in subshrub/woody herbaceous species (henceforth
collectively termed "subshrubs"), while larger shrub/treelet species have nodules with
intermediate structures between FTs and symbiosomes (Naisbitt et al., 1992).

The nodule anatomy of most species of *Chamaecrista* is so far undescribed, as
is the nodulation status of the majority of the genus. In addition, there have been
significant advances in microscopy techniques since Naisbitt et al., (1992), particularly in
methods to determine the composition of membranes and cell walls that could be
associated with the FTs and symbiosomes (Faria et al. 2022). Therefore, the first aim of
this study was to record the nodulation status of a wide variety of *Chamaecrista* species
covering their whole range of growth habits, from 20 m-high tree species to small
subshrubs at only 20-30 centimeters in height. Plant size variation in *Chamaecrista* is
also related to the biomes within which they occur: large trees are distributed in tropical
rainforests where soils are nitrogen (N) rich, while shrubs and subshrubs are distributed
in tropical savannas and in xeric formations with N-poor soils. Moreover, the phylogenetic
relations within the genus *Chamaecrista* also shows that monophyletic groups share
similar types of habitats and growth habit. The few arborescent species in the genus, all
native to tropical rainforests, are grouped in the basal monophyletic section *Apoucouita*
(Coutinho et al., 2016; Souza et al. 2021), while the considerably more numerous species
from open areas are shrubs and subshrubs that are scattered widely across the taxonomy
of the genus (Conceição et al., 2009, Souza et al., 2021). A second aim was to evaluate
the anatomy and ultrastructure of root nodules from *Chamaecrista* species of all growth
habits, focusing on the chemical composition of structures (symbiosomes, FTs, and
intermediates) enclosing the bacteroids within the infected tissue of the nodules.

The third aim was to explore the identity of the rhizobial symbionts of a wide
variety of *Chamaecrista* species covering the whole range of growth habits from trees
through to shrubs and subshrubs, and from nodules with FTs to those with symbiosomes.
Current data suggest that non-mimosoid Caesalpinioideae are preferentially nodulated

by *Bradyrhizobium* strains (Fonseca et al., 2012; Yao et al., 2014, 2015; Parker 2015;
Sprent et al., 2017; Rathi et al., 2018; Cabral Michel et al., 2021). Indeed, in the specific
case of the largest nodulating genus in this group, *Chamaecrista*, several studies have
shown that the North American subshrub *C. fasciculata* (Michx.) Greene (partridge pea)
has nodules that are associated with *Bradyrhizobium* (Parker 2012, Parker & Rousteau
2014; Urquiaga et al., 2019; Klepa et al., 2019), and a recent molecular analysis of 47
strains from nine shrub and subshrub *Chamaecrista* species in Brazil described all the
symbiotic strains as belonging to genotypes of *Bradyrhizobium* (Santos et al., 2017).
Outside the New World, reports on native shrubby *Chamaecrista* species in India and
Africa suggest that they are also mainly nodulated by bradyrhizobia (de Lajudie et al.,
1998; Beukes et al., 2016; Rathi et al., 2018). Although evidence to date suggests that a
strong association between *Chamaecrista* and bradyrhizobia is consistent, nothing is
known about the diversity of rhizobial symbionts in nodules of tree species which have
their bacteroids enclosed in FTs as opposed to those with symbiosomes, nor whether
there is a link between rhizobial genotypes and the different biomes within which their
hosts occur.
Using these data plus unpublished anatomical data obtained from *Chamaecrista*
nodules sampled during the expedition of Sprent et al. (1996), combined with data from
the literature (Naisbitt et al. 1992; Santos et al. 2017; Faria et al. 2022), we then test the
hypothesis that the distribution of FTs and symbiosomes in the genus is not random and
may be the result of co-evolution between groups (sections) of *Chamaecrista* that are
native to particular environments (biomes), and the rhizobial microsymbionts that live
within them. This was done by constructing a phylogeny with ITS-*TrnL-F* sequences from
119 separate *Chamaecrista* taxa, and then plotting onto it plant growth habit, nodule
ultrastructure (occurrence of FTs or symbiosomes, if known), and the *nodC* genotypes of
the rhizobial microsymbionts.
160
Materials and methods
162
Botanical material
164
To sample root nodules in the field, we first located the 17 *Chamaecrista* species studied
here in different biomes of Bahia State, Brazil (Fig. S1). Samples of the root system
bearing nodules were collected from each individual plant, mainly during the rainy
season, when nodules are most active (dos Reis Junior et al., 2010). Aerial parts of each

species were collected, dried, and deposited in the Herbarium of the Universidade
Estadual de Santa Cruz (UESC) (Table S1) for confirmation of their identities.

On average, 12 nodules were collected from each of the 17 *Chamaecrista* species,
totaling 204 nodules. These samples were used to characterize the infected tissues and
to verify the presence of FTs using scanning electron microscopy (SEM), transmission
electron microscopy (TEM), light and fluorescence microscopy (see details below).

Anatomy, histochemistry, ultrastructure and immunocytochemistry of nodules

The nodules were separated from the root system and cut into 1-2 mm^3 pieces with a
razor blade, before being fixed in a solution of 2.5% glutaraldehyde in 0.1 M sodium
cacodylate buffer (pH 7.2). Sample processing followed Santos et al. (2017) for
anatomical and ultrastructural characterization.

Histochemistry using the fluorescent compound calcofluor white, which binds to
beta 1-3 and beta 1-4 polysaccharides, such as those found in cellulose (Wood et al.,
1983), was used to provide evidence for the presence of invasive infection threads (ITs)
and FTs within the infected tissue of *Chamaecrista* nodules. Briefly, sections of nodules
were incubated with calcofluor white (1 g L^{-1}) in Evans blue as a background stain (0.5 g
L^{-1}) (Sigma-Aldrich) according to Wood et al. (1983), and the sections were then observed
under a Leica DM2500 equipped with ebq100-04 fluorescence coupled to a Leica
DFC310 Fx digital camera.

For transmission electron microscopy (TEM), serial ultrathin sections were collected
on nickel grids for immunogold tests using the monoclonal antibody JIM5 to verify the
presence of a homogalacturonan (HG) epitope in FTs inside the nodule infected cells;
this HG epitope is an essential component of pectin in cell walls (VandenBosch et al.,
1989; Fonseca et al., 2012). For species from which *Paraburkholderia* were isolated as
potential symbionts, nodules were also tested for the *in-situ* presence of symbiotic strains
using a polyclonal antibody against *P. phymatum* STM815^T according to dos Reis Junior
et al. (2010).

For scanning electron microscopy (SEM), fixed samples were dehydrated in a
graded acetone series to absolute, and completely dried using a Bal-Tec CPD 030 critical
point drier. Dried samples were mounted on stubs, coated with gold in a Bal-Tec SCD
050 sputter coater and viewed with a FEI Quanta 250 at the Centro de Microscopia
Eletrônica (CME) at UESC.

Rhizobia isolation, cultivation and characterization

Potential rhizobia were isolated from root nodules of seven of the 17 *Chamaecrista*
species sampled in the field from native undisturbed environments. An intensive effort
was made to isolate rhizobia from tree species of *Chamaecrista* because they often have
woody and lignified nodules, and hence the symbiotic bacteria are more difficult to isolate.
Accordingly, approximately 230 nodules were used for rhizobial isolation from *C.*
ensiformis var. *plurifoliolata*, *C. duartei*, *C. eitenorum* and *C. bahiae*. Rhizobia were
isolated, cultivated and characterized following the procedures of Rhem et al. (2021).

Bacterial DNA extraction, amplification, sequencing and phylogenetic analysis

For samples from each plant, bacterial isolates were grouped and selected according to
similarities in their phenotypical (colony) characteristics. Genomic DNA from selected
isolates was extracted according to Santos et al. (2017). The DNA was resuspended in
ultrapure water and stored at -20 °C. The yield and purity of the extracted DNA was
measured in a spectrophotometer (Shimadzu, SPD-M6A) by the ratio of absorbance at
260 and 280 nm. For *Bradyrhizobium* strains, the 16S rRNA gene and ITS region were
amplified. A multilocus sequence analysis (MLSA) was performed following Rhem et al.
(2021) to identify the phylogenetic positions of symbiont strains from different
Chamaecrista species. For MLSA, DNA sequences of four housekeeping genes were
used, i.e., *recA* encoding recombinase A, *dnaK* encoding the Hsp70 chaperone, *rpoB*
encoding RNA polymerase beta subunit, and *glnII* encoding glutamine synthetase isoform
II. We also amplified two symbiotic genes, i.e. approximately 930 bp of the *nodC*
(nodulation N-acetylglucosaminyltransferase) and 780 bp of the *nifH* (nitrogenase
reductase) genes. Thermal cycler programs and primers used are described in Rhem et
al. (2021). For *Paraburkholderia* strains, the 16S rRNA, *recA*, *nifH* and *nodC* genes were
amplified with the same primers and PCR conditions cited by Silva et al. (2018). Amplified
DNA was verified by horizontal electrophoresis in 1% (w/v) agarose gels and PCR
products were purified following a cold salt precipitation and resuspended in sterile
ultrapure water. All amplicons obtained were sequenced by ACTGene Análises
Moleculares Ltda (Alvorada, RS, Brazil). Nucleotide sequences of strains were analyzed
for percentage sequence similarity using BLASTn of the National Center for
Biotechnology Information (NCBI). Sequences from the present study and those of
closely related, reference and type strains (as per the LPSN list of valid and not validly

published type strains) were downloaded from NCBI in FASTA format, and then aligned
using CLUSTAL X (Thompson et al. 1997) of MEGA X (Kumar et al. 2018). Phylogenetic
trees were constructed with the sequence alignments of all the tested genes. Maximum-
likelihood trees were built in MEGA 7 using the Tamura 3-parameter correction method.
The robustness of the branches of the trees was estimated with 1000 bootstrap
replications. The partial sequences of all genes derived from the present study were
deposited in GenBank, and their Accession Numbers are listed in Table S2. Please note
that it was not possible to amplify all of the examined gene loci from all of the strains.
247
248 *Plant DNA extraction, amplification, sequencing and phylogenetic analysis*
249
DNA was sampled from 106 species of *Chamaecrista* (119 taxa) covering all sections
and including all species with root nodules analyzed in the present study plus others
accessed from the literature and from the dataset associated with Sprent et al. (1996),
with only *C. zygophylloides* (Taub.) H.S. Irwin & Barneby not included (Table S3).
Additionally, four species of *Cassia* and *Senna* were included as an outgroup. Most of the
sequence data were accessed from GenBank (http://www.ncbi.nlm.nih.gov/genbank/),
but seven new sequences of the plastid *trnL-F* and four of the nuclear internal transcribed
spacer (ITS) were generated for a further seven *Chamaecrista* species. All DNA
sequences and associated voucher information are deposited in GenBank (Table S3).
Total DNA extraction, amplification, PCR product purification, and sequencing were as
described in Conceição et al. (2009). Electropherograms were assembled and edited
using the Geneious platform (Drummond et al., 2012). Alignments of all sequences were
performed using Muscle (Edgar, 2004) with default settings. Manual edition to correct
obvious alignment errors and to remove sections with dubious alignments were inspected
using the Geneious platform (Drummond et al., 2012). Bayesian analyses (BAs) were
performed with MrBayes 3.1 (Ronquist & Huelsenbeck, 2003) using a combined data set
with four partitions (*trnL-F*, ITS1, 5.8S and ITS2). Nucleotide-substitution models were
selected, on the basis of the Akaike information criterion (AIC) values, with JModeltest
2.1 (Guindon & Gascuel, 2003; Darriba et al., 2012). The substitution models selected for
trnL-F was GTR+G and for ITS1 was GTR+I+G, for 5.8S was SYM+I and for ITS2 was
GTR+G. Indels were coded as the standard characters "variable". We run all phylogenetic
analyses via the CIPRES Science Gateway v. 3.3 online portal (Miller et al., 2010). We
used FigTree 1.4.2 (Rambaut, 2014) to view and edit the final tree.
273

Nodulation tests with rhizobial strains

We selected strains isolated from *Chamaecrista* species to evaluate their nodulation

capacity. Six *Chamaecrista* species, ranging in size/habit from trees (*C. bahiae, C.*

ensiformis var. *plurifoliolata* and *C. duartei*), treelets (*C. blanchetii*) to subshrubs (*C.*

desvauxi, C. rotundifolia), as well as the promiscuous papilionoid legume Siratro

(*Macroptilium atropurpureum*) were used as host plants in the nodulation tests. In

addition, *Mimosa pudica* was used as a promiscuous mimosoid legume to test the two

Paraburkholderia isolates. Siratro seedlings were inoculated with the different strains

according to Santos et al. (2017), while the *Chamaecrista* spp. and *M. pudica* were

inoculated according to Silva et al. (2018). At harvest (3 months after inoculation),

nodulation of the root system was evaluated for each plant, and the presence of the

symbiosis-essential protein leghemoglobin (Lb), as indicated by a pinkish coloration in

their interior, was scored.

Results

Anatomy, ultrastructure and immunocytochemistry of nodules

Chamaecrista species have a wide range of growth habits and habitats, but most are

shrubs and subshrubs. The few tree species are usually found in rainforests, such as the

Mata Atlântica (Fig. 1A) and Amazon Forest; an exception is *C. eitenorum* which has its

habitat in seasonally dry tropical forests (SDTF) in the Chapada Diamantina. Most of the

tree species have characteristic ramiflorous racemes (Fig. 1B) and are placed in the small

section *Apoucouita* at the base of the genus (Coutinho et al., 2016). Woody shrub species

occur in the *Cerrado* (savannah) and in open rocky fields often at altitudes >1000 m

(denoted *Campo Rupestre*) (Fig. 1C, D); these belong in sect. *Absus*, members of which

have terminal or axillary racemes. The *Cerrado* and *Caatinga* (a semiarid Brazilian biome)

are also the habitat for smaller, but still woody, sub-shrub and herbaceous species (Fig.

1E, F) which have axillary and supra-axillary reduced racemes (fascicles); these are

mostly contained in sect. *Chamaecrista*.

A complete list of nodulating *Chamaecrista* species is given in Table S4. Nineteen

Chamaecrista taxa were examined in the present study for nodulation, plus twelve from

the study of Sprent et al. (1996), and a further eight from de Faria (unpublished). All were

nodulated, including 26 new reports.

The nodules sampled in the present study (varying from 0.1 to 2 cm in length) were
usually found on the secondary superficial roots (Fig 2A) of all the *Chamaecrista* species
examined. In general, nodules were cylindrical when young, but became lobed with age
with a few branches, and were a dark brown surface color (Fig. 2B); most were viable
and active as evidenced by the pink color in their interior due to Lb production (inset Fig.
2B). Nodule morphology and anatomy characterized all of the samples as indeterminate
as they have a persistent meristem at the distal end (Fig. 2C). *Chamaecrista* nodules
have a central infected tissue containing the microsymbionts which is surrounded by an
uninfected cortex consisting of a parenchymatous outer cortex composed of several
layers (4 to 6) of isodiametric cells with phenolic compounds being found scattered
throughout the outer cortical region; this is separated from the inner cortex by an
endodermis (Fig. 2C). Several vascular bundles are located at the periphery of the inner
cortex (Fig. 2C). In all *Chamaecrista* species, the central region of nodules contains a
combination of large infected parenchymatic cells and smaller uninfected interstitial cells
(Fig. 2C). In the invasion zone, which was located distally to the infected tissue, the
rhizobia were contained within ITs that invaded cells (Fig. 2D). Depending on the growth
habit of the species, these ITs either released the bacteria into symbiosomes (Fig. 2G,
S2), or the ITs developed into thinner-walled FTs which occupied almost the entire
cytoplasmic volume of the infected cells (Fig. 2D, S2). In mature nodules, senescent cells
were observed in the proximal part of the infected tissue, with a collapsing mass of FTs
in their interior (Fig. 2E). The small uninfected interstitial cells were vacuolated, and
distinct from the infected cells (Fig. 2F, G, S2). In subshrub *Chamaecrista* species, the
root nodules had infected cells with symbiosomes completely occupying the available
cytoplasm (Fig. 2G, S2I, K).

The occurrence of nodules with conspicuous FTs was confined to tree
Chamaecrista species in the section *Apoucouita* (Table 1, S4, Figs. 3, 4, S2), which are
generally restricted to tropical forests (Coutinho et al. 2016; Souza et al. 2021). The
composition of the FT cell walls was examined using a combination of three methods: (1)
histochemical staining of cellulose using Calcofluor white, (2) anatomical observations
with light microscopy and electron microscopy (TEM and SEM), and (3) immunogold
labelling of pectin HG using JIM5 combined with TEM.

Analysis of cellulose deposition in nodule infected cells using calcofluor white
showed a similar pattern to the pectin observations *i.e.* with primary cell wall components
evident in nodules from trees, treelets and large shrubs, but not in mature infected cells
in nodules of subshrubs (Table 2). For example, cellulosic material was detected in FTs

in nodules of *C. bahiae* (Fig. 3A), *C. duartei* (Fig. 3B), *C. ensiformis* var. *plurifoliolata*, *C. brachystachya, C. confertifomis* (Fig. 3C) and *C.* x *blanchetiformis* (Fig. 3D). Fluorescence indicating cellulose was also observed in the infected zone of a *C. arrojadoana* nodule (Fig. 3E) where it was localized to an IT in cells recently invaded by rhizobia and, more generally, in immature invaded cells, but not in mature infected cells of this subshrub species (Fig. 3F).

Under the SEM, conspicuous FTs were observed in nodules of the trees *C. bahiae* (Fig. 4A), *C. duartei*, *C. eitenorum* and *C. ensiformis* var. *plurifoliolata*, and in the shrub species *C. blanchetiformis*, *C. brachystachya* and *C. confertiformis*, all of which had FTs with well-defined walls like those of the "standard" ITs of many legume types. The mature infected cells in the infected zone of the nodules were densely occupied by FTs, and these differed in structure from those observed in nodules of the subshrub species *C. repens* (Fig. 4B), *C. arrojadoana* and *C. ramosa*, wherein bacteroids were enclosed in FTs with thinner cell walls (so-called FT-SYM; Table S4) which allowed for the bacterial outline with associated material on the bacterial surface to be observed under the SEM (inset Fig. 4B). For nodules on the subshrub species, *C. belemi, C. blanchetii, C. zygophylloides, C. desvauxii, C. pascuorum, C. rotundifolia* (Fig. 4C, Table S4) and *C. supplex*, the bacteroids were free in the cells, although they are most likely to be enclosed within symbiosome membranes, most of which did not survive processing for SEM. The SEM observations were confirmed using light microscopy and TEM *i.e.* that tree species in the section *Apoucouita*, such as *C. bahiae*, *C. duartei* and *C. ensiformis* have FTs (Fig. 4D, S2A - D), that symbiosomes in treelets from the section *Baseophyllum* have an intermediate (FT-SYM) structure and/or contain both FTs and symbiosomes (Fig. S2E – H), while shrubs and subshrubs from the sections *Absus* and *Chamaecrista* (Fig. 4E, F, S2I – L) generally contain symbiosomes only. However, there are several exceptions in sections *Absus* and *Chamaecrista* that have an FT-SYM type, such as C. *arrojadoana* (Fig. 4F), *C. mucronata* (Fig. S2L), and *C. desvauxii* (Naisbitt et al. 1992).

Immunocytochemistry using the monoclonal antibody JIM5 provided further information about the nature of FTs and symbiosomes in *Chamaecrista* nodules. It indicated the presence of an unesterified HG epitope in the pectic component of both FTs (Fig. 4D, S2B, D) and symbiosomes (Fig. 4E, S2H, L) in infected cells of nodules from several species of *Chamaecrista* examined (Table 2; Fig. S2). For nodules on tree species in the section *Apoucouita*, such as *C. bahiae*, *C. duartei* and *C. ensiformis*, JIM5 immunogold labeled the walled FTs (Fig. 4D, S2B, D). JIM5 also labelled the thinner walled FTs of treelets and large shrubs in the section *Baseophyllum*, such *as C. blanchetii*

(Fig. S2F), but in nodules on other *Baseophyllum* species, such as *C. cytisoides*, the
infected cells contained a combination of JIM5-labelled symbiosomes and FTs, with the
FTs being more densely labeled (Fig. S2H). Similarly, JIM5 labeling was observed in
nodules of some subshrub species in the highly speciose sections *Absus* and
Chamaecrista, including those that had their bacteroids enclosed in either symbiosomes
(Fig. 4E, S2J) or were intermediate (FT-SYM), such as *C. arrojadoana* and *C. mucronata*
(Fig. 4F, S2L). It should be noted, however, that this was not a uniform observation across
these *Chamaecrista* sections, as nodules of many species had no JIM5 labelling, such
as *C. chapadae* (Fig. S2J), and of those that did have it the JIM5 epitope was located
within the symbiosomes themselves (Fig. 4E). Image analysis of the infected N-fixing
cells in sections from nodules on the species shown in Fig. S2 indicated that the FT-type
nodules from section *Apoucouita* were less densely colonized by bacteroids than those
of the SYM-type nodules in sections *Absus* and *Chamaecrista*, while the FT-SYM-type
nodules from section *Baseophyllum* were intermediately colonized (Fig. S3).

Light microscopy combined with immunohistochemistry using polyclonal antibodies
against *P. phymatum* STM815[T] and *Cupriavidus taiwanensis* LMG19424[T] indicated the
presence of *Paraburkholderia* as symbionts in nodules of the tree species *C. eitenorum*
(Fig. 5A, B). Specific immunogold localization of the *P. phymatum* antibody to the
bacteroids was confirmed under the TEM (Fig. 5C, D). Further immunogold localization
with JIM5 revealed HG epitopes on the cell walls surrounding both the FTs (Fig. 5E) and
the invasive ITs (5F) in the *C. eitenorum* nodules containing their *Paraburkholderia*
symbionts.

Phylogenetic analysis of nitrogen-fixing symbionts: housekeeping genes and ITS region

The diversity of rhizobia nodulating the various *Chamaecrista* species in the present study
was examined by MLSA to establish the phylogenetic relationships between symbiont
strains associated with tree and treelet *Chamaecrista* species *vis-a-vis* strains deposited
in the databases plus those isolated in our earlier study of symbionts from subshrub
Chamaecrista spp. (Santos et al. 2017) (Fig. 6). Most strains were grouped in the genus
Bradyrhizobium on the basis of close similarity to sequences of type strains. This genus
comprises various supergroups, including two mega-clades: I, the *B. japonicum* group,
and II, the *B. elkanii* group (Menna et al. 2009; Avontuur et al. 2019; Ormeno-Orillo and
Martinez-Romero, 2019). One cluster of *Chamaecrista* strains (Tree cluster I; Fig. 6)
consisting of isolates from the trees *C. ensiformis* var. *plurifoliolata, C. bahiae* and *C.*

duartei were located within the *B. japonicum* supergoup, while the other cluster of
Chamaecrista tree strains (Tree cluster II; Fig. 6), consisting only of isolates from *C.*
ensiformis var. *plurifoliolata* were located within the *B. elkanii* supergroup. Mega-clade I
also contained a large cluster of strains formed exclusively of bradyrhizobia isolated by
Santos et al. (2017) from root nodules of subshrub *Chamaecrista* species (Subshrub-
Shrub cluster; Fig. 6).

Analysis of the ITS region (16S–23S rDNA) (Fig. S4) suggested congruency with
the MLSA phylogeny, with two clusters containing strains from nodules of tree
Chamaecrista spp. plus one Subshrub-Shrub cluster. Strains from subshrubs formed a
separate cluster from all the *Bradyrhizobium* groups (Fig. S4). The ITS analysis indicates
the large differences in the DNA sequences of these *Chamaecrista* isolates, as it is one
of the strongest tools for discriminating between bradyrhizobial populations owing to its
high degree of variation providing greater powers of resolution (Willems et al., 2003). In
general, the *rrs* (16S rRNA) phylogeny (Fig. S5) was congruent with the MLSA and ITS
phylogenies, but with less precision and with modification in the position of some
bradyrhizobial isolates. None of the Bahia *Chamaecrista* strains clustered with the
described species from *C. fasciculata*, *B. frederickii* (Urquiaga et al., 2019) and *B. niftali*
(Klepa et al., 2019), both of which were isolated in the USA.

Confirming the immunogold microscopy observations for this species (Fig. 5A – D),
both strains isolated from nodules on the tree *C. eitenorum* sampled in Chapada
Diamantina belonged to the genus *Paraburkholderia*. A concatenated (16S rDNA + *recA*)
phylogenetic analysis demonstrated that they are potentially a new species (Fig. S6)
related to *P. nodosa* that was originally isolated from nodules of *Mimosa scabrella* (Chen
et al., 2007).

Phylogenetic analysis of nitrogen-fixing symbionts: symbiotic genes

The *nodC* (Fig. 7) and *nifH* (Fig. S7) bradyrhizobia phylogenies were relatively congruent
with the concatenated housekeeping and ITS phylogenies, with both forming the same
two Tree I and Tree II clusters as the MLSA, but the single subshrub-shrub cluster (SSC)
comprising strains from this study and that of Santos et al. (2017) was divided into two
sub-groups (SSCI & SSCII) in the *nodC* phylogeny (Fig. 7). SSCI corresponded to Cluster
1 of Santos et al. (2017) but now also incorporated *C. rotundifolia* and *C. ramosa*
symbionts from the present study, while SSCII was more heterogenous, grouping several
Santos et al. (2017) strains with a *C. blanchetii* strain from the present study

(BRUESC623) plus strains from various other legumes. A single strain (BRUESC1034)

isolated from the shrub *C. repens* occupied a unique position outside any of the main

nodC clusters. For the tree symbionts, Tree cluster II (TCII) constituted a tight group of

six strains from *C. ensiformis* var. *plurifoliolata*, while Tree cluster I (TCI) constituted the

main group of *Chamaecrista* tree strains; these were related to *Bradyrhizobium* type

species that were isolated in Brazil from various legumes (except for *B. iriomotense*

EK05[T] from Japan). The *nifH* phylogeny differed from the *nodC* one in that the two SSC

sub-groups were not apparent, with all the strains being grouped into a single cluster with

Bradyrhizobium sp. Tv2a-2 2 (isolated from the caesalpinioid legume *Tachigali versicolor*

in Barro Colorado Island of Panama; Tian et al., 2015) and *B. ganzhouense*, a symbiont

of *Acacia melanoxylon* (Lu et al. 2014).

The two strains of *Paraburkholderia* isolated from *C. eitenorum* (BRUESC1092,

BRUESC1093) formed a tight cluster in the *nodC* phylogeny with *Paraburkholderia* sp.

BRUESC684 (Fig. S8), a strain isolated from *Calliandra viscidula* collected in Chapada

Diamantina in Bahia State (Silva et al., 2018), but it should also be noted that both *C.*

eitenorum strains had symbiosis gene sequences which were closely related to *P. nodosa*

and *P. mimosarum* which are widely isolated symbionts of *Mimosa* species in Brazil

(Bontemps et al., 2010). Strains BRUESC1092 and BRUESC1093 also grouped with the

P. nodosa type strain Br3437[T] in the *nifH* gene phylogeny (Fig. S9).

Nodulation ability and host range of the Chamaecrista *rhizobia*

From the 28 strains of *Bradyrhizobium* isolated from nodules of seven *Chamaecrista*

species (*C. bahiae, C. duartei, C. eitenorum, C. ensiformis* var. *plurifoliolata, C. blanchetii,*

C. ramosa, C. repens and *C. rotundifolia*) 23 were tested for their nodulation ability on

Siratro, and 15 were tested on six *Chamaecrista* spp. The tested strains represented the

whole range of growth habits in the genus, from trees to subshrubs (Table 2), but also

the four *nodC* genotype clusters identified in Fig. 7 (SSCI, SSCII, TCI and TCII). Five of

the strains were isolated from the tree *C. ensiformis* var. *plurifoliolata*; three of these,

representing *nodC* TCI (BRUESC1010) and TCII (BRUESC964), and BRUESC1011 (not

included in the *nodC* phylogeny) formed nodules on their homologous host (the other two

strains, BRUESC956 and BRUESC967 were not tested on *C. ensiformis*), and four of the

five strains nodulated the subshrub *C. rotundifolia* (BRUESC964 was not tested).

However, no nodules were observed when one of the *C. ensiformis* var. *plurifoliolata*

strains (BRUESC1011) was inoculated onto another *Chamaecrista* tree species (*C.*

bahiae). A *Bradyrhizobium* strain isolated from *C. bahiae* (BRUESC952) nodulated *C.*

rotundifolia, but not *C. blanchetii* and *C. desvauxii*, while strain BRUESC1107 from *C.*
duartei formed effective nodules on *C. desvauxii*. The majority of the 19 *Bradyrhizobium*
strains tested nodulated Siratro; the exceptions were BRUESC623 from *C. blanchetii*
BRUESC964 from *C. ensiformis* var. plurifoliolata, BRUESC1033 and BRUESC1034
from *C. repens*, and BRUESC1106 from *C. duartei* (Table 2). Finally, the two
Paraburkholderia strains isolated from the tree *C. eitenorum* (BRUESC1092 and
BRUESC1093) effectively nodulated other tree species in the section *Apoucouita*, *C.*
duartei and *C. ensiformis*, but nodulated *Mimosa pudica* ineffectively, and both failed to
nodulate Siratro.

Phylogenetic analysis of the genus Chamaecrista

The individual phylogenetic analyses of *Chamaecrista* did not show apparent
incongruence between the ITS and *trnL-F* regions (Fig. 8), but as already described for
Chamaecrista (Conceição et al., 2009; Rando et al., 2016), the ITS region gives higher
resolution within the sections. In the *Chamaecrista* phylogeny (Fig. 8), three main large
clades emerged with high support (PP=1). These groups corresponded to the section
Apoucouita embracing the arborescent rainforest species, the section *Baseophyllum*
consisting of treelets and shrubs from Campo Rupestre and Caatinga, and the sections
Absus and *Chamaecrista*, which contain the highest diversity of species in the genus,
with most species being shrubs and subshrubs. Interestingly, these main clades
highlighted here are congruent with the phylogeny of the nitrogen-fixing microsymbionts,
and also demonstrate that the FT phenotype in which bacteroids are enclosed in cell wall
material, is apparently confined to the arborescent species in section *Apoucouita*,
whereas in species from the other clades/sections the microsymbionts are mainly
contained within symbiosomes. Interestingly, however, there are also a number of
species, particularly in section *Baseophyllum*, but also scattered amongst the other
sections, that are intermediate (FT-SYM) with regard to their bacteroid phenotype (Fig.
8).

Discussion

The occurrence of fixation threads (FTs), symbiosomes or intermediates depends on the
taxonomy, growth habit, and habitat of the Chamaecrista *host*

The 26 new reports in the present study raises the number of confirmed nodulating

Chamaecrista species to 74 (Sprent, 2009; Santos et al. 2017; Faria et al. 2022; Table

S4), now representing nearly a quarter of this highly speciose genus. In general, tissue

distribution in the interior of *Chamaecrista* nodules is similar to that of indeterminate

nodules from other neotropical Caesalpinioid trees, such as *Anadenanthera peregrina*

(Gross et al., 2002), *Mimosa* spp. (dos Reis Junior et al., 2010), and *Dimorphandra* spp.

(Fonseca et al., 2012). However, the present study has also confirmed that *Chamaecrista*

nodules are unique within the Caesalpinioideae since their infected cells can, depending

on species, have their bacteroids either retained within FTs, as occurs in nodules on all

other non-Mimosoid nodulated Caesalpinioideae, or have their rhizobial symbionts

released into symbiosomes (SYM-type nodules), as occurs in all mimosoid and most

papilionoid nodules so far reported (Naisbitt et al., 1992; Sprent, 2001; Sprent 2009;

Sprent et al., 2017; Faria et al. 2022). The distribution of FT- versus SYM-type nodules

in *Chamaecrista* appears to depend on the growth habit of the plant *i.e.,* current

knowledge, reinforced by considerable additional data from the present study, indicates

that tree species are all FT-type, whereas smaller shrub, subshrub and herbaceous

species, which represent the majority of *Chamaecrista* species, tend to be SYM-types

(Naisbitt et al. 1992). Interestingly, the present study has also confirmed using SEM and

TEM that some treelet and large shrub species exhibit an intermediate (FT-SYM) type of

nodule in which the FTs are less distinct (Naisbitt et al. 1992), suggesting a transitional

stage between FTs and symbiosomes in *Chamaecrista* that has not so far been observed

in other non-Mimosoid grade Caesalpinioideae symbioses (Faria et al. 2022).

Composition of the FT in Chamaecrista species

As with other non-mimosoid Caesalpinioideae trees, such as *Dimorphandra* (Fonseca et

al., 2012), *Erythrophleum* and *Moldenhawera* species (De Faria et al. 2022), the FTs in

Chamaecrista tree species comprise a cell wall containing homogalacturonan (HG)

epitopes as recognized by the monoclonal antibody, JIM5, which labels unesterified

pectin. The present study has demonstrated that *Chamaecrista* FTs also contain

cellulose; this concurs with the study of Faria et al. (2022) who have recently

demonstrated that the cell wall of FTs in nodules on a range of non-mimosoid

Caesalpinioideae contain several pectic components, but also cellulosic ones. Indeed,

there is mounting evidence that the wall of the FT differs little from the IT cell wall (and

plant cell walls in general) except that the FT wall is generally thinner and contains less

unesterified pectin (JIM5 epitope) (Fonseca et al. 2012; Faria et al. 2022; this study). It

should also be noted that FTs are not simple cell wall-bound structures, but are similar to

"conventional" symbiosomes in possessing a host membrane that surrounds the cell wall

(Faria et al. 2022), and hence the relative absence of unesterified pectin in FTs, as well

as their relatively thin walls compared to ITs is likely to be an adaptation to allow the

easier exchange of nutrients and O_2 between the rhizobial bacteroids and the host

cytoplasm across this membrane (Fonseca et al. 2012; Faria et al. 2022). Such nutrient

and gaseous exchange would most likely be impeded by the thick cell wall of a typical IT

containing pectin that has been stiffened via cross-linking through the action of pectin

methylesterases (PME) (Su, 2023).

Our study has also revealed more about the ultrastructure of the FT-SYM- and SYM-

type *Chamaecrista* nodules, including the histochemical nature of the symbiosomes. For

example, JIM5-labelled material could be observed in a matrix that surrounds the

symbiosomes in mature infected cells from nodules of some SYM and FT-SYM-type

Chamaecrista shrub and subshrub species in the present study from Brazil, but also in

the Asian subshrub species *C. pumila* (Rathi et al. 2018). The JIM5 labelling observed in

the FT-SYM-type nodules is most likely because these are essentially thinner versions of

FTs, and still have obvious cell walls surrounding their bacteroids (Naisbitt et al. 1992;

this study), but it is curious that some SYM-type nodules also possess pectin without any

obvious walls. In pea nodules, although glycoproteins and glycolipids were observed,

symbiosomes contained neither polysaccharides nor cell wall material (Perotto et al.,

1991, 1995). This implies that our observations of bacteroids in some SYM-type

Chamaecrista nodules surrounded by partially methyl-esterified and unesterified HG

epitopes (JIM5) are different from the symbiosomes housing bacteroids within nodules

on species in the "advanced" Inverse Repeat-Lacking Clade (IRLC); the latter contain

greatly-enlarged endo-reduplicated bacteroids that have completely lost their ability to

divide and to proliferate outside the host plant i.e. they are essentially organelles (Brewin,

1991; Gage, 2004; Oono et al. 2010; Sprent et al. 2017; Ardley & Sprent, 2021;

Mathesius, 2022). On the other hand, such terminally differentiated bacteroids are not the

norm outside the IRLC, so they are clearly not essential for legume nodule functioning,

although they may be more efficient at fixing N_2 (Mathesius, 2022). In most other legume

nodules, undifferentiated bacteroids are housed within symbiosomes, and it could be

argued that the *Chamaecrista* SYM-type nodules are essentially like these, especially

those in the Mimosoid clade (De Faria et al. 2022 and references therein). Therefore, as

suggested in the earlier study on a much narrower range of species (Naisbitt et al. 1992),

it would indeed appear that *Chamaecrista* contains a range of nodule phenotypes from

full FTs in the trees in the section *Apoucouita* through intermediates (FT-SYM) in treelets

and large shrubs (e.g. in the section *Baseophyllum*, although not exclusively) to
"standard" symbiosomes (SYM-type) in the majority of *Chamaecrista* species in the
sections *Absus* and *Chamaecrista*. The comparative efficiencies of the different nodule
types to fix N are not yet clear, but the number of bacteroids per cell (Fig. S3, this study)
are significantly reduced in the FT-type compared to the SYM-type nodule, with the FT-
SYM-types being intermediate in both parameters, which suggests that the possession
of a cell wall around the bacteroids reduces the degree to which host cells can be packed
with N-fixing rhizobia. This, together with and reduced proportion of infected cells per
nodule (Naisbitt et al. 1992) is likely to reduce the overall efficiency of the FT-type nodule.
598
599 *The influence of plant habit and biome on the diversity of nitrogen-fixing symbionts in*
600 *Chamaecrista*
601
Our study plus that of Santos et al. (2017) has demonstrated that in common with almost
all non-Mimosoid legumes of the subfamily Caesalpinioideae (Fonseca et al., 2012;
Parker, 2015; Cabral Michel et al., 2021; Avontuur et al. 2021), symbiotic bacteria isolated
from nodules across the genus *Chamaecrista* belong almost exclusively to the genus
Bradyrhizobium. Both housekeeping genes and ITS sequences of *Chamaecrista* strains
were congruent and taken together with the two symbiosis related genes (*nifH* and *nodC*)
showed that at least a dozen strains could represent putative novel species of
Bradyrhizobium. Also, *C. ensiformis,* a widespread arborescent species from tropical
forests, had the highest diversity of *Bradyrhizobium* symbionts, differing from its closely-
related "cousin" in the section *Apoucouita*, *C. eitenorum,* which has a more restricted
distribution in SDTF.
The phylogenetic analysis of both the *nodC* and *nifH* genes showed that the
Bradyrhizobium strains obtained from *Chamaecrista* nodules sampled in Bahia formed
clearly separated branches suggesting that symbiotic genes were probably vertically
transmitted, as was already indicated by studies on bradyrhizobia from other legumes
(Moulin et al., 2004; Parker, 2015, Fonseca et al., 2012; Stepkowski et al., 2007, 2018).
These results also suggest a very high diversity of bradyrhizobia nodulating
Chamaecrista trees, shrubs and subshrubs native to Brazil, and to Bahia in particular.
The *nodC* phylogeny supported the likelihood of there being co-evolution and specificity
of the *Chamaecrista-Bradyrhizobium* symbiosis since the rhizobia of the subshrub
Chamaecrista species in sections *Absus* and *Chamaecrista* that are highly divergent from
their distant cousins in the rainforest tree section *Apoucouita* also had highly divergent

nodC sequences. On the other hand, the different environments that these plant species
occupy, especially their soils (e.g. pH), will also play a part in their selection of particular
symbionts, as demonstrated for *Mimosa* in the Cerrado (Pires et al. 2018), and for
Chamaecrista in India (Rathi et al. 2018). Indeed, although the cross-inoculation
experiments demonstrated some degree of specificity, particularly for *Chamaecrista* tree
and treelet species, the fact that the widely distributed subshrub *C. rotundifolia* is
nodulated by *Bradyrhizobium* strains from all the observed *nodC* clades, including those
from the section *Apoucouita* species *C. bahiae* and *C. ensiformis*, as well as the ability of
most of the tested isolates to nodulate Siratro, suggests that Brazilian *Chamaecrista*
symbionts can be cosmopolitan in their selection of hosts outside their normal native
ranges, despite their apparent co-evolution with their (often) endemic hosts. A similar
observation was made with *Mimosa*, another large nodulated legume genus that has
radiated in Bahia, albeit with Beta-rhizobial symbionts rather than bradyrhizobia
(Bontemps et al. 2010).

Although bradyrhizobia are clearly the principal symbionts of *Chamaecrista* in Brazil
and elsewhere (see Introduction), other rhizobial types can nodulate the genus, such as
Sinorhizobium (*Ensifer*) in alkaline soils in India (Rathi et al. 2018). In the present study,
the tree species *C. eitenorum*, which is native to the SDTF, had the apparently unique
property of being nodulated by strains of *Paraburkholderia* related to those that nodulate
Mimosa and *Calliandra* spp. in the same environment (Bontemps et al. 2010; Silva et al.
2018); indeed, the fact that these isolates were also capable of nodulating *C. duartei* and
C. ensiformis in cross-inoculation studies suggests that *Chamaecrista-Paraburkholderia*
symbioses might be relatively abundant, at least among the tree species. Further to this,
the *C. eitenorum- Paraburkholderia* interaction is the first confirmed report of a Beta-
rhizobial symbiosis in nodules from a Caesalpinioideae species outside the Mimosoid
clade (LPWG, 2017), and it is also the first report of FTs being occupied by this type of
microsymbiont, which demonstrates that the FT phenotype is not controlled by any
particular rhizobial type (e.g. *Bradyrhizobium*), but is an inherent plant characteristic,
which might be expected if the "host controls the party" (Ferguson et al. 2019).

Concluding remarks

Our molecular phylogeny of *Chamaecrista* exhibits the same relationship among the
monophyletic clades already observed in other studies (Conceição et al., 2009; Rando et
al., 2016; Mendes et al., 2020; Souza et al., 2021), even with the incorporation of new

taxa. The small section *Apoucouita* remains a monophyletic group with the addition of six new arborescent species. The section also appears as a sister group of all remaining species of the genus, corroborating that its habitat and habit diverged distinctly from the others, and that it may be basal within the genus. The shifts of growth habit appear to be correlated with the habitat changes in *Chamaecrista*. In the genus, the diversification occurred from rain forests with arborescent habits to open areas (savannahs and SDTF) with smaller shrub and subshrub growth habits (Conceição et al., 2009; Coutinho et al. 2016; Souza et al., 2021).

We propose that the lower nutrient soils (and several other stresses, such as seasonal drought) associated with savannahs and SDTF not only drove a reduction in growth habit, but also necessitated a greater dependency on a more reliable nodulating symbiosis, as the smaller root systems of shrub and subshrub *Chamaecrista* species have a reduced access to a much more limited pool of soil N compared to large rainforest-dwelling trees that also have a higher capacity to (re)cycle their N. Therefore, the shrub and subshrub *Chamaecrista* species have largely rejected the FT-type symbiosis of their arboreal cousins in the section *Apoucouita*, and have gradually adopted (via the intermediate FT-SYM-type nodule) the more intimate SYM-type nodule, which with its full incorporation of the bacteroids into the host cytoplasm exhibits the compartmentalization that is linked with more stable and efficient symbioses (Parniske, 2018; Chomicki et al. 2020; Faria et al. 2022; Libourel et al. 2023; James 2023; Mohd-Radzman & Drapek, 2023). In this respect, the FT versus SYM pattern revealed by the present study across the highly speciose genus *Chamaecrista* mirrors that across the Caesalpinioideae subfamily as a whole i.e. that the hugely diverse (in terms of both growth habits and number of genera) Mimosoid clade has retained nodulation by rejecting the less intimate and relatively unstable FT-type nodule of its few nodulating cousins in the Caesalpinioideae grade that subtends it (with the notable exception of *Chamaecrista*), and by adopting the SYM-type nodule has avoided the massive losses of nodulation evident in the non-Mimosoid Caesalpinioideae (Faria et al. 2022).

ACKNOWLEDGEMENTS

The authors thank Dr Alan Prescott and Tamanatha Jithesh for assistance with image analysis, and Professor A. Nogueira and Dr Chrizelle Beukes for critical reading and suggestions. We would like to thank FAPESB for a grant to PAC Alves via the postgraduate Program of Biology and Biotechnology of Microorganisms, and the Center

for Electron Microscopy at UESC for infrastructure. EK James acknowledges funding
from CAPES via the Ciençia sem Fronteiras programme.

REFERENCES

Avontuur JR, Palmer M, Beukes CW, Chan WY, Coetzee MP, Blom, J, ... &
Steenkamp, E. T. 2019 Genome-informed *Bradyrhizobium* taxonomy: where to from
here? *Systematic and Applied Microbiology*, **42:** 427-439.

Beukes CW, Stepkowski T, Venter SN, Cłapa T, Phalane FL, le Roux MM, Steenkamp
ET. 2016. Crotalarieae and Genisteae of the South African great escarpment are
nodulated by novel *Bradyrhizobium* species with unique and diverse symbiotic loci.
Molecular Phylogenetics and Evolution **100**: 206-218.

Bontemps C. Elliott GN, Simon MF, dos Reis Junior, F.B., Gross, E., Lawton, R.C.,
Neto, N.E., Loureiro, M.F., de Faria, S.M., Sprent, J.I., James, E.K., Young, J.P.W.
2010. *Burkholderia* species are ancient symbionts of legumes. *Molecular Ecology* **19:** 44-
52.

708
Brewin NJ. 1991. Development of the legume root nodule. *Annual Review of Cell Biology*
7: 191-226. 1991.

Cabral-Michel D, Costa EM, Guimarães AA, Carvalho TS, Castro Caputo PS,
Willems, A, Moreira FMS. 2021. *Bradyrhizobium campsiandrae* sp. nov.,
a nitrogen-fixing bacterial strain isolated from a native leguminous tree from the Amazon
adapted to flooded conditions. *Archives of Microbiology* **203:** 233–240.

Chomicki G, Werner GDA, West SA, Kiers ET. 2020. Compartmentalization drives the
evolution of symbiotic cooperation. *Philosophical Transactions Royal Society of London.*
Series B: Biological Sciences **375:** 20190602.

Conceição AS, Queiroz LP, Lewis GP, de Andrade MJG, de Almeida PRM,
Schnadelbach AS, Van Den Berg C. 2009. Phylogeny of *Chamaecrista* (Leguminosae
Caesalpinioideae) based on nuclear and chloroplast DNA regions. *Taxon* **58:** 1168-1180.

Coutinho IAC, Rando, JG, Conceição AS, Meira RMSA. 2016. A study of the
morphoanatomical characters of the leaves of *Chamaecrista* (L.) Moench sect.
Apoucouita (Leguminosae-Caesalpinioideae). *Acta Botanica Brasilica* doi: 10.1590/0102-

33062016abb0035

Darriba. D, Taboada GL, Doallo R, Posada D. 2012. JModelTEst 2: more models, new heuristics and parallel computing. *Nature Methods* **9:** 772.

de Lajudie P, Willems A, Nick G, Moreira F, Molouba F, Hoste B, Torck U, Neyra M, Collins MD, Lindström K, Dreyfus B, Gillis M. 1998. Characterisation of tropical tree rhizobia and description of *Mesorhizobium plurifarum* sp. nov. *International Journal of Systematic Bacteriology* **48:** 369–382.

Delaux PM, Radhakrishnan G, Oldroyd G. 2015. Tracing the evolutionary path to nitrogen-fixing crops. *Current Opinion in Plant Biology* **26:** 95-99.

Diabate M, Munive A, de Faria SM, Ba A, Dreyfus B, Galiana A. 2005. Occurrence of nodulation in unexplored leguminous trees native to the West African tropical rainforest and inoculation response of native species useful in reforestation. *New Phytologist* **166:** 231-239.

dos Reis Junior FB, Simon MF, Gross E, Boddey RM, Elliott GN, Neto NE, Loureiro MF, Queiroz LP, Scotti MR, Chen W-M et al. 2010. Nodulation and nitrogen fixation by *Mimosa* spp. in the Cerrado and Caatinga biomes of Brazil. *New Phytologist* **186:** 934-946.

Doyle JJ. 1998. Phylogenetic perspective on nodulation: evolving views of plants and symbiotic bacteria. *Trends in Plant Science* **3:** 473-478.

Drummond AJ, Ashton B, Cheung M, Heled J, Moir R, Stones-Havas S, Thierer T, Wilson A. 2012. Geneious Version 6.1.6. Available from: <http://www.geneious.com>.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32: 1792-1797.

Estrada-de los Santos, P., Palmer, M., Chávez-Ramírez, B., Beukes, C., Steenkamp, E.T., Briscoe, L., Khan, N., Maluk, M., Lafos, M., Humm, E. et al. 2018. Whole genome analyses suggest that *Burkholderia* sensu lato contains two additional novel genera (*Mycetohabitans* gen. nov., and *Trinickia* gen. nov.): implications for the evolution of diazotrophy and nodulation in the *Burkholderiaceae. Genes* **9:** 389 doi:10.3390/genes9080389

Faria SM, McInroy S, Sprent JI. 1987. The occurrence of infected cells, with persistent
infection threads, in legume root nodules. *Canadian Journal of Botany* **65:** 553-558.

Faria SM, Diedhiou AG, de Lima HC, Ribeiro RD, Galiana A, Castilho AF, Henriques
JC. 2010. Evaluating the nodulation status of leguminous species from the Amazonian
forest of Brazil. *Journal of Experimental Botany* **61:** 3119-3127.

Faria SM, Ringelberg JJ, Gross E, Koenen EJM, Cardoso D, Ametsitsi GKD,
Akomatey J, Maluk M, Tak N, Gehlot HS, Wright KM, Teaumroong N, Songwattana
P, de Lima HC, Prin Y, Zartman CE, Sprent JI, Ardley J, Hughes CE, James EK 2022.
The innovation of the symbiosome has enhanced the evolutionary stability of nitrogen
fixation in legumes. *New Phytologist* **235:** 2365-2377.

Ferguson BJ, Mens C, Hastwell AH, et al. 2019. Legume nodulation: The host controls
the party. *Plant Cell and Environment* **42:**41–51. https://doi.org/10.1111/ pce.13348

Flora do Brasil 2020 em construção. 2019. *Jardim Botânico do Rio de Janeiro.* [WWW
document] URL http://floradobrasil.jbrj.gov.br/. [accessed 30 January 2018].

Fonseca MB, Peix A, de Faria SM, Mateos PF, Rivera LP, Simões-Araujo JL, França
MGC, Isaias RMS, Cruz C, Velázquez E et al. 2012 Nodulation in *Dimorphandra wilsonii*
Rizz. (Caesalpinioideae), a threatened species native to the Brazilian Cerrado. *PlosOne*
7: e49520.

Gage DJ. 2004. Infection and invasion of roots by symbiotic, nitrogen-fixing rhizobia
during nodulation of temperate legumes. *Microbiology and Molecular Biology Review* **68:**
280-300.

Gross E, Cordeiro L, Caetano FH. 2002. Nodule ultrastructure and initial growth of
Anadenanthera peregrina (L.) Speg. var. *falcata* (Benth.) Altschul plants Infected with
rhizobia. Annals of Botany **90:** 175-183.

Guindon S, Gascuel O. 2003. A simple, fast and accurate method to estimate large
phylogenies by maximum-likelihood. Systematic. *Biology* **52:** 696-704.

Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis
program for Windows 95/98/NT. *Nucleic Acids Symposium Series* **41:** 95-98.

Han SZ, Wang, ET, Chen WX. 2005. Diverse bacteria isolated from root nodules of

Phaseolus vulgaris and species within the genera *Campylotropis* and *Cassia* grown in
China. *Systematic and Applied Microbiology* **28:** 265-276.

Irwin HS, Barneby RC. 1982. The American Cassiinae, a synoptical revision of
Leguminosae tribe Cassieae subtribe Cassiinae in the New World. *Memoirs of New York*
Botanical Garden **35:** 1-918.

James EK. 2023. Intimacy stabilizes symbiotic nodulation. *Nature Plants*
https://doi.org/10.1038/s41477-023-01438-5
789

Klepa MS, Urquiaga MC, Somasegaran P, Delamuta JRM, Ribeiro RA, Hungria MA.
2019. *Bradyrhizobium niftali* sp. nov., an effective nitrogen-fixing symbiont of partridge
pea [*Chamaecrista fasciculata* (Michx.) Greene], a native caesalpinioid legume broadly
distributed in the USA. *International Journal of Systematic and Evolutionary Microbiology*
69: 3448–3459.

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular Evolutionary
Genetics Analysis across computing platforms *Molecular Biology and Evolution* **35:**1547-
1549; doi: 10.1093/nar/25.24.4876.

Lafay B, Burdon JJ. 2007. Molecular diversity of legume root-nodule bacteria in Kakadu
National Park, Northern Territory, Australia. *PloS One* **2:**e277.

Lewis GP, Schrire B, Mackinder B, Lock M. 2005. *Legumes of the World*. London, UK:
Royal Botanic Gardens, Kew.

Legume Phylogeny Working Group (LPWG) 2021. The World Checklist of Vascular
Plants (WCVP): Fabaceae, vers. June 2021 (R. Govaerts, Ed.).
http://sftp.kew.org/pub/data_collaborations/Fabaceae/DwCA

Legume Phylogeny Working Group (LPWG). 2017. A new subfamily classification of
the Leguminosae based on a taxonomically comprehensive phylogeny. Taxon 66: 44-77

Libourel C, Keller J, Brichet L. *et al.* 2023. Comparative phylotranscriptomics reveals
ancestral and derived root nodule symbiosis programmes. *Nature Plants*
https://doi.org/10.1038/s41477-023-01441-w

Lu JK, Dou YJ, Zhu YJ, Wang SK, Sui XH, Kang LH. 2014. *Bradyrhizobium*
ganzhouense sp. nov., an effective symbiotic bacterium isolated from *Acacia*

melanoxylon R. Br. nodules. *International Journal of Systematic and Evolutionary Microbiology* **64:** 1900-1905.

MAPA. 2011. Secretaria de Defesa Agropecuária. *Instrução Normativa no 13*, de 24 de março de 2011.

Mendes TP, Souza AOD, Silva MJD. 2017. A new species hidden in the lowlands of Tocantins, Brazil: *Chamaecrista tocantinensis* (Fabaceae). *Systematic Botany* **42:** 326-337.

Mendes, TP, Oliveira, AOS, Silva, MJD 2020. Molecular phylogeny and diversification timing of the *Chamaecrista* sect. *Absus* subsect. *Absus* ser. *Paniculatae*, a newly circumscribed and predominantly endemic of the Cerrado Biome group. Phytotaxa **446:** 159-182.

Menna P, Barcellos FG, Hungria M. 2009. Phylogeny and taxonomy of a diverse collection of *Bradyrhizobium* strains based on multilocus sequence analysis of the 16S rRNA gene, ITS region and *glnII, recA, atpD* and *dnaK* genes. *International Journal of Systematic and Evolutionary Microbiology* **59:** 2934-2950.

Miller MA, W. Pfeiffer W, Schwartz T. 2010. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. *Gateway Computing Environments Workshop (GCE)*, New Orleans: LA, USA, pp. 1-8. doi: 10.1109/GCE.2010.5676129.

Mohd-Radzman NA, Drapek C. 2023. Compartmentalisation: A strategy for optimising symbiosis and tradeoff management. *Plant Cell and Environment* **46:** 2998–3011.

Morgulis A, Coulouris G, Raytselis Y, Madden TL, Agarwala R, Schäffer AA. 2008. Database indexing for production MegaBLAST searches. *Bioinformatics* **15:** 1757-1764.

Moulin L, Béna G, Boivin-Masson C, Stepkowski T. 2004. Phylogenetic analyses of symbiotic nodulation genes support vertical and lateral gene co-transfer within the *Bradyrhizobium* genus. *Molecular Phylogenetics and Evolution* **30:** 720-732.

Naisbitt T, James EK, Sprent JI. 1992. The evolutionary significance of the genus *Chamaecrista*, as determined by nodule structure. *New Phytologist* **122:** 487-492.

Oldroyd GED, Downie JA. 2008. Coordination of nodule morphogenesis with rhizobial infection in legumes. *Annual Review of Plant Biology* **59:** 519-546.

Oono R, Schmitt I, Sprent JI, Denison RF. 2010. Multiple evolutionary origins of legume traits leading to extreme rhizobial differentiation. *New Phytologist* **187:** 508–520.

Parker MA. 2012. Legumes select symbiosis island sequence variants in *Bradyrhizobium. Molecular Ecology* **21:** 1769-1778.

Parker MA. 2015. The spread of *Bradyrhizobium* lineages across host legume clades: from *Abarema* to *Zygia. Microbiology Ecology* **69:** 630-640.

Parker MA, Kennedy DA. 2006. Diversity and relationships of bradyrhizobia from legumes native to eastern North America. Canadian Journal of Microbiology **52:** 1148-1157.

Parker MA, Rousteau A. 2014. Mosaic origins of *Bradyrhizobium* legume symbionts on the Caribbean island of Guadeloupe. *Molecular Phylogenetics and Evolution* **77:** 110-115.

Parniske M. 2018. Uptake of bacteria into living plant cells, the unifying and distinct feature of the nitrogen-fixing root nodule symbiosis. *Current Opinion in Plant Biology* **44:** 164–174.

Perotto S, Donovan N, Drobak BJ, Brewin NJ. 1995. Differential expression of a glycosyl inositol phospholipid antigen on the peribacteroid membrane during pea nodule development. *Molecular Plant-Microbe Interactions* **8:** 560-568.

Perotto S, VandenBosch KA, Butcher GW, Brewin NJ. 1991. Molecular composition and development of the plant glycocalyx associated with the peribacteroid membrane of pea root nodules. *Development* **112:** 763-773.

Rambaut A. 2014. FigTree 1.4.2 [online]. Available from http://tree.bio.ed.ac.uk/software/figtree/ [accessed 17 October 2023].

Rando, JG, Zuntini, AR, Conceição, AS, van den Berg, C, Pirani, JR & Queiroz, LP. 2016. Phylogeny of *Chamaecrista* ser. *Coriaceae* (Leguminosae) unveils a lineage recently diversified in Brazilian Campo Rupestre vegetation. *International Journal of Plant Sciences* **177:** 3–17.

Rando, J.G.; Cota, M.M.T.; Conceição, A.S.; Barbosa, A.R.; Barros, T.L.A. 2020. *Chamaecrista* **in Flora do Brasil 2020.** Jardim Botânico do Rio de Janeiro. Available at:

<http://floradobrasil.jbrj.gov.br/reflora/floradobrasil/FB22876>. Accessed on: 25 Oct. 2021

Rathi, S., Nisha Tak, N., Bissa, G., Chouhan, B., Ojha, A., Adhikari, D., Barik, S.K., Satyawada, R.R., Sprent, J.I., James, E.K., et al. 2018. Selection of *Bradyrhizobium* or *Ensifer* symbionts by the native Indian caesalpinioid legume *Chamaecrista pumila* depends on soil pH and other edaphic and climatic factors. *FEMS Microbiology Ecology* 10.1093/femsec/fiy180

Rhem, M. F. K., Silva, V. C., dos Santos, J. M. F., Zilli, J. É., James, E. K., Simon, M. F., & Gross, E. 2021. The large mimosoid genus *Inga* Mill. (tribe Ingeae, Caesalpinioideae) is nodulated by diverse *Bradyrhizobium* strains in its main centers of diversity in Brazil. *Systematic and Applied Microbiology*, 126268.

Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19:** 1572–1574

Salmi A, Boulila F, Bourebaba Y, Le Roux C, Belhadi D, De Lajudie P. 2018. Phylogenetic diversity of *Bradyrhizobium* strains nodulating *Calicotome spinosa* in the Northeast of Algeria. *Systematic and Applied Microbiology* **41:** 452-459.

Santos JMF, Alves PAC, Silva VC, Rhem MFK, James, EK, Gross E. 2017. *Bradyrhizobium* diversity and structural aspects of nodulation on herbaceous *Chamaecrista* (Moench) species. Systematic and Applied Microbiology **40:** 69-79.

Silva VC, Alves PAC, Rhem MFK, dos Santos JMF, James EK, Gross E. 2018. Brazilian species of *Calliandra* Benth. (tribe Ingeae) are nodulated by diverse strains of *Paraburkholderia*. *Systematic and Applied Microbiology* **41:** 241-250.

Singer SR, Maki SL, Farmer AD, Ilut D, May GD, Cannon SB, Doyle JJ. 2009. Venturing beyond beans and peas: what can we learn from *Chamaecrista*? *Plant Physiology* **151:** 1041-1047.

Souza, AO, Lewis GP, Silva MJ. 2021. A new infrageneric classification of the pantropical genus *Chamaecrista* (Fabaceae/ Caesalpinioideae) based on a comprehensive molecular phylogenetic analysis and morphology. Botanical Journal of the Linnean Society 197: 350-395.

Sprent JI. 2009 *Legume Nodulation: A Global Perspective*. Wiley-Blackwell, USA.

Sprent, J. I. 2001. *Nodulation in legumes*. London, UK: Royal Botanic Gardens, Kew.

Sprent JI, Ardley JK, James EK. 2013. From North to South: A latitudinal look at legume
nodulation processes. *South African Journal of Botany* **89:** 31-41.

Sprent JI, Ardley JA, James EK. 2017. Biogeography of nodulated legumes and their
nitrogen fixing symbionts. *New Phytologist* **215:** 40-56.

Sprent JI, Geoghegan IE, Whitty PW, James EK. 1996. Natural abundance of ^{15}N
and ^{13}C in nodulated legumes and other plants in the Cerrado and neighbouring regions
of Brazil. *Oecologia* **105:** 440-446.

Stępkowski T, Hughes CE, Law IJ, Markiewicz Ł, Gurda D, Chlebicka A, Moulin, L.
2007. Diversification of lupine *Bradyrhizobium* strains: evidence from nodulation gene
trees. *Applied and Environmental Microbiology* **73:** 3254-3264.

Stępkowski T, Banasiewicz J, Granada CE, Andrews M, Passaglia LMP. 2018.
Phylogeny and phylogeography of rhizobial symbionts nodulating legumes of the tribe
Genisteae. *Genes* **9:** 163; doi:10.3390/genes9030163

Su C. 2023. Pectin modifications at the symbiotic interface. *New Phytologist* **238:** 25–32.

Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Thompson DGH. 1997. The
CLUSTAL_X Windows Interface: Flexible Strategies for Multiple Sequence Alignment
Aided by Quality Analysis Tools. Nucleic Acids Research, 25, 4876-4882; doi:
10.1093/nar/25.24.4876.

Tian R, Parker M, Seshadri R, Reddy TB, Markowitz V, Ivanova N, Pati A, Woyke T,
Baeshen MN, Baeshen NA, Kyrpides N. 2015. High-quality permanent draft genome
sequence of *Bradyrhizobium* sp. Tv2a. 2, a microsymbiont of Tachigali versicolor
discovered in Barro Colorado Island of Panama. *Standards in Genomic Sciences.* **10:** 27.

Tsyganova AV, Brewin NJ, Tsyganov VE. 2021. Structure and development of the
legume-rhizobial symbiotic interface in infection threads. *Cells* **10:** 1050.
https://doi.org/10.3390/cells10051050

Urquiaga MC, Klepa MS, Somasegaran P, Ribeiro RA, Delamuta JRM, Hungria MA.
2019. *Bradyrhizobium frederickii* sp. nov., a nitrogen-fixing lineage isolated from nodules

of the caesalpinioid species *Chamaecrista fasciculata* and characterized by tolerance to
high temperature in vitro. *International Journal of Systematic and Evolutionary*
Microbiology

Weisburg WG, Barns SM, Pelletier DA, Lane DJ. 1991. 16S ribosomal DNA
amplification for phylogenetic study. *Journal of Bacteriology* **173:** 697-703.

Willems A, Munive A, de Lajudie P, Gillis M. 2003. In most *Bradyrhizobium* groups
sequence comparison of 16S-23S rDNA internal transcriber spacer regions corroborates
DNA-DNA hybridizations. *Systematic and Applied Microbiology* **26:** 203-210.

Wood PJ, Fulcher RG, Stone BA. 1983. Studies on the specificity of interaction of
cereal cell wall components with Congo Red and Calcofluor. Specific detection and
histochemistry of $(1\rightarrow 3),(1\rightarrow 4),-\beta$-D-glucan. *Journal of Cereal Science* **1:** 95-110.

Yao Y, Wang R, Lu JK, Sui XH, Wang ET, Chen WX. 2014. Genetic diversity and
evolution of *Bradyrhizobium* populations nodulating *Erythrophleum fordii*, an evergreen
tree indigenous to the southern subtropical region of China. *Applied and Environmental*
Microbiology **80:** 6184-6194.

Yao Y, Sui XH, Zhang XX, Wang ET, Chen WX. 2015. *Bradyrhizobium erythrophlei* sp.
nov. and *Bradyrhizobium ferriligni* sp. nov., isolated from effective nodules of
Erythrophleum fordii. *International Journal of Systematic and Evolutionary Microbiology*
65: 1831-1837.

Table 1. Plant growth habit (and taxonomic section within *Chamaecrista*), plus symbiosome type (fixation threads, symbiosomes or intermediate), bacteroid size, reaction to calcofluor white and JIM5 observed in infected cells of root nodules from *Chamaecrista* spp. native to Bahia State, Brazil, and number of rhizobia strains isolated from each sample.

Species	Predominant habit (Section)	Symbiosome type*	Bacteroid size (μm)	Calcofluor white	JIM5	Number of strains
C. bahiae	Tree (*Apoucouita*)	FT	-	+	+	2
C. duartei	Tree (*Apoucouita*)	FT	-	+	+	4
C. ensiformis var. *plurifoliolata*	Tree (*Apoucouita*)	FT	-	+	+	15
C. eitenorum	Tree (*Apoucouita*)	FT	-	+	+	2
C. blanchetii	Treelet (*Baseophyllum*)	FT	-	+	+	1
C. × blanchetiformis	Treelet (*Baseophyllum*)	FT	-	+	+	-
C. brachystachya	Treelet (*Baseophyllum*)	FT	-	+	+	-
C. confertiformis	Treelet (*Baseophyllum*)	FT	-	+	+	-
C. zygophylloides	Shrub (*Absus*)	FT-SYM	5	-	+	-
C. belemii	Shrub (*Absus*)	FT-SYM	5	-	+	-
C. arrojadoana	Shrub (*Chamaecrista*)	FT-SYM	3	-	+	-
C. repens	Shrub (*Chamaecrista*)	FT-SYM	3-4	-	+	2
C. ramosa	Shrub (*Chamaecrista*)	FT-SYM	8	-	+	1
C. desvauxii	Shrub (*Chamaecrista*)	FT-SYM	3	-	+	-
C. pascuorum	Shrub (*Chamaecrista*)	SYM	3-4	-	+	-
C. rotundifolia	Shrub (*Chamaecrista*)	SYM	5	-	+	2
C. supplex	Subshrub (*Chamaecrista*)	SYM	4	-	+	-
C. serpens	Subshrub (*Chamaecrista*)	SYM	4	-	+	-
C. flexuosa	Subshrub (*Chamaecrista*)	SYM	3-4	-	+	-

* FT = Fixation Thread, SYM = Symbiosome; FT-SYM = Intermediate between FT and SYM.

Table 2. Nodulation ability on various plant hosts of rhizobial strains isolated from different *Chamaecrista* host species native to Bahia state, Brazil. All strains tested were *Bradyrhizobium* except for those indicated as *Paraburkholderia**. The *nodC* group that the *Bradyrhizobium* strains belonged to (Fig. 7) are indicated in parentheses: Subshrub-Shrub Cluster I (SSCI), Subshrub-Shrub Cluster II (SSCII), Tree Cluster I (TCI), Tree Cluster II (TCII), or not known.

Strain	Host	*Chamaecrista* spp. tested*	Siratro nodulation
BRUESC623 (SSCII)	*C. blanchetii* (Treelet)	*C. blanchetii* (+)	(-)
BRUESC956 (TCII)	*C. ensiformis* var. *plurifoliolata* (Tree)	*C. rotundifolia* (+)	(+)
BRUESC1010 (TCI)	*C. ensiformis* var. *plurifoliolata* (Tree)	*C. rotundifolia* (+) *C. ensiformis* (+)	nt
BRUESC964 (TCII)	*C. ensiformis* var. *plurifoliolata* (Tree)	*C. ensiformis* (+)	(-)
BRUESC967 (TCI)	*C. ensiformis* var. *plurifoliolata* (Tree)	*C. rotundifolia* (+)	(+)
BRUESC1011 (not known)	*C. ensiformis* var. *plurifoliolata* (Tree)	*C. bahiae* (-) *C. rotundifolia* (+) *C. ensiformis* (+)	(+)
BRUESC952 (not known)	*C. bahiae* (Tree)	*C. rotundifolia* (+) *C. blanchetii* (-) *C. desvauxii* (-)	(+)
BRUESC1033 (not known)	*C. repens* (Shrub)	*C. rotundifolia* (+) *C. blanchetii* (-)	(-)
BRUESC1034 (singleton)	*C. repens* (Shrub)	*C. bahiae* (+) *C. rotundifolia* (+)	(-)
BRUESC1106 (not known)	*C. duartei* (Tree)	*C. blanchetii* (-)	(-)
BRUESC1107 (not known)	*C. duartei* (Tree)	*C. desvauxii* (+)	(+)
BRUESC1102 (SSCI)	*C. rotundifolia* (Subshrub)	*C. desvauxii* (+)	nt
BRUESC1103 (SSCI)	*C. rotundifolia* (Subshrub)	*C.bahiae* (-) *C. blanchetii* (-)	(+)
*BRUESC1092 (*Paraburkholderia*)	*C. eitenorum* (Tree)	*C. duartei* (+) *C. desvauxii* (-) *C. ensiformis* (+)	(-)
*BRUESC1093 (*Paraburkholderia*)	*C. eitenorum* (Tree)	*C. duartei* (+) *C. desvauxii* (-) *C. ensiformis* (+)	(-)

nt = not tested; ns = not sequenced. (+), (-) = positive and negative nodulation. *Also nodulated *Mimosa pudica.*
BRUESC955 (TCI), BRUESC961 (TCI), BRUESC963 (TCII), BRUESC965 (TCII), BRUESC966 (not known), BRUESC968 (TCII), BRUESC969 (TCII), and BRUESC1091 (TCI) nodulated Siratro, but were not tested on any *Chamaecrista* species.

 Figure legends

Fig. 1. *Chamaecrista* species and their morphological variation across the various
sections of the genus. (A) Tree species of *C. bahiae* (Section *Apoucouita*) in the tropical
rainforest; (B) Flower and initial developing fruit of *C. duartei*; (C) General view of a *C.*
confertiformis plant in *campo rupestre* habitat; (D) Detail of a *C. blanchetii* (Section
Baseophyllum) fruit; (E) Supra-axillary fascicles of *C. repens* (Section *Chamaecrista*); (F)
Solitary flowers of *C. ramose* (Section *Chamaecrista*). Scale bars: (B) = 3 cm; (D, E, F) =
1 cm.

Fig. 2. Morphology and anatomy of nodules from species in the various sections of the
genus *Chamaecrista*. **A.** Nodules of *C. bahiae* (Section *Apoucouita*) under natural soil
conditions; **B.** View of nodule morphologies. An arrow indicates a branched nodule; **C.**
General view of a longitudinal section of a *C. bahiae* nodule showing infected tissue (it)
in the center surrounded by the inner (ic) and outer cortex (oc); **D.** Sector of a *C. bahiae*
nodule showing the invasion zone and part of the nitrogen (N) fixation zone; **E.** Proximal
part of a *C. bahiae* nodule showing infected cells in different stages of senescence; **F.**
Detail of mature infected cells of *C. bahiae* occupied by fixation threads and interstitial
uninfected cells; **G.** Detail of mature infected cells of *C. rotundifolia* (Section
Chamaecrista) occupied by symbiosomes and some interstitial non infected cells. Scale
bars: **A, B** = 1 cm; **C** = 500 µm; **D, E** = 50 µm; **F, G** = 20 µm.

Fig. 3. Fluorescence micrographs of Calcofluor White-stained semi-thin sections of
nodules from species in the various sections of the genus *Chamaecrista*. Fluorescence
was detected in the infected cells of: **A.** *C. bahiae* (Section *Apoucouita*); **B.** *C. duartei*
(Section *Apoucouita*); **C.** *C. confertiformis* (Section *Baseophyllum*) and **D.** *C.*
blanchetiformis (Section *Baseophyllum*). **E.** No fluorescence was detectable in the
infected cells of *C. rotundifolia* (Section *Chamaecrista*). **F.** Fluorescence was detected in
the newly-invaded cells of the invasion zone and in the early infected cells, but not in the
mature infected cells of *C. belemii* (Section *Absus* subs. *Zygophyllum*). Scale bars: **A, B,**
D, F = 50 µm; **C, E**= 20 µm.

Fig. 4. Scanning electron micrographs (SEMs) (**A – C**) and Transmission electron
micrographs (TEMs) (**D – F**) of infected cells in *Chamaecrista* nodules from species in
the various sections of the genus demonstrating the structure and morphology of the

bacteroids across the genus. **A.** *C. bahiae* (FT-type, Section *Apoucouita*); **B.** *C. repens* (FT-SYM-type, Section *Chamaecrista*)*;* **C.** *C. rotundifolia* (SYM-type, Section *Chamaecrista*)*.* **D – F.** TEMs of infected cells after immunogold localization of homogalacturonan epitopes with the monoclonal antibody JIM5, which recognizes a pectin epitope in plant cell walls. **D** *C. bahiae*; **E** *C. arrojadoana* (FT-SYM-type, Section *Chamaecrista*) showing gold particles in symbiosome membrane (wall) and matrix (arrows); **F** *C. supplex* (SYM-type, Section *Chamaecrista*) Scale bars: **A, C** = 10 μm; **E** = 10 μm; **B, D, F**= 1 μm.

Fig. 5. Light micrographs (**A, B)** and TEMs (**C – F**) of sections of *Chamaecrista eitenorum* var. *eitenorum* (Section *Apoucouita*) nodules. **A.** Infected tissue immunogold labelled with an antibody against *Paraburkholderia phymatum* STM815[T] plus silver enhacement; **B.** Infected tissue immunolabelled with an antibody against *Cupriavidus taiwanensis* LMG19434[T] plus silver enhancement (negative control). **C.** Immunogold localization on the bacteroid surface using an antibody against *Paraburkholderia phymatum* STM815[T] (arrows). **D.** Negative control using IGL buffer. **E.** Immunogold localization of homogalacturonan epitopes (arrows) with the monoclonal antibody JIM5, which recognizes a pectin epitope in plant cell walls, in an infected cell. **F.** A "classical" invasion infection thread strongly immunogold labelled with JIM5 (arrows). Scale bars: **A, B** = 50 μm, **C, D, E, F** = 0.5 μm.

Fig. 6. Maximum-likelihood phylogeny for the genus *Bradyrhizobium* based on the concatenated dataset consisting of sequences of the genes *atpD, dnaK, glnII, gyrB, recA* and *rpoB*. The isolates examined in this study are indicated in bold, and those from dos Santos et al. (2017) are indicated by *. The scale bar indicates the number of nucleotide substitutions per site. Numbers on branches are bootstrap values for 1000 replications (shown only when ≥70%).

Fig. 7. Maximum-likelihood phylogeny of the genus *Bradyrhizobium* based on *nodC*. Host association and specific geographic origin is listed in Table 1. The isolates examined in this study are indicated in bold, and those from dos Santos et al. (2017) are indicated by *. The scale bar indicates the number of nucleotide substitutions per site. Numbers on branches are bootstrap values for 1000 replications (shown only when ≥70%).

Fig. 8. A. Phylogeny of the Leguminosae family adapted from LPWG (2017) showing the genera in the Caesalpinioideae subfamily associated with symbionts. **B.** Phylogeny of *Chamaecrista* based on DNA sequences of nuclear ITS and plastidial *trnL-trnF* loci. Majority-rule consensus tree derived from Bayesian analysis; the values of posterior probability (PP) are indicated above the branches in decimal form. Taxonomic groups of associated symbionts are colored. Species indicated by *(*C. chapadae*, *C. zygophylloides*) were not included in the phylogenetic analysis, but based on morphology, it was possible to place them in their appropriate taxonomic group. Arrows indicate the ages of nodes as estimated by Rando et al. (2016). **C.** *C. bahiae*; **D.** *C. desvauxii*; **E.** *C. arrojadoana.* (photos Juliana Rando)

Fig. S1. Location and distribution of *Chamaecrista* nodule sampling in Bahia State (NE Brazil).

Fig. S2. Light micrographs (**A, C, E, G, I, K**) and transmission electron micrographs (TEMs) plus immunogold labelling with JIM5 (**B, D, F, H, J, L**) of nodules of *Chamaecrista* across four sections of the genus illustrating the range of nodule anatomical types from tree species with their bacteroids enclosed in fixation threads (FT) labelled with JIM5 (**A – D**) through intermediate types (FT-SYM) on treelets (**E – H**) to membrane-bound symbiosomes (SYM) on shrub and subshrub species that have little or no JIM5 signal (**I – L**). **A, B.** *Chamaecrista duartei* (section *Apoucouita*). **C, D.** *Chamaecrista ensiformis* (section *Apoucouita*). **E, F.** *Chamaecrista blanchettii* (section *Baseophyllum*). **G, H.** *Chamaecrista cytisoides* (section *Baseophyllum*). **I, J.** *Chamaecrista chapadae* (section *Absus*, subsect. *Absus*). **K, L.** *Chamaecrista mucronata* (section *Chamaecrista*). The FTs in the section *Apoucouita* tree species *C. duartei* (**A**) and *C. ensiformis* (**C**) are discernible in infected cells (*) at the light microscope level. The walls of the FTs are indicated by arrows in TEMs in **B** and **D,** but note that the FTs in *C. ensiformis* (**D**) are more densely labelled with JIM5 than those in *C. duartei* (**B**). However, in the latter species the electron dense cell walls are more apparent, thus illustrating that there is no direct relationship between cell wall density and the presence/absence of unesterified pectin (JIM5) in FTs. Bacteroids in nodules on the section *Baseophyllum* treelet species *C. blanchetii* (**E**) and *C. cytisoides* (**G**) are less clearly defined at the light microscope level and this is because at the TEM level (**F, H**) it is observed that bacteroids either have thin FT walls with sparse JIM5 labelling (arrows in **F, H**) or no walls at all (i.e., most of the bacteroids in *C. cytisoides*, **H**); however, more densely labelled walls surrounding bacteroids were also

occasionally observed within *C. cytisoides* infected cells (double arrows in **H**), suggesting that this species harbours both FTs and symbiosomes. The bacteroids in nodules on shrub and subshrub species *C. chapadae* (**I**) and *C. mucronata* (**K**) were even less distinct under the light microscope compared to the *Baseophyllum* treelet species, and this was also confirmed under the TEM where they were shown to be enclosed in symbiosomes Bars = 20 μm (**A, C, E, G. I, K**), 500 nm (**B, D, F, L**), 1 μm (**H, J**), b = bacteroid, w = plant cell wall in **B, D, F, H, J, L.**

Fig. S3. Proportional (%) occupation of infected cells by symbiotic rhizobia estimated by counting pixels from light micrographs similar to those presented in Fig S2A and C for FT-type, Fig. S2E and G for FT-SYM-type, and Fig. S2I and K for SYM-type nodules. Data are presented as the mean proportion (± s.d.) of each infected cell profile filled with toluidine blue-stained structures representing bacteroids; 7-9 sections were examined per symbiosome type (FT, FT-SYM, SYM).

Fig. S4. Maximum-likelihood phylogeny for the genus *Bradyrhizobium* based on the ITS region. Host associations and specific geographic origins are listed in Table 1. The isolates examined in this study are indicated in bold, and those from dos Santos et al. (2017) are indicated by *. The scale bar indicates the number of nucleotide substitutions per site. Numbers on branches are bootstrap values for 1000 replications (shown only when ≥70%).

Fig. S5. Maximum-likelihood phylogeny for the genus *Bradyrhizobium* based on 16S rDNA. Host association and specific geographic origin is listed in Table 1. The isolates examined in this study are indicated in bold, and those from dos Santos et al. (2017) are indicated by *. The scale bar indicates the number of nucleotide substitutions per site. Numbers on branches are bootstrap values for 1000 replications (shown only when ≥70%).

Fig. S6. Maximum-likelihood phylogeny for the genus *Paraburkholderia* based on the concatenated dataset consisting of sequences for the genes 16S rDNA and *recA*. The isolates examined in this study are indicated in bold. The scale bar indicates the number of nucleotide substitutions per site. Numbers on branches are bootstrap values for 1000 replications (shown only when ≥70%).

Fig. S7. Maximum-likelihood phylogeny for the genus *Bradyrhizobium* based on the *nifH* gene. Host association and specific geographic origin is listed in Table 1. The isolates examined in this study are indicated in bold, and those from dos Santos et al. (2017) are indicated by *. The scale bar indicates the number of nucleotide substitutions per site. Numbers on branches are bootstrap values for 1000 replications (shown only when ≥70%).

Fig. S8. Maximum-likelihood phylogeny for the genus *Parburkholderia* based on the *nodC* gene. Host association and specific geographic origin is listed in Table 1. The isolates examined in this study are indicated in bold. The scale bar indicates the number of nucleotide substitutions per site. Numbers on branches are bootstrap values for 1000 replications (shown only when ≥70%).

Fig. S9. Maximum-likelihood phylogeny for the genus *Paraburkholderia* based on the *nifH* gene. Host association and specific geographic origin is listed in Table 1. The isolates examined in this study are indicated in bold. The scale bar indicates the number of nucleotide substitutions per site. Numbers on branches are bootstrap values for 1000 replications (shown only when ≥70%).

Table S1. Accession numbers of aerial parts of *Chamaecrista* species sampled in Bahia (BA), Brazil and deposited in the Herbarium of the Universidade Estadual de Santa Cruz (UESC). Also included is information about growth habit, vegetation type and locality where the *Chamaecrista* species were sampled in BA specifically for this study.

Table S2. Genbank accession numbers of gene sequences from strains isolated in the present study.

Table S3. Voucher information and GenBank accession numbers of the nuclear ITS and plastidial *trnL-trnF* loci sequences included in the phylogeny of *Chamaecrista* (Fig. 8). New sequences are marked with an asterisk (*).

Table S4. Confirmed positive global nodulation reports for *Chamaecrista*; data are taken from Sprent (2009) unless noted otherwise. Growth habit and habitat information were either noted by the authors during sample collection in Brazil or were extracted from *Flora e Funga do Brasil* (https://floradobrasil.jbrj.gov.br/FB22876) and/or *Plants of the World*

1137   *Online* (https://powo.science.kew.org/). Nodule anatomy data, including the presence of

1138   either Symbiosome (SYM), Fixation Threads (FT), or Intermediate (FT-SYM) are taken

1139   from the literature, from the study of Sprent et al. (1996), or from the present study.

1140   Taxonomic information with regards to Sections and Subsections were extracted from

1141   Souza et al. (2021).

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6 Concatenated

Fig. 7. *nodC*

FIG 8 A. Phylogeny of the Leguminosae family adapted from LPWG (2017) showing the genera in the Caesalpinioideae subfamily associated with symbionts. **B.** Phylogeny of *Chamaecrista* based on DNA sequences of nuclear ITS and plastidial *trnL-trnF* loci. Majority-rule consensus tree derived from Bayesian analysis; the values of posterior probability (PP) are indicated above the branches in decimal form. Taxonomic groups of associated symbionts are colored. Species indicated by *(*C. chapadae*, *C. zygophylloides*) were not included in the phylogenetic analysis, but based on morphology, it was possible to place them in their appropriate taxonomic group. Arrows indicate the ages of nodes as estimated by Rando et al. (2016). **C.** *C. bahiae*; **D.** *C. desvauxii*; **E.** *C. arrojadoana.* (photos Juliana Rando)