Translationally active soil microorganisms during substrate-induced respiration

Nina Rose Camillone,[*,1,2] **Mary Ann Victoria Bruns,**[1,2,3] **Raúl Román,**[1,2,4] **Daniel Wasner,**[5] **Estelle Couradeau**[1,2,3]

[1]Department of Ecosystem Science and Management, The Pennsylvania State University, University Park, PA, USA

[2]One Health Microbiome Center, Huck Institutes of the Life Sciences, The Pennsylvania State University, University Park, PA, USA

[3]Intercollege Graduate Degree Program in Ecology, The Pennsylvania State University, University Park, PA, USA

[4]Instituto Multidisciplinar para el Estudio del Medio Ramon Margalef, Universidad de Alicante, Alicante, Spain

[5]Department of Environmental Systems Science, ETH Zurich, Zurich, Switzerland

[*]Address correspondence to E. Couradeau, estelle@psu.edu

M. A. Bruns and E. Couradeau are co-senior authors.

Word counts: Abstract 230, Importance 68, Body text 5386

ABSTRACT

Soil microorganisms carry out many processes that are fundamental to soil functions. Among the millions of microbial cells present in a gram of soil, however, less than 2% are commonly estimated to be active at any point in time. Because the respiratory response of a bulk soil to carbon substrate addition would be expected to reflect the number of active cells, we hypothesized a positive correlation between active cells and soil respiration rates during substrate-induced respiration (SIR) assays. To test this, we monitored respiration and active cell counts during 24-h incubations of agricultural soil subsamples after treating with two carbon

substrates or a water-only control. We enumerated active cells with the Bioorthogonal Non-canonical Amino Acid Tagging (BONCAT) method. BONCAT provides a labeled amino acid for active cells to incorporate into newly synthesized proteins, which can then be tagged with a fluorescent dye to enable enumeration by flow cytometry. Both respiration rates and active cell counts increased over time and were positively correlated with each other after 6 h of incubation. After 24 h, increases in active cells were proportionally greater than increases in respiration. Additionally, carbon-amended soils had higher respiration rates than water-only soils with similar active cell counts, suggesting differences in carbon use efficiency. Our study documents for the first time the respiratory response from *in-situ* microbial activation induced by substrate amendment of soil within 6 h, a short enough timescale to exclude most cell replication. This study also demonstrates that the correlation between active cell numbers and respiration is substrate-dependent.

IMPORTANCE

While many critical ecosystem services provided by soil are known to rely on microbial activity, the soil microbial community largely remains a black box. While respiration is a common indicator of bulk soil microbial activity, this study demonstrates that the relationship between respiration and the number of active cells differs based on available carbon substrates. Advancing knowledge in this area will both enable better interpretation of biological soil tests by land managers and inform researchers modeling contributions of soil microbial respiration to global carbon dynamics.

INTRODUCTION

Soil ecosystem services are intrinsically dependent on microbial activity through the many vital processes carried out by soil microbes. These include organic matter decomposition, plant nutrient provisioning, soil aggregate formation, nitrogen fixation, antibiotic production, and more[1], with global implications for nutrient cycling and climate change. Eighty percent of the 3200 Pg carbon in terrestrial ecosystems is stored in soil, with a flux of around 60 Pg per year due to microbial decomposition[2]. Microbial carbon use efficiency (CUE) is considered low when biomass production is accompanied by high carbon mineralization as atmospheric CO_2,

whereas high-CUE indicates more soil carbon retention in biomass per unit respiration.

Variations in CUE significantly impact global carbon cycling[3].

The relationship between the soil microbiome and macroscale nutrient cycles is

complicated by the soil's high percentage of cells that are resting, inactive, or dormant[4, 5].

Since soil microenvironments are typically low in carbon and energy sources, many microbes

remain dormant to survive and reactivate pending resource availability[6]. Early studies

enumerating redox-active soil bacterial cells report active fractions of 5-20% or as low as 1%[7,

4, 8, 9]. More recently, however, other methods have found over 40%[10], 50%[11], or 90%[12]

active bacteria in soil samples. In a laboratory incubation study, transcriptionally active soil

bacterial fraction increased from near zero to 15% after 48 h[13]. This highlights that cell activity

status is dynamic, with a potentially activatable fraction that may be difficult to classify in a

dormant-active binary. Going forward, we define active microbes as those synthesizing new

proteins and dormant microbes as those with no detectable translation to focus on potential for

metabolic activity while excluding residual extracellular enzymes.

In light of the numerous beneficial functions of soil microbes, increased soil microbial

activity is often considered desirable overall. For example, bulk soil respiration is a common

proxy for potential microbial activity used as an indicator of agricultural soil health [14, 15].

Additionally, multiple substrate-induced respiration (MSIR) assays, such as MicroResp$^{©}$,

measure soil microbial responses to diverse C substrates, so that a broader variety of respired

substrates indicates a more active and dynamic community [16, 17]. It is well established that

carbon amendments initially stimulate soil respiration to a greater extent than cell division[18].

Particularly within 6 h, no shifts in community composition were observed to suggest substrate-

specific reproductive selection[19]. However, it remains unknown whether reactivation of

dormant microbes is an important contributor to SIR. Otherwise, bulk soil respiration may be

controlled by a small percentage of already-active cells through fluctuations in their cell-specific

respiration. This would indicate a major limitation for whole-community microbiome analysis in

understanding and predicting C cycling. Further, experimentally and conceptually decoupling

bulk community activity from the number of active microbes allows differentiation of active,

efficient community members from and less efficient community members having similar
respiration rates.

In this study, we investigate whether the SIR response is linked to increases in active cells
relative to the total number of cells in the soil. Specifically, we employed bioorthogonal non-
canonical amino acid tagging (BONCAT) and flow cytometry to enumerate active cells in tandem
with measuring SIR. BONCAT enables the detection of translationally active microbes through
the incorporation of a modified amino acid that is later tagged with a fluorescent dye[13, 20].
We considered that the relationship between bulk respiration and number of active cells may
be dependent on the simplicity of available C substrates. Thus, as in MSIR, we used two carbon
substrate treatments (glucose and galactose) to induce different respiration rates along with a
water-only treatment. We measured activity after four incubation times up to 24 h. Having
simultaneously monitored soil microbial activity at the bulk and cell levels, we calculated
respiration rate per active cell. We hypothesized that variation in respiration rate among
substrate treatments and across time would reflect changes in the number of active cells and
not the number of total cells, suggesting shifts in cellular activity status. This finding would
support the consideration of the fluidity of microbial activity status as a factor in modeling and
managing soil ecosystem processes. Further, if bursts in respiration are strongly linked with
dormant cell activation, respiration tests can be developed as proxies for the potentially active
microbial fraction, facilitating future collection and interpretation of soil microbial data.

MATERIALS AND METHODS

Soil collection and experimental design

All soil samples were collected at the Pennsylvania State University's Russell E. Larson
Agricultural Research Center near Pennsylvania Furnace, PA (40.72°N, -77.92°W), from a long-
term tillage experiment established in 1978[21]. A composite sample of bulk soil was collected
in May 2023 from the top 10 cm of a long-term no-till plot planted with wheat, representing the
Ap horizon of a Hagerstown silt loam (fine, mixed, semiactive, mesic Typic Hapludalfs in USDA
soil taxonomy). The soil was sieved to 2 mm and stored at 4°C for less than 4 months before
experimentation. Gravimetric water content was determined to be 16.6% by oven-drying a

subsample at 105°C for 24 h. Available water holding capacity was determined to be 62.2% by

the funnel method[42]. Incubations of soil subsamples with cycloheximide to inhibit fungi or

chloramphenicol to inhibit bacteria for 6 h showed that respiration was around 60% attributable

to bacteria and 40% to fungi (Supplemental Table 1)[22].

The experimental timeline is summarized in Figure 1. Each experimental replicate

consisted of 1.2 g field-moist soil (1 g oven-dried soil equivalent). Replicates were placed into

12-mL glass tubes sealable via screw-on caps with silicone septa (Exetainer®12ml Vial - Flat

Bottom 738W, Labco Limited, High Wycombe, UK). The septa also allowed needle syringe access

for headspace sampling. For substrate treatments, 100 µL of liquid was added to each tube:

either an aqueous substrate solution containing 3.0 mol L^{-1} carbon (in the form of glucose or

galactose, Sigma-Aldrich, Saint Louis, MO, USA) or ultrapure water. This resulted in a final

concentration equivalent to 30 mg glucose g^{-1} soil water, which is the concentration commonly

used with the MicroResp^{TM} system[16, 17]. The gravimetric water content of the soil after

addition of the substrate treatments was 30%, which was 50% of the available water holding

capacity. The tubes were sealed to limit moisture evaporation and incubated in the dark at 22°C

for 2, 6, 12, or 24 h. The time intervals reflect previous studies where respiration rates in surface

horizon soil microcosms amended with glucose peaked from 5-10 h (depending on glucose

concentration) and returned to near initial rates after 24 h[23]. Galactose was selected as a

substrate treatment because it stimulates a soil respiration rate intermediate to those of

glucose and water-only[16]. Initial pilot experiments included lignin and vanillic acid as

additional substrates, but these treatments resulted in very low BONCAT activity levels (on

average less than 0.5%) compared to the false positive discovery rate (0.2%) even after 24 h

incubation. These substrates were not included, and further research will be necessary to

understand why the BONCAT signal was so low in these treatments. Six replicate tubes were

included per substrate treatment per incubation length.

Activity monitoring: respiration measurement and cell extraction

Two hours prior to the end of the incubation period (i.e. concurrently with the start of

the substrate treatment for the 2-h replicates), the tubes were opened, flushed with ambient

air, and resealed to begin a CO_2 accumulation period of equal length for all replicates. While the

tubes were open, 250 µL of 800 µmol L^{-1} L-homopropargylglycine (HPG), a synthetic amino acid

homologous to methionine was also added to 5 replicates per substrate treatment per

incubation length. The sixth replicate in each group received 250 µL of ultrapure water as a

control for the BONCAT dye signal. As a result, the moisture content of all replicates reached

90% of available water holding capacity. The addition of HPG initiated the first stage of the

BONCAT active cell enumeration method, where translationally active cells incorporate HPG into

their proteins such that they can later be distinguished from HPG-negative inactive cells[13].

FIG 1   Experimental design as a timeline of substrate incubations, headspace

sampling, and cell collection. L-homopropargylglycine (HPG) is a synthetic amino acid

homologous to methionine used to label translationally active cells in the BONCAT workflow.

Two hours later at the conclusion of the incubation period, 2 mL of headspace air was

collected with a needle syringe from each tube and immediately analyzed for CO_2 content using

a LI-7000 CO_2/H_2O Gas Analyzer (LI-COR). A calibration curve based on standards with known

CO_2 concentration was used to determine µmol CO_2 in experimental samples. The CO_2 content

of the ambient air was recorded and later subtracted from the headspace concentrations to

calculate the soil respiration rate. Immediately after headspace sampling, 5 mL 0.02% Tween 20

in phosphate buffer saline (PBS) was added to each replicate, and tubes were shaken

horizontally for 5 min at high speed to extract cells from the soil. At this point, a killed soil

replicate was added to the workflow. This subsample had been autoclaved three times at 1-day

intervals (15 min, 121°C) in order to provide a control for staining with no cells. After shaking,

the tubes were centrifuged for 5 min at 30 x g to separate soil particles without pelleting cells.

The supernatant cell suspensions were stored at -20°C in 1 mL aliquots to which 500 µL 30%

glycerol was added (final concentration 10% glycerol) and were processed within one week.

Three replicates (one each from 6-h glucose, 12-h glucose, and 6-h water) had to be removed

from analysis because respiration rates could not be quantified due to leaky tube seals.

Active microbial cell labeling

The methods for collecting cells and BONCAT staining with a click chemistry reaction

were adapted from previously published methods [13, 24]. After thawing and briefly vortexing

the frozen cell suspensions, they were centrifuged at 50 x g for 5 min to pellet any larger

remaining soil particles. Previous studies have used much higher centrifuge speeds while

maintaining high cell recovery[25], but for this soil we had very low cell yields when higher

speeds were used to pellet soil particles, perhaps due to cell clumping. The supernatant was

recovered and passed through a 35 µm filter. Then, cells were pelleted from the suspension by

centrifuging at 16,000 x g for 5 min. The supernatant was poured off and discarded, and the

cells were resuspended in 70.4 µL 1x PBS in 2-mL snap-cap tubes by briefly vortexing. A dye

mixture was prepared and added to each tube for a final reaction concentration of 5 µmol L^{-1}

amino guanidine hydrochloride, 5 µmol L^{-1} sodium L-ascorbate, 100 µmol L^{-1} copper sulfate

pentahydrate (ThermoFisher Scientific, Eugene, OR, USA), 500 µmol L^{-1} tris-

hydroxypropyltriazolylmethylamine (THPTA), and 5 µmol L^{-1} FAM picolyl-azide dye in a final

volume of 80 µL. Unless otherwise noted, all BONCAT reagents including HPG were purchased

from Click Chemistry Tools (Scottsdale, AZ, USA). The tubes were incubated in the dark for 60

min to permit the click reaction between the azide dye and the HPG incorporated into the

proteins of translationally active cells. Next, samples were washed to remove excess dye by

centrifuging at 16,000 x g for 5 min, resuspending in 1 mL 1x PBS by brief vortexing, and

repeating these steps twice more. The resulting cell suspension was diluted in 1x PBS and

passed through a 35 µm filter (Falcon Round-Bottom Tubes with Cell Strainer Cap, 5 mL). SYTO™

59 Red Fluorescent Nucleic Acid Stain (ThermoFisher Scientific, Invitrogen, Eugene, OR, USA)

was added as a counterstain to distinguish cells from clay particles. Cell suspensions were

incubated in the dark for 15 min.

Flow cytometry

Cell enumeration was performed using the LSR Fortessa flow cytometer (BD Biosciences, Franklin Lakes, NJ, USA) at the Pennsylvania State University Flow Cytometry Core, which is equipped with 405 nm (violet), 488 nm, (blue), 532 nm (green), and 640 nm (red) lasers. The BONCAT dye, FAM-picolyl azide (490 nm excitation, 510 nm emission), was captured in a green channel ("FITC") off the 488 nm blue laser. The DNA counterstain, SYTO59 (622 nm excitation, 645 nm emission), was captured in a red channel ("APC") off the 640 nm red laser. Forward and side light scatter were also collected for all samples. All samples were run on the low flow rate setting (which we measured to be 22.6 µL/min for our samples). Data was collected for 120 s. Knowing the flow rate of the instrument enabled calculation of the original concentration of cells in the soil using the number of cells detected, the volume of extract processed, and the mass of soil extracted.

As a negative control, a "cell extract" from a soil sample that had been autoclaved 3 times at 1-day intervals (killed control) was counterstained with SYTO59 to set the threshold for distinguishing cells from soil mineral particles. Similarly, soil samples incubated with water instead of HPG solution (HPG-negative control) were used to set the threshold for BONCAT positive cells, since they should not retain any of the BONCAT dye.

Data analysis was performed using FlowJo Software (Tree Star, Ashland, OR, USA). First, forward and side scatter fluorescence thresholds were set to constrain the size of particles examined. Secondly, a threshold for SYTO positive (SYTO+) particles, which are assumed to be cells rather than clay particles, was determined based on fluorescence intensity in the APC channel from the killed (autoclaved) control samples counterstained with SYTO59. This step accounts for any potential autofluorescence from or staining of soil particles. Finally, a threshold for BONCAT positive (BONCAT+) cells was determined based on the fluorescence intensity in the FITC channel from the SYTO+ population (cells) in the HPG-negative control. To account for electronic noise from the flow cytometer or any potential retention of dyes on soil particles, the number of SYTO+ events detected in the killed control sample was subtracted from the SYTO+ count in other samples within each batch. Additionally, the number of BONCAT+ events

detected in the HPG-negative control was subtracted from the BONCAT+ population in

experimental samples. Any sample where the number of BONCAT+ events was less than or

equal to the number of BONCAT+ events in the counterstain-only control was recorded to have

zero active cells. Preliminary work using the LIVE/DEAD™ *Bac*Light™ Bacterial Viability and

Counting Kit, for flow cytometry (ThermoFisher Scientific, Invitrogen, Eugene, OR, USA)

detected negligible percentages of dead microbes recovered using our extraction method

(<0.65% on average; see Supplemental Table 2), likely due to our cell extraction method failing

to preserve cells without intact membranes, i.e. those deemed dead by the staining kit used.

Thus, we report total cells as representative of the living community.

Statistical analysis

Statistical analysis was conducted using R version 4.2.1[26], and figures were generated

using R packages ggplot2[27] and ggbreak[28]. Substrate treatments within each incubation

length and incubation lengths within each substrate were compared using one-way ANOVAs.

Incubation length was treated as a categorical variable due to the non-linear relationship of

response variables and time. Tukey HSD post-hoc tests were performed for variables with

significant ANOVAs ($p < 0.05$). The rate of change of numbers of active cells in each substrate

treatment was calculated per unit time as if it were a growth rate, but we do not refer to this as

a "growth rate" since new active cells could be the result either of cell division or activation of

dormant cells. Instead, this variable will be referred to simply as the rate of change of numbers

of active cells. The cell-specific respiration rate was calculated by dividing the bulk respiration

rate by the number of active cells in the same experimental replicate as an indicator of

resource-use efficiency. The relationship between respiration rate and log-transformed number

of active cells was analyzed using linear regression.

RESULTS

Cell counts and bulk respiration

241 The total number of microbial cells ranged from 3.1×10^5 to $2.2 \times 10^7\,g^{-1}$ soil throughout

242 the experiment (Figure **2**A). The number of active cells ranged from 1.7×10^3 to $4.4 \times 10^6\,g^{-1}$ soil

243 (bars in Figure **2**B, left y-axis), or 0.1% to 27% of the total (percentages noted below data bars in

Figure 2A). Generally, the number and percentage of active cells increased with time for all

treatments. For the glucose treatment, numbers of total and active cells were highest after 12 h

of incubation. In galactose, total and active cells were highest at 24 h. For the water treatment,

total cells were highest at 12 h while active cells were highest at 24 h. Among the treatments,

the glucose treatment had the highest total number of cells throughout the experiment while

the total cells in the galactose and water treatments never significantly differed from each

other. As for active cells, they likewise were highest in glucose compared to the other

treatments at 6 h, 12 h, and 24 h. In water, active cell numbers averaged slightly lower than in

galactose at 6 h and 24 h, but they were significantly higher than in galactose at 12 h.

Contrastingly, galactose had the most active cells at 2 h, although it was still low at 2% of total

cells.

The bulk respiration rate increased after 24 h in all 3 treatments (lines in Figure 2B, 267

right y-axis). The respiration rate of the substrate-amended treatments did not significantly

differ from their initial rates until 24 h. Glucose consistently had a higher respiration rate than

galactose, and galactose consistently had a higher respiration rate than water-only.

FIG 2   Total cells, active cells, and respiration rate. (A) Total cells over time for

different substrate treatments, reported in cells per g dry soil. The lower (orange) portion of

each bar represents the active cell fraction. Numbers below each bar indicate the

percentage of active cells. (B) Active cells (bars, left y-axis) and respiration rate (lines, right y-

axis). Note that 0 for the right y-axis is elevated. For both (A) and (B), uppercase letters

represent statistically significant differences found by Tukey's HSD test ($p < 0.05$) among

different substrate treatments within the same time point, indicated by the x axis.

Lowercase letters represent statistically significant differences across time points within the

same substrate treatment, indicated by color. The absence of a letter of a particular case

above a bar indicates that Tukey's test was not conducted due to an insignificant ANOVA

result. Bars indicate standard error.

The largest absolute increases in numbers of total and active cells coincided with each

other for all three treatments. In glucose and water, they were between 6 and 12 h, but in

galactose they were later, between 12 and 24 h. For glucose, the largest increase in respiration

rate occurred between 12 and 24 h, after the large increase in number of active cells between 6

and 12 h. For galactose, the largest changes in number of active cells and respiration coincided

between 12 and 24 h. For water, the largest increase in respiration occurred first, between 2 and

6 h, before the increase in active cells at 12 h. Glucose had the largest changes in total and

active cells and in respiration rate compared to the other treatments.

Cell-specific respiration rate

Cell-specific respiration was calculated for each replicate by dividing bulk respiration by

the number of active cells. In all three treatments, the cell-specific respiration rate was highest

after 2 h and decreased thereafter. However, this decrease was only statistically significant for

glucose (Figure 3). At 2 h, the glucose treatment had the highest cell-specific respiration, driven

by low active cell counts and a high respiration rate. At 12 h, cell-specific respiration was highest

for galactose, caused by lower active cell counts than glucose and higher respiration rates than

water. After 6 h and after 24 h, there were no statistically significant differences in cell-specific

respiration rate among treatments.

FIG 3   Cell-specific respiration rate, calculated by dividing bulk respiration by the number of

active cells. Uppercase letters represent statistically significant differences found by Tukey's

HSD test ($p<0.05$) among substrate treatments within the same time point, indicated by the x

axis. Lowercase letters represent statistically significant differences across time points within

the same substrate treatment, indicated by color. The absence of a letter of a particular case

above a bar indicates that Tukey's test was not conducted due to an insignificant ANOVA

result. Bars indicate standard error.

DISCUSSION

Methodological insights and comparisons

Our study documents for the first time *in-situ* soil microbial cell activation induced by

substrate amendment within 6 h, a short enough timescale to exclude most cell replication. This

is possible using the BONCAT method because it detects protein translation rather than cell

growth, which has previously been used to monitor soil microbial activity via methods such as

SIP[29]. A longer activity labeling period can increase the detection of low-activity cells but does

not give a snapshot for time-resolved changes in cell activity status. Other methods for

enumeration of active soil microbes involve dyeing active cells post extraction[5, 9, 10, 11],

whereas in BONCAT active cells are tagged *in situ* with a synthetic amino acid prior to extraction

and subsequently dyed through a click-chemistry reaction. This could lead to discrepancies

between detected and actual activity if dormant cells are reawakened or if activity is suppressed

through the disturbance of extraction, e.g. aerobic respiration decreasing upon saturation.

Differences among soil active microbes labeling methods are summarized in Table **1**.

Comparative studies are needed to determine if variation in reported numbers of active cells

reflects actual cell physiology (e.g. some cells contain active enzymes but are not currently

translating new ones), natural differences in soil samples across time and space, or inefficiencies

in some methods compared to others.

Some remaining methodological challenges in active cell enumeration are specific to

BONCAT, while others apply more broadly. For example, BONCAT does not detect the activity of

microbes with access to native methionine (outcompeting analog HPG) or microbes with

adequate protein reserves without need for translation. Regardless of staining method,

extraction cannot capture all microbes present in the soil, especially filamentous microbes such

as fungi. Typically, soils contain 10^7-10^{10} prokaryotic cells per gram[30], compared to our counts

of 5×10^6 after 2 h of incubation (Figure **2**A). We confirmed that our cell extracts had very low

numbers of dead microbes (Supplemental Figure 1), which may contribute to our lower cell

counts considering previous research reporting 40% dead microbial cells in soil[7]. We assumed

that our percentages of active cells were representative of the whole live community. In the

end, any binary divide between active and inactive cells is inherently a simplification, yet it can

still yield insights into overall community status for the purpose of this discussion.

TABLE 1 Comparison of methods to enumerate active microbial enumeration in soil.

Abreviations: 5-cyano-2,3-ditolyl tetrazolium chloride (CTC, an artificial electron acceptor),

5(6)-carboxyfluorescein diacetate (CFDA, a substrate hydrolyzed by intracellular esterases),

fluorescence *in situ* hybridization (FISH), stable-isotope probing (SIP).

Method	CTC	CFDA	FISH	DNA-SIP	BONCAT
Activity type	Aerobic respiration (redox)	Metabolic enzymes	rRNA content	DNA replication	Protein translation
Time scale	30+ min	30+ min	Fixed cells	1+ days	30+ min
Labeling location	Extracted cells	Extracted cells	Extracted cells	*in situ*	*in situ*
Reported active fraction	2-6%[9] <1%[5]	42%[10]	55%[11]	91%[12]	25%[13]

Glucose amendment reactivated dormant cells

Active cell numbers increased most rapidly from 2-6 h of incubation with glucose (Figure

4). Notably, this was our only observation of an increase in total cells that was outpaced by a

larger increase in active cells. In other words, the number of dormant cells (estimated by

subtracting active from the total) decreased, suggesting cell activation. Alternatively, or in

addition, this could indicate a death rate among inactive cells outpaced by reproduction among

active cells; however, this seems less likely considering known soil microbial growth rates

(depicted in Figure 4 as rate of change per h). Assuming no growth or reactivation of dormant

cells, active cells would have been doubling every 37 min to achieve the observed increase

between 2-6 h for the glucose treatment. This approaches the theoretical maximum growth rate

under optimum soil conditions with as a generation time of 0.5 h[31]. For comparison,

Eschedrichia coli has a generation time around 0.3 h *in vitro*[32]. In actuality, soil microbes have

been clocked with doubling times only as low as 7.2 h in the rhizosphere[33] and 14 to 45 days

in bulk soil[34] (depicted in Figure 4 as rate of change per h). This suggests that reactivation of

dormant cells largely contributed to the increase in active cells. With the same reasoning of a

high rate of increase in active cells, dormant microbes were likely reactivating in the water

treatment between 2-12 h (Figure 4). The rate of change in active cell counts in the galactose

treatment did not measurably change throughout the experiment and remained comparable to

a highly active rhizosphere environment. The growth rates for total cells were generally much

lower than active cells' (Supplemental Figure 1).

FIG 4   Rate of change in number of active cells over time. Bars indicate standard error,

propagated from the standard error in the enumerated community sizes. For comparison,

dashed lines indicate several known growth rates: a laboratory culture of *Escherichia coli*

(generation time 0.3 h)[32], an active rhizosphere community (generation time 7.2 h)[33],

and bulk forest soil (generation time 14.1 days) [34]. Note the break in the y axis between

2.5 and 7.

Cell-specific activity differed by treatment

As hypothesized, respiration rates and active cell counts were positively correlated,

although the relationship differed by substrate. Regression analysis revealed statistically

significant relationships between these variables within each treatment and across all

treatments together, although in the glucose treatment it was only significant at the p<0.06

level (equations reported in Figure 5A). Differences among the regression models' slopes or

intercepts result in the regression line for glucose being higher than that of galactose, which is

in turn higher than water's. This pattern reflects overall trends in the treatments' cell-specific

respiration across the experiment. Lower respiration and active cell counts in the galactose

treatment than the glucose treatment are consistent with the ability of cells to utilize glucose

directly. By contrast, galactose requires multiple enzyme-catalyzed reactions before a

downstream metabolite enters the glycolytic pathway [35].

Without any carbon addition, the water-only treatment has the lowest regression

coefficient (Figure 5A). This indicates that this moist soil environment supported an expanding

active community with minimal changes in respiration rate. This finding is consistent with the

Microbial Efficiency-Matrix Stabilization (MEMS) framework, which theorizes that respiration is

highest where inputs are low quality (excess ratio of C to other nutrients) and have a low

capacity to be stabilized in the mineral soil matrix [36], e.g. glucose supports lower CUE than

lignin. Active microbes in the water-only, carbon-poor treatment may use carbon more

efficiently, i.e. prioritize its assimilation as biomass relative to mineralization. This is also

consistent with previous studies that found that soil microbes metabolized complex substrates

with a higher carbon use efficiency[37]. (also check decreased CUE with excess C in [38])

A

B

380   Active microbial community (log cells/g soil)   B

FIG 5   Regressions of bulk respiration onto active cell counts in log scale. The shaded regions show the 95% confidence region for each regression line. (A) Regression lines for all data and for each substrate individually. P-values for coefficients and intercepts respectively are <0.001 and 0.08 for all data, 0.06 and 0.9 for glucose, <0.0001 and 0.009 for galactose, and 0.03 and 0.9 for water. (B) Regression lines for each time point individually. P-values for coefficients and intercepts respectively are 0.9 and 0.6 for 2 h, 0.03 and 0.2 for 6 h, 0.08 and 0.2 for 12 h, and 0.003 and 0.008 for 24 h.

Lag between respiration response and increase in active cells

Within 2 h of substrate addition, the number of active cells did not differ among treatments (Figure **2**), while the two carbon-amended treatments already had much higher bulk

respiration rates than water-only (Figure 2B). Thus, our results indicate an initial lag where

changes in translational activity are decoupled from the respiration response. As hypothesized,

we observed that active cell counts and respiration rates followed similar patterns across

treatments when cell-specific respiration was consistent, i.e. after 6 and 24 h of substrate

incubation (Figure 3). Regression models calculated within each time point also indicated

positive correlations between respiration and active cell counts with statistically significant

regression coefficients at both 6 and 24 h, but not at 2 or 12 h (Figure 5B). At these time points,

by contrast cell-specific respiration differed among treatments, reflecting weak or no correlation

between respiration and number of active cells. The glucose treatment had the highest cell-

specific respiration at 2 h, and galactose had the highest at 12 h (Figure 3). These differences

suggest that glucose-amended soil microbes began rapidly metabolizing with a burst of

respiration per cell within 2 h before the activation of dormant cells by 6 h. Meanwhile,

galactose induced increases in respiration and active cell counts at a slower but steadier rate. In

fact, the galactose treatment exhibited the least variation in cell-specific respiration and the

strongest positive correlation (highest regression R^2) between bulk respiration and active cell

counts (Figure 5A). A low level of respiration from BONCAT-negative cells also would have

contributed to low cell-specific respiration and a strong correlation between respiration and

active cells. This could be the case if few cells already contained the enzymes necessary for

galactose metabolism. Regardless, for all substrate treatments, cell-specific respiration tended

to decrease over time, suggesting an initial dip in CUE with excess carbon and a lag in cell

activation post substrate addition. In the field, this could contribute to increased carbon

emissions from more frequently disturbed soils.

Implications for soil testing

Our results demonstrate the utility of SIR as a proxy for soil microbial activation in an

agricultural soil. Soil testing already utilizes SIR for crop yield prediction[39, 40]. Thus, future

research in more soils establishing a correlation between SIR and numbers of active microbes

would also implicate a correlation with agricultural yields. This may seem intuitive, considering

the microbial basis for many critical agricultural soil functions such as decomposition and

aggregate formation. However, other factors complicate the relationship between microbial

activity and optimal soil functionality. Frequent tillage disturbance, for example, could over-

stimulate microbial activity and cause faster depletion of soil organic matter with time.

Additionally, CUE can affect the relationship between respiration and microbial functionality. A

high respiration rate in an SIR test can indicate a responsive microbial community or simply a

less efficient one.

However, microbial researchers have described activity status not simply as a dichotomy

between dead and alive or between active and dormant, but as a spectrum of activity levels,

defining potentially active microbes as those that resume activity after 4-12 h of stimulation [7].

While a large active microbial fraction could achieve a high respiration rate and rapidly deplete

soil carbon stores, a large *potentially* active fraction would be a better indicator of a responsive

and resilient microbiome supporting soil functions. The present study found rapid increases in

active cell counts but not total cell counts for 6 h of substrate incubation. Previous MSIR studies

have found 6-h incubation time to be optimal for observing the existing community while

limiting time for shifts in community composition[16, 17, 19], which our study corroborates.

Additionally, our results showed a similar effect of substrate treatment on respiration rate and

active cell counts at this time (Figure **2**B), and the earliest correlation between these two

indicators (Figure **5**B). These results support the interpretation of SIR tests as measures of

potentially active microbes responsive to different substrates.

At the same time, active cell numbers alone could not consistently predict respiration

due to differences among substrate treatments. Both carbon substrates induced higher cell-

specific respiration than the water-only treatment at different times (Figure **3**), and carbon-

amended replicates had higher respiration than the water replicates with similar numbers of

active cells (Figure **5**A). Previously, a field study found that tilled soils with incorporation of

organic residues had higher respiration rates with lower microbial biomass compared to no-till

soils[41], similarly pointing toward higher activity per cell with access to carbon.

Overall, this project demonstrates a relationship between respiration and numbers of

active microbes. This provides a basis for interpreting respiration-based soil tests in microbial

terms and targeting research efforts on active microbes due to their contribution to essential

soil functions. Future research coupling BONCAT with cell sorting and metagenomic sequencing can reveal which microbes are activated under different soil conditions and amendments. Linking other bulk soil tests to active cell counts can further inform soil management decisions for promoting desired outcomes of microbial activity.

ACKNOWLEDGMENTS

We acknowledge that gas analysis for CO_2 content was conducted in the lab of Jason Kaye with training and support from Brosi Bradley. We would also like to acknowledge that cell enumerations were conducted at the Huck Institutes' Flow Cytometry Core Facility (RRID:SCR_024460) on the BD Fortessa flow cytometer with training and support from multiple staff members.

DATA AVAILABILITY STATEMENT

Original flow cytometry data are available as FCS files at doi.org/10.26208/G8KX-GG32 through Penn State's DataCommons.

FUNDING

This work was funded through a Flower Grant awarded by the Ecology Institute at the Penn State Huck Institutes of Life Sciences to authors Bruns, Camillone, and Couradeau. This work was also supported by the USDA National Institute of Food and Agriculture and Hatch Appropriations under Project #PEN04949 and Accession #7006508.

CONFLICTSOFINTEREST

The authors declare no conflict of interest.

REFERENCES

References

[1] **Hartmann M, Six J**. Nov 2022. Soil structure and microbiome functions in agroecosystems. *Nature Reviews Earth & Environment* 4 (1):4–18. doi: 10.1038/s43017-022-00366-w.

[2] **Lal R**. 2008. Sequestration of atmospheric CO2 in global carbon pools. *Energy &*

Environmental Science 1 (1):86. doi:10.1039/b809492f.

[3] **Tao F, Huang Y, Hungate BA, Manzoni S, Frey SD, Schmidt MWI, Reichstein M,**

Carvalhais N, Ciais P, Jiang L, Lehmann J, Wang YP, Houlton BZ, Ahrens B, Mishra

U, Hugelius G, Hocking TD, Lu X, Shi Z, Viatkin K, Vargas R, Yigini Y, Omuto C,

Malik AA, Peralta G, Cuevas-Corona R, Di Paolo LE, Luotto I, Liao C, Liang YS,

Saynes VS, Huang X, Luo Y. May 2023. Microbial carbon use efficiency promotes global soil

carbon storage. *Nature* 618 (7967):981–985. doi:10.1038/s41586-023-06042-3.

[4] **Lennon JT, Jones SE**. 2 2011. Microbial seed banks: the ecological and evolutionary

implications of dormancy. *Nature Reviews Microbiology* 9:119–130. doi: 10.1038/nrmicro2504.

[5] **Salazar A, Lennon JT, Dukes JS**. May 2019. Microbial dormancy improves predictability of

soil respiration at the seasonal time scale. *Biogeochemistry* 144(1):103–116.

doi:10.1007/s10533-019-00574-5.

[6] **Hobbie JE, Hobbie EA**. 2013. Microbes in nature are limited by carbon and energy: the

starving-survival lifestyle in soil and consequences for estimating microbial rates. *Frontiers in*

Microbiology 4. doi:10.3389/fmicb.2013.00324.

[7] **Blagodatskaya E, Kuzyakov Y**. 12 2013. Active microorganisms in soil: Critical review of

estimation criteria and approaches. *Soil Biology and Biochemistry* 67:192–211. doi:

10.1016/j.soilbio.2013.08.024.

[8] **Wang G, Mayes MA, Gu L, Schadt CW**. Feb 2014. Representation of Dormant and Active

Microbial Dynamics for Ecosystem Modeling. *PLoS ONE* 9 (2):e89252.

doi:10.1371/journal.pone.0089252.

[9] **Winding A, Binnerup SJ, Sørensen J**. Aug 1994. Viability of Indigenous Soil Bacteria Assayed

by Respiratory Activity and Growth. *Applied and Environmental Microbiology* 60 (8):2869–2875.

doi:10.1128/aem.60.8.2869-2875.1994.

[10] **Espina L**. Aug 2020. An approach to increase the success rate of cultivation of soil bacteria

based on fluorescence-activated cell sorting. *PLOS ONE* 15 (8):e0237748.

doi:10.1371/journal.pone.0237748.

[11] **Bertaux J, Gloger U, Schmid M, Hartmann A, Scheu S**. Jun 2007. Routine fluorescence

in situ hybridization in soil. *Journal of Microbiological Methods* 69 (3):451–460.

doi:10.1016/j.mimet.2007.02.012.

[12] **Ding C, Xu X, Liu Y, Huang X, Xi M, Liu H, Deyett E, Dumont MG, Di H, Hernández M, Xu J,**

Li Y. Mar 2023. Diversity and assembly of active bacteria and their potential function along soil

aggregates in a paddy field. *Science of The Total Environment* 866:161360.

doi:10.1016/j.scitotenv.2022.161360.

[13] **Couradeau E, Sasse J, Goudeau D, Nath N, Hazen TC, Bowen BP, Chakraborty R,**

Malmstrom RR, Northen TR. 2019. Probing the active fraction of soil microbiomes using

BONCAT-FACS. *Nature Communications* 10:2770. doi:10.1038/s41467-019-10542-0.

[14] **Rieke EL, Cappellazzi SB, Cope M, Liptzin D, Bean GM, Greub KLH, Norris CE, Tracy PW,**

Aberle E, Ashworth A, Tavarez OB, Bary AI, Baumhardt RL, Gracia AB, Brainard DC, Brennan JR,

Reyes DB, Bruhjell D, Carlyle CN, Crawford JJW, Creech CF, Culman SW, Deen B, Dell CJ, Derner

JD, Ducey TF, Duiker SW, Dyck MF, Ellert BH, Solorio AE, Fonte SJ, Fonteyne S, Fortuna AM,

Foster JL, Fultz LM, Gamble AV, Geddes CM, Griffin-LaHue D, Grove JH, Hamilton SK, Hao X,

Hayden ZD, Honsdorf N, Howe JA, Ippolito JA, Johnson GA, Kautz MA, Kitchen NR, Kumar S,

Kurtz KSM, Larney FJ, Lewis KL, Liebman M, Ramirez AL, Machado S, Maharjan B, Gamiño

MAM, May WE, McClaran MP, McDaniel MD, Millar N, Mitchell JP, Moore AD, Moore PA,

Gutiérrez MM, Nelson KA, Omondi EC, Osborne SL, Alcalá LO, Owens P, Pena-Yewtukhiw EM,

Poffenbarger HJ, Lira BP, Reeve JR, Reinbott TM, Reiter MS, Ritchey EL, Roozeboom KL, Rui Y,

Sadeghpour A, Sainju UM, Sanford GR, Schillinger WF, Schindelbeck RR, Schipanski ME,

Schlegel AJ, Scow KM, Sherrod LA, Shober AL, Sidhu SS, Moya ES, Luce MS, Strock JS, Suyker

AE, Sykes VR, Tao H, Campos AT, Eerd LLV, Verhulst N, Vyn TJ, Wang Y, Watts DB, William BB,

Wright DL, Zhang T, Morgan CLS, Honeycutt CW. May 2022. Linking soil microbial community

structure to potential carbon mineralization: A continental scale assessment of reduced tillage.

Soil Biology and Biochemistry 168:108618. doi:10.1016/j.soilbio.2022.108618.

[15] **Franzluebbers A**. Jul 1999. Potential C and N mineralization and microbial biomass from

intact and increasingly disturbed soils of varying texture. *Soil Biology and Biochemistry* 31

(8):1083–1090. doi:10.1016/s0038-0717(99)00022-x.

[16] **Creamer RE, Stone D, Berry P, Kuiper I**. 2016. Measuring respiration profiles of soil microbial

communities across Europe using MicroResp™ method. *Applied Soil Ecology* 97:36–43.

doi:10.1016/j.apsoil.2015.08.004.

[17] **Campbell CD, Chapman SJ, Cameron CM, Davidson MS, Potts JM**. Jun 2003. A

Rapid Microtiter Plate Method To Measure Carbon Dioxide Evolved from Carbon Substrate

Amendments so as To Determine the Physiological Profiles of Soil Microbial Communities by

Using Whole Soil. *Applied and Environmental Microbiology* 69 (6):3593–3599.

doi:10.1128/aem.69.6.3593-3599.2003.

[18] **NGUYEN TT, MARSCHNER P**. Feb 2017. Soil Respiration, Microbial Biomass and Nutrient

Availability in Soil After Addition of Residues with Adjusted N and P Concentrations.

Pedosphere 27 (1):76–85. doi:10.1016/s1002-0160(17)60297-2.

[19] **Gao F, Fan H, Chapman SJ, Yao H**. 4 2022. Changes in soil microbial community

activity and composition following substrate amendment within the MicroResp™ system.

Journal of Soils and Sediments 22:1242–1251. doi:10.1007/s11368-022-03143-w.

[20] **Hatzenpichler R, Connon SA, Goudeau D, Malmstrom RR, Woyke T, Orphan VJ**.

Jun 2016. Visualizing in situ translational activity for identifying and sorting slow-growing

archaeal-bacterial consortia. *Proceedings of the National Academy of Sciences* 113 (28).

doi:10.1073/pnas.1603757113.

[21] **Duiker SW, Beegle DB**. Jul 2006. Soil fertility distributions in long-term no-till, chisel/disk

and moldboard plow/disk systems. *Soil and Tillage Research* 88 (1-2):30–41.

doi:10.1016/j.still.2005.04.004.

[22] **Anderson JPE, Domsch KH**. Mar 1975. Measurement of bacterial and fungal contributions

to respiration of selected agricultural and forest soils. *Canadian Journal of Microbiology* 21

(3):314–322. doi:10.1139/m75-045.

[23] **Drobník J**. 4 1960. Primary oxidation of organic matter in the soil. *Plant and Soil* 12:199–

211. doi:10.1007/BF01343649.

[24] **Reichart NJ, Jay ZJ, Krukenberg V, Parker AE, Spietz RL, Hatzenpichler R**. 11 2020. Activity-

based cell sorting reveals responses of uncultured archaea and bacteria to substrate

amendment. *ISME Journal* 14:2851–2861. doi:10.1038/s41396-020-00749-1.

[25] **Khalili B, Weihe C, Kimball S, Schmidt KT, Martiny JBH**. Oct 2019. Optimization of a

Method To Quantify Soil Bacterial Abundance by Flow Cytometry. *mSphere* 4 (5).

doi:10.1128/msphere.00435-19.

[26] **R Core Team**. 2022. R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

[27] **Wickham H**. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

https://ggplot2.tidyverse.org.

[28] **Xu S, Chen M, Feng T, Zhan L, Zhou L, Yu G**. Nov 2021. Use ggbreak to Effectively

Utilize Plotting Space to Deal With Large Datasets and Outliers. *Frontiers in Genetics* 12.

doi:10.3389/fgene.2021.774846.

[29] **Papp K, Hungate BA, Schwartz E**. 1 2020. Glucose triggers strong taxon-specific responses

in microbial growth and activity: insights from DNA and RNA qSIP. *Ecology* 101:e02887.

doi:10.1002/ECY.2887.

[30] **Frossard A, Hammes F, Gessner MO**. Jun 2016. Flow Cytometric Assessment of Bacterial

Abundance in Soils, Sediments and Sludge. *Frontiers in Microbiology* 7.

doi:10.3389/fmicb.2016.00903.

[31] **Borer B, Or D**. Feb 2022. Bacterial age distribution in soil – Generational gaps in

adjacent hot and cold spots. *PLOS Computational Biology* 18 (2):e1009857.

doi:10.1371/journal.pcbi.1009857.

[32] **Tuttle AR, Trahan ND, Son MS**. Jan 2021. Growth and Maintenance of *Escherichia coli*

Laboratory Strains. *Current Protocols* 1 (1). doi:10.1002/cpz1.20.

[33] **Bowen GD, Rovira AD**. Sep 1976. Microbial Colonization of Plant Roots. *Annual Review of*

Phytopathology 14 (1):121–144. doi:10.1146/annurev.py.14.090176.001005.

[34] **Caro TA, McFarlin J, Jech S, Fierer N, Kopf S**. Apr 2023. Hydrogen stable isotope probing of

lipids demonstrates slow rates of microbial growth in soil. *Proceedings of the National Academy*

of Sciences 120 (16). doi:10.1073/pnas.2211625120.

[35] **Wang H, Sun T, Zhao Z, Gu S, Liu Q, Wu T, Wang D, Tian C, Li J**. Apr 2021. Transcriptional

Profiling of Myceliophthora thermophila on Galactose and Metabolic 580 Engineering for

Improved Galactose Utilization. *Frontiers in Microbiology* 12. doi:10.3389/fmicb.2021.664011.

[36] **Cotrufo MF, Wallenstein MD, Boot CM, Denef K, Paul E**. Feb 2013. The Microbial Efficiency-

Matrix Stabilization (MEMS) framework integrates plant litter decomposition with soil organic

matter stabilization: do labile plant inputs form stable soil organic matter? *Global Change*

Biology 19 (4):988–995. doi:10.1111/gcb.12113.

[37] **Bölscher T, Wadsö L, Börjesson G, Herrmann AM**. Feb 2016. Differences in substrate use

efficiency: impacts of microbial community composition, land use management, and substrate

complexity. *Biology and Fertility of Soils* 52 (4):547–559. doi:10.1007/s00374-016-1097-5.

[38] **Manzoni S, Taylor P, Richter A, Porporato A, Ågren GI**. Jul 2012. Environmental and

stoichiometric controls on microbial carbon-use efficiency in soils. *New Phytologist* 196 (1):79–

91. doi:10.1111/j.1469-8137.2012.04225.x.

[39] **Franzluebbers AJ**. Jan 2016. Should Soil Testing Services Measure Soil Biological Activity?

Agricultural I& Environmental Letters 1 (1). doi:10.2134/ael2015.11.0009.

[40] **Stenberg B**. May 1998. Soil attributes as predictors of crop production under

standardized conditions. *Biology and Fertility of Soils* 27 (1):104–112.

doi:10.1007/s003740050407.

[41] **Mangalassery S, Mooney SJ, Sparkes DL, Fraser WT, Sjögersten S**. May 2015. Impacts of

zero tillage on soil enzyme activities, microbial characteristics and organic matter functional

chemistry in temperate soils. *European Journal of Soil Biology* 68:9–17.

doi:10.1016/j.ejsobi.2015.03.001.

[42] **Nelson, JT, Adjuik, TA, Moore, EB, VanLoocke, AD, Ramirez Reyes, A, & McDaniel, MD.** Dec

2023. A Simple, Affordable, Do-It-Yourself Method for Measuring Soil Maximum Water Holding

Capacity. *Communications in Soil Science and Plant Analysis* 55(8):1190–1204.

doi:10.1080/00103624.2023.2296988.