SpottedPy quantifies relationships between spatial transcriptomic hotspots and uncovers new environmental cues of epithelial-mesenchymal plasticity in cancer

Eloise Withnell[1], Maria Secrier[1,*]

[1]UCL Genetics Institute, Department of Genetics, Evolution and Environment, University College London, London, WC1E 6BT, UK

* To whom correspondence should be addressed (m.secrier@ucl.ac.uk)

Keywords: spatial transcriptomics, epithelial-to-mesenchymal transition, tumour microenvironment, cell plasticity, hotspot analysis

ABSTRACT

Spatial transcriptomics is revolutionising our ability to explore intratissue heterogeneity in cancer, but methods that can effectively capture cancer cell niches and explore their relationships with the tumour microenvironment at various spatial scales remain limited. Here we present SpottedPy, a Python package designed to identify tumour hotspots and map spatial interactions within the cancer ecosystem. We employ SpottedPy to examine epithelial-mesenchymal plasticity in breast cancer and highlight locally stable niches associated with angiogenic and hypoxic regions, and shielded by myCAFs, macrophages and perivascular cell populations. Hybrid and mesenchymal hotspot distribution followed transformation gradients within the tissue reflecting progressive immunosuppression. Our method offers the flexibility to explore spatial relationships at different scales, from immediate neighbours to broader tissue modules, providing new insights into the spatial dynamics of the tumour microenvironment.

BACKGROUND

The spatial organisation of cancer cells and their interactions with immune and stromal cells in their environment are instrumental in dictating tumour behaviour and progression[1]. Recent developments in sequencing technologies that are able to profile large areas of the entire tissue in minute detail, such as spatial transcriptomics[2,3], are increasingly enabling us to gain a more comprehensive understanding of the complex ecosystem of tumour microenvironments (TMEs)[4,5]. Profiling the cancer tissue spatially allows us to explore the tumour architecture and its heterogeneity in detail, paving the way to deciphering the crosstalk between tumour cells and their surroundings and opening up new therapeutic opportunities[6,7].

Several studies to date have demonstrated the advantages of spatial transcriptomics in delineating major tissue domains with distinct cell composition[8], cancer hallmarks[9], immunosuppressive hubs[10,11], entire tumour ecotypes with divergent clinical outcomes[12] or the impact of specific drugs on inhibiting tumour progression[13]. However, focusing on areas of the tissue that are relevant for a specific biological question and surveying relationships between cell populations at the right scale remains a challenge in these datasets. To delineate biologically meaningful tissue areas using spatial transcriptomics, some of the current analytical methods, such as SpaGCN[14] and BayesSpace[15] focus on unsupervised clustering of gene expression. Approaches like NeST[16] or GASTON[17] take this one step further and incorporate a nested structure or topography metrics to outline hierarchically organised co-expression hotspots aligning with tissue histology. Given that similar cells often cluster together[18,19], methods that can reliably detect statistically significant clusters are important in reinforcing the accuracy of cell states determined from continuous signatures. CellCharter[20] capitalises on this concept and uses Gaussian mixture models to identify stable clusters representing spatial niches exhibiting distinct shapes and cell plasticity. More targeted clustering approaches employing user-defined signatures or cell types have recently been implemented in Voyager[21] and Monkeybread[22]. Such methods can play a significant role in enhancing the interpretation of cell types that are inferred through spatial transcriptomic deconvolution techniques, but flexibly exploring spatial units at different scales remains difficult.

When it comes to assessing the spatial proximity of different clusters, methods that infer this through co-enrichment within the immediate neighbourhood have been implemented in packages like Squidpy[23]. However, these approaches lack analytical methods to define and compare distances between specific areas or cell populations of interest that might exert influence on each other across different ranges. Given that the scale of certain biological processes in cancer, such as hypoxia, remains elusive, relying solely on conventional spot neighbourhood-centric methods might obscure complex, spatial interactions. This is known as the Modifiable Areal Unit Problem (MAUP) in geostatistics[24], where spatial data patterns are observed to shift contingent on the size and shape of the spatial analysis units. Whilst a growing number of methods address the need for multi-scale analysis [25,23,26,27], the effect of changing spatial units has generally been underexplored in spatial biology[18].

Here, we build on the use of key ideas in the geostatistics field within spatial biology as previously demonstrated by Voyager[21] to devise an analytical method tailored to interrogate spatial

relationships at various scales within 10X Visium transcriptomic datasets. Our approach defines
areas densely inhabited by particular cell types or marked by user-defined gene signatures
(hotspots) and areas depleted of cell types or gene signatures (coldspots), and statistically
evaluates the proximity of such areas to other predefined hotspots or coldspots. We additionally
compare the spatial relationships detected using the hotspot approach to those observed when
looking only at the immediate neighbourhood of individual spots. We assess how the relationships
between these variables change when varying the hotspot size or the neighbourhood size
surrounding a spot. We have implemented this method in the Python package SpottedPy, available
at https://github.com/secrierlab/SpottedPy.
84
To highlight the potential of our method, we focus on a key process underlying cancer progression,
the epithelial-to-mesenchymal transition (EMT). During EMT, polarised epithelial cells undergo
multiple molecular changes and lose their identity to acquire a mesenchymal phenotype[28]. The
interplay between EMT and the tumour microenvironment (TME) is multifaceted: while the TME is
believed to be an inducer of EMT, mesenchymal tumour cells potentially influence the TME[29,30]. This
dynamic is further complicated by the nature of EMT, which is not merely a dichotomous event.
Current research suggests that EMT is a spectrum, varying from a continuous gradient to distinct,
discrete stages[31,32]. Depending on the context, cells undergoing this transition can be locked in an
EMT state, or alternate between a large landscape of EMT states, a phenomenon called epithelial-
mesenchymal plasticity (EMP)[33]. We have previously shown that the tumour spatial organisation
follows EMP gradients and that hybrid and mesenchymal cells establish distinct interactions with the
TME in small datasets of spatially-profiled breast tumours[34]. However, the spatial organisation and
interactions of cells across this EMP spectrum within the tumour milieu remain largely undefined at
a larger scale and could offer great potential in developing therapies that exploit cell intrinsic or
microenvironmental vulnerabilities linked with this process. Here, we showcase the capability of
SpottedPy to unveil new relationships between EMT hotspots and the TME in breast cancer spatial
transcriptomics data. By facilitating multi-scale comparisons of tumour and TME relationships, we
yield rigorous evidence of spatial dynamics and offer an interpretable and intuitive measure of
interactions between tumour cells and their environment at flexible scales.
104
105
106 **RESULTS**
107
108 **SpottedPy: a tool to investigate biological modules and spatial relationships at different**
109 **scales**
110
From direct cell-cell interactions to immediate neighbourhoods and even across larger modules,
cancer cell evolution is impacted at different scales by its environment (**Fig. 1a**). However, exploring
this landscape flexibly and determining the areas within the tissue where these effects are most
prominent is not straightforward. Whilst neighbourhood enrichment is commonly used in the
field[23,25,29], analysing continuous expression signatures and examining of how the size of the
neighbourhood influences spatial relationships are areas that remain underdeveloped. Furthermore,
inspecting the immediate neighbourhood of cells of interest versus broader hotspots within the

tissue will yield different insights into the tissue architecture and organisation, and this warrants
further investigation.

We introduce SpottedPy, a Python package that enables a statistically principled interrogation of
biological modules and cell-cell relationships at different scales in spatially profiled tissue (**Fig. 1b**).
This method encompasses:

• **Neighbourhood enrichment analysis:** Our method introduces a function designed to examine
correlations between cell states, populations or processes within individual spatial
transcriptomics spots and their immediate neighbourhood (**Fig. 1bi**). In this context, a
'neighbourhood' is delineated as a ring encompassing six Visium spots around a designated
central spot, which we calculate by treating the spatial spots as a network.

• **Hotspot identification:** We have implemented the Getis-Ord G* statistic to identify spatial
clusters of continuous signatures across the tumour slide. Users can flexibly filter for specific
regions within the slide to focus on creating hotspots. By comparing defined regions with high or
low expression or cell type signatures against a null hypothesis of random distribution, this
analysis reveals statistically significant 'hotspots' or 'coldspots'. Hotspots represent areas with a
high concentration of a particular cell type or signature, suggesting an aggregation that is
unlikely to be due to chance. Conversely, coldspots indicate regions where the target cells or
signatures are scarce, also beyond what would be expected in a random distribution. We
provide functionality to test enrichment of specific signatures within hot and coldspots.

• **Distance metrics:** SpottedPy includes analytical tools to measure and interpret the distances
between identified clusters, such as between tumour and immune hotspots. The primary method
for this calculates the *shortest path to a hotspot*, defined as the minimum distance from any
point within a defined hotspot to the nearest point within a specified comparison hotspot (**Fig.
1bii**). Additionally, SpottedPy introduces two supplementary methods for distance analysis. The
centroid to centroid approach compares the centroids of hotspots (**Fig. 1biii**) for a macro-level
view of spatial distribution. The *gene signature correlation* method correlates tumour signatures
within a defined region (therefore without hotspot creation) with the distance to other hotspots of
interest, allowing an investigation of continuous tumour state signatures (**Fig. 1 iv**). Importantly,
SpottedPy assigns a statistical significance to these proximity measures to determine if
observed patterns are likely to occur just by chance.

• **Sensitivity analysis:** We provide the ability to systematically evaluate how cell-cell
relationships evolve within the tissue as we vary the size of the neighbourhood or of the hotspot
of interest. For the neighbourhood enrichment approach, this can be assessed by varying the
number of concentric rings around the central spot. For the hotspot approach, SpottedPy
recalculates the Getis-Ord G* statistic with varying neighbourhood sizes, enabling the
identification of clusters at different spatial scales. By analysing how the distances between
hotspots change with neighbourhood size, the package can illuminate shifting spatial
relationships, providing insights into how biological entities interact across different scales.

Additionally, SpottedPy allows users to examine how cluster relationships change when
modifying the significance threshold for identifying hotspots with the Getis-Ord G* statistic.

To highlight the potential of our method, we employ SpottedPy to investigate the relationships
between tumour cells undergoing EMT and the TME in 12 breast cancer slides profiled using the
10x Genomics Visium spatial transcriptomics platform, integrated from Wu et al[18], Barkley et al[24]
and the 10x Genomics website[25]. To infer individual cell types within the slides, we performed cell
deconvolution using the Cell2location method[36] and a scRNA-seq reference of annotated breast
cancer cell population profiles from 123,561 cells[13]. To more precisely identify tumour cells, which
tend to show vast expression variability, we employed the copy number inference tool STARCH[37]
and only kept spots that showed evidence of copy number changes, which are likely to be tumour-
specific. Next, we delineated tumour hotspots using SpottedPy and scored them using an EMT
hallmark signature which has been linked with patient outcomes in the clinic[38,39]. Furthermore, to
explore the heterogeneity of EMT stable states established during the development and progression
of breast cancer, we employed the discrete EMT states recently defined by Brown et al[40],
encompassing an epithelial phenotype, two intermediate (hybrid) states (EM2 and EM3), a late
intermediate quasi-mesenchymal state (M1) and a fully mesenchymal state (M2).

The spatial landscape of EMT and associated tumour hallmarks

Firstly, we aimed to explore the cellular environment that is conducive to E/M progression. To do so,
we focused solely on tumour hotspots and two key cellular hallmarks that have been previously
linked with EMT, hypoxia and angiogenesis. Hypoxia, characterised by low oxygen levels, has been
long recognized as a key enabler of tumorigenic processes[41]. Under hypoxic conditions, tumour
cells stabilise hypoxia-inducible factors (HIFs), primarily HIF-1α, which promotes angiogenesis[42],
the formation of new blood vessels from pre-existing vasculature, to re-establish oxygen supply.
Hypoxia has been shown to induce EMT and resistance to therapy[43], and therefore understanding
how such relationships develop spatially within the tissue can help devise localised therapies that
can interrupt these interactions in breast cancer. Within individual spatial transcriptomics slides, we
used SpottedPy to delineate tumour areas, and further identified EMT hotspots within these areas
using using the EMT hallmark signature (**Fig. 2a**). To confirm and further explore the emergence of
additional cancer hallmarks in the context of EMT, we also defined proliferative, hypoxic and
angiogenic hotspots within the same slides (**Fig. 2a**). When visually inspecting the slides, we find
angiogenesis and hypoxia frequently accompanying EMT hotspots (**Fig. 2a**). When quantifying
hotspot distances using SpottedPy, we confirm that EMT hotspots tend to be closer to angiogenic
and hypoxic hotspots compared to EMT coldspots, proliferative hotspots, or the average tumour cell
(**Fig. 2b-c**). In contrast, proliferative hotspots were significantly closer to EMT coldspots (p<0.001,
Fig. 2c). These relationships were consistently observed across all breast cancer slides
(**Supplementary Fig. 1a**).

To further grasp the positioning of these EMT "hot" and "cold" areas within the tumour, we used
SpottedPy to determine the tumour perimeter (**Fig. 2d**) and calculated distances to it. We find EMT
hotspots closer to the tumour perimeter compared to EMT coldspots, suggesting a state with
significant interaction with the surrounding microenvironment (**Fig. 2e**). As expected, angiogenesis

hotspots were located closest to the tumour perimeter, followed by hypoxia hotspots (**Fig. 2f**). The
spatial localisation of angiogenesis near the tumour perimeter aligns with its function in supplying
nutrients and oxygen to rapidly growing tumours[44]. The prominence of hypoxic regions succeeding
angiogenic zones is consistent with the understanding that rapid tumour growth often outpaces its
vascular supply, leading to hypoxic conditions[41]. These hypoxic conditions are alleviated in the
angiogenic areas, as we find hypoxic coldspots are closest to the tumour perimeter (**Fig. 2f**). Cell
proliferation hotspots were observed furthest from the tumour perimeter, and located at spatially
distinct locations to EMT hotspots, in line with tumour growth studies outlining a proliferative
epithelial core and EMT transformation at the periphery facilitating cancer cell intravasation and
migration[45,46].
213
214 **EMT hotspots are immunosuppressed and shielded by myCAFs and macrophages**
215
Having confirmed that SpottedPy is able to recapitulate expected spatial hallmarks of EMT within
the breast cancer tissue, we next expanded our analysis to dissect the interplay between tumour
cells undergoing EMT and other immune and stromal cells in the microenvironment. Alongside EMT
hotspots, we calculated hotspots for 21 cell types in the TME, including different lymphocyte,
myeloid and fibroblast populations (**Fig. 3a-c**). When visually inspecting these hotspots, we
observed myCAF hotspots and EMT hotspots tended to co-localise (**Fig. 3d**). Quantifying hotspot
distances using SpottedPy allowed us to confirm that tumour EMT hotspots were indeed closest to
myCAF hotspots (**Fig. 3e**). This is well in line with existing studies, as myCAFs have been shown to
produce TGF-β, which is a well-known EMT trigger[47]. They have also been linked to ECM
deposition and suppression of antitumor immunity[48,49,50,51]. In contrast, iCAFs appeared to be the
most distant cell type to EMT hotspots (**Fig. 3e**). This is in keeping with observations in pancreatic
cancer by Ohlund et al[52], showing that direct juxtracrine interactions with cancer cells are required
for myCAF formation, whereas iCAFs are induced by secreted factors from cancer cells and thus
are not found in their proximity.
230
myCAF hotspots appeared significantly closer to EMT hotspots compared to EMT coldspots,
followed by macrophages, cycling perivascular-like cells (PVL), monocytes and dendritic cells (DCs)
(p<0.001, **Fig. 3f**). Monocytes, and particularly tumour-associated macrophages (TAMs) derived
from them, are known to modulate the environment of tumour cells undergoing EMT, usually by
promoting immune suppression in the TME, which facilitates tumour progression and metastasis[53].
PVL cells, a type of stromal cell characterized by a perivascular-like profile yet not associated with
blood vessels, represent a recently identified cell type whose functions are largely unknown[54].
Differentiated PVL cells have been linked to immune evasion in breast cancer[54]. Natural Killer (NK)
and NK T cells, the immune cells that can directly kill transformed cells, were the least associated
with EMT hotspots when averaged across slides, potentially reflecting a mechanism of immune
evasion employed by tumour cells that have undergone EMT (**Fig. 3f**).
242
The cohort-level associations between EMT hotspots, myCAFs and macrophages were maintained
in individual tumours, suggesting that these are universal patterns of EMT transformation in breast
cancer and not subtype specific (**Supplementary Fig. 1b**). In contrast, levels of inter-patient
heterogeneity, often even within the same breast cancer subtype, were evident for a variety of cells

including memory B-cells, naïve B-cells, iCAFs, NK cells, NKT cells, CD4+ T-cells and CD8+ T-
cells. Distances to EMT hotspots were consistent across subgroups of cells (**Supplementary Fig.**
1c), suggesting that, within individual patients, these cells share common response patterns
irrespective of the broader heterogeneity observed across the patient cohort.
251
Due to the close relationship with potential immunosuppressive actors, we next sought to test
whether EMT hotspots were indeed likely to be immunosuppressed. We found that EMT hotspots
displayed a significantly increased expression of immunosuppression and exhaustion markers[35]
compared to EMT coldspots (**Fig. 3g-h**), with the key immune checkpoints *B7-H3* (*CD276)*, *OX40*
(*TNFRSF4*) and *TIM3* (*HAVCR2*) displaying significantly higher expression (p<0.05) in the majority
on tumour slides (**Fig. 3h**). Thus, it appears that the chronic nature of immune activation nearby
EMT hotspots leads to the exhaustion of these cells, potentially suggesting opportunities for
immune reactivation through checkpoint blockade strategies. To verify this hypothesis, we
examined the expression of an interferon-gamma signature that has been associated with response
to immunotherapy[55] (**Fig. 3g**, **Fig. 3i**). We found that EMT hotspots had significantly increased
expression of genes within the signature, most notably of *HLA-DRA*, a marker of improved response
to anti-PD1 therapy[56].
264
265 Overall, these results showcase the range of hotspot analyses enabled by the SpottedPy package
266 and the potential to uncover useful biological insights.
267
268 **Sensitivity analysis of hotspots**
269
Determining how the hotspot size, governed by the parameter specifying the number of nearest
neighbours for the Getis-Ord Gi* metric, affects spatial relationships is crucial for meaningful spatial
analysis. We systematically increased hotspot dimensions (**Fig. 4a-b**) to assess the consistency
and robustness of identified spatial associations. We find that associations between EMT hotspots,
hypoxia and angiogenesis as well as mutual exclusivity with proliferative hotspots are robust and
consistent features of the tumour microenvironment, with such relationships remaining remarkably
stable across a range of hotspot dimensions (**Fig. 4c**). The cell populations that we previously
identified as having the nearest proximity to EMT hotspots at a fixed parameter size (myCAFs,
macrophages, cycling PVL cells and monocytes) also maintained this relationship when varying
hotspot sizes. Cells that were farther apart presented less stable associations, such as dendritic
cells where the relationship broke down at a hotspot size of 200, and naive B-cells where the
relationship changed multiple times with increasing hotspot size (**Fig. 4c**, **Supplementary Fig. 2a**).
These findings suggest that interactions with certain cells in the TME may be more pronounced and
relevant at a smaller scale. We found that NK cells were the only cells consistently adjacent to EMT
coldspots at various hotspot sizes. Adjusting the p-value cut-off used to detect spatial clusters using
the Getis-Ord Gi* highlighted similar relationships (**Supplementary Fig. 2b**). We also see that the
number of hotspots decreases when increasing the hotspot parameter size, indicative of the
hotspots becoming more clustered together as the neighbourhood expands, as expected (**Fig. 4d**).
288
289
290

Other distance metrics

All the results described so far have evaluated distances between hotspots using the shortest path method (**Fig. 1bii**). We thus wondered whether the stability of these relationships would be affected by the method chosen to measure hotspot distances. To investigate this, we used the "centroid to centroid" approach to calculate the distance from centroids of each EMT hotspot to centroids for cancer hallmarks and immune/stromal hotspots, with the resultant rankings displayed in **Supplementary Fig. 2c-d**. Reassuringly, we observe a consistent relationship with tumour hallmark signatures, with EMT hotspots located closer to hypoxic and angiogenic regions compared to EMT coldspots and further away from proliferative regions. We also see cycling PVLs and myCAFs remaining in the top cell types associated with EMT hotspots, while macrophages and monocytes appear more distal to these hotspots compared to our previous shortest path distance approach.

Whilst the "centroid to centroid" methodology offers a practical and straightforward way to approximate the distances between hotspots, it is crucial to acknowledge its simplicity. As depicted in **Supplementary Fig. 2e**, the size of a hotspot significantly influences its centroid location. Consequently, a larger hotspot, despite being physically closer, may appear farther away when measuring centroid distances, due to its centroid being located farther from the point of interest compared to a smaller, neighbouring hotspot. Thus, the centroid approach may miss local variation that the shortest path method can capture. This could suggest that applying it on our breast cancer slides could potentially miss the more complex relationship observed between EMT hotspots and macrophage or monocyte-enriched areas.

Neighbourhood enrichment analysis

The neighbourhood enrichment technique captures more localised, shorter-range relationships with the TME, by correlating phenotypes across a central tumour spot and the direct neighbourhood surrounding it (a ring encompassing six Visium spots) (**Fig. 1bi**). Varying the number of rings around the central spot enables us to assess whether the observed hotspot relationships shift with the unit of analysis.

Our analysis revealed that angiogenesis, myCAFs and monocytes exhibited the most significant correlation, in descending order, with EMT hallmark signatures ($p < 0.001$) across the 12 slides (**Fig. 5a**). This finding reinforces the spatial association between these cell types and signatures with EMT that we previously identified using the hotspot method. A significant association was not evident for naive B-cells or CD4+ T-cells (**Fig. 5a**). This lack of a discernible relationship using the neighbourhood enrichment approach for CD4+ T-cells, despite the hotspot approach capturing a weak significant relationship, suggests that the relationship is more indirect, potentially occurring over a broader spatial expanse than our neighbourhood enrichment approach can accurately capture.

We found that these spatial relationships were stable across various neighbourhood sizes (**Fig 5b-c**). Among the most consistent were the hypoxia and angiogenic signatures. Conversely, relationships between EMT and plasmablasts, iCAFs or proliferative hotspots were the most

335 variable (**Fig 5c**), suggesting that this relationship might be more strongly determined by the
336 environment in a local neighbourhood rather than at a broader scale.
337
338 **EMT state fluctuations shape distinct immune niches within the same tumour**
339
As mentioned previously, EMT is not a binary process – instead, cells are found to occupy multiple
hybrid states during the E/M transition. We sought to investigate the spatial distribution of tumour
hotspots occupying epithelial (EPI), early intermediate (EM2, EM3), late intermediate quasi-
mesenchymal (M1) and fully mesenchymal (M2) states using our multi-scale approach. We
captured distinct gene programs representing these states using Non-Negative Matrix Factorisation
(NMF) via the CoGAPs workflow[57] (**Supplementary Fig. 3a**). The corresponding hotspots occupied
distinct spatial locations within the tissue (**Supplementary Fig. 3b**). When visually inspecting these
hotspots, we detected a progressive transformation in the tumour, highlighted in Slide 4 (**Fig. 6a**).
This transition was marked by a spatial shift from EPI into the M1 state, with EM3 serving as an
intermediate stage. EM2 displayed volatility in this progression, while M2 was predominantly co-
localised with the EPI state. The experimental study by Brown et al[40] had detected that M2 cell
clones gained integrin β4 (a key epithelial marker) when cultured, which might have played a
significant role in steering these cells towards adopting characteristics more akin to an epithelial
phenotype. This would possibly explain the co-localization of these two states within the spatial
transcriptomics slide.
355
To investigate the relationship between these states, we correlated them with each other and the
generic EMT hallmarks signature employed previously for each tumour spot (**Supplementary Fig.
4a**). The lack of significant positive correlation between EPI, EM2, EM3, M1, M2 suggests that these
are discrete EMT states. We found a significant correlation between the EMT hallmark signature
and the quasi-mesenchymal M1 state, suggesting these are likely capturing a similar state. The EPI
state was negatively correlated with EMT hallmarks signature, as expected. EM2, EM3 and M2 are
likely distinct states, as evidenced by the lack of correlations across the spots. In terms of spatial
distribution, the EMT hallmark hotspots were located closest to the M1 hotspots, and furthest away
from the EPI hotspots (**Supplementary Fig. 4b**), in line with the correlation analysis and confirming
the hypothesised identities of these states.
366
When investigating how tumour cells occupying distinct EMT states relate to their
microenvironment, we found that the EPI state has negative correlations with immune and stromal
cells within the TME, suggestive of a state that is not directly being shaped by the TME (**Fig. 6b**).
Interestingly, the M1 state had stronger associations overall with a wide range of cells within the
TME, with the strongest relationships established with myCAFs, macrophages and monocytes. We
observed similar but weaker correlations with the EM3 state, and considerably weaker correlations
with the EM2 state. The progressive loss of association with cells in the EM3 and EM2 states is in
line with the idea of these states representing intermediate, more plastic states preceding the
apparently more stable M1 state. The M1 state is observed in proximity to natural killer (NK) cells,
which is a distinct deviation from the EMT hallmark signature. This observation suggests that while
there are resemblances between the M1 and EMT hallmark signatures, the M1 state is likely

representative of a unique cellular phenotype with the potential to recruit cells capable of directly
eliminating cancer cells.
380
Furthermore, the quasi-mesenchymal M1 state presented an enrichment of markers linked with
immunosuppression and positive response to checkpoint inhibitors (**Fig. 6c**, **Supplementary Fig.**
4c), with a subset of genes, most notably *OX40 (TNFRSF4), TIM3* (*HAVCR2*), *HLA-DRA, CXCL9*
and *CXCL10*, driving these relationships (**Fig. 6d-e**). The intermediate states are to a certain extent
on the way to adopting this immune suppressive phenotype, with weaker relationships observed
with the EM2 state, a slightly stronger enrichment with EM3 and the strongest score with M1 (**Fig.**
6c-e, **Supplementary Fig. 4d**). In contrast, the M2 state has a unique phenotype, and displays both
positive and negative relationships with genes within these signatures. We compared these states
to the proliferative signature and found that proliferative hotspots mirror the relationship of the EPI
state, suggesting that it is a tumour state that is not linked to immunosuppression.
391
392 Overall, this analysis sheds light on the changing landscape of tumour-TME interactions during EMT
393 progression in breast cancer, highlighting both intratumour heterogeneity as well as universal
394 interactions that could be exploited for therapy.
395
396
397 **DISCUSSION**
398
In this study we introduce SpottedPy, a Python package that identifies tumour hotspots in spatial
transcriptomics slides and explores their interplay with the TME at varying scales. We show that the
Getis-Ord G* statistic can be successfully applied to delineate cellular hotspots and provide
meaningful biological insights into the spatial organisation of the tumour tissue in its immune and
stromal contexture. Whilst various studies have recently applied hotspot analysis to spatial
transcriptomic data[58,59,21], these methods do not analyse the distances between hotspots. We build
on these approaches to analyse the spatial relationships between hotspots, with the additional
ability of anchoring hotspot identification to specific regions of interest, such as tumour cells or non-
tumour cells, enabling us to delve into the spatial dynamics within the TME via a more targeted
approach. By computing and statistically comparing distances, we offer an interpretable and intuitive
measure of relationship between spatial variables. Our method additionally assesses the effect of
hotspot size on spatial relationships and compares hotspot spatial trends to the relationships
captured using neighbourhood approaches, more frequently applied in spatial transcriptomic
analysis, to build layers of spatial evidence.
413
By adopting our SpottedPy methodology to explore the spatial dynamics of tumour plasticity
phenotypes in breast cancer, we have uncovered key differences between tumour regions
undergoing EMT and those lacking evidence for this transformation. Our approach illuminates the
pronounced spatial correlations of EMT with key cancer hallmarks, notably hypoxia and
angiogenesis, in line with findings by He et al[11] in breast cancer spatial transcriptomics detecting
these signatures overlapping certain niches. As tumour cells undergo EMT in response to hypoxic
stimuli, they are likely to gain a survival advantage in a nutrient-deprived environment and be better

equipped to invade and migrate towards regions with better oxygenation, potentially following
angiogenic gradients[60,43,46].
423
We find a strong relationship between EMT and myCAFs across all slides. CAFs have been linked
to tumour cells undergoing EMT in our previous bulk and spatial transcriptomic analyses[34], and
have been shown to induce EMT in endometrial cancer cells[61] and hepatocellular carcinoma[62].
However, the relationship between the recently defined CAF subtypes, myCAFs and iCAFs, and
EMT has been less well studied and here we show specific interactions with the former and not the
latter. It is worth noting that CAFs share similar genes with EMT signatures and therefore
differentiating between these two cell types can be challenging, notably in bulk tumour settings[63].
Here we have leveraged scRNA-seq for deconvolution, alongside detecting copy number
aberrations (CNAs) to confirm the presence of tumour cells, which adds an additional layer of
confidence to the accurate delineation of these cell populations. However, the accuracy of this
separation cannot be fully guaranteed and future research investigating this relationship using
single-cell resolution spatial transcriptomic data would allow us to confirm this relationship more
confidently.
437
In addition to the expected myCAF associations, we observe a strong relationship with
macrophages and monocytes across multiple spatial scales. Macrophages secrete TGF-β, TNF-α,
IL-6 and IL-8, which are all well characterised EMT stimuli[64,65]. The relationship has been observed
in bulk transcriptomics[66], in spatial analysis of mouse models of skin carcinoma where depletion of
macrophages inhibited EMT progression[31] and in specific niches within breast cancer spatial
transcriptomics slides[11].
444
We uncovered heterogeneous relationships across breast tissue slides with other key immune cells
such as NK cells, NKT cells and T cells, highlighting the multifaceted interplay between these
components. T-cells have been shown to induce EMT in breast cancer[67,68], and this relationship has
further been highlighted in bulk transcriptomics[69] and smaller scale spatial transcriptomic
analyses[34,70] however there is also evidence showing the exclusion of these cells, linked to the
relationship between EMT, macrophages and CAFs promoting an immune suppressed
environment[64,51,71]. Indeed, our findings uncover notable associations between EMT hotspots and
immune suppression, alongside signatures indicative of a response to checkpoint therapy, building
upon evidence that EMT may offer crucial insights for existing strategies in immunotherapy[72,11].
454
Our analysis reveals that EMT occurs in discrete spatial locations distinct from proliferative
signatures. This finding is in line with a recent analysis of breast cancer by Jia et al[51] utilising a more
focused spatial transcriptomic dataset, previous research by Tsai et al[73] demonstrating that a
departure from a mesenchymal-like state is a prerequisite for tumour cell proliferation in mouse
models, and a recent study by Chen et al[74] investigating EMT states in scRNA-seq data. Such
spatial characterisations at various scales were largely unexplored.
461
Delving deeper into EMT, we observed that hybrid EMT states exhibit more heterogeneous and
weaker associations with the TME in comparison to the quasi-mesenchymal M1 state. This disparity
might be indicative of the inherent plasticity of hybrid EMT states[75,76], complicating our ability to

delineate clear relationships, but also of a directed trajectory towards an M1 state. The M2 state
demonstrates relatively weaker connections with the TME, which may be attributed to the
experimental gain of integrin β4 (a key epithelial marker) when cultured, a limitation mentioned in
the original study which potentially transformed the state towards a more epithelial phenotype[40]. We
found that the epithelial state exhibits a notable degree of independence from the TME which may
be due to the ability of cancer cells to successfully avoid immune recognition in the incipient stages
of neoplasia.
472
These results point to a highly dynamic and plastic nature of tumour cells in navigating the
complexities of their microenvironment. The interactions likely extend beyond a linear framework.
Hypoxia, a known catalyst for both angiogenesis and EMT[77], can set off a cascade of events that
not only amplify these processes but also create a conducive environment for the recruitment of
immunosuppressive cells such as macrophages[78,79]. These cells can in turn bolster angiogenesis,
thereby fuelling a self-perpetuating cycle that further complicates the tumour landscape. These
insights provide a further understanding of the cellular interactions and environmental factors that
underpin tumour progression and metastasis, and could in the future pave the way for the
development of targeted interventions aimed at disrupting these complex networks for therapeutic
benefit.
483
The consistency of the relationships we observed across different hotspot sizes and neighbourhood
scales further strengthens our confidence in the findings. The neighbourhood ring approach
predominantly captured TME cells that have infiltrated the tumour, offering insights into the
immediate cellular interplay at the tumour periphery. In contrast, the hotspot methodology provided
a broader view, encompassing interactions at more distal locations. By pinpointing statistically
significant cellular hotspots, we bolster the reliability of our observations, especially considering the
inherent inaccuracies that can arise from deconvolution algorithms applied to non-single cell
transcriptomic datasets such as those from the Visium platform. Furthermore, we offer three
methods to analyse hotspot distances, which allows flexibility in the analyses depending on the
biological questions that the user intends to answer.
494
In the future, a comprehensive benchmarking of different feature-based spatial clustering
approaches, such as kernel density estimation, and different statistical approaches to quantify the
distances between different hotspots would be valuable. This would enable us to determine how
sensitive spatial relationships are to methodological choice. Moreover, extending these analyses to
single cell-resolved spatial datasets and incorporating ligand-receptor signalling information into the
evaluation of spatial effects on cell populations will increase the confidence in the identified
relationships.
502
Our insights into the spatial organisation of tumours during EMT progression are limited by the
significant amount of uncertainty surrounding EMP programmes[32,33] and their incomplete
characterisation in different types of breast cancer and other neoplasms. In the future, integrating
further hybrid states characterised in other breast cancer studies[74] will help expand our
understanding of this complex process alongside its multiple locally stable peaks and valleys. The
heterogeneous relationships observed with CD8+ and CD4+ T-cells and NK cells require further

experimental validation and exploration. As spatial transcriptomics datasets become more widely
available, expanding this analysis beyond the current 12 slides could help clarify the perceived
spatial heterogeneity and better distinguish universal relationships from local, patient-specific
effects.

Overall, our findings confirm expected spatial effects of EMT progression in tumours, demonstrating
that SpottedPy is an adequate tool to capture complex associations between tumour cells and their
microenvironment. Such insights can help unveil local effects of the TME and linked tumour cell
vulnerabilities that could ultimately be exploited for therapeutic benefit. While the analyses
presented here illustrate insights into breast cancer tissue organisation, we note that SpottedPy can
be applied to discern spatial relationships in other cancer types as well as other diseases and even
within healthy tissue. SpottedPy has been developed on spatial transcriptomics data from the 10x
Visium platform; however, we note it can be easily extended to other spatially-resolved platforms
and future releases will provide further functionality to enable this.

Conclusion

In conclusion, SpottedPy provides a detailed and multifaceted analysis of the spatial dynamics
within individual spatially-profiled tumours. By rigorously investigating the proximities of various
cellular components, we have underscored the significant influence exerted by cells undergoing
EMT in sculpting the TME and highlight SpottedPy as a package that can be applied to answer
other spatial biology questions.

531
532
533 **METHODS**

534
Spatial transcriptomic datasets

Breast cancer Visium slides were obtained from Barkley et al.[29] (slides 0-2), from 10x Genomics
(slides 3-5)[35] and Wu et al.[12] (slides 6-12).   Slide annotations, if available, are displayed in
Supplementary Fig. 1b. We combined the three datasets of breasts cancer 10X Genomics Visium
spatial transcriptomic datasets into a common *anndata* Python format for analysis. Pre-processing
and normalisation were conducted using the ScanPy (Single-Cell Analysis in Python) package[80].
We analysed a total of 32,845 spatially profiled spots, and retained spots if they exhibited at least
100 genes with at least 1 count in a cell, had more than 250 counts per spot and less than 20% of
total counts for a cell which are mitochondrial.

Spatial data deconvolution

545
Due to the imperfect near-single cell resolution of current spatial transcriptomic methods, we require
a method to deconvolve each spot in order to infer the cellular populations enriched in each spot.
Cellular deconvolution was carried out using Cell2location[36]. Cell2location decomposes the spatial
count matrix into a predefined set of reference cell signatures by modelling the spatial matrix as a
negative binomial distribution, given an unobserved gene expression level rate and gene- and
batch-specific over-dispersion. A scRNA-seq breast cancer dataset containing 100,064 cells from

26 patients and 21 cell types from Wu et al[12] was chosen to perform the deconvolution. Cell types
in the chosen breast dataset consisted of cancer epithelial cells (basal, cycling, Her2, LumA, LumB),
naïve and memory B cells, myCAF-like and iCAF-like cancer-associated fibroblasts, perivascular-
like cells (PVL), including immature, cycling and differentiated, cycling T-cells, cycling myeloid cells,
dendritic cells (DCs), endothelial cells expressing *ACKR1*, *CXCL12* or *RGS5*, endothelial lymphatic
LYVE1-expressing cells, luminal progenitors and mature luminal cells, macrophages, monocytes,
myoepithelial cells, natural killer (NK) cells, natural killer T (NKT) cells, plasmablasts, CD4+ T cells,
and CD8+ T cells. The scRNA regression model was trained with 500 epochs, and the spatial
transcriptomic model trained with 20,000 epochs using a GPU. To delineate the tumour cells within
our spatial transcriptomics dataset, we used the STARCH Python package designed to infer copy
number alterations (CNAs)[37]. STARCH identifies tumour clones (setting K=3 clones) and non-
tumour spots. It confirms identification of normal spots by clustering the first principal component
into two clusters using K-means (setting K=2). This approach is based on the principle that the
direction of maximum variance in the expression data typically reflects the division between non-
cancerous and cancerous spots.
567
568   **EMT state and hallmark signature scoring**
569
To identify distinct EMT states, we employed data from Brown et al[40], consisting of seven RNA-seq
sequenced cell clones, derived from SUM149PT inflammatory breast cancer cell line with 3 repeats
spanning the EMT spectrum from epithelial-like (EPI), quasi-mesenchymal (M1), fully mesenchymal
(M2) and three distinct intermediates (EM1, EM2, EM3) . We used these data to derive a weighted
gene signature to represent the EMT states. We captured EMT gene patterns from this data using
non-negative matrix factorisation (NMF) by applying the CoGAPs workflow[57]. We used ProjectR's
implementation of *lmfit* R function to map the captured EMT patterns onto the spatial transcriptomic
spots[81]. This transfer learning approach assumes that if datasets share common latent spaces, a
feature mapping exists between them and can measure the extent of relationships between the
datasets. The final states were captured with 20 patterns and 10,000 training iterations. The number
of patterns were chosen based on capturing the discrete states with the highest accuracy. The EM1
state was not distinguishable from the EPI state, so we merged the two states. Thus, overall we
obtained scores for one epithelial, two intermediate, a quasi-mesenchymal and a fully mesenchymal
state for each spot.
584
585   The EMT, hypoxia and angiogenic hallmark signatures[38], response to immunotherapy signature[82],
586   immunosuppressive signature[12] and proliferative signature[83] were scored using
587   *scanpy.tl.score_genes* function.
588
589   **Graph construction**
590
591   The SquidPy[23] (Spatial Single-Cell Analysis in Python) package was used for graph construction
592   using *sq.gr.spatial_neighbors* and slide visualisation of the Visium spatial slides. NetworkX[84] was
593   used for further analysis of the networks derived from the spatial transcriptomic spots. The
594   deconvolved spot results were used to assign node labels. Edges were assigned based on the spot
595   neighbours.

Neighbourhood enrichment analysis

We calculated neighbours for each spot by summing the deconvolution results in a ring surrounding the spot of interest, and normalising by the number of spots assigned as a neighbour, using the adjacency matrix of the graph to calculate the interacting cells.

Neighbourhood enrichment was calculated by correlating signatures across a central spot of interest and the direct neighbourhood of spots surrounding it (a ring encompassing six Visium spots), after filtering for tumour spots only. Correlations were calculated using Pearson's correlation coefficient. To perform the sensitivity analysis, we increased the number of rings surrounding a spatial transcriptomic spot to consider as spot neighbours and compared the change in correlation coefficient.

We additionally compared this neighbourhood enrichment strategy to an approach solely correlating the central spot with the cellular composition in the neighbourhood ring around the spot, for which found less stable spatial relationships. Consequently, we kept the formed as default neighbourhood enrichment method, although the latter is also available to employ in the SpottedPy package.

Hotspot analysis

Hotspots were calculated using The Getis-Ord G* statistic as implemented using the PySAL package[85], using 10 as neighbourhood size parameter by default and a p-value of 0.05, unless otherwise stated.

The Getis-Ord G* equation is defined as follows:

$$G_i^* = \frac{\sum_{j=1}^{n} w_{ij} x_j - \bar{x} \sum_{j=1}^{n} w_{ij}}{s \sqrt{\frac{n \sum_{j=1}^{n} w_{ij}^2 - (\sum_{j=1}^{n} w_{ij})^2}{n-1}}}$$

Where w_{ij} is the spatial weight between location i and j, $\bar{x}$ is the mean of the variable of interest across all locations, s is the standard deviation of the variable of interest across all locations and n is the total number of locations.

A high positive value at location i suggests a hotspot for the attribute, while a negative value indicates a coldspot. The significance of G^* is determined by comparing the observed G_{obs}^* to a distribution of G^* values generated under the assumption of spatial randomness. This distribution is obtained by permuting the attribute values across locations and recalculating G^* for each permutation. The p-value for a hotspot (when G^* is positive) or a coldspot (when G^* is negative) is then derived from this distribution. This approach provides a non-parametric method to evaluate the significance of spatial clusters, offering a robust measure against potential spatial randomness in the data. Hotspots for EMT states and cell proliferation were calculated after filtering for tumour cells as labelled by STARCH. at location i suggests a hotspot for the attribute, while a negative value indicates a coldspot. The significance of G^* is determined by comparing the observed G_{obs}^* to a distribution of G^* values generated under the assumption of spatial randomness. This distribution is

obtained by permuting the attribute values across locations and recalculating G^* for each
permutation. The p-value for a hotspot (when G^* is positive) or a coldspot (when G^* is negative) is
then derived from this distribution. This approach provides a method to evaluate the significance of
spatial clusters, offering a robust measure against potential spatial randomness in the data.
Hotspots for EMT states and cell proliferation were calculated after filtering for tumour cells as
labelled by STARCH.
642
643 **Distance metrics**
644
645 We used the ***shortest path*** approach to calculate distances between hotspots as follows:
646 ● Let H represent the set of coordinates of spots in the hypoxia hotspot.
647 ● Let M represent the set of coordinates of spots in the mesenchymal tumour hotspot.
648 ● Let E represent the set of coordinates of spots in the epithelial tumour hotspot.
649
650 For a spot m in M and a spot e in E, we calculated the Euclidean distance to a spot h as $d(m,\ h)$.
651 The minimum distances from M to H and E to H are:
652

$$d_{min}(M, H) = \min_{m \in M, h \in H} d(m, h)$$

653
654

$$d_{min}(E, H) = \min_{e \in E, h \in H} d(e, h)$$

655
656
657 We can then compare the distribution of distances between the different spots in the hotspots.
658
659 The ***centroid approach*** is calculated as follows. The centroid C_H of a set of spots H with
660 coordinates x_h, y_h is the arithmetic mean of the coordinates. This point represents the centre of the
661 mass of the points in the set H.
662 For set H :

$$C_H = \left(\frac{\sum_{h \in H} x_h}{|H|}, \frac{\sum_{h \in H} y_h}{|H|} \right)$$

663
664
665 Similar calculations are employed for M and E. We then calculated the Euclidian distance between
666 the centroids.
667
668 We calculated the ***gene signature correlation*** metric by correlating each tumour signature value
669 with the distance from that spot to the nearest hotspot of interest. We then compared how different
670 signatures, using a continuous score to represent them, are correlated to distances to other
671 hotspots of interest.
672
673 **Tumour perimeter calculation**
674
Any spot was considered part of the tumour perimeter if it had more than two neighbouring spots
(nodes in the graph) that were not classified as tumour spots. This approach helped us to delineate
the boundary of the tumour accurately by focusing on the transitional area where tumour and non-

tumour spots meet. To quantify the number of tumour hotspots, we calculated the number of
connected components within the graph that were labelled as hotspots. This calculation was crucial
for understanding the distribution and clustering of tumour cells.
681
682 **Sensitivity analysis**
683
684 The sensitivity analysis to evaluate the impact of varying hotspot sizes on the spatial relationships
685 was achieved by incrementally adjusting the neighbourhood parameter for the Getis-Ord statistic,
686 which directly influenced the size of identified hotspots. As we expanded the neighbourhood
687 parameter, we compared the distances between the newly defined hotspots and other existing
688 hotspots of interest.
689
690 **Statistical analysis**
691
692 Groups were compared using a two-sided Student's t test. Multiple testing correction was performed
693 where appropriate using the Bonferroni method. Graphs were generated using the seaborn and
694 Matplotlib Python packages.
695

FIGURES

Figure 1: SpottedPy provides a multi-scale approach to analyse spatial transcriptomic relationships. (a) Overview of spatial scales captured in the SpottedPy workflow, from direct cellular contacts to broader cellular hotspots. Figure created with BioRender.com. **(b)** SpottedPy workflow overview. Visium spatial transcriptomic data is loaded as a pre-processed *AnnData* object where there is the option to select the region of interest within the slide e.g., *AnnData.obs* column labelled with tumour cells. Hotspots are identified based on transcriptional signatures or cell types. Key spatial measurements are highlighted in purple, with the additional functionality in blue. The default distance metrics include: (i) Neighbourhood enrichment, which correlates cell prevalence or signatures in individual spots with their immediate neighbourhood, and (ii) Shortest path to hotspot,

calculating the minimum distance between each spot within a selected hotspot and the nearest spot
in other hotspots. Additional features for analysing hotspot distances in the package include: (iii)
Centroid-to-centroid, which calculates distances between the centroids of two hotspots, and (iv)
Gene signature correlation, correlating gene signatures with distances to hotspots. Sensitivity
analysis allows us to compare relationships defined at different scales in both approaches, either by
increasing the number of rings included for neighbourhood enrichment or increasing the hotspot
size. The outputs for the different modules include various plots to highlight the relationships. Figure
created with BioRender.com.
716

717
Figure 2: The spatial interplay between EMT progression and cancer hallmarks. (a) Spatial
transcriptomics slide (Slide 0) highlighting from left to right: tumour spots, the EMT gradient across
these tumour spots, EMT hotspots, hypoxic hotspots and angiogenic hotspots identified by
SpottedPy. The black square indicates a representative area where the close proximity of EMT,
angiogenic and hypoxic hotspots is depicted. **(b)** Bubble plot illustrating the distance from EMT
hotspots, EMT coldspots, proliferative hotspots and the average tumour cell to angiogenic and
hypoxic hotspots, averaged across all the 12 samples. Smaller bubbles represent increased vicinity.
(c) Differences in proximity between EMT hotspots/coldspots and hypoxic, proliferative and
angiogenic regions, summarised across the 12 slides. The dashed line represents no difference in
relative distance to EMT hotspots or EMT coldspots. The dots situated above the dashed line
indicate hallmarks that are significantly closer to EMT hotspots. The colours indicate the p-value
ranges obtained from the Student's t test for differences in distance to EMT hot/cold areas. **(d)**
Spatial plot depicting the tumour perimeter in red, tumour cells in blue and the microenvironment in
grey. **(e)** Distance from the tumour perimeter to EMT hotspots and EMT coldspots, respectively.
Smaller bubbles represent closer hotspots, and the data are summarised across all 12 slides. **(f)**
Distances from selected hotspots to the tumour perimeter, ordered by increasing proximity, across
the 12 cases. The dashed line represents no significant difference. The colours depict p-value
ranges obtained from Student's t tests for differences in distance to the tumour perimeter.
736

Figure 3: The spatial interplay between EMT progression and the TME. (a) Spatial transcriptomics plots highlighting tumour cell spots (left), the EMT gradient through these tumour

spots (middle) and EMT hotspots identified by SpottedPy (right) in Slide 5. **(b)** Spatial localisation of
macrophage-enriched spots (left) and SpottedPy-defined macrophage hotspots (right) in Slide 5. **(c)**
Spatial localisation of myCAF-enriched spots (left) and SpottedPy-defined myCAF hotspots in Slide
5. **(d)** EMT hotspots, EMT coldspots and myCAF hotspots overlayed on the same slide (left), and
illustrated within the context of all tumour areas (right). The zoomed in area of interest (black
square) highlights the close proximity of EMT hotspot to myCAFs. **(e**) Distance between EMT
hotspots and different TME cell hotspots, ranked by proximity. Smaller bubbles represent shorter
distances to EMT hotspots. Results are averaged over 12 slides. **(f)** Distances from various cells in
the TME to EMT hotspots/coldspots. The dashed line represents no difference in proximity to either
EMT hotspots or EMT coldspots. The dots situated to the right of the dashed line indicate cell
populations that are significantly closer to EMT hotspots, ordered by decreasing proximity. The
colours indicate the p-value ranges obtained from the Student's t test for differences in distance to
EMT hot/cold areas. Results are averaged across 12 slides. **(g)** Barplots showing signature scores
of immune suppression[12] and checkpoint inhibitor response[82] scored within EMT hotspots and
coldspots. **(h)** Differences in the average expression of genes in the immune suppression
signature[12] between EMT hotspots and EMT cold spots for each slide (row). Red depicts genes
significantly upregulated in EMT hotspots and blue indicates genes significantly upregulated in EMT
coldspots (Student's t test p<0.05, adjusted for multiple testing using the Bonferroni correction).
White indicates a non-significant relationship. **(i)** Similar to (g) but focusing on the genes in the
checkpoint inhibitor response signature[82].
760

Figure 4: Sensitivity analysis of hotspot relationships. (a) EMT hotspot generation using a hotspot neighbourhood parameter of 2, 10, 50, 100 and 300, respectively. Increasingly larger neighbourhoods are highlighted in different colours as indicated in the legend. **(b)** Hypoxia and epithelial hotspot generation using a hotspot neighbourhood parameter of 2, 10, 50, 100 and 300, respectively. **(c)** Sensitivity plots highlighting the distance from EMT hotspots (blue) and EMT

767 coldspots (green) to regions enriched in various cancer hallmarks and TME components as the
768 hotspot size increases. The distances to the average of all tumour cells are used as a reference
769 (orange). Distances are averaged over all 12 slides. **(d)** Number of distinct hotspots identified as the
770 hotspot neighbourhood parameter is increased, averaged over 12 slides.
771

Figure 5: Neighbourhood enrichment analysis of EMT spatial dynamics. (a) Neighbourhood
enrichment analysis results displaying correlations between EMT within tumour cell spots and the

abundance of various cell populations within the immediate TME (surrounding spots only). Red
indicates a positive correlation, blue a negative correlation and white a non-significant correlation
(Pearson p>0.05). 1 ring is used to define the neighbourhood. **** p<0.0001, *** p<0.001, ** p<0.01,
* p<0.05. **(b)** Line plots illustrating the impact of progressively expanding the number of concentric
rings - from 1 to 100- around a Visium spot on the correlation between the EMT signature and
various cells in the TME. **(c)** Bar plot highlighting the change in correlation strength as ring size is
increased from 1 to 100. Red indicates a positive correlation change, blue a negative change.
782

Figure 6: EMT state dynamics uncovered from the spatial exploration of breast cancer tissue.
(a) Spatial plots depicting epithelial (EPI), intermediate (EM2, EM3), quasi-mesenchymal (M1) and fully mesenchymal (M2) hotspots in Slide 4. **(b)** Neighbourhood enrichment analysis depicting the association between tumour cells occupying distinct EMT states and other cells in the immediate TME, summarised across all 12 slides. Red indicates a significant positive correlation (Pearson, p<0.05), blue a significant negative correlation (p<0.05) and white a non-significant correlation (p>0.05). **** p<0.0001, *** p<0.001, ** p<0.01, * p<0.05. **(c)** Scaled immune suppression[12] and

immunotherapy signature[82] scores calculated using Gene Set Enrichment Analysis (GSEA) for each
EMT state hotspot and proliferative hotspot, summarised across the 12 samples. **(d)** Enrichment
and depletion of expression for genes in the immune suppression signature within EMT state
hotspots for each slide (row). Red depicts genes significantly upregulated in EMT state hotspots
compared to the average of all tumour cells and blue represents genes significantly downregulated
in the EMT state hotspot (Student's t test $p<0.05$). White indicates a non-significant relationship. P-
values were adjusted for multiple testing using the Bonferroni correction. **(e)** Similar to (d), focusing
on genes in the checkpoint inhibitor response signature.
799
800
801

DECLARATIONS

Ethics approval and consent to participate

All datasets employed in this study are publicly available and comply with ethical regulations, with approval and informed consent for collection and sharing already obtained by the relevant organizations.

Consent for publication

Not applicable.

Availability of data and materials

Breast cancer Visium slides were obtained from Barkley et al.[29], Wu et al.[12] and the 10x Genomics website[35].

SpottedPy is implemented as a Python package available at https://github.com/secrierlab/SpottedPy, released under a GNU General Public License v3.0 with an accompanying tutorial. Processed spatial transcriptomics data have been deposited at Zenodo: https://zenodo.org/records/10371890.

Competing interests

The authors declare no competing interests.

Funding

EW was supported by a studentship award from the Health Data Research UK-The Alan Turing Institute Wellcome PhD Programme in Health Data Science (218529/Z/19/Z). MS was supported by a UKRI Future Leaders Fellowship (MR/T042184/1). Work in MS's lab was supported by a BBSRC equipment grant (BB/R01356X/1) and a Wellcome Institutional Strategic Support Fund (204841/Z/16/Z).

Authors' contributions

MS designed the study and supervised the analyses. EW developed the SpottedPy package and performed all the analyses.

REFERENCES

1. Yuan, Y. Spatial Heterogeneity in the Tumor Microenvironment. *Cold Spring Harb Perspect Med* **6**, a026583 (2016).

2. Ståhl, P. L. *et al.* Visualization and analysis of gene expression in tissue sections by spatial transcriptomics. *Science* (2016) doi:10.1126/science.aaf2403.

3. Asp, M., Bergenstråhle, J. & Lundeberg, J. Spatially Resolved Transcriptomes—Next Generation Tools for Tissue Exploration. *BioEssays* **42**, 1900221 (2020).

4. Moses, L. & Pachter, L. Museum of spatial transcriptomics. *Nat Methods* **19**, 534–546 (2022).

5. Marx, V. Method of the Year: spatially resolved transcriptomics. *Nat Methods* **18**, 9–14 (2021).

6. Zhang, L. *et al.* Clinical and translational values of spatial transcriptomics. *Sig Transduct Target Ther* **7**, 1–17 (2022).

7. Andersson, A. *et al.* Spatial deconvolution of HER2-positive breast cancer delineates tumor-associated cell type interactions. *Nat Commun* **12**, 6012 (2021).

8. Kumar, T. *et al.* A spatially resolved single-cell genomic atlas of the adult human breast. *Nature* **620**, 181–191 (2023).

9. Sibai, M. *et al.* The spatial landscape of Cancer Hallmarks reveals patterns of tumor ecology. *bioRxiv* 2022.06.18.496114 (2023) doi:10.1101/2022.06.18.496114.

10. Multimodal Analysis of Composition and Spatial Architecture in Human Squamous Cell Carcinoma. *Cell* **182**, 497-514.e22 (2020).

11. He, S. *et al.* Starfysh reveals heterogeneous spatial dynamics in the breast tumor microenvironment. 2022.11.21.517420 Preprint at https://doi.org/10.1101/2022.11.21.517420 (2022).

12. Wu, S. Z. *et al.* A single-cell and spatially resolved atlas of human breast cancers. *Nat Genet* **53**, 1334–1347 (2021).

13. Lengrand, J. *et al.* Pharmacological targeting of netrin-1 inhibits EMT in cancer. *Nature* **620**, 402–408 (2023).

14. Hu, J. *et al.* SpaGCN: Integrating gene expression, spatial location and histology to identify

spatial domains and spatially variable genes by graph convolutional network. *Nat Methods* **18**,

1342–1351 (2021).

15. Zhao, E. *et al.* Spatial transcriptomics at subspot resolution with BayesSpace. *Nat Biotechnol*

39, 1375–1384 (2021).

16. Walker, B. L. & Nie, Q. NeST: nested hierarchical structure identification in spatial

transcriptomic data. *Nat Commun* **14**, 6554 (2023).

17. Chitra, U. *et al.* Mapping the topography of spatial gene expression with interpretable deep

learning. *bioRxiv* 2023.10.10.561757 (2023) doi:10.1101/2023.10.10.561757.

18. Palla, G., Fischer, D. S., Regev, A. & Theis, F. J. Spatial components of molecular tissue

biology. *Nat Biotechnol* **40**, 308–318 (2022).

19. Seferbekova, Z., Lomakin, A., Yates, L. R. & Gerstung, M. Spatial biology of cancer evolution.

Nat Rev Genet **24**, 295–313 (2023).

20. Varrone, M., Tavernari, D., Santamaria-Martínez, A., Walsh, L. A. & Ciriello, G. CellCharter

reveals spatial cell niches associated with tissue remodeling and cell plasticity. *Nat Genet* 1–11

(2023) doi:10.1038/s41588-023-01588-4.

21. Moses, L. *et al.* Voyager: exploratory single-cell genomics data analysis with geospatial

statistics. 2023.07.20.549945 Preprint at https://doi.org/10.1101/2023.07.20.549945 (2023).

22. Bernstein, M. N. *et al.* Monkeybread: A Python toolkit for the analysis of cellular niches in single-

cell resolution spatial transcriptomics data. 2023.09.14.557736 Preprint at

https://doi.org/10.1101/2023.09.14.557736 (2023).

23. Palla, G. *et al.* Squidpy: a scalable framework for spatial omics analysis. *Nat Methods* **19**, 171–

178 (2022).

24. Fotheringham, A. S. & Wong, D. W. S. The Modifiable Areal Unit Problem in Multivariate

Statistical Analysis. *Environ Plan A* **23**, 1025–1044 (1991).

25. Dries, R. *et al.* Giotto: a toolbox for integrative analysis and visualization of spatial expression

data. *Genome Biology* **22**, 78 (2021).

26. He, Y. *et al.* ClusterMap for multi-scale clustering analysis of spatial gene expression. *Nat*

Commun **12**, 5909 (2021).

27. Li, Z. & Zhou, X. BASS: multi-scale and multi-sample analysis enables accurate cell type

clustering and spatial domain detection in spatial transcriptomic studies. *Genome Biol* **23**, 168

(2022).

28. Yang, J. *et al.* Guidelines and definitions for research on epithelial–mesenchymal transition. *Nat*

Rev Mol Cell Biol **21**, 341–352 (2020).

29. Barkley, D. *et al.* Cancer cell states recur across tumor types and form specific interactions with

the tumor microenvironment. *Nat Genet* **54**, 1192–1201 (2022).

30. Aggarwal, V., Montoya, C. A., Donnenberg, V. S. & Sant, S. Interplay between tumor

microenvironment and partial EMT as the driver of tumor progression. *iScience* **24**, 102113

(2021).

31. Pastushenko, I. & Blanpain, C. EMT Transition States during Tumor Progression and

Metastasis. *Trends in Cell Biology* **29**, 212–226 (2019).

32. Cook, D. P. & Vanderhyden, B. C. Context specificity of the EMT transcriptional response. *Nat*

Commun **11**, 2142 (2020).

33. Haerinck, J., Goossens, S. & Berx, G. The epithelial–mesenchymal plasticity landscape:

principles of design and mechanisms of regulation. *Nat Rev Genet* **24**, 590–609 (2023).

34. Malagoli Tagliazucchi, G., Wiecek, A. J., Withnell, E. & Secrier, M. Genomic and

microenvironmental heterogeneity shaping epithelial-to-mesenchymal trajectories in cancer. *Nat*

Commun **14**, 1–20 (2023).

35. Home Page - 10x Genomics. https://www.10xgenomics.com/.

36. Kleshchevnikov, V. *et al.* Cell2location maps fine-grained cell types in spatial transcriptomics.

Nat Biotechnol **40**, 661–671 (2022).

37. Elyanow, R., Zeira, R., Land, M. & Raphael, B. J. STARCH: copy number and clone inference

from spatial transcriptomics data. *Phys Biol* **18**, 035001 (2021).

38. Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection.

Cell Syst **1**, 417–425 (2015).

39. Vasaikar, S. V. *et al.* EMTome: a resource for pan-cancer analysis of epithelial-mesenchymal

transition genes and signatures. *Br J Cancer* **124**, 259–269 (2021).

40. Brown, M. S. *et al.* Phenotypic heterogeneity driven by plasticity of the intermediate EMT state

governs disease progression and metastasis in breast cancer. *Science Advances* **8**, eabj8002

(2022).

41. Muz, B., de la Puente, P., Azab, F. & Azab, A. K. The role of hypoxia in cancer progression,

angiogenesis, metastasis, and resistance to therapy. *Hypoxia (Auckl)* **3**, 83–92 (2015).

42. Weidemann, A. & Johnson, R. S. Biology of HIF-1α. *Cell Death Differ* **15**, 621–627 (2008).

43. Tam, S. Y., Wu, V. W. C. & Law, H. K. W. Hypoxia-Induced Epithelial-Mesenchymal Transition

in Cancers: HIF-1α and Beyond. *Front Oncol* **10**, 486 (2020).

44. Nishida, N., Yano, H., Nishida, T., Kamura, T. & Kojiro, M. Angiogenesis in Cancer. *Vasc Health*

Risk Manag **2**, 213–219 (2006).

45. Almagro, J., Messal, H. A., Elosegui-Artola, A., Rheenen, J. van & Behrens, A. Tissue

architecture in tumor initiation and progression. *Trends in Cancer* **8**, 494–505 (2022).

46. Saxena, K., Jolly, M. K. & Balamurugan, K. Hypoxia, partial EMT and collective migration:

Emerging culprits in metastasis. *Translational Oncology* **13**, 100845 (2020).

47. Biffi, G. *et al.* IL-1-induced JAK/STAT signaling is antagonized by TGF-β to shape CAF

heterogeneity in pancreatic ductal adenocarcinoma. *Cancer discovery* **9**, 282 (2019).

48. Ghahremanifard, P., Chanda, A., Bonni, S. & Bose, P. TGF-β Mediated Immune Evasion in

Cancer—Spotlight on Cancer-Associated Fibroblasts. *Cancers* **12**, 3650 (2020).

49. Principe, D. R., Timbers, K. E., Atia, L. G., Koch, R. M. & Rana, A. TGFβ Signaling in the

Pancreatic Tumor Microenvironment. *Cancers (Basel)* **13**, 5086 (2021).

50. Xu, J., Lamouille, S. & Derynck, R. TGF-β-induced epithelial to mesenchymal transition. *Cell*

Res **19**, 156–172 (2009).

51. myCAFs are better than yours: targeting myofibroblasts potentiates immunotherapy. *Trends in*

Cancer **9**, 1–2 (2023).

52. Öhlund, D. *et al.* Distinct populations of inflammatory fibroblasts and myofibroblasts in

pancreatic cancer. *Journal of Experimental Medicine* **214**, 579–596 (2017).

53. Amer, H. T., Stein, U. & El Tayebi, H. M. The Monocyte, a Maestro in the Tumor

Microenvironment (TME) of Breast Cancer. *Cancers* **14**, 5460 (2022).

54. Wu, S. Z. *et al.* Stromal cell diversity associated with immune evasion in human triple-negative

breast cancer. *The EMBO Journal* **39**, e104063 (2020).

55. Johnson, A. M. *et al.* Cancer Cell-Intrinsic Expression of MHC Class II Regulates the Immune

Microenvironment and Response to Anti-PD-1 Therapy in Lung Adenocarcinoma. *J Immunol*

204, 2295–2307 (2020).

56. Senosain, M.-F. *et al.* HLA-DR cancer cells expression correlates with T cell infiltration and is

enriched in lung adenocarcinoma with indolent behavior. *Sci Rep* **11**, 14424 (2021).

57. Sherman, T. D., Gao, T. & Fertig, E. J. CoGAPS 3: Bayesian non-negative matrix factorization

for single-cell analysis with asynchronous updates and sparse data structures. *BMC*

Bioinformatics **21**, 453 (2020).

58. Spatial and single-nucleus transcriptomic analysis of genetic and sporadic forms of Alzheimer's

Disease | bioRxiv. https://www.biorxiv.org/content/10.1101/2023.07.24.550282v1.

59. Ospina, O. E. *et al.* spatialGE: quantification and visualization of the tumor microenvironment

heterogeneity using spatial transcriptomics. *Bioinformatics* **38**, 2645–2647 (2022).

60. Emami Nejad, A. *et al.* The role of hypoxia in the tumor microenvironment and development of

cancer stem cell: a novel approach to developing treatment. *Cancer Cell Int* **21**, 62 (2021).

61. Wang, X., Zhang, W., Sun, X., Lin, Y. & Chen, W. Cancer-associated fibroblasts induce

epithelial-mesenchymal transition through secreted cytokines in endometrial cancer cells. *Oncol*

Lett **15**, 5694–5702 (2018).

62. Jia, C. *et al.* Cancer-associated Fibroblasts induce epithelial-mesenchymal transition via the

Transglutaminase 2-dependent IL-6/IL6R/STAT3 axis in Hepatocellular Carcinoma. *Int J Biol*

Sci **16**, 2542–2558 (2020).

63. Tyler, M. & Tirosh, I. Decoupling epithelial-mesenchymal transitions from stromal profiles by

integrative expression analysis. *Nat Commun* **12**, 2592 (2021).

64. Li, X., Chen, L., Peng, X. & Zhan, X. Progress of tumor-associated macrophages in the

epithelial-mesenchymal transition of tumor. *Front Oncol* **12**, 911410 (2022).

65. Brabletz, S., Schuhwerk, H., Brabletz, T. & Stemmler, M. P. Dynamic EMT: a multi-tool for

tumor progression. *The EMBO Journal* **40**, e108647 (2021).

66. Johansson, J., Tabor, V., Wikell, A., Jalkanen, S. & Fuxe, J. TGF-β1-Induced Epithelial–

Mesenchymal Transition Promotes Monocyte/Macrophage Properties in Breast Cancer Cells.

Frontiers in Oncology **5**, (2015).

67. Kmieciak, M., Knutson, K. L., Dumur, C. I. & Manjili, M. H. HER-2/neu antigen loss and relapse

of mammary carcinoma are actively induced by T cell-mediated anti-tumor immune responses.

Eur J Immunol **37**, 675–685 (2007).

68. Santisteban, M. *et al.* Immune-induced epithelial to mesenchymal transition in vivo generates

breast cancer stem cells. *Cancer Res* **69**, 2887–2895 (2009).

69. Galsky, M. D. *et al.* 850PD - Epithelial-mesenchymal transition (EMT), T cell infiltration, and

outcomes with nivolumab (nivo) in urothelial cancer (UC). *Annals of Oncology* **28**, v297 (2017).

70. Chockley, P. J. *et al.* Epithelial-mesenchymal transition leads to NK cell–mediated metastasis-

specific immunosurveillance in lung cancer. *J Clin Invest* **128**, 1384–1396.

71. Datar, I. & Schalper, K. A. Epithelial–Mesenchymal Transition and Immune Evasion during Lung

Cancer Progression: The Chicken or the Egg? *Clin Cancer Res* **22**, 3422–3424 (2016).

72. Terry, S. *et al.* New insights into the role of EMT in tumor immune escape. *Mol Oncol* **11**, 824–

846 (2017).

73. Tsai, J. H., Donaher, J. L., Murphy, D. A., Chau, S. & Yang, J. Spatiotemporal regulation of

epithelial-mesenchymal transition is essential for squamous cell carcinoma metastasis. *Cancer*

Cell **22**, 725–736 (2012).

74. Cheng, Y.-C. *et al.* Reconstruction of single cell lineage trajectories and identification of diversity

in fates during the epithelial-to-mesenchymal transition. *bioRxiv* 2023.09.19.558325 (2023)

doi:10.1101/2023.09.19.558325.

75. Sinha, D., Saha, P., Samanta, A. & Bishayee, A. Emerging Concepts of Hybrid Epithelial-to-

Mesenchymal Transition in Cancer Progression. *Biomolecules* **10**, 1561 (2020).

76. Subhadarshini, S., Markus, J., Sahoo, S. & Jolly, M. K. Dynamics of Epithelial–Mesenchymal

Plasticity: What Have Single-Cell Investigations Elucidated So Far? *ACS Omega* **8**, 11665–

11673 (2023).

77. Zhang, Y. *et al.* Hypoxia in Breast Cancer—Scientific Translation to Therapeutic and Diagnostic

Clinical Applications. *Frontiers in Oncology* **11**, (2021).

78. Fu, Z., Mowday, A. M., Smaill, J. B., Hermans, I. F. & Patterson, A. V. Tumour Hypoxia-

Mediated Immunosuppression: Mechanisms and Therapeutic Approaches to Improve Cancer

Immunotherapy. *Cells* **10**, 1006 (2021).

79. Lee, C.-T., Mace, T. & Repasky, E. A. Hypoxia-Driven Immunosuppression: A new reason to

use thermal therapy in the treatment of cancer? *Int J Hyperthermia* **26**, 232–246 (2010).

80. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data

analysis. *Genome Biology* **19**, 15 (2018).

81. Stein-O'Brien, G. L. *et al.* Decomposing Cell Identity for Transfer Learning across Cellular

Measurements, Platforms, Tissues, and Species. *Cell Systems* **8**, 395-411.e8 (2019).

82. Ayers, M. *et al.* IFN-γ-related mRNA profile predicts clinical response to PD-1 blockade. *J Clin*

Invest **127**, 2930–2940 (2017).

83. Nielsen, T. O. *et al.* A Comparison of PAM50 Intrinsic Subtyping with Immunohistochemistry

and Clinical Prognostic Factors in Tamoxifen-Treated Estrogen Receptor–Positive Breast

Cancer. *Clinical Cancer Research* **16**, 5222–5232 (2010).

84. Proceedings of the Python in Science Conference (SciPy): Exploring Network Structure,

Dynamics,   and   Function   using   NetworkX.

https://conference.scipy.org/proceedings/SciPy2008/paper_2/.

85. Rey, S. J. & Anselin, L. PySAL: A Python library of spatial analytical methods. *The Review of*

regional studies **37**, 5–27 (2007).

1032