Unveiling *Cryptosporidium parvum* Sporozoite-Derived Extracellular Vesicles: Profiling, Origin, and Protein Composition

Lucia Bertuccini[1†], Zaira Boussadia[2†], Anna Maria Salzano[3†], Ilaria Vanni[4†], Ilaria Passerò[5], Emanuela Nocita[5], Andrea Scaloni[3], Massimo Sanchez[1], Massimo Sargiacomo[6], Maria Luisa Fiani[6] and Fabio Tosini[5]*

[1] Core Facilities, Istituto Superiore di Sanità, Rome, Italy.

[2] National Center for Drug Research and Evaluation, Istituto Superiore di Sanità, Rome, Italy.

[3] Proteomics, Metabolomics and Mass Spectrometry laboratory, ISPAAM Consiglio Nazionale delle Ricerche, Portici, Italy.

[4] Department of Food Safety, Nutrition and Veterinary Public Health (SANV), Istituto Superiore di Sanità, Rome, Italy.

[5] Department of Infectious Diseases, Istituto Superiore di Sanità, 00161 Rome, Italy.

[6] National Center for Global Health, Istituto Superiore di Sanità, 00161 Rome, Italy.

[†] These authors contributed equally to this work and share first authorship.

* Correspondence:

Fabio Tosini

fabio.tosini@iss.it

Keywords: *Cryptosporidium*, sporozoite, excystation, rhomboid, extracellular vesicles, exosome, aspartyl protease, GP60.

Abstract

Cryptosporidium parvum is a common cause of a zoonotic disease and a main cause of diarrhea in

newborns around the world. Effective drugs or vaccines are still lacking. Oocyst is the infective

form of the parasite; after the ingestion the oocyst excysts and releases four sporozoites into the

intestine that rapidly attack the enterocytes. The membrane protein CpRom1 is a large rhomboid

protease that is expressed by sporozoites and recognized as antigen by the host immune system. In

this study, we observed the release of CpRom1 with extracellular vesicles (EVs) not previously

described. To investigate this phenomenon, we isolated and resolved EVs from the excystation

medium by differential ultracentrifugation. By fluorescence flow cytometry and transmission

electron microscopy (TEM), we identified two types of sporozoite-derived vesicles: large

extracellular vesicles (LEVs) and small extracellular vesicles (SEVs), having different a mean size

of 150 nm and 60 nm, respectively. Immunodetection experiments proved the occurrence of

CpRom1 and the Golgi protein CpGRASP in LEVs, while immune-electron microscopy

experiments demonstrated localization of CpRom1 on LEVs surface.

TEM and scanning electron microscopy (SEM) showed the generation of LEVs by budding of the

outer membrane of sporozoites; conversely, the origin of SEVs remained uncertain.

Differences between LEVs and SEVs were observed for protein composition as proved by the

corresponding electrophoretic profiles. Indeed, a dedicated proteomic analysis identified 5 proteins

unique to LEVs composition and 16 proteins unique to SEVs. Overall, 60 proteins were identified

in the proteome of both types of vesicles and most of these proteins (48 in number) were already

identified in the molecular cargo of extracellular vesicles from other organisms. Noteworthy, we

identified 12 proteins unique to *Cryptosporidium* spp. that had never been associated with EVs.

This last group included the immunodominant parasite antigen glycoprotein GP60, which is one of

the most abundant proteins in both LEVs and SEVs.

1   Introduction

Cryptosporidium genus consists of 38 species of obligate parasites of the phylum Apicomplexa that infect vertebrates of many species (Feng et al., 2018). Several species of *Cryptosporidium* are important pathogens affecting mammals' intestine and cause cryptosporidiosis, a diarrhoeal disease that usually lasts several days. Two closely related species, namely *Cryptosporidium parvum* and *Cryptosporidium hominis*, are responsible for most cases of human cryptosporidiosis with watery diarrhoea as the main symptom (Bouzid et al., 2013). In immunocompetent adults, cryptosporidiosis usually has a limited duration (7-10 days), but differently, this infection poses a serious risk to people with immunodeficiency, who may have a long-term and life-threatening illness due to severe dehydration (Checkley et al., 2015). Remarkably, *C. parvum* and *C. hominis* are also responsible for most cases of neonatal cryptosporidiosis, a major cause of diarrhoea in children in endemic regions, where transmission is favoured by poor hygiene conditions (Kotloff et al., 2013). Transmission of the parasite occurs via the oral-fecal route either directly (animal to human or human to human) or indirectly through contaminated food and, above all, water (Xiao and Feng, 2017).

Oocyst is the parasitic stage released with feces by infected subjects, and it is responsible for the transmission of the infection to other subjects. Oocysts are characterized by a thick "shell", the oocyst wall, constituted by polysaccharides, lipids and peculiar proteins defined as *Cryptosporidium* oocyst wall proteins (COWPs), which make oocysts particularly resistant to environmental and chemical stresses (Templeton et al., 2004). Noteworthy, oocysts are also resistant to treatments to disinfect drinkable water; this is a significant risk factor for water plant contamination, which can cause large outbreaks with thousands of infected people (MacKenzie et al., 1994). Despite the relevance of *Cryptosporidium* for human health, effective drug therapies are still lacking, and vaccines have not yet been developed.

Contrarily to other apicomplexan parasites such as Plasmodium spp. and Toxoplasma

gondii, Cryptospordium completes its life cycle in a single host (monoxenous). Excystation is the

first stage of the infectious process and consists of rupture of the oocyst wall and the egress of four

sporozoites; this is triggered first the passage of the oocyst in the stomach then by its subsequent

contact with the bile salts in the small intestine (Fayer and Xiao, 2007). The released sporozoites

adhere to the luminal side of enterocytes in the small intestine. The close contact between the

parasite and the host cell membrane induces the formation of a parasitophorous vacuole (PV). PV

protrudes toward the intestinal lumen, but the intracellular stages of the parasite are completely

wrapped by the host cell membrane. The motile stages of *Cryptosporidium* (*i. e.*, sporozoites and

merozoites), like other apicomplexans, are characterized by the apical complex at the anterior end

of the cell. The apical complex includes specialized subcellular organelles, such as rhoptry,

numerous micronemes and various dense granules, which are vesicular structures specialized for the

parasitic function. These organelles discharge their content sequentially from the egress of

sporozoites during the attachment and invasion of the host cell, and until the formation of a

parasitophorous vacuole at the luminal side of the cell (Guerin et al., 2023).

Oocyst excystation as well as the organelle discharge can be easily replicated *in vitro* even in

the absence of host cells. In fact, free sporozoites maintained at 37 °C in an adequate medium,

release most of their organelle content within two hours (Chen et al., 2004). Free sporozoites,

among other things, carry antigens that can be successfully counterbalanced by circulating host

antibodies (Tosini et al., 2019). By investigating the antigenic proteins involved in the preliminary

phases of the excystation, we identified a rhomboid protease, namely CpRom1, which occurs in

sporozoites (Trasarti et al., 2007). Rhomboids are ubiquitous serine-proteases that are entirely

embedded in the lipid bilayer of cell membranes (Kühnle et al., 2019). Rhomboids in Apicomplexa

play a unique role before and during the invasion of the host cell by cleaving adhesins, surface

proteins exposed in the motile stages such as sporozoites, during their path towards the host cell and

in the phase of penetration through the host cell membrane (Carruthers and Blackman 2005; Dowse

et al., 2008).

In this study, we observed that CpRom1 is released by sporozoites as bound to extracellular vesicles

(EVs). This coincidence led us to investigate the release of microvesicles by sporozoites.

Nowadays, EVs are recognized as fundamental elements for intercellular communication both in

unicellular and in multicellular organisms (Doyle and Wang, 2019). They also serve as effectors of

intercellular communication in protozoan pathogens, mediating interactions between parasites and

between parasites and host cells (Wang et al., 2022). Here we describe that *C. parvum* sporozoites

release two types of EVs that, based on their size, we have here classified as large extracellular

vesicles (LEVs) and small extracellular vesicles (SEVs). These vesicles exhibited significant

differences in their protein composition, but also presented common molecules including one of the

most antigenic *C. parvum* component, namely glycoprotein GP60 (Allison et al., 2011), also known

as GP40/GP15.

2   Materials and Methods

2.1   Parasites and excystation procedure

Fresh *C. parvum* oocysts (Iowa strain) were supplied by Bunch Grass Farm (Deary, Idaho USA),

stored at 4 °C in PBS with penicillin (1000 U.I./ml) and streptomycin (1 mg/ml). The excystation

procedure was performed as follows: aliquots of 1×10^7 oocysts per ml were pelleted at 2000 rpm ,

for 5 min, resuspended in 1 ml of 10 mM HCl, and then incubated at 37 °C for 10 min. Oocysts

were pelleted again as above and resuspended in 1 ml of excystation medium (D-MEM containing 2

mM sodium taurocholate); they were maintained at 15 °C, for 10 min, and then moved to 37 °C to

induce the excystation. Excystation mixtures were sampled at various times after the induction.

2.2   Expression of recombinant CpRom1 and CpGRASP and production of the

corresponding antisera

The CpRom1 and CpGRASP coding seqence (monoexonic genes without introns) were directly

amplified from *C. parvum* genomic DNA (Iowa strain).

2.2.1   Cloning, expression and purification of recombinant CpRom1

The recombinant 6His-CpRom1 was obtained as follows: the CpRom1 coding sequence was

amplified with AAC<u>GAGCTC</u>GATATGTCCGATTTTGTTTTCA as the forward primer including

the   *Sac*I   (underlined)   restriction   site,   and

TCCC<u>CCCGGG</u>TCATCAAGAAAAATCATATCCAAATA as the reverse primer including the

*Sma*I (underlined) restriction site. Amplification with 2X Phusion Flash High fidelity PCR Master

Mix (Finnzymes) was performed using 80 ng of genomic DNA as template as follows: 95 °C for 5

min, 35 cycles of 94 °C for 15 sec, 50 °C for 15 sec, 72 °C for 4 min, and a final extension at 72 °C

for 10 min in a Veriti 96 well thermal cycler (Applied Biosystem). Amplicons were purified with

QIAquick PCR Purification Kit (Qiagen GmbH, Hilden, Germany) and digested with *Sac*I and

*Sma*I restriction enzymes (New England Biolabs). This fragment was then ligated in the *Sac*I and

*Sma*I digested pQE80 vector (Qiagen GmbH, Hilden, Germany) using the Quick Ligase kit (New

England Biolabs), and the ligation mix was used to transform the *Escherichia coli* M15 strain.

Positive clones were selected on LB agar plates with 100 µg/ml ampicillin and 25 µg/ml kanamycin

by PCR screening; recombinant plasmid with recombinant CpRom1 ORF was sequenced to check

the fusion with the histidine-tag coding sequences at its N-terminus. For 6His-CpRom1 purification,

20 ml of an overnight culture of recombinant bacteria were inoculated in 1 l of LB with 100 µg/ml

ampicillin and 25 µg/ml kanamycin, and cultured at 37 °C, with vigorous shacking, until a 0.6 OD

was reached; then, 1 mM IPTG was added to the culture and the bacterial growth continued for

additional 3 h. Bacteria were pelleted first at 4,000 rpm for 10 min, resuspended in 100 ml cold PBS

and centrifuged again at 6,000 rpm for 20 min. Then, the pellet was resuspended in 20 ml of

denaturing buffer A (100 mM NaH$_2$PO4, 10 mM Tris-HCl, 6 M guanidine-HCl, pH 8.0), and stirred

at 25 °C, overnight; the corresponding lysate was clarified by centrifugation at 10,000 rpm, for 30

min. Then, 5 ml of 50% Ni-NTA resin was added and the mix was gently stirred for 60 min, at 25

°C; the slurry was loaded slowly on a 10 ml column to pack the resin and then washed with 40 ml

(8 x 5 ml wash) of buffer C (100 mM NaH$_2$PO4, 10 mM Tris-HCl, 8 M urea, pH 6.3). Purified

protein was eluted with 4 aliquots of 2.5 ml of buffer D (100 mM NaH$_2$PO4, 10 mM Tris-HCl, 8 M

urea, pH 5.9) and with 4 aliquots of 2.5 ml of buffer E (100 mM NaH$_2$PO4, 10 mM Tris-HCl, 8 M

urea, pH 4.5). Pooled fractions from buffer D and pooled fractions from buffer E were extensively

dialyzed against PBS with increasing concentrations of glycerol (from 10% to 50%) in Slide-A-

Lizer^{tm} (ThermoFisher) dialysis cassettes with a cut-off of 10 kDa.

2.2.2   Cloning, expression and purification of recombinant CpGRASP

To clone the CpGRASP coding sequence, the following forward primer

CC<u>GGATCC</u>GGAGGTGCGCAAACCAAAC including *Bam*HI restriction site (underlined) and

reverse primer CT<u>CCCGGG</u>TTATATTTCTCCTTGGTCTGTG including *Sma*I restriction site

(underlined) were used. PCR amplification was performed with Hot Star Taq Plus (Qiagen) using 5

ng of genomic DNA as template and these PCR conditions: 95 °C for 5 min, 35 cycles of 94 °C for

15 sec, 55 °C for 15 sec, 72 °C for 4 min, and a final extension at 72 °C for 10 min. Reactions were

conducted in a Veriti 96 well thermal cycler (Applied Biosystem). The amplified fragment was

digested with *Bam*HI and *Sma*I restriction enzymes and ligated in pQE80 vector (Qiagen) digested

with the same enzymes, using Quick Ligase kit (New England Biolabs). Ligation was used to

transform *E. coli* M15 host strain, positive colonies were selected by PCR screening, and amplicons

were sequenced to verify the fusion of histidine tag at the 5'-end of the inserted CpGRASP

sequence. The histidine-tagged CpGRasp (6His-CpGRASP) was then purified as above.

2.2.3   Production of specific mouse antisera for 6His-CpRom1 and 6His-CpGRASP

To produce specific antisera for 6His-CpRom1 and 6His-CpGRASP, Balb-C mice were immunized

with the following schedule: i) 100 mg of protein plus complete Freund adjuvant as first inoculum;

ii) 100 mg of protein plus incomplete Freund adjuvant as second inoculum, 30 days after the first

inoculum; iii) a final inoculum with 100 mg of protein in PBS, 30 days after the second inoculum.

Mice were bled 15 days after the last inoculum and 150-300 ml of serum was obtained from each

mouse.

2.3   Isolation of extracellular vesicles from excystation medium by differential

centrifugation

Sporozoite microvesicles were prepared from the excystation medium of fresh oocyst aliquots after

2 h incubation at 37 °C; excystation was blocked by placing samples on ice for 5 min. All

subsequent manipulations were performed on ice and centrifugation steps were performed at 4 °C.

Excystation medium was centrifuged at 2000 rpm, for 20 min, to remove most of oocysts and

sporozoites and the resulting supernatant was centrifuged again at 5000 rpm, for 10 min, to remove

residual oocysts, sporozoites, and excystation debris. To isolate EVs, supernatant was transferred to

1.5 ml tubes (Eppendorf), centrifuged for 20 min at 10,000 g and the pellet containing LEVs

resuspended in 100 μl of ice-cold PBS. To isolate SEVs, the supernatant was transferred to new

tubes (Beckman Coulter) and ultracentrifuged on a TLA 120.2 rotor (Beckman Coulter) at 100,000

g for 3 hours. The pellet was resuspended in 1.2 ml of ice-cold PBS and recentrifuged as above. The

final pellet containing SEVs was resuspended in 100 μl of ice-cold PBS. All the procedure is

schematized in Figure 1.

2.4   Western blot analysis

Proteins in the excystation medium and supernatants after ultracentrifugation were precipitated with

10% ice-cold trichloroacetic acid (TCA)/0.015% (w/v) sodium deoxycholate (DOC). After 10 min

on ice, samples were microfuged and the pellets were washed twice with 0.5 ml of ice-cold acetone,

air dried, resuspended in 50 μl of lysis buffer (1% w/v SDS, 1% v/v Triton X-100, 0.5% w/v DOC,

10 mM 1,4-dithiothreitol (DTT), 1 mM EDTA) containing 1 μl/ml of Protease Inhibitor Cocktail

(Sigma-Aldrich), and incubated for 5 min at 70 °C in ThermoMixer® (Eppendorf).

Pellets after centrifugation (LEVs) and ultracentrifugation (SEVs) were resuspended in 50 µl of

lysis buffer containing 1 µl/ml of Protease Inhibitor Cocktail (Sigma-Aldrich) as above. Samples

were boiled in Laemmli sample buffer before separation by SDS-PAGE on 4-20% TGX™ precast

gels (Bio-Rad, Hercules, CA, USA). Gels were transferred to nitrocellulose membranes (Bio-Rad),

which were then blocked in PBS with 5% non-fat milk in 0.1% Tween-20. Blots were incubated

with primary antibodies, at 4 °C, overnight. For recombinant 6His-CpRom1 and 6His-CpGRASP,

blots were incubated with mouse monoclonal RGS-His antibody (Qiagen) diluted 1:1,000. Blots for

CpRom1 were probed with mouse anti-CpRom1 serum 1 diluted 1:250, while blots for CpGRASP

were probed with mouse anti-CpGRASP serum diluted 1:100. For all blots incubation with

secondary antibodies was conducted for 1 h, at 25 °C, with goat anti-mouse IgG-HRP conjugate

(Bio-Rad, Hercules, CA USA) diluted 1:3,000. Detection of proteins was performed using Pierce

ECL substrate (ThermoFisher) and bands were visualized with a ChemiDoc MP Imaging System

(Bio-Rad, Hercules, CA, USA).

2.5　Ultrastructural analysis and immunolocalisation of extracellular vesicles by

transmission electron microscopy

For negative staining in transmission electron microscopy (TEM), samples of LEVs and SEVs were

suspended in 50 µl of 2.5% w/v glutaraldehyde in 0.1 M sodium cacodylate and incubated at 4 °C,

overnight. Samples (20 µl) were deposited by successive applications on carbon-coated grids for

electron microscopy; they were left to adsorb for 20 min, and the excess fluid was blotted with filter

paper. A 2% w/v phosphotungstic acid contrasting solution (5 µl) was added on grids and air dried.

Samples were observed by a PHILIPS Morgagni 268 TEM (FEI - Thermo Fisher) (Thery et al.,

2006). For immunolocalization, LEVs and SEVs pellets were suspended in 100 µl of PBS, and

aliquots of 20 µl were adsorbed on carbon-coated grids for electron microscopy, for 20 min.

Samples were air dried and transferred to a drop of 2% w/v paraformaldehyde, for 5 min. After two

washes with PBS, the grids were floated on PBS drops containing 0.1 M glycine for 30 min,

washed with PBS, blocked with PBS containing 5% goat serum and 1% w/v BSA, for 30 min, and

washed with PBS containing 0.05% w/v TWEEN 20 and 1% w/v BSA (PBS/BSA/TW). Then grids

were floated on rabbit polyclonal anti-ID2 serum (1:10) in PBS/BSA/TW, for 1 h, at 25 °C, rinsed

in the same buffer, incubated on 10 nm gold-conjugated goat anti-rabbit IgG (SIGMA) (1:50), for 1

h, rinsed again in buffer PBS/BSA/TW and treated with 1% glutaraldehyde, for 5 min. Finally, the

samples were contrasted and embedded in a mixture 1:9 mixture of 4% uranyl acetate and 2%

methyl cellulose respectively for 10 min on ice. The samples were air dried and analyzed by a

PHILIPS Morgagni 268 TEM (FEI - Thermo Fisher) (Thery et al., 2006).

For ultrathin sections, 1 ml of excystation mixture (taken 30 min after induction) was loaded onto a

1 ml syringe and filtered with a 5 μm disposable filter Millex-SV (Millipore) to remove residual

non-excysted and empty oocysts. Purified sporozoites were fixed in 2.5% w/v glutaraldehyde, 2%

paraformaldehyde, 2 mM $CaCl_2$ in 0.1 M sodium cacodylate, pH 7.4, and processed according to

Perry and Gilbert (1979). Parasites were washed in cacodylate buffer and post-fixed with 1% OsO_4

in 0.1 M sodium cacodylate, for 1 h, at 25 °C; then, they were treated with 1% tannic acid in 0.05 M

cacodylate buffer, for 30 min, and rinsed in 1% sodium sulphate, 0.05 M cacodylate, for 10 min.

Post-fixed specimens were washed, dehydrated through a graded series of ethanol solutions (30 to

100% ethanol) and embedded in Agar 100 (Agar Scientific Ltd, UK). Ultrathin sections, obtained

by an UC6 ultramicrotome (Leica), were stained with uranyl acetate and Reynolds lead citrate and

examined at 100 kV with a PHILIPS EM208S TEM (FEI - ThermoFisher) equipped with the

Megaview III SIS camera (Olympus).

2.6   Scanning electron microscopy

1×10^7 *C. parvum* oocysts (see above) were induced to excystation. At various times (from 0 to 30

min), the samples were fixed with 2.5% glutaraldehyde in 0.1 M sodium cacodylate buffer and 30

μl were left to adhere to polylysine-treated round glass coverslips (Ø 10mm), for 1 h, at 25 °C.

Samples were processed for scanning electron microscopy (SEM) as previously described with

slight modifications (Shively and Miller, 2009). Briefly, samples were post-fixed with 1% OsO_4 in

0.1 M sodium cacodylate, for 1 h, at 25 °C, and were dehydrated through a graded series of ethanol

solutions (from 30% to 100%). Then, absolute ethanol was gradually replaced by a 1:1 solution of

hexamethyldisilazane (HMDS) and absolute ethanol, for 30 min, and successively by pure HMDS,

for 1 h. Finally, HMDS was completely removed, and samples were left to dry in a desiccator, at

room temperature, for 2 h, overnight. Then dried samples were mounted on stubs, carbon coated

and analyzed by a FE-SEM Quanta Inspect F (FEI, Thermo Fisher Scientific).

2.7   Fluorescent labelling of extracellular vesicles and flow cytometric analysis

Fluorescent labelling of proteins of LEVs and SEVs was conducted as follows: pellets were

resuspended in 50 µl PBS containing 10 µM Alexa Fluor 647 NHS Ester (Life Technologies) and

incubated for 30 min, at 25 °C. Then pellets were resuspended again in 50 µl PBS containing 10 µM

CFDA-SE (CFSE) (Life Technologies), for 30 min at room temperature. Reactions were stopped by

adding 2 µl of 100 mM L-glutamine. Fluorescent LEVs and SEVs from the corresponding pellets

were analysed by flow cytometry (FC) on a Gallios Flow Cytometer (Beckman Coulter) using an

optimised procedure, as previously described (Coscia et al., 2016). Briefly, the instrument was set

using the Flow Cytometry Sub-micron Particle Size Reference Kit providing green-fluorescent

beads of different sizes (Invitrogen™) to establish the correct threshold value to apply on the

525/40 nm fluorescent channel (FL1). This procedure allows to create gates of the following

dimensions: ≤200 nm, 500 nm and 1 µm. Sample analysis was performed by mixing 5 µl of

fluorescent vesicles with 20 µl of Flow-Count Fluorospheres (Beckman Coulter) in a final volume

of 200 µl of PBS to determine absolute counts. Fluorescent populations were analysed by plotting

fluorescence at 525/40 nm (FL1) versus log scale side scatter (SSarea). Sample acquisition was

stopped at 2,000 Flow-Count Fluorospheres events. The total number of fluorescent vesicles was

established according to the formula: $x = (((y \times a)/b)/c) \times d$, where y = events counted at 2,000

counting beads; a = number of counting beads in the sample; b = number of counting beads

registered (2,000); c = volume of sample analysed; and d = total volume of exosome preparation.

Kaluza Software v. 2.0 (Beckman Coulter) was used for FC analysis.

2.8   Iodixanol gradient separation of extracellular vesicles

Freshly labelled fluorescent LEVs and SEVs from 1×10^7 oocsts were diluted in 0.3 ml of PBS.

Then, fluorescent samples were mixed with 1 ml of 60% iodixanol solution (OptiprepTM, Sigma-

Aldrich), overlaid with 0.5 mL of 40%, 0.5 mL of 30% and 1.8 mL 10% of iodixanol solutions and

floated into the gradient by ultracentrifugation in a SW60Ti rotor (Beckman) at 192,000 x g, for 18

hours, stopping without brake. After centrifugation, 12 fractions of 330 µl were collected from the

top of the tube, diluted 40- to 80-fold with PBS and analysed by FC. Fraction densities were

determined by refractometry. Gradient solutions were produced from the working solution (WS) by

dilution with the HM solution. WS is prepared by mixing 5 volumes of OptiPrep™ with 1 volume

of 0.25 M sucrose, 6mM EDTA, 60 mM Tris-HCl, pH 7.4; while HM solution contained 0.25 M

sucrose, 1 mM EDTA, 10 mM Tris-HCl, pH 7.4.

Gradient fractions of fluorescent LEVs and SEVs were analysed by FC on a Gallios Flow

Cytometer (Beckman Coulter) using an optimised procedure, as previously described (Coscia et al.,

2016).

2.9   Proteomic analysis

Proteins from LEVs and SEVs were separated on a precast 4-15 % SDS-PAGE (Bio-Rad, Hercules,

CA USA) and stained with Coomassie G-250. Gel lanes were cut into 9 slices and each one was *in-*

gel reduced, alkylated with iodoacetamide and digested with trypsin, as previously reported

(Salzano et al., 2013). Peptide mixtures were then desalted by µZipTipC18 tips (Millipore) before

nano-liquid chromatography-electrospray ionization-Orbitrap tandem mass spectrometry (nLC-ESI-

Q-Orbitrap-MS/MS) analysis, which was performed on a Q-Exactive Plus mass spectrometer

equipped with UltiMate 3000 HPLC RSLC nano system (Thermo Fischer Scientific, USA).

Peptides were separated on an Easy C18 column (100×0.075 mm, 3 µm) at a flow rate of 300

nl/min using a linear gradient from 5 to 40% acetonitrile in 0.1% formic acid, over 60 min. Mass

spectra were acquired at nominal resolution 70,000 in the range m/z 350-1500, and data-dependent

automatic MS/MS acquisition was applied to the ten most abundant ions (Top10), enabling dynamic

exclusion with repeat count 1 and exclusion duration 30 s. The mass isolation window and the

collision energy for peptide fragmentations were set to *m/z* 1.2 and 32%, respectively. Raw data

from nLC-ESI-Q-Orbitrap-MS/MS analysis were searched by MASCOT v2.6.1, (Matrix Science,

UK) node within Proteome Discoverer suite, against a *C. parvum* (strain Iowa II) database of

protein sequences (4076 sequences) retrieved from CryptoDB site (https://cryptodb.org), and results

were merged into a single mgf file to obtain proteins identified in LEVs and SEVs fractions. The

following parameters were used for protein identification: mass tolerance values of 20 ppm and

0.05 Da for precursor and fragment ions, respectively; trypsin as proteolytic enzyme with maximum

missed-cleavage sites of 2; Cys carbamidomethylation as fixed modification, Met oxidation,

Asn/Gln deamidation and pyroglutamate formation from Gln/Glu, as variable modifications. A

significance threshold of $p<0.05$ was set for protein identification, and molecular candidates with at

least 2 significantly matched peptide sequences and protein Mascot score >50 were further

considered for definitive assignment, after manual spectra visualization and verification.

2.10   Bioinformatic analyis

The sequence of CpRom1 (CryptoDB: cgd6_760) and CpGRASP (CryptoDB: cgd7_340) were

downloaded at CryptoDB site (https://cryptodb.org). Similarity searches with DNA and protein

sequences were conducted on non-redundant GenBank databases using the BLAST program

(https://blast.ncbi.nlm.nih.gov/Blast.cgi). The prediction of transmembrane domains was performed

at   TMHMM-2.0   (http://www.cbs.dtu.dk/services/TMHMM/)   and   at   Phobius

(https://phobius.sbc.su.se/index.html). Prediction of structural and functional domains was

performed   at   Pfam   (http://pfam.xfam.org)   and   Prosite   (https://prosite.expasy.org).   The

bidimensional representation of CpRom1 in Figure 2 was realized with TMRPres2D software

(http://bioinformatics.biol.uoa.gr/TMRPres2D/download.jsp). Searches for potential cleavage sites

by proteases were performed at PeptideCutter (https://www.expasy.org/resources/peptidecutter) and

at ProP 1.0 server (http://www.cbs.dtu.dk/services/ProP/) for Arg and Lys propeptide cleavage sites

(furin-like). Searches for potential GPI-anchor were performed at GPI Modification Site Prediction

(https://mendel.imp.ac.at/gpi/gpi_server.html). Subcellular localization was predicted by DeepLoc-

1.0 (http://www.cbs.dtu.dk/services/DeepLoc/) and was reported only when the probability was

higher than 50%. Protein-protein interaction networks were obtained with STRING v.12.0

(https://string-db.org). Searches for proteins associated with extracellular vesicles were performed

in the following databases: ExoCarta (http://www.exocarta.org) and Vesiclepedia

(http://www.microvesicles.org).

3 Results

3.1 Characterization of CpRom1 expressed in oocysts and sporozoites

The CpRom1 rhomboid has been previously identified starting from the 46 amino acid-long ID2 peptide (Trasarti et al., 2007). A prediction based on the protein sequence showed that CpRom1 consists of 990 residues with a theoretical molecular mass of 109.3 kDa; this makes CpRom1 the largest rhomboid ever described. Computational predictions showed that CpRom1 has a large N-terminal portion of approximately 500 amino acids, which includes the ID2 peptide (Figure 2). This large portion of the protein is exposed towards the extracellular environment and does not show any recognizable domain (figure 2). A well-distinguishable rhomboid domain, which consists of six transmembrane helices surrounding the proteolytic site, is contained in the final 450 amino acids at the C-terminus.

To identify the native CpRom1 rhomboid, we prepared a mouse polyclonal serum reacting with the full-length protein. To this end, a fusion protein with six histidine-tag (6His-CpRom1) was expressed in bacteria, purified by Ni-NTA chromatography and used to immunize Balb-C mice. The polyclonal mouse antiserum was used in Western blot experiments on sporozoite lysates, in which a band of approximately 110 kDa was recognized, in agreement with an expected molecular mass of 109.3 kDa (Figure 3). However, the serum also detected a smaller band migrating at about 75 kDa, which might represent a cleaved form of the 110 kDa protein. These CpRom1 forms were present in resting oocyst at time 0 of excystation (Figure. 3A), with a similar amount of the 110 kDa form in unexcysted oocyst and sporozoites, whereas a lower amount of the 75 kDa species was detected after excystation. Probably, the latter protein form is quickly degraded after excystation.–A significant amount of CpRom1 was also detectable in the supernatant of excystation medium after eliminating sporozoites and residual oocysts. In fact, TCA precipitates of excystation medium after 30 min of incubation, probed with anti-CpRom1 serum revealed a distinctive band of 110 kDa (Figure 3B). Given the presence of six transmembrane domain in CpRom,1 which eventually

hampers its release in the excystation medium, we hypothesized that CpRom1 is released in

association with membranous vesicles.

3.2   CpRom1 is associated with released extracellular vesicles during the excystation

To test this the hypothesis reported above, we developed an ultracentrifugation protocol to separate

extracellular vesicle populations from the excystation medium, after preventive removal of oocyst-

sporozoite residues. A schematic representation of the procedure is shown in Figure 1. Depending

on centrifugation conditions, two types of *C. parvum* EVs were selectively recovered that were

tentatively classified as large extracellular vesicles (LEVs) and small extracellular vesicles (SEVs).

Protein extracts from LEVs and SEVs were then assayed for the presence of CpRom1 by dedicated

immunoblotting. Results showed that a protein band of 110 kDa was observed in the LEVs extracts,

but not in the SEVs counterparts (Figure 4A).

To demonstrate that CpRom1 is associated with EVs, we assayed *C. parvum* LEVs and SEVs by

negative staining immunoelectron microscopy using a polyclonal rabbit serum directed against the

ID2 peptide (Trasarti et al., 2007). In agreement with the immunoblotting results, numerous strong

immunogold signals were detected on the surface of large vesicles (> 150 nm) occurring in LEVs

(Figure 4B). A negative control with pre-immune rabbit serum showed sporadic signals with one or

two gold particles for microscopic field (Figure 4C). We also tested SEVs by negative staining

immunoelectron microscopy with the same anti-ID2 serum, and we observed small vesicles that

were not labelled by immunogold particles (Figure 4D).

All together, these results suggested that sporozoites release two different types of extracellular

vesicles, among which those present in LEVs are generally larger than those in SEVs.

Immunolabelling for CpRom1 showed the presence of this protein only in LEVs.

3.3   Characterization of sporozoite extracellular vesicles by flow cytometry and electron

microscopy

To further characterize sporozoite EVs, we labelled vesicles in LEVs and SEVs sequentially with

two different fluorescent probes: at first, we used the hydrophilic dye NHS-AF647, which labels

proteins on the external surface of the vesicles; then, we utilized with the lipophilic dye CFSE,

which passively diffuse into vesicles labelling also internal vesicular proteins. Successively, vesicles

contained in LEVs and SEVs were separated by density gradient centrifugation on an iodixanol

gradient, and the fluorescent content of fractions was analysed by flow cytometry. FC analysis

showed that LEVs comprised a heterogeneous population of intact vesicles ranging in size from 50

nm to over 250 nm (Figure 5A, B). Notably, LEVs were often clumped together (Figure 6A),

contributing to the highest density of extracellular vesicles according to FACS analysis. In contrast,

SEVs displayed a more homogeneous composition (Figure 5C), consisting of smaller intact vesicles

with sizes below 150 nm (Figure 6B).

EVs were also quantified by FC. We observed that LEVs are more numerous than SEVs; moreover,

both vesicle populations presented a peak in fractions with density ranging between 1.08-1.14 g/ml

(Figure 6D). To establish more specifically the size of vesicles occurring in LEVs and SEVs, the

diameters of a hundred vesicles were measured through electron microscopy, and the corresponding

micrographs were subjected to statistical analysis. This experiment revealed that LEV (figure 6A)

and SEV (figure 6B) had a different medium size of 150 nm and 60 nm respectively. (Figure 6C).

3.4   Ultrastructural characterization of vesicles during the excystation

To capture images of vesicles at the time of their release from sporozoites, we analysed by SEM and

TEM the excystation mixture at various time points (from 0 to 30 min) after the induction. Hence,

we observed some sporozoites immediately after the egress from two oocysts (Figure 7A). Oocysts

appeared as thin, empty envelopes, and oocysts and sporozoites are surrounded by numerous

vesicles of varying sizes scattered throughout the microscopic field. Similarly, we captured images

at higher magnification showing two vesicles of different sizes at the very moment of their release

from the sporozoite membrane (Fig. 7B). Finally, we observed emerging vesicles from the apical

part of a sporozoite (Figure 7C) as well as a vesicle of approximately 200 nm just released from the

posterior part of another sporozoite (Figure 7D). Together, these images supported the origin of

EVs, at least the larger ones, by budding of the external membrane.

Since SEVs have physical parameters like exosomes, which originate from intracellular multi-

vesicular bodies (MVBs) (Van Niel et al., 2018; Barreca et al., 2023), we looked for a similar

structure in the sporozoite cytoplasm. Electron microscopy images showed a vacuolar structure with

internal membranes that resemble MVBs (Figure 8A and 8B). A similar MVB-like structure has not

yet been described so far in *Cryptosporidium* spp.

3.5   The Golgi protein CpGRASP is associated with LEVs but not with SEVs

No homologues of the most common mammalian proteins associated with exosomes, namely

tetraspanins (eg, CD63, CD81 or CD9) or endosomal sorting complex required for transport

(ESCRT) proteins (eg, Alix and Tsg101) (Théry et al. 2018), can be identified in *Cryptosporidium*

proteome (data not shown). On the other hand, well-conserved *Cryptosporidium* homologues of

proteins involved in vesicle trafficking can be traced back among those associated with the Golgi

apparatus. In all eukaryotes, the Golgi complex is a central hub for the vesicle trafficking (Doyle

and Wang, 2019), and Golgi proteins could represent markers for tracking various types of vesicles.

The Golgi reassembly and stacking proteins (GRASPs) belong to a protein family with a conserved

similarity in all eukaryotes, and the *Cryptosporidium* homolog is easily identifiable in CryptoDB.

Moreover, following a search in protein databases of extracellular vesicles and in the literature,

GRASPs have been found in different types of EVs (Chua et al., 2012; Peres da Silva et al., 2018).

Therefore, we proceeded to clone the gene for the *C. parvum* homolog of GRASPs to express a

histidine-fusion protein in bacteria. The purified recombinant protein (6his-CpGRASP) was then

used to generate a specific antiserum in mice. In Western blot experiments, when probed with the

anti-GRASP serum, an 80 kDa band was identified (Figure 9) in the whole oocyst-sporozoite lysate,

which corresponded to the expected molecular mass of 82.49 kDa. The same 80 kDa band was also

observed in LEVs extract, while no band was detected in the SEVs extract or in the supernatant

following the ultracentrifugation. This last result suggested that the biogenesis of LEVs involves

Golgi-derived vesicles unlike SEVs, which conversely did not show any evidence for thre

occurrence of CpGRASP.

3.6   Comparison of protein profiles of LEVs and SEVs by fluorescent labelling and SDS-PAGE electrophoresis

A first analysis regarding the protein content in sporozoite vesicles was performed by comparing the electrophoretic profiles of the two types of vesicles. To this end, we double-labeled the LEVs and SEVs with NHS-AF647 (red) and CFSE (green) and analysed the corresponding protein profiles by SDS-PAGE (Dehghani and Gaborski, 2020). Figure 10 shows the image obtained by overlaying the green and red channels onto the electrophoretic protein profiles of LEVs and SEVs. Upon comparing the electrophoretic profiles, we have identified distinct common bands (black arrows) as well as some prominent unique bands (white arrows) in SEVs. This method allowed a rapid generation of distinctive patterns for LEVs and SEVs and confirmed the occurrence of differences in the protein content of them. Of note, the SEVs profile exhibited a greater number of red bands indicating the likely presence of a greater number of surface proteins.

3.7   Proteomic analysis of LEVs and SEVs content

To identify *C. parvum* proteins present in above-reported vesicles, components of LEVs and SEVs were resolved by SDS-PAGE and stained with Coomassie G250. The resulting whole gel lanes (see Supplemental Figure S1) were divided into 9 slices that were further subjected to trypsinolysis and then to nLC-ESI-Q-Orbitrap-MS/MS analysis. On the whole, we identified 60 *C. parvum* proteins of which 39 were in common between the two vesicle types; conversely, 5 proteins were identified only in LEVs, while 16 were found exclusively in SEVs. All identified proteins are listed in Table 1, and details of the corresponding proteomic analysis are reported in Supplemental Table S1. The full annotation from the CriptoDB database for all assigned proteins is reported in Supplemental Table S2, which also includes presumptive information on the corresponding function, gene ontology (GO terms ID), assignment as membrane/soluble component, and cellular localisation.

According to functional predictions, the largest number (14) of annotated proteins included enzymes related to cell metabolism, while the second group (12 in number) consisted of components involved in protein folding, such as heat shock proteins (HSPs) and molecular

chaperones (Figure 11, panel A). This latter group included 4 heat shock proteins 70 (HSP70s) and

2 heat shock proteins 90 (HSP90s) that may have a different subcellular localization, as inferred by

the variable presence of signals and/or transmembrane domains. Additional protein categories were

related to protein synthesis (9 in number), regulatory function (6 in number), cytoskeleton (3 in

number), redox homeostasis (2 in number) and proteolysis (2 in number) (Figure 10, panel A). The

presumptive function remained unknown for 12 proteins.

All the proteins were also analysed in silico to identify signal peptide, transmembrane domain and

endoplasmic reticulum retention signal, which are indicative traits of their subcellular localization.

Thus, the largest group was composed of soluble components (34 in number) with a putative

cytoplasmic localization (Figure 11, panel B). A second group of 7 proteins showed a signal for

sorting and/or retention in the endoplasmic reticulum. Six secretory proteins were identified due to

signal peptide at their N-terminus, whereas three showed a transmembrane domain. Finally, 2

proteins were predicted as nuclear components. However, the location of eight proteins remained

unclear, as the potential location is assigned to subcellular compartments (*i.e.* mitochondrion,

plastid, and lysosome), the existence of which is still unknown in *C. parvum*.

A putative interactome of EVs proteins was modelled by STRING interaction analysis. This search

revealed a big network connecting 51 components (Supplemental figure S2, panel A and

Supplemental Table S3), which was identified with a with medium confidence (0.4). The

involvement of most (85%) of the identified proteins in this network emphasized the occurrence of

a functional assembly bridging different molecular processes, which seemed highly represented in

EVs. When higher confidence (0.9) was used for the analysis, this interaction network was limited

to include only 34 proteins (57.3%) and three main sub-networks (Supplemental Figure S2, panel

B) related to carbohydrate metabolism. Therefore, the above-reported interaction data highlighted

the possible existence of functional macromolecular complexes in the *C. parvum* EVs

involved in specific biological processes, as evidenced by the coherence with the most

represented predicted functional categories (Figure 11, panel A).

The occurrence of highly related protein homologues assigned to EVs in other organisms was

verified for 48 out of the 60 species identified in this study. Conversely, this proteomic investigation

originally identified 12 proteins never assigned to EVs so far, among which all the ones (4 in

number) here predicted being membrane components (Supplemental Table S2). An *in-silico*

prediction of the schematic structure of these 12 proteins is shown in Figure 12. The occurrence of a

signal peptide was evidenced in 10 proteins, among which 4 also contained a transmembrane

domain, thus suggesting an extracellular/membrane localization for these components; the

remaining 2 proteins did not contain localization motifs, as typical of cytoplasmic/luminal proteins.

These 12 proteins included: i) three large molecules, such as the cysteine-rich secretory protein

(cgd7_4310-RA-p1) including repeated CAP domains related to secretory proteins of metazoans

(Gibbs et al., 2008) and allergen V5/Tpx-1-related (cgd5_2020_RA-p1); ii) a protein

(cgd6_1630_RA) containing a CS domain, which is a binding module for HSP90, possibly involved

in recruiting heat shock proteins to multiprotein assembly (Lee YT, et al 2004); iii) membrane-

linked aspartyl-protease (cgd1_3690-RA-p1) that was detected only in the SEVs proteome.

Additional proteins were uncharacterized proteins (cgd1_660-RA, cgd2_3780-RA, cgd3_1540-RA,

cgd3_3370-RA, cgd6_1080-RA, cgd6_4460-RA, cgd6_710-RA, cgd7_400-RA) that do not show

similarity with already known protein families. This group of *Cryptosporidium*-specific proteins

also included the glycoprotein GP60 (cgd6_1080-RA-p1), which is a well-known immunodominant

antigen of the parasite (Wanyiri et al., 2007) and was detected in both types of EVs.

4   Discussion

The excystation in *Cryptosporidium* spp. represents the beginning of a new infectious cycle and implies the release of intact vesicles from sporozoites. It has been previously demonstrated that micronemes and dense granules excrete the most of their soluble contents in a couple of hours after the excystation in the external medium (Chen et al., 2004). We here show that *C. parvum* sporozoites also release extracellular vesicles and that can be distinguished in large and small types, here named LEVs and SEVs.

Once the excystation starts, the release of EVs proceeds autonomously without any additional requirements, such as host factors, and can be readily achieved also *in vitro*. On this basis, differential centrifugation of the excystation medium resolved *C. parvum* EVs in two sediments that, based on their features, were related to LEVs and SEVs, respectively. Dedicated TEM experiments definitively demonstrated that these vesicles have a different size and, accordingly, can be distinguished into LEVs (medium size150 nm) and SEVs (medium size 60 nm). Furthermore, these vesicles are partially different for protein composition as initially shown by the western blotting analysis, which demonstrated the presence of CpRom1 and CpGRASP exclusively in LEVs, as well as by fluorescent labelling combined with SDS-PAGE analysis that revealed distinct protein banding.

Furthermore, the two types of vesicles are partially different in compositions as shown by the presence of CpRom1 and CpGRASP exclusively on LEVs as well as by the distinct fluorescent protein banding showed by the SDS-PAGE analysis. This partial difference was finally verified by dedicated proteomic experiments that traced some proteins only in one of the two sediments; in particular, 39 proteins were in common between two vesicle types, while 5 and 16 proteins were identified only in LEVs and SEVs, respectively.

The finding of *C. parvum* EVs reported in the present study was fortuitous, and it was the consequence of our investigation on CpRom1 rhomboid, which belongs to a peculiar family of serine proteases integrally embedded in the plasma membrane. Apicomplexa have different types of

rhomboids that act cleaving specific membrane proteins in a sequential manner; some of these molecules have fundamental roles in the motility of the parasite, and in its penetration capacity into the host cell (Dowse et al., 2008; Rugarabamu et al., 2015). At present, the specific role of CpRom1 in sporozoite invasion of enterocytes is still unclear. Nevertheless, this study confirmed that mature CpRom1is the largest rhomboid described so far, with a measured mass of 110 kDa. The two-dimensional model of this protein predicted a distinctive feature, with a large N-terminal hydrophilic portion extending outward from the plasma membrane (Figure 2). Consistent with this model, the gold particles used in immune electron microscopy experiments labelled the external membrane of LEVs (Figure 4B). These results were obtained using antibodies targeting the ID2 antigen, which is part of the above-reported N-terminal region exposed on the external surface of the membrane (Figure 2). It is worth noting that, except for those with mitochondrial localization (PARLs), all the rhomboids described so far are associated with the external side of cell membrane (Kandel and Neal, 2020). In this context, this study is original since it demonstrates for the first time the association of a rhomboid with extracellular vesicles.

To determine the subcellular origin of the sporozoite EVs, we sought to trace proteins that are undoubtedly involved in vesicular trafficking within the cell. Among these proteins, we observed that the GRASP proteins from *Cryptosporidium* spp. are strictly related to other members of this protein family. Indeed, immunoblotting experiments demonstrated that CpGRASP, a member of the conserved Golgi-associated protein family, is present on LEVs and not on SEVs similarly to CpRom1. Golgi reassembly stacking protein family are tethered to the external membrane of Golgi vesicles, forming the Golgi stack network that coordinates the sorting of secreted proteins (Rabouille and Lindsted, 2016). In addition, GRASPs regulate the unconventional secretion of proteins directed toward the cell membrane, tagging vesicles originating directly from the ER through a Golgi-independent pathway (Ahat et al., 2019). Therefore, the exclusive presence of CpGRASP on LEVs suggests that the biogenesis of these vesicles is associated with the ER- and/or the Golgi pathway.

TEM and SEM micrographs clearly showed the generation of EVs budding from the

external membrane of sporozoites (Figure 7). Budding from the plasma membrane is the most

common mechanism for the generation of EVs, except for exosomes. Among other things, a

specific difference between budding-generated EVs (or ectosomes) and exomes is their size, which

ranges from 100 to 1000 nm for the ectosomes and from 30 to 150 nm for the exosomes (Meldolesi,

2018). Indeed, the newly formed vesicles as shown in Figure 7B and 7D were larger than

150 nm, in agreement with what expected for vesicles generated through the budding

process. Therefore, we can conclude that outward budding of *C. parvum* sporozoite membrane is

responsible for the generation of LEVs.

Tentatively determining the origin of smaller SEVs was more challenging, as these vesicles

were only observed after their release into the excystation medium. Since SEVs share some

physical parameters with exosome like density (ranging from 1.08 to 1.2 g/ml) and size (ranging

from 50 to 150 nm) it was reasonable to consider SEVs as exosome-like vesicles. Exosomes are

generally distinguished from other vesicles for their origin within the endosomal pathway through

the formation of MVBs; upon fusion with the plasma membrane these intracellular bodies release

exosomes outside the cell (Doyle and Wang, 2019). After assuming the existence of organelles like

MVBs in sporozoites, we accurately examined the corresponding TEM micrographs finally

identifying novel cytoplasmic structures that highly resemble MVBs; accordingly, we here defined

them MVB-like organelles. Analogously to MVBs, MVB-like organelles are made of a large

globular vesicle enclosed by a lipid membrane, which in turn contains various smaller vesicles

(Figure 8). These novel organelles occur in proximity of the sporozoite membrane. However, since

MVBs of proven endosomal origin have not yet been described in *Cryptosporidium* spp., our

assimilation of SEVs to exosomes remains speculative.

Integrated Western blotting, fluorescent labelling and proteomics experiments have shown

that *C. parvum* LEVs and SEVs also differ slightly in protein composition. Of the 60 proteins

assigned to proteomics, 39 were common in two types of vesicles, while 5 and 16 components were

unique to LEVs and SEVs respectively. However, proteins identified by proteomic analysis

represent the most abundant proteins of the vesicles and other methods (*i.e.* Western blotting) can be

used to track lower-enriched proteins such as CpRom1 and CpGRASP that selectively occur in

LEVs.

Based on a specific analysis for presumptive cellular localization, most (35 out of 60) of

these vesicle proteins were predicted to occur in the cell cytoplasm. Various types of cytoplasmic

proteins have already been described in extracellular vesicles (Gurung et al., 2021), even if their

function in these particles is not yet clear. However, it is conceivable that some cytoplasmic proteins

are dragged from the cytoplasm during the formation of the emerging vesicles. On the other hand, 6

vesicle proteins here identified show a consensus sequence for the endoplasmic reticulum (ER),

while other 6 ones contain a signal peptide, suggesting their sorting towards EVs through specific

ER and/or Golgi-mediated secretory pathways.

Based on gene ontology information, most (48 out of 60) of the proteins here identified were

tentatively assigned to a known functional category, while 12 ones were considered peculiar of

Cryptosporidium spp. Among the former group, the largest number of annotated molecules included

enzymes related to cellular metabolism and components involved in protein folding and

biosynthesis.

Bioinformatic analysis of the corresponding protein interactions revealed the existence of a single

network connecting most (51 out of 60) of the assigned molecular entries, suggesting the existence

of unique functional machinery interconnecting different processes.

Regarding the remaining 12 *Cryptosporidium*-specific vesicle proteins, 10 have never been

described so far, whereas the protein identified as cgd1_3690-RA was clearly homologous to an

aspartyl-protease and glycoprotein GP60 (cgd6_1080-RA) have been already mentioned in several

studies. Figure 12 outlines some of the molecular features of these *Cryptosporidium*-specific

proteins, including their presumptive localization with respect to plasma membrane, which is based

on the presence of a signal peptide and transmembrane or other specific domains. Noteworthy, 10 of

these proteins show a signal peptide, suggesting that reach the vesicle through a secretive pathway,

while 4 of them, such as GP60, have a transmembrane domain essential for their exposure on the

surface of EVs.

With respect to aspartyl protease (cgd1_3690-RA), which we will call membrane protease CpMAP,

it has only been detected in SEVs. *C. parvum* aspartyl-proteases were previously demonstrated

being crucial for a successful infection. In this context, the aspartyl-protease inhibitor Indinavir was

previously shown reducing parasite proliferation both *in vitro* and *in vivo* when administered in the

early phase of infection (Mele et al., 2003). The present study demonstrates that CpMAP is

expressed in the early stage of parasite infection, and it is conveyed by vesicles outside the

sporozoites, thus possibly playing a role in the proteolysis of one or more host proteins.

Otherwise, GP60 was detected in LEVs and SEVs. This glycoprotein, also referred to as

GP40/15, is one of the most relevant antigens of the parasite, which may determine an

immunodominant response in the host (Sestak et al., 2002). It occurs exclusively in the

Cryptosporidium genus and has no homologs even in other apicomplexans. This protein is

synthesized as a large precursor of 60 kDa (GP60), which may undergo a proteolytic cleavage by a

furin-like protease to generate a soluble fragment of approximately 40 kDa (GP40) and a smaller

membrane-anchored segment of 15 kDa (GP15) (Wanyiri et al., 2007). Despite its nature of

membrane protein, GP15 is released during the sporozoite gliding (O'Connor et al., 2007). It is

remarkable that this fact is in perfect accordance with the discharge of EVs after the excystation.

Extracellular vesicles constitute important elements of the host-parasite interaction and

mediate cell-to-cell communication in different directions: between the parasite and its host, among

the parasites, and among the host cells in response to the parasitic infection (Wu et al., 2019). In

most of the cases, parasite-generated vesicles modulate the host immune response by transferring

parasite molecules (*i.e.* mRNA, various types of non-coding RNA, DNA and proteins) that act on

host cells like macrophages (Olajide and Cai, 2020).

In apicomplexan parasites, extracellular vesicles were studied in *Plasmodium* spp. and

Toxoplasma gondii. In *Plasmodium* spp. host-generated exosomes have been described containing

parasite proteins. In *Plasmodium yoelii*, these exosomes inoculated in mice can elicit an IgG

response and a protective immunity (Martin-Jaular et al., 2011). In *Plasmodium falciparum*, host-

generated exosomes can also coordinate gametocytogenesis among infected red blood cells by

delivering parasitic DNA (Regev-Rudzki et al., 2013). Importantly, microvesicles induced by

Plasmodium spp. contribute to the onset of inflammatory symptoms of malaria. In *P. falciparum,*

the number of extracellular vesicles excreted by blood cells increases in cerebral malaria (Pankoui

Mfonkeu et al., 2010; Sahu et al., 2013), and a high number of them correlates with high fever in

Plasmodium vivax infection (Campos et al., 2010).

Host-generated extracellular vesicles are induced also by *T. gondii* in infected fibroblasts (Pope and

Lässer , 2013), and the parasitic antigens in microvesicles stimulate dendritic cells leading to a Th1

protective response (Dlugońska and Gatkowska, 2016). It was also demonstrated that *T. gondii*

generates its own exosomes and extracellular vesicles (Wowk PF et al., 2017).

With this study, we have added novel information in the complex scenario of extracellular vesicles

secreted by apicomplexan parasites. We discovered that also *C. parvum* excysted sporozoites

release EVs. Based on their size, these extracellular vesicles were resolved in LEVs and SEVs.

Integrated approaches allowed defining common and peculiar characteristics of both vesicle

typologies. CpRom1 and CpGRASP were selectively identified in LEVs, and thus can be

considered markers of this vesicle type. Conversely, the abundant antigenic glycoprotein GP60 was

detected in both type of vesicles; this fact may play a role in directing the host's immune response.

Further studies are needed to decipher the specific function of these extracellular vesicles and their

relationship with *C. parvum* infection.

Author contributions

LB: Investigation, Methodology, Writing – review & editing. ZB: Investigation, Methodology, Writing – review & editing. AMS: Investigation, Methodology, Writing – review & editing. IV: Investigation, Methodology, Writing – review & editing. IP: Investigation, Writing – review & editing. EM: Investigation, Writing – review & editing. AS: Conceptualization, Formal Analysis, Writing – review & editing. MS[1]: Conceptualization, Formal Analysis, Writing – review & editing. MS[6]: Conceptualization, Formal Analysis, Writing – review & editing. MLF: Supervision, Conceptualization, Formal Analysis, Writing – review & editing. FT: Resources, Supervision, Conceptualization, Investigation, Methodology, Writing – review & editing.

Funding

This work was supported by grants from the National Recovery and Resilience Plan, mission 4, component 2, investment 1.3, MUR call n. 341/2022 funded by the European Union-NextGenerationEU for the project "One Health Basic and Translational Research Actions addressing Unmet Needs on Emerging Infectious Diseases (IN-FACT)", D.C. 1554/2022, PE00000007.

Acknowledgments

We are thankful to CryptoDB for making available all the genomic data used in this study.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

Ahat, E., Li, J., Wang, Y. (2019). New Insights Into the Golgi Stacking Proteins. Front. Cell. Dev. Biol. 7:131. doi: 10.3389/fcell.2019.00131

Allison, G. M., Rogers, K. A., Borad, A., Ahmed, S., Karim, M. M., Kane, A.V., et al. (2011). Antibody responses to the immunodominant *Cryptosporidium* gp15 antigen and gp15 polymorphisms in a case-control study of cryptosporidiosis in children in Bangladesh. Am. J. Trop. Med. Hyg. 85(1):97-104. doi: 10.4269/ajtmh.2011.11-0043

Barreca, V., Boussadia, Z., Polignano, D., Galli, L., Tirelli, V., Sanchez, M., et al. (2023). Metabolic labelling of a subpopulation of small extracellular vesicles using a fluorescent palmitic acid analogue. J. Extracell. Vesicles, e12392. doi.org/10.1002/jev2.12392

Bouzid, M., Hunter, P.R., Chalmers, R.M., Tyler, K.M. (2013). *Cryptosporidium* pathogenicity and virulence. Clin. Microbiol. Rev. 26:115-34. doi: 10.1128/CMR.00076-12

Campos, F.M., Franklin, B.S., Teixeira-Carvalho, A., Filho, A.L., de Paula, S.C., Fontes, C. J., et al. (2010). Augmented plasma microparticles during acute *Plasmodium vivax* infection. Malar. J. 9:327. doi: 10.1186/1475-2875-9-327

Carruthers, V.B., Blackman, M.J. (2005). A new release on life: emerging concepts in proteolysis and parasite invasion. Mol. Microbiol. 55(6):1617-30. doi: 10.1111/j.1365-2958.2005.04483

Checkley, W., White, A.C.Jr., Jaganath, D., Arrowood, M.J., Chalmers, R.M., Chen, X.M., et al. (2015). A review of the global burden, novel diagnostics, therapeutics, and vaccine targets for cryptosporidium. Lancet. Infect. Dis. 15:85-94. DOI: 10.1016/S1473-3099(14)70772-8

Chen, X.M., O'Hara, S.P., Huang, B.Q., Nelson, J.B., Lin, J.J., Zhu, G., et al. (2004). Apical organelle discharge by *Cryptosporidium parvum* is temperature, cytoskeleton, and intracellular calcium dependent and required for host cell invasion. Infect. Immun. 72(12):6806-16. doi: 10.1128/IAI.72.12.6806-6816.2004

Chua, C.E., Lim, Y.S., Lee, M.G., Tang, B.L. (2012). Non-classical membrane trafficking processes galore. J. Cell. Physiol. 227(12):3722-30. doi: 10.1002/jcp.24082

Coscia, C., Parolini, I., Sanchez, M., Biffoni, M., Boussadia, Z., Zanetti, C., et al. (2016).

Generation, Quantification, and Tracing of Metabolically Labeled Fluorescent Exosomes. Methods.

Mol. Biol. 1448:217-35. doi: 10.1007/978-1-4939-3753-0_16

Dehghani, M., Gaborski, T. R. (2020). Fluorescent labeling of extracellular vesicles. Methods.

Enzymol. 645:15-42. doi: 10.1016/bs.mie.2020.09.002

Długońska, H., Gatkowska, J. (2016). Exosomes in the context of *Toxoplasma gondii* – host

communication. Ann. Parasitol. 62(3):169–174. doi: 10.17420/ap6203.50

Doyle, L.M., Wang, M.Z. (2019). Overview of Extracellular Vesicles, Their Origin, Composition,

Purpose, and Methods for Exosome Isolation and Analysis. Cells. 8(7):727. doi:

10.3390/cells8070727

Dowse, T.J., Koussis, K., Blackman, M.J., Soldati-Favre, D. (2008). Roles of proteases during

invasion and egress by *Plasmodium* and *Toxoplasma*. Subcell. Biochem. 47:121-39. doi:

10.1007/978-0-387-78267-6_10

Dowse, T.J., Soldati, D. (2005). Rhomboid-like proteins in Apicomplexa: phylogeny and

nomenclature. Trends. Parasitol. 21(6):254-8. doi: 10.1016/j.pt.2005.04.009

Fayer, R., and Xiao, L. (2007). *Cryptosporidium* and Cryptosporidiosis (CRC Press).

Feng, Y., Ryan, U.M., Xiao, L. (2018). Genetic diversity and population structure of

Cryptosporidium. Trends. Parasitol. 34(11):997-1011. doi: 10.1016/j.pt.2018.07.009

Gibbs, G.M., Roelants, K., O'Bryan, M.K. (2008). The CAP superfamily: cysteine-rich secretory

proteins, antigen 5, and pathogenesis-related 1 proteins--roles in reproduction, cancer, and immune

defense. Endocr. Rev. 29(7):865-97. doi: 10.1210/er.2008-0032

Guérin, A., Strelau, K.M., Barylyuk, K., Wallbank, B.A., Berry, L., Crook, O.M., et al. (2023).

Cryptosporidium uses multiple distinct secretory organelles to interact with and modify its host cell.

Cell. Host. Microbe. 31(4):650-664.e6. doi: 10.1016/j.chom.2023.03.001

Gurung, S., Perocheau, D., Touramanidou, L., Baruteau, J. (2021) The exosome journey: from

biogenesis to uptake and intracellular signalling. Cell. Commun. Signal. 19(1):47. doi:

10.1186/s12964-021-00730-1

Kandel, R.R., Neal, S.E. (2020). The role of rhomboid superfamily members in protein

homeostasis: Mechanistic insight and physiological implications. Biochim. Biophys. Acta Mol.

Cell. Res. 1867(10):118793. doi: 10.1016/j.bbamcr.2020.118793

Kotloff, K.L., Nataro, J. P., Blackwelder, W.C., Nasrin, D., Farag, T.H., Panchalingam, S., et al.

(2013). Burden and aetiology of diarrhoeal disease in infants and young children in developing

countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. Lancet.

382:209-22. doi: 10.1016/S0140-6736(13)60844-2

Kühnle, N., Dederer, V., Lemberg, M.K. (2019). Intramembrane proteolysis at a glance: from

signalling to protein degradation. J. Cell. Sci. 132(16):jcs217745. doi: 10.1242/jcs.217745

Lee, Y.T., Jacob, J., Michowski, W., Nowotny, M., Kuznicki, J., Chazin, W. J. (2004). Human Sgt1

binds HSP90 through the CHORD-Sgt1 domain and not the tetratricopeptide repeat domain. J. Biol.

Chem. 279(16):16511-7. doi: 10.1074/jbc.M400215200

MacKenzie, W.R., Hoxie, N.J., Proctor, M.E., Gradus, M.S., Blair, K.A., Peterson, D.E., et al.

(1994). A massive outbreak in Milwaukee of *Cryptosporidium* infection transmitted through the

public water supply. N. Engl. J. Med. 331:161-7. doi: 10.1056/NEJM199407213310304

Martin-Jaular, L., Nakayasu, E.S., Ferrer, M., Almeida, I.C., Del Portillo, H.A. (2011). Exosomes

from *Plasmodium yoelii*-infected reticulocytes protect mice from lethal infections. PLoS One.

6(10):e26588. doi: 10.1371/journal.pone.0026588

Meldolesi, J. (2018). Exosomes and Ectosomes in Intercellular Communication. Curr .Biol.

28(8):R435-R444. doi: 10.1016/j.cub.2018.01.059

Mele, R., Gomez Morales, M.A., Tosini, F., Pozio, E. (2003). Indinavir reduces *Cryptosporidium*

parvum infection in both in vitro and in vivo models. Int. J. Parasitol. 33(7):757-64. doi:

10.1016/s0020-7519(03)00093-6

O'Connor, R.M., Wanyiri, J.W., Cevallos, A.M., Priest, J.W., Ward, H.D. (2007). *Cryptosporidium*

parvum glycoprotein gp40 localizes to the sporozoite surface by association with gp15. Mol

Biochem Parasitol. 156(1):80-3. doi: 10.1016/j.molbiopara.2007.07.010

Olajide, J.S., Cai, J. (2020). Perils and Promises of Pathogenic Protozoan Extracellular Vesicles.

Front. Cell. Infect. Microbiol. 10:371. doi: 10.3389/fcimb.2020.00371

Pankoui Mfonkeu, J.B., Gouado, I., Fotso Kuaté, H., Zambou, O., Amvam Zollo, P.H., Grau, G.E.,

et al. (2010). Elevated cell-specific microparticles are a biological marker for cerebral dysfunctions

in human severe malaria. PLoS One. 5(10):e13415. doi: 10.1371/journal.pone.0013415

Peres da Silva, R., Martins, S.T., Rizzo, J., Dos Reis, F.C.G., Joffe, L.S., Vainstein, M., et al. (2018).

Golgi Reassembly and Stacking Protein (GRASP) Participates in Vesicle-Mediated RNA Export in

Cryptococcus Neoformans. Genes (Basel). 9(8):400. doi: 10.3390/genes9080400

Perry, M.M., Gilbert, A.B. (1979). Yolk transport in the ovarian follicle of the hen (*Gallus*

domesticus): lipoprotein- like particles at the periphery of the oocyte in the rapid growth phase. J.

Cell. Sci. 39: 257–272. doi: 10.1242/jcs.39.1.257

Pope, S.M., Lässer, C. (2013). *Toxoplasma gondii* infection of fibroblasts causes the production of

exosome-like vesicles containing a unique array of mRNA and miRNA transcripts compared to

serum starvation. J. Extracell. Vesicles. Dec 11;2. doi: 10.3402/jev.v2i0.22484

Rabouille, C., Linstedt, A.D. (2016). GRASP: A Multitasking Tether. Front. Cell. Dev. Biol. 4:1.

doi: 10.3389/fcell.2016.00001

Regev-Rudzki, N., Wilson, D.W., Carvalho, T.G., Sisquella, X., Coleman, B.M., Rug, M., et al.

(2013). Cell-cell communication between malaria-infected red blood cells via exosome-like

vesicles. Cell. 153(5):1120-33. doi: 10.1016/j.cell.2013.04.029

Rugarabamu, G., Marq, J.B., Guérin, A., Lebrun, M., Soldati-Favre, D. (2015). Distinct

contribution of *Toxoplasma gondii* rhomboid proteases 4 and 5 to micronemal protein protease 1

activity during invasion. Mol. Microbiol. 97(2):244-62. doi: 10.1111/mmi.13021

Sahu, U., Sahoo, P.K., Kar, S.K., Mohapatra, B.N., Ranjit, M. (2013). Association of TNF level

with production of circulating cellular microparticles during clinical manifestation of human

cerebral malaria. Hum. Immunol. 74(6):713-21. doi: 10.1016/j.humimm.2013.02.006

Salzano, A.M., Novi, G., Arioli, S., Corona, S., Mora, D., Scaloni, A. (2013). Mono-dimensional

blue native-PAGE and bi-dimensional blue native/urea-PAGE or/SDS-PAGE combined with nLC-

ESI-LIT-MS/MS unveil membrane protein heteromeric and homomeric complexes in *Streptococcus*

thermophiles. J. Proteomics. 94:240-261. doi: 10.1016/j.jprot.2013.09.007.

Sestak, K., Ward, L.A., Sheoran, A., Feng, X., Akiyoshi, D.E., Ward, H.D., et al. (2002). Variability

among *Cryptosporidium parvum* genotype 1 and 2 immunodominant surface glycoproteins. Parasite

Immunol. 24(4):213-9. doi: 10.1046/j.1365-3024.2002.00455.x

Shively, S., Miller, W.R. (2009). The use of HMDS (hexamethyldisilazane) to Replace Critical

Point Drying (CPD) in the Preparation of Tardigrades for SEM (Scanning Electron Microscope)

Imaging.   Transactions   of   the   Kansas   Academy   of   Science.   112(4):198-200.

doi:10.1660/062.112.0407

Templeton, T.J., Lancto, C.A., Vigdorovich, V., Liu, C., London, N.R., Hadsall, K.Z., et al. (2004).

The *Cryptosporidium* oocyst wall protein is a member of a multigene family and has a homolog in

Toxoplasma. Infect. Immun. 72(2):980-7. doi: 10.1128/IAI.72.2.980-987.2004

Théry, C., Clayton, A., Amigorena, S., Raposo, G. (2006). Isolation and Characterization of

Exosomes from Cell Culture Supernatants and Biological Fluids, in the Current Protocols in Cell

Biology. 3.22.1-3.22.29.

Théry, C., Witwer, K.W., Aikawa, E., Alcaraz, M.J., Anderson, J.D., Andriantsitohaina, R., et al.

(2018). Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position

statement of the International Society for Extracellular Vesicles and update of the MISEV2014

guidelines. J. Extracell. Vesicles. 7(1):1535750. doi: 10.1080/20013078.2018.1535750

Tosini, F., Ludovisi, A., Tonanzi, D., Amati, M., Cherchi, S., Pozio, E., et al. (2019). Delivery of

SA35 and SA40 peptides in mice enhances humoral and cellular immune responses and confers

protection against *Cryptosporidium parvum* infection. Parasit. Vectors. 12(1):233. doi:

10.1186/s13071-019-3486-8

Trasarti, E., Pizzi, E., Pozio, E., Tosini, F. (2007). The immunological selection of recombinant

peptides from *Cryptosporidium parvum* reveals 14 proteins expressed at the sporozoite stage, 7 of

which are conserved in other apicomplexa. Mol. Biochem. Parasitol. 152:159-69. doi:

10.1016/j.molbiopara.2006.12.010

Van Niel, G., D'Angelo, G., Raposo, G. (2018). Shedding light on the cell biology of extracellular

vesicles. Nat. Rev. Mol. Cell. Biol. 19(4): 213–228). doi: 10.1038/nrm.2017.125

Wang, X., Chen, J., Zheng, J. (2022). The state of the art of extracellular vesicle research in

protozoan infection. Front. Genet. 13:941561. doi: 10.3389/fgene.2022.941561.

Wanyiri, J.W., O'Connor, R., Allison, G., Kim, K., Kane, A., Qiu, J., et al. (2007). Proteolytic

processing of the *Cryptosporidium* glycoprotein gp40/15 by human furin and by a parasite-derived

furin-like protease activity. Infect. Immun. 75(1):184-92. doi: 10.1128/IAI.00944-06

Wowk, P.F., Zardo, M.L., Miot, H.T., Goldenberg, S., Carvalho, P.C., Mörking, P.A. (2017).

Proteomic profiling of extracellular vesicles secreted from *Toxoplasma gondii*. Proteomics. 17(15-

16). doi: 10.1002/pmic.201600477

Wu, Z., Wang, L., Li, J., Wang, L., Wu, Z., Sun, X. (2019). Extracellular Vesicle-Mediated

Communication Within Host-Parasite Interactions. Front. Immunol. 9:3066. doi:

10.3389/fimmu.2018.03066

Xiao, L., Feng, Y. (2017). Molecular epidemiologic tools for waterborne pathogens

Cryptosporidium spp. and *Giardia duodenalis*. Food Waterborne Parasitol. 8-9:14-32. doi:

10.1016/j.fawpar.2017.09.002

Figure captions

Figure 1. Diagram of the procedure used for the isolation of extracellular vesicles from the

excystation medium.

Figure 2. The two-dimensional model of CpRom1 in relation to the cell membrane: the diagram

shows protein six transmembrane regions, two cytoplasmic loops (above the membrane), the large

N-terminal extracellular region including the ID2 antigen, and the small extracellular region at the

C-terminus (above the membrane). The small arrows show the position of the ID2 peptide (see

text), and the amino acid sequence of the ID2 is expanded below.

Figure 3. CpRom1 identification by Western blot. A, Western blotting on oocysts and sporozoite

lysates probed with mouse anti-CpRom1 serum: 1, lysate from $5x10^6$ unexcysted oocysts; 2, lysate

from $5x10^6$ oocysts after 30 min of excystation. B, Western blotting on excystation medium and

TCA-precipitated supernatant of probed with mouse anti-CpRom1 serum: 1, lysate from $1x10^7$

oocysts after 30 min of excystation; 2, TCA-precipitated supernatant before the excystation; 3,

TCA-precipitated supernatant of sample 1.

Figure 4. Immunolocalization of CpRom1 by Western blot on microvesicle extracts and

immunoelectron microscopy of the corresponding extracellular vesicles. A, Immunoblotting with

anti-CpRom1 mouse serum: 1, unexcysted oocysts lysate; 2, excysted soporozoites; 3, LEVs

extract; 4, SEVs extract; 5, TCA-precipitated supernatant of SEVs. B, Negative staining

immunoelectron microscopy of LEVs labelled with anti-CpRom1 rabbit serum. C, Negative

staining immunoelectron microscopy of LEVs labelled with pre-immune rabbit serum. D, Negative

staining immunoelectron microscopy of SEVs labelled with anti-CpRom1 rabbit serum.

Figure 5. Flow cytometry analysis of gradient fractions of LEVs and SEVs. A, PBS buffer used to

resuspend EVs as negative control. B, Dot plot of LEVs fraction at 1.1 g/ml density. C, Dot plot of

SEVs at 1.1 g/ml density. D, diagram comparing the distribution of LEVs and SEVs in the gradient

fractions. Left panels indicate gate dimensions (A≤200 nm, B=500 nm and C=1 µm). Right dot plot

panels show vesicles distribution in terms of fluorescence.

Figure 6. Physical characterization of LEVs and SEVs by electron-microscopy. A, TEM negative

staining of LEVs. B, TEM negative staining of SEVs. C, diagram showing percentage distribution

of vesicle size in LEVs (green) and in SEVs (orange).

Figure 7. Electron microscopy images of EVs at their release from sporozoites. A, Backscattered

electron SEM micrograph of two excysted oocysts showing the release of vesicles during the

sporozoites egress. B, High magnification of an egressed sporozoite which shows two budding

vesicles (head arrows). C, TEM micrograph showing the plasmamembrane budding of a vesicle

from the apical region of a sporozoite (white arrow). D, TEM micrograph of a vesicle budding from

the posterior region of a sporozoite. DG: dense granules; mn: micronemes; Cr: crystalloid; Nu:

nucleus.

Figure 8. Ultrathin sections of excysted sporozoites (A, B) showing possible MVB-like organelles

inside the cells (white arrows).

Figure 9. Western blot probed with mouse anti-CpGRASP serum. 1, lysate oocyst-sporozoite

probed with pre-immune serum; 2, lysate oocyst-sporozoite; 3, LEVs extract; 4, SEVs extract; 5,

SEVs supernatant precipitated with TCA. 4-20% SDS-PAGE, lane 1 probed with 1:500 with mouse

serum before the immunization; lane 2-5 probed with 1:500 mouse serum after the immunization

with recombinant CpGRASP.

Figure 10. SDS-PAGE of two independent excystation experiments followed by labelling with the fluorescent dyes NHS-AF647 (red) and CFSE (green) of the different stages of centrifugation: 1 with $1x10^7$oocysts, 2 with 5×10^6 oocysts duplicate sporozoite lysate; LEVs and SEVs after labelling with NHS-AF647 (red) and CFSE (green) molecular probes. Black arrows indicate common green bands, open arrows indicate red bands present only in SEVs extract.

Figure 11. Quantitative representation of the different categories of the proteins identified in the extracellular vesicles. A, graphic of the EVs proteins distributed on the basis of their presumptive functions. The classification was based on the sequence homologies with characterized proteins. B, graphic of the EVs proteins distributed based on their presumptive sub-cellular localization. Prediction was made with DeepLoc-1.0 (http://www.cbs.dtu.dk/services/DeepLoc/).

Figure 12. Cartoon showing some structural features of the *Cryptosporidium* characteristic proteins (see text). UniProtKB IDs are reported in brackets. Proteins were divided into three categories according to the presence of hydrophobic motifs (*i. e.* presence/absence of a signal peptide and/or a transmembrane domain) and their presumptive localization respect to the cell membrane. Prediction of signal peptides and transmembrane domains was performed at Phobius (https://phobius.sbc.su.se/index.html). Prediction of other structural and functional domains was performed at Pfam (http://pfam.xfam.org) and Prosite (https://prosite.expasy.org). List of domains in the figure: Peptidase 1, PEPTIDASE_A1, PS51767; PLP: PROKAR_LIPOPROTEIN, PS51257; CAP, Cysteine-rich secretory protein family, CL0659; GP60, Glycoprotein GP60 of Cryptosporidium, PF11025; PPASE Tensin, PPASE_TENSIN, PS51181; Chase, CHASE, PS50839; DegV, DEGV, PS51482; MyB-L, MYB_LIKE, PS50090; LRR, Leucin Rich Repeat, LRR, PS51450; Thr, Threonin Rich Region, THR_RICH, PS50325; ARM, Armadillo/plakoglobin ARM repeat, ARM_REPEAT, PS50176;CS, CS, PS51203.

Functional Categories

- (3) Cytoskeleton 5%
- (1) Membrane protease 2%
- (14) Metabolism 23%
- (12) Unknown 20%
- (6) Regulatory 10%
- (2) Redox homeostasis 3%
- (1) Proteolysis 2%
- (9) Protein synthesis 15%
- (12) Chaperons and protein folding 20%

Subcellular localisation

- (4) Cell Membrane 7%
- (8) Uncertain 13%
- (2) Nucleus 3%
- (6) Extracellular 10%
- (6) Endoplasmatic reticulum 10%
- (34) Cytoplasm 57%

Cell membrane

CYTOPLASMIC

EXTRACELLULAR

GSPTSPGSPIFGAGTSGGGSRAPMPKFSGGSGSSPRSSITPLSPQR

ID2 antigenic peptide