Sequential Transitions of Male Sexual Behaviours Driven by Dual Acetylcholine-Dopamine Dynamics

Ai Miyasaka[1,2], Takeshi Kanda[1,3#], Naoki Nonaka[2#], Yuka Terakoshi[1], Yoan Cherasse[1], Yukiko Ishikawa[1], Yulong Li[4], Hotaka Takizawa[5,6], Jun Seita[2], Masashi Yanagisawa[1,7,8], Takeshi Sakurai[1,9,10*], Katsuyasu Sakurai[1*], Qinghua Liu[1,11,12*]

[1]International Institute for Integrative Sleep Medicine, University of Tsukuba, Tsukuba, Ibaraki 305-8575, Japan.

[2]Advanced Data Science Project, RIKEN Information R&D and Strategy Headquarters, RIKEN, Tokyo 103-0027, Japan.

[3]Department of Neurophysiology, Nara Medical University, Kashihara, Nara 634-8521, Japan

[4]State Key Laboratory of Membrane Biology, Peking University School of Life Sciences, Beijing 100871, China.

[5]Institute of Systems and Information Engineering, University of Tsukuba, Tsukuba, Ibaraki 305-8573, Japan.

[6]Academic Computing and Communications Center, University of Tsukuba, Tsukuba, Ibaraki 305-8577, Japan.

[7]Department of Molecular Genetics, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA

[8]Life Science Center for Survival Dynamics, Tsukuba Advanced Research Alliance, University of Tsukuba, Tsukuba, Ibaraki 305-8575, Japan

[9]Institute of Medicine, University of Tsukuba, Tsukuba, Ibaraki 305-8575, Japan.

[10]Life Science Center for Tsukuba Advanced Research Alliance (TARA), University of Tsukuba, Tsukuba, Ibaraki 305-8575, Japan.

[11]National Institute of Biological Sciences, Beijing (NIBS), Beijing 102206, China.

[12]Tsinghua Institute of Multidisciplinary Biomedical Research (TIMBR), Tsinghua University, Beijing 102206, China.

[#]These authors contributed equally to this work

[*]Correspondence: sakurai.takeshi.gf@u.tsukuba.ac.jp

sakurai.katsuyasu.gm@u.tsukuba.ac.jp

liuqinghua@nibs.ac.cn

Abstract

In mammals, males execute a stereotypical and organized sequence of sexual behaviours, such as mounting, intromission, and ejaculation, to successfully complete copulation. However, the neural mechanisms that govern the sequential transitions of male copulatory behaviours remain unclear. Here, we report that dopamine (DA) and acetylcholine (ACh) dynamics in the ventral shell of the nucleus accumbens (vsNAc) closely align with serial transitions of sexual behaviours in male mice. In particular, the vsNAc exhibits a unique pattern of 1.5–2.2 Hz dual ACh-DA rhythms that correspond to the pelvic thrust rhythm during intromission. The dual ACh-DA rhythms are generated locally by reciprocal regulation between ACh and DA signalling via nicotinic acetylcholine (nAChR) and dopamine D2 (D2R) receptors, respectively. Knockdown of choline acetyltransferase (ChAT) and D2R expression in the vsNAc diminishes intromission and ejaculation. We showed that ACh signalling promotes the initiation of intromission, whereas DA signalling sustains intromission by inhibiting the activities of D2R-expressing neurons in the vsNAc. Moreover, optogenetic activation of ChAT[vsNAc] neurons during intromission slows down the DA rhythm, a specific activity signature that precedes ejaculation, and leads to immediate ejaculation. Taken together, dual ACh-DA dynamics in the vsNAc coordinate sequential transitions of male copulatory behaviours from intromission to ejaculation.

Keywords: ventral shell of the nucleus accumbens (vsNAc), acetylcholine (ACh), dopamine (DA), dual ACh-DA rhythms, intromission, ejaculation

Introduction

In mammals, males engage in a stereotypical sequence of sexual behaviours when they encounter females in oestrous. This sequence of male sexual behaviours consists of courtship (appetitive) behaviours, such as sniffing and chasing, and followed by copulatory (consummatory) behaviours, such as mounting, intromission, and ejaculation in rodents[1–6]. Appropriate expression of sequential sexual behaviours is not only considered the attractivity of males, but also is important for the successful completion of sexual behaviours[7]. The neural mechanisms that regulate the sequential transitions of male sexual behaviours in the mammalian brain are largely unclear. For example, the nucleus accumbens (NAc) is thought to be involved in courtship behaviours, whereas the preoptic area (POA) has been implicated in copulatory behaviours[1,6]. However, recent studies indicated that POA primarily controls mounting[8–11]. Thus, the specific brain regions that regulate intromission and ejaculation are not yet identified, which hinders our understanding of the neural basis for controlling the intromission-to-ejaculation transition.

Accumulating studies have suggested that DA signalling functions as a primary driver for sexual motivation and behaviours[12–19]. Administration of cholinergic agonists in the preoptic area or substantia nigra changes the number or frequency of intromissions preceding ejaculation in rodents, implicating a regulatory role of ACh signalling during male copulatory behaviours[14,20,21]. It remains unclear, however, where and how precisely DA/ACh dynamics regulate male sexual behaviours[14–17,22–24]. Early microdialysis studies have detected an increase in extracellular DA concentration in the nucleus accumbens (NAc) during male sexual behaviours in rats[25–27]. Recent fibre photometry studies have recorded DA transients during intromission and ejaculation in male mice using the G protein-coupled receptor activation-based (GRAB)$_{DA}$ sensor with a high temporal resolution[15–17]. However, the regulatory mechanism and functional significance of DA dynamics in the NAc during male sexual behaviours remain unclear.

In this study, we hypothesise that ACh signalling modulates DA dynamics in the NAc to

regulate male sexual behaviours, based on recent reports that ACh signalling regulates DA release

from the dopaminergic axon terminals in the striatum[28–32]. By monitoring DA dynamics in various

subregions of the NAc using the $GRAB_{DA2m}$ sensor[15,16], we found that real-time DA dynamics in

the ventral shell of NAc (vsNAc) closely aligned with the sequential transitions of sexual

behaviours in male mice. Consistent with our hypothesis, we observed a unique signature of dual

ACh-DA rhythms in the vsNAc during intromission. Ex vivo brain slice imaging and

computational modelling revealed that the dual ACh-DA rhythms were generated by reciprocal

regulation between ACh and DA signalling via nAChR and D2R receptors, respectively. By

optogenetic manipulations of ACh and DA release into the vsNAc, we demonstrated that dual ACh

and DA dynamics in the vsNAc cooperatively drive the sequential transitions of male copulatory

behaviours.

Results

DA dynamics in the vsNAc closely align with serial transitions of male sexual behaviours

After sniffing and chasing receptive female mice, male mice engage in repeated cycles of

mounting and intromission, which culminate in an ejaculation followed by a brief falling and

immobile phase (Fig. 1a). To study DA dynamics in the NAc during male sexual behaviours, we

performed fibre photometry recording at the NAc of C57BL/6J male mice following injection in

the NAc with adeno-associated virus (AAV) expressing $GRAB_{DA2m}$ sensor in neurons under human

synapsin promotor (hSyn) (Fig. 1b). Since the NAc is anatomically heterogeneous[33–36], we

examined DA dynamics in different subregions of the NAc by inserting optic fibres into the core

(c), medial shell (ms), or ventral shell (vs) of the NAc, which was classified based on the

expression pattern of substance P (Fig. 1c, d)[37]. Distinct patterns of extracellular DA dynamics

were detected in the cNAc, msNAc, and vsNAc during male sexual behaviours (Fig. 1e-j). Both

vsNAc and cNAc, but not msNAc, exhibited an increase of $GRAB_{DA2m}$ signals that corresponded to

all of the sequential transitions of male sexual behaviours, from the initial encounter of female mice

to the final intromission-to-ejaculation transition (Fig. 1e-g; Supplementary Table 1). Among these,

only the vsNAc exhibited a 1-2 Hz rhythmic fluctuation of $GRAB_{DA2m}$ signals during intromission,

which appeared to be concurrent with the intromission thrust rhythm (Fig. 1h-j; Supplementary

Table 2). These observations identify the vsNAc as a candidate brain region where DA dynamics

may regulate the sequential transitions from intromission to ejaculation.

Local regulation of DA rhythm in the vsNAc during intromission

The ventral tegmental area (VTA) comprises a heterogeneous population of dopaminergic

neurons (DA^{VTA} neurons) that, depending on their genetic and anatomical characteristics, send

axonal projections to different target regions, such as in the NAc[33,35]. By two-colour cholera toxin

B subunit (CTB)-mediated retrograde labelling, we found that the vsNAc and msNAc received

projections from the anterolateral (al) VTA and the posteromedial (pm) VTA, respectively

(Extended Data Fig. 1a-c).

To compare the activities of the dopaminergic neurons in the alVTA and pmVTA during

male sexual behaviours, we unilaterally injected AAV-hSyn-DIO-GCaMP6s and inserted optical

fibres into the alVTA or pmVTA of *DAT-ires-Cre* (DAT^{Cre}) mice, respectively (Extended Data Fig.

1d). Consistent with the different projection patterns, fibre photometry recording revealed that

alVTA, but not pmVTA, neurons exhibited GCaMP (Ca^{2+}) dynamics in accordance with

$GRAB_{DA2m}$ dynamics in the vsNAc during male sexual behaviours (Extended Data Fig. 1e-h;

Supplementary Table 3). However, we did not observe rhythmic fluctuation of Ca^{2+} dynamics in

the alVTA or pmVTA during intromission (Extended Data Fig. 1e).

We also performed fibre photometry to monitor the neuronal activity in DA^{VTA} axons

projecting to the vsNAc ($DA^{VTA \to vsNAc}$) by injecting AAV-hSyn-DIO-GCaMP6s into the VTA and

inserting optical fibres into the vsNAc of DAT^{Cre} mice (Extended Data Fig. 1i). Unlike neuronal

activity in the somata of alVTA, the Ca^{2+} dynamics in $DA^{VTA \rightarrow vsNAc}$ axons was similar to

$GRAB_{DA2m}$ dynamics in the vsNAc, both in terms of high signal intensity and rhythmic fluctuations

during intromission (Extended Data Fig. 1j, k; Supplementary Table 4). By recording the 405 nm

isosbestic GCaMP control signals, we confirmed that the rhythmic GCaMP signals in the vsNAc

reflected real neuronal activity rather than motion noises during intromission (Extended Data Fig.

1k). These observations suggest that DA dynamics in the vsNAc is regulated locally within the

vsNAc and, to some extent, independent of the somata localised in the VTA.

Dual ACh-DA rhythms in the vsNAc during intromission

Previous studies have shown that ACh released from cholinergic interneurons promote DA

release from DA^{VTA} axons in the striatum[28–32]. By immunohistochemistry of choline

acetyltransferase (ChAT) in DAT^{Cre} mice, in which DA neurons were marked by GFP expression,

we confirmed the existence of cholinergic interneurons ($ChAT^{vsNAc}$) in the proximity of the DA^{VTA}

axons in the vsNAc (Extended Data Fig. 2a-c). To test our hypothesis that ACh signalling regulate

DA dynamics in the vsNAc during male sexual behaviours, we unilaterally injected AAV-hSyn-

DIO-GCaMP6s and inserted optic fibres into the vsNAc of *ChAT-ires-Cre* ($ChAT^{Cre}$) male mice

(Extended Data Fig. 2d-f; Supplementary Table 5). Fibre photometry recording revealed that

$ChAT^{vsNAc}$ neurons exhibited rhythmic activity during intromission (Extended Data Fig. 2f). Next,

we examined the ACh dynamics in the vsNAc by injecting AAV-hSyn-$GRAB_{ACh3.0}$ into the vsNAc

of C57BL/6J mice (Extended Data Fig. 2g-i; Supplementary Table 5). Similar to the rhythmic

fluctuation of $GRAB_{DA2m}$ signals, we also observed the rhythmic fluctuation of $GRAB_{ACh3.0}$ signals

during intromission (Extended Data Fig. 2i). These results are consistent with the idea that the

rhythmic activity of cholinergic interneurons may regulate the rhythmic release of ACh and

possibly DA in the vsNAc during intromission.

Next, we performed dual fibre photometry imaging to simultaneously monitor ACh and DA

dynamics after AAV-mediated expression of green fluorescent GRAB$_{ACh3.0}$ and red fluorescent

GRAB$_{rDA2m}$ in the vsNAc of C57BL/6J male mice (Fig. 2a). Importantly, we observed a specific

signature of dual ACh and DA rhythms during intromission (Fig. 2b). By the wavelet coherence

analysis, we found that the coherence between ACh and DA rhythms around 1-4 Hz band was

significantly increased during intromission (Fig. 2c, d, Extended Data Fig. 2j; Supplementary Table

6, 7). Both ACh and DA rhythms were characterised by a frequency of 1.5–2.2 Hz by fast Fourier

transform analysis (Fig. 2e). The ACh rhythm, but not DA rhythm, appeared just before the onset

of intromission: the 1.5–2.2 Hz power of ACh rhythm was significantly higher than that of DA

rhythm from 6-s before to 2-s after the intromission onset (Fig. 2f; Supplementary Table 6).

To assess the time-lag between ACh and DA rhythms during intromission, we compared the

ACh/DA rhythm with the pelvic thrust rhythm by analysing the dynamic changes in the body size

of male and female mice in the video (Fig. 2g). By plotting the ACh and DA rhythms around every

thrusting event, we found the trough of ACh rhythm was ~0.1 s before, whereas that of DA rhythm

was ~0.1–0.2 s after the thrusting (Fig. 2h). Moreover, the ACh rhythm, but not DA rhythm,

persisted during the short pauses of pelvic thrusting during long intromissions (Fig. 2g). Taken

together, these observations suggest that the ACh rhythm may function as the primary driver to

generate the dual ACh-DA rhythm during intromission (Fig. 2e-h).

Signatures of the intromission preceding ejaculation (IPE)

While the majority of intromissions do not lead to ejaculation (intromission only (IO)), the

last intromission typically transitioned into ejaculation (intromission preceding ejaculation (IPE))

during a successful cycle of sexual behaviours (Fig. 2i). Thus, a comparison of dual ACh-DA

dynamics between IO and IPE may provide insights into the neural basis for controlling the

intromission-to-ejaculation transition. Interestingly, the coherence of the dual ACh-DA rhythms

was significantly enhanced at the intromission onset of IPE relative to IO (Extended Data Fig. 2k, l;

Supplementary Table 7). While the dual 1.5–2.2 Hz ACh-DA rhythms continued till the

termination of IO (Fig. 2j), there was a specific slowdown of the 1.5–2.2 Hz DA rhythm just before

ejaculation that coincides with the big surge of DA release in the IPE (Fig. 2k). Importantly, this

brief suppression of DA rhythm was only observed in the vsNAc (Fig. 2l; Supplementary Table 6),

not in the msNAc or cNAc, during the offset of IPE (Extended Data Fig. 2m; Supplementary Table

8), suggesting that it may be a prerequisite for the intromission-to-ejaculation transition.

ACh signalling regulates DA release in the vsNAc

To examine whether ACh signalling could regulate DA release from DA^{VTA} axons in the

vsNAc, we injected AAV-hSyn-FLEX-ChrimsonR and AAV-hSyn-GRAB$_{DA2m}$ into the vsNAc of

ChAT^{Cre} mice followed by ex vivo NAc brain slice imaging coupled with optogenetics and

pharmacological treatments (Fig. 3a-c). Optogenetic activation of ChAT^{vsNAc} neurons evoked an

immediate and robust increase in GRAB$_{DA2m}$ signals in the NAc slices (Fig. 3b, c). This cholinergic

stimulation of DA release was suppressed by nAChR antagonist DhβE at all frequencies (Fig. 3b,

c). Similarly, the cholinergic activation of DA release was suppressed by mAChR antagonist

atropine during 20 or 40 Hz stimulation of ChAT^{vsNAc} neurons (Fig. 3b, c). In response to 1 or 10

Hz stimulation of ChAT^{vsNAc} neurons, however, the blockade of mAChR induced a sharp increase

of GRAB$_{DA2m}$ signals (Fig. 3b, c), possibly owing to the disinhibition of mAChR-mediated

suppression of ACh release[38]. Accordingly, in situ hybridisation detected the expression of all five

mAChR1-5 (*Chrm1-5*) in the majority of ChAT^{vsNAc} neurons (Extended Data Fig. 3;

Supplementary Table 9). In addition, DA^{VTA} neurons expressed both nAChR[39] and mAChR1-5

(Extended Data Fig. 3; Supplementary Table 9). These results suggest that ACh can promote DA

release via nAChR signalling, but may also inhibit DA release in some contexts via mAChR

signalling in the vsNAc.

DA signalling reciprocally regulates ACh release in the vsNAc

To examine whether DA signalling reciprocally regulated the activity of ChAT^{vsNAc} neurons, we performed ex vivo Ca^{2+} imaging with NAc slices following injection of AAV-Syn-FLEX-ChrimsonR into the VTA and AAV-hSyn-DIO-GCaMP6s into the vsNAc of *ChAT^{Cre};DAT^{Cre}* mice (Fig. 3a, d, e). Optogenetic (1 Hz) stimulation of DA$^{VTA \rightarrow NAc}$ axons evoked different responses in the ChAT^{NAc} neurons: ~20% increased, ~30% decreased, and ~50% exhibited no change of Ca^{2+} dynamics (Fig. 3e). By contrast, higher ($\geq$10 Hz) frequency stimulation of DA$^{VTA \rightarrow NAc}$ axons increased the activity of ~40% ChAT^{NAc} neurons and decreased the activity of ~30% ChAT^{NAc} neurons (Fig. 3e). Notably, a small subset of ChAT^{NAc} neurons displayed rhythmic fluctuation of Ca^{2+} dynamics in the NAc slices (Fig. 3d), which was consistent with the rhythmic fluctuation of Ca^{2+} dynamics in ChAT^{vsNAc} neurons and rhythmic release of ACh in the vsNAc observed by in vivo fibre photometry during intromission (Extended Data Fig. 2d-i).

The observation of both *Drd1^{+}*ChAT^{+} and *Drd2^{+}*ChAT^{+} neurons in the vsNAc (Fig. 3f, g; Supplementary Table 10) raised two potential mechanisms for the regulation of ChAT^{vsNAc} neurons by DA signalling: 1) inhibition of ChAT^{vsNAc} neurons via D2R-mediated Gi/o signalling; 2) activation of ChAT^{vsNAc} neurons via D1R-mediated Gs signalling[40]. To probe these possibilities, we injected AAV-Syn-FLEX-ChrimsonR and AAV-hSyn-GRAB$_{ACh3.0}$ into the vsNAc of *ChAT^{Cre}* mice (Fig. 3a, h-k). Optogenetic activation of ChAT^{vsNAc} neurons increased GRAB$_{ACh3.0}$ signals in the NAc slices, which was suppressed by D2R agonist bromocriptine under all conditions except for 40 Hz stimulation (Fig. 3h, i). Thus, the D2R-mediated inhibition of ChAT^{NAc} neurons could suppress the release of ACh from ChAT^{NAc} neurons following $\leq$20 Hz optogenetic stimulation (Fig. 3i). Next, we investigated whether ACh release was also modulated by D1R signalling in the NAc slices by application of D1R agonist SKF38398. Activation of D1R signalling caused distinct changes in ACh release in the NAc slices, with 12.1% (up), 50.4% (down), and 37.5% (no change)

in GRAB$_{Ach3.0}$ signals (Fig. 3j, k). Taken together, these results suggest the reciprocal regulation of

ACh release by DA signalling in the vsNAc, which is potentially mediated by D1R or D2R

expressed on ChAT^{vsNAc} neurons.

Modelling of dual Ach-DA rhythms by reciprocal regulation between ACh and DA signalling

To further study how the dual ACh-DA rhythms were generated by reciprocal regulation

between ACh and DA in the vsNAc during intromission, we performed computational modelling

using a simplified neuronal assemblage consisting of the dopaminergic ($DA_{sim}^{VTA \to vsNAc}$) axon

terminals, cholinergic neurons ($ChAT_{sim}^{vsNAc}$), and the extracellular environment (Fig. 4a, Extended

Data Fig. 4a-e). In this modelling, we defined the release probability measurement (RPM) of the

neuronal assemblage as a measure of the intensity of neuronal activity (Fig. 4b). We set

$Activation$ as a constant value to mimic the external stimuli, such as the spontaneous firing of

each neuronal assemblage or stimulation by other neurons (Extended Data Fig. 4a, b). Additionally,

the $Receptor$ function represents the effect of each receptor upon binding of ACh or DA (Fig. 4b).

Both $Activation$ and $Receptor$ constantly change the RPM that determines the volume of

neurotransmitter (NT) released to the extracellular environment (Fig. 4b).

The binding of DA to different dopamine receptors activates distinct intracellular G protein

signalling to regulate neuronal activity: D1R triggers Gs signalling to activate neuronal activity,

whereas D2R activates G protein-gates inwardly rectifying potassium (GIRK) channels-mediated

signalling to inhibit neuronal activity[40–42]. On the other hand, the binding of ACh to nAChR causes

a rapid influx of cations, such as Na$^+$ and Ca^{2+}, to activate neuronal activity[43,44]. Herein, we focused

on nAChR rather than mAChR1-5 to simplify our modelling to explore the characteristics of D1R

and D2R relative to nAChR. We hypothesised that the characteristics of each receptor consist of

two axes: the effect intensity (effect: RE) with which the receptor changes neuronal activity; the

processing speed (speed: RS) of the receptor in affecting the neuronal activity upon binding of ACh

or DA (Fig. 4b).

Cholinergic activation is required for modelling dual ACh-DA rhythms

The goal of the modelling is to estimate the RE and RS values of each receptor to minimize

the $Diff(NT_{sim}, NT_{real})$, i.e., the difference between simulated signals (NT_{sim}) and

experimentally recorded in vivo GRAB signals (NT_{real}). To simplify the model, both the RS and

RE values of the nAChR were defined as 1, and the RS and RE values of D1R and D2R relative to

the nAChR were explored (Fig. 4a, b). Based on the ex vivo imaging (Fig. 3) and in situ

hybridisation results (Extended Data Fig. 3) and previous report of D2R expression in

dopaminergic neurons[45], we placed D1R and D2R on $ChAT_{sim}^{vsNAc}$ neurons and nAChR and D2R on

$DA_{sim}^{VTA \to vsNAc}$ axon terminals in this model (Fig. 4a). Importantly, we found that the simulated dual

rhythms of ACh and DA closely resembled the experimentally recorded ACh and DA rhythms

(Fig. 4c).

To investigate the relative contribution of $ChAT_{sim}^{vsNAc}$ neurons and $DA_{sim}^{VTA \to vsNAc}$ axon

terminals, we compared the similarity between NT_{real} and NT_{sim} ($Diff(NT_{sim}, NT_{real})$) in three

activation patterns: $ChAT_{sim}^{vsNAc}$ and $DA_{sim}^{VTA \to vsNAc}$ ($Activation_{(1,1)}$); only $ChAT_{sim}^{vsNAc}$

($Activation_{(1,0)}$); only $DA_{sim}^{VTA \to vsNAc}$ ($Activation_{(0,1)}$) (Extended Data Fig.4a, b). The similarity

between NT_{real} and NT_{sim} rhythms suggested that the $Activation_{(1,1)}$ and $Activation_{(1,0)}$

conditions were much better than the $Activation_{(0,1)}$ condition to simulate dual ACh-DA rhythms

(Fig. 4d; Supplementary Table 11). These observations suggest that the activation of cholinergic

neurons is necessary to generate dual ACh-DA rhythms.

D2R, but not D1R, is required for modelling dual ACh-DA rhythms

We visualised the explored values of the RS and RE for each assumed receptor during the simulation. Notably, the RS of D2R was in the range of 5–50 ms, whereas the RS of D1R was approximately 350–400 ms (Fig. 4e), indicating that the processing speed of D2R is approximately 7–80 fold faster than that of D1R in our model. This finding is consistent with the fact that the GIRK-mediated D2R signalling is much faster than Gs-mediated D1R signalling. On the other hand, the RE of D1R or D2R on $ChAT_{sim}^{vsNAc}$ neurons are approximately 1/100 or 1/50 of the RE of nAChR on $DA_{sim}^{VTA \to vsNAc}$ axon terminals, respectively. By contrast, the RE of D2R on $DA_{sim}^{VTA \to vsNAc}$ axon terminals is about 2 to 4 times of the RE of nAChR (Fig. 4f), such that the effect intensity of D2R on $DA_{sim}^{VTA \to vsNAc}$ axon terminals is hundreds fold larger than that of D2R on $ChAT_{sim}^{vsNAc}$ neurons in our model.

To determine the receptors essential for generating the dual ACh-DA rhythms, we performed simulations to evaluate the similarity between NT_{sim} and NT_{real} following ablation of each receptor (Fig. 4g, Extended Data Fig. 4f, g). Consistent with estimation that D2R has larger RE than D1R, the ablation of D2R, but not D1R, on $ChAT_{sim}^{vsNAc}$ neurons worsens the similarity between NT_{sim} and NT_{real} as compared with the full receptor condition (Fig. 4g; Supplementary Table 12). Similarly, the ablation of D2R or nAChR on $DA_{sim}^{VTA \to vsNAc}$ axon terminals significantly worsened the similarity between NT_{sim} and NT_{real}. These observations underscore the importance of D2R-mediated DA signalling and nAChR-mediated ACh signalling in the generation of dual ACh-DA rhythms.

Knockdown of ChAT and D2R in the vsNAc diminishes intromission and ejaculation

To validate our simulation results in vivo, we performed RNAi-mediated knockdown of ChAT, D1R, or D2R expression by injecting the vsNAc of male mice with AAV expressing short hairpin (sh)RNA targeting *Chat*, *Drd1* or *Drd2* mRNA (Fig. 4h). In situ hybridisation verified

efficacy of knockdown of target gene expression in the vsNAc (Fig. 4i-k, Extended Data Fig. 5a;

Supplementary Table 13). After 3 weeks post AAV-shRNA injection, the sexual behaviour tests

were conducted over a period of 3 weeks (Fig. 4h, Extended Data Fig. 5b-h). Consistent with the

simulation results of in silico receptor ablation (Fig. 4g), knockdown of ChAT or D2R, but not

D1R, expression in the vsNAc diminished the frequency of sexual behaviours, such as intromission

and ejaculation, relative to the scrambled shRNA-expressing mice (Fig. 4l, m; Supplementary

Table 14). Moreover, knockdown of ChAT or D2R prolonged the average latency of mounting,

intromission, and ejaculation, whereas knockdown of D2R reduced the average number of bouts

and total duration of mounting and intromission (Extended Data Fig. 5; Supplementary Table 15).

Importantly, neither ChAT nor D2R knockdown affected the motor coordination of the mice as

measured by the rotarod test (Fig. 4n; Supplementary Table 16), excluding the possibility that male

sexual behaviours were affected indirectly by impairing motor coordination. These results indicate

that both ACh and DA signalling play important roles during male copulatory behaviours. Based on

the results of ex vivo brain slice imaging, computational modelling, and RNAi knockdown

experiments, we concluded that the dual ACh-DA rhythms are likely generated in the vsNAc by

reciprocal ACh-DA regulations mediated by nAChR and D2R signalling in $DA^{VTA \rightarrow vsNAc}$ axon

terminals and D2R signalling in $ChAT^{vsNAc}$ neurons, respectively.

High level of DA signalling in the vsNAc is important for the maintenance of intromission

A high level of extracellular DA was observed in the vsNAc, and only the vsNAc exhibited

the 1.5–2.2 Hz DA rhythm during intromission (Fig. 1). To examine whether DA signalling plays a

crucial role during intromission, we optogenetically manipulated the $DA^{VTA \rightarrow vsNAc}$ axon terminals

to investigate whether changing DA dynamics in the vsNAc could affect male copulatory

behaviours. Specifically, we bilaterally injected AAV-Ef1a-DIO-ChR2 or AAV-Ef1a-DIO-

eNpHR3.0 into the VTA and inserted optic fibres into the vsNAc of DAT^{Cre} mice (Fig. 5a). Notably,

optogenetic activation of the $DA^{VTA \to vsNAc}$ axons either before intromission (during sniffing, chasing, or mounting) or during intromission (after two pelvic thrust movements) increased the total duration of intromission (Fig. 5b-e, Extended Data Fig. 6; Supplementary Table 17, 18). Additionally, optogenetic activation before intromission also increased the number of intromission bouts, although it did not immediately trigger intromission (Fig. 5b, Extended Data Fig. 6b). Conversely, optogenetic suppression of the $DA^{VTA \to vsNAc}$ axon terminals before or during intromission did not affect male sexual behaviours, possibly owing to incomplete suppression of DA signalling (see discussion). These results suggest that high level of DA signalling in the vsNAc promotes the initiation and maintenance of intromission.

DA-mediated inhibition of $D2R^{vsNAc}$ neurons sustains male sexual behaviours, whereas suppression of $D1R^{vsNAc}$ neurons ensures forward progression of sexual behaviours

To elucidate how DA dynamics regulate activity of D1R- and D2R-expressing neurons in the vsNAc during male sexual behaviours, we unilaterally injected AAV-hSyn-DIO-GCaMP6s and inserted optic fibres in the vsNAc of *Drd1-ires-Cre* (*Drd1^Cre*) and *Drd2-ire-Cre* (*Drd2^Cre*) mice, respectively (Fig. 5f-i, Extended Data Fig. 7a, b; Supplementary Table 19, 20). Fibre photometry recording revealed that the activity of both $D1R^{vsNAc}$ and $D2R^{vsNAc}$ neurons were suppressed during intromission (Extended Data Fig. 7a, b), despite the opposite effects of D1R (excitatory) and D2R (inhibitory) signalling on neuronal activity[40–42]. The simplest explanation for this paradoxical observation is that DA signalling preferentially activates the D2R-mediated inhibition of neuronal activities during intromission owing to the different thresholds for D1R and D2R activation[46,47,12] (see discussion).

To study the functional importance of suppression of $D1R^{vsNAc}$ and $D2R^{vsNAc}$ neurons during intromission, we specifically expressed ChR2 in the $D1R^{vsNAc}$ or $D2R^{vsNAc}$ neurons by injecting AAV-Ef1a-DIO-ChR2 into the vsNAc of *Drd1^Cre* and *Drd2^Cre* mice, respectively (Fig. 5j-

p, Extended Data Fig. 7c-f; Supplementary Table 21, 22). While optogenetic activation of D1R^{vsNAc}

neurons before intromission did not affect male sexual behaviours, optogenetic stimulation of these

neurons during intromission increased the total mounting duration by inducing repetitive mounting

behaviours (Fig. 5k-p, Extended Data Fig. 7c, d). Thus, the suppression of D1R^{vsNAc} neurons during

intromission is critical for ensuring the forward progression of male sexual behaviours.

On the other hand, optogenetic activation of D2R^{vsNAc} neurons during sniffing, chasing,

mounting, and intromission immediately stopped any of these sexual behaviours (Fig. 5k, l).

Consequently, opto-stimulation of D2R^{vsNAc} neurons before intromission increased total mounting

duration and prolonged ejaculation latency (Fig. 5m, n), whereas opto-stimulation of D2R^{vsNAc}

neurons during intromission decreased total intromission duration (Fig. 5o, Extended Data Fig. 7e,

f). These observations indicate that DA-mediated suppression of D2R^{vsNAc} neurons is necessary for

sustaining male sexual behaviours including intromission (Fig. 5q).

ACh signalling in the vsNAc promotes sequential transitions of intromission and ejaculation

To study how ACh signalling in the vsNAc regulated male sexual behaviours, we bilaterally

injected AAV-Ef1a-DIO-ChR2 or AAV-Ef1a-DIO-eNpHR3.0 and inserted optic fibres into the

vsNAc of *ChAT*^{Cre} male mice (Fig. 6a). Next, we optogenetically activated or inhibited the ChR2-

or eNpHR3.0-expressing ChAT^{vsNAc} neurons, respectively, before or during intromission and

examined the effects on male copulatory behaviours (Fig. 6b-g, Extended Data Fig 8). While

optogenetic activation of ChAT^{vsNAc} neurons before intromission did not affect mating behaviours,

optogenetic inhibition of ChAT^{vsNAc} neurons before intromission significantly prolonged the

latency of mounting, intromission, and ejaculation (Fig. 6b, d, e, Extended Data Fig 8a;

Supplementary Table 23). These results suggest that ACh signalling in the vsNAc is required for

the initiation of mounting and intromission.

Notably, optogenetic activation of ChAT^{vsNAc} neurons during intromission induced rapid

ejaculation by shortening the ejaculation latency and increasing the ejaculation likelihood (Fig. 6c,

f, g; Supplementary Table 24). Moreover, the number of mounting or intromission bouts and the

mounting total duration were also decreased (Extended Data Fig 8g-i). In contrast, optogenetic

inhibition of ChAT^{vsNAc} neurons during intromission did not affect mating behaviours (Fig. 6f, g,

Extended Data Fig 8g-k; Supplementary Table 24). These results indicate that activation of ACh

signalling in the vsNAc triggers the intromission-to-ejaculation transition.

To study how activation of ACh signalling changed DA dynamics during intromission, we

injected AAV-hSyn-GRAB$_{DA2m}$ and AAV-Syn-FLEX-ChrimsonR into the vsNAc of *ChAT^{Cre}* mice

and monitored DA dynamics by fibre photometry during optogenetic stimulation of ChAT^{vsNAc}

neurons (Fig. 6h-j). Interestingly, optogenetic activation of ChAT^{vsNAc} neurons during intromission

resulted in a decrease of the 1.5–2.2 Hz power of the DA rhythm followed by a brief suppression of

DA release, which resembled the characteristic slowdown of DA rhythm just before ejaculation

during IPE (Fig. 6k, l; Supplementary Table 25). Taken together, these results suggest that

activation of ACh signalling in the vsNAc during intromission leads to a slowdown of DA rhythm

followed by immediate ejaculation (Fig. 6m).

Discussion

Dual ACh-DA dynamics in the vsNAc drive sequential transitions of male copulatory behaviours

The neural mechanisms that regulate the sequential transitions of male copulatory

behaviours, such as mounting, intromission, and ejaculation, remain unclear. Previous studies have

identified specific brain regions that control mounting[8–11], but the brain regions that regulate

intromission and ejaculation are to be elucidated. Here, we identified the vsNAc as a critical brain

region where dual ACh-DA dynamics aligned with and coordinate the sequential transitions from

intromission to ejaculation. In particular, the vsNAc exhibits a unique signature of 1.5–2.2Hz dual

ACh-DA rhythms that are concurrent with the pelvic thrust rhythm during intromission. The ACh

rhythm, but not DA rhythm, appears before the intromission onset and persists during the short

pauses of pelvic thrusting during intromission. These observations suggest that ACh rhythm

functions as the primary driver for generating the dual ACh-DA rhythms during intromission. We

proposed that dual ACh-DA rhythms play a crucial role in the maintenance of intromission possibly

by reinforcing the pelvic thrust rhythm. By optogenetic manipulations of ACh and DA release in

the vsNAc, we showed that ACh signalling promoted the initiation of intromission, whereas DA

signalling was important for sustaining intromission through the suppression of $D2R^{vsNAc}$ neurons.

Furthermore, optogenetic activation of $ChAT^{vsNAc}$ neurons during intromission elicits immediate

ejaculation by triggering a slowdown of DA rhythm–a specific activity signature that typically

precedes ejaculation during IPE. Taken together, these results demonstrate that dual ACh-DA

dynamics harmonise in the vsNAc to drive the sequential transitions of male copulatory behaviours

from intromission to ejaculation.

Dual ACh-DA rhythms are generated by reciprocal regulations between ACh and DA release

Both $ChAT^{vsNAc}$ neurons and $DA^{VTA \to vsNAc}$ axons, but not DA^{VTA} somata, exhibited

rhythmic neuronal activity during intromission, suggesting that the dual ACh-DA rhythms are

regulated locally by mutual interaction between $ChAT^{vsNAc}$ neurons and $DA^{VTA \to vsNAc}$ axons.

Consistent with this idea, we showed reciprocal regulation between ACh and DA release mediated

by nAChR and D1R or D2R signalling in the NAc brain slices by ex vivo (GRAB or GCaMP)

imaging with optogenetic and pharmacological manipulations. Computational modelling suggests

that the dual ACh-DA rhythms in the vsNAc are generated by reciprocal ACh-DA regulations via

D2R signalling in cholinergic neurons and D2R and nAChR signalling in dopaminergic axon

terminals. Accordingly, the shRNA-mediated knockdown of ChAT and D2R, but not D1R,

expression in the vsNAc diminished the probability of sexual behaviours, such as intromission and

ejaculation. Although we primarily focused on nAChR, D1R, and D2R to simplify our modelling

of the dual ACh-DA rhythms, we could not exclude a potential role for mAChRs in the generation

of dual ACh-DA rhythms during intromission. A recent study also reported the dual ACh-DA (~2

Hz) rhythms in the dorsal striatum during reward delivery, but these rhythms were not generated by

direct interaction between ACh and DA release within the striatum[28]. Thus, it is possible that the

dual ACh-DA rhythms can be generated by different mechanisms to regulate diverse biological

processes.

Distinct functions of D1R and D2R signalling during male sexual behaviours

Our finding that D1R^{vsNAc} and D2R^{vsNAc} neurons, which are largely distinct neuron

populations, were both suppressed during intromission is consistent with the proposed "threshold"

hypothesis for D1R and D2R-mediated DA signalling during sexual behaviours[12,47]. Since D2R has

a higher affinity than that of D1R[46], DA signalling may preferentially cause the D2R-mediated

suppression of D2R^{vsNAc} neurons, while D1R^{vsNAc} neurons are not activated during intromission

(Supplementary Table 26). Alternatively, it is plausible that suppression of D2R^{vsNAc} neurons may

indirectly lead to active suppression of D1R^{vsNAc} neurons through unknown mechanisms.

Importantly, optogenetic activation of D2R^{vsNAc} neurons either before or during intromission

rapidly stopped the sexual behaviours. It is plausible that activation of D2R^{vsNAc} neurons is

associated with aversive emotions[48,49], which may function as an "off" switch to pause male sexual

behaviours. Thus, the DA-mediated suppression of D2R^{vsNAc} neurons is necessary for sustaining

male sexual behaviours such as intromission.

The highest level of DA release was observed at the onset of female encounter or during

ejaculation, which might cause the activation of D1R^{vsNAc} neurons to regulate the initiation and

termination of male sexual behaviours. Optogenetic activation of D1R^{vsNAc} neurons during

intromission increased the total mounting duration by inducing repetitive mounting behaviour. It is

plausible that optogenetic stimulation of D1R^{vsNAc} neurons mimics the activation of D1R^{vsNAc}

neurons at female encounter and, thus, resetting the sexual behaviours. Therefore, the suppression

of D1R^{vsNAc} neurons during intromission ensures the forward progression of male sexual

behaviours by preventing animals from returning to the previous behavioural sequences, such as

mounting.

ACh signalling regulates ejaculation timing by triggering slowdown of DA rhythm

We observed a specific slowdown of DA rhythm just before ejaculation that coincides with

a big surge of DA release during IPE. Accordingly, optogenetic activation of ChAT^{vsNAc} neurons

resulted in a similar slowdown of DA rhythm followed by immediate ejaculation. This

optogenetics-induced suppression of DA rhythm was even stronger than the suppression of DA

rhythm just before ejaculation during IPE (Fig. 6l). By contrast, optogenetic inhibition of

DA$^{VTA\rightarrow vsNAc}$ axons during intromission did not elicit ejaculation (Supplementary Table 27). In the

striatum, cholinergic interneurons make monosynaptic or polysynaptic axo-axonic contact on

DA^{VTA} axons[30,38]. ACh released from cholinergic interneurons can evoke action potentials at the

DA^{VTA} axons and promote DA release from proximal and distant DA axons via the action potential

propagation[29,31]. Therefore, optogenetic activation of ChAT^{vsNAc} neurons may have a larger effect

on DA dynamics in the vsNAc than direct optogenetic manipulation of DA$^{VTA\rightarrow vsNAc}$ axons.

Moreover, optogenetic stimulation of ChAT^{vsNAc} neurons results in a high level of ACh signalling,

which then leads to a brief suppression of DA signalling, both of which may be required for

inducing ejaculation.

It should be noted that pharmacological inhibition of mAChRs enhanced the release of DA

following 1 or 10 Hz opto-stimulation of ChAT^{vsNAc} neurons in the NAc brain slices, suggesting

that mAChRs could suppress DA release in specific contexts of ACh signalling. Optogenetic

stimulation of ChAT^{vsNAc} neurons may cause a surge of ACh signalling, leading to the suppression

of DA release by activating the inhibitory mAChRs on DA^{VTA} axons. Similarly, we observed that

the coherence of ACh-DA rhythms between 1–4 Hz frequency was significantly higher at the onset

of IPE than IO. This increased coherence of dual ACh-DA rhythms probably results in a higher

level of ACh signalling, which in turn may cause the suppression of DA rhythm by activating the

inhibitory mAChRs. Future studies are warranted to elucidate the detailed mechanism for why dual

ACh-DA rhythms specifically results in the slowdown of DA rhythm and triggers the intromission-

to-ejaculation transition during IPE, but not during IO.

To the best of our knowledge, this study is the first report of ejaculation induction by

manipulating a small population of genetically defined neurons in freely behaving animals. Our

results reveal, for the first time, a crucial role of ACh signalling in regulating the timing of

ejaculation in mammals. Premature ejaculation (PE) is a prevalent sexual disorder affecting

approximately 20–30% of sexually active men[50]. Moreover, PE has been observed in relatively

high percentages in Parkinson's disease patients, which is thought to be the result of the

medications and the disease itself, with DA signalling being the common contributing factor[51].

Selective serotonin reuptake inhibitors are also used to treat patients with PE[52]. Although the exact

mechanism is not yet fully understood, serotonin can affect the activity of dopamine neurons and

DA release[53]. Therefore, these drugs may affect DA dynamics and, in turn, regulate ejaculation

latency. We anticipate that our findings will be a starting point for more sophisticated studies into

the molecular and neural mechanisms that govern ejaculation timing and the potential development

of new therapeutics for sexual dysfunctions in humans.

Methods

Mice

All animal experiments were performed at the International Institute for Integrative Sleep Medicine, University of Tsukuba, in accordance with the guidelines for animal experimentation. The experiments were approved by the animal experiment committee of each institute and were conducted in accordance with NIH guidelines. Mice were given food and water *ad libitum* and maintained at a temperature of 23 °C and relative humidity of 50%, with a 12-h light/12-h dark cycle. C57BL/6J mice (ID# 000664 from Charles River), male *ChAT*[Cre] mice (B6;129S6-Chattm2(cre)Lowl/J; ID# 006410, Jaxson Laboratory)[54], male *DAT*[Cre] mice (B6.SJLSlc6a3tm1.1(cre)Bkmn/J; ID# 006660, Jaxson Laboratory)[55], male *Drd1*[Cre] mice (B6.FVB(Cg)-Tg(Drd1-cre)EY262Gsat/Mmucd; ID# 030989-UCD from MMRRC originally from GENSAT BAC Tg project), and male *Drd2*[Cre] mice (B6.FVB(Cg)-Tg(Drd2-cre)ER44Gsat/Mmucd; ID# 032108-UCD from MMRRC originally from GENSAT BAC Tg project) were used in this study. The ages of the mice used for the experiments were 8 to 32 weeks.

Viruses

The plasmids of AAV-hSyn-DIO-GCaMP6s (serotype 9), AAV-Syn-FLEX-rc[ChrimsonR-tdTomato] (serotype 1), AAV-hSyn-DIO-EGFP (serotype 9), AAV-EF1a-double floxed-hChR2(H134R)-EYFP-WPRE-HGHpA (serotype 9), AAV-Ef1a-DIO-EYFP (serotype 9), and AAV-Ef1a-DIO eNpHR3.0-EYFP (serotype 9) were purchased from Addgene. The plasmids of AAV[shRNA]-CMV>mCherry-U6>Scramble_shRNA (serotype 9), AAV[shRNA]-mCherry-U6>mChat[shRNA#1] (serotype 9), and AAV[shRNA]-mCherry-U6>*Drd1*[shRNA#1] (serotype 9) were purchased from vector-builder. AAV-H1-shScramble-4-CMV-mCHERRY (serotype 10) and AAV-H1-shD2R-4-CMV-mCHERRY (serotype 10) were shared by the University of Tsukuba. The plasmids of AAV-hSyn-GRAB_DA2m (GRAB$_{DA2m}$) (serotype 9), AAV-hSyn-GRAB_ACh3.0

496 (GRAB$_{ACh3.0}$) (serotype 9), and AAV-hSyn-rDA2m (GRAB$_{rDA2m}$) (serotype 9) were provided by

497 Yulong Li. All AAVs were prepared at the University of Tsukuba. AAV titres were 9.79×10^{11} –

498 6.24×10^{14} genomic copies/mL.

Stereotaxic surgery

AAVs were produced using the triple-transfection helper-free method and purified as

previously described[56]. The titres of AAVs were determined by qPCR and purified viruses were

stored in aliquots at -80°C. Details of AAVs are listed in Supplementary Table 28. Adult male mice

(aged > 8 weeks) were stereotactically injected with AAVs and implanted with optical fibres after

at least one week post AAV injection. The animals were anaesthetised with isoflurane (2–4%) and

placed in a stereotaxic frame (David Kopf Instruments, Tujunga, CA). Viruses were stereotaxically

delivered to the brains of mice at 2–5 months of age as described previously[57]. In brief, virus was

delivered unilaterally (0.2 µL of AAV-hSyn-GRAB$_{DA2m}$, AAV-hSyn-DIO-GCaMP6s, AAV-hSyn-

GRAB$_{ACh3}$, and AAV-hSyn-GRAB$_{rDA2m}$) or bilaterally (0.3 µL of AAV-Syn-FLEX-rc[ChrimsonR-

tdTomato], AAV-hSyn-DIO-EGFP, AAV-EF1a-double floxed-hChR2(H134R)-EYFP-WPRE-

HGHpA, AAV-Ef1a-DIO EYFP, AAV-Ef1a-DIO eNpHR3.0-EYFP. Immediately after surgery,

mice were placed on a heated pad individually and then returned to their home cage following

recovery from anaesthesia. Animals were allowed at least 2 weeks of recovery in a reversed

light/dark room following surgery prior to being tested in behavioural assays.

For retrograde tracing, we then injected 0.3 µL of the retrograde tracer cholera toxin subunit

B (CTB) conjugated with AlexaFluor-594 and AlexaFluor-488 (Thermo Fisher Scientific) into the

msNAc and vsNAc. Mice were perfused 5 days after CTB injection. Whole brain sections were

prepared.

Immunohistochemistry

For histological analysis, animals were perfused with 4% paraformaldehyde, and the brains

were dissected and post-fixed overnight in 4% paraformaldehyde. Brains were sectioned at 60 μm

thickness using a vibrating microtome (Leica) and immunostaining was performed as described

previously[58]. In brief, sections were collected in PBS and were treated with permeabilisation

solution (1% Triton in PBS) at room temperature for 3 h, followed by incubation with the

blocking buffer (10% Blocking One [Nacalai Tesque] with 1% Triton X-100 in PBS), and

incubated overnight at 4 °C using primary antibodies in the blocking buffer. After washing with

PBS three times, brain slices were incubated with secondary (second) antibody in blocking buffer

at 4 °C overnight. The slices were stained with NeuroTrace fluorescent Nissl stain (Invitrogen, N-

21279) or 4', 6-diamidino-2-phenylinodole (DAPI) (Dojindo, D523), washed, mounted, and

coverslipped.

Primary antisera used were chicken anti-GFP (Novus Biologicals; 1:4,000), rat anti-GFP

(Nacalai Tesque; 1:2,000), rat anti-Substance P (Merck & Co., Inc.; 1:500), rabbit anti-Substance P

(ImmunoStar Inc.; 1:1500), goat anti-mCherry (Sicgen; 1:2,000), rat anti-mCherry (Thermo Fisher

Scientific; 1:500), goat anti-ChAT (Merck & Co., Inc.; 1:500), mouse anti-TH (Santa Cruz

Biotechnology; 1:250), sheep anti-Digoxigenin-POD (Roche Applied Science; 1:2,000), and sheep

anti-Fluorescein-POD (Roche Applied Science; 1:2,000).

Secondary antisera used were as follows: Alexa Fluor 488 donkey anti-rat (Jackson

ImmunoResearch; 1:500), Alexa Fluor 488 donkey anti-rabbit (Jackson ImmunoResearch; 1:500),

Alexa Fluor 488 donkey anti-chicken (Jackson ImmunoResearch; 1:500), Alexa Fluor 647 donkey

anti-rat (Jackson ImmunoResearch; 1:500), Alexa Fluor 647 donkey anti-rabbit (Jackson

ImmunoResearch; 1:500), Alexa Fluor 647 donkey anti-goat (Jackson ImmunoResearch; 1:500),

Alexa Fluor 647 donkey anti-goat (Thermo Fisher Scientific; 1:1,000), Cy3 donkey anti-rat

(Jackson ImmunoResearch; 1:500), Cy3 donkey anti-rabbit (Jackson ImmunoResearch; 1:500),

Cy3 donkey anti-goat (Jackson ImmunoResearch; 1:500), and Cy3 donkey anti-guinea pig (Jackson

ImmunoResearch; 1:500).

In situ hybridisation

The cDNA fragments of mouse *Drd1*, *Drd2*, *Chrm1*, *Chrm2*, *Chrm3*, *Chrm4*, and *Chrm5*

were amplified by PCR using antisense primers containing the T7 promoter sequence. In vitro

transcription was performed with PCR-amplified templates using T7 RNA polymerase (Hoffman

La Roche AG, Basel, Switzerland) to synthesise antisense RNA probes (Supplementary Table 29).

Two-color in situ hybridisation was performed as previously described[59]. Briefly, the mice were

deeply anaesthetised with isoflurane and transcardially perfused with ice-cold 10% sucrose in

Milli-Q water, followed by ice-cold 4% PFA. The collected brain samples were post-fixed with 4%

PFA at 4 °C overnight, followed by displacement with 30% sucrose in PBS containing 0.1%

diethylpyrocarbonate at 4 °C overnight. Coronal brain sections (40 μm) were prepared using a

cryostat (Leica Biosystems, Nussloch, Germany). Brain sections were treated with proteinase K

(Roche), acetylated, and then incubated with hybridisation buffer containing antisense RNA probes

at 60 °C for 16 h. After stringent washing, the brain sections were incubated with peroxidase

(POD)-conjugated anti-FITC antibody (Roche Applied Science, Germany; 1:2,000) or POD-

conjugated anti-Digoxigenin antibody (Roche Applied Science, Germany; 1:500) at 4 °C overnight.

To sequentially use a POD-conjugated antibody, a 2% sodium azide solution was used to inactivate

HRP. TSA systems (TSA-FITC and TSA-Biotin) and NeutrAvidin Protein, DyLight™ 550

(Thermo Fisher Scientific; 1:500) conjugate (Thermo Fisher Scientific) were used to visualise

mRNA signals.

 Analysis of the brain slice images

Imaging of brain sections was conducted using a LSM800, NanoZoomer XR (Hamamatsu

Photonics, Hamamatsu City, Japan) and Axio Zoom V16 on-axis zoom microscope (Carl Zeiss AG,

Oberkochen, Germany). To confirm the AAV-infected sites in the fibre photometry (Fig. 1, 2, 5, 6,

Extended Data Fig.1, 2, 7), RNAi silencing (Fig. 4i-k, Extended Data Fig. 5), and optogenetics

experiments (Fig. 5, 6 and Extended Data Fig. 6-8), coronal brain sections were immunostained for

substance P (SP)[37]. The recording site was identified based on fibre tracking of the brain

histological section sample. Each subregion of the NAc was defined as follows. While the cNAc

shows a low level of SP expression, the msNAc and vsNAc are characterised by high levels of SP

expression. Moreover, the msNAc is located on the medial side of the cNAc, and the vsNAc is

sandwiched between the cNAc and the dense SP axons (ventral pallidum) (Fig. 1c).

Quantification of mAChR1-5, Drd1, *and* Drd2 *on* ChAT^{vsNAc} *and* DA^{VTA} *neurons in the brain slice*

images

To confirm the expression patterns of ChAT and dopamine receptors (*Drd1* and *Drd2*) in

the NAc (Fig. 3g), ChAT and *mAChR1-5* (*mAChR1*, *mAChR2*, *mAChR3*, *mAChR4*, *mAChR5*) in the

vsNAc (Extended Data Fig. 3c), and TH and *mAChR1-5* in the VTA (Extended Data Fig. 3d), we

manually annotated the target marker^{+} somata using a multi-point function and extracted the NAc

subregions by polygon selection in ImageJ/Fiji. The number of signals in each subregion was

counted using schemes implemented in Python 3.7.2. Expression ratios were calculated based on

the number of regions of each target marker. The *Drd1* or *Drd2* ratio in the ChAT^{vsNAc} neurons was

calculated as the target marker (*Drd1* or *Drd2*)$^{+}$ChAT^{+} / ChAT^{+} (%) (Fig. 3g). To calculate the

mAChR1-5 ratio in the ChAT^{vsNAc} neurons, the target marker (*mAChR1-5*)$^{+}$ChAT^{+} / ChAT^{+} (%)

was calculated (Extended Data Fig. 3c). To calculate the *mAChR1-5* ratio in DA^{VTA} neurons, the

target marker $(mAChR1\text{-}5)^+TH^+ / TH^+$ (%) was calculated (Extended Data Fig. 3d). The mean

values for each mouse are shown (Fig. 3g, Extended Data Fig. 3c, d).

The shRNA-mediated knockdown in the vsNAc

AAV[shRNA]-CMV>mCherry-U6>Scramble_shRNA,   AAV[shRNA]-mCherry-

U6>mChat[shRNA#1],   AAV[shRNA]-mCherry-U6>*Drd1*[shRNA#1],   AAV-H1-shScramble-4-

CMV-mCHERRY, AAV-H1-shD2R-4-CMV-mCHERRY) at 100 nL/min with a Hamilton syringe

using a micropump. To calculate the knockdown ratio of target marker by shRNA, we prepared

three types of images in which shRNA (mCherry), the target marker (GFP: *Drd1, Drd2*, or ChAT),

and DAPI were observed around the vsNAc. The target$^+$ (mCheery$^+$, GFP$^+$, or DAPI$^+$) somata were

extracted from each image using OpenCV 4.5.5. First, cv2.medialBlur with a kernel size of 9 was

applied to the image to reduce noise. Next, the histogram of the image was calculated using

cv2.calcHist, and the peak of the histogram was obtained using scipy.signal.find_peaks with a

distance of 5. V_{pk} was set to the pixel value corresponding to the maximum peak. The image was

binarized with a threshold of V_{pk} minus 10, and the contours of the target$^+$ somata were extracted

by applying cv2.findContours to the binarized image. The mode and method were set to

becv2.RETR_EXTERNAL and cv2.CHAIN_APPROX_SIMPLE, respectively. We selected the

contours of DAPI and the target markers with areas between 10 and 200 in the images. The

knockdown ratios were calculated from the region numbers of the target markers, shRNA, and

DAPI. The shRNA knockdown ratio was calculated as the target marker (*Drd1, Drd2* or

ChAT)$^+$shRNA$^+$ / DAPI$^+$shRNA$^+$ (%). The mean values for each mouse are shown in Fig. 4i-k.

Sexual behaviour test

Hormone priming

Stimulus females in oestrus were induced as described previously[8]. C57BL/6J female mice

were ovariectomised at > 8 weeks of age. One week after the surgery, estradiol benzoate dissolved

in sesame oil (10 µg/0.1 mL; EB) was administered to all female mice as a stimulus. More than 1 d

after the first EB injection, all female mice were hormonally primed with subcutaneous injections

of EB 48 and 24 h before testing, and they received progesterone dissolved in sesame oil (500

µg/0.1 mL) 4–6 h before testing to ensure high sexual receptivity. Male mice were exposed to all

female mice in their home cages. Female mice were examined for ejaculation using a copulatory

plug. Female stimuli that experienced ejaculation one time were used in the sexual behaviour test.

Male mice were exposed to a different receptive female mouse from that used in the previous test.

Behavioural analysis

Sexual behaviours, including mounting, intromission, ejaculation, and falling, were

analysed based on videos obtained using cameras. The video frames were manually annotated with

these sexual behaviours using BORIS[60]. The duration and number of mountings, intromissions, and

ejaculations were recorded for each mouse. The latency of each behaviour was calculated by

subtracting the female onset from the onset of each behaviour. For optogenetic manipulation before

intromission, appetitive behaviours, including sniffing and chasing a female mouse, were manually

annotated (Fig. 5b, k, 6b).

Visualisation of the intromission thrust movements

The body regions of two male and female mice during intromission were extracted using

Python 3.7.2 and OpenCV 4.5.5. to visualise their thrusting movements. First, video frames with

intromission behaviours were extracted from the original videos, and the intromission frames were

then converted into grayscale images. Next, the contrast-limited adaptive histogram equalisation

(CLAHE) operator was applied to the grayscale images to improve their contrast (clipLimit = 2.0,

tileGridSize = (3, 3)). The histogram of each CLAHE image was calculated using cv2.calcHist. The

histogram had several pixel values with a height of 500 pixels. With V_{500} representing the

minimum pixel value, the binarization operator was applied to the CLAHE image using threshold

V_{500}. Conversely, the createBackgroundSubstractorMOG2 (BSMOG2) was applied to the grayscale

images, and the median smoothing filter with a kernel size of 25 was applied to the BSMOG2

images. The binarized CLAHE images were multiplied by the smoothed BSMOG2 images, and

black (zero) pixels were extracted from the multiplied images as mouse body regions (Fig. 2g). The

Z-scored lines of mouse body region sizes and the bottom point as thrusting behaviour were plotted

using MATLAB (Fig. 2h).

Fibre photometry experiment

i)   *Fibre photometry recording*

For fibre photometry imaging of neurotransmitters, AAV-hSyn-GRAB$_{DA2m}$, AAV-hSyn-

GRAB$_{ACh3.0}$, or AAV-hSyn-GRAB$_{rDA2m}$ was injected unilaterally into the NAc of C57BL/6J mice.

For fibre photometry imaging of Ca^{2+} dynamics, AAV-hSyn-DIO-GCaMP6s was injected

unilaterally into the VTA of *DAT^{Cre}* mice or the vsNAc of *ChAT^{Cre}* mice, *Drd1^{Cre}* mice, or *Drd2^{Cre}*

mice. More than 1 week after virus injection, a 200-µm core, 0.50 NA optical fibre (Thorlabs and

Lymyth) was implanted unilaterally into the vsNAc or VTA. The mice were allowed to recover for

at least 2 weeks before the experiments.

To monitor animal behaviour during fibre photometry imaging, (method 1) a web camera

(Logicool HD Webcam C525r, Logitech International S.A., Switzerland) was used. To synchronise

the fibre photometry recording and the video recording that captures mouse behaviours, (method 2)

a USB camera (FLIR USB3.0 Vision Camera # CM3-U3-13Y3M-CS; Teledyne FLIR, Wilsonville,

OR) was used. The USB camera released the shutter at a rate of 10 fps based on the Fibre

Photometry Console signals. On the day of the experiment, mice were recorded in their home cages

with regular chow feeding for 2 h in the dark phase. More than 10 min after the onset of recording, an oestrous female mouse was introduced into the home cages of male mice. The recording was stopped more than 2 h after introducing the female. After the test, we reviewed the video footage and examined the female mouse's vagina to identify any signs of ejaculation by the male mouse.

For DA, ACh, or GCaMP imaging, the fluorescence emitted by $GRAB_{DA2m}$, $GRAB_{ACh3.0}$, or GCaMP was collected under a blue LED (CLED465), and the fluorescence emitted by $GRAB_{rDA2m}$ was collected under a green LED (CLED560). The $GRAB_{DA2m}$, $GRAB_{ACh3.0}$, and GCaMP signals were recorded at an acquisition rate of > 10 Hz, and those of $GRAB_{ACh3.0}$ and $GRAB_{rDA2m}$ were recorded at an acquisition rate of 20 Hz using Doric Neuroscience Studio (version 5. 4. 1. 5, Doric Lenses Inc.). The LED output intensity was adjusted so that the light excitation of the optical fibre cable was approximately 30 $\mu W/mm^2$. The signals of $GRAB_{DA2m}$, $GRAB_{ACh3.0}$, and GCaMP between 500–540 nm and those of $GRAB_{rDA2m}$ between 580–680 nm were recorded.

ii) Data analysis for fibre photometry signals

All fibre photometry signals, except for the two-colour imaging of GCaMP6s, $GRAB_{ACh3.0}$, and $GRAB_{rDA2m}$, were transformed into 10 Hz by a MATLAB interp1 function to align with the behavioural events. As peri-event time plots (PETP), To analyse the signal dynamics around the onset and offset of each sexual behaviour, we extracted the signal intensities for 20 s before and after the behavioural events (for a total of 40 s). The ΔF/F ratio was calculated by dividing the signal intensity in each frame by the mean value for 20 s before the behavioural events. The ΔF/F ratio was z-scored against the extracted 40-s signal intensities. To compare the intensity changes of each signal, mean ΔF/F every 10 s in each behavioural event, in which the baseline was the mean for the first 10 s were plotted (Fig. 1e-g, Extended Data Fig. 1g, h, j, 2e, h, 7a, b).

To analyse the DA rhythm (Fig. 1h-j, 2l, Extended Data Fig. 2m) or dual ACh-DA rhythms (Fig. 2e, j, k) during intromission, we calculated the power spectrum using MATLAB FFT with a 3 s window allowing a 0.3 s overlap or 0.25 s overlap, respectively. To show that ACh rhythm, but

not DA rhythm, started before intromission, we analysed the sum of the power spectrum of the 1.5–2.2 Hz band of ACh/DA rhythms every 2 s observed from 10 s before to 10 s after the intromission onset (Fig. 2f). To analyse the DA rhythm during intromission, we collected the sum of the power spectrum every 0.2 Hz band during intromission behaviour (Fig. 2l, Extended Data Fig. 2m).

To analyse the coherence between $GRAB_{ACh3.0}$ and $GRAB_{rDA2m}$, first we prepared the extracted signals around intromission. The $GRAB_{ACh3.0}$ and $GRAB_{rDA2m}$ signals were extracted 20 s before the intromission onset to 10 s before the intromission onset and 10 s before the intromission offset to 10 s after the intromission offset. These two signals were combined, and 50 s length signals were prepared. Using these 50 s length signals, MATLAB wavelet coherence was applied. Average values of the wavelet coherence over all the target events were calculated and it was visualised by heatmap plot (Fig. 2c). To compare the coherence between $GRAB_{ACh3.0}$ and $GRAB_{rDA2m}$ around the intromission, we calculated the mean of the wavelet coherence around <1 Hz, 1–4 Hz, and >4 Hz every 10 s (Fig. 2d, Extended Data Fig. 2j) and the mean of the wavelet coherence in IO and IPE around 1-4 Hz every 5 s (Extended Data Fig. 2k, l).

To visualise ACh and DA dynamics in the vsNAc during each thrust in the intromission phase, we extracted the time point that showed the bottom of the area of the mouse body, which was assumed to be the pelvic thrusting behaviour. Based on the bottom time point, we extracted $GRAB_{ACh3.0}$ and $GRAB_{rDA2m}$ around the thrusting behaviour ± 0.5 s (Fig. 2g, h).

To analyse the $GRAB_{DA2m}$ dynamics with optogenetic activation of $ChAT^{vsNAc}$ neurons using ChrimsonR, we calculated the power spectrum using MATLAB FFT with a 3 s window allowing 0.3 s overlap and collected the power spectrum of the 1.5–2.2 Hz band observed from 10 s before the ejaculation onset to the ejaculation onset during intromission (Fig. 6l).

Ex vivo imaging and optogenetics

We used 8–21-week-old $ChAT^{Cre}$ or $ChAT^{Cre}$; DAT^{Cre} male mice for imaging and

optogenetic experiments during acute brain histological section preparation. These mice were

injected with the following AAV vectors into the target brain areas at least 3 weeks prior to the

experiments: (i) for imaging the DA response to photostimulation of cholinergic neurons in the

vsNAc, AAV-Syn-FLEX-ChrimsonR-tdTomato and AAV-hSyn-GRAB$_{DA2m}$ were injected into the

vsNAc of $ChAT^{Cre}$ mice (Fig. 3b, c); (ii) for imaging Ca^{2+} in cholinergic neurons in response to

photostimulation of dopaminergic axons from the VTA, AAV-Syn-FLEX-ChrimsonR-tdTomato

and AAV-hSyn-DIO-GCaMP6s were injected into the VTA and vsNAc of $ChAT^{Cre}$;DAT^{Cre} double

transgenic mice, respectively (Fig. 3d, e); (iii) for imaging the ACh response to photostimulation of

cholinergic neurons in the vsNAc, AAV-Syn-FLEX-ChrimsonR-tdTomato and AAV-hSyn-

GRAB$_{ACh3.0}$ were injected into the vsNAc of $ChAT^{Cre}$ mice (Fig. 3h-k). Acute brain histological

sections were prepared using a slightly modified N-methyl-d-glucamine (NMDG) protective

recovery method[61,62]. Briefly, mice were anaesthetised with isoflurane and then cardiovascularly

perfused with ice-cold NMDG-artificial cerebrospinal fluid (aCSF) consisting of 93 mM NMDG,

93 mM HCl, 2.5 mM KCl, 1.2 mM NaH_2PO_4, 30 mM $NaHCO_3$, 20 mM HEPES, 25 mM D-glucose,

5 mM Na-ascorbate, 2 mM thiourea, 3 mM Na-pyruvate, 10 mM $MgSO_4$, and 0.5 mM $CaCl_2$ (pH

7.4, adjusted with HCl, oxygenated with 95% O_2/5% CO_2). The brains were sliced to a thickness of

200 μm in ice-cold NMDG-aCSF using a vibratome (Leica VT1200S). Slices containing the

striatum were incubated in NMDG-aCSF at 32 °C for 12 min and then in HEPES-aCSF consisting

of 124 mM NaCl, 2.5 mM KCl, 1.2 mM NaH_2PO_4, 24 mM $NaHCO_3$, 5 mM HEPES, 12.5 mM D-

glucose, 2 mM $CaCl_2$, and 1 mM $MgSO_4$ (~310 mOsm, oxygenated with 95% O_2/5% CO_2) at room

temperature for at least 1 h. Slices were transferred into a chamber under an upright fluorescence

microscope (Zeiss AxioExaminer D1, HXP120, 20x or 40x water-immersion objective lens) and

submerged in perfused HEPES-sCSF at 32–35 °C. Fluorescence signals were acquired using an

EM-CCD camera (Andor iXon Ultra 888, 1024×1024 pixels, 16 bit) and iQ software (Andor), in

which green and red fluorescence were observed using a Zeiss filter set 38HE (excitation 470/40

nm, emission 525/50 nm) and Zeiss filter set 43 (excitation 545/25 nm, emission 605/70 nm),

respectively. Green fluorescence included $GRAB_{DA2m}$, $GRAB_{ACh3.0}$, and GCaMP6s. The frame rate

was 5 fps, except for experiments with Ca^{2+} imaging or bath application of SKF38398 (frame rate

of 0.2 fps). For optical stimulation, an optical fibre was placed just above the target area and

outside the field of view, and optical stimuli (579 nm, 10-ms pulses for 30 s, Shanghai Laser) were

controlled using a function generator (Master-9, AMPI). A data acquisition system (Digidata®

1440A Low-noise data acquisition system, Molecular Devices) was employed to synchronise the

timing of optical stimuli and image acquisition. This system allowed for precise monitoring of the

timing of both processes.

Image analysis was performed using Fiji and R. All images were subtracted from the

background signal, which was the mean fluorescence intensity outside the field of view of the

objective. Bleaching was corrected by linear interpolation of the medians of fluorescence intensity

for 5–10 s before and after photostimulation. For optogenetics and imaging with $GRAB_{DA2m}$ or

$GRAB_{ACh3.0}$, the edges of the 1024×1024-pixel images were removed to obtain 1008×1008-pixel

images because the edges were outside the field of view of the objective lens. A total of 196 (16 ×

16) square ROIs with 63 pixels per side were created in a 1008×1008 image, and the mean

fluorescence intensity within each ROI was calculated. ROIs without $GRAB_{DA2m}$ or $GRAB_{Ach3.0}$

expression (mean intensity <100) were excluded from the analysis. The mean fluorescence

intensity was z-scored against the value obtained for 20 s before photostimulation. For experiments

with SKF38398, bleaching was corrected using the median values for 60 s before and 120–180 s

after bath application, and the fluorescence intensity was z-scored against the values for 60 s before

the application. For Ca^{2+} imaging with GCaMP6s, ROIs were set on the somata of the neurons

expressing GCaMP6s.

ROIs were classified using the Mann–Whitney U test, with values for 5 s before and 10 s

after photostimulation for the GCaMP experiments (Fig. 3e), and 60 s before and 120 s after the

application of SKF38398 for the GRAB$_{ACh3.0}$ experiments (Fig. 3k).

Computational modelling of DA and ACh dynamics

As we observed similar dynamics in GRAB and GCaMP (GRAB$_{DA2m}$ and GCaMP6s of

dopaminergic axons in the vsNAc [Fig 1 and Extended Data Fig. 1i-k], GRAB$_{ACh3.0}$ and GCaMP6s

of cholinergic neurons in the vsNAc [Extended Data Fig. 2d-i]) by fibre photometry imaging, we

hypothesised that (1) the neuronal activity of the neuronal ensemble reflects the amount of NT

(neurotransmitter) released; (2) the GCaMP dynamics reflect the release probability of the NT.

Based on these hypotheses, we developed a computational model to estimate the cellular

environment in which neurons and receptors are involved in generating dual ACh-DA rhythms

during intromission. In this model, we set neuronal assembly ($ChAT_{sim}^{vsNAc}$ and $DA_{sim}^{VTA \rightarrow vsNAc}$), the

internal activity of neurons (release probability measurement, RPM), NT release (ACh_{sim} and

DA_{sim}) based on the RPM, receptors expressed on neurons (D1R, D2R, and nAChR), and the

extracellular environment in which the NT can be released. To mimic the change of the amount of

the extracellular NT over time, in this model, we assumed the three steps of cycle: (i) the activation

of target neurons by the external stimuli; (ii) NT released from the target neurons; (iii) the effect of

the receptor by binding NT to the receptor on the target neurons. These cycles were repeated a

certain number of times and, during this period, we recorded the RPM and amount of NT_{sim}

released from the neurons which was determined based on the RPM value. Through this model, we

attempted to estimate the appropriate cellular environment, including the type of receptors or

neurons involved, by minimising $Diff(NT_{sim}, NT_{real})$, the difference between the actual ACh-DA

dynamics in vivo (NT recorded by fibre photometry: NT_{real}) and model-simulated ACh-DA

dynamics (NT generated by the model: NT_{sim}) (Fig. 4a, b, Extended Data Fig. 4).

Hypothetical neurons and receptors used in this simulation were identified based on ex vivo

imaging (Fig. 3), the in situ hybridisation results (Extended Data Fig. 3), and previous reports[45]. As

neurons and receptors, we placed D1R and D2R in $ChAT_{sim}^{vsNAc}$ neurons, as well as D2R and

nAChR in $DA_{sim}^{VTA \to vsNAc}$ axon terminals. From our ex vivo imaging, we found that inhibitory

mAChR may function as an autoreceptor (Fig. 3b, c). However, since all subtypes of *mAChR* were

expressed on ChAT[vsNAc] and DA[VTA] neurons (Extended Data Fig. 3), to simplify our modelling, we

focused on D1R, D2R, and nAChR receptors in this experiment (Fig. 4a).

(i) Activation of target neurons by the external stimuli

To mimic the cellular environment that neurons are activated by the external stimuli, in this

modelling, we assumed that the RPM which reflects the intensity of neuronal activity was changed

by the external stimuli or the effect of the receptor expressed on the neurons. $Activation$ includes

the spontaneous activity of the target neurons and stimulation of neurons by substances other than

ACh and DA. Here, we defined that the $Activation$ continuously adds a certain value to the target

neuron. Thus, the value added as an external stimulus to the RPM of the target neurons every cycle

was constant.

(ii) Release of NT from the target neurons

To mimic the manner of NT release, we defined that the volume of NT_{sim} released to the

extracellular environment from the neurons was determined by the RPM value of the neurons. In

this model, the volume of NT_{sim} released is equivalent to the RPM value if RPM is positive and 0

if RPM is negative (Fig. 4b). Additionally, we defined the diffusion function of DA to mimic the in

vivo extracellular DA diffusion mechanism[63]. The time course and intensity of the DA diffusion

were investigated using this model (Extended Data Fig. 4c-e).

(iii) Effect of the receptor by binding NT to the receptor on the target neurons

813    Next, we considered the strategy to mimic the behaviour of nAChR, D1R, and D2R. As

814 reported in previous studies, DA receptors are GPCRs that are thought to respond more slowly than

815 nAChR, the ionotropic receptor[40–44]. Incorporating these insights, we defined the characteristics of

816 each receptor using two variables: RE (receptor efficacy), the effect intensity which changes RPM;

817 RS (receptor speed), the latency where the $Receptor$ function affects RPM from the time that

818 NT_{sim} binds to the receptor. In this experiment, to simplify the setting, we set the nAChR related

819 variables as constant values (nAChR's RE=1, and the RS=1), and estimated the characteristics of

820 D1R and D2R by exploring their RE and RS. RE of D1R and D2R was searched over a range of

821 positive values, and their RS was searched over a range of natural numbers.

822    For example, when ACh_{sim} binds to the nAChR expressed on the $DA_{sim}^{VTA \to vsNAc}$ axon

823 terminals, there is an increase in the RPM of the $DA_{sim}^{VTA \to vsNAc}$ axon terminal $ACh_{sim} \times 1$

824 (nAChR's RE=1) one step later (nAChR's RS=1). Since the D1R receptor is an excitatory GPCR,

825 we defined its operation as follows: when D1R receives DA_{sim}, "D1R's RS" step later, the RPM is

826 increased to $DA_{sim} \times$ "D1R's RE." By contrast, we defined operation of the D2R receptor, an

827 inhibitory GPCR, receiving DA_{sim} as "D2R's RS" step later, RPM is decreased by a factor of

828 $DA_{sim} \times$ "D2R's RE". With these receptors' settings, we conducted computational modelling to

829 estimate the RE and RS values.

830    To estimate receptor efficacy (RE) and receptor speed (RS) of DA receptors, we explored

831 RE and RS values which minimise $Diff(NT_{sim}, NT_{real})$, the difference between NT_{sim} and NT_{real}

832 using Bayesian optimisation (Fig. 4b). We used Optuna[64], an automatic hyperparameter

833 optimisation software framework, to determine the best values for RE and RS using

834 $Diff(NT_{sim}, NT_{real})$ as an indicator. We used the following values as a fixed value throughout

835 exploration. The value added by $Activation$ is 1. The number of trials conducted to estimate RE

836 and RS of D1R and D2R was approximately 1,000 to 10,000, depending on the experimental

837 design. In each experimental design, out of these trials, the one representing the smallest

838 $Diff(NT_{sim}, NT_{real})$, the condition where NT_{sim} and NT_{real} are most similar, was used to define

839 RE and RS of D1R and D2R. To express the effect speed of each receptor in seconds, the RS of

840 each receptor and DA diffusion step were divided by 200, the sampling rate (Fig. 4e, Extended

841 Data Fig. 4e).

Experiment 1: Performance comparison by the different activation patterns

844   To prove which neurons should be activated in the generation of dual ACh-DA rhythms, we

845 compared the $Diff(NT_{sim}, NT_{real})$ under the following three conditions: (1) $ChAT_{sim}^{vsNAc}$: on /

846 $DA_{sim}^{VTA \rightarrow vsNAc}$: on (= $Activation_{(1,1)}$); (2) $ChAT_{sim}^{vsNAc}$: on / $DA_{sim}^{VTA \rightarrow vsNAc}$: off (= $Activation_{(1,0)}$);

847 (3) $ChAT_{sim}^{vsNAc}$: off / $DA_{sim}^{VTA \rightarrow vsNAc}$: on (= $Activation_{(0,1)}$) (Fig. 4d, Extended Data Fig. 4a, b).

Experiment 2: Estimation of RE and RS of dopamine receptors

850   To reveal the RE and RS values of D1R and D2R generating dual ACh-DA rhythms, we

851 estimated the RE and RS values under the $Activation_{(1,1)}$ with full receptors condition (D1R and

852 D2R on $ChAT_{sim}^{vsNAc}$, and D2R and nAChR on $DA_{sim}^{VTA \rightarrow vsNAc}$). We focused on trials showing the

853 converged $Diff(NT_{sim}, NT_{real})$ because otherwise the trial indicated the failure of the variable

854 search (Fig. 4e, f).

Experiment 3: Performance comparison by deletion of each candidate receptor

To find which candidate receptor is important for the generation of dual ACh-DA rhythms,

we estimated the RE and RS values under five conditions which delete a target receptor; deletion of

(1) D1R on $ChAT_{sim}^{vsNAc}$, (2) D2R on $ChAT_{sim}^{vsNAc}$, (3) D2R on $DA_{sim}^{VTA \rightarrow vsNAc}$, (4) D2R on both

$ChAT_{sim}^{vsNAc}$ and $DA_{sim}^{VTA \rightarrow vsNAc}$, and (5) nAChR on $DA_{sim}^{VTA \rightarrow vsNAc}$. Then, we compared the

$Diff(NT_{sim}, NT_{real})$ among these five conditions and the condition with full receptor using the

estimated each candidate RE and RS of D1R and D2R (Fig. 4g, Extended Data Fig. 4f, g).

Gene knockdown experiments

Sexual behaviour test with shRNA treatment

To silence the gene expression of ChAT, D1R, or D2R, AAV-CMV-mCherry-U6-shmChAT, AAV-CMV-mCherry-U6-shD1R, or AAV-H1-shD2R-CMV-mCherry was injected into the vsNAc of C57BL/6J male mice. In the control group, AAV-H1-shScramble-CMV-mCherry or AAV-CMV-mCherry-U6-shScramble was injected into the vsNAc of C57BL/6J male mice. Three weeks (± 1 day) following AAV injection, male mice underwent the sexual behaviour test once a week (± 1 day) for 3 weeks. Male mice were relocated to the test room 30 min before starting the test. After the acclimation period, a female stimulus mouse was introduced into the home cage of a male mouse. Video recordings were conducted over the next 2 h. After the test, we reviewed the video footage and inspected the female mice to determine whether male mice exhibited sexual behaviours. The intromission and ejaculation rates were then calculated based on the frequency of these behaviours observed across the three tests (Fig. 4h, l, m). The average latency, the number of bouts, and total duration of each behaviour were calculated across the tests (Extended Data Fig. 5).

Rotarod test with shRNA treatment

A rotarod test was conducted to confirm the motor coordination of mice expressing shRNA in the vsNAc. To silence the gene expression of ChAT, D1R, or D2R, AAV-CMV-mCherry-U6-shmChAT, AAV-CMV-mCherry-U6-shD1R, or AAV-H1-shD2R-CMV-mCherry was injected into the vsNAc of C57BL/6J male mice. For the control group, AAV-H1-shScramble-CMV-mCherry or AAV-CMV-mCherry-U6-shScramble was injected into the vsNAc of C57BL/6J male mice. More than 3 weeks after the AAV injection, male mice were moved to the test room. After more than 15 min, we initiated a habituation process for the rotarod machine. During this phase, mice were placed in the stationary state of the rotarod machine for 2 min. After acclimatisation, the male mice

were allowed to practice on a rotarod for 2 min (12 rpm). Subsequently, we conducted three rotarod

tests (each gradually increasing from 5 to 40 rpm for 300 s). We recorded the latency period before

the mice were dropped from the rotarod machine. Between each test, the mice were allowed a 5-

min break. The median value was plotted as an effect of motor coordination (Fig. 4n).

Optogenetics experiments

To activate $DA^{VTA \rightarrow vsNAc}$ axons, $ChAT^{vsNAc}$ neurons, $D1R^{vsNAc}$ or $D2R^{vsNAc}$, AAV-Ef1a-

DIO-ChR2-EYFP, or AAV-hSyn-DIO-EGFP was injected bilaterally into the VTA of DAT^{Cre} mice

or the vsNAc of $ChAT^{Cre}$ mice, $Drd1^{Cre}$ mice, or $Drd2^{Cre}$ mice. To inhibit $DA^{VTA \rightarrow vsNAc}$ axon

terminals or $ChAT^{vsNAc}$ neurons, AAV-Ef1a-DIO-eNpHR3.0-EYFP or AAV-Ef1a-DIO-EGFP was

injected bilaterally into the VTA of DAT^{Cre} mice or the vsNAc of $ChAT^{Cre}$ mice. After more than 2

weeks, 200-μm core and 0.50 NA optical fibres (Thorlabs and Lymyth) were bilaterally implanted

into the vsNAc. The mice were allowed to recover for at least 2 weeks before conducting the

experiments. Before the tests, it was confirmed that the male mice ejaculated twice. This was

determined by verifying the presence of a plug in the vagina of the female mouse.

Two tests were performed: manipulation before (20 Hz, 10 ms) and during intromission (20

Hz, 10 ms) (Fig. 5b-e, k-p, 6b-g, Extended Data Fig. 6, 7c-f, 8). The interval between each test was

at least 4 days. On the test day, the mice were moved to the experimental room 1 h before the

experiments in the dark phase and optical fibres were connected to their heads. The axonal

terminals of dopaminergic and cholinergic neurons in the vsNAc were activated (462 nm, Shanghai

Laser) or inhibited (579 nm, Shanghai Laser). We created a system using an electrical board to

record the stimulation timing and manage the stimulation frequency using the Arduino UNO and

Doric Neuroscience Studio. Video recordings and recordings of the optogenetic stimulation

schedule were initiated simultaneously using Doric Neuroscience Studio. In each test, the

experimenter observed the behaviour of the mice and turned on the switch to shine the laser. The

switch was immediately turned off if the target behaviour was prevented. If the male mice did not

exhibit any sexual behaviour within 10 min of starting the test, the test was terminated and repeated

on a different day. If the subject male mice did not ejaculate within 10 min after the onset of

stimulation, we stopped the stimulation and confirmed the timing of ejaculation until 20 min after

the onset of stimulation (Fig. 6b).

For manipulation before the intromission (20 Hz, 10 ms) test, the experimenter turned on

the laser while observing sniffing, chasing, and mounting. When mounting was not prevented, the

mice showed intromission, the stimulation was stopped, and the timing of ejaculation was

confirmed. In the other case, the target behaviour was prevented within 10 min of starting the

stimulation; accordingly, we stopped the stimulation and confirmed the timing of ejaculation. If the

male mice did not ejaculate within 20 min of starting the stimulation, the test was terminated, and

the ejaculation latency was recorded as 1,200 s.

For manipulation during the intromission tests (20 Hz, 10 ms), the experimenter turned on

the laser when seeing the intromission, specifically when thrusting occurred more than twice. In

cases where ejaculation was not observed within 10 min of starting the stimulation, the stimulation

ceased, and we confirmed the timing of ejaculation within 20 min of starting the stimulation. The

recorded ejaculation latency in such cases was 1,200 s.

Fibre photometry with optogenetics

To monitor DA dynamics when stimulating cholinergic neurons, AAV-Syn-FLEX-

ChrimsonR-tdTomato and AAV-hSyn-GRAB$_{DA2m}$ were injected into the vsNAc of *ChAT^{Cre}* mice.

Subsequently, optical fibres with a 200-μm core and 0.50 NA (Thorlabs and Lymyth) were

bilaterally implanted in the vsNAc (Fig. 6h, i). The mice were allowed to recover for at least 2

weeks before performing the experiments. GRAB$_{DA2m}$ recordings were conducted without

optogenetics to confirm the GRAB$_{DA2m}$ signals. After confirmation, bilateral optogenetic

manipulations without fibre photometry recordings were conducted to confirm optogenetic effects

on behaviour. Target neurons were activated using a laser (579 nm, Shanghai Laser).

After confirming the behavioural expression, DA dynamics were recorded during

optogenetic activation of cholinergic neurons in the vsNAc. In this experiment, unilaterally DA

dynamics were monitored, and target neurons were activated during intromission. We created a

system using an electrical board to record the stimulation timing and manage the stimulation

frequency with the help of the Arduino UNO and Doric Neuroscience Studio. Fibre photometry

recordings, video recordings using a USB camera (FLIR USB3.0 Vision Camera # CM3-U3-

13Y3M-CS), and recordings of the optogenetic stimulation schedule were synchronised using

Doric Neuroscience Studio. In each test, the experimenter observed the behaviour of the mice.

When observing intromission, particularly more than two intromission-thrusting movements, the

experimenter turned on the switch button to shine the laser. The fluorescence emitted by

GRAB$_{DA2m}$ was recorded under a blue LED (CLED465) using a Doric fibre photometry system for

DA imaging. GRAB$_{DA2m}$ was recorded at 10 Hz using Doric Neuroscience Studio (version 5.4.1.5).

The LED output intensity was adjusted such that the amount of light exiting the optical fibre cable

was approximately 30 μW/mm^2. The GRAB$_{DA2m}$ signals within the 500–540 nm range were

recorded (Fig. 6j).

Quantification and statistical analysis

Recorded videos and histological samples were analysed blind to relevant variables,

including administrated solution, genotype, prior housing condition, surgical procedure, and virus

injection. Videos were manually annotated using BORIS. In particular, anogenital investigation

(sniffing and chasing), mounting, repeated pelvic thrust (intromission), and ejaculation were scored

for male sexual behaviours. After annotations, various parameters including latency, number, and

duration were calculated for further analysis.

Data were processed and analysed using MATLAB, R, Python 3.7.2, and GraphPad PRISM

9.4.1 (GraphPad Software, San Diego, CA). All results are expressed as mean ± SEM. Data of non-

paired samples were analysed with two-tailed non-parametric Mann–Whitney U test or Kruskal–

Wallis test. All multiple comparisons were corrected with Dunn's multiple comparisons correction.

Data of paired samples were analysed with two-way ANOVA followed by Šídák's post-hoc

multiple comparison test. The significance threshold was held at $\alpha=0.05$, two-tailed (not significant

[ns], $p > 0.05$; $*p < 0.05$; $**p < 0.01$; $***p < 0.001$; $****p < 0.0001$). Details are available in

Supplemental Supplementary Table 1-25.

Data and code availability

Further information and requests for resources should be directed to and will be fulfilled by

the Lead Contact, Qinghua Liu (liuqinghua@nibs.ac.cn). This study did not generate new unique

reagents. The code and software used in this study will be freely available through GitHub or

Zenodo once the paper is published.

References

1. Everitt, B. J. Sexual motivation: A neural and behavioural analysis of the mechanisms underlying appetitive and copulatory responses of male rats. *Neurosci. Biobehav. Rev.* **14**, 217–232 (1990).

2. Ågmo, A. Male rat sexual behavior. *Brain Res. Protoc.* **1**, 203–209 (1997).

3. Pfaus, J. G., Kippin, T. E. & Centeno, S. Conditioning and Sexual Behavior: A Review. *Horm. Behav.* **40**, 291–321 (2001).

4. Paredes, R. G. & Ågmo, A. Has dopamine a physiological role in the control of sexual behavior?: A critical review of the evidence. *Prog. Neurobiol.* **73**, 179–225 (2004).

5. Hull, E. M. & Dominguez, J. M. Sexual behavior in male rodents. *Horm. Behav.* **52**, 45–55 (2007).

6. Balthazart, J. & Ball, G. F. Topography in the preoptic region: Differential regulation of appetitive and consummatory male sexual behaviors. *Front. Neuroendocrinol.* **28**, 161–178 (2007).

7. Beach, F. A. Sexual attractivity, proceptivity, and receptivity in female mammals. *Horm. Behav.* **7**, 105–138 (1976).

8. Sano, K., Tsuda, M. C., Musatov, S., Sakamoto, T. & Ogawa, S. Differential effects of site-specific knockdown of estrogen receptor α in the medial amygdala, medial pre-optic area, and ventromedial nucleus of the hypothalamus on sexual and aggressive behavior of male mice. *Eur. J. Neurosci.* **37**, 1308–1319 (2013).

9. Wei, Y.-C. *et al.* Medial preoptic area in mice is capable of mediating sexually dimorphic behaviors regardless of gender. *Nat. Commun.* **9**, 279 (2018).

10. Karigo, T. *et al.* Distinct hypothalamic control of same- and opposite-sex mounting behaviour in mice. *Nature* **589**, 258–263 (2021).

11. Bayless, D. W. *et al.* A neural circuit for male sexual behavior and reward. *Cell* **186**, 3862-

3881.e28 (2023).

12. Hull, E. M. *et al.* Opposite influence of medial preoptic D1 and D2 receptors on genital

reflexes: Implications for copulation. *Life Sci.* **51**, 1705–1713 (1992).

13. Hull, E. M., Eaton, R. C., Moses, J. & Lorrain, D. Copulation increases dopamine activity in

the medial preoptic area of male rats. *Life Sci.* **52**, 935–940 (1993).

14. Bitran, D. & Hull, E. M. Pharmacological analysis of male rat sexual behavior. *Neurosci.*

Biobehav. Rev. **11**, 365–389 (1987).

15. Sun, F. *et al.* A Genetically Encoded Fluorescent Sensor Enables Rapid and Specific Detection

of Dopamine in Flies, Fish, and Mice. *Cell* **174**, 481-496.e19 (2018).

16. Sun, F. *et al.* Next-generation GRAB sensors for monitoring dopaminergic activity in vivo. *Nat.*

Methods **17**, 1156–1166 (2020).

17. Dai, B. *et al.* Responses and functions of dopamine in nucleus accumbens core during social

behaviors. *Cell Rep.* **40**, 111246 (2022).

18. Fiorino, D. F. & Phillips, A. G. Facilitation of sexual behavior and enhanced dopamine efflux

in the nucleus accumbens of male rats after D-amphetamine-induced behavioral sensitization. *J.*

Neurosci. Off. J. Soc. Neurosci. **19**, 456–463 (1999).

19. Holstege, G. *et al.* Brain Activation during Human Male Ejaculation. *J. Neurosci.* **23**, 9185–

9193 (2003).

20. Hull, E. M. *et al.* Brain localization of cholinergic influence on male sex behavior in rats:

Agonists. *Pharmacol. Biochem. Behav.* **31**, 169–174 (1988).

21. Winn, P. Cholinergic stimulation of substantia nigra: effects on feeding, drinking and sexual

behaviour in the male rat. *Psychopharmacology (Berl.)* **104**, 208–214 (1991).

22. Qian, T. *et al.* A genetically encoded sensor measures temporal oxytocin release from different

neuronal compartments. *Nat. Biotechnol.* **41**, 944–957 (2023).

23. Hull, E. M., Muschamp, J. W. & Sato, S. Dopamine and serotonin: influences on male sexual

behavior. *Physiol. Behav.* **83**, 291–307 (2004).

24. Hillegaart, V. & Ahlenius, S. Facilitation and inhibition of male rat ejaculatory behaviour by

the respective 5-HT1A and 5-HT1B receptor agonists 8-OH-DPAT and anpirtoline, as

evidenced by use of the corresponding new and selective receptor antagonists NAD-299 and

NAS-181. *Br. J. Pharmacol.* **125**, 1733–1743 (1998).

25. Pleim, E. T., Matochik, J. A., Barfield, R. J. & Auerbach, S. B. Correlation of dopamine release

in the nucleus accumbens with masculine sexual behavior in rats. *Brain Res.* **524**, 160–163

(1990).

26. Pfaus, J. G. *et al.* Sexual behavior enhances central dopamine transmission in the male rat.

Brain Res. **530**, 345–348 (1990).

27. Damsma, G., Pfaus, J. G., Wenkstern, D., Phillips, A. G. & Fibiger, H. C. Sexual behavior

increases dopamine transmission in the nucleus accumbens and striatum of male rats:

Comparison with novelty and locomotion. *Behav. Neurosci.* **106**, 181–191 (1992).

28. Krok, A. C. *et al.* Intrinsic dopamine and acetylcholine dynamics in the striatum of mice.

Nature **621**, 543–549 (2023).

29. Liu, C. *et al.* An action potential initiation mechanism in distal axons for the control of

dopamine release. *Science* **375**, 1378–1385 (2022).

30. Threlfell, S. *et al.* Striatal dopamine release is triggered by synchronized activity in cholinergic

interneurons. *Neuron* **75**, 58–64 (2012).

31. Wang, L. *et al.* Modulation of dopamine release in the striatum by physiologically relevant

levels of nicotine. *Nat. Commun.* **5**, 3925 (2014).

32. Wang, L. *et al.* Temporal components of cholinergic terminal to dopaminergic terminal

transmission in dorsal striatum slices of mice. *J. Physiol.* **592**, 3559–3576 (2014).

33. Beier, K. T. *et al.* Circuit Architecture of VTA Dopamine Neurons Revealed by Systematic
Input-Output Mapping. *Cell* **162**, 622–634 (2015).

34. Al-Hasani, R. *et al.* Distinct Subpopulations of Nucleus Accumbens Dynorphin Neurons Drive
Aversion and Reward. *Neuron* **87**, 1063–1077 (2015).

35. Poulin, J.-F. *et al.* Mapping projections of molecularly defined dopamine neuron subtypes
using intersectional genetic approaches. *Nat. Neurosci.* **21**, 1260–1271 (2018).

36. Chen, R. *et al.* Decoding molecular and cellular heterogeneity of mouse nucleus accumbens.
Nat. Neurosci. **24**, 1757–1771 (2021).

37. Voorn, P., Gerfen, C. R. & Groenewegen, H. J. Compartmental organization of the ventral
striatum of the rat: Immunohistochemical distribution of enkephalin, substance P, dopamine,
and calcium-binding protein. *J. Comp. Neurol.* **289**, 189–201 (1989).

38. Cachope, R. *et al.* Selective Activation of Cholinergic Interneurons Enhances Accumbal Phasic
Dopamine Release: Setting the Tone for Reward Processing. *Cell Rep.* **2**, 33–41 (2012).

39. Jones, I. W., Bolam, J. Paul. & Wonnacott, S. Presynaptic localisation of the nicotinic
acetylcholine receptor β2 subunit immunoreactivity in rat nigrostriatal dopaminergic neurones.
J. Comp. Neurol. **439**, 235–247 (2001).

40. Neve, K. A., Seamans, J. K. & Trantham-Davidson, H. Dopamine receptor signaling. *J. Recept.*
Signal Transduct. Res. **24**, 165–205 (2004).

41. Greif, G. J., Lin, Y. J., Liu, J. C. & Freedman, J. E. Dopamine-modulated potassium channels
on rat striatal neurons: specific activation and cellular expression. *J. Neurosci.* **15**, 4533–4544
(1995).

42. Cruz, H. G. *et al.* Bi-directional effects of GABAB receptor agonists on the mesolimbic
dopamine system. *Nat. Neurosci.* **7**, 153–159 (2004).

43. Galzi, J.-L. *et al.* Mutations in the channel domain of a neuronal nicotinic receptor convert ion
selectivity from cationic to anionic. *Nature* **359**, 500–505 (1992).

44. Fucile, S. Ca2+ permeability of nicotinic acetylcholine receptors. *Cell Calcium* **35**, 1–8 (2004).

45. De Mei, C., Ramos, M., Iitaka, C. & Borrelli, E. Getting specialized: presynaptic and

postsynaptic dopamine D2 receptors. *Curr. Opin. Pharmacol.* **9**, 53–58 (2009).

46. Richfield, E. K., Penney, J. B. & Young, A. B. Anatomical and affinity state comparisons

between dopamine D1 and D2 receptors in the rat central nervous system. *Neuroscience* **30**,

767–777 (1989).

47. Bazzett, T. J. *et al.* Dose dependent D2 effects on genital reflexes after MPOA injections of

quinelorane and apomorphine. *Life Sci.* **48**, 2309–2315 (1991).

48. Lin, Y.-H. *et al.* Accumbal D2R-medium spiny neurons regulate aversive behaviors through

PKA-Rap1 pathway. *Neurochem. Int.* **143**, 104935 (2021).

49. Soares-Cunha, C., Coimbra, B., Sousa, N. & Rodrigues, A. J. Reappraising striatal D1- and D2-

neurons in reward and aversion. *Neurosci. Biobehav. Rev.* **68**, 370–386 (2016).

50. Raveendran, A. V. & Agarwal, A. Premature ejaculation - current concepts in the management:

A narrative review. *Int. J. Reprod. Biomed.* **19**, 5–22 (2021).

51. Bronner, G., Royter, V., Korczyn, A. D. & Giladi, N. Sexual Dysfunction in Parkinson's

Disease. *J. Sex Marital Ther.* **30**, 95–105 (2004).

52. Althof, S. E. *et al.* An Update of the International Society of Sexual Medicine's Guidelines for

the Diagnosis and Treatment of Premature Ejaculation (PE). *Sex. Med.* **2**, 60–90 (2014).

53. Dremencov, E., Mansari, M. E. & Blier, P. Effects of sustained serotonin reuptake inhibition on

the firing of dopamine neurons in the rat ventral tegmental area. *J. Psychiatry Neurosci.* **34**,

223–229 (2009).

54. Rossi, J. *et al.* Melanocortin-4-receptors Expressed by Cholinergic Neurons Regulate Energy

Balance and Glucose Homeostasis. *Cell Metab.* **13**, 195–204 (2011).

55. Bäckman, C. M. *et al.* Characterization of a mouse strain expressing Cre recombinase from the

3' untranslated region of the dopamine transporter locus. *Genes. N. Y. N 2000* **44**, 383–390

(2006).

56. Lazarus, M. *et al.* Arousal effect of caffeine depends on adenosine A2A receptors in the shell of

the nucleus accumbens. *J. Neurosci. Off. J. Soc. Neurosci.* **31**, 10067–10075 (2011).

57. Inoue, S. *et al.* Periodic Remodeling in a Neural Circuit Governs Timing of Female Sexual

Behavior. *Cell* **179**, 1393-1408.e16 (2019).

58. Liu, C. *et al.* Posterior subthalamic nucleus (PSTh) mediates innate fear-associated

hypothermia in mice. *Nat. Commun.* **12**, 2648 (2021).

59. Ishii, K. K. *et al.* A Labeled-Line Neural Circuit for Pheromone-Mediated Sexual Behaviors in

Mice. *Neuron* **95**, 123-137.e8 (2017).

60. Friard, O. & Gamba, M. BORIS: a free, versatile open-source event-logging software for

video/audio coding and live observations. *Methods Ecol. Evol.* **7**, 1325–1330 (2016).

61. Ting, J. T., Daigle, T. L., Chen, Q. & Feng, G. Acute brain slice methods for adult and aging

animals: application of targeted patch clamp analysis and optogenetics. *Methods Mol. Biol.*

Clifton NJ **1183**, 221–242 (2014).

62. Takahashi, T. M. *et al.* Optogenetic induction of hibernation-like state with modified human

Opsin4 in mice. *Cell Rep. Methods* **2**, 100336 (2022).

63. Floresco, S. B., West, A. R., Ash, B., Moore, H. & Grace, A. A. Afferent modulation of

dopamine neuron firing differentially regulates tonic and phasic dopamine transmission. *Nat.*

Neurosci. **6**, 968–973 (2003).

64. Akiba, T., Sano, S., Yanase, T., Ohta, T. & Koyama, M. Optuna: A Next-generation

Hyperparameter Optimization Framework. in *Proceedings of the 25th ACM SIGKDD*

International Conference on Knowledge Discovery & Data Mining 2623–2631 (Association for

Computing Machinery, 2019). doi:10.1145/3292500.3330701.

1124

1125

Acknowledgements

We thank M. Jing for sharing reagents; A. Yoshida, M. Endo, N. Fujii-Sekine, T. Lou, and A. Watanabe for the behavioural analysis; Editage (www.editage.com) for English language editing; K. Tanaka, N. Takata, T. Miyazaki, and H. Yamada for discussing the experiment; A. □gmo, S. Inoue, and A. Hirano for valuable discussions and comments on the manuscript. This study was supported by JSPS KAKENHI (21J2255 to A.M.), JSPS KAKENHI Grant-in-Aid for Scientific Research on Innovative Areas, "Willdynamics" (19H05006 to K.S.); AMED under Grant JP21zf0127005 (to Q.L.); the WPI program from Japan's MEXT.

Author Contributions

A.M., K.S., T.S., and Q.L. conceived the project and designed the experiments. A.M. performed all stereotaxic surgeries and conducted experiments except as noted. A.M. wrote code for photometry and optogenetics data acquisition and analyses. Y.T. performed tissue samplings. A.M. and Y.T. performed histology analyses and behavioural analyses. K.S. performed retrograde tracing. T.K., A.M., and Y.I. performed ex vivo brain slice imaging. N.N., A.M., and J.S. performed computational modelling. Y.C., A.M., and K.S. prepared AAV vectors. A.M. and T.K. performed the statistical analyses. Y.L. provided the GRAB sensors rDA2.0m and ACh3.0. A.M. and Q.L. wrote original manuscript with input from all co-authors and A.M., T.K., N.N., Y.C., M.Y., H.T., K.S., T.S., and Q.L. edited and reviewed the manuscript.

Competing interests

The authors declare that they have no competing interests.

Figure legends

Figure 1. DA dynamics in the vsNAc closely align with serial transitions of male sexual behaviours.

a, Schematic of sexual behaviours in male mice.

b, Schematic of fibre photometry recording of GRAB$_{DA2m}$ signals in the vsNAc of C57BL/6J mice.

c, Representative image showing immunostaining of ChAT and substance P expression in the NAc and surrounding regions (scale bar = 500 μm).

d, Schematics of recording sites of GRAB$_{DA2m}$ imaging in various NAc subregions.

e-g, Top: Peri-event time plot (PETP) of z-scored $\Delta F/F$ of $GRAB_{DA2m}$ fluorescence around each transition of male sexual behaviour in the (**e**) cNAc, (**f**) msNAc, and (**g**) vsNAc. Middle: Heatmap of PETP, with individual events/row; all data from males. Bottom: Mean $\Delta F/F$ of $GRAB_{DA2m}$ of the three recording sites every 10 s in each behavioural event. See Supplementary Table 1.

h, Representative traces of $GRAB_{DA2m}$ dynamics during intromission and spectrum band calculated by FFT analysis in the (top) cNAc, (middle) msNAc, and (bottom) vsNAc.

i, Comparative power spectrum analysis of DA dynamics in the cNAc, msNAc, and vsNAc during intromission.

j, Mean power spectrum of 1–2 Hz frequency of DA dynamics before, during, and after intromission in the cNAc, msNAc, and vsNAc. See Supplementary Table 2.

Mean ± SEM n = 5 (vsNAc), 10 (msNAc), and 11 (cNAc) (**e, f, g**).

See also Extended Data Fig. 1.

1172 Figure 2. Dual ACh-DA rhythms are observed in the vsNAc during intromission.

1173 **a**, Strategy for simultaneous fibre photometry recordings of ACh and DA dynamics in the vsNAc.

1174 **b**, Representative traces of GRAB_ACh3.0 and GRAB_rDA2m before, during, and after intromission.

1175 **c**, Heatmap plot of mean wavelet coherence between GRAB_ACh3.0 and GRAB_rDA2m dynamics

1176 around the extracted 50 s intromission.

d, Comparison of 1–4 Hz wavelet coherence between $GRAB_{Ach3.0}$ and $GRAB_{rDA2m}$ among every 10 s around intromission.

e, Representative Z-scored traces of $GRAB_{ACh3.0}$ and $GRAB_{rDA2m}$ in the vsNAc (top); heatmap showing the power spectrum processed by the FFT analysis of $GRAB_{ACh3.0}$ (centre) and $GRAB_{rDA2m}$ (bottom) around intromission onset.

f, Comparison of the total amounts of 1.5–2.2 Hz power collected every 2 s around intromission onset.

g, Thrusting was determined by the region size change of the extracted male and female mice.

h, $GRAB_{ACh3.0}$ and $GRAB_{rDA2m}$ around one thrusting during one intromission. (Top) The average traces (bottom) heatmap plot.

i, Representative laser illustration of sexual behaviour.

j, k, Same as (**e**), FFT analysis around IO offset (**j**), and IPE offset (**k**).

l, Power spectrum of $GRAB_{DA2m}$ in the vsNAc during 10 s offset of IO and IPE. Same as Fig. 1, n = 5 mice (vsNAc). Two-way ANOVA Šídák's multiple comparisons test over IO offset and IPE offset every 0.2 Hz. See Supplementary Table 6 (**d, f, l**).

Figure 3. Ex vivo NAc brain slice imaging reveals mutual regulation of ACh and DA

signalling.

a, Diagram of ex vivo experiments. Letters in parentheses indicate the corresponding figures.

b, c, Optogenetic stimulation of ChAT[NAc] neurons and imaging of GRAB$_{DA2m}$. Representative

traces of z-scored ΔF/F of GRAB$_{DA2m}$ signals in response to 10 ms-photostimulation at 20 Hz for

30 s with neither DHβE or atropine (left), with only DHβE (centre), and with only atropine (right).

Bottom orange bars indicate the duration of photostimulation (**b**). Cumulative probability plots of

mean z-scored ΔF/F of GRAB$_{DA2m}$ signals for 30 s after the onset of photostimulation. Plot colours

indicate frequencies of the photostimulation. The experimental conditions are the same as those of

the figures directly above. n = 2332 ROIs (left), 776 ROIs (centre), and 570 ROIs (right) at any

photostimulation frequency (**c**).

d, e, Optogenetic stimulation of DA$^{VTA \to NAc}$ axon terminals and Ca^{2+} imaging with GCaMP.

Representative traces of GCaMP signals in response to 10 ms-photostimulation at 20 Hz for 10 s 20

Hz (**d**). Classification of ChAT^{NAc} neurons by the response to photostimulation (**e**, n = 181 cells).

f, g, ChAT$^+$ neurons in the vsNAc are colocalised with *Drd1* and *Drd2* (scale bars = 10μm) (**f**). The

ratio of ChAT$^+$*Drd1*$^+$ neurons/ ChAT$^+$ neurons and that of ChAT$^+$*Drd2*$^+$ neurons/ ChAT$^+$ neurons

(**g**). See Supplementary Table 10.

h, i, Optogenetic stimulation of ChAT^{nAc} neurons and imaging of GRAB$_{ACh3.0}$. Same as (**b**), but for

GRAB$_{ACh3.0}$ signals in aCSF with (left) or without bromocriptine (right) (**h**). Same as (**c**), but for

GRAB$_{ACh3.0}$ signals. n = 657 ROIs (1 Hz), 490 ROIs (10 Hz), 325 ROIs (20 Hz), and 654 ROIs (40

Hz) (left). N = 657 ROIs (1 Hz), 165 ROIs (10 Hz), 654 ROIs (20 Hz), and 489 ROIs (40 Hz)

(right) (**i**).

j, k, Bath application of a D1R agonist and imaging of GRAB$_{ACh3.0}$. All traces of z-scored ΔF/F of

GRAB$_{ACh3.0}$ signals in response to bath application of SKF38398 (**j**). Pie chart of the response

pattern of the ROI (**k**, n = 685 ROIs).

1218

1219
1220 **Figure 4. Computational modelling reveals that nAChR and D2R signalling are necessary for**

1221 **the generation of dual ACh-DA rhythms.**

1222 **a**, Schematic representation of the model.

1223 **b**, Setting of receptors distribution in the model.

1224 **c**, Representative traces of NT_{sim} and NT_{real} of ACh and DA.

d, Activation of cholinergic neurons required to generate dual ACh-DA rhythms. $Diff(NT_{sim}, NT_{real})$ for different activation input patterns with four full receptors. See Supplementary Table 11.

e, RS (Receptor speed) of D1R is larger than that of D2R in the case of generation of dual ACh-DA rhythms.

f, RE (Receptor efficacy) of D1R in $ChAT_{sim}^{vsNAc}$ is smaller than those of D2R and D2R in $ChAT_{sim}^{vsNAc}$ is smaller than D2R in $DA_{sim}^{VTA \rightarrow vsNAc}$ in the case of generation of dual ACh-DA rhythms. The RE of nAChR is shown as a green dashed line.

g, $Diff(NT_{sim}, NT_{real})$ in the condition of ablation of each receptor under the $ChAT_{sim}^{vsNAc}$:on / $DA_{sim}^{VTA \rightarrow vsNAc}$:off activation condition. See Supplementary Table 12.

Mean ± SEM data from all 59 events of 3 males.

See also Extended Data Fig. 4.

h, Schematic of gene knockdown in the vsNAc.

i-k, Coronal brain sections from shRNA targeting D1R (**i**), D2R (**j**), and ChAT (**k**) injected mice. White arrows indicate ChAT$^+$ cells (**k**). Scale bars = 50μm.

l, m, The effects of in vivo knockdown of D1R, D2R, or ChAT in the vsNAc on intromission rate (**l**) and ejaculation rate (**m**).

n, The effects of in vivo knockdown of D1R, D2R, or ChAT in the vsNAc on the rotarod test.

Mean ± SEM n = 4 (shRNA-scramble), 6 (shRNA-ChAT) (**k**); n = 7 (shRNA-scramble), 6 (shRNA-ChAT) (**l, m**); n = 7 (shRNA-scramble), 7 (shRNA-ChAT) (**n**); n = 4 (shRNA-scramble), 3 (shRNA-D1R) (**i, l-n**); n = 4 (shRNA-scramble), 5 (shRNA-D2R) (**j**); n = 8 (shRNA-scramble), 9 (shRNA-D2R) (**l, m**); n = 8 (shRNA-scramble), 9 (shRNA-ChAT) (**n**).

*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

See also Extended Data Fig. 5 and Supplementary Table 13-16.

1250
Figure 5. Activation of DA release into vsNAc increases intromission behaviour.

a, Strategy for optogenetic manipulation. Experimental schedule (top) and surgery design (bottom).

b, **d**, Laser illustration of sexual behaviour and optogenetic manipulation before intromission (**b**)

and during intromission (**d**).

c, e, The effect of the optogenetic inhibition (left) and activation (right) of DA release into the

vsNAc before (**c**) and during (**e**) intromission on the intromission total duration.

Mean ± SEM n = 8 (GFP, control for eNpHR3.0), 10 (eNpHR3.0), n = 6 (GFP, control for ChR2),

9 (ChR2) (**c**); n = 8 (GFP, control for eNpHR3.0), 10 (eNpHR3.0), n = 6 (GFP, control for ChR2)

10 (ChR2) (**e**). $*p < 0.05$.

See also Extended Data Fig. 6 and Supplementary Table 17, 18.

f, in situ staining of *Drd1* and *Drd2* around the NAc.

g, Schematic of fibre photometry recording of GCaMP6s signals expressed in the D1R or D2R-

expressing neurons in the vsNAc of *Drd1^Cre* or *Drd2^Cre* mice, respectively.

h, i, Neuronal activity of D1R^vsNAc (**h**) and D2R^vsNAc neurons (**i**) are suppressed during intromission.

See also Extended Data Fig. 7a, b and Supplementary Table 19, 20.

j, Strategy for optogenetic manipulation of D1R^vsNAc and D2R^vsNAc neurons. See Extended Data

Fig.7.

k, l, Laser illustration of sexual behaviour and optogenetic manipulation of D1R^vsNAc and D2R^vsNAc

neurons before (**k**) and during intromission (**l**).

m, n, The effect of the optogenetic activation of D1R^vsNAc (left) and D2R^vsNAc neurons (right)

before intromission on the mounting total duration (**m**) and ejaculation latency (**n**).

o, p, The effect of the optogenetic activation of D1R^vsNAc (left) and D2R^vsNAc neurons (right) during

intromission on the intromission total duration (**o**) and mounting total duration (**p**).

q, Suppressed D1R^vsNAc and D2R^vsNAc during intromission is important for the forwarding the

sexual behaviour.

Mean ± SEM n = 5 (GFP in D1R^vsNAc), 4 (ChR2 in D1R^vsNAc), n = 4 (GFP in D2R^vsNAc), 4 (ChR2

in D2R^vsNAc) (**m, n**); n = 4 (GFP in D1R^vsNAc), 5 (ChR2 in D1R^vsNAc), n = 5 (GFP in D2R^vsNAc), 6

(ChR2 in D2R^vsNAc) (**o, p**). $*p < 0.05$, $**p < 0.01$.

See also Extended Data Fig. 7c-f; Supplementary Table 21, 22.

Figure 6. Optogenetic activation of ACh release into vsNAc during intromission induces

ejaculation by triggering slowdown of Da rhythm.

a, Strategy for optogenetic manipulation. Experimental schedule (top) and surgery design (bottom).

b, **c**, Laser illustration of sexual behaviour and optogenetic manipulation before intromission (**b**)

and during intromission (**c**).

d, **e**, The effect of the optogenetic inhibition (left) and activation (right) of ChAT^{vsNAc} neurons

before intromission on the intromission latency (**d**) and ejaculation latency (**e**).

f, **g**, The effect of the optogenetic inhibition (left) and activation (right) of ChAT^{vsNAc} neurons

during intromission on the ejaculation latency (**f**) and ejaculation likelihood (**g**).

Mean ± SEM n = 6 (GFP, control for eNpHR3.0), 5 (eNpHR3.0), n = 8 (GFP, control for ChR2), 6

(ChR2) (**d, e**); n = 7 (GFP, control for eNpHR3.0), 5 (eNpHR3.0), n = 9 (GFP, control for ChR2), 7

(ChR2)) (**f, g**). $*p < 0.05$, $**p < 0.01$.

h-l, Activation of ChAT^{vsNAc} neurons induces a slowdown of the DA rhythm.

m, ACh rhythm appears before intromission, while the optogenetic inhibition of ChAT^{vsNAc}

neurons before intromission may prevent these ACh rhythms and the intromission onset. The

optogenetic activation of ChAT^{vsNAc} neurons during intromission may mimic the increased

coherence of ACh and DA and that ACh precedes DA more in IPE than in IO, which may trigger a

slowdown of DA rhythm and ejaculation.

See also Extended Data Fig. 8, Supplementary Table 23-25.

Extended Data Figures

 Extended Data Fig. 1. Rhythmic neuronal activity of dopaminergic neurons is observed only

 in axon terminals in vsNAc (related to Figure 1).

 a-c, The anterior-lateral VTA projects to the vsNAc, while the posterior-medial VTA projects to

 the msNAc (scale bars = 500 μm).

d-h, The neuronal activity of dopaminergic neurons in VTA subregions correspond to the

projection sites in NAc but did not fluctuate during intromission. See Supplementary Table 3.

d, Recording sites of GCaMP6s imaging in the VTA of DAT^{Cre} mice: alVTA and pmVTA. €

Strategy for imaging of GCaMP6s around the VTA subregions in DAT^{Cre} mice.

e, Representative traces for ΔF/F of GCaMP6s around the VTA subregions during intromission

with laser illustration of sexual behaviour.

f, PETP of GCaMP6s fluorescence at each recording site around each sexual behaviour.

g, h, The neuronal activity of dopaminergic axon terminals in vsNAc shows rhythmic fluctuation

during intromission.

i, Strategy for imaging of GCaMP6s of the axon terminals of VTA in vsNAc of DAT^{Cre} mice.

j, PETP of GCaMP6s fluorescence of dopaminergic axon terminals in vsNAc around each sexual

behaviour. See Supplementary Table 4.

k, Representative traces for ΔF/F of GCaMP6s excited by 465 nm (Ca^{2+} dependent) and 405 nm

(Ca^{2+} independent) LED around the VTA subregions during intromission with laser illustration of

sexual behaviour.

Mean ± SEM n = 5 (alVTA), 4 (pmVTA) (**d-h**); n = 5 (VTA→vsNAc) (**i-k**).

1322

1323

Extended Data Fig. *2*. ACh dynamics and neuronal activity of cholinergic interneurons are

rhythmic during intromission (related to Figure 2).

a-c, Dopaminergic axons project to the vsNAc and cholinergic interneurons in the vsNAc (scale bar

= 500 μm). Strategy for labelling of dopaminergic neurons using AAV (**a**). Immunostained coronal

brain sections for GFP, ChAT, substance P, and DAPI around the VTA (**b**) and NAc (**c**).

d-f, The neuronal activity of cholinergic neurons in vsNAc is rhythmic during intromission.

g-i, ACh release into vsNAc is rhythmic during intromission.

(**d, g**) Strategy for imaging of GCaMP6s in cholinergic neurons in the vsNAc in $ChAT^{Cre}$ mice (**d**)

and $GRAB_{ACh3.0}$ in the vsNAc in C57BL/6J mice (**g**). Coronal brain sections for GCaMP6s (**d**) and

$GRAB_{ACh3.0}$ (**g**) immunostained for ChAT (scale bars = 500 μm).

(**e, h**) PETP of GCaMP6s (**e**) and $GRAB_{ACh3.0}$ (**h**) fluorescence at each recording site around each

sexual behaviour. See also Supplementary Table 5.

(**f, i**) Representative traces for $\Delta F/F$ of GCaMP6s (**f**) and $GRAB_{ACh3.0}$ (**i**) during intromission with

laser illustration of sexual behaviour.

Mean ± SEM n = 5 (**d-f**); n = 4 (**g-i**).

j, Mean wavelet coherence between $GRAB_{ACh3.0}$ and $GRAB_{rDA2m}$ every 10 s around intromission of

<1Hz IO (left) and >4Hz IO (right).

k, Heatmap plot of mean wavelet coherence between $GRAB_{ACh3.0}$ and $GRAB_{rDA2m}$ around IO (top)

and IPE (bottom).

l, Comparison of 1–4 Hz wavelet coherence between $GRAB_{ACh3.0}$ and $GRAB_{rDA2m}$ every 10 s

around intromission among IO and IPE. See Supplementary Table 7.

m, Power spectrum of $GRAB_{DA2m}$ in the cNAc (top) and msNAc (bottom) during 10 s offset of IO

and IPE. n = 10 mice (cNAc), n = 9 mice (msNA). Two-way ANOVA Šídák's multiple

comparisons test over IO offset and IPE offset every 0.2 Hz. *ns* in all frequency in the cNAc and

msNAc. See Supplementary Table 8.

1349

Extended Data Fig. 3. *mAChR1-5* on cholinergic neurons in the vsNAc and on dopaminergic

neurons in the VTA (related to Figure 3).

a, Coronal brain sections around the NAc stained with anti-ChAT and in situ hybridisation of

mAChR subtype1–5 (*mAChR1, mAChR2, mAChR3, mAChR4,* and *mAChR5* represent *Chrm1*,

Chrm2, *Chrm3*, *Chrm4*, and *Chrm5*, respectively). Scale bars = 200 µm (left). Enlarged images of

white squares surrounding areas in left images (scale bars = 10 µm) (right).

b, Coronal brain sections around the VTA stained with anti-TH and in situ hybridisation of *mAChR*

subtypes 1–5 (scale bars = 200 µm). White arrows indicate the double positive cells such as

mAChR1$^+$ChAT$^+$ cells (**a**, **b**).

c, The ratio of ChAT$^+$ cells expressing each *mAChR* subtype to the total number of ChAT$^+$ cells in

the vsNAc (%).

1361 **d**, The ratio of TH$^+$ cells expressing each *mAChR* subtype to the total number of TH$^+$ cells in the

1362 VTA (%).

1363 Mean ± SEM n = 5 in *mAChR1,2,4,5* and n = 4 in *mAChR3* (**a, c**); n = 5 in each *mAChR* (**b, d**). See

1364 also Supplementary Table 9.

1365

1366
Extended Data Fig. 4. Modelling explored activation pattern of each neuron (related to

Figure 4).

a, b, The expected release patterns of ACh and DA depend on the different activation inputs on (**a**)

$ChAT_{sim}^{vsNAc}$ ($Activation_{(1,0)}$), and (**b**) $DA_{sim}^{VTA \to vsNAc}$ ($Activation_{(0,1)}$).

c, Distribution setting of released DA_{sim} toward D1R and D2R on $ChAT_{sim}^{vsNAc}$ neurons and D2R on

$DA_{sim}^{VTA \rightarrow vsNAc}$ axon terminals (D1R on $ChAT_{sim}^{vsNAc}$: D2R on $ChAT_{sim}^{vsNAc}$: D2R on

$DA_{sim}^{VTA \rightarrow vsNAc}$=1:1:2).

d, Relationship between $Diff(NT_{sim}, NT_{real})$ and explored DA diffusion rate.

e, Relationship between $Diff(NT_{sim}, NT_{real})$ and the explored DA diffusion step.

f, g, $Diff(NT_{sim}, NT_{real})$ in the condition of ablation of each receptor in the condition of (**f**)

$Activation_{(1,1)}$ and (**g**) $Activation_{(0,1)}$. For (**f**) and (**g**), the Kruskal–Wallis test was used for

statistical analysis, and data are shown as the mean ± SEM. See Supplementary Table 12.

1379

1380

1381

Extended Data Fig. 5. Gene knockdown of ChAT, D1R, and D2R in the vsNAc (related to

Figure 4).

a, The AAV-shRNA is injected into the vsNAc.

b-h, The results of gene knockdown of ChAT (left), D1R (centre) and D2R (right). (**b-d**) The knockdown effects on mounting behaviour, its latency (**b**), its number of bouts (**c**), and its total duration (**d**). (**e-g**) The knockdown effects on intromission behaviour, its latency (**e**), its number of the bout (**f**), and its total duration (**g**). (**h**) The knockdown effect on ejaculation. See also Supplementary Table 15.

Extended Data Fig. 6. Optogenetic manipulation of dopaminergic axons in the vsNAc around intromission (related to Figure 5).

a-h, The effects of optogenetic inhibitory (left, eNpHR3.0) and excitatory (right, ChR2) manipulation of DA$^{VTA \to vsNAc}$ axons before intromission (**a-d**) and during intromission (**e-h**).

a-d, Activation of DA$^{VTA \to vsNAc}$ axons before intromission increases the number of intromission bouts, while inhibition does not affect sexual behaviour. The duration of optogenetic stimulation is not different depending on each group.

e-h, Manipulation of DA$^{VTA \to vsNAc}$ axons during intromission does not affect sexual behaviour. The duration of optogenetic stimulation is not different depending on each group.

i, Optogenetic activation of DA$^{VTA \to vsNAc}$ axons either before or during intromission increases intromission behaviour, while inhibition does not affect the sexual behaviour.

1403   See also Supplementary Table 17, 18.

1404

1405

1406

Extended Data Fig. 7. Optogenetic manipulation of D1R^{vsNAc} and D2R^{vsNAc} around

intromission (related to Figure 5).

a, b, PETP of GCaMP fluorescence at D1R^{vsNAc} (**a**) and D2R^{vsNAc} (**b**) around each sexual behaviour.

c-f, The effects of optogenetic excitatory (ChR2) manipulation of D1R^{vsNAc} (**c, d**) and D2R^{vsNAc} (**e,**

f) before intromission (**c, e**) and during intromission (**d, f**).

Mean ± SEM n = 5 (**a**); n = 5 (**b**). See Supplementary Table 19, 20.

Mean ± SEM n = 5 (GFP in D1R^{vsNAc}), 4 (ChR2 in D1R^{vsNAc}) (**c**); n = 4 (GFP in D1R^{vsNAc}), 5

(ChR2 in D1R^{vsNAc}) (**d**); n = 4 (GFP in D2R^{vsNAc}), 4 (ChR2 in D2R^{vsNAc}) (**e**); n = 5 (GFP in

D2R^{vsNAc}), 6 (ChR2 in D2R^{vsNAc}) (**f**). $^{*}p < 0.05$, $^{**}p < 0.01$.

See Supplementary Table 21, 22.

1417

1418

Extended Data Fig. 8. Optogenetic manipulation of cholinergic neurons in the vsNAc around

intromission (related to Figure 6).

a-k, The effects of optogenetic inhibitory (left, eNpHR3.0) and excitatory (right, ChR2) manipulation of ChAT^{vsNAc} neurons before intromission (**a-f**) and during intromission (**g-k**).

a-f, Inhibition of ChAT^{vsNAc} neurons before intromission prolongs the latency of mounting behaviour (**a**), while inhibition does not affect the sexual behaviour (**a-g**). The duration of optogenetic stimulation is not different depending on each group (**f**).

g-k, Activation of ChAT^{vsNAc} neurons during intromission decreases the number of mounting bouts (**g**), intromission bouts (**i**) and the total duration of mounting (**h**), while inhibition does not affect

sexual behaviour (**g-j**). The duration of optogenetic stimulation is not different depending on each

group (**k**).

l, Optogenetic activation of ChAT^{vsNAc} neurons before intromission does not affect sexual

behaviour, while inhibition increases the latency of mounting, intromission, and ejaculation.

Optogenetic activation of ChAT^{vsNAc} neurons during intromission decreases mounting, the number

of intromission bouts, and ejaculation latency, and increases ejaculation probability which the

optogenetic manipulation induces ejaculation.

See Supplementary Table 23, 24.