An agarose disk electroporation method for *ex vivo* retinal tissue cultured at the air-liquid interface reveals electrical stimulus-induced cell cycle reentry in retinal cells

Megan L. Stone[1], Hannah H. Lee[2] , Edward M. Levine[1,2,*]

[1]Department of Cell and Developmental Biology
Vanderbilt University, Nashville TN 37232

[2]Department of Ophthalmology and Visual Sciences
Vanderbilt University Medical Center, Nashville TN 37232

*Corresponding author
Ed.levine@vumc.org

Funding was generously provided by a Ruth L. Kirschstein Predoctoral Individual National Research Service Award (F31EY035554) (MLS), Janet and Jim Ayers Research Fund in Regenerative Visual Science (EML), the William A. Black Chair in Ophthalmology (EML), and through an unrestricted grant from Research to Prevent Blindness, Inc. to the Department of Ophthalmology and Visual Sciences.

Keywords:

Ethics statement: All procedures and experiments with mice were approved under protocol M1600035 by the Vanderbilt Institutional Animal Care and Use Committee and conformed to the ARVO guidelines for the use of animals in vision research. The research was conducted in concordance with the NIH guidelines for Responsible Conduct in Research (RCR).

The authors declare no conflicts of interest

Motivation

Tissues cultured at the air-liquid interface, such as retinal tissue, are adhered to a filter membrane with media underneath but not fully submerged. If tissues are fully submerged in liquid, they detach from the membrane and become damaged. Electroporation typically requires tissue submergence for electric current flow. We have developed a submergence-free electroporation method using an agarose disk for electric current delivery that can efficiently and consistently electroporate *ex vivo* retinal tissue cultured at an air-liquid interface without the need to submerge the tissue and disrupt the culture system.

Summary

It is advantageous to culture the *ex vivo* murine retina along with many other tissue types at the air-liquid interface. However, gene delivery to these cultures can be challenging. Electroporation is a fast and robust method of gene delivery, but typically requires submergence in a liquid buffer to allow electric current flow. We have developed a submergence-free electroporation technique using that allows for efficient gene delivery to the *ex vivo* murine retina. This method advances our ability to use *ex vivo* retinal tissue for genetic studies and can easily be adapted for any tissue cultured at an air-liquid interface. Use of this method has revealed valuable insights on the state of *ex vivo* retinal tissues and the effects of electrical stimulation on retinal cells.

Introduction

23  The retina is a highly structured organ that relies on intercellular communication

24 to respond to injury and maintain homeostasis[1–3]. The field of retinal research relies on

25 various model systems to study the structure, function, and pathology of the retina. *Ex*

26 *vivo* explant systems involve isolating and culturing tissue samples, which allows for the

27 preservation of the complex structure and physiological characteristics of organs like the

28 retina[4–6]. *Ex vivo* retinal tissue cultures offer several advantages over other models,

29 particularly in the study of retinal function and degeneration. *Ex vivo* models allow the

30 study of cellular changes within the context of the tissue structure compared to *in vitro,*

31 which lack the 3-dimensional context, or organoid models, which lack prior organismal

32 physiological integration and external environmental exposures. Furthermore, *ex vivo*

33 models provide the opportunity for more advanced manipulations and analyses that

34 may be difficult to perform and interpret *in vivo,* such as live imaging, small molecule

35 delivery and cell reprogramming studies[7,8].

36  *Ex vivo* retinal tissues experience cell death and activation of Müller glia over

37 time in culture. This makes *ex vivo* retinal explant models particularly valuable as an

38 injury model[6,9–12]. It is of interest to improve the health of these cultures to study the

39 retina under normal physiological conditions, however. Innovations in culture technique

40 and supplementation has been shown to promote the survival of *ex vivo* retinal

41 explants. One such innovation is to culture tissue on PTFE membrane inserts with

42 media below the membrane filter. This technique maintains an air-liquid interface that is

43 valuable for gas exchange in the tissue[9,13,14]. Optimization studies to preserve the

44 health of *ex vivo* tissues are ongoing, but as is, long term culture of *ex vivo* retinal tissue

45 is a useful injury context to observe live changes in response to genetic manipulation.

Gene delivery to *ex vivo* retinal tissue can be challenging. Viral methods, such as

AAV and lentivirus, are limited by size constraints for DNA cargo and have delayed

expression time compared to non-viral methods[15,16]. Electroporation is a robust method

for plasmid delivery that promotes quick expression. However, electroporation requires

that cells or tissues be submerged in buffer with an electrode to allow electric current

flow. Submersion of tissue is feasible prior to culture, but submerging tissues at the air-

liquid interface causes detachment from the support membrane, which restricts the

ability to transfect after cultures are established.

To overcome the obstacles to delivery of genetic material to *ex vivo* retinal

tissues, we developed an electroporation method that avoids tissue submergence in a

large volume by generating a confined liquid interface between the electrode and

sample (Figure 1). With this adaptation, we characterized the efficiency of

electroporation at 1- and 14-days *ex vivo* (DEV) in the air-liquid interface culture format.

We also characterized the effects of electrical stimulation on cell proliferation in *ex vivo*

retinal cultures. Lastly, we investigate the limitations in assessing transfection efficiency

when relying on a fluorescent reporter construct as a readout.

Results

Agarose disk electroporation of *ex vivo* retinal tissue efficiently delivers genetic
material with minimal tissue submergence
66
The components required for this electroporation system are: (1) an arm

electrode that can be fixed to a ring stand and lowered onto cultured tissues in cell

culture insert, (2) a 0.5% agarose disk, 5mm thick and the width of the arm electrode,

(3) DNA with an electroporation dye containing glycerol to increase viscosity, and (4) an

anode dish underneath the tissue to complete the electrical circuit. Electroporation was

performed at 24 hours (1DEV) or 14 days (14DEV). All *ex vivo* retinal explant tissues

were dissected from mice between 8-12 weeks of age.

To create a complete electric circuit, chilled PBS (4°C) is placed on the anode

surface and the cell culture insert containing the retinal tissue is placed on top (Figure

1A, B). A 0.5% agarose disk roughly 10mm wide is adhered to the arm electrode

surface (Figure 1C). The disk should be adhered with no air bubbles formed between it

and the electrode, which could impede electrical flow. Once the agarose disk is adhered

to the electrode, 10-15uL of DNA-glycerol mixture (1-3 pMol DNA, 2.5% methyl green

dye and 7.5% glycerol) is pipetted on top of the agarose disk facing upward (Figure 1C,

D). The electrode is rotated 180 degrees and the DNA mixture will stay adhered to the

agarose as a suspended drop. The arm electrode is lowered until the DNA mixture

contacts the tissue, at which point current is delivered (Figure 1E). Fluorescent reporter-

expressing plasmids with a general promoter, such as pCMV-eGFP, were used to

assess transfection.

Differences in cell types transfected at the edge and center of the tissue at 1DEV

Since tissues were cultured with their basal surface toward the air-liquid

interface, the cells most accessible to plasmid uptake were astrocytes, retinal ganglion

cells, displaced amacrine cells, and Müller glia[17,18]. Electroporation of pCMV-eGFP at

1DEV revealed a consistent difference in morphology from cells transfected at the

center of the retina compared to the edge that is defined as 100μm in from tissue

periphery around perimeter (Figure 2B). Cells transfected at the edge of the tissue

resembled Müller glia in morphology (Figure 2D, G). Colocalization of eGFP with SOX9,

a marker of Müller glia[19–21], is consistent with this observation (Figure 2D). In contrast,

transfected cells toward the center of the tissue had a retinal ganglion cell-like

morphology, which is distinctly recognizable by long axons converging at the optic nerve

head (Figure 2C, F). Electroporation of pCMV-mCherry into retinas from Thy1:YFP-16

mice (Thy1:YFP), which express YFP in a large cohort of RGCs in addition to other

inner retinal neurons[22], provided further evidence that RGCs were transfected (Figure

2G). Quantifications revealed that the distributions of transfected cells were skewed

toward SOX9+ cells at the tissue edge and toward Thy1:YFP+ cells toward the center

(Figure 2H). Overall, there is equivalent efficiency in transfection of Müller glia and

Thy1:YFP+ cells at 1DEV, but their distribution is location-dependent (Figure 2H).

Müller glia are preferentially transfected at 14DEV

Changes in cell survival, location, size, and morphology occur over time in

culture[10–12]. Retinal tissue was electroporated at 14DEV to assess differences in

transfection profile over time. Cells transfected at the edge of tissue resembled Müller

glial morphology (Figure 3D, G), but the characteristic retinal ganglion cell morphology

visible in the center at 1DEV was not observed (Figure 3C, D, F, G). Staining for SOX9

revealed an increase in transfected Müller glia compared to inner retinal neurons as

assessed with Thy1:YFP regardless of location (Figure 3D, G). These observations

suggest that there is a decrease in neurons transfected at 14DEV overall, and an

increase in transfection of Müller glia at the tissue center (Figure 3H).

To visualize tissue structure with respect to inner neurons and Müller glia at 1

and 14DEV, orthogonal slice projections were constructed from Z-stack confocal

images of Thy1:YFP or Rlbp1:GFP transgenic retinas, the latter serving as a reporter for

Müller glia[23]. An ROI "slice" was cropped from each image in the same orientation to

observe center and edge regions (Figure 4). At 1DEV, YFP+ cells were observed in the

ganglion cell layer and inner nuclear layer (Figure 4A). Also observed was a line of

fluorescence in the inner plexiform layer, suggesting maintained laminar organization.

However, fewer YFP+ cells were observed in the ganglion cell layer at the retinal edge

(Figure 4A, left). At 14DEV, YFP expression in the ganglion cell layer was diminished in

the central region but interestingly was retained in the inner plexiform and nuclear layers

(Figure 4B). In Rlbp1:GFP tissue, GFP expression was also detectable at the ganglion

cell surface in the center of tissue at 1DEV (Figure 4C), consistent with their radial

morphology[24]. At 14DEV, there was an increase in GFP expression throughout the

tissue including at the electroporation surface (Figure 4D), most likely the result of the

extended time in culture. The loss of RGCs and accumulation of Müller glia at the basal

surface of the tissue could explain the shift toward preferential electroporation of Müller

glia at 14DEV.

Electrical stimulation induces BrdU incorporation in *ex vivo* retina

The adult mouse retina is normally quiescent, but proliferation can occur to a

limited extent in non-neuronal cells such as microglia and Müller glia in response to

injury and/or inflammation, two contexts that are likely occurring in the *ex vivo* retina

[6,10,12,25–30]. Proliferation has also been shown to increase in response to electrical

stimulation[31–35], though this has yet to be shown in the retina. We therefore assessed

proliferation in the cultures with and without electroporation to obtain baseline

measurements. Three conditions were tested: not electroporated (*-EP*), electroporated

with DNA loading solution but lacking DNA (*+EP*), or electroporated with pCMV-eGFP

plasmid (*+EP^DNA*). These treatments were applied to *ex vivo* retinas at 1DEV or 14DEV.

At 72 hours post-treatment,16uM BrdU was delivered in the cell culture media for 24

hours (Figure 5A). At both times in culture, there were significant increases in BrdU+

cells per mm^3 in the *+EP* and *+EP^DNA* conditions compared to the *-EP* condition (Figure

5B,C). The similar levels of BrdU incorporation in the *+EP* and *+EP^DNA* conditions were

expected since eGFP was being expressed. These results indicate that electrical pulse

application is sufficient to promote proliferation in *ex vivo* retinal tissue.

The proportion of BrdU+ cells that are Müller glia increases in 14DEV compared

to 1DEV cultures and following electrical stimulus

153   We predicted that microglia and Müller glia were re-entering the cell cycle in

response to electrical stimulation. To assess this, *ex vivo* retinal tissues were

immunostained with the microglial marker IBA1 or with SOX9 in combination with BrdU

detection, and colocalization with BrdU was quantified to determine the number of these

cells undergoing cell cycle entry (Figure 6). At 1 and 14DEV, BrdU+ microglia and

Müller glia were observed in *-EP* and *+EP* conditions (Figure 6A, B). Quantification

revealed a significantly increased proportion of BrdU+ Müller glia, indicated by SOX9

labeling, at 14DEV in *+EP* tissues compared to 1DEV *+EP* (Figure 6C). The number of

BrdU+ Müller glia significantly increased in *+EP* tissues at 14DEV compared to 1DEV

+EP tissues and 14DEV *-EP* tissues, whereas the number of BrdU+ microglia remained

steady (Figure 6D). These results suggest an increase in cell cycle entry in Müller glia

that correlates with time in culture and electrical stimulation.

Transfection efficiency is greater than what is observable through fluorescent

protein expression when delivering a fluorescent reporter construct

A limitation of using fluorescent reporter expression constructs such as pCMV-

eGFP is that cells could have been transfected but expression of the fluorescent protein

was below detection. Since as few as 4 molecules of Cre recombinase are sufficient for

recombination[36–38], we predicted that transfection of a Cre expression construct into a

mouse harboring a recombination reporter allele would reveal a higher transfection

efficiency than what can be detected with pCMV-eGFP (Figure 7A). To test this, pCMV-

Cre and pCMV-eGFP were co-transfected at equimolar concentrations into *ex vivo*

retinal tissues from mice with a Cre-inducible *Rosa^{AI14}* allele. *Ex vivo* tissues were

imaged every 24 hours and fixed at 72 hours post-electroporation.

In cultures transfected at 1DEV (Figure 7B) or 14DEV (Figure 7C), similar trends

were observed at the tissue center and edge. Whereas the majority of eGFP+ cells

were tdTomato (tdTom)+, many tdTom+ cells were not eGFP+ (Venn diagrams in

Figure 7B,C). Quantifications revealed a significant increase in the number of co-labeled

cells per the eGFP population compared to the tdTom population at both timepoints

(Figure 7D). The transfection efficiency and level of expression of these constructs is

expected to be similar based on the promoter elements shared between them.

Additionally, there were significant increases in the percentages of all transfected cells

expressing tdTom compared to eGFP in the 1DEV and 14DEV cultures (Figure 7E).

These data suggest that the transfection efficiency is greater than what can be

observed by transfection of expression plasmids for fluorescent proteins.

Discussion

This study introduces an agarose disk submergence-free method of

electroporation for *ex vivo* retinal tissue cultured at the air-liquid interface. This

electroporation method offers a rapid and time-dependent transfection efficiency while

maintaining tissue integrity. This method is cost effective and does not require

specialized equipment, making it generally accessible. Though we have only tested

retinal tissue, it could easily be adapted for any tissue type that is cultured using an air

liquid interface method, including lung, intestine, or skin[13,14,39,40].

We observed a significant difference in the cell types transfected at different

times in culture. This finding suggests that the timing of the transfection is useful for

targeting specific cell populations within the retina. Müller glia undergo hypertrophy and

migration in response to injury signals from retinal neurons[41–45]. Likely, the increase in

availability of Müller glia at the electroporation surface is due to the loss of retinal

ganglion cells and changes in Müller glial morphology associated with tissue

remodeling[46].

We found that the application of electrical stimulus alone leads to an increase in

proliferation in Müller glia as indicated by BrdU incorporation. The mammalian retina is

quiescent after retinal development, but Müller glia and microglia are known to exhibit

limited proliferation in response to injury or inflammation. This proliferation occurs

typically in young postmitotic mice and diminishes after injury onset[6,10,12,25–27,30,42,45].

Postmitotic Müller glia have been shown to proliferate in culture, however this effect was

observed in *ex vivo* retinal tissues from P10 mice and diminished by p14[10]. The mice

used here were adults between 8-12 weeks of age. This indicates a change in cell

behavior following electrical stimulus that is not typical under normal physiological

conditions.

Proliferation resulting from electrical stimulation has been reported in multiple cell

types, including neural stem cells, human embryonic stem cells, osteoblasts, and

fibroblasts[31,47–50]. Electroporation can also stimulate neural progenitor cells differentiate

into neurons[50–52]. Amongst the BrdU+ cell population, there was a significant increase in

BrdU+ Müller glia at later timepoints and following electrical stimulus whereas the

presence of BrdU+ microglia was not dependent on time in culture. This could indicate

an increase in Müller glia in the tissue at this timepoint or a change in cell state that

leads to increased proliferation of Müller glia. Further studies are needed to determine

whether proliferating Müller glia in this context also acquire neurogenic properties.

Lastly, deliver of a Cre expression construct into a recombination reporter mouse

strain predicts that that true transfection efficiency is greater than observed with a

fluorescent reporter expression construct, meaning more cells may receive gene

delivery than accounted for. It is advantageous with this system to use a sensitive

reporter for electroporation, like Cre-mediated reporter expression, or to boost the signal

for the fluorescent reporter with antibodies to enhance detection.

Limitations of Study

While this study provides valuable insights into the overall efficiency of transfection, especially on some retinal cell types, further research is necessary to fully characterize the effects on every cell type. This method has only been tested on retinal tissue, though it could be adapted for other tissue types cultured in an air-liquid interface format. Proliferation observed in this study was measured using BrdU incorporation, which labeled cells during S-phase of mitosis. BrdU could incorporate in cells that are then arrested in S-phase and do not fully divide. Further testing is required to understand the extent of proliferation following electrical stimulation.

Although certain cell types may be more amenable to electroporation, the method is not inherently cell type specific. Other methods of transfection should be employed if a study requires expression to only occur in certain cell types. We demonstrated that BrdU incorporation is induced following electrical stimulation with this method. If a study is designed to test the effects of an electroporated construct on proliferation, an important control would be the inclusion of an *EP* condition without DNA in addition to the negative control plasmid.

Acknowledgments

E.M.L received funding from M.L.S was supported by Janet and Jim Ayers Research Fund in Regenerative Visual Science (EML), the William A. Black Chair in Ophthalmology, and through an unrestricted grant from Research to Prevent Blindness, Inc. to the Department of Ophthalmology and Visual Sciences. M.L.S was supported by a Ruth L. Kirschstein Predoctoral Individual National Research Service Award through the National Eye Institute (F31EY035554).

Author Contributions

255 Conceptualization, E.M.L and M.L.S; Methodology E.M.L, H.H.L, and M.L.S; Formal
256 Analysis, M.L.S; Investigation, M.L.S, Resources, E.M.L, Data Curation, M.L.S, Writing -
257 Original Draft - M.L.S, Writing - Review & Editing, E.M.L and M.L.S, Visualization,
258 M.L.S, Supervision, E.M.L, Project Administration, E.M.L, Funding Acquisition, E.M.L.
259

Declaration of Interests

The authors declare no competing interests.

Figure Legends

Figure 1. Visual methodology of agarose disk electroporation for *ex vivo* retinal

cultures. (A) Murine retinal tissue cultured at the air-liquid interface on PFTE

membrane after 5 days *ex vivo* (5 DEV) (1X, Brightfield). (B) Agarose disk

electroporation set-up. 5mm high, 10mm diameter 0.5% agarose disk adhered to an

arm electrode above an anode dish containing *ex vivo* retinal tissue on PTFE

membrane with chilled PBS underneath. (C) 12uL of DNA-glycerol mixture is pipetted

onto the agarose disk facing upward. (D) DNA-Glycerol mixture on top of agarose disk

facing upward. (E) 180° rotation electrode arm containing agarose disk and DNA-

glycerol mixture lowered onto *ex vivo* retinal tissue.

Figure 2. Preferential transfection at 1DEV of Müller glia and neurons at the edge

and center of tissue, respectively. (A) Timeline of experimental design for 1DEV

transfection of wild type tissue with pCMV-eGFP reporter plasmin and staining for SOX9

to detect Müller glia. (B) Schematic indicating edge (100μm from the retina periphery)

and center of tissue. (C) pCMV-eGFP transfected cells in wild-type tissue transfected at

1DEV. Scale bar: Left, 200μm. Right, 50μm. (D) Left: pCMV-eGFP transfected cells,

middle: SOX9 staining, and right: overlap of eGFP and SOX9 at the center (top) and

edge (bottom) of tissue at 1DEV. Arrows indicate double-labeled cells. Scale Bar:

50μm.(E) Timeline of experimental design for 1DEV transfection of Thy1:YFP neuronal

reporter tissue with pCMV-mCherry reporter plasmid. (F) pCMV-mCherry transfected

cells in Thy1:YFP tissue transfected at 1DEV (YFP not shown). Scale bar: Left, 200μm,

Right, 50μm. (G) Left: pCMV-mCherry transfected cells, middle: Thy1:YFP reporter, and

right: overlap of mCherry and YFP at the center (top) and edge (bottom) of tissue after

electroporation at 1DEV. Arrows indicate double-labeled cells. Scale Bar: 50μm. (H)

Quantification of percentage of cells transfected with pCMV-eGFP that are SOX9+ and

cells transfected with pCMV-mCherry that are Thy1:YFP+ at the edge, center and

combined edge and center of tissue after electroporation at 1DEV. Mean ± SEM is

shown. N=4-6. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001. No bracket indicates p

> 0.05. All statistical tests performed on arcsine transformed values.

Figure 3. Preferential transfection of Müller glia at 14DEV. (A) Timeline of

experimental design for 14DEV transfection of wild type tissue with pCMV-eGFP

reporter plasmin and staining for SOX9 to detect Müller glia. (B) Schematic indicating

edge (100μm from the retina periphery) and center of tissue. (C) pCMV-eGFP

transfected cells in wild-type tissue transfected at 14DEV. Scale bar: Left, 200μm. Right,

50μm.(D) Left: pCMV-eGFP transfected cells, middle: SOX9 staining, and right: overlap

of eGFP and SOX9 at the center (top) and edge (bottom) of tissue at 14DEV. Arrows

indicate double-labeled cells. Scale Bar: 50μm. (E) Timeline of experimental design for

14DEV transfection of Thy1:YFP neuronal reporter tissue with pCMV-mCherry reporter

plasmid. (F) pCMV-mCherry transfected cells in Thy1:YFP tissue transfected at 14DEV

(YFP not shown). Scale bar: Left 200μm, Right, 50μm. (G) Left: pCMV-mCherry

transfected cells, middle: Thy1:YFP reporter, and right: overlap of mCherry and YFP at

the center (top) and edge (bottom) of tissue after electroporation at 14DEV. Arrows

indicate double-labeled cells. Scale Bar: 50μm. (H) Quantification of percentage of cells

transfected with pCMV-eGFP that are SOX9+ and cells transfected with pCMV-mCherry

that are Thy1:YFP+ at the edge, center and combined edge and center of tissue after

electroporation at 14DEV. Mean ± SEM is shown. N=4-5. *p < 0.05, **p < 0.01, ***p <

0.001, ****p < 0.0001. No bracket indicates p > 0.05. All statistical tests performed on

arcsine transformed values.

Figure 4. Changes in tissue structure at electroporation surface in orthogonal

slice projections of 1DEV & 14DEV Thy:YFP and Rlbp1:GFP tissues favor

electroporation of Müller glia at 14DEV. Orthogonal slice projections of center and

edge created from 3D projections of ROIs (354.25 pixels by 11.76 pixels) to visualize

cross sections within *ex vivo* retinal tissue from transgenic mouse models,.ONL: Outer

nuclear layer. INL: Inner nuclear layer. GCL: Ganglion cell layer. IPL: Inner plexiform

layer. (*) ONH: optic nerve head. (A) Orthogonal slice projections of the edge (left) and

center (right) of Thy1:YFP *ex vivo* tissue fixed at 1DEV. Scale Bar:100μm. (N=4). (B)

Orthogonal slice projections of the edge (left) and center (right) of Thy1:YFP *ex vivo*

tissue fixed at 14DEV. (N=4). (C) Orthogonal slice projections of the edge (left) and

center (right) of Rlbp1:GFP *ex vivo* tissue fixed at 1DEV. (N=4) (D) Orthogonal slice

projections of the edge (left) and center (right) of Rlbp1:GFP *ex vivo* tissue fixed at

14DEV. (N=4).

Figure 5. BrdU incorporation increases with electrical stimulation in the *ex vivo*

retina. (A) Timeline of experimental design for tissue electroporated at 1 and 14DEV.

BrdU is delivered in culture medium for 24 hours. Following fixation, pieces of retinal

tissue were cut, stained for BrdU incorporation, and imaged using confocal microscopy.

No Electroporation (*-EP*) tissues were given BrdU and fixed at the same timepoints

indicated for electroporated tissues (*+EP, +EP^{DNA}*). (B) BrdU incorporation *-EP* (left)*,*

+EP (center) and *+EP^{DNA}* (right) tissues at 1DEV (top) and 14DEV (bottom). Scale Bar:

100µm. (C) Quantification of the number of BrdU+ cells detected per tissue volume

(mm^3). Mean ± SEM is shown. N=5-10. *p < 0.05, **p < 0.01, ***p < 0.001, ****p <

0.0001. No bracket indicates p > 0.05. All statistical tests performed on arcsine

transformed values.

Figure 6. Increase in BrdU+ Müller glia at 14DEV vs. 1DEV and following electrical

stimulus. (A) Left: 1DEV *-EP* tissues stained for IBA1 (top left) or SOX9 (bottom left)

and BrdU (top middle & bottom middle). Overlap of IBA1 & BrdU (top right) or SOX9 &

BrdU (bottom right). Arrows indicate double labeled cells. Scale bar: 50µm. Right: 1DEV

+EP tissues stained for IBA1 (top left) or SOX9 (bottom left) and BrdU (top middle &

bottom middle). Overlap of IBA1 & BrdU (top right) or SOX9 & BrdU (bottom right).

Arrows indicate double labeled cells. Scale bar: 50µm. (B) Left: 14DEV *-EP* tissues

stained for IBA1 (top left) or SOX9 (bottom left) and BrdU (top middle & bottom middle).

Overlap of IBA1 & BrdU (top right) or SOX9 & BrdU (bottom right). Arrows indicate

double labeled cells. Scale bar: 50μm. Right: 14DEV *+EP* tissues stained for IBA1 (top

left) or SOX9 (bottom left) and BrdU (top middle & bottom middle). Overlap of IBA1 &

BrdU (top right) or SOX9 & BrdU (bottom right). Arrows indicate double labeled cells.

Scale bar: 50μm. (C)Left: Quantification of the percent of BrdU+ cells that are IBA1+ in -

EP and *+EP* conditions at 1DEV and 14DEV. Mean ± SEM is shown. No bracket

indicates $p > 0.05$. All statistical tests performed on arcsine transformed values. Right:

Quantification of the percent of BrdU+ cells that are SOX9+ in *-EP* and *+EP* conditions

at 1DEV and 14DEV. Mean ± SEM is shown. N=4-5. *$p < 0.05$. No bracket indicates $p >$

0.05. All statistical tests performed on arcsine transformed values. (D) Left:

Quantification of colocalized SOX9+, BrdU+ cells per tissue volume (mm^3) in *-EP* and

+EP conditions at 1DEV and 14DEV. Mean ± SEM is shown. N=5. No bracket indicates

$p > 0.05$. All statistical tests performed on arcsine transformed values. Right:

Quantification of colocalized SOX9+, BrdU+ cells per tissue volume (mm^3) *-EP* and

+EP conditions at 1DEV and 14DEV. Mean ± SEM is shown. N=5-6. **$p < 0.01$, ***$p <$

0.001. No bracket indicates $p > 0.05$. All statistical tests performed on arcsine

transformed values.

Figure 7. Transfection efficiency is greater than what is observed through

expression of a fluorescent reporter construction. (A) Schematic of detection of

pCMV-eGFP expression vs. pCMV-Cre expression in a Rosa[AI14] reporter tissue when

few or many copies of the construct are transfected in a cell. (B) Left: Cells transfected

with pCMV-CRE (left) or pCMV-eGFP (middle) and the overlap of these channels (right)

at the center (top) or edge (bottom) of tissues electroporated at 1DEV with both

constructs at equal molar concentration. Scale bar: 100µm. Right: Euler plot

demonstrating the number of cells detected as eGFP+. tdTom+ or double labeled at the

center (top) and edge (bottom) of tissue at 1DEV. (C) Left: Cells transfected with pCMV-

CRE (left) or pCMV-eGFP (middle) and the overlap of these channels (right) at the

center (top) or edge (bottom) of tissues electroporated at 14DEV with both constructs at

equal molar concentration. Scale bar: 100µm. Right: Euler plot demonstrating the

number of cells detected as eGFP+. tdTom+ or double labeled at the center (top) and

edge (bottom) of tissue at 14DEV. (D) Left: Quantification of the percent of colocalized

cells detectable as transfected with pCMV-eGFP (eGFP+) and pCMV-Cre (tdTom+)

within eGFP+ or tdTom+ populations at 1DEV. Mean ± SEM is shown. N=4. **p < 0.01.

No bracket indicates p > 0.05. All statistical tests performed on arcsine transformed

values. Right: Quantification of the percent of colocalized cells detectable as transfected

with pCMV-eGFP (eGFP+) and pCMV-Cre (tdTom+) within eGFP+ or tdTom+

populations at 14DEV. Mean ± SEM is shown. N=4. *p < 0.05. No bracket indicates p >

0.05. All statistical tests performed on arcsine transformed values. (E) Left:

Quantification of the percent of cells detectable as transfected with pCMV-eGFP

(eGFP+) and pCMV-Cre (tdTom+) within the population of all transfected cells at 1DEV.

Mean ± SEM is shown. N=4. **p < 0.01. No bracket indicates p > 0.05. All statistical

tests performed on arcsine transformed values. Right: Quantification of the percent of

cells detectable as transfected with pCMV-eGFP (eGFP+) and pCMV-Cre (tdTom+)

within the population of all transfected cells at 14DEV. Mean ± SEM is shown. N=4. *p <

0.05. No bracket indicates p > 0.05. All statistical tests performed on arcsine

transformed values.

STAR Methods

I.   Resource Availability

Lead Contact

Further information and requests for resources and reagents should be directed

to and will be fulfilled by the lead contact: Edward M. Levine, ed.levine@vumc.org

Materials Availability

This study did not generate new unique reagents.

Data and code availability

Microscopy and quantification data reported in this paper will be shared by the

lead contact upon request.

II.   Experimental model and participant details

Mice

B6/129 mice (Jackson Laboratory (#101043)): 8 weeks old, female.

Thy-1 YFP-16 (B6.Cg-Tg(Thy1-YFP)16Jrs/J) (Jackson Laboratory (#003709)): 8 to 12

weeks old, female and male.

Rlbp1-GFP (Tg(Rlbp1-GFP)1Eml (MGI: 6195229)): 8 to 12 weeks old, female and male.

Rlbp1-CreERT2; Rosa^{Ai14} (129S6.Cg-Tg(Rlbp1-cre/ERT2)1Tfur (MGI:Pending); B6.Cg-

Gt(ROSA)26Sor$^{tm14(CAG-tdTomato)Hze}$/J (Jackson Laboratory (#007914))), 8 to 12 weeks

old, female and male.

All mice are co-housed in a 12h/12h light-dark cycle with water and food ad libitum. All

transgenic mice are backcrossed to B/129 mice. All animal experiments with mice were

approved under the protocol M1600235 by the Vanderbilt Institutional Animal Care and

Use committee and conform to the ARVO guidelines for the use of animals in vision

research. All possible efforts were made to minimize animal suffering and the number of

animals used.

III.   Method Details

Tamoxifen administration

To activate Cre recombination in transgenic mice, 2 doses of 200µg per gram

body weight (GBW) of tamoxifen (Sigma Aldrich, T2859) were delivered in corn oil

(Sigma Aldrich, C8267) via oral gavage. Mice were euthanized for tissue collection at

least 5 days after tamoxifen treatment.

Retinal dissection

Prior to tissue harvest, dissection tools were sterilized with 70% ethanol and UV

light for 15 minutes, and workstation was sterilized with 70% ethanol. 1X HBSS with

Ca^{2+} and Mg^{2+} (Gibco, 14025) with 10mM HEPEs (Sigma Aldrich, H0887) and 1X

Antibiotics-Antimycotics (Gibco, 15240062) (referred to as HBSS hereon) was prepared

and kept on ice. Mice were euthanized by CO_2 asphyxiation and cervical dislocation.

Each eye was enucleated, rinsed in cold HBSS, then transferred to a fresh HBSS dish.

Under a dissection scope and on a cold plate to keep tissues chilled, an incision was

made in the cornea. The cornea was removed using spring scissors. The sclera,

choroid and RPE were removed, and the retina and lens were transferred to a fresh

HBSS dish. The lens was removed. Using spring scissors, 4 incisions were cut into

retinal tissue from periphery through 25-40% of tissue toward optic nerve head, creating

a flower-shaped retinal tissue that can be flattened.

***Ex vivo* retinal explant tissue culture**

On the day of culture, plating media with supplements was prepared:

NeuroCult™ Plating Media (Stem Cell Technologies, 05713), 1X SM1 neuronal

supplement (Stem Cell Technologies, 05711) , 500µM L-Glutamine (Gibco, 25030149),

and 100unit Penicillin/100µg Streptomycin (Pen/Strep) (ThermoFisher, 15140) (referred

to as plating media hereon).

Retinal tissue was dissected as described above. A hydrophilic PTFE cell culture

insert (Millicell, PICM0RG50) was prepared by pipetting 150uL of plating media onto

insert membrane. Retinal tissue was mounted onto the PTFE membrane with the retinal

apical surface (photoreceptor side) in juxtaposition. Tissue was placed into a 6-well

plate (Corning, 353004) containing 1mL plating media and incubated at 37°C with 5%

CO^2. After 48 hours, 500uL of initial plating media was replaced with BrainPhys™

Neuronal Medium (Stem Cell Technologies, 05790) containing 1X SM1 neuronal

supplement, 1X N2 Supplement-A (Stem Cell Technologies, 07152) and Pen/Strep

(referred to as BrainPhys media hereon). 500μL of media was removed and replaced

with fresh BrainPhys media every 48 hours for the extent of the culture.

Agarose Disk Electroporation

0.5% agarose (RPI, A20090) in 1X PBS was heated to dissolve. 500μL was

pipetted into multiple wells of a 24-well plate(Corning, CLS3516) and left to solidify at

room temperature for 30 minutes in a cell culture hood. An anode dish was made by

lining a 100mm dish (Sigma Aldrich, Z358762) with aluminum foil, sterilized under UV

light for 15 minutes, and connected to the electroporator (BTX, 45-0052) by clamping

the anode cable onto the aluminum foil. Sterile, chilled PBS was pipetted onto the

anode dish. Tools and workspace were sterilized with 70% ethanol. The PTFE insert

containing the tissue culture was washed by gently dipping the membrane in fresh,

sterile PBS. The PTFE insert was then placed on the PBS in the anode dish.

An arm electrode (Bulldog Bio, CUY700P7L) connected to the electroporator was

mounted on a micromanipulator (Narishige, UMM-3C) with the electrode surface facing

up. Using a plastic transfer pipet (Fisherbrand, 13-711-7M) cut above the tapered tip, an

agarose disk was stamped from the solidified agarose and transferred using a spatula

to the arm electrode face. The disk was gently pressed to ensure adherence without

bubbles underneath.

The 12μL DNA-glycerol transfection solution was prepared using a combination

of 3pMol plasmid DNA diluted in sterile PBS with 7.5% glycerol (Sigma Aldrich, C8267),

and 2.5% methyl green dye (MCE, HYD0163). The DNA-Glycerol solution was pipetted

on top of the agarose disk and the arm electrode was rotated 180° while the DNA-

glycerol solution and the agarose disk stayed adhered via surface tension. The arm

electrode was lowered until the DNA solution was in contact with the tissue. Once there

is contact between all components of the system, a closed circuit is formed.

Each tissue was electroporated with five 50V pulses for 50ms each with a 250ms

break between pulses. The arm electrode was slowly raised to disconnect from the

tissue, and the tissue was returned to the incubator.

The following plasmids were used in this study: pCMV-eGFP (Levine Lab),

pCMV-mCherry (ClonTech, 632524), pCMV-Cre (Addgene, 123133).

Retinal explant culture tissue fixation

Culture medium was removed and replaced with room temperature PBS to wash

3 times for 10 minutes each. 100µL 1X PBS was pipetted on top of tissue for each

wash. Tissues were fixed in 4% PFA for 2 hours at room temperature or at 4°C for 24

hours. PFA was removed and fixed tissues were washed 3 times with 1 mL 1X PBS for

30 minutes at room temperature. Fixed tissues were stored at 4°C in 1 mL 1X PBS with

0.01% sodium azide (Sigma Aldrich, 08591).

Whole mount immunohistochemistry

Each quadrant of the retinal tissue was separated and cut in half to generate 8

tissue fragments for staining. Individual tissue fragments were transferred into a 96-well

plate (Corning, CLS3628). Tissues were washed 3 times with 1mL 1X PBS containing

1% Triton-X (1% PBST), followed by 100µL 10% normal donkey serum (NDS)

(SouthernBiotech, 003001) in 1% PBST (blocking buffer) for 2 hours at room

temperature. Blocking buffer was removed and tissues were stained with primary

antibodies in 1% NDS blocking buffer for 2 hours at room temperature or 24 hours at

4°C. Tissues were washed 3 times with 200µL 1% PBST for 30 minutes each.

Secondary antibodies in 1% NDS blocking buffer were applied for 2 hours at room

temperature. The secondary antibody was removed, and tissue was washed 3 times

with 200µL 1% PBST for 30 minutes per wash. Stained tissues were mounted on

Colorfrost plus slides (Fisherbrand, 12-550-17) in Fluoromount G (Invitrogen, 00-4958-

02), coversliped, allowed to dry at room temperature overnight, then stored at 4°C until

imaged.

The following primary antibodies were used: α-BrdU (1:200, Abcam, ab6326), α-

IBA1 (1:500, FUJIFILM, 019-19741), and α-SOX9 (1:500, Chemicon, ab5535).

Secondary antibodies conjugated to Alexa 488 (Invitrogen, A21206), Alexa 568

(Invitrogen, A10042), and Alexa 647 (Intvitrogen, 21247) were each diluted at 1:500.

BrdU treatment and detection

16µM BrdU was added to culture media at varying timepoints in culture per

experiment. BrdU media was replenished daily until fixation. *Ex vivo* tissues were

treated with BrdU for 24-72 hours depending on the experiment. Following treatment,

tissues were fixed with PFA as described above. A 1/8 piece of tissue was selected at

random and cut using as scalpel from the retinal culture to stain. 2N HCl (Fisher

Scientific, SA541) was used for antigen retrieval for 45 minutes at room temperature,

followed by a 20-minute neutralization with 0.1M sodium borate (pH 8.5) (Millipore

Sigma, 1066690010). Tissues were then immunostained as described above.

Imaging

Cultures were imaged using a fluorescent stereoscope (Nikon SMZ1270i) every

48 hours post-culture establishment, and every 24 hours post-electroporation monitor

for morphological, structural, or fluorescent protein expression changes within the

tissue. The Zeiss LSM 710 confocal microscope was used to image tissue cultures after

fixation alone or fixation and staining. Images were analyzed using FIJI version 2.9.0.

Orthogonal slice projections

To observe structural changes in *ex vivo* retinal tissue cultures, 40X Z-Stack

images were obtained from the Zeiss LSM 710 confocal microscope with 1µm steps

between Z slices. Using FIJI version 2.9.0, 354.25 by 11.76 pixel ROIs were selected at

random but angled from tissue periphery toward optic nerve head. These ROIs were

converted to 3D projections and rotated to view as a cross section.

IV.   **Quantification and Statistical Analysis**

All cell counting and tissue volume calculations were calculated from 20X Z-

stack, tile confocal images using Imaris 10.1 software. No statistical methods were used

to predetermine sample sizes. GraphPad Prism 10 was used for the following statistical

tests: All proportion values were arcsine transformed prior to statistical analysis.

Comparisons between more than two groups were made by two-way ANOVA with

Tukey post-hoc test. Comparisons between two groups within the same tissue, as in

figure 7, were made using paired two-tailed t tests. Differences were considered

statistically significant at $p < 0.05$. Data are presented as mean ± SEM. Statistical

significance is indicated with asterisks: *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p <$

0.0001.

References

1. Lucas RJ. Mammalian Inner Retinal Photoreception. *Curr Biol*. 2013;23(3):R125-R133.
doi:10.1016/j.cub.2012.12.029

2. Vecino E, Rodriguez FD, Ruzafa N, Pereiro X, Sharma SC. Glia–neuron interactions in the
mammalian retina. *Prog Retin Eye Res*. 2016;51:1-40. doi:10.1016/j.preteyeres.2015.06.003

3. Hu Y, Wang X, Hu B, et al. Dissecting the transcriptome landscape of the human fetal neural
retina and retinal pigment epithelium by single-cell RNA-seq analysis. *PLoS Biol*.
2019;17(7):e3000365. doi:10.1371/journal.pbio.3000365

4. Cornwall MC, Fein A, Macnichol Jr. EM. Cellular mechanisms that underlie bleaching and
background adaptation. *J Gen Physiol*. 1990;96(2):345-372.

5. Sparrow JR, Parish CA, Hashimoto M, Nakanishi K. A2E, a Lipofuscin Fluorophore, in Human
Retinal Pigmented Epithelial Cells in Culture. *Invest Ophthalmol Vis Sci*. 1999;40(12):2988-
2995.

6. Ueki Y, Karl MO, Sudar S, et al. P53 is required for the developmental restriction in Müller
glial proliferation in mouse retina. *Glia*. 2012;60(10):1579-1589. doi:10.1002/glia.22377

7. Lee JE, Liang KJ, Fariss RN, Wong WT. Ex vivo Dynamic Imaging of Retinal Microglia using
Time-lapse Confocal Microscopy. *Invest Ophthalmol Vis Sci*. 2008;49(9):4169-4176.
doi:10.1167/iovs.08-2076

8. Jones MK, Lu B, Chen DZ, et al. In Vitro and In Vivo Proteomic Comparison of Human Neural
Progenitor Cell-Induced Photoreceptor Survival. *PROTEOMICS*. 2019;19(3):1800213.
doi:10.1002/pmic.201800213

9. Ueki Y, Reh TA. EGF stimulates müller glial proliferation via a BMP-dependent mechanism.
Glia. 2013;61(5):778-789. doi:10.1002/glia.22472

10. Löffler K, Schäfer P, Völkner M, Holdt T, Karl MO. Age-dependent Müller glia neurogenic
competence in the mouse retina. *Glia*. 2015;63(10):1809-1824. doi:10.1002/glia.22846

11. Schnichels S, Kiebler T, Hurst J, et al. Retinal Organ Cultures as Alternative Research Models.
Altern Lab Anim. 2019;47(1):19-29. doi:10.1177/0261192919840092

12. Schaeffer J, Delpech C, Albert F, Belin S, Nawabi H. Adult Mouse Retina Explants: From ex
vivo to in vivo Model of Central Nervous System Injuries. *Front Mol Neurosci*.
2020;13:599948. doi:10.3389/fnmol.2020.599948

13. Hong Yee C, Aoki S, Uchihashi K, et al. The Air Liquid-interface, a Skin Microenvironment,
Promotes Growth of Melanoma Cells, but not Their Apoptosis and Invasion, through
Activation of Mitogen-activated Protein Kinase. *Acta Histochem Cytochem*. 2010;43(1):1-7.
doi:10.1267/ahc.09036

14. Jiang D, Schaefer N, Chu HW. Air–Liquid Interface Culture of Human and Mouse Airway
Epithelial Cells. In: Alper S, Janssen WJ, eds. *Lung Innate Immunity and Inflammation:*
Methods and Protocols. Methods in Molecular Biology. Springer; 2018:91-109.
doi:10.1007/978-1-4939-8570-8_8

15. Venkatesh A, Ma S, Langellotto F, Gao G, Punzo C. Retinal gene delivery by rAAV and DNA
electroporation. *Curr Protoc Microbiol*. 2013;0 14:Unit-14D.4.
doi:10.1002/9780471729259.mc14d04s28

16. Ail D, Malki H, Zin EA, Dalkara D. Adeno-Associated Virus (AAV) - Based Gene Therapies for
Retinal Diseases: Where are We? *Appl Clin Genet*. 2023;16:111-130.
doi:10.2147/TACG.S383453

17. Jeon CJ, Strettoi E, Masland RH. The Major Cell Populations of the Mouse Retina. *J Neurosci*.
1998;18(21):8936-8946. doi:10.1523/JNEUROSCI.18-21-08936.1998

18. Mahabadi N, Khalili YA. Neuroanatomy, Retina. In: *StatPearls [Internet]*. StatPearls
Publishing; 2023. Accessed December 3, 2023.
https://www.ncbi.nlm.nih.gov/books/NBK545310/

19. Kato M, Sudou N, Nomura-Komoike K, Iida T, Fujieda H. Age- and cell cycle-related
expression patterns of transcription factors and cell cycle regulators in Müller glia. *Sci Rep*.
2022;12(1):19584. doi:10.1038/s41598-022-23855-w

20. Pellissier LP, Hoek RM, Vos RM, et al. Specific tools for targeting and expression in Müller
glial cells. *Mol Ther - Methods Clin Dev*. 2014;1. doi:10.1038/mtm.2014.9

21. Poché RA, Furuta Y, Chaboissier MC, Schedl A, Behringer RR. Sox9 Is Expressed in Mouse
Multipotent Retinal Progenitor Cells and Functions in Müller Glial Cell Development. *J Comp*
Neurol. 2008;510(3):237-250. doi:10.1002/cne.21746

22. Feng G, Mellor RH, Bernstein M, et al. Imaging neuronal subsets in transgenic mice
expressing multiple spectral variants of GFP. *Neuron*. 2000;28(1):41-51. doi:10.1016/s0896-
6273(00)00084-2

23. Vazquez-Chona FR, Clark AM, Levine EM. *Rlbp1* Promoter Drives Robust Müller Glial GFP
Expression in Transgenic Mice. *Investig Opthalmology Vis Sci*. 2009;50(8):3996.
doi:10.1167/iovs.08-3189

24. Wang J, O'Sullivan ML, Mukherjee D, Puñal VM, Farsiu S, Kay JN. Anatomy and spatial
organization of Müller glia in mouse retina. *J Comp Neurol*. 2017;525(8):1759-1777.
doi:10.1002/cne.24153

25. Rashid K, Akhtar-Schaefer I, Langmann T. Microglia in Retinal Degeneration. *Front Immunol*.
2019;10:1975. doi:10.3389/fimmu.2019.01975

26. Wohl SG, Schmeer CW, Witte OW, Isenmann S. Proliferative Response of Microglia and
Macrophages in the Adult Mouse Eye after Optic Nerve Lesion. *Investig Opthalmology Vis*
Sci. 2010;51(5):2686. doi:10.1167/iovs.09-4537

27. Vázquez-Chona FR, Swan A, Ferrell WD, et al. Proliferative reactive gliosis is compatible with
glial metabolic support and neuronal function. *BMC Neurosci*. 2011;12(1):98.
doi:10.1186/1471-2202-12-98

28. Nomura-Komoike K, Saitoh F, Komoike Y, Fujieda H. DNA Damage Response in Proliferating
Müller Glia in the Mammalian Retina. *Investig Opthalmology Vis Sci*. 2016;57(3):1169.
doi:10.1167/iovs.15-18101

29. Goldman D. Müller glia cell reprogramming and retina regeneration. *Nat Rev Neurosci*.
2014;15(7):431-442. doi:10.1038/nrn3723

30. Hoang T, Wang J, Boyd P, et al. Gene regulatory networks controlling vertebrate retinal
regeneration. *Science*. 2020;370(6519):eabb8598. doi:10.1126/science.abb8598

31. Chen C, Bai X, Ding Y, Lee IS. Electrical stimulation as a novel tool for regulating cell
behavior in tissue engineering. *Biomater Res*. 2019;23:25. doi:10.1186/s40824-019-0176-8

32. Hernández-Bule ML, Paíno CL, Trillo MÁ, Úbeda A. Electric Stimulation at 448 kHz Promotes
Proliferation of Human Mesenchymal Stem Cells. *Cell Physiol Biochem*. 2014;34(5):1741-
1755. doi:10.1159/000366375

33. Katoh K. Effects of Electrical Stimulation of the Cell: Wound Healing, Cell Proliferation,
Apoptosis, and Signal Transduction. *Med Sci*. 2023;11(1):11. doi:10.3390/medsci11010011

34. Liu M, Xie D, Zeng H, Zhai N, Liu L, Yan H. Direct-current electric field stimulation promotes
proliferation and maintains stemness of mesenchymal stem cells. *BioTechniques*.
2023;74(6):293-301. doi:10.2144/btn-2022-0112

35. Martín D, Bocio-Nuñez J, Scagliusi SF, et al. DC electrical stimulation enhances proliferation
and differentiation on N2a and MC3T3 cell lines. *J Biol Eng*. 2022;16(1):27.
doi:10.1186/s13036-022-00306-8

36. Jullien N. Regulation of Cre recombinase by ligand-induced complementation of inactive
fragments. *Nucleic Acids Res*. 2003;31(21):131e-1131. doi:10.1093/nar/gng131

37. Tian X, Zhou B. Strategies for site-specific recombination with high efficiency and precise
spatiotemporal resolution. *J Biol Chem*. 2021;296:100509. doi:10.1016/j.jbc.2021.100509

38. Stachowski K, Norris AS, Potter D, Wysocki VH, Foster MP. Mechanisms of Cre recombinase
synaptic complex assembly and activation illuminated by Cryo-EM. *Nucleic Acids Res*.
2022;50(3):1753-1769. doi:10.1093/nar/gkac032

39. Stollmeier M, Kahlert S, Zuschratter W, et al. Air-liquid interface cultures trigger a metabolic
shift in intestinal epithelial cells (IPEC-1). *Histochem Cell Biol*. 2023;159(5):389-400.
doi:10.1007/s00418-023-02180-x

40. Gueven N, Glatthaar B, Manke HG, Haemmerle H. Co-cultivation of rat pneumocytes and
bovine endothelial cells on a liquid-air interface. *Eur Respir J*. 1996;9(5):968-975.
doi:10.1183/09031936.96.09050968

41. Vera MS, Simón MV, Prado Spalm FH, et al. Ceramide-1-phosphate promotes the migration
of retina Müller glial cells. *Exp Eye Res*. 2021;202:108359. doi:10.1016/j.exer.2020.108359

42. Goldman D. Müller glial cell reprogramming and retina regeneration. *Nat Rev Neurosci*.
2014;15(7):431-442. doi:10.1038/nrn3723

43. Tackenberg MA, Tucker BA, Swift JS, et al. Müller cell activation, proliferation and migration
following laser injury. *Mol Vis*. 2009;15:1886-1896.

44. Bringmann A, Iandiev I, Pannicke T, et al. Cellular signaling and factors involved in Müller
cell gliosis: Neuroprotective and detrimental effects. *Prog Retin Eye Res*. 2009;28(6):423-
451. doi:10.1016/j.preteyeres.2009.07.001

45. Nomura-Komoike K, Saitoh F, Fujieda H. Phosphatidylserine recognition and Rac1 activation
are required for Müller glia proliferation, gliosis and phagocytosis after retinal injury. *Sci
Rep*. 2020;10(1):1488. doi:10.1038/s41598-020-58424-6

46. Jones BW, Watt CB, Frederick JM, et al. Retinal remodeling triggered by photoreceptor
degenerations. *J Comp Neurol*. 2003;464(1):1-16. doi:10.1002/cne.10703

47. Kumar A, Nune K, Misra R. Understanding the response of pulsed electric field on osteoblast
functions in three-dimensional mesh structures. *J Biomater Appl*. 2016;31(4):594-605.
doi:10.1177/0885328216658376

48. Zhu W, Ye T, Lee SJ, et al. Enhanced neural stem cell functions in conductive annealed
carbon nanofibrous scaffolds with electrical stimulation. *Nanomedicine Nanotechnol Biol
Med*. 2018;14(7):2485-2494. doi:10.1016/j.nano.2017.03.018

49. Snyder S, DeJulius C, Willits RK. Electrical Stimulation Increases Random Migration of
Human Dermal Fibroblasts. *Ann Biomed Eng*. 2017;45(9):2049-2060. doi:10.1007/s10439-
017-1849-x

50. Zhao H, Steiger A, Nohner M, Ye H. Specific Intensity Direct Current (DC) Electric Field
Improves Neural Stem Cell Migration and Enhances Differentiation towards βIII-Tubulin+
Neurons. *PLoS ONE*. 2015;10(6):e0129625. doi:10.1371/journal.pone.0129625

51. Dong Z yong, Pei Z, Li Z, Wang Y ling, Khan A, Meng X ting. Electric field stimulation induced
neuronal differentiation of filum terminale derived neural progenitor cells. *Neurosci Lett*.
2017;651:109-115. doi:10.1016/j.neulet.2017.05.001

52. Ryan CNM, Doulgkeroglou MN, Zeugolis DI. Electric field stimulation for tissue engineering
applications. *BMC Biomed Eng*. 2021;3:1. doi:10.1186/s42490-020-00046-0

A

Culture wild-type retinas — Electroporate with **pCMV-eGFP** plasmid — Fix *ex vivo* retinas — Immunostain for **anti-SOX9**

0DEV   14DEV   17DEV   Post-Fixation

B

Edge (100 μm)
Center

C   pCMV-eGFP

14DEV

Scale Bar: 200uM

D

| pCMV-eGFP | SOX9 | MERGE |

14DEV Center

14DEV Edge

Scale Bar: 50uM

E

Culture **Thy1:YFP** retinas — Electroporate with **pCMV-mCherry** plasmid — Fix and image retinas

0DEV   14DEV   17DEV

F   pCMV-mCherry

14DEV

Scale Bar: 200uM

G

| Thy1:YFP | CMV-mCherry | Merge |

14DIV Center

14DIV Edge

Scale Bar: 50uM

H

Cell types transfected at 14DEV

○ Transfected Müller glia (Sox9+eGFP+/eGFP+)

■ Transfected neurons (Thy1:YFP+ mCherry+/mCherry+)

Y-axis: % Transfected Cells

X-axis: Edge, Center, Edge & Center

**** (for each group)

Edge   Center

A **1DEV THY1:YFP**

DAPI YFP

Scale Bar 100uM

ONL
INL
GCL

IPL →

ONH

B **14DEV THY1:YFP**

ONL
INL
GCL

IPL →

ONH

C **1DEV RLBP1:GFP**

DAPI GFP

ONL
INL
GCL

*

ONH

D **14DEV RLBP1:GFP**

ONL
INL
GCL

*

ONH

A

0DEV

Begin Culture	Electroporation	Deliver BrdU	Fix Tissues
0DEV	**1DEV**	**4DEV**	**5DEV**

14DEV

Begin Culture	Electroporation	Deliver BrdU	Fix Tissues
0DEV	**14DEV**	**17DEV**	**18DEV**

40X
Imaging

B

No Electroporation (*-EP*) | Electroporation without DNA (*+EP*) | Electroporation with DNA (*+EP^{DNA}*)

BrdU

Electroporated at 1DEV

Electroporated at 14DEV

Scale Bar 100 uM

C

BrdU+ cells following electroporation

Scale Bar 100 uM