**Human telomere length is chromosome specific and conserved across**
**individuals**

Kayarash Karimian[1,2], Aljona Groot[3], Vienna Huso[1,2], Ramin Kahidi[4], Kar-Tong Tan[5,6,7],
Samantha Sholes[1,2,8], Rebecca Keener[9], John F. McDyer[10], Jonathan K. Alder[10], Heng
Li [11,12], Andreas Rechtsteiner[3], Carol W. Greider[1,3, *]

[1]Department of Molecular Biology and Genetics, Johns Hopkins University School of
Medicine, Baltimore, MD, 21205, USA

[2]Biochemistry, Cellular and Molecular Biology Graduate Program, Johns Hopkins
University School of Medicine, Baltimore, MD, 21205, USA

[3]Department of Molecular Cell and Developmental Biology, University of California,
Santa Cruz,

[4] University of Calgary, Calgary, AB, Canada

[5] Harvard Medical School, Department of Genetics, Boston, MA

[6] Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA

[7] Broad Institute, Cancer Program, Cambridge, MA,

[8] Present address Merck & Co., 770 Sumneytown Pike, West Point, PA 19486

[9] Department of Biomedical Engineering, Johns Hopkins University,

[10] Pulmonary, Allergy, Critical Care, and Sleep Medicine Division, Department of
Medicine, University of Pittsburgh

[11] Dana-Farber Cancer Institute, Department of Data Sciences, Boston, MA,

[12] Harvard Medical School, Department of Biomedical Informatics, Boston, MA

*Corresponding Author

Carol W Greider, PhD
Department of Molecular, Cell & Developmental Biology
University of California, Santa Cruz
1156 High Street
Santa Cruz, CA 95064
E-mail: cgreider@ucsc.edu

Short telomeres cause age-related disease and long telomeres predispose to cancer;

however, the mechanisms regulating telomere length are unclear. Current methods for

telomere length measurement are not direct, precise, or widely accessible. Here we

describe a direct nanopore telomere profiling assay using an Oxford Nanopore

Technology (ONT) MinION that is easy to implement, precise, and cost effective with

broad applications in research, and the clinic. Telomere length measurement is

currently used in clinical settings to make highly consequential treatment decisions.

Using patient samples our method returned similar results to the clinical, FlowFISH

assay.  Telomere profiling enables mapping of the telomere to specific chromosomes,

and we identified both chromosome-specific and haplotype-specific telomere length

distribution with a remarkable 6kb difference between some telomere lengths.  Further,

we found that specific chromosome ends were consistently shorter or longer than the

average length across 150 individuals. The presence of conserved chromosome end-

specific telomere lengths suggests there are new paradigms in telomere biology that are

yet to be explored. Understanding these mechanisms will allow deeper insights into

telomere biology that can lead to new approaches to disease.

**Introduction**

Human health is profoundly affected by telomere length, yet the detailed mechanism of

length regulation is poorly understood. Telomere length is maintained as an equilibrium

distribution with constant shortening at each round of DNA replication, which is

counterbalanced by *de novo* addition of new telomere repeats by the enzyme

telomerase [1]. Failure to maintain the length distribution leads to inherited Short

Telomere Syndromes and age-related degenerative disease such pulmonary fibrosis,

immunodeficiency, and bone marrow failure [2]. Conversely long telomeres predispose

people to cancer [3], and a cluster of mutations that increases telomerase activity is one

of the most common mutational signatures in cancer [4,5].

Precisely how telomerase action maintains a length equilibrium is of great interest. The

prevailing 'protein counting' model for length maintenance [6,7] proposes that proteins that

bind telomeric TTAGGG repeats negatively regulate telomere elongation in *cis.* This is

supported by evidence that telomerase stochastically elongates short telomeres more

frequently than long telomeres [8]. Together these studies propose that the length

equilibrium is maintained by telomerase lengthening short telomeres to precisely

counterbalance shortening of all telomeres. An implication of this model is that all

telomeres will be regulated around a similar mean length distribution.

Methods for measuring telomere length have had significant influence on length

regulation models. Southern blotting was first established to measure the length of all

telomeres in a large population of cells simultaneously [9] Szostak, 1982 #170}.

Telomere Southerns probed with the telomere repeat sequence reveal a distribution of

lengths of every chromosome end in every cell in the population. The distribution of

lengths is difficult to quantitate, and absolute lengths differ significantly between labs [10].

The protein counting model was based on data from Southern blots which explains the

focus on the regulation of the distribution of lengths across all telomeres [11-13]. The

adoption of FlowFISH as the clinical standard for diagnosis of telomere diseases [14-16]

may have reinforced the thinking in the field that telomeres are maintained around a

global length distribution. FlowFISH measures the average of all the TTAGGG repeats

in patients' lymphocytes and is normalized to the median length of all telomeres on a

Southern blot [17]. The fact that this method is robust and accurately identifies telomere

mediated disease may imply to some that the global telomere length average is the

biologically relevant measurement, when in fact it might not be.

qFISH allows measurement of individual telomeres by fluorescence by *in situ*

hybridization on each telomere in a metaphase spread [18].  qFISH experiments have

suggested that telomeres on all chromosome arms are not globally regulated around a

common length distribution [19-21], however this data in not yet reconciled with the

prevailing protein counting model of length regulation.  This may be because

sophisticated microscopy and image analysis is required for qFISH and so this method

is not readily accessible. Here we describe nanopore telomere profiling, which

measures the length of each individual telomere in the cell at nucleotide resolution. With

this new method we establish that indeed individual telomeres on specific chromosome

ends are maintained around their own unique length distributions, and the means of

these distributions can differ by more than 6kb. Nanopore telomere profiling using the

ONT MinION is easy to implement, precise, cost effective, and will be of broad use in

research, clinical, and biotechnology settings. Telomere profiling represents a paradigm

shift in telomere analysis and will enable exploration of entirely new areas of telomere

biology.

**Results**

To determine whether human telomeres are maintained around a common length

distribution across all chromosomes or if specific chromosome ends maintain their own

unique length distributions, we developed a method to enrich telomere so that we could

sequence them by Oxford Nanopore Technology (ONT) long read

sequencing. We ligated the telomeric ends with a biotinylated oligonucleotide

(TeloTag) that contains a multiplexing barcode and restriction enzyme sites. Following

ligation, we pulled down the tagged telomeres with streptavidin beads and released

them by restriction enzyme digestion prior to sequencing (Fig. 1a). To assess the

enrichment efficiency, we prepared libraries from both telomere enriched and non-

enriched samples and sequenced them on an ONT MinION. Enrichment was efficient,

recovered ~17% of total telomere input (Extended Data Fig. 1. b, c), and resulted in a

~3400-fold increase in sequenced telomeres (Extended Data Fig.1d and methods). We

routinely multiplexed samples and generated ~50,000 telomere reads per flow cell with

an average fragment length (subtelomere + telomere) of ~20kb.  The cost per sample

when multiplexing was approximately $75 (see methods).

**Bioinformatic analysis of telomere length**

We developed a bioinformatic pipeline to determine both "bulk length" (all telomeres) as

well as chromosome end-specific telomere length. We used ONT Guppy base calling

and filtered for reads containing telomere repeats (Methods). We initially determined

telomere length using a method we used in yeast [22] that has been previously used in

human cells [23-25].  Reads are first mapped to a reference genome with a defined

telomere/subtelomere boundary and telomere length is defined as base pairs from the

boundary to the end of the read (Methods).  However, we found heterogeneity in

subtelomere sequences among individuals, where sometimes the subtelomere was

slightly longer or shorter than in the reference genome which caused overcalling or

under calling of telomere lengths (Extended Data Fig. 2). To overcome this, we

developed TeloNP, an algorithm to define the subtelomere boundary and measure

telomere length directly from the nanopore sequencing reads while taking systematic

nanopore basecalling errors [26] into account.  The subtelomere boundary was set at the

start of a sustained discontinuity in the telomere repeat pattern when scanning from the

end of the read towards the subtelomere (Fig. 1b and Extended Data Fig. 3) (see

methods). Telomere length was defined as the base pairs from the TeloTag to the

subtelomere boundary on the reads.

To examine whether telomere length determined by TeloNP after Guppy base calling

accurately represents the true length of the telomere repeats, we examined the

electrical current signals from the flow cells. We developed TeloPeakCounter to count

the repeated current peaks from the (TTAGGG)n in the nanopore Fast5 signal data to

test base calling by Guppy (Extended Data Fig. 4a, b) (see methods). Length

determined by TeloNP after Guppy base calling was in good agreement with

TeloPeakCounter.  We note the new ONT Dorado base caller overestimated telomere

length compared with both TeloPeakCounter and Guppy (Extended Data Fig. 4 c, d).

We therefore adopted Guppy base calling for our analyses.

**Nanopore telomere profiling accurately and reproducibly reports telomere length**

We performed telomere profiling by sequencing DNA from PBMCs (see methods) of

people ranging from 0 years to 90 years of age (Fig. 1d) and found agreement with

telomere lengths on a Southern blot (Fig. 1c). To test reproducibility, we measured

telomere length of DNA from one individual on the same flow cell (intra-assay) (Fig. 1e

and Extended Data Fig. 1e) (correlation of variation, CV 1.3%) or on different flow cells

(inter-assay) Fig. 1f and Extended Data Fig. 1f (CV 2.4%). This low variability compares

well with FlowFish, the that has an inter-assay CV of 2.2%, and considerably

outperformed the frequently used qPCR assay that has an inter-assay CV of 25.0% [15].

In addition, we tested inter-lab variability by measuring telomere length of seven

samples where the same DNA was enriched and sequenced by two different people in

two different labs (Johns Hopkins and UCSC) and found highly reproducible results

(mean difference of 104.7 bp with SEM of +/- 34 bp) (Fig. 1g). To determine whether

any fragment length bias of nanopore sequencing could skew telomere length

determination, we compared restriction enzyme cutting with a combination of *BamHI*

and *EcoRI* which generates fragments ~9 kb, or with *AsiSI* and *PvuI,* which generate

fragments ~25 kb. We found similar telomere lengths (Extended Data Fig. 5a, b)

indicating fragment length in these size ranges did not have detectable bias on telomere

length determination.

**Telomere profiling determines telomere shortening with age at nucleotide**

**resolution**

Telomere length is known to shorten with age [15,27-30], however previous methods could

not measure telomere length at nucleotide resolution.  To test the dynamic range, we

first applied nanopore telomere profiling to DNA samples of 11 individuals from 0-84

years of age and ranked them based on descending order for telomere length. We then

did a Southern blot on the same DNA. Telomere profiling predicted the rank order of

telomere lengths and captured the wide dynamic range of the Southern blot (Fig. 2a and

b). Southern blotting does not measure the shortest telomeres because telomere

repeats are required for probe hybridization on a Southern. We plotted the 1st, 10th and

50th percentile of telomere length as determined by telomere profiling and observed a

decrease of the 50th and 10th percentile as the mean length shortened. However, the

1st percentile telomere length did not decrease suggesting there is a threshold length in

PBMC's of ~1,100 bp below which telomeres cannot be maintained (Fig 2c).

To directly compare nanopore telomere profiling to FlowFISH, we conducted nanopore

telomere profiling on whole blood and PBMCs (see methods) of 132 healthy individuals

ranging from 0 to 90 years of age (Fig. 2d).  Using the mean telomere length for each

individual, we defined $90^{th}$, $50^{th}$ and $10^{th}$ intervals of telomere length at each age using

statistical methods used for FlowFISH [15].  While the shapes of the curves are very

similar between telomere profiling and FlowFISH, the absolute lengths of the telomeres

are longer for FlowFISH. Nanopore telomere profiling of cord blood showed a telomere

length distribution with a mean of 7,986 +/- 245 bp (Fig. 2d) across 18 samples. This is

shorter than the average for FlowFISH and with less variance (Fig. 2e) [15]. Average cord

blood telomere length estimates measured by FlowFISH vary from ~18kb [31] to ~9kb [32]

to ~11 kb [15]. FlowFISH fluorescence signal is normalized to Southern blots, which

includes some subtelomeric sequences and this may account for the longer telomere

lengths of FlowFISH. Furthermore, Southern blot estimated telomere lengths are known

to vary between laboratories [10]. In contrast, nanopore telomere profiling offers a precise

readout in base pairs that can be directly compared between laboratories.

To compare our method directly to FlowFISH, we sequenced 5µg of archived DNA from

blinded samples of IPF patients previously diagnosed with Short Telomere Syndromes.

Telomere profiling showed that bulk telomere length in most IPF samples were similar

to the FlowFISH measurement (Fig. 2f). FlowFISH uses flow cytometry and can

distinguish telomere lengths in specific cell types from whole blood samples [33,34] and

some samples have discordant lymphocyte and granulocyte telomere lengths [15,35,36],

Nanopore telomere profiling will report the average length from all cell types in the

samples.   Thus, while nanopore telomere profiling likely can be used for diagnosing

short telomere patients in the future, additional development such as isolation of specific

cell types may help to capture heterogeneity of clinical samples.

**Human telomeres have chromosome end-specific length and haplotype-specific**

**length differences**

To determine whether humans have chromosome end-specific telomere length, we first

examined telomeres from the diploid HG002 cell line for which a high-quality reference

genome is available [37]. Human subtelomeres contain many blocks of homology shared

between different telomeres (paralogy blocks) [23,38]. Simulation of long read data from

CHM13 references genome showed that *minimap2* [39,40] can assign simulated reads to

the correct telomere with high accuracy using 10kb of subtelomere sequence [26]. We

isolated DNA from HG002 cell line, sequenced the telomeres and mapped reads with

an average total length of 16.4 kb (4.6 kb telomere repeats and 11.8 kb sub-telomeric

sequence, on average) to the HG002 reference genome using *minimap2* using a

customized filtering pipeline (methods). Seventy-seven chromosome ends passed our

quality filters, and we found 66 ends had significant differences in length distribution

from the grand mean (Fig. 3. a, b). In addition to chromosome end-specific lengths, we

also found that some telomeres showed significant differences between the maternal

and paternal haplotypes. In some cases, remarkably, there was more than 6kb

difference in mean length, for example for chromosome 1p Maternal (1pM) and 1p

Paternal (1pP). Thus, like in yeast [22], humans have chromosome end-specific telomere

length distributions.

**Chromosome specific telomere lengths are conserved across individuals.**

To determine whether chromosome end-specific differences were conserved across a

broad population, we used *minimap2* to map telomere reads from 150

individuals to the subtelomere sequences from the recently released pangenome

containing 47 high quality T2T assemblies representing 94 haploid genomes [41] and

filtered for reads with >2kb of alignment (see methods).

We removed the acrocentric and X Y chromosome ends because the

high rate of meiotic recombination between these ends across a population would not

allow them to map uniquely [42].

*Minimap2* map quality score (mapq) reports quality of alignment when mapping reads to

a single reference genome but is not optimized for mapping to the multiple genomes

present in the pangenome as most reads have multiple near identical alignments and

thus get low mapq. We compared the pangenome alignment of reads from our 300

haploid genomes to the alignment in three high quality haploid reference T2T genomes

CHM13, HG002 maternal and HG002 paternal to establish if reads reproducibly

mapped to the same subtelomere in each reference genome. The reads were mapped

to the respective haploid genome using mapq of 60 (Fig. 4 a, b, c).  Pangenome

chromosomes are shown by the columns in the matrix heatmaps and T2T haploid

genome chromosomes are shown by the rows. The color in each square indicates what

fraction of reads mapping to chromosome in the respective haploid T2T genome.  The

diagonal indicates the fraction of reads mapping to the same chromosome end in the

pangenome and the respective haploid genomes. Of all reads, 87% mapped to the

same chromosome end in the pangenome and CHM13, 90% in the pangenome and

HG002 maternal and 88% in the pangenome and HG002 paternal. We also quantified

the percent of reads that mapped to the same chromosome end in the pangenome and

all three haploid reference genomes (Fig. 4d and Extended Data Fig. 5 a, b, c). For 33

of the 39 chromosomes ends, 100-60% of the reads mapped to the same chromosome

end in the pangenome and all three haploid genomes.  Six chromosome ends had

between 10-20% of reads map to the same chromosome end in the pangenome and all

three haploid genomes. When we added back the acrocentric chromosomes, we found

0 reads amongst this group mapped back to the same chromosome end in all three

references (Extended Data Fig 5d), as expected for reads known to map to different

chromosome ends across a population. Together this data suggests that the reads we

found mapping to a certain pangenome chromosome map with high confidence.

To compare the telomere length of each chromosome end across the aging population,

we established the relative mean telomere length.  For each individual, their grand

mean telomere length was subtracted from the their chromosome specific telomere

length. The resulting mean centered chromosome telomere lengths were plotted with

zero indicating no difference from the individual's grand mean for that chromosome (Fig

4e). We ranked the chromosome ends by their relative telomere lengths and found that

17p, 20q and 12p tended to be the shortest telomeres in the population while 4q, 12q

and 3p tended to be the longest (Fig. 4e).   Thus, while haplotype specific differences in

telomere length are seen in a single individual (Fig. 3a), across a population, on

average, certain chromosome ends are more likely to be shorter while others are more

likely to be longer than the grand mean. Remarkably, previous work using qFISH to

measure telomere length on metaphase spreads in 10 individuals also found 17p, 20q

and 12p among the top 4 shortest and 4q, 12q and 3p among the top 8 longest ends [20]

strengthening the conclusion that some chromosome ends are reproducibly shorter or

longer than the grand mean.

To determine whether chromosome end-specific telomere lengths are present at birth,

we mapped the reads from cord blood to the pangenome and calculated the relative

mean telomere lengths as described above (Fig. 4f).  While we had fewer cord blood

samples, and therefore fewer chromosome ends met our quality filters, we found again

that 17p, 20q and 12p were shorter while 4q, 12q and 3p were longer than the grand

mean. This supports previous work [43] that suggested that telomere length at birth is

maintained with age.

293
294
**Discussion**

A fundamental understanding of the mechanisms that regulate telomere length is

essential to develop future disease treatments. When the telomere length distribution

shifts to shorter lengths, some telomeres become critically short, initiating senescence,

[44-47] and can cause age-related degenerative disease in humans [16]. Inherited mutations

that shift to a longer equilibrium predispose people to cancer [3,48] and the most frequent

somatic mutations in cancer increase telomerase levels and lengthen telomeres [4,5].

Nanopore telomere profiling will enable the dissection of how individual telomere

lengths on specific chromosomes are maintained and may play a role triggering

senescence, and ultimately in disease.

**Chromosome end-specific telomere length equilibria imply new regulatory**

**mechanisms**

The predominant protein counting model for telomere length maintenance proposes that

telomere proteins that bind TTAGGG repeats repress the elongation of a given telomere

in *cis* [6,49] and longer telomeres have more repression, allowing shorter telomeres to be

preferentially elongated [8]. This model represents a robust way to maintain a length

equilibrium [50]. However, since all telomeres have the same TTAGGG repeats, the

model predicts that all telomeres would be regulated around a shared equilibrium

length. The demonstration of end-specific lengths indicates that other, yet unknown

factors, can play a key role modifying the set point for each unique telomere length

distribution.

In yeast that lack telomerase, all chromosome end-specific length distributions

shortened at similar rates [22], suggesting telomere elongation, not shortening, is the

major influence on chromosome specific length. Telomere elongation is the sum of the

frequency of elongation of any given end (telomerase recruitment) and number of

repeats added per elongation event (telomerase processivity). When the sum of these

events, on average, equals the rates of telomere shortening, the equilibrium point is set.

However, given end-specific length distributions, it is clear that this simple view does not

represent the full complexity of the system. There must be factors at specific

chromosome ends that regulate telomerase recruitment, processivity, or both, to

establish end specific lengths. In addition, stochastic shortening such as telomere rapid

deletion [51] or replication fork collapse [52,53] may play yet unknown roles in establishing

telomere length equilibrium.

**Mechanisms that may influence end specific telomere length**

Subtelomeric sequences are obvious candidates to regulate end-specific telomere

lengths. In yeast, subtelomere DNA binding proteins can affect telomere length [54],

although the mechanism is not yet understood. The subtelomeric TAR1 element [55]

present in paralogy block 23 [25,38] was proposed to regulate telomere length, possibly

through binding CTCF and regulating expression of the lncRNA, TERRA [56-58]. Previous

studies suggested that the absence of TAR1 may correlate with shorter telomeres [25].

However, we did not find a direct relationship of the shortest telomere with those ends

described by Dubocanin *et. al.* that lack TAR1 (8q,13p,14p, 17p, 21p, 22p Xp) in our

data set. Future comprehensive analysis of the subtelomere sequences adjacent to long

and short telomeres will lead to new testable models for establishment of telomere

length equilibria.

Epigenetic modifications of DNA or histones may influence telomere length [59]. Human

and mouse subtelomeric regions are known to be methylated at CpG sites [60] and

experiments in mice suggest that loss of DNA methyltransferases results in shorter

telomeres [61]. Sequences in the subtelomere could recruit chromatin modifying enzymes

that might influence length regulation. Subtelomere sequences may also influence other

mechanisms that have been proposed to regulate telomere length including replication

timing and tethering to the nuclear periphery [62]. The availability of nanopore telomere

profiling will allow exploration of the role of these factors in establishing telomere length

equilibria.

**Chromosome end-specific length differences are present at birth and maintained**

**as telomeres shorten with age**

Telomere length is inherited from parent to child. Evidence of this comes from the

genetic anticipation in patients with Short Telomere Syndromes; short telomeres are

passed down to each generation, and the severity of disease increases across

generations [63]. Similarly, in mice heterozygous for telomerase deletion, short telomeres

are progressively passed down across 6 generations causing progressive severity of

disease [64]. Twin studies have also documented the inheritance of telomere length in

humans [65].

Analysis of chromosome end-specific telomere lengths across 150 individuals showed

specific telomeres tend to be the longest or shortest, supporting a previous study using

qFISH on 10 individuals that identified a similar set of chromosomes as the longest and

shortest [20]. Cord blood also showed that 17p, 20q and 12p were among the shortest

and 4q, 12q and 3p were among the longest ends suggesting that telomere length

differences present at birth are maintained over decades. This establishment of

chromosome-end specific telomere length equilibria at birth [43] and maintenance of the

equilibria after birth leaves little room for proposed effects of life history, psychological,

or environmental exposures [66] on telomere length. The similarity of our data with

Martens *et al.* qFISH analysis is remarkable, and our method will enable future studies

to explore the biological significance of this finding. We did not prospectively choose our

samples to be representative of the diversity of the human population, but rather to span

a wide age range. However, future studies could be powered to examine whether

certain chromosome ends are consistently the shortest or longest more broadly in a

diverse human population.

**Implications for human disease**

Being able to accurately measure chromosome-end specific telomere length has

important implications for human disease. Nanopore telomere profiling determines

nucleotide resolution of the length distribution and can distinguish the length of specific

chromosome ends unlike Southern blots, qPCR, or FlowFISH assay. Telomere profiling

employs the accessible MinION instrument that can be used in-house in any research

or clinical lab, with very low start-up costs, allowing for equitable access to telomere

length determination methods. This method provides the opportunity to prospectively

develop clinical standards analogous to those for FlowFISH and may allow clinical

length measurements in samples other than blood. In addition, having a highly

reproducible assay that can be easily automated will enable experimental approaches

to define new regulators of telomere length. The role telomere elongation in the

immortalization of cancer cells has been known since 1990 [28,67,68]. Having a precise tool

that can be automated, will allow new approaches that may exploit telomere length

modulation in cancer treatment. Finally, the identification of conserved chromosome

end-specific telomere lengths implies that new, undiscovered biological mechanisms

influence telomere length. Nanopore telomere profiling will empower the field as a whole

to dissect these mechanisms, leading to new discoveries in telomere biology.

**Methods**

**Cells and Cell Lines**

PMBCs were purchased from Stem Cell Technologies, ZenBio Inc, and Precision for Medicine.

Samples were chosen from the repository based on age to span the distribution from 0 to 95.

Cord blood was purchased from Stem Cell Technologies to represent the youngest individuals

in the population. Blood samples used to initially calibrate the assay were de-identified excess

samples from Johns Hopkins Hospital, certified as exempt by the John Hopkins IRB. HG002

cells were cultured in RPMI 1640 media (Gibco, Cat.11875093) supplemented with 2g/L

glucose 2mM L-glutamine (Glutamax, Gibco, Cat.35050061) 15% fetal bovine serum (Gibco,

Cat.26140079) and 1% penicillin-streptomycin (Gibco, Cat.15140122). PBMCs were counted

using the Luna II hemocytometer (VitaScientific, Cat.LGBD10029).

**Telomere Southern blot analysis**

Genomic DNA was isolated using the Promega Wizard gDNA kit (Cat.A1120, Promega) and

quantified by QuBit 3.0 (Thermo Fisher) using the DNA kit (Q32853; Thermo Fisher).

Approximately 1 µg of genomic DNA was restricted with *Hinf*I (NEB, Cat.R0155M) and *Rsa*I

(NEB, Cat.R0167L,) and resolved by 0.8% Tris-acetate-EDTA (TAE) agarose gel

electrophoresis (Invitrogen, Cat.EA0375BOX). 10 ng of a 1kB Plus DNA ladder (NEB,

Cat.N3200) was included as a size reference. Following denaturation (0.5 M NaOH, 1.5M NaCl)

and neutralization (1.5 M NaCl, 0.5 M Tris-HCL, pH 7.4) the DNA was transferred in 10x SSC

(3M NaCl, 0.35 M NaCitrate) to a Nylon membrane (GE Healthcare, Cat. RPN303B) by vacuum

blotting (Boekel Scientific). The membrane was UV crosslinked (Stratagene), prehybridized in

Church buffer (0.5M Na2HP04, pH7.2, 7% SDS, 1mM EDTA, 1% BSA), and hybridized

overnight at 65°C using a radiolabeled telomere fragment and ladder, as previously described

(Morrish and Greider 2009). The membrane was washed twice with a high salt buffer (2x SSC,

0.1% SDS) and twice with a low salt buffer (0.5X SSC, 0.1% SDS) at 65°C, exposed to a

Storage Phosphor Screen (GE Healthcare), and scanned on a Storm 825 imager (GE

Healthcare). The images were copied from ImageQuant TL (GE Life Sciences) to Adobe

PhotoShop CS6, signal was adjusted across the image using the curves filter, and the image

was saved as a .tif file.

**Preparation of HMW DNA**

A modified DNA extraction protocol was used to produce high molecular weight DNA based on

the Lucigen/EpiCentre's MasterPureTM Complete DNA and RNA Purification Kit A (Biosearch

Technologies, Cat MC85200).  For HG002 cell line, fresh or frozen cell pellets were osmotically

lysed in presence of 150mL of Nuclei Prep Buffer (NEB, Cat.T3052) supplemented with 5.5 mL

of Rnase A (NEB, Cat. T3018L) and 5.5 mL of RNase If (NEB, Cat. M0243L) per million cells for

15 seconds and mixed by flicking.  For PBMC or fresh blood samples, an optional PBS wash

followed by Red Blood Cell lysis step was included (10 mins at RT) prior to hypotonic lysis with

Nuclei Prep Buffer (NEB, Cat.T3052) and Rnase digestion.  Nuclei from 1 million cells were

lysed with 300 mL of lysis buffer supplemented with 20 mL of Proteinase K (20mg/mL)

(ThermoFisher, Cat. 25530049).  Lysates were incubated at 50 degrees C for a minimum of 24

hours overnight with periodic vortexing at low speeds (minimum speed to achieve swirling of the

solution).  150 mL of MPC Protein Precipitation Reagent solution from the Lucigen/EpiCentre's

MasterPureTM Complete DNA and RNA Purification Kit A was added to precipitate proteins

followed by centrifugation at 2000 x g for 30 mins.  DNA was precipitated by adding 500 mL of

cold isopropanol (100%) (Supply Store, Cat.100209) and pelleted by centrifugation (2000 x g for

20 mins).  DNA pellets were washed 3X with 70% ethanol and hydrated in pre-warmed (37°C)

Elution buffer (Qiagen, 10 mM Tris-Cl, pH 8.5. Cat. 19086) and incubated on HulaMixer™

Sample Mixer (Thermo Fisher Scientific, Cat. 15920D) at 37°C incubator overnight at 1rpm end

over end mixing.

**Annealing of TeloTags for duplex barcode assembly:**

TeloTags were prepared in 100µIL reactions with 5mM of each of the 6 permutations of

telomere splint Extended Data Fig.1A) and 30 mM of biotinylated adapter in HiFi Taq DNA

Ligase Reaction Buffer (NEB, Cat. M0647S). Annealing was done by heating to 99 degrees and

slowly decreasing the temperature 1°C /min in a Veriti™ 96-Well Thermal Cycler (Applied

Biosystems, Cat. 4375786). After annealing, reactions were diluted 1:100 in 1x Taq buffer and

kept at 4°C. The sequences of the TeloTag and splint adapter is listed in Extended Data Table

2)

**Telomere Tagging**

High molecular weight genomic DNA (gDNA) was quantified using the Qubit dsDNA BR assay

kit (Thermo Fisher Scientific, Cat.Q32850).  A total of 40 µg of gDNA was incubated with 3µl of

ClaI (NEB, Cat. R0197S) or AsiSI (NEB, Cat.R0630L) or PmeI (NEB, Cat.R0560L), or BamHI

(NEB, Cat.R0136M) for 2 hours at 37°C, with gentle flicking every 20 mins. Subsequently, the

enzyme was heat-inactivated at 65°C for 20 mins. Ligations were carried out using 4 ug of DNA

per 50 µl reaction.

Tagging reactions were done in 50 µL volume for each reaction in a MicroAmp™ TriFlex Well

PCR Reaction Plate (Applied Biosystems, Cat. A32811), with 4µl/reaction of 0.3µM duplex

TeloTag adapter, 5µl/reaction of 10X HiFi Taq DNA Ligase Reaction Buffer (NEB, Cat.

M0647S), and 1µl/reaction of HiFi Taq DNA Ligase (NEB, Cat. M0647S). The TeloTagging

reactions were incubated for 5 mins at 65°C in a Veriti™ 96-Well Thermal Cycler (Applied

Biosystems™, Cat.4375786). Ligations were done through 15 cycles of denaturing at 65°C for 1

min, followed by annealing and ligating at 45°C for 3 mins with a 15% ramp down of rate

between steps.

**Telomere Enrichment and Nanopore Sequencing**

For chromosome-specific telomere length measurements, we typically used 30-40 µg of DNA

per sample. A standard 3 mL tube of blood or 30 million PBMC produced ~200 ug of DNA. For

bulk telomere length measurements, as little as 5-10 µg of starting gDNA was employed. All

pipetting was performed using wide bore pipette tips to minimize DNA shearing, except for

addition of SPRI beads where accurate volume ratios are extremely important for successful

cleanups.  All the Telomere Tagging reactions were pooled in DNA LoBind (Eppendorf,

Cat.0030108523) tube. Cleanup and removal of excess TeloTag adapters was done using SPRI

beads (Beckman Coulter, Cat. B23318) a ratio SPRI beads to DNA of 45 µL:100 µL was used.

The samples were incubated with SPRI beads rotating end over end on a Hula mixer for 20

mins at 10rpm. SPRI beads were then separated using a DynaMag™-2 Magnet (Thermo Fisher

Scientific, Cat. 12321D) and washed while on the magnet twice with freshly made 85% ethanol.

DNA was eluted using heated (65 °C) 1X rCutsmart Buffer. The volume of elution volume was

calculated to achieve 150 ng/ml final concentration based on input DNA amount.  The eluting

SPRI beads were incubated for 20 mins at 65°C with gently flicking every 5 mins. SPRI beads

were removed using a DynaMag™-2 Magnet.

The gDNA recovery was quantified using the Qubit dsDNA BR assay kit. Tagged gDNA was

enriched using Dynabeads™ MyOne™ Streptavidin C1 (Thermo Fisher Scientific, Cat. 65001).

The beads were allowed to room temperature while being resuspended on a HulaMixer™

Sample Mixer (Thermo Fisher Scientific) at 3 rpm for 1h. A ratio streptavidin to DNA of 1

µg:250ng was used. The beads were washed once in Binding Buffer from Dynabeads™

kilobaseBINDER™ Kit and resuspended in equal volume binding buffer as eluted DNA volume.

The beads were then added to the gDNA sample and incubated at room temp at 1 rpm on a

HulaMixer™ Sample Mixer for 20 mins. Reactions can be scaled up or down as needed, though

the maximum volume of beads + gDNA + binding buffer should not exceed 1.4 ml for a single

1.5 ml Protein LoBind tube (Eppendorf, Cat. 30108442). Multiple tubes can be used and pooled

at the restriction enzyme digest step.   After binding to streptavidin, the beads were washed

using the following sequence to remove background genomic DNA: 2x kilobaseBinder wash

buffer, 2x Elution buffer (Qiagen, 10 mM Tris-Cl, pH 8.5), 1x rCutsmart Buffer.

To release telomeres, the streptavidin bead-telomere complex was resuspended in 72 µl of 1X

rCutsmart, 3 µl of PvuI (NEB, Cat. R3150S) PacI (NEB, Cat. R0547L) or EcoRI (NEB,

Cat.R0101M), and incubated at 37°C for 30 min, with periodic gentle flicking. The sample was

then heated at 65°C for 20 mins to release any bound telomeres. If multiple tubes were used,

sequential rounds of digestion can be used by adding restriction enzyme to the eluted telomere

solution from the first step and incubating with streptavidin-telomere beads in the second tube.

Recovered tagged gDNA was quantified using the Qubit dsDNA HS assay kit. The expected

recovery was approximately 0.1-0.01% of the starting gDNA sample.

Enriched telomeres were carried forward into the standard Nanopore library prep protocol from

ONT. All reactions were prepared using Ligation Sequencing Kit V10 (SQK-LSK114) kits and

sequenced on R9.4.1 (Oxford Nanopore Technologies, FLO-MIN106D) flow cells. Libraries

were eluted in 40 ml of elution buffer (Qiagen, 10 mM Tris-Cl, pH 8.5) with optional 15 mins of

incubation at 37°C to recover long molecules. Each library was split into 3 reactions. Each

reaction was sequenced on a flow cell for ~18 hours before flow cell flushing/washing using flow

cell wash kit (Oxford Nanopore Technologies EXP-WSH004) and loading of the remaining

fraction.  Reads were collected using MinKNOW software (5.7.5) without live basecalling.

**Determination of telomere/subtelomere boundary position and telomere length**

To determine the length of the telomere repeats, we tested two methods. One method is based

on determining the junction of the telomeres and subtelomeres in the respective reference

genome (CHM13, HG002 maternal and HG002 paternal) and the second method determines

the subtelomere to telomere junction in every read. For method 1, to determine the junction in a

reference genome, we developed a Python algorithm named TeloBP (Telomere Boundary

Point). TeloBP employs a rolling window approach, scanning from the telomere into the

chromosome, identifying the telomere-subtelomere junction by detecting a discontinuity in a

user defined telomeric pattern. The algorithm's default telomeric pattern is a sequence where at

least 50% of nucleotides are "GGG". As the window moves along at six nucleotide intervals, it

scores telomere similarity in 100-nucleotide segments. Variants of the telomere repeats known

to be in the subtelomere do not significantly change sequence content. The junction is defined

when the sequence content no longer matches a telomere like sequence content. This is

calculated by averaging the similarity of a sequence with a 500 bp window, marking the start of

a 50% deviation, then scanning until the increase in discontinuity plateaus, marking the

subtelomere boundary. After reads are mapped to the reference genome, for each read the

telomere length is determined as the number of base pairs from subtelomere junction in the

reference to the TeloTag. This method incorporated many variant telomere repeats into the

telomere that are not incorporated by identifying the boundary as 4X TTAGGG (Extended Data

Fig 3).

In the second method we determined the subtelomeres/telomeres boundary in each read. We

developed a version of TeloBP that considers common errors in the nanopore Guppy base

calling. These patterns are set by default based on findings in [26],

"[^GGG]GGG|[^AAA]AAA|TTAGG." for G strand and "CTTCTT|CCTGG|CCC…" C strands. But

the patterns can be user defined as new base callers are developed. We named this algorithm

TeloNP (Telomere NanoPore). Both TeloBP and TeloNP are available on Github

(https://github.com/GreiderLab).

**Custom genome for mapping telomeres**

For mapping reads to the T2T genomes CHM13 and HG002 we generated custom reference

genomes. We first extracted the terminal 500kb of chromosome end for each genome, then

removed the telomere repeats (as determined by TeloBP) from the reference genome to allow

for maximized weighing of subtelomere information for read mapping.

**Bioinformatic filtering of telomere reads**

Reads were first filtered for any of the following telomere patterns ["TTAGGGTTAGGG",

"TTAAAATTAAAATTAAAA", "CCCTCCGATA", "TGGCCTGGCCTGGCC"] based on findings in

previous literature [26]. To identify reads with a TeloTag at the end and to demultiplex samples we

performed a pairwise alignment of the 24bp barcodes with the terminal 300bp of each read

using the pairwise Alignment function in the Bio Strings package of Bioconductor

(doi:10.18129/B9.bioc.Biostrings , R package version 2.68.1,

https://bioconductor.org/packages/Biostrings). The alignment score cutoff was set so the false

discovery rate for our nanopore reads was < 1% based on random 24bp barcode sequences

and unused ONT barcode sequences. We used Minimap2 with the -x map-ont option to map

our reads to the custom genomes HG002 and CHM13 [40]. We only considered primary

alignments that started within 1 kb of the subtelomere boundary.

**Peak calling to measure telomere length with** TeloPeakCounter

To examine whether the Guppy and Dorado base caller accuracy call the telomere length in

correctly, we developed an algorithm, TeloPeakCounter, to count repeated peaks, or waves, in

the electrical signal data measured by the nanopore device. These distinct repeated waves

found in the telomere region of reads correspond to the TTAGGG telomere repeat sequences.

TeloPeakCounter analyzes and counts these distinctive, periodic wave patterns in the electric

signal data, and enables a direct measurement of telomere length. Assuming each wave

represents a 6-nucleotide telomere repeat, we can compute estimated telomere lengths for a

read. The code for TeloNP and TeloPeakCounter is available at GitHub

(https://github.com/GreiderLab)

**Mapping HG002 subtelomere to maternal and paternal alleles**

For the diploid HG002 genome some maternal and paternal subtelomere sequences are very

similar and correct assignment of reads becomes difficult. We developed a two-step mapping

procedure for mapping HG002 reads. In the first step, reads are mapped to the HG002 diploid

genome. Mapq mapping confidence scores are set low for these mappings, as the mapper can

have difficulty deciding between very similar maternal and paternal subtelomere sequences. We

applied a relatively low mapq filter cutoff of 10 to the diploid mapping. In a second step we

mapped the reads also to the maternal and paternal haploid genomes separately. Mapq scores

generally increase for the alignments to the haploid genomes. To identify high confidence

alignments, we applied a mapq cutoff of 30 to the haploid genome alignments. A read needed to

map to the same chr end in both mappings and pass the two mapq cutoffs to be considered

correctly assigned.  There were different numbers of reads for specific chromosome ends due to

the restriction enzyme sometimes cutting very near a telomere. To minimize this, we used two

different restriction enzymes for both the initial cutting and for the release. This allowed mapping

of more reads for some chromosome ends.

**Pangenome based mapping for chromosome assignment of telomeres from diverse**

**individuals**

We mapped reads to the pangenome to efficiently capture telomere length across the diverse

population. A references file of 500kb of subtelomere sequences was assembled from each of

the genomes [41] in the pangenome.  We mapped our reads from 150 individuals to this

reference. We filtered for reads that had a minimum of 2kb alignment to the pangenome

reference. To compare telomere length across individuals we removed acrocentric

chromosomes (13p, 14p, 15p, 21p, 22p) and X and Y subtelomeres (XpYp, and Xq Yq) which

recombine in the population.

**Statistical Analysis**

To determine whether HG002 chromosome specific telomere lengths were significantly different

from the individual's grand mean telomere length, we used Analysis of the Mean (ANOM).

Statistics were calculated using the R package rstatix (v0.7.0) and ANOM (v0.2)

(https://cran.rproject.org/web/packages/rstatix/index.html).

**Data Availability**

Data generated during the study will be made available in public sequence repository Sequence

Read Archive (SRA)  https://www.ncbi.nlm.nih.gov/sra and is available upon request.

**Code availability**

The python code for TeloBP, TeloNP and TeloPeakCounter is available at
https://github.com/GreiderLab
615
616
**Acknowledgements**

We thank Dr. Rachel Green for provision of laboratory space at Johns Hopkins as well as

discussions. Drs. Brendan Cormack and Christine Gao and for reviewing the manuscript. Carla

Connelly, Julie Brunelle, and Margaret Strong provided experimental and logistical assistance.

We thank Dr. Mary Armanios and her lab for help in the early stages of assay development and

Dr. Ludmilla Danilova for statistical analysis. This work was supported by NIH grant

R35CA209974 to CWG and Johns Hopkins Bloomberg Distinguished Professorship to CWG.

**Author contributions**

These authors contributed to the following aspects of this work. Conceptualization CWG, KK,
SS; Data curation, AG, KK, RK, RWK, AR;  Formal analysis, KK, RK, AR; Funding acquisition
CWG, HL; Investigation AG, VH, KK, RWK, Methodology,CWG, VH, KK, RK, RWK, AR, SS;
Project administration CWG; Resources JA, CWG, JFM, KK; Software KK, RK, HL, AR ,K-TT;
Supervision CWG, HL; Validation CWG, AG, KK, RK, AR; Visualization JA, KK, RK, AR; Writing
original draft CWG, KK Writing – review & editing JA, CWG, AG, VH, KK, AR, SS
632
633 **Competing interest declaration**

634 CWG and KK are inventors of US Patent PCT/US2023/073375 titled "methods for telomere
635 length measurement".

**References**

1    Greider, C. W. & Blackburn, E. H. Identification of a specific telomere terminal
transferase activity in *Tetrahymena* extracts. *Cell* **43**, 405-413 (1985).

2    Armanios, M. Syndromes of telomere shortening. *Annu Rev Genomics Hum Genet* **10**,
45-61 (2009).

3    Armanios, M. The Role of Telomeres in Human Disease. *Annu Rev Genomics Hum*
*Genet* **23**, 363-381 (2022).

4    Horn, S. *et al.* TERT promoter mutations in familial and sporadic melanoma. *Science*
**339**, 959-961 (2013).

5    Huang, F. W. *et al.* Highly recurrent TERT promoter mutations in human melanoma.
*Science* **339**, 957-959 (2013).

6    Marcand, S., Gilson, E. & Shore, D. A protein-counting mechanism for telomere length
regulation in yeast. *Science* **275**, 986-990 (1997).

7    Smogorzewska, A. *et al.* Control of human telomere length by TRF1 and TRF2. *Mol Cell*
*Biol* **20**, 1659-1668 (2000).

8    Teixeira, M. T., Arneric, M., Sperisen, P. & Lingner, J. Telomere length homeostasis is
achieved via a switch between telomerase- extendible and -nonextendible states. *Cell*
**117**, 323-335 (2004).

9    Blackburn, E. H. & Chiou, S. S. Non-nucleosomal packaging of a tandemly repeated
DNA sequence at termini of extrachromosomal DNA coding for rRNA in Tetrahymena.
*Proc Natl Acad Sci U S A* **78**, 2263-2267 (1981).

10   Aviv, A. *et al.* Impartial comparative analysis of measurement of leukocyte telomere
length/DNA content by Southern blots and qPCR. *Nucleic acids research* **39**, e134
(2011).

11   Marcand, S., Wotton, D., Gilson, E. & Shore, D. Rap1p and telomere length regulation in
yeast. *Ciba Found Symp* **211**, 76-93; discussion 93-103 (1997).

12   Wotton, D. & Shore, D. A novel Rap1p-interacting factor, Rif2p, cooperates with Rif1p to
regulate telomere length in Saccharomyces cerevisiae. *Genes Dev* **11**, 748-760 (1997).

13   Levy, D. L. & Blackburn, E. H. Counting of Rif1p and Rif2p on Saccharomyces
cerevisiae telomeres regulates telomere length. *Molecular and cellular biology* **24**,
10857-10867 (2004).

14   Aubert, G., Hills, M. & Lansdorp, P. M. Telomere length measurement-caveats and a
critical assessment of the available technologies and tools. *Mutation research* **730**, 59-
67 (2012).

15   Alder, J. K. *et al.* Diagnostic utility of telomere length testing in a hospital-based setting.
*Proc Natl Acad Sci U S A* **115**, E2358-E2365 (2018).

16   Armanios, M. Telomeres and age-related disease: how telomere biology informs clinical
paradigms. *J Clin Invest* **123**, 996-1002 (2013).

17   Rufer, N., Dragowska, W., Thornbury, G., Roosnek, E. & Lansdorp, P. M. Telomere
length dynamics in human lymphocyte subpopulations measured by flow cytometry [see
comments]. *Nat Biotechnol* **16**, 743-747 (1998).

18   Lansdorp, P. M. *et al.* Heterogeneity in telomere length of human chromosomes. *Hum
Mol Genet* **5**, 685-691 (1996).

19   Zijlmans, J. M. *et al.* Telomeres in the mouse have large inter-chromosomal variations in
the number of T2AG3 repeats. *Proc Natl Acad Sci U S A* **94**, 7423-7428 (1997).

20   Martens, U. M. *et al.* Short telomeres on human chromosome 17p. *Nat Genet* **18**, 76-80
(1998).

21   Londono-Vallejo, J. A., DerSarkissian, H., Cazes, L. & Thomas, G. Differences in
telomere length between homologous chromosomes in humans. *Nucleic Acids Res* **29**,
3164-3171 (2001).

22   Sholes, S. L. *et al.* Chromosome-specific telomere lengths and the minimal functional
telomere revealed by nanopore sequencing. *Genome Res* **32**, 616-628 (2022).

23   Grigorev, K. *et al.* Haplotype diversity and sequence heterogeneity of human telomeres.
*Genome Res* (2021).

24   Tham, C. Y. *et al.* High-throughput telomere length measurement at nucleotide
resolution using the PacBio high fidelity sequencing platform. *Nat Commun* **14**, 281
(2023).

25   Dubocanin, D. & Stergachis, A. B. Single-molecule architecture and heterogeneity of
human telomeric DNA and chromatin. *BioRxiv* (2022).

26   Tan, K. T., Slevin, M. K., Meyerson, M. & Li, H. Identifying and correcting repeat-calling
errors in nanopore sequencing of telomeres. *Genome Biol* **23**, 180 (2022).

27   Harley, C. B., Futcher, A. B. & Greider, C. W. Telomeres shorten during ageing of
human fibroblasts. *Nature* **345**, 458-460 (1990).

28   Hastie, N. D. *et al.* Telomere reduction in human colorectal carcinoma and with ageing.
*Nature* **346**, 866-868 (1990).

29   Vaziri, H. *et al.* Loss of telomeric DNA during aging of normal and trisomy 21 human
lymphocytes. *Am. J. Hum. Genet* **52**, 661-667 (1993).

30   Aubert, G., Baerlocher, G. M., Vulto, I., Poon, S. S. & Lansdorp, P. M. Collapse of
telomere homeostasis in hematopoietic cells caused by heterozygous mutations in
telomerase genes. *PLoS Genet* **8**, e1002696 (2012).

31   Rufer, N. *et al.* Telomere fluorescence measurements in granulocytes and T lymphocyte
subsets point to a high turnover of hematopoietic stem cells and memory T cells in early
childhood. *J Exp Med* **190**, 157-167 (1999).

32   Martens, U. M. *et al.* Telomere maintenance in human B lymphocytes. *Br J Haematol*
**119**, 810-818 (2002).

33   Van Ziffle, J. A., Baerlocher, G. M. & Lansdorp, P. M. Telomere length in subpopulations
of human hematopoietic cells. *Stem Cells* **21**, 654-660 (2003).

34   Baerlocher, G. M. & Lansdorp, P. M. Telomere length measurements in leukocyte
subsets by automated multicolor flow-FISH. *Cytometry A* **55**, 1-6 (2003).

35   Alder, J. K. & Armanios, M. Telomere-mediated lung disease. *Physiol Rev* **102**, 1703-
1720 (2022).

36   Lakota, K. *et al.* Short lymphocyte, but not granulocyte, telomere length in a subset of
patients with systemic sclerosis. *Ann Rheum Dis* **78**, 1142-1144 (2019).

37   Garg, S. *et al.* Chromosome-scale, haplotype-resolved assembly of human genomes.
*Nat Biotechnol* **39**, 309-312 (2021).

38   Stong, N. *et al.* Subtelomeric CTCF and cohesin binding site organization using
improved subtelomere assemblies and a novel annotation pipeline. *Genome Res* **24**,
1039-1050 (2014).

39   Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094-
3100 (2018).

40   Li, H. New strategies to improve minimap2 alignment accuracy. *Bioinformatics* **37**, 4572-
4574 (2021).

41   Liao, W. W. *et al.* A draft human pangenome reference. *Nature* **617**, 312-324 (2023).

42   Antonarakis, S. E. Short arms of human acrocentric chromosomes and the completion of
the human genome sequence. *Genome Res* **32**, 599-607 (2022).

43   Graakjaer, J. *et al.* The relative lengths of individual telomeres are defined in the zygote
and strictly maintained during life. *Aging Cell* **3**, 97-102 (2004).

44   Hemann, M. T., Strong, M. A., Hao, L.-Y. & Greider, C. W. The Shortest Telomere, Not
Average Telomere Length, Is Critical for Cell Viability and Chromosome Stability. *Cell*
**107**, 67-77 (2001).

45  d'Adda di Fagagna, F. *et al.* A DNA damage checkpoint response in telomere-initiated
senescence. *Nature* **426**, 194-198 (2003).

46  Enomoto, S., Glowczewski, L. & Berman, J. MEC3, MEC1, and DDC2 are essential
components of a telomere checkpoint pathway required for cell cycle arrest during
senescence in Saccharomyces cerevisiae. *Mol Biol Cell* **13**, 2626-2638 (2002).

47  IJpma, A. & Greider, C. W. Short telomeres induce a DNA damage response in
Saccharomyces cerevisiae. *Mol Biol Cell* **14**, 987-1001 (2003).

48  McNally, E. J., Luncsford, P. J. & Armanios, M. Long telomeres and cancer risk: the
price of cellular immortality. *J Clin Invest* **130**, 3474-3481 (2019).

49  Smogorzewska, A. & de Lange, T. Regulation of telomerase by telomeric proteins. *Annu
Rev Biochem* **73**, 177-208 (2004).

50  Shore, D. & Bianchi, A. Telomere length regulation: coupling DNA end processing to
feedback regulation of telomerase. *Embo J* **28**, 2309-2322 (2009).

51  Lustig, A. J. Clues to catastrophic telomere loss in mammals from yeast telomere rapid
deletion. *Nat Rev Genet* **4**, 916-923 (2003).

52  Zimmermann, M., Kibe, T., Kabir, S. & de Lange, T. TRF1 negotiates TTAGGG repeat-
associated replication problems by recruiting the BLM helicase and the TPP1/POT1
repressor of ATR signaling. *Genes Dev* **28**, 2477-2491 (2014).

53  Paschini, M. *et al.* Spontaneous replication fork collapse regulates telomere length
homeostasis in wild type cells. *bioRxiv* (2020).

54  Arneric, M. & Lingner, J. Tel1 kinase and subtelomere-bound Tbf1 mediate preferential
elongation of short telomeres by telomerase in yeast. *EMBO reports* **8**, 1080-1085
(2007).

55  Brown, W. R. A. *et al.* Structure and polymorphism of human telomere-associated DNA.
*Cell* **63**, 119-132 (1990).

56  Nergadze, S. G. *et al.* CpG-island promoters drive transcription of human telomeres.
*RNA* **15**, 2186-2194 (2009).

57  Deng, Z. *et al.* A role for CTCF and cohesin in subtelomere chromatin organization,
TERRA transcription, and telomere end protection. *Embo J* **31**, 4165-4178 (2012).

58  Feretzaki, M., Renck Nunes, P. & Lingner, J. Expression and differential regulation of
human TERRA at several chromosome ends. *RNA* **25**, 1470-1480 (2019).

59  Achrem, M., Szucko, I. & Kalinka, A. The epigenetic regulation of centromeres and
telomeres in plants and animals. *Comp Cytogenet* **14**, 265-311 (2020).

60    Toubiana, S. & Selig, S. Human subtelomeric DNA methylation: regulation and roles in
telomere function. *Curr Opin Genet Dev* **60**, 9-16 (2020).

61    Gonzalo, S. *et al.* DNA methyltransferases control telomere length and telomere
recombination in mammalian cells. *Nat Cell Biol* **8**, 416-424 (2006).

62    Arnoult, N. *et al.* Replication timing of human telomeres is chromosome arm-specific,
influenced by subtelomeric structures and connected to nuclear localization. *PLoS Genet*
**6**, e1000920 (2010).

63    Armanios, M. & Blackburn, E. H. The telomere syndromes. *Nat Rev Genet* **13**, 693-704
(2012).

64    Hao, L. Y. *et al.* Short telomeres, even in the presence of telomerase, limit tissue
renewal capacity. *Cell* **123**, 1121-1131 (2005).

65    Slagboom, P. E., Droog, S. & Boomsma, D. I. Genetic determination of telomere size in
humans: a twin study of three age groups. *Am J Hum Genet* **55**, 876-882 (1994).

66    Conklin, Q. A., Crosswell, A. D., Saron, C. D. & Epel, E. S. Meditation, stress processes,
and telomere biology. *Curr Opin Psychol* **28**, 92-101 (2019).

67    Greider, C. W. Telomeres, telomerase and senescence. *Bioessays* **12**, 363-369 (1990).

68    de Lange, T. *et al.* Structure and variability of human chromosome ends. *Mol. Cell. Biol.*
**10**, 518-527 (1990).

792
793

**Fig.1 | Nanopore telomere profiling is accurate and precise. a** Schematic depicting nanopore telomere profiling enrichment strategy. Telomeres are tagged with a biotin adapter (TeloTag), enriched by streptavidin pull down, and sequenced. **b** Subtelomere boundary is identified using an algorithm that detects significant deviation from the telomere repeat pattern. **c.** Southern blot of telomere lengths from 6 individuals. **d.** Telomere length measured by Nanopore telomere profiling for the same individuals as in **c**. The dashed line represents the mean telomere length of each distribution. Each point represents a single telomere read. **e** Telomere length measurements from a single donor measured 14 times on a single flow cell. **f** Telomere length measurements from a single donor measured across 6 flow cells **g.** Telomere length profiles from the same samples generated in two different laboratories JH=Johns Hopkins (blue), SC=UC Santa Cruz (gold) showing reproducible length profiles for 7 individuals. The difference in telomere length in base pairs is shown at the top.

**Fig. 2 | Nanopore telomere profiling measures telomere length dynamics in the aging population and separates short telomere patients from healthy individuals. a.** Nanopore telomere length profiles of 11 samples ranked in decreasing order. Age of individual noted at bottom. **b.** Southern blot of 11 samples loaded according to the order calculated in **a**, each lane is a different person. Each dot is a read. **c.** The mean telomere length was determined for 132 individuals aged 0 to 90 (blue dots). Cord blood lengths are shown in purple. The population distribution shows the 90th percentile (blue), the median (green line) and 10th percentiles (red) for telomere length in this population. Intervals were derived using parameters established previously for FlowFISH [15]. **d.** The 50th 10th and 1st percentile of telomere lengths for the data in **a**. are plotted with age on the X axis as in **b**. The dotted line represents the mean length of the 1st percentile for all samples **e.** Lymphocyte telomere length from FlowFISH data from Alder et al. (gray dots) and cord blood (purple dots). The lengths for 15 short telomere syndrome patients with IPF (red) were determined by FlowFISH. **f.** Nanopore telomere length profiles from the same 15 patients with short telomere syndrome shown in **e.** plotted against population distribution from **c**. One point represents two individuals aged 50 have nearly identical length and are indistinguish-able in the figure.

**Fig. 3 | Chromosome end-specific telomere lengths a.** Violin plots of the distribution of telomere lengths for 77 telomeres from HG002 that mapped with confidence and passed our filters (see methods). Each end is labeled with the chromosome number and p for the short and q for long arms. The haplotypes for each chromosome end are labeled Maternal (M) and Paternal (P). The mean, 90th and 10th percentile for each distribution are shown with short horizontal black lines in each violin plot. The grand mean of all telomeres is at the far right (all). The dashed line represents the grand mean of all telomeres **b**. Analysis of the means (ANOM) multiple contrast test of each telomere length distribution against the grand mean of all telomere lengths for data in **a.** The number of reads for each chromosome end is shown at the top. P-values were adjusted for multiple hypotheses testing using the Bonferroni method. Chromosome ends with length profiles reaching outside of the shaded gray region between the upper decision limit (UDL) and lower decision limit (LDL) are considered significantly different from the grand mean. (*) p ≤ 0.05, (**), p ≤ 0.01. (***), p ≤ 0.001; nonsignificant differences have no stars.

**Fig. 4 | Chromosome-specific telomere lengths are conserved and established at birth.** We used the pangenome reference to assign chromosome end status for (~720,000) telomere reads obtained from 150 individuals. Telomere reads were mapped to the pangenome with a requirement of 2kb of alignment. **a.** Matrix heatmap showing pairwise comparison of chromosome assignments of telomere reads when mapped to the pangenome or to the CHM13 reference genome. Heatmap shows what fraction of reads mapping to a given chromosome end in the pangenome (column) map to each chromosome end in CHM13 (rows). Light yellow indicates 0% and dark red indicates 100% of reads mapping to the respective CHM13 chr. end. **b.** As in **a**. but reads mapped to the HG002 paternal reference **c.** As in **a**. but reads mapped to the HG002 maternal reference genome. **d.** Bar graph showing the fraction of reads that mapped for each Chr end in the pangenome to the same Chr end in all three haploid genomes (CHM13, HG002 maternal and HG002 paternal). Colors are the same as in the heatmaps **a**. **b**. and **c**. **e**. For any given individual, we calculated Relative telomere length. For each chromosome end in each individual, we calculated the mean telomere length and subtracted it from the individual's grand mean telomere length. Zero indicates no difference between the chromosome end's mean telomere length and the individual's grand mean telomere length. Bars in the violin plots represent the mean length of all of the specific chromosome end in the population and the whiskers represent the standard error of the mean. **f**. Same as in **e**. but for cord blood samples only**.**

# Extended Data Figures

**a**

**Extended Data Fig. 1. Quantitation of enrichment and assay reproducibility. a.** Sequence of one representative TeloTag adapter. The barcoded adaptor (top strand) is annealed to a mixture of splints that have all 6 permutations of the CCCTAA sequence to improve chances of in-frame annealing to the telomere 3' overhang. **b.** Southern blot of telomeres recovered after biotin pull down using different volumes of streptavidin bead enrichment. **c.** Quantification of the efficiency of enrichment using increasing ratio of streptavidin beads to DNA. **d.** Enrichment of telomeric reads using biotin pull down relative to WGS. **e.** Intra-assay coefficient of variation (CV) of one single with different barcodes measured multiple times on the same flow cell. **f.** Inter-assay coefficient of variation (CV) of one sample measured multiple times across different flow cells. Mean telomere length of a single sample measured on multiple different runs.

**Extended Data Fig. 2: Heterogeneity in human subtelomere sequence means the telomere subtelomere boundary point can differ in sequence reads from diverse genomes and the reference genome a**. Telomere reads from the DNA identical to the reference genome will align at the boundary point in the reference. However, for some individuals a telomeric read will map well but there is extra sequence past the reference boundary point. For others there may be less subtelomere sequence on the read b. When telomere length is determined by mapping to the reference sequence boundary point, this can lead to incorrectly longer (II) or incorrectly shorter (III) telomere length distributions.

**Extended Data Fig. 3: Establishing telomere boundary points with TeloBP algorithm a**.
Representation of the nucleotide offsets for several different parameters as a rolling window
scanning from telomere end on right (see methods). **b.** IGV view of the telomere sequence and
where the boundary is called **c.** Example of where TeloBP incorporates variant repeats into the
telomere, compared to method setting a boundary of 4 consecutive repeats of TTAGGG.

**Extended Data Fig 4: Analysis of telomere length by TeloPeakCounter**. **a**. Representation of the subtelomere and telomere sequence electrical signal **b**. High resolution image of peaks in the telomere repeats in electrical signal. **Comparison of Guppy versus Dorado base caller**. Each blue dot represents an individual telomere read. 2435 read were examined from one data set (F63) from Fig 2. **c**. Comparison of telomere length determined by the peak counting vs Dorado base calling.  **d**. Comparison of telomere length determined by the peak counting vs Guppy base calling. **e**. Comparison of Guppy telomere length by TeloNP vs Dorado. **f.** Comparison of subtelomere length with Guppy vs Dorado.

**Extended Data Fig 5**. **Length of fragments does not affect telomere length determination.** **a.** The length of the fragments when genomic DNA is cut with AsiSI and PvuI is shown in blue. The length of the fragments when cut with BamHI and EcoRI is shown in orange. **b.** The telomere length of fragments cut with AsiSI and PvuI is in blue and BamHI and EcoRI is shown in orange. **c.** The distribution of fragment lengths for all reads: the Y axis is the number of reads and the X axis is the length in base pairs.

**Extended Data Fig. 6. Concordance of reads mapped to the pangenome with mapping to CHM13 and HG002 Mat and HG002 Pat.** We used different mapq scores to quantitate the fraction of reads that mapped to the same chromosome ends as the pangenome and the three referenced genomes **a.** Mapq score of 1 **b.** Mapq score of 30 **c.** Mapq score of 60. **d**. In previous analysis the acrocentric were omitted. Here they were included 13p,14p,15p, 21p and 22p and show less than 0 reads mapped to the same chromosome ends for these acrocentric.

**Extended Data Table 1. Samples included in comparison of FlowFISH and Telomere Profiling**

| Sample | age | FlowFISH lymphocyte length (bps) | FlowFISH granulocyte length (bps) | Telomere profiling length (bps) |
|---|---|---|---|---|
| 1 | 29 | 6210 | 5480 | 5045 |
| 2 | 45 | 3790 | 3860 | 4054 |
| 3 | 49 | 2790 | 3640 | 3678 |
| 4 | 51 | 3420 | 4020 | 3609 |
| 5 | 55 | 4520 | 4860 | 5545 |
| 6 | 56 | 3690 | 3990 | 3519 |
| 7 | 57 | 3670 | 2830 | 3759 |
| 8 | 59 | 3300 | 3600 | 2954 |
| 9 | 65 | 3580 | 4250 | 4945 |
| 10 | 66 | 3630 | 4430 | 3774 |
| 11 | 69 | 3670 | 3990 | 3541 |
| 12 | 69 | 3670 | 4300 | 3883 |
| 13 | 70 | 3480 | 4500 | 4203 |
| 14 | 72 | 3060 | 3090 | 3236 |
| 15 | 72 | 3530 | 4000 | 4003 |