**Stable chromosome configuration and loop-based polarization in animal symbionts**

Tobias Viehboeck[1,2,3§], Philipp M. Weber[1§], Nicole Krause[1,3], Nelle Varoquaux[4], Frédéric
Boccard[5], Ivan Junier[4], Silvia Bulgheresi[1,6]* and Virginia S. Lioy[5,6]*

[1] Department of Functional and Evolutionary Ecology, Environmental Cell Biology Group,
University of Vienna, Vienna, Djerassiplatz 1, A-1030 Vienna, Austria
[2] Division of Microbial Ecology, Center for Microbiology and Environmental Systems Science,
University of Vienna, Djerassiplatz 1, 1030 Vienna, Austria
[3] University of Vienna, Vienna Doctoral School of Ecology and Evolution
[4] Univ. Grenoble Alpes, CNRS, UMR 5525, VetAgro Sup, Grenoble INP, TIMC, 38000
Grenoble, France
[5] Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC),
91198 Gif-sur-Yvette, France
[6] Correspondence: silvia.bulgheresi@univie.ac.at, virginia.lioy@i2bc.paris-saclay.fr
§ These authors contributed equally and are ordered in alphabetical order
* These authors contributed equally and are ordered in alphabetical order

**ABSTRACT**
**Chromosome partitioning precedes the division of the cytoplasm, and its evolution is**
**linked with the positioning of the division plane. So far, bacterial chromosome biology**
**has heavily focused on transversally dividing, free-living ones. Here, we determined the**
**chromosome organization of three longitudinally dividing *Neisseriaceae* exclusively**
**inhabiting the oral cavity of mammals. We showed that in all three multicellular bacteria**
**the origin of DNA replication is invariably located at the host-attached (proximal) pole.**
**Next, 3C-seq revealed loop-based folding of the *ori* region in *Alysiella filiformis* and**
***Simonsiella muelleri*. Moreover, genes involved in cell motility, piliation and signal**
**transduction mechanisms were specifically looped when transcriptionally and**
**translationally active cells adhered to a substrate, but not when cultured in liquid.**
**Overall, we propose that proximal positioning of the *ori* and loop-based folding of its**
**surrounding DNA may mediate localized translation of proteins involved in host**
**colonization.**

**INTRODUCTION**

In rod-shaped bacteria, chromosomes have been shown to adopt two main spatial
configurations: left-*ori*-right (transverse) or *ori-ter* (longitudinal) [1,2]. In the transverse
configuration, the origin of DNA replication (*ori*) and the terminus of replication (*ter*) are at mid-

cell, so that the left and right arms reside in opposite cell halves. In the longitudinal

configuration, *ori* is located at one cell pole and *ter* at the opposite pole so that both left, and

right chromosomal arms are parallel to the long axis of the cell. While model rods such as

*Escherichia coli* and *Bacillus subtilis* can switch between transverse and longitudinal

arrangements depending on the growth conditions or the cell cycle stage [3–7], the chromosome

in *Caulobacter crescentus*, *Pseudomonas aeruginosa* and *Vibrio cholerae* maintains a

longitudinal configuration throughout generations [8–10]. In these three monoflagellated and

vertically polarized bacteria, after the replication of the *ori*, the ParABS system actively

segregates the duplicated *parS* regions towards the opposite pole [11,12]. A variation on the

theme of the longitudinal configuration is the so-called *ori-ter-ter-ori* configuration reported for

the diploid *Actinomycetales Corynebacterium glutamicum* [13]. In this configuration, the two *ori*

are tethered to the poles in non-dividing cells, but both new-born *ori* are segregated to mid-

cell, where they remain until fission is completed. Only then, they get tethered to the newly

forming cell poles. While in all the aforementioned bacteria the intracellular localization of the

*ori* can drastically vary during the cell cycle, in the longitudinally dividing *Candidatus*

Thiosymbion oneisti the *ori* is maintained in the central third of the cell during its replication

and lateral segregation, likely by the ParABS system [14]. Taken together, irrespective of how

the chromosome is configured, during replication and concomitant segregation, the subcellular

localization of at least one of the sister *ori* along the long axis changes substantially in all

bacteria studied so far [15], but the longitudinally dividing symbiont *Ca.* T. oneisti [14].

Besides displaying specific spatial dispositions and segregation modes, bacterial

chromosomes may differ in their 3D internal organization, as revealed by chromosome

conformation capture techniques coupled to deep sequencing (e.g., 3C-seq or Hi-C; [7,16–21]. In

all cases, Hi-C contact maps showed that chromosomes are segmented in sub-megabase

chromosomal interaction domains (CIDs) whose boundaries often contain long and highly

expressed gene clusters, indicating that intense transcription may limit the interactions

between adjacent chromatin domains [17,22]. Although CIDs are reminiscent of the eukaryotic

topologically associated domains (TADs), they differ in their formation mechanisms. In

mammalian cells, TADs are often delimited by chromosomal loops [23]. These loops are

observed at TADs boundaries as corner peaks and result from the structural maintenance of

chromosomes (SMC) cohesin complexes and the CCCTC-binding factor (CTCF) binding the

DNA at specific sites [24–26]. In bacteria, chromosomal loops have been described in *B. subtilis.*

Initially, some of these loops were proposed to be involved in the regulation of *ori* firing [7,27].

More recently, loops were shown to be involved in some CID establishment during stationary

phase via the action of the transcriptional factor Rok [28]. Specifically, Rok was shown to form

large nucleoprotein complexes that interact with each other over large distances, which

eventually leads to anchored chromosomal loops [28]. Finally, chromosomal loops have also

been detected in euryarchaeota [29] and involved in the aggregation of high expressed insulated

domains in the crenarchaeon *Aeropyrum pernix* [30]. In the former case they are SMC-

dependent but transcription independent, whereas in the latter, SMC-less organisms loops are

transcription sensitive [29,30].

Despite the importance of chromosome orientation, segregation and conformation for

cell functioning, only a limited number of bacteria have been studied so far [12,19,31]. Here, we

investigated the chromosomes of three multicellular *Neisseriaceae* commonly found attached

to the oral mucosa of warm-blooded vertebrates [32]: a canine strain of *Conchiformibius steedae*

[33,34], a caprine strain of *Alysiella filiformis* [35,36] and a human strain of *Simonsiella muelleri* [36,37]

(Fig. 1a-c). By applying DNA fluorescence in situ hybridization on fixed cells (FISH), we

showed that the chromosome configuration of these oral bacteria is stable, with the *ori*

invariably localized at the host-attached (proximal) pole throughout their cell cycle, likely

involving the centromeric protein ParB. Moreover, 3C-seq revealed the systematic presence

of CIDs in exponential phase, most of them being delimited by long and highly expressed

genes. Besides domains, we observed chromosomal loops in the host-proximal region of both

*A. filiformis* and in *S. muelleri*, what we refer to as chromatin polarization hereafter. In *S.*

*muelleri*, some loops delineated CIDs, similarly to mammal-like mechanisms associated with

the formation of TADs. Strikingly, the abundance and the position of chromosomal loops were

sensitive to the growing conditions (solid or liquid medium), as well as to transcription or

translation inhibitors. Namely, growth in liquid medium led, in *A. filiformis*, to the enlargement

of an *ori*-containing chromosomal loop and, in *S. muelleri*, to a ~10-fold decrease in the

number of chromosomal loops. Strikingly, in both oral bacteria, genes putatively involved in in

motility and piliation were specifically looped when cells adhered to agar dishes, suggesting

that loops might be associated with transertion phenomena involved in host colonization.

Altogether, we propose that stable chromosome configuration and loop-based folding of

chromosomes are cell biological mechanisms underlying host-bacterium interactions.

**RESULTS**

**Stable longitudinal chromosome configuration in three oral cavity symbionts.**

The three multicellular bacteria studied in this work are vertically polarized, as indicated by the

host-proximal localization of their fimbriae [32,37–39]. We will henceforth refer to the host-attached,

fimbriae-rich pole of *A. filiformis* as the proximal pole and to the free pole as the distal pole.

Regarding their division modes, *A. filiformis* divides in a distal-to-proximal fashion [32], whereas

in the crescent-shaped *S. muelleri* and *C. steedae*, both poles are attached to the mouth

epithelium which makes their mid-cell correspond to the region which is furthest away from

the animal surface [32,37]. Moreover, in *S. muelleri* and *C. steedae* the fimbriae localize to the

side of the filament facing the oral mucosa, also referred to as the proximal (or ventral) side
[32,37]. Concerning their division mode, *S. muelleri* and *C. steedae* both septate from their host-
attached poles toward mid-cell [32].

To determine their ploidy and chromosome configuration, we fixed exponentially
growing *C. steedae, A. filiformis* and *S. muelleri* and subjected them to DNA fluorescence in
situ hybridization (FISH) with specific probes specifically targeting their predicted *ori* and *ter*
(Supplementary Fig. 1).

When *C. steedae* was subjected to DNA FISH, almost 50% of the cells (44%; n = 215)
displayed two *ori* foci (Fig. 2a, Supplementary Table 1), and 76% (n = 236) displayed at least
two *ter* foci (Fig. 2d, Supplementary Table 1), with *ori* invariably localized proximally (Fig. 2b-
c), and the *ter* localized at mid-cell (Fig. 2e-f). Furthermore, lateral segregation of the *ori* was
discernable (Supplementary Fig. 2a-c). As observed for *C. steedae*, most *S. muelleri* cells
(47%; n = 188; Fig. 2g) had two *ori*, each at one pole (Fig. 2h-i). As for the *ter*, we detected
one or two foci at mid-cell in most of the cells (61% and 39%, respectively; n = 293; Fig. 2j-l).

Multifork replication, as observed in fast-growing bacteria [40], could explain the relatively
high *ori* count observed in *C. steedae* and *S. muelleri*. To exclude multifork replication in
exponentially growing *C. steedae* and *S. muelleri*, we subjected them to marker frequency
analysis (MFA), together with *A. filiformis* and *Escherichia coli*. We found that the *ori-ter* ratios
of all three multicellular *Neisseriaceae* did not significantly change between exponentially
growing and stationary phase cells, with values in the 1.3-1.9 range (Supplementary Fig. 3a-
f). In *E. coli*, instead, the *ori-ter* ratio changed from 2.5 to 0.97 between exponentially growing
and stationary phase cells (Supplementary Fig. 3g-h), as previously observed [41]. This
indicates that, contrary to *E. coli*, oral cavity symbionts do not undergo multifork replication
and that *C. steedae* and *S. muelleri* are likely diploid.

Applied to *A. filiformis*, DNA FISH revealed that 98% of the cells (n = 379; Fig.3 a) have
one *ori* and 96% (n=183; Fig. 3d) have one *ter*, indicating that this bacterium is likely
monoploid. As for the configuration of the *A. filiformis* chromosome, we observed that,
independently of the cell division stage, *ori* localized proximally (Fig. 3b-c and Supplementary
Fig. 2d-e), whereas *ter* was distal (Fig. 3e-f and Supplementary Fig. 2f-g). Moreover,
localization pattern analysis of *A. filiformis* containing either one or two *ori* foci (Supplementary
Fig.2d-e) or either one or two *ter* foci (Supplementary Fig. 2f-g) indicated lateral chromosome
segregation. Finally, after identification of the *A. filiformis* ParB protein using phylogenetic
analysis (Supplementary Fig. 4a-b), we generated specific anti-ParB antibodies and used
them for immunostaining (Supplementary Fig. 4c). The results were consistent with this
centromeric protein mediating longitudinal chromosome configuration and segregation (Fig. 3
g-i).

Collectively, our results show that multicellular *Neisseriaceae*, irrespective of their
ploidy and irrespective of their cell cycle stage, localize their *ori* proximally and their *ter* distally
resulting in stable *ori-ter-ter-ori* (*C. steedae* and *S. muelleri,* Fig. 2m) or *ori-ter* (*A. filiformis,*
Fig. 3j) configurations. Furthermore, based on its localization pattern, the centromeric protein
ParB might mediate proximal positioning of the *ori* in *A. filiformis* (Fig.3 j). Finally, the finding
of stable chromosome configurations in bacterial symbionts living attached to marine
nematodes [14] (Supplementary Fig. 5) suggests that they might have convergently evolved to
serve the symbiotic lifestyle.
156
**Starkly different chromosome architectures in three evolutionary related *Neisseriaceae***
Next, we used 3C-seq and RNA-seq to study the folding properties of the
chromosomes of *C. steedae*, *A. filiformis*, and *S. muelleri* grown on solid medium and their
putative relationships with gene expression. Despite the phylogenetic relatedness of these
oral bacteria [32], genome-wide contact maps at a 5 kb-bin resolution revealed starkly different
chromosome architectures (Fig. 4a-d). Firstly, only the maps of *C. steedae* and *S. muelleri*
displayed a secondary diagonal, indicating the presence of low frequency, but significant,
inter-arm DNA contacts (Supplementary Fig. 6a-b). This was consistent with the longitudinal
organization of their chromosome (see previous section). By contrast, the contact map of *A.*
*filiformis*, revealed rare inter-arm contacts despite the apparent longitudinally configuration of
its chromosome and the presence of the condensin complex Smc-ScpAB and the ParABS
system (Fig.4b; Supplementary Fig. 6c).
169
Along the main diagonal, the 3C contact matrices revealed the presence of nested
domains. To detect the chromosomal loci implicated in domains that span a wide range of
sizes, we used the Frontier Index (FI), a tool that has been developed specifically to deal with
the presence of multiple scales in the domain organization of 3C contact matrices [42].
Specifically, the FI detects bins associated with either a significant upwards or a significant
downwards interaction bias at any scale (up- or downstream peaks, respectively, Figure 4). In
this context, we will henceforth refer to (1) loci corresponding to an up- or to a downstream
peak as frontiers, (2) loci where an upstream peak is followed by a downstream peak (±3 bins)
as domain boundaries and (3) loci between two domain boundaries as CIDs. In total, we
identified 18, 16 and 37 frontiers in *C. steedae*, *A. filiformis*, and *S. muelleri*, respectively, with
<50% of them corresponded to domain boundaries (Table 1, Fig.4a-d; Supplementary Fig. 7a,
Supplementary Fig. 8a-b and Supplementary Fig. 9a-b). In the case of substrate-attached *C.*
*steedae* and *A. filiformis*, 80% (4 out of 5) and 100% (5 out of 5) of the domain boundaries
matched long and/or highly expressed genes (HEGs, including *rrna* operons; Table 1;
Supplementary Fig. 10a-b), consistent with previously observations in other bacteria [7,13,16–

[18,22,43]. Notably, piliation genes (*pilMNOPQ*) located in the domain boundary situated at 1.7 Mb

appeared strongly insulated from the rest of the chromosome (Fig. 4a). As for *S. muelleri*, the

correlation between domain boundaries and HEGs was 45% in replicate 1 (only five out of the

11 boundaries matched HEGs), but was 69% in replicate 2 (Table 1, Supplementary Fig.10a-

b).

190           Collectively, we unveiled the large-scale organization of the chromosomes of three

multicellular *Neisseriaceae*. Our results showed the absence of frequent inter-arm contacts in

*A. filiformis* despite a longitudinal organization similar to *C. steedae and S. muelleri.* We also

unveiled the presence of chromatin constrains that define multiple frontiers which are not

linked to transcriptional activity in any of the studied symbionts. Finally, we have uncovered

the existence of CIDs whose boundaries are not associated to HEGs, suggesting that intense

transcription of long transcriptional units (or of a succession of them) is not the sole

mechanism of chromatin organization in *S. muelleri*.

**Loop-based polarization of the chromosomes in substrate-attached *A. filiformis* and *S.***

***muelleri***

201           Apart from domains, the contact maps of *A. filiformis* and *S. muelleri* revealed peaks

of genetic interaction that we ascribed to chromosomal loops (indicated as black circles, Fig.

4c,e,f and Table 2). To identify their number unambiguously, we developed and applied a

loop-detection algorithm to two contact maps established independently for each oral

bacterium (replicate 1 and replicate 2). In the case of *A. filiformis*, we detected one 140 kb-

long loop on the right side of the *ori* (Fig. 4b,e, Fig. 6a and Tables 1 and 2). Although this loop

did not match any domain boundary (Tables 1 and 2), it resulted from the interaction of loci

involved in host colonization (see last section of the Results).

209           In the case of *S. muelleri*, our loop-detection algorithm revealed 36 distinct loops (Fig.

4c-d, black circles) engaging 25 different chromosomal regions (referred to as looping regions

hereafter), with each looping region being connected to 2.4 other looping regions on average.

These loops were not distributed homogenously throughout the chromosome. Instead, they

generated a highly condensed region around *ori* (between positions 0.8 and 1.4 Mb), a second

one on its right side (between positions 0.3 and 0.7 Mb) and a third one on its left side (between

1.7 and 2.1 Mb) (Fig. 4c-d,f, Fig.6b), leaving a ≈1.4 Mb loopless region around *dif*. Of note,

50% of the loops matched domain boundaries (Fig. 4c-d, f and Tables 1 and 2).

217           Altogether, contact maps of the human symbiont *S. muelleri* unveiled a loop-based

organization of its chromatin resembling that seen in mammalian cells. Furthermore, the

identified loop in *A. filiformis* close to *ori* together with the strong bias of *S. muelleri* loops for

the proximal (*ori*-containing) region imply that the chromosomes of these bacteria are not

homogeneously folded along the cell long axis. We will hereafter refer to this phenomenon as
loop-based polarization of the *A. filiformis* and *S. muelleri* chromosomes.

**Loss of substrate attachment affects loop-based chromosome folding**
To assess whether losing the attachment to a substrate would result in a depolarization
of chromosome folding (i.e., in a loss of chromosomal loops), we performed 3C-seq in *A.*
*filiformis* and *S. muelleri* grown in liquid culture and fixed either in exponential or stationary
phase (Fig. 5b-f, Supplementary Fig. 8c-e, Supplementary Fig. 9c-e). Furthermore, we also
performed 3C-seq on liquid-grown *C. steedae* (Fig. 5a) to know whether changes in growing
conditions or growth phase would affect the number of frontiers.
Overall, the global 3D organization of the chromosomes of liquid grown *C. steedae*, *A.*
*filiformis* and *S. muelleri* was conserved, e.g., we observed a secondary diagonal in *C. steedae*
and *S. muelleri*, and rare inter-arm contacts in *A. filiformis.* In addition*,* FI detected a similar
number of frontiers in the three symbionts (Fig. 5a-c, Supplementary Fig. 7b, Supplementary
Fig. 8c-d and Supplementary Fig. 9c-d), albeit exponential phase *C. steedae* and liquid-grown
*A. filiformis* (irrespective of the growth phase), displayed more domain boundaries compared
to bacteria grown on agar dishes (Table 1). Also of note, more domain boundaries
corresponded to HEGs in liquid-grown *C. steedae*.
Most strikingly, in liquid growing *S. muelleri* the number of chromosomal loops
decreased from 36 to seven irrespective of the growth phase (Fig. 5c, e-f, Supplementary
Fig.9c-e and Table 2). Moreover, only one of the seven loops detected in liquid-grown *S.*
*muelleri* matched a frontier (Table 1). Concerning *A. filiformis*, the 140 kb-long *ori*-containing
loop detected in solid-grown cells (Fig. 4b,e and Fig. 6a top panel) became 215 kb-long in
cells grown in liquid culture, irrespective of the growth phase (Fig. 5b,d, Fig. 6a bottom panel
and Supplementary Fig. 8c-e).
All in all, loss of substrate attachment mainly affected the loop-based organization of
the chromatin in *A. filiformis* and *S. muelleri*. Namely, loss of substrate attachment led to the
formation of a longer, *ori*-containing chromosomal loop in *A. filiformis* and to an 80% loss of
chromosomal loops in *S. muelleri*, with only seven left in the immediate vicinity of its *ori*.

**Loop-based polarization of the chromosomes in substrate-attached symbionts may**
**facilitate localized translation of proteins involved in host colonization.**
To assess the function of chromosomal loops, we first tested the transcriptional levels of the
genes they involved in substrate-attached *A. filiformis* and *S. muelleri* (Supplementary Fig.10c;
Supplementary Data Set 1). In *A. filiformis*, we did not find any significant difference in the
median level of transcription of looped genes versus the median level of transcription of all the
genes. In substrate-attached *S. muelleri*, however, transcripts of looped genes were scarce

(p<0.01), except for those encoding for translation, ribosomal structure and ribosomal
biogenesis proteins (p<0.05) (Supplementary Data Set 1). Of note, growth of *S. muelleri* in
liquid medium did not increase the abundance of transcripts corresponding to looped genes
suggesting that looping was not responsible of the low abundance of their transcripts in
substrate-attached cells.

In our quest to understand the biological function of chromosomal loops, we then
resorted to automatic annotation of the looping regions (based on the eggNOG database). In
substrate-attached *A. filiformis*, the two looping regions contained genes putatively needed for
colonization of the host, including a signal recognition particle (SRP)-containing,15 kb-long
adhesin and a transcription repressor involved in efflux mediated antimicrobial resistance (Fig.
6a top, Supplementary Tables 5 and 6 and Supplementary Data Set 1); [44]. However, when *A.*
*filiformis* was grown in liquid medium, none of the looped genes belonged to the COG category
*Host adhesion and/or immune evasion*. Instead, looped genes belonged to either *Information*
*storage and processing* or to *Metabolism* (Fig. 6a bottom, Supplementary Fig. 11a and
Supplementary Data Set 1).

We then annotated the genes situated in the looping regions in *S. muelleri* growing on
solid medium using the eggNOG database (Fig. 6b, Supplementary Fig. 11a and
Supplementary Data Set 1). Firstly, we found the following solid medium-specific COG
categories: *Cell motility* (including *pilA* and a gene encoding for a prepilin-related protein),
*Defense mechanisms* (including the antibiotic resistance gene *ampD*) and *Signal transduction*
*mechanisms*. Of note, 40% of cell motility genes were looped when *S. mulleri* was grown on
agar plates. Concerning the "second most-looped" COG category, *Intracellular*
*trafficking/Secretion*, although this category is not solid substrate-specific, the Type 4
Secretion System genes virB3-6 and the gene encoding for the SRP receptor *ftsY* were only
looped when cells were substrate-attached. Also noteworthy were the COG categories *Cell*
*wall/membrane/envelope biogenesis* and *Coenzyme transport and metabolism*. Although
these two categories were not solid medium-specific, *wecA* and *neuABC* genes (encoding for
capsule or lipopolysaccharide sialyating enzymes), two spermidine synthases and an adhesin
were only looped in substrate-attached *S. muelleri* (Fig. 6b, Supplementary Fig. 11a and
Supplementary Data Set 1). Finally, among the genes which could not be classified into any
COG category, but that were only looped in substrate-attached *S. muelleri* and that may be
involved in host adhesion and/or immune evasion, we found three genes encoding for
immunoglobulin-binding proteins (*tspB*) and three copies of a gene encoding for zonula
occludens toxins (*zot*) (Fig. 6b, Supplementary Fig. 11a and Supplementary Data Set 1).

In summary, functional annotation of highly frequently interacting (looped) genes
suggests that, at least some of the loops may result from transertion, i.e., from DNA-
membrane interaction due to the coupled transcription and translation of membrane or

secreted proteins associated with host colonization. Consistent with this hypothesis, blocking
transcription or translation in substrate-attached *A. filiformis* had the same effect as growing
them in liquid culture (Table 1, Fig.6a, Supplementary Fig.12). Along the same line, blocking
translation in substrate-attached *S. muelleri* led to the loss of loops engaging genes encoding
for cell motility proteins, pili, the SRP receptor and hemolysin (Table 1, Fig. 6b, Supplementary
Fig.11 and Supplementary Fig.12).
301
302   **DISCUSSION**
303           During the past decade, studies on symbiont cell biology revealed that rod-shaped
bacteria may specifically localize their fimbriae to their host-attached poles [39]. Moreover,
bacterial symbionts orient their cytokinetic machineries orthogonal to the animal surface and
septate longitudinally, so that both daughter cells stay attached to the host by one pole [39,45,46].
Recently, we found that, thanks to a bidimensional segregation mode, a longitudinally dividing
symbiont could keep its chromosome orientation toward its host throughout generations [14].
Here, we expanded on the capacity of symbiotic cells to polarize their components by focusing
on the spatial disposition and architecture of their genetic material.
311           Firstly, we found that all the symbionts studied here configure their chromosomes
longitudinally throughout the cell cycle, i.e., the *ori* stays proximal irrespective of the ploidy
and of chromosome segregation stage. This is achieved by lateral chromosome segregation,
likely active and ParABS-mediated. Indeed, in *A. filiformis*, we identified seven *ori*-adjoining
*parS* sites, and ParB signal was proximal throughout the cell cycle. Similarly, in the rod-shaped
and nematode symbiont *Candidatus* Thiosymbion hypermnestrae, ParB foci are proximal at
all cell cycle stages, but the very last one (Supplementary Fig.5). As for the nematode
symbiont *parS* sites, we identified two of them in the vicinity of the *ori*. Therefore, in *A. filiformis*
and *Ca.* T. hypermnestrae, ParA might drive lateral segregation of *ori*-ParB. As for *C. steedae*
and *S. muelleri*, the former has two *parS* sites in the vicinity of the *ori* and four in the vicinity
of the *ter*, whereas the latter has six *ori*- and two *ter*-adjoining *parS* sites. Further, in *C. steedae*
and *S. muelleri*, 3C contact maps revealed the presence of inter-arm contacts (Supplementary
Fig.6), suggesting that ParB mediates the loading of Smc-ScpAB at the *parS* sites situated in
the vicinity of the *ori*. Interestingly, *B. subtilis* harbors eight *parS* sites in the *ori* region and
each of them presents a specific ParB and Smc-ScpAB recruitment efficiency [47].
Characterization of the affinity of ParB for the different *parS* sites is necessary to determine
whether ParB has a higher affinity for *parS* sites situated in the vicinity of the *ori* than for the
distal ones, and whether this may play a role in differential recruitment of the Smc-ScpAB
complex in *S. muelleri* and *C. steedae.*
330           Secondly, DNA FISH-based count of *ori* and *ter* foci and MFA indicate that *S. muelleri*
331   and *C. steedae* are likely diploid, i.e., newborn cells have two *ori* and two *ter* and therefore

two copies of the same chromosome. Although about one third of *S. muelleri* or *C. steedae* displayed only one *ori* focus (instead of two), we ascribe this to an insufficient detection efficiency of the DNA FISH technique. Likely for the same reason, most *S. muelleri* cells displayed only one *ter* focus instead of two foci. Diploidy has also been reported in pathogenic *Neisseriaceae* [48], in the actinomycetes *C. glutamicum* [13] and in the Euryarchaeon *Methanothermobacter thermautotrophicus* [49]. Because bearing two (or more) chromosome copies increases the chance of correctly repairing the DNA by homologous recombination, we hypothesize that *C. steedae* and *S. muelleri* evolved diploidy to withstand stress [50] including that exerted by bacteriophages or transposable elements [51]. In this regard, the identification of prophage gene clusters and three transposase-encoding genes is noteworthy (Supplementary Fig. 9b). Alternatively, or additionally, diploidy may have evolved to maintain the intracellular position of genetic loci in longitudinally dividing curved bacteria, which adhere to the animal surface with both poles.

A third point is the spatial disposition of the genetic material inside the cell. Up to this work, all studied bacterial chromosomes displayed highly variable chromosome configurations: either *ori*, *ter* or both drastically changed their intracellular localization pattern during DNA replication and segregation (or when experiencing different growth phases) [1,31], except for the longitudinally dividing *Ca.* T. oneisti, where the *ori* is kept at mid-cell throughout generations and irrespective of the cell cycle stage [14]. We currently do not know what pushed the oral symbionts to evolve formidably stable chromosome configurations. One evolutionary driver could have been the necessity to keep the genes where their products are needed, especially if these products are essential for the symbiont survival. Apart from possessing a genetic repertoire typical of aerobic chemoorganoheterotrophs, our annotations indicate that the oral cavity symbionts cannot synthesize vitamin $B_{12}$ and, intriguingly, two genes encoding for transporters mediating its uptake (the TonB-dependent-vitamin $B_{12}$ transporter BtuB and a vitamin $B_1$ transporter) are situated close to the *ori* in *S. muelleri* and are therefore, thanks to the stable chromosome configuration, invariably proximal (Supplementary Data Set 1). The hypothesis that stable chromosome configurations evolved to facilitate the intracellular localization of membrane or secreted proteins where they are most needed relies on the assumption that mRNAs may be kept where they are transcribed, as shown for *C. crescentus* [52]. An additional piece of data linking the intracellular localization of bacterial genes to the place they have to function is that translation of membrane proteins may trigger chromosomal locus repositioning [53,54]. Intriguingly, incubation in 25 µg/ml chloramphenicol led to nucleoid compaction in both *A. filiformis* and *S. muelleri* (data not shown). However, localization pattern analysis of selected *A. filiformis* and *S. muelleri* mRNAs is needed to find out whether its chromosome serves as a template for the proximal localization of mRNAs involved, for

example, in the uptake of essential nutrients from the host (e.g., the mRNA encoding for the
vitamin transporter Btub).

Finally, in the case of *C. steedae* and *S. muelleri* (adhering to a solid substrate or
planktonic) a secondary diagonal was visible in the contact maps consistent with the stable
longitudinal organization revealed by DNA FISH. However, a secondary diagonal does not
seem to be a prerequisite of chromosome longitudinal organization, as we detected rare inter-
arm contacts in *A. filiformis* which also has an *ori-ter* configured chromosome. This suggests
that in this monoploid symbiont the two chromosome arms are laying longitudinally, but further
away from each other than in the thinner, longer and apparently diploid *C. steedae* and *S.*
*muelleri*. Alternatively, the lack of inter-arm contacts in *A. filiformis* might be due to the absence
of an SMC-based, *ori-ter* oriented loop extrusion mechanism [55].

Prior to this study, intensively transcribed loci were found in at least 60% of the CID
boundaries [7,13,17–19,22,43]. The correlation between domain boundaries and HEGs is consistent
with the previous findings except for stationary phase *A. filiformis* and *S. muelleri* (Table 1),
possibly because the number of frontiers was flawed by the intrinsic noise associated to 3C
matrices (Supplementary Figs. 7-9). Furthermore, in these symbionts, we observed biased
interactions in a single direction around certain loci, meaning that these loci do not separate
two consecutive CIDs. Interestingly, these frontiers are not associated with HEGs, but with
chromatin constrains that remain to be identified.

Concerning chromosomal loops, in substrate-attached *S. muelleri* that were actively
transcribing and translating, they were frequently found at domain boundaries (Tables 1 and
2), resembling the corner peaks demarcating mammalian TADs. Their association with TADs
can be explained by the following model: the cohesin complex binds the chromatin at a random
place, extrudes DNA bi-directionally through a loop and is eventually blocked by the CCCTC-
binding factor CTCF bound at the borders of the TAD [56]. The blocking indeed promotes the
stabilization of the loop, which thus manifests as a corner peak in the contact map [26,57]. In *B.*
*subtilis*, the transcriptional factor Rok, likely assisted by the SMC complex, is associated to
chromosomal loop formation delineating CIDs in stationary phase [28]. Whether a similar system
is responsible for chromosomal looping and domain formation in substrate-attached *S.*
*muelleri* remains to be demonstrate.

Functional annotation suggested that, at least some *S. muelleri* chromosomal loops,
may result from localized transcription and translation of proteins involved in cellular motility,
piliation and immune evasion. Localized transcription and translation of host colonization
factors, facilitated by stable chromosome configurations, would allow, when cells are attached
to the oral mucosa, to efficiently translate and assemble (or transert) membrane-bound
molecular complexes required for host colonization. Consistent with this hypothesis, in *A.*
*filiformis*, an adhesin gene and a gene involved in efflux-mediated antibiotic resistance were

looped only when actively transcribing and translating cells adhered to a substrate, suggesting that, like in *S. muelleri*, chromosomal loops may be involved in host adhesion and/or immune evasion. (Fig.6b, Supplementary Tables 5 and 6 and Supplementary Data Set 1). Along this line, transertion of the type 3 secretion system was recently shown in *Vibrio parahaemolyticus* incubated with bile acids [54].

Given that 3C-seq did not reveal any chromosomal loop in *C. steedae* (Fig. 4a, Fig. 5a and Supplementary Fig. 7), irrespective of the culture conditions, chromosomal loops do not seem a *sine qua non* feature of bacteria thriving on animal surfaces. Instead, loop-based chromosome polarization might be an exclusive feature of multicellular *Neisseriaceae* belonging to the M1 lineage (*Alysiella* and *Simonsiella* spp.) [32]. Although chromosome conformation capture did not reveal chromosomal loops in the unicellular *Neisseriaceae Neisseria elongata* (data not shown), we must study more *Neisseriaceae* species and isolates to test this hypothesis.

In conclusion, it is possible that stable chromosome configurations, together with the presence of chromosomal loops in *A. filiformis* and *S. muelleri*, mediate localized (proximal) translation of proteins involved in host colonization (e.g., pili, enzymes for extracellular polysialic acid synthesis, adhesins, zonula occludens toxins, immunoglobulin-binding proteins). More generally, we propose that specific chromosome positioning and folding may represent cellular adaptations to an obligate symbiotic lifestyle.

**AUTHOR CONTRIBUTIONS**

T.V. did 3C experiments, marker frequency analysis, RNA sequencing, genome sequencing and assembly, bioinformatic and formal analyses, visualization, wrote and revised the manuscript. P.M.W. did DNA staining, DNA FISH, formal analyses and visualization, wrote and revised the manuscript; N.K. did DNA staining, DNA FISH and formal analyses. I.J. performed computational analyses of 3C-seq data, developed the loop detection algorithm and revised the manuscript. N.V. performed preliminary FI analysis. B.D. did DNA FISH. F.B. acquired funding. V.S.L. performed preliminary 3C analysis, conceptualized and supervised the work, acquired funding, provided resources, wrote, and revised the manuscript. S.B. conceptualized and supervised the work, acquired funding, provided resources, wrote, and revised the manuscript.

**ACKNOWLEDGEMENTS**

This work was supported by the Austrian Science Fund (FWF) project P28593 (T.V. P.M.W, N.K.), FWF project P28743 (T.V., P.M.W), FWF doc.funds MAINTAIN (T.V.), a DOC-fellowship from the Austrian Academy of Science (P.M.W.), and a PhD completion grant from the University of Vienna (P.M.W.), by the French National Research Agency: grant number

ANR-20-CE35-005 (V.S.L), by the 80 Prime CNRS project MIMIC (F.B. and I.J.) and SIRIG (N.V and V.S.L.) through the MITI interdisciplinary exploratory research porgramms. We thank Bocar Diallo for performing some DNA FISH experiments, to Frédéric Veyrier (INRS, Quebec, CA) for introducing *Conchiformibius steedae* to us and for sharing the RNAseq protocol, the members of the F.B. and S.B. laboratories for fruitful discussions and to S. Bury-Moné for careful reading and feedback on the article. We acknowledge the sequencing and bioinformatics expertise of the I2BC High-throughput sequencing facility, supported by France Génomique (ANR-10-INBS-09). We are also indebted to the staff of the VBCF NGS Unit (Laura-Maria Bayer) for assistance with Oxford Nanopore MinION sequencing. The computational results of this work have been achieved using the Life Science Compute Cluster (LiSC) of the University of Vienna.

**DECLARATION OF INTEREST**

The authors declare no competing interests.

**REFERENCES**

1. Toro, E. & Shapiro, L. Bacterial Chromosome Organization and Segregation. *Cold Spring Harb. Perspect. Biol.* **2**, a000349 (2010).

2. Wang, X., Llopis, P. M. & Rudner, D. Z. Organization and segregation of bacterial chromosomes. *Nat. Rev. Genet. 2013 143* **14**, 191–203 (2013).

3. Niki, H., Yamaichi, Y. & Hiraga, S. Dynamic organization of chromosomal DNA in Escherichia coli. *Genes Dev.* **14**, 212–223 (2000).

4. Bates, D. & Kleckner, N. Chromosome and replisome dynamics in E. coli: Loss of sister cohesion triggers global chromosome movement and mediates chromosome segregation. *Cell* **121**, 899–911 (2005).

5. Wang, X., Llopis, P. M. & Rudner, D. Z. Bacillus subtilis chromosome organization oscillates between two distinct patterns. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 12877–12882 (2014).

6. Cass, J. A., Kuwada, N. J., Traxler, B. & Wiggins, P. A. Escherichia coli Chromosomal Loci Segregate from Midcell with Universal Dynamics. *Biophys. J.* **110**, 2597–2609 (2016).

7. Marbouty, M. *et al.* Condensin- and Replication-Mediated Bacterial Chromosome Folding and Origin Condensation Revealed by Hi-C and Super-resolution Imaging. *Mol. Cell* **59**, 588–602 (2015).

8. Viollier, P. H. *et al.* Rapid and sequential movement of individual chromosomal loci to specific subcellular locations during bacterial DNA replication. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 9257–9262 (2004).

9.    Vallet-Gely, I. & Boccard, F. Chromosomal Organization and Segregation in
Pseudomonas aeruginosa. *PLOS Genet.* **9**, e1003492 (2013).

10.   David, A. *et al.* The Two Cis-Acting Sites, parS1 and oriC1, Contribute to the
Longitudinal Organisation of Vibrio cholerae Chromosome I. *PLOS Genet.* **10**,
e1004448 (2014).

11.   Surovtsev, I. V. & Jacobs-Wagner, C. Subcellular Organization: A Critical Feature of
Bacterial Cell Replication. *Cell* **172**, 1271–1293 (2018).

12.   Jalal, A. S. B. & Le, T. B. K. Bacterial chromosome segregation by the ParABS
system. *Open Biology* vol. 10 (2020).

13.   Böhm, K. *et al.* Novel chromosome organization pattern in Actinomycetales-
overlapping replication cycles combined with diploidy. *MBio* **8**, (2017).

14.   Weber, P. M. *et al.* A Bidimensional Segregation Mode Maintains Symbiont
Chromosome Orientation toward Its Host. *Curr. Biol.* **29**, 3018-3028.e4 (2019).

15.   Kleckner, N. *et al.* The bacterial nucleoid: Nature, dynamics and sister segregation.
*Curr. Opin. Microbiol.* **22**, 127–137 (2014).

16.   Wang, X. T., Dong, P. F., Zhang, H. Y. & Peng, C. Structural heterogeneity and
functional diversity of topologically associating domains in mammalian genomes.
*Nucleic Acids Res.* **43**, 7237–7246 (2015).

17.   Le, T. B. K., Imakaev, M. V., Mirny, L. A. & Laub, M. T. High-resolution mapping of the
spatial organization of a bacterial chromosome. *Science (80-. ).* **342**, 731–734 (2013).

18.   Lioy, V. S. *et al.* Multiscale Structuring of the E. coli Chromosome by Nucleoid-
Associated and Condensin Proteins. *Cell* **172**, 771-783.e18 (2018).

19.   Lioy, V. S. *et al.* Dynamics of the compartmentalized Streptomyces chromosome
during metabolic differentiation. *Nat. Commun. 2021 121* **12**, 1–14 (2021).

20.   Ren, Z. *et al.* Centromere Interactions Promote the Maintenance of the Multipartite
Genome in Agrobacterium tumefaciens. *MBio* **13**, (2022).

21.   Szafran, M. J. *et al.* Spatial rearrangement of the Streptomyces venezuelae linear
chromosome during sporogenic development. *Nat. Commun. 2021 121* **12**, 1–15
(2021).

22.   Le, T. B. & Laub, M. T. Transcription rate and transcript length drive formation of
chromosomal interaction domain boundaries. *EMBO J.* **35**, 1582–1595 (2016).

23.   Hansen, A. S., Cattoglio, C., Darzacq, X. & Tjian, R. Recent evidence that TADs and
chromatin loops are dynamic structures. *Nucleus* **9**, 20–32 (2018).

24.   Busslinger, G. A. *et al.* Cohesin is positioned in mammalian genomes by transcription,
CTCF and Wapl. *Nature* **544**, 503–507 (2017).

25.   Wutz, G. *et al.* Topologically associating domains and chromatin loops depend on
cohesin and are regulated by CTCF, WAPL, and PDS5 proteins. *EMBO J.* **36**, 3573–

3599 (2017).

26.   Sanborn, A. L. *et al.* Chromatin extrusion explains key features of loop and domain

formation in wild-type and engineered genomes. *Proc. Natl. Acad. Sci. U. S. A.* **112**,

E6456–E6465 (2015).

27.   Matthey-Doret, C. *et al.* Computer vision for pattern detection in chromosome contact

maps. *Nat. Commun. 2020 111* **11**, 1–11 (2020).

28.   Dugar, G., Hofmann, A., Heermann, D. W. & Hamoen, L. W. A chromosomal loop

anchor mediates bacterial genome organization. *Nat. Genet.* **54**, 194–201 (2022).

29.   Cockram, C., Thierry, A., Gorlas, A., Lestini, R. & Koszul, R. Euryarchaeal genomes

are folded into SMC-dependent loops and domains, but lack transcription-mediated

compartmentalization. *Mol. Cell* **81**, 459-472.e10 (2021).

30.   Badel, C. & Bell, S. D. Chromosome architecture in an archaeal species naturally

lacking structural maintenance of chromosomes proteins. *Nat. Microbiol. 2023* 1–11

(2023) doi:10.1038/s41564-023-01540-6.

31.   Toro, E., Shapiro, L., Badrinarayanan, A., Le, T. B. K. & Laub, M. T. Bacterial

Chromosome Organization and Segregation. *Annu. Rev. Cell Dev. Biol.* **31**, 171–199

(2015).

32.   Nyongesa, S. *et al.* Evolution of longitudinal division in multicellular bacteria of the

Neisseriaceae family. *Nat. Commun.* **13**, 4853 (2022).

33.   Kuhn, D. A. & Gregory, D. A. D. A. Emendation of Simonsiella muelleri Schmid and

description of Simonsiella steedae sp. nov., with designations of the respective

proposed neotype and holotype strains. *Curr. Microbiol. An Int. J.* **1**, 11–14 (1978).

34.   Xie, C. H. & Yokota, A. Phylogenetic analysis of Alysiella and related genera of

Neisseriaceae: Proposal of Alysiella crassa comb. nov., Conchiformibium steedae

gen. nov., comb. nov., Conchiformibium kuhniae sp. nov. and Bergeriella denitrificans

gen. nov., comb. nov. *J. Gen. Appl. Microbiol.* **51**, 1–10 (2005).

35.   Tønjum, T. Alysiella. *Bergey's Man. Syst. Archaea Bact.* 1–5 (2015)

doi:10.1002/9781118960608.gbm00973.

36.   Simons, H. Saprophytische Oscillarien des Menschen und der Tiere. *Zentralbl.*

*Bakteriol. Parasitenkd. Infekt. Hyg. Abt* **1**, 501–510 (1922).

37.   Hedlund, B. P. & Tønjum, T. Simonsiella. *Bergey's Man. Syst. Archaea Bact.* 1–12

(2015) doi:10.1002/9781118960608.gbm00983.

38.   Kaiser, G. E. & Starzyk, M. J. Ultrastructure and cell division of an oral bacterium

resembling Alysiella filiformis. *Can. J. Microbiol.* **19**, 325–327 (1973).

39.   Leisch, N. *et al.* Asynchronous division by non-ring FtsZ in the gammaproteobacterial

symbiont of Robbea hypermnestra. *Nat. Microbiol.* **2**, 1–5 (2016).

40.   Reyes-Lamothe, R. & Sherratt, D. J. The bacterial cell cycle, chromosome inheritance

and cell growth. *Nat. Rev. Microbiol. 2019 178* **17**, 467–478 (2019).

41.  Ivanova, D. *et al.* Shaping the landscape of the Escherichia coli chromosome:

Replication-transcription encounters in cells with an ectopic replication origin. *Nucleic*

*Acids Res.* **43**, 7865–7877 (2015).

42.  Varoquaux, N., Lioy, V. S., Boccard, F. & Junier, I. Computational Tools for the

Multiscale Analysis of Hi-C Data in Bacterial Chromosomes. *Methods Mol. Biol.* **2301**,

197–207 (2022).

43.  Val, M. E. *et al.* A checkpoint control orchestrates the replication of the two

chromosomes of Vibrio cholerae. *Sci. Adv.* **2**, (2016).

44.  Colclough, A. L., Scadden, J. & Blair, J. M. A. TetR-family transcription factors in

Gram-negative bacteria: Conservation, variation and implications for efflux-mediated

antimicrobial resistance. *BMC Genomics* **20**, 1–12 (2019).

45.  Leisch, N. *et al.* Growth in width and FtsZ ring longitudinal positioning in a

gammaproteobacterial symbiont. *Curr. Biol.* **22**, R831–R832 (2012).

46.  Pende, N. *et al.* Host-Polarized Cell Growth in Animal Symbionts. *Curr. Biol.* **28**, 1039-

1051.e5 (2018).

47.  Anchimiuk, A. *et al.* A low smc flux avoids collisions and facilitates chromosome

organization in bacillus subtilis. *Elife* **10**, (2021).

48.  Tobiason, D. M. & Seifert, H. S. Genomic content of Neisseria species. *J. Bacteriol.*

**192**, 2160–2168 (2010).

49.  Hildenbrand, C., Stock, T., Lange, C., Rother, M. & Soppa, J. Genome copy numbers

and gene conversion in methanogenic archaea. *J. Bacteriol.* **193**, 734–743 (2011).

50.  Soppa, J. Polyploidy in archaea and bacteria: About desiccation resistance, giant cell

size, long-term survival, enforcement by a eukaryotic host and additional aspects. *J.*

*Mol. Microbiol. Biotechnol.* **24**, 409–419 (2014).

51.  Zborowsky, S. & Lindell, D. Resistance in marine cyanobacteria differs against

specialist and generalist cyanophages. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 16899–

16908 (2019).

52.  Llopis, P. M. *et al.* Spatial organization of the flow of genetic information in bacteria.

*Nature* **466**, 77–81 (2010).

53.  Libby, E. A., Roggiani, M. & Goulian, M. Membrane protein expression triggers

chromosomal locus repositioning in bacteria. *Proc. Natl. Acad. Sci. U. S. A.* **109**,

7445–7450 (2012).

54.  Kaval, K. G. *et al.* Membrane-localized expression, production and assembly of Vibrio

parahaemolyticus T3SS2 provides evidence for transertion. *Nat. Commun.* **14**, (2023).

55.  Wang, X., Brandão, H. B., Le, T. B. K., Laub, M. T. & Rudner, D. Z. Bacillus subtilis

SMC complexes juxtapose chromosome arms as they travel from origin to terminus.

*Science (80-. ).* **355**, 524–527 (2017).

56.    Dekker, C., Haering, C. H., Peters, J. M. & Rowland, B. D. How do molecular motors

fold the genome? *Science (80-. ).* **382**, 646–648 (2023).

57.    Chang, L. H., Ghosh, S. & Noordermeer, D. TADs and Their Borders: Free Movement

or Building a Wall? *J. Mol. Biol.* **432**, 643–652 (2020).

58.    Livny, J., Yamaichi, Y. & Waldor, M. K. Distribution of centromere-like parS sites in

bacteria: Insights from comparative genomics. *J. Bacteriol.* **189**, 8693–8703 (2007).

59.    Carnoy, C. & Roten, C. A. The dif/Xer recombination systems in proteobacteria. *PLoS*

*One* **4**, e6531 (2009).

60.    Barrero-Canosa, J., Moraru, C., Zeugner, L., Fuchs, B. M. & Amann, R. Direct-

geneFISH: a simplified protocol for the simultaneous detection and quantification of

genes and rRNA in microorganisms. *Environ. Microbiol.* **19**, 70–82 (2017).

61.    Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of

image analysis. *Nat. Methods 2012 97* **9**, 671–675 (2012).

62.    Vischer, N. O. E. *et al.* Cell age dependent concentration of Escherichia coli divisome

proteins analyzed with ImageJ and ObjectJ. *Front. Microbiol.* **6**, 135279 (2015).

63.    Ducret, A., Quardokus, E. M. & Brun, Y. V. MicrobeJ, a tool for high throughput

bacterial cell detection and quantitative analysis. *Nat. Microbiol.* **1**, 1–7 (2016).

64.    Valero-Mora, P. M. *ggplot2: Elegant Graphics for Data Analysis . Journal of Statistical*

*Software* vol. 35 (Springer-Verlag New York, 2010).

65.    R Core Team. R: A Language and Environment for Statistical Computing. (2021).

66.    Koren, S. *et al.* Canu: Scalable and accurate long-read assembly via adaptive κ-mer

weighting and repeat separation. *Genome Res.* **27**, 722–736 (2017).

67.    Vaser, R., Sović, I., Nagarajan, N. & Šikić, M. Fast and accurate de novo genome

assembly from long uncorrected reads. *Genome Res.* **27**, 737–746 (2017).

68.    Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, 1–4

(2021).

69.    Walker, B. J. *et al.* Pilon: An integrated tool for comprehensive microbial variant

detection and genome assembly improvement. *PLoS One* **9**, e112963 (2014).

70.    Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler

transform. *Bioinformatics* **26**, 589–595 (2010).

71.    Matsen, F. A., Kodner, R. B. & Armbrust, E. V. pplacer: linear time maximum-

likelihood and Bayesian phylogenetic placement of sequences onto a fixed reference

tree. *BMC Bioinformatics* **11**, 538 (2010).

72.    Mistry, J., Finn, R. D., Eddy, S. R., Bateman, A. & Punta, M. Challenges in homology

search: HMMER3 and convergent evolution of coiled-coil regions. *Nucleic Acids Res.*

**41**, e121–e121 (2013).

73. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
Assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res.* **25**, 1043–1055 (2015).

74. Seemann, T. Prokka: Rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–
2069 (2014).

75. Huerta-Cepas, J. *et al.* Fast Genome-Wide Functional Annotation through Orthology
Assignment by eggNOG-Mapper. *Mol. Biol. Evol.* **34**, 2115–2122 (2017).

76. Huerta-Cepas, J. *et al.* eggNOG 5.0: a hierarchical, functionally and phylogenetically
annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic*
*Acids Res.* **47**, D309–D314 (2019).

77. Vallenet, D. *et al.* MicroScope: a platform for microbial genome annotation and
comparative genomics. *Database* **2009**, (2009).

78. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol* **30**, 772–80 (2013).

79. Zimmermann, L. *et al.* A Completely Reimplemented MPI Bioinformatics Toolkit with a
New HHpred Server at its Core. *J. Mol. Biol.* **430**, 2237–2243 (2018).

80. Gabler, F. *et al.* Protein Sequence Analysis Using the MPI Bioinformatics Toolkit.
*Curr. Protoc. Bioinforma.* **72**, e108 (2020).

81. Robert, X. & Gouet, P. Deciphering key features in protein structures with the new
ENDscript server. *Nucleic Acids Res.* **42**, W320–W324 (2014).

82. Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

83. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., Von Haeseler, A. & Jermiin, L. S.
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods*
*2017 146* **14**, 587–589 (2017).

84. Hoang, D. T., Chernomor, O., Von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBoot2:
Improving the Ultrafast Bootstrap Approximation. *Mol. Biol. Evol.* **35**, 518–522 (2018).

85. Warnes, G. R. *et al.* gplots: Various R Programming Tools for Plotting Data. (2016).

86. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat.*
*Methods* **9**, 357–359 (2012).

87. Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing
data analysis. *Nucleic Acids Res.* **44**, W160–W165 (2016).

88. Krzywinski, M. *et al.* Circos: An information aesthetic for comparative genomics.
*Genome Res.* **19**, 1639–1645 (2009).

89. Wall, L., Christiansen, T. & Orvant, J. *Programming Perl*. (O'Reilly Media, Inc, 2000).

90. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

91.  Schmieder, R. & Edwards, R. Quality control and preprocessing of metagenomic
datasets. *Bioinformatics* **27**, 863–864 (2011).
92.  Bushnell, B. *BBMap: A Fast, Accurate, Splice-Aware Aligner. Ernest Orlando*
*Lawrence Berkeley National Laboratory, Berkeley, CA (US)* (2014). doi:10.1186/1471-
2105-13-238.
93.  Imakaev, M. *et al.* Iterative correction of Hi-C data reveals hallmarks of chromosome
organization. *Nat. Methods* **9**, 999–1003 (2012).
94.  Kopylova, E., Noé, L. & Touzet, H. SortMeRNA: Fast and accurate filtering of
ribosomal RNAs in metatranscriptomic data. *Bioinformatics* **28**, 3211–3217 (2012).
95.  Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930
(2014).
96.  Hunter, J. D. Matplotlib: A 2D graphics environment. *Comput. Sci. Eng.* **9**, 90–95
(2007).
97.  Van Rossum, G. & Drake, F. L. *Python 3 Reference Manual*. (CreateSpace, 2009).
98.  Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).
99.  Xie, Z. & Tang, H. ISEScan: automated identification of insertion sequence elements
in prokaryotic genomes. *Bioinformatics* **33**, 3340–3347 (2017).

683
684

685  **LEAD CONTACT AND MATERIALS AVAILABILITY**
686  Further information and requests for resources and reagents should be directed to and will
687  be fulfilled by the lead contact, Silvia Bulgheresi (silvia.bulgheresi@univie.ac.at). This study
688  did not generate new strains or unique reagents.

689
**DATA AND CODE AVAILABILITY**
Data for genome sequencing has been deposited in Bioproject PRJNA687213 (*A. filiformis*),
PRJNA687218 (*S. muelleri*), and PRJNA779076 (*C. steedae*), and genomes have been
deposited in GenBank as GCF_020162295.1 (*A. filiformis*), GCF_020162275.1 (*S. muelleri*),
and GCF_023547005.1 (*C. steedae*). Data for marker frequency analysis has been deposited
in Bioproject PRJNA795449 (*A. filiformis*), PRJNA795451 (*S. muelleri*), and PRJNA795450
(*C. steedae*), and corresponding non-normalized, 1kb-binned read count files in Gene
Expression Omnibus (GEO) GSE194131 (*A. filiformis*), GSE194134 (*S. muelleri*), and
GSE205139 (*C. steedae*). Data for RNA-Seq has been deposited in Bioproject PRJNA795348
(*A. filiformis*) and PRJNA795350 (*S. muelleri*), and corresponding raw read count files in GEO
GSE194132 (*A. filiformis*), and GSE194135 (*S. muelleri*). Data for 3C-seq has been deposited

in Bioproject PRJNA837061 (*A. filiformis*), PRJNA837062 (*S. muelleri*), PRJNA1052629 (*C.*
*steedae),* and corresponding matrices in GEO GSE250400 (*A. filiformis*), GSE250402 (*S.*
*muelleri*), and GSE250401 (*C. steedae*). SRA accession numbers for sequencing reads and
GEO accession numbers for data files are listed in Supplementary Table 4. The
documentation for the ImageJ plugin Fil-Tracer can be accessed here:
https://sils.fnwi.uva.nl/bcb/objectj/examples/Fil-Tracer/MD/Fil-Tracer.html.
707
708

**EXPERIMENTAL MODEL AND SUBJECT DETAILS**
The bacterial strains *Alysiella filiformis* (DSM 16848), *Simonsiella muelleri* (DSM 2579) and
*Conchiformibius steedae* (DSM 2580) have been received from the German Collection of
Microorganisms and Cell Cultures GmbH (DSMZ). All *Neisseriaceae* strains were prepared
for experimentation as described below unless otherwise noted. *A. filiformis* cultures were pre-
cultured by inoculating liquid peptone-yeast (PY) medium with glycerol stocks stored at -70°C
and grown overnight at 37°C at 120 rpm, transferred to fresh PY medium the next day and
grown until the specified $OD_{600}$ (see each section). *S. muelleri* and *C. steedae* cultures were
streaked out from glycerol stocks onto BSTSY agar plates, grown overnight at 37°C, and
colonies transferred into liquid BSTSY medium, and grown at 37°C at 120 rpm until the
specified $OD_{600}$ (see each section).
For sampling symbiotic nematodes, sediment samples were collected on multiple field
trips (2015-2019) in 1 m depth from a sand bar off Carrie Bow Cay, Belize (16°48'11.01''N,
88°4'54.42''W). Specimens of *Robbea hypermnestrae* were extracted from the sediment by
stirring the sand in seawater and pouring the supernatant through a 63 µm mesh sieve. The
retained material was transferred into a Petri dish and single nematodes were handpicked
using pipettes under a dissecting microscope. For DNA FISH, whole symbiotic nematodes
were fixed in 3 or 4% PFA for 12-14 h at 4°C, washed with 70% ethanol and stored in 70%
ethanol at -20°C. Fixed symbiotic nematodes were transported from Carrie Bow Cay to the
University of Vienna deep-frozen.
729

**METHOD DETAILS**
**Generation of DNA fluorescence in situ hybridization (FISH) probes.** We used the
genome draft of *Ca.* T. hypermnestrae (Genbank accession number GCA_020443945.1) to
design specific primers against the *Ca.* T. hypermnestrae *ori* and *ter* regions, as well as
against DNA stretches containing the *parS2* (on a different contig than *ori* and *parS1)* [58] or the
*ftsQAZ* operon (Supplementary Table 1; *Ca.* T. hypermnestrae *dif* site sequence:
ACTTTACATAATATACATTATGCGTAA) [59]. *Robbea hypermnestrae* nematodes were
rehydrated in phosphate-buffered saline (PBS) and bacteria were detached from the worms

by sonication. Subsequently, 1 µl of bacterial suspension was used as template in each 25 µl
PCR reaction (primer sequences and PCR conditions are listed in Supplementary Tables 2
and 3). A 2,858 nt-long fragment containing the *dnaA* and *dnaN* genes (targeting the *ori*), a
3,690 nt-long fragment containing the *ftsQAZ* operon, a 3,524 nt-long fragment containing the
*parS2* site (targeting an *ori*-proximal site) and a 3,239 nt-fragment containing the predicted *dif*
site (targeting a *ter*-proximal site) were amplified. Each purified fragment was then used as
template to PCR-amplify dsDNA polynucleotide probes (referred to as *ori*, *ftsQAZ*, *parS2* and
*ter*, respectively) (see Supplementary Tables 2 and 3 for primers and PCR conditions).

For the oral cavity symbionts *Alysiella filiformis* and *Simonsiella muelleri*, we used
Gene-PROBER (Moraru, 2021) to design specific primers hybridizing with the origin or
terminus of DNA replication or in their immediate vicinity (Supplementary Fig.2). For gDNA
extraction, cells were lysed using lysozyme and Proteinase K, DNA extracted using Phenol-
Chloroform and precipitated using ammonium acetate (2.5 M final concentration), two volumes
of 100% ice-cold Ethanol, and 5 µg/ml glycogen. RNA was degraded by incubation with 10
mg/ml RNase A for 30 min at 37°C, and DNA cleaned up using DNA Clean & Concentrator-5
(Zymo). Extracted gDNA was used as a template in each 25 µl PCR reaction (primers
sequences and PCR conditions are listed in Supplementary Tables 2 and 3), amplifying a
9,232 nt-long *ori* fragment and a 6,778 nt-long *ter* fragment from *A. filiformis*, an 8,158 nt-long
*ori* and an 8,187 nt-long *ter* fragment from *S. muelleri*, and a 7,353 nt-long *ori* and a 4,498 nt-
long *ter* fragment from *C. steedae.* Each purified fragment was then used as template to PCR-
amplify dsDNA polynucleotide probes (referred to as Af/Sm/Cs-*ori*, and as Af/Sm/Cs-*ter*,
respectively) (see Supplementary Table 2 and 3 for primers and PCR conditions).

All polynucleotide probes were chemically labeled with the Alexa Fluor 594 using the
Ulysis Nucleic Acid Labeling Kit (ThermoFisher) following the same modifications to the
manufacturers' protocol as in (Barrero-Canosa et al., 2016).

**DNA fluorescence in situ hybridization (FISH)**

Single *Robbea hypermnestrae* nematodes were rehydrated in PBS and nematode
symbionts *Ca.* T. hypermnestrae were detached by sonication. Exponentially growing oral
cavity symbionts were fixed in 3% or 4% PFA for 12-14 h at 4°C or for 1 h at room temperature,
washed with PBS and stored in 1:1 ethanol/PBS. For the hybridization procedure, we followed
a slightly modified version of the direct-gene FISH protocol [60]. After letting the cell suspension
dry onto a well of a Poly-L-lysine coated Epoxy-slide, cells were dehydrated in a series of
increasing ethanol concentrations and permeabilized with freshly prepared lysozyme solution
for 1 h on ice or at 37°C. Probes were diluted in hybridization buffer containing 35 or 45%
formamide to final concentrations ranging 62-246 pg/ml and each probe was applied to the
cells individually. Slides were transferred into a hybridization chamber and incubated for 40
min at 85°C and subsequently at 46°C for 2 h. Washing buffer was applied to the cells once

briefly and once for 15 min at 48°C and, finally, cells were incubated in PBS and PBS
supplemented with 5 µg/ml DNA stain Hoechst 33342 for 10 min at room temperature. Upon
a quick wash in ddH$_2$O and, subsequently, in 100% ethanol, cells were air-dried and mounted
in 4.5 µl Vectashield mounting medium (Vector Labs) per microscopic slide well.

**DNA staining.** Methanol-fixed nematodes or oral-cavity symbionts (fixed in 3% or 4%
PFA for 12-14 h at 4°C or for 1 h at room temperature) were rehydrated and washed in PBS
containing 0.1% Tween 20 (PBT), followed by incubation in 5 µg/ml Hoechst 33342 PBT for
15 min. After two washing steps to remove unbound DNA stain, worms were sonicated for 45
s to dissociate *Ca.* T. hypermnestrae from its host prior mounting. 1.5 ml of the bacterial
suspension was mixed with 0.75 ml of Vectashield mounting medium (Vector Labs) and
applied to a 1% agarose covered microscopy slide.

**Fluorescence microscopy.** Slides containing symbiont cells subjected to DNA FISH
were imaged using a Nikon Eclipse NI-U microscope equipped with a MFCool camera
(Jenoptik). Images were acquired using the ProgRes Capture Pro 2.8.8 software (Jenoptik).
Microscopic images were processed using the public domain program ImageJ [61] in
combination with plugin ObjectJ [62] and Fil-Tracer [32] or MicrobeJ 5.13J [63]. Cell outlines were
traced and morphometric measurements recorded. Positions of the fluorescence foci (i.e.,
points of maximal fluorescent emissions) within the cell boundaries were measured and
plotted as fraction of the normalized cell width and length of the cell that contained them.
Automatic cell recognition was double-checked manually. For representative images, the
background subtraction function of ImageJ was used, and brightness and contrast were
adjusted for better visibility. Data analysis was performed using Excel 2021 (Microsoft
Corporation), density plots were created with the ggplot2 [64] in R v.3.6.1 [65] and figures were
compiled using Illustrator 2021 (Adobe Systems).

**Western blotting.** The western blot against ParB was done as previously mentioned
in Weber et al., 2019, with slight amendments. The primary antibody used was a custom
peptide rabbit polyclonal anti- *A. filiformis* ParB antibody (Eurogentec) with a 1:1000 dilution.
The secondary antibody used was a horseradish peroxidase-conjugated anti-rabbit secondary
antibody (Amersham Biosciences) with a 1:10,000 dilution.

**Immunostaining.** Deep-frozen methanol-fixed nematodes were rehydrated and
washed in PBS containing 0.1% Tween 20 (PBT), followed by permeabilization of the bacterial
peptidoglycan by a 15 min incubation with 0.1% (wt/vol) lysozyme at room temperature.
Blocking was carried out for 1 hr in PBT containing 2% (wt/vol) bovine serum albumin (blocking
solution) at room temperature. *Ca.* T. hypermnestrae was incubated with a 1:500 dilution of
peptide rabbit polyclonal anti-*Ca.* T. oneisti ParB antibody (Eurogentec). All primary antibodies
were incubated in blocking solution overnight at 4°C. Upon incubation with primary antibody
(or without, in the case of the negative control) samples were washed three times in PBT and

incubated with a 1:500 dilution of secondary Alexa 488-conjugated anti-rabbit antibody

(Jackson ImmunoResearch, USA) in blocking solution for 1 hr at room temperature. Unbound

secondary antibody was removed by three washing steps in PBT and thereupon incubated in

5 mg/ml Hoechst 33342 PBT for 15 min. After two washing steps to remove unbound DNA

stain, worms were sonicated for 45 s to dissociate *Ca*. T. hypermnestrae from its host prior

mounting. 1.5 ml of the bacterial suspension was mixed with 0.75 ml of Vectashield mounting

medium (Vector Labs) and applied to a 1% agarose covered microscopy slide.

**Genome sequencing and assembly.** For whole-genome sequencing, cells were

lysed using lysozyme and Proteinase K, DNA extracted using phenol-chloroform and

precipitated in two volumes of 100% ice-cold ethanol, 2.5 M ammonium acetate and 5 µg/ml

glycogen. RNA was degraded by incubation with 10 mg/ml RNase A for 30 min at 37°C, and

DNA cleaned up using DNA Clean & Concentrator-5 (Zymo). The library for Oxford Nanopore

Technologies (ONT) sequencing was prepared using the ONT 1D ligation sequencing kit

(SQK-LSK110) and sequenced on an R9.4 flow cell (FLO-MIN106) on a MinION for 48 h. Base

calling was performed locally with ONT's Guppy Basecalling Software guppy

5.0.11+2b6dbffa5 (dna_r9.4.1_450bps_sup.cfg), and resulting fastq-files were trimmed using

using qcat (https://github.com/nanoporetech/qcat). For short read sequencing, sequencing

libraries were prepared using the DNA NEB Ultra II FS (NEB) and sequenced on an Illumina

NovaSeq SP with 100 bp single-end reads at the VBC NGS facility. The ONT data was

assembled using canu v2.1.1 with genomeSize=2.5m [66], and polished with the long reads

using 5 rounds of racon v1.4.3 (-m 8 -x -6 -g -8 -w 500, [67], reads mapped with minimap2 v2.7

-ax map-ont (Li, 2018) and samtools v1.9 [68], and one round of medaka v1.74.0 using the

'consensus'    (--model    r941_min_sup_g507)    and    'stitch'    modules

(https://github.com/nanoporetech/medaka). The resulting assemblies were polished with the

short reads using 5 rounds of pilon v1.22 [69] with bwa v0.7.16a mem [70] mapped reads. The

genome completeness was assessed using CheckM version 1.0.18 [71–73] with the

gammaproteobacterial marker gene set using the taxonomy workflow. Assembled contigs

were annotated using Prokka v1.14.6 [74]. Cluster of orthologous genes (COGs) were annotated

using eggnog mapper v2.1.10 [75] against eggNOG 5.0 [76]. Additional annotations were obtained

through the MicroScope platform [77]. For the ParB phylogeny, the alignment was performed

with mafft v7.427 (L-INS-I mode) [78], and secondary structures predicted using ALi2D [79,80] and

ESPript 3 [81] using the crystal structure of *Caulobacter crescentus* ParB (PDB# 6T1F). The

maximum likelihood phylogeny was reconstructed using IQ-TREE v2.1.2 [82] with the best-fit

model automatically selected by ModelFinder Plus [83] and 1,000 ultrafast bootstraps [84]. The

phylogeny was visualized in FigTree v1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/).

**Prediction of the origin and of the terminus of DNA replication.** The *ori* and *ter*

positions    were    calculated    based    on    GC    skew    and    using    Genskew

(https://genskew.csb.univie.ac.at/) and smoothed using the lowess function (smootherspan =
0.005) of gplots v3.1.1 [85] in R 3.6.1 [65]. *ParS* sequences were predicted using the conserved
consensus *parS* (5'-NGTTNCANGTGNAACN-3') [58] and *dif* sites using the betaproteobacterial
consensus *dif* sequence (5'-HNNNBNNAYVAYNNDBVTTATGTHAANT-3', [59]. Sequencing
coverage was obtained by mapping the raw genomic reads obtained for genome sequencing
against the reference genomes using bowtie2 v2.4.4 [86] and samtools v1.9 [68]. Mapped reads
were counted and binned in 5 kb non-overlapping windows using bamCoverage of
(standalone) deeptools v3.5.1 [87], and the resulting bigwig file converted to human-readable
format using the bigWigToWig of UCSC v391. The coverage data was square transformed to
better depict differences between *ori* and *ter*, and smoothed using the lowess function
(smootherspan = 0.05). For illustration purposes (circular plots and 3C-seq heatmaps), the
assemblies were reoriented to have the GC-skew-predicted *ori* in the middle of the genome
sequence. Circular plots were generated using circos v0.69-8 [88] with Perl 5.028003 [89].

**Marker frequency analysis (MFA).** Cells were grown until an $OD_{600}$ 0.6 (late
exponential phase) and 1 (stationary phase) and collected by centrifugation at 16,000 g for 5
min at room temperature. Cells were lysed using lysozyme and Proteinase K, DNA extracted
using Phenol-Chloroform and precipitated using ammonium acetate (2.5M final
concentration), two volumes of 100% ice-cold Ethanol, and 5 µg/ml glycogen. RNA was
degraded by incubation with 10 mg/ml RNase A for 30 min at 37°C, and DNA cleaned up using
DNA Clean & Concentrator-5 (Zymo).

Sequencing libraries were prepared at the VBC NGS facility, and 150bp paired-end
sequenced on an Illumina NovaSeq 6000 machine. Reads were adapter-trimmed and filtered
using trimmomatic v0.39 (ILLUMINACLIP:adapters.fa:2:30:10 LEADING:3 TRAILING:3
MINLEN:80 SLIDINGWINDOW:4:15 [90] and prinseq-lite v0.20.4 (-min_qual_mean 30 -min_len
25) [91]. PhiX contaminant sequences were removed using bbduk.sh of bbmap v38.90 (k=31
hdist=1) [92], as well as further contaminants (human, animal, plant, fungi) removed against
repeat-masked references (minid=0.95 maxindel=3 bwr=0.16 bw=12 quickmatch fast
minhits=2 qtrim=rl trimq=10 untrim). Reads were mapped against the reference chromosomes
generated in this study using bowtie2 v2.4.4 [86], and index-sorted using samtools v1.9 [68].

Reads were counted using the bamCoverage program of (standalone) deeptools
v3.5.1 [87] in non-overlapping 1kb windows and the resulting bigwig file converted to human-
readable format using the bigWigToWig of UCSC v391. The count data was smoothed using
the lowess function (smootherspan = 0.1) and plotted using ggplot2 [64] in R v.3.6.1 [65], and the
ratio of the GC-skew predicted *ori* to *ter* coverage (number of non-normalized mapped reads)
calculated.

**Chromosome conformation capture (3C-seq).** 3C libraries were prepared as
previously described [7,19]. For growth on solid medium, *C. steedae, A. filiformis* and *S. muelleri*

were plated from glycerol stock onto BSTSY agar plates and grown at 37°C for 24 h and 48
h, after which they were scraped off and transferred to BSTSY/FBS containing 6% f.c
formaldehyde, fully resuspended, and fixed at room temperature at 120 rpm, followed by 30
min more at 4°C. Glycine (250 mM f.c.) was added, and the cells incubated 30 min at 4°C at
120 rpm. After fixation, cells were collected by centrifugation for 10 min at 3,500 g at 4°C,
resuspended in fresh medium, collected again by centrifugation, and resuspended in 1 ml
fresh medium before final collection by centrifugation. Cell pellets were stored at -70°C. For
liquid cultures, *A. filiformis* was grown until $OD_{600}$ of 0.3 (exponential phase) or 1 (stationary
phase). For blocking transcription or translation, cells grown overnight on plate were incubated
with Rifampicin (25 µg/ml f.c.) or Chloramphenicol (25 µg/ml) for 30 min prior to cell fixation
and collection.

For digestion, cell pellets were thawed on ice for 10 min, resuspended in 500 µl Tris
10 mM EDTA 0.5 mM (TE) (pH 8), and lysed using 4 µl Ready-lyse lysozyme (Epicentre) for
20 min at 37°C, 300 rpm. SDS (0.5% f.c.) was added and incubated for 10 min at room
temperature. 500 µl lysed cells were then transferred to a tube containing 4.5 ml digestion
buffer (1% Triton X-100, 1x CutSmart buffer (New England Biolabs)), and 100 µl cells (non-
digested control) were transferred to a tube containing 0.9 ml of digestion buffer. 100U of HpaII
(New England Biolabs) were added to the 5 ml digestion mix. Both tubes were incubated for
3 h at 37°C. After digestion, 1 ml of digestion mix was put on ice (digested control) together
with the non-digested control. The remaining digestion mix was split into 4 parts, centrifuged
for 20 min at 20,000 g, and pellets were carefully resuspended in 1 ml water each, and pooled
into 7 ml ligation buffer (1X ligation buffer 3 (New England Biolabs; without ATP), 1 mM ATP,
1 mg/ml BSA, 500 U of T4 DNA ligase 5 U/µl (ThermoFisher)). Ligation was carried out at
16°C for 4 h, followed by incubation together with the two control tubes overnight at 65°C with
proteinase K (0.25 mg/ml f.c.) and EDTA (6 mM f.c.). The next day, DNA of the ligated sample
was precipitated with 1/10 volume of 3 M sodium acetate (pH 5.2) and two volumes of
isopropanol, and after one hour at -80°C, DNA was pelleted by centrifugation for 30 min at
10,000 g at 4°C and resuspended in 500 µl 1X TE buffer. DNA in all tubes (ligation and two
controls) were extracted once with 400 µl phenol-chloroform pH 8.0 (vortexed for 1 min and
centrifuged for 5 min 16,000 g at 4°C), precipitated (see above; 30 min at -80°C, followed by
centrifugation for 20 min at 16,000 x g at 4°C), washed with 400 µl 80% cold ethanol, and
diluted in 30 µl 1X TE buffer supplemented with RNase A (1 mg/ml f.c.; ThermoFisher). Tubes
containing ligated DNA were pooled. A 1% agarose gel was run with aliquots of the ligated
and two control DNA to check for efficiency of digest and ligation.

Ligated DNA (5 µg) was adjusted with water to 130 µl, and sheared using a Covaris
S220 (Duty cycle 5, Intensity 5, cycles/burst 200, time 60 s for 4 cycles), and the DNA purified
using the QIAquick PCR purification kit (40 µl elution buffer; Qiagen), and DNA end-repaired

with 1X T4 DNA ligase buffer (New England Biolabs), 0.4 mM dNTP mix, 15 U T4

polynucleotide kinase (New England Biolabs), 5 U Klenow DNA polymerase (Roche), and

incubated for 30 min at RT. DNA was purified using MinElute columns (Qiagen). A-tailing was

done with 1X NEBuffer 2 (New England Biolabs), 0.2 mM dATP, and 15 U Klenow exo (New

England Biolabs), for 30 min at 37°C before inactivation for 20 min at 65°C, and DNA again

purified using MinElute columns (Qiagen). Sequencing adapters were ligated with 0.5 mM

custom-made adapters, $2.10^6$ U T4 DNA ligase, and 1X T4 DNA ligase buffer, for 2 hours at

RT, followed by inactivation for 20 min at 65°C. DNA fragments were purified in a size from

400 to 900 bp using a PippinPrep apparatus (SAGE Science). Each library was amplified in

12 cycles with 1X Phusion Buffer (ThermoFisher) , 0.2 mM dNTPs, 0.2 µM Illumina primers

PE1.0     (5'-AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGA-3')     and

PE2.0 (5'-CAAGCAGAAGACGGCATACGAGAT-3'), 3 µl of 3C library, and 1U of Taq Phusion

polymerase (ThermoFisher; 98 °C 30 s, 12 cycles of 65°C 30 s, and 72°C 30 s, and 72°C for

7 min). The PCR reaction was purified using MinElute PCR purification kit (Qiagen), and

libraries sequenced on an Illumina NextSeq500 with 75 bp paired-end reads.

**Generation of contact maps.** Raw contact maps were built as described previously

[18]. Briefly, each read was assigned to a restriction fragment. Non- informative events such as

self-circularized restriction fragments, or uncut co- linear restriction fragments were discarded.

The chromosome of each bacterium was divided into 5 kb bins and the frequencies of contacts

between genomic loci for each bin were assigned. Contact frequencies were visualized as

heatmaps. Raw contact maps used for comparative analyses were built using 1 million valid

reads. Raw contact maps were normalized using Iterative correction and eigenvector

decomposition (ICE) [93]To facilitate visualization, normalized contact matrices are visualized

as log matrices. A Gaussian filter of H=0.5 is used when plotting the matrices. To quantify the

frequency of interactions in the secondary diagonals (Supplementary Figure 6), we applied to

our data a method based on the filtering of significant 3C contacts and developed as described

[55].

**Identification of boundaries in contact maps.** To identify boundaries we applied the

Frontier Index (FI) method as described in detail in [42] and summarized in [19]. Briefly, the FI

method allows to identify loci associated with demarcations, or frontiers, in the 3C contact

maps that can occur at any scale, meaning that the method is useful to identify domain

organization irrespective of the underlying scale. Specifically, at a given locus (let say *i*), two

frontiers can be found at most, each one corresponding to a side of the locus – one frontier

indicates that the neighboring loci on the same side of the locus *i* tend to make more contacts

with each other with respect to neighboring loci separated by the same genomic distances but

located on each side of the locus. Visually, identified frontiers can be vertical (or upstream) or

horizontal (downstream) (see e.g. Fig. 4) [42].

**Loop analysis**. To facilitate analysis, normalized bins interacting with less than 10%
of the total number of bins were considered empty due to multi-mapping reads such as multiple
rRNA operons and removed from the analysis. This corresponded to approximately 10% of
the total number of bins in S. muelleri. In practice, we removed the corresponding row and
columns of the 3C-seq matrices and analyzed the resulting smaller matrices.

Loop detection was performed as following. We first slightly smoothed 3C-seq matrices
using a Gaussian filter with width 0.5. Next, we computed a 4C profile for each bin. Peaks in
these profiles were good candidates for loops such that we extracted the highest peaks of
each profile, where the height of a peak was defined as the difference of 4C value between
the peak and the maximum value of the two flanking minima. The resulting histogram of the
heights could be divided into an exponential distribution for small values and a strong deviation
from this exponential distribution for large values. In this context, we considered a conservative
threshold value to distinguish the two sets of values. 4C peaks higher than this threshold were
then considered as loop. Eventually, looping regions were defined as maximal sets of
contiguous bins such that any two consecutive bins were involved in at least one loop with a

**Transcriptomics.** All organisms were grown in the same conditions as those applied
for the 3C-seq experiments in three biological replicates each as for the 3C-seq experiments.
500 µl liquid culture were mixed well with 1 ml RNAprotect Bacteria Reagent (Qiagen) and
incubated for 5 min, and cells grown on solid medium scraped off and resuspended in 1 ml
thereof. Cells were collected by centrifugation for 10 min at 5,000 g, and cell pellets stored at
-70°C until the next day.

Total RNA was extracted using the RNeasy Mini Kit (Qiagen) for low cell numbers,
including the on-column DNA digestion. Eluted RNA (30 µl) was treated with 2 U of DNase for
30 min using the TURBO DNA-free kit (Invitrogen). RNA was cleaned up using the RNA Clean
& Concentrator-25 kit (Zymo) including the on-column DNA digestion. Absence of DNA was
checked via PCR (35 cycles) against a ~200 bp fragment of *gyrB*. RNA was quality checked
at the VBC NGS facility (all samples had a RIN > 9), rRNA depleted using an in-facility
developed protocol, and reverse-stranded sequencing libraries were prepared using the
NEBNext Ultra II Directional RNA Library Prep Kit (New England Biolabs). Libraries were
sequenced on an Illumina NovaSeq 6000 machine, using 100 bp single-end reads (*A.
filiformis*) and 150 bp paired-end reads (*S. muelleri, C. steedae*). Reads were adapter-trimmed
and filtered using trimmomatic v0.39 (ILLUMINACLIP:adapters.fa:2:30:10 LEADING:3
TRAILING:3 MINLEN:80 SLIDINGWINDOW:4:15; (Bolger et al., 2014)) and prinseq-lite
v0.20.4 (-min_qual_mean 30 -min_len 25) [92], as well as further contaminants (human, animal,
plant, fungi) removed against repeat-masked references (minid=0.95 maxindel=3 bwr=0.16
bw=12 quickmatch fast minhits=2 qtrim=rl trimq=10 untrim). Reads mapping to rRNA were
removed using sortmerna v4.1.0 [94] against the pre-built sortmerna databases and the

organism-specific rRNA sequences. Reads were mapped against the reference genome
generated in this study using bowtie2 v2.4.4 [86], Q>30 filtered and index-sorted using samtools
v1.9 [68]. Reads were counted using featureCounts of the subread-package v2.0.0 [95] in single-
end mode (-s 2) or in paired-end mode (-s 2 -p -B -C) against the "gene" features of the
reference genomes' gff-files.

To show transcription levels along the chromosome, trimmed reads were mapped
against the reference chromosomes using bowtie2 v2.4.4 [86], and index-sorted using samtools
v1.9 [68]. Per-basepair coverage was determined using samtools depth (-a -d 0) [68], and binned
in 5 kb bins using using a custom script in R v3.6.1 [65]. Counts were normalized to the sum of
counts per sample and square-root transformed. All plots were made using matplotlib v3.5.1
[96] in python v3.7.12 [97].

For differential expression analysis, raw per-gene counts and metadata (i.e.,
conditions) were imported using DESeqDataSetFromMatrix in DESeq2 package v1.30.1 [98] in
R v4.0.3 [65], and differential expression analysis carried out using the function 'DESeq', and
results extracted using the function 'results' with padj.cutoff = 0.05 and lfc.cutoff = log2(2). For
each sample, raw per-gene counts were divided by gene length, and TPM values calculated
by multiplying each normalized count value by $10^6$, divided by the sum of all counts per
sample. Per-sample TPM values were averaged per condition. Genes were classified into
expression level categories by considering the distribution parameters as indicated in
Supplementary Fig. 10.

1017

1018

1019
1020
**Figure 1. DNA localization in *C. steedae*, *A. filiformis* and *S. muelleri* cells.** Schematic representations and confocal microscope images of *C. steedae* (a), *A. filiformis* (b) and *S. muelleri* labeled with the peptidoglycan precursor TADA for 60 min (green) and stained with Hoechst 33342 (magenta). (a) Viewed from the top, *C. steedae* DNA is either homogenously distributed (cell #1), excluded from midcell (cell #2) or X-shaped in cells undergoing late septation stages (cell #3). Rightmost panel displays a side view of cells 1-3. (b) Frontal view of an *A. filiformis* filament reveals a V-shaped DNA localization pattern. Rightmost panel displays a 90 degrees rotated view of the cells marked with an asterisk. (c) Viewed from the top, *S. muelleri* DNA is either excluded from midcell (cell #1), displaying a X-shaped pattern in cells undergoing late septation stages (cell #2) or homogenously distributed (cell #3). Rightmost panel displays a side view of cells 1-3. Scale bars are 1 μm.

1032

**Figure 2. Stable *ori-ter-ter-ori* chromosome configuration in the diploid *C. steedae* and**

***S. muelleri.*** (a, j) Histograms showing number of *ori* or (d,j) *ter* foci per cell. Confocal

microscope images of representative filaments of *C. steedae* (b,e) and *S. muelleri* (h,k)

subjected to either *ori* DNA FISH (b,h) or *ter* DNA FISH (e,k); phase contrast images are

displayed in upper panels and corresponding FISH (green) and DNA (magenta) fluorescence

images in bottom panels. (c,i) Subcellular localization of 215 *ori* foci in *C. steedae* and of 188

*ori* foci in *S. muelleri*. (f, j) Subcellular localization of 236 *ter* foci in *C. steedae* cells and of 293

*ter* foci in *S. muelleri*. (m) Model of *ori-ter-ter-ori* chromosome configuration and lateral

chromosome segregation in the diploid crescent-shaped *S. muelleri* and *C. steedae*. Scale

bars are 1 µm.

1044

1045
1046

**Figure 3. Stable *ori-ter* configuration in the monoploid *A. filiformis.*** (a) Histograms

showing number of *ori* foci (n = 748) and (d) *ter* foci (n = 447) per cell. (b,e) Confocal

microscope images of representative filaments of *A. filiformis* subjected to *ori* DNA FISH (b)

or *ter* DNA FISH (e); phase contrast images are displayed in upper panels and corresponding

FISH (green) and Hoechst 33342 (magenta) fluorescence images in bottom panels. (c,f)

Subcellular localization of 748 *ori* foci and 447 *ter* foci in *A. filiformis*. (g) Histogram showing

number of ParB foci, (h) epifluorescence microscope image of an *A. filiformis* filament

immunostained with anti-ParB antibody and (i) subcellular localization of 611 ParB foci. (j)

Model of *ori-ter* chromosome configuration and lateral chromosome segregation in the

monoploid *A. filiformis*. Scale bars are 1 µm.

1057

1058

1059

1060

**Figure 4. Normalized chromosome contact maps of *C. steedae*, *A. filiformis* and *S. muelleri* grown on solid medium.** (a, b, c) *ori*-centered contact maps (green arrows denote *ori*), (d) *dif*-centered *S. muelleri* contact map (red arrows denote *dif*) of cells grown on solid medium, highlighting the loops observed in both replicates (black circles) in the lower half of each contact map. Below each map, the black or red circles show the position of highly expressed genes (HEGs) and rRNA operons if their presence correlates with a CID border. Below each track, the Frontier Index (FI) shows frontiers with upstream (green) or downstream (orange) bias of interaction and the normalized RNA-Seq counts in 5 kb bins show the transcriptomic data obtained in the same conditions that the contact map. The dashed line represents the threshold use to identify highly expressed regions. (e) Detail of (a) showing the *A. filiformis* loop right of *ori*. (f) Detail of (c) showing the *S. muelleri ori* region (left panel) and detail of (c) showing corner peaks of CIDs (right panel). (g) Detail of (d) showing the *S. muelleri* region around *dif*, with CID borders. The remaining replicates are shown in Supplementary Figs. 6-8.

1075

1076

1077

1078

**Figure 5. Normalized chromosome contact maps of *C. steedae*, *A. filiformis* and *S. muelleri* grown in liquid culture.** (a, b, c) *ori*-centered contact maps (green arrows denote *ori*), highlighting the loops observed in both replicates (black circles) in the lower half of each contact map. Below each map, the black or red circles show the position of highly expressed genes (HEGs) and rRNA operons if their presence correlates with a CID border. Below each track, the Frontier Index (FI) shows frontiers with upstream (green) or downstream (orange) bias of interaction and the normalized RNA-Seq counts in 5 kb bins show the transcriptomic data obtained in the same conditions that the contact map. The dashed line represents the threshold use to identify highly expressed regions. (d) Detail of (b) showing the absence of the loop shown in Fig. 4e (dashed circle) and the emergence of a new loop containing the *ori* (closed circle). (e) Detail of (c) showing the absence of many loops (dashed circles) and the emergence of six loops (closed circles) in the vicinity of *S. muelleri* ori. (f) Detail of (c) showing the absence of the two corner peaks shown in Fig.4 f (right panel).

1092

1093

**Figure 6. Circular representations of the *A. filiformis* and *S. muelleri* chromosomes and eggNOG-based annotation of genes contained in looping regions.** (a, b) Circular representation of the *A. filiformis* and *S. muelleri* chromosomes when grown on solid medium (top panels) or in liquid medium (bottom panels), respectively. Looping regions are indicated

as black lines on the white circle. The three inner colored circles represent the position of genes classified as one of the three COG categories 'Cellular Processes and Signaling' (red), 'Information Storage and Processing' (blue), and 'Metabolism' (green). Connections between looping regions (loops) are shown in black. Green arrows indicate the position of *ori*, and the red arrows indicate the position of *dif*.

**Table 1.** Number of frontiers, domain boundaries and co-occurrence with either highly expressed genes (HEGs) or chromosomal loops in different growing conditions.

| | Number of frontiers (up- or downstream peak) | Number of domain boundaries (overlapping up- and downstream peak ± 3 bins) | % of domain boundaries containing highly expressed genes (HEGs) | Number of detected chromosomal loops overlapping with frontiers | Number of detected chromosomal loops overlapping with domain boundaries |
|---|---|---|---|---|---|
| **C. steedae** | | | | | |
| **Solid medium** | 18 | 5 | 80 | NA | NA |
| **Liquid medium (exponential phase)** | 20 | 12 | 92 | NA | NA |
| **Liquid medium (stationary phase)** | 13 | 4 | 100 | NA | NA |
| **A. filiformis** | | | | | |
| **Solid medium** | 16 | 4 | 100 | 1 | 0 |
| **Liquid medium (exponential phase)** | 16*/25** | 7*/13** | 71*/62** | 1*/1** | 0 |
| **Liquid medium (stationary phase)** | 22 | 7 | 29 | 1 | 0 |
| **Solid medium + Rifampicin** | 20 | 5 | n.a. | 1 | 0 |
| **Solid medium + Chloramphenicol** | 15 | 5 | n.a. | 1 | 0 |
| **S. muelleri** | | | | | |
| **Solid medium** | 32*/37** | 11*/16** | 45*/69** | 19*/17** | 12*/18** |
| **Liquid medium (exponential phase)** | 28 | 6 | 67 | 1 | 0 |
| **Liquid medium (stationary phase)** | 28 | 7 | 43 | 2 | 0 |

| | | | | | |
|---|---|---|---|---|---|
| **Solid medium + Rifampicin** | 18 | 2 | n.a. | 1 | 0 |
| **Solid medium + Chlorampheni col** | 30 | 10 | n.a. | 1 | 4 |

1108

1109    *Replicate 1; **Replicate 2

1110

1111    HEGs: Highly expressed genes are genes that are found in regions in which the level of
1112    transcription is classified as Cat_4 (Supplementary Fig. 10 and see main text).

1113 **Table 2.** Number of looping regions, of loops and of genes interacting via loops in *A. filiformis*

1114 *and S. muelleri.*

| | Solid medium (repl. 1) | Solid medium (repl. 2) | **Solid medium (repl. 1 and repl.2 )** | Liquid medium, exponential phase (repl. 1) | Liquid medium, exponential phase (repl. 2) | **Liquid medium, exponential phase (repl.1 and repl.2)** | **Liquid medium, stationary phase** | Solid medium + Rifampicin | Solid medium.+ Chloramphenicol |
|---|---|---|---|---|---|---|---|---|---|
| *A. filiformis* | | | | | | | | | |
| **Number of looping regions** | 2 | 6 | 2 | 2 | 6 | 2 | 4 | 2 | 2 |
| **Average looping region size (kb)** | 7.5 | 5 | 5 | 5 | 8.75 | 5 | 10 | 5 | 5 |
| **Number of loops** | 1 | 3 | 1 | 1 | 2 | 1 | 2 | 1 | 1 |
| **Number of genes interacting via loops** | 22 (1%) | 35 (1.5% ) | 10 (0.5%) | 19 (1%) | 25 (1%) | 12 (0.5%) | 27 (1%) | 12 (0.5%) | 12 (0.5%) |
| *S. muelleri* | | | | | | | | | |
| **Number of looping regions** | 41 | 40 | 25 | 12 | n.a. | n.a. | 11 | 17 | 17 |
| **Average looping region size (kb)** | 11.5 | 11 | 13 | 9 | n.a. | n.a. | 7 | 7.5 | 8 |
| **Number of loops** | 87 | 79 | 36 | 10 | n.a. | n.a. | 7 | 11 | 13 |
| **Number of genes interacti** | 554 (21%) | 510 (20%) | 372 (15%) | 127 (5%) | n.a. | n.a. | 100 (4%) | 158 (6%) | 166 (6%) |

| ng via loops | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| | | | | | | | | | |

1115
1116