Cryo-EM structure of the bacterial effector protein SipA bound to F-actin reveals a unique mechanism for filament stabilization

Emily Z. Guo[1], Steven Z. Chou[2], Maria Lara-Tejero[1], and Jorge E. Galán[1]*

[1]Department of Microbial Pathogenesis, Yale University School of Medicine, New Haven, CT 06536, and [2]Department of Molecular Biology and Biophysics, UConn Health, Farmington, CT 06030

Key words: bacterial pathogenesis, *Salmonella* pathogenesis, actin dynamics, type III protein secretion, host-pathogen interactions

* For correspondence: jorge.galan@yale.edu

ABSTRACT

The bacterial pathogen *Salmonella* spp. modulates cellular processes by delivering effector proteins through its type III secretion systems. Among these effectors, SipA facilitates bacterial invasion and promotes intestinal inflammation. The mechanisms by which this effector carries out these functions are incompletely understood, although SipA's ability to modulate actin dynamics is central to some of these activities. Here, we report the cryo-EM structure of SipA bound to filamentous actin. The structure shows that this effector stabilizes actin filaments through unique interactions of its carboxy-terminal domain with four actin subunits. Furthermore, our structure-function studies revealed that SipA's actin-binding activity is independent of its ability to stimulate intestinal inflammation. Overall, these studies illuminate critical aspects of *Salmonella* pathogenesis and provide unique insight into the mechanisms by which a bacterial effector modulates actin dynamics.

MAIN

Salmonella enterica serovars can cause a range of human illnesses, from self-limiting gastroenteritis (i.e. food poisoning) to systemic, life-threatening infections (i.e. typhoid fever), resulting in an estimated 200,000 annual deaths worldwide (1-8). Central to *Salmonella* pathogenesis is their ability to invade intestinal epithelial cells and cause intestinal inflammation (4, 5, 7, 8). Epithelial cell invasion is orchestrated by the combined activity of several effectors of a type III protein secretion system (T3SS) encoded within its pathogenicity island 1 (SPI-1) (9-12). These effectors trigger cellular responses that result in the reorganization of the actin cytoskeleton, leading to the formation of large lamellipodia-like structures or membrane ruffles that internalize the bacteria into a membrane-bound compartment, the *Salmonella*-containing vacuole.

Three of these effector proteins, SopE, SopE2, and SopB, play a central role in the internalization process by stimulating Rho-family GTPases, which initiate downstream signaling leading to the WASH/WAVE-dependent activation of endogenous actin nucleators, such as the Arp2/3 complex (13-18). SopE, and its close homolog SopE2, directly target the Rho-family GTPases Rac1 and Cdc42, acting as potent exchange factors (13, 19). In contrast, SopB activates endogenous cellular exchange factors by fluxing phosphoinositides through its phosphoinositide-phosphatase activity (15).

Two other effectors, SipC and SipA, contribute to bacterial invasion by directly targeting actin (20, 21). Besides its role in T3SS-mediated protein translocation (22), SipC can nucleate and bundle actin filaments (21). A SipC mutant defective in actin-binding but proficient in T3SS-mediated protein translocation, showed a relatively small but significant defect in bacterial internalization (23), indicating an accessory but important role in bacterial cell entry. SipA plays a complex role in *Salmonella* pathogenesis. Through its carboxy-terminal actin-binding domain, SipA contributes to bacterial entry by reducing the critical concentration of actin required for its

polymerization (20, 24). Furthermore, SipA stabilizes actin filaments and prevents the depolymerizing action of F-actin-severing proteins such as ADF/Cofilin and gelsolin (25). Consequently, the absence of SipA results in a reduced size of the bacterial-stimulated actin ruffles and a decrease in epithelial cell invasion (20, 25). In addition, through its amino-terminal domain, SipA modulates the intestinal inflammatory response to *Salmonella* by mechanisms that are not yet fully understood (26, 27). However, it remains unclear whether the actin-binding activity of SipA is related to this additional function since, despite previous efforts (28), the lack of high-resolution structural information on the SipA-actin interface has hampered the construction of SipA mutants that are essential to clarify this issue.

Similarly, due to the lack of structural information, the mechanisms by which SipA modulates actin dynamics are poorly understood. Although the crystal structure of the actin-binding domain of SipA in isolation is available (29), its atomic structure bound to F-actin is not. Previous low-resolution negative stained cryo-EM structures led to the proposal that SipA "staples" actin filaments through the deployment of two non-globular arms that reach out to opposing actin strands, presumably engaging actin in a similar manner to nebulin (29, 30). However, the lack of high-resolution structures has prevented the rigorous testing of such a model. Furthermore, machine learning–based approaches, such as that afforded by Alpha Fold (31, 32), are inadequate to predict the structure of SipA bound to F-actin. In this study, we report the cryo-EM structure of SipA bound to F-actin at 3.6 Å resolution. Our structure, validated by functional studies, clarifies the function of SipA, reveals a unique mechanism of F-actin stabilization by an actin-binding protein, and provides unique insight into the mechanisms of *Salmonella* invasion.

Results

SipA binding changes the symmetry of the decorated filaments

The *S.* Typhimurium effector protein SipA is 685 amino acids in length and is composed of at least three domains: an N-terminal chaperone-binding domain, a central ~200 amino acid domain modeled by AlphaFold as long unstructured loops, and a C-terminal actin-binding globular domain (Supplementary Fig. 1). We first attempted to purify full-length SipA using different tags and purification conditions. However, consistent with previous reports (29), all constructs containing the central region were insoluble or unstable. Therefore, to study the actin-stabilizing function of SipA, we concentrated our efforts on its C-terminal domain (amino acids 447 to 685), which in previous studies was found to contain all its actin-modulating activities (20, 29). Consistent with previous reports (20, 29), we found that purified $SipA_{447-685}$ interacts with skeletal muscle actin with high affinity (0.54 µM) as determined by a co-sedimentation assay (Supplementary Fig. 2). Also, consistent with previous observations (20), negatively stained actin filaments polymerized in the presence of SipA observed under the electron microscope showed a thicker and straighter appearance than filaments polymerized in its absence (Supplementary Fig. 3).

To determine the structure of the SipA-actin complex, purified G-actin was first polymerized to form filaments, mixed with purified $SipA_{447-685,}$ and examined by single particle cryo-EM. We collected 4,740 movies, of which 2,370 were used for manual filament picking (see Methods). In total, 278,740 filament segments were boxed out for further analysis. The helical parameters of $SipA_{447-685}$-decorated actin filaments were determined by helical indexing of the averaged power spectrum. The rise and translation of actin subunits along the helix of bare actin filaments is about 27.4 Å. Notably, the rise of SipA-decorated actin filaments is 109.9 Å, roughly four times longer than that of bare actin filaments. The rise line (the first layer line that crosses the meridian), although weak, is visible in the averaged power spectrum (Supplementary Fig. 4). By applying helical symmetry in the EM reconstruction, we obtained a map with extra densities (colored in magenta and yellow in Fig. 1) alongside the grooves between the two strands of the actin filaments (colored light green and dark green in Fig. 1), which were not present in actin filaments alone and therefore assumed to correspond to SipA (Fig. 1). Notably, in 2D class

averages, the SipA densities were much weaker than that of the actin filaments and displayed low resolution and poor map connectivity in initial 3D reconstructions (Supplementary Fig. 5). These findings indicate a sub-stoichiometric presence of SipA on the actin filaments. We processed the particles with SipA binding at both grooves and generated a final map with a global resolution of 3.6 Å determined by the Fourier shell correlation (FSC) at 0.143 cutoff (Supplementary Fig. 6). To improve the map connectivity and interpretability, DeepEMhancer (33) was applied to post-process the refined 3D map. Local resolution analysis revealed that the cryo-EM map for the actin filament exhibits a higher resolution of approximately 3.5 Å, whereas the SipA map displays a lower resolution, ranging from 3.5 Å to 5.5 Å (Supplementary Fig. 7). This map showed densities corresponding to the side chains of most actin residues, as well as the bound nucleotide and the associated Mg2+ in polymerized actin (Supplementary Fig. 8).

The density corresponding to $SipA_{447-685}$ contains a compact globular domain with dimensions (30 Å x 40 Å x 40 Å) that correspond well with its previously solved crystal structure (PDB ID: 1Q5Z) (29) (Fig. 1), and a poorly resolved non-globular extension at the C-terminus that inserts deeply into F-actin (Supplementary Fig. 9). However, contrary to what was previously reported (29), we did not observe the N-terminal non-globular protrusion or arm, even though the same SipA construct was used in both studies.

SipA binds to actin filaments with a stoichiometry of 2:4

The atomic model of the SipA-F-actin complex was built by docking the crystal structure of SipA (PDB ID: 1Q5Z) (29) and the Mg-ADP-actin filament (PDB ID: 6DJON) (34) onto the reconstructed EM map using Chimera (35) and several rounds of model building and refinement with COOT (36) and Phenix (37). Each asymmetric unit of the SipA-decorated actin filaments contains four actin subunits and two SipA subunits (Figure 2A, actin subunits are colored in light green and dark green, and SipA subunits are colored in magenta and yellow). As a result, SipA binds to actin filaments with a stoichiometry of 2:4 (Supplementary Movie 1). A globular domain that

encompasses residues 514-657 and a C-terminal extension that comprises residues 658-675 of SipA could be resolved in our map (Fig. 2A). The absence of the N-terminal region (residues 447-513) in our structure suggests that this region is flexible, aligning with earlier observations that the central domain of SipA, situated in immediate proximity to the amino-terminal end of its actin-binding domain, lacks a defined structure, which was confirmed by AlphaFold modeling (31) (Supplementary Fig. 1).

The alignment of the globular domain of SipA within our cryo-EM structure with its crystal structure (PDB ID: 1Q5Z) highlighted minimal conformational changes induced by actin binding, evident in a Cα root mean square deviation (RMSD) over residues 514-658 of 0.843 Å (Fig. 2B). Only minor local conformational changes in SipA were observed at its actin-binding surface. Similarly, the actin subunit in our structure and a previously solved EM structure of ADP-actin filaments (PDB ID: 6DJO) showed minimal differences (backbone RMSD: 0.610 Å) (Fig. 2C). Minor local conformational changes were observed at the actin-SipA interfaces, particularly involving the D-loop region. No significant conformational alterations were observed in the bound nucleotide, ADP, the associated cation, Mg^{2+}, or the surrounding residues of the actin subunit (Fig. 2C). The actin filament in our structure aligns well with the published Mg-ADP-actin filament (PDB ID: 6DJO), suggesting that the binding of SipA does not alter the assembly of actin subunits in this filament (Fig. 2D), but only changes its helical symmetry.

In addition to the globular domain, we observed a poorly resolved carboxy-terminal extension of SipA, which inserts deeply into the actin filament (Fig. 3A and 3D and Supplementary Fig. 9). We established that, as previously shown (29), this extension is required for the ability of SipA to polymerize G-actin (Supplementary Fig. 10A and 10B) although it appears dispensable for its ability to stabilize F-actin (Supplementary Fig. 10C and 10D). Furthermore, consistent with the functional importance of this extension, its deletion also affected the ability of *S*. Typhimurium to invade cultured epithelial cells (Supplementary Fig. 10E).

Each SipA molecule binds to four actin subunits on either side of the filament

The two strands of the actin filament create grooves running along opposite sides of the filament axis. Each SipA molecule binds to four actin subunits on either side of the two actin strands. The interface between the globular domain of SipA and the actin filament is unprecedented in that a single globular domain engages four actin subunits on both strands (N, N+1, N+2, and N+3) (Fig. 3A) through hydrophobic interactions and ionic bonds. Its interface with the actin N subunit is relatively small (~155 $Å^2$), involving residues in the SD4 domain in actin (Fig. 3B, interface residues are highlighted in magenta) and in helix 4 (H4) and loop 4 (L4) in $SipA_{447-685}$ (Fig. 3C and 3D, interface residues are highlighted in cyan)(all the interface areas were calculated by PDBePISA). The interface between SipA and actin N+1 is large and buries ~746 $Å^2$. This interface is stabilized by more substantial interactions involving residues in the SD1 and SD2 loops of the actin subunit (Fig. 3B, interface residues are highlighted in magenta) and H1, L2, H2, L3, H3, L4, H4, and L5 in $SipA_{447-685}$ (Fig. 3C and 3D, interface residues are highlighted in blue). The interface between SipA and actin N+2 is also substantial, burying 531.6 $Å^2$ and involving actin residues mostly in the helices of SD3 and SD4 domain (Fig. 3B, interface residues are highlighted in magenta), engaging residues in L2, H1, L2, H2, L7, and H7 in $SipA_{447-685}$ (Fig. 3C and 3D, interface residues are highlighted in orange). Finally, the interface between SipA and actin #N+3 is mainly between the loops of the SD1 and SD3 domain in actin (Fig. 3B, interface residues are highlighted in magenta), and H2, L3, H7 and L8 in $SipA_{447-685}$ (Fig. 3C and 3D, interface residues are highlighted in green) and buries ~214 $Å^2$. Overall, the combined interface between the globular domain of SipA and F-actin is substantial thus explaining its relatively high affinity for F-actin.

Comparison of SipA with other actin-binding molecules

Because of their affinities for polymerized actin, the fungal toxins phalloidin and jasplakinolide are frequently used to visualize actin filaments. SipA targets actin at a similar site to where these fungal toxins bind filamentous actin (Fig. 4A and 4B). However, superimposition of the structures

of SipA-decorated actin filament with the toxin-actin complexes, revealed that the C-terminal extension domain of SipA would clash with the two toxin molecules, indicating the binding of SipA may block the binding of phalloidin and jasplakinolide (Fig. 4A and 4B). Pyrene labeling has been widely employed to measure the kinetics of actin polymerization and the interaction between actin and actin-binding proteins, including SipA (20, 29). Notably, pyrene binds to actin close to the SipA-binding site (38). Alignment of our structure with the structure of pyrene-bound actin filament (PDB ID:7K21) shows that the binding of pyrene alters the conformation of the D-loop region of the actin subunit in a manner that would clash with SipA. More specifically, V45 on the D-loop region of actin clashes with H539 of SipA (Fig. 4C). These findings therefore suggest that actin polymerization assays employing pyrene actin may not be suitable for studying the effect of SipA on actin dynamics.

The actin-binding site of SipA also overlaps with the binding site of other well-characterized actin-regulatory proteins. For example, SipA binds the same groove in F-actin that serves as a binding site for the muscle protein nebulin (Fig. 4D) (39, 40). Based on low-resolution structures of SipA and Nebulin bound to actin, it was previously proposed that SipA mimics nebulin in its mechanism of actin-binding (30). Our structure is consistent with this notion since several actin residues engaged by SipA (e.g. K68, E117, E224, E276, and R372) are also engaged by nebulin (41).

The Arp2/3 complex forms branches by nucleating the daughter filament from the side of the mother actin filaments (42). Superimposition of our structure and the previous structure of the actin branch junction (PDB ID: 8E9B) (43) shows that SipA and Arp3 bind at the same site on actin filaments (Fig. 4E). Thus, binding of SipA likely inhibits the binding of Arp2/3 complex thus potentially preventing the branching of actin filaments.

It was previously reported that SipA interferes with the F-actin severing activity of ADF/cofilin (25). Although SipA and cofilin (44) bind actin at different sites, the SipA-bound filaments have different helical symmetries therefore potentially interfering with cofilin binding. In

fact, superimposing the SipA- and cofilin-bound (5YU8) filamentous actin structures using the inner domain (subdomains 3 and 4) of one actin subunit shows that Cofilin would clash with the D-loop region and the C-terminal loop (S348-F352) of actin in its SipA-bound configuration (Fig. 4F). These observations provide a plausible mechanistic explanation for the observation that actin filaments bound to SipA are resistant to the severing activity of ADF/cofilin (25).

Bacterial actin homologs lack structural features that are essential for SipA binding

Many bacteria, including *Salmonella*, encode highly conserved cytoskeletal proteins such as MreB and ParM, that are structural homologs of actin (45, 46). However, despite the close structural similarities between mammalian actin and the bacterial counterparts, SipA should have evolved to be unable to target these bacterial actin homologs as such binding would likely be detrimental to bacterial physiology and could potentially trap SipA within the bacterial cytoplasm. Consistent with this notion, comparison of the structures of MreB and ParM with actin shows that several actin structural features that are crucial for the formation of its interface with SipA are partially or totally absent from MreB and ParM (Fig. 5). These results indicate that, as previously suggested (30), SipA has evolved to avoid binding to the bacterial actin homologs.

Structure-function analyses of SipA indicate that its actin-modulating and pro-inflammatory activities are independent of one another

SipA has been associated with two different phenotypes, bacterial invasion of intestinal epithelial cells through its actin-modulating activity (20), and the stimulation of intestinal inflammation through poorly understood mechanisms that involve its amino-terminal half (26, 27, 47). Whether the ability of SipA to stimulate intestinal inflammation requires its actin-modulating activity is unclear since the lack of structural information has prevented the generation of a mutant specifically defective in this function while preserving the proper folding and integrity of the entire protein. This issue is pertinent since it has been shown that actin polymerization, when triggered

by microbial pathogens, can activate innate immune signaling (48). To address this issue, we introduced mutations in SipA residues that are implicated in the formation of its interfaces with actin and examined their effect on F-actin-binding and G-actin polymerization. Reflecting the redundant nature of the SipA-actin interface, none of the individual mutations introduced significantly affected the ability of SipA to bind actin (Fig. 6A). However, introduction of changes in two or more SipA residues involved in forming its interface with actin significantly reduced its binding affinity. Notably, simultaneous mutations in S537 and K636, which interact with the D-loop and C-terminus of actin (Supplementary Fig. 11), resulted in a drastic reduction in its binding affinity (Fig. 6A).

To examine the contribution of the actin-binding activity of SipA to the interaction of *S.* Typhimurium with host cells, we constructed a mutant strain expressing the *sipA*$^{S537E/K636E}$ allele from its native chromosomal context. We then tested the resulting strain's ability to stimulate actin-cytoskeletal rearrangements and invade cultured epithelial cells. We found that, in comparison to the wild type, the actin cytoskeletal changes stimulated by the *S.* Typhimurium *sipA*$^{S537E/K636E}$ mutant strain were more diffuse, with less defined ruffles (Fig. 6B). Consistent with the more limited actin rearrangements, the invasion ability of the mutant strain was significantly impaired when measured with two independent assays (Fig. 6C-6E). Notably, the phenotype of the *S.* Typhimurium *sipA*$^{S537E/K636E}$ mutant strain was similar to the phenotype of the *S.* Typhimurium Δ*sipA* mutant, further demonstrating that the introduced mutations largely eliminated its ability to bind actin (Fig. 6C-6E).

To investigate the role of SipA's actin-modulating activity in promoting inflammation, we compare the intestinal inflammatory response of mice infected with different *S.* Typhimurium strains. Specifically, we compared mice infected with a strain lacking *sipA*, a strain expressing the actin-binding defective mutant SipA$^{S537E/K636E}$, and a strain expressing wild-type SipA. To enhance the detection of SipA-dependent inflammation (27), we introduced a Δ*spiA* mutation in all strains, rendering them deficient for the SPI-2 T3SS. We found that mice infected with *S.* Typhimurium

expressing either wild type SipA or the actin-binding defective mutant displayed smaller cecum sizes and lighter weights indicative of reduced cecal content due to inflammation (Fig. 7A and 7B). In addition, these mice exhibited substantial intestinal inflammation, as observed by histopathology (Fig. 7C). In contrast, mice infected with the *sipA*-deficient *S*. Typhimurium strain exhibited cecum sizes similar to uninfected controls (Fig. 7A and 7B) and much reduced intestinal inflammation (Fig. 7C). These results indicate that the actin-binding activity of SipA is not necessary for inducing intestinal inflammation. Thus, the actin-modulating and pro-inflammatory activities of SipA are independent of each other.

Discussion

Bacterial effector proteins that can modulate cellular processes are often evolutionary designed to mimic the activities of eukaryotic proteins that carry out a similar function (49, 50). This mimicry often involves convergent evolution since many of these bacterial mimics do not exhibit either primary sequence or structural similarity to their eukaryotic counterparts (49). The actin-binding effector protein SipA has been postulated to mimic the actin-binding protein nebulin (30). In addition, it has been proposed to exert its actin-stabilizing function by working as a "molecular staple", using two flexible arms to bind actin subunits on opposing actin strands (29). However, the lack of a high-resolution structure of SipA bound to actin filaments has prevented the rigorous testing of these hypotheses. To address these issues, we solved the cryo-EM structure of the actin-binding carboxy-terminal domain of SipA bound to actin filaments. We found that each SipA molecule simultaneously engages four actin subunits through interactions involving only its globular carboxy-terminal domain. The SipA binding sites on actin overlap those of nebulin, a very large muscle skeletal protein, which through the tandem arrangement of multiple actin-binding domain repeats can engage up to 200 actin subunits (39). Our structure showed that SipA and each nebulin repeat bind the same groove in the actin filaments and share some actin residues

necessary for their interaction. This observation is therefore consistent with the notion that SipA and the nebulin repeat exhibit a similar actin-binding mode.

Our structure also showed that the SipA actin-binding site overlaps with the binding site of a well-characterized actin nucleator, the Arp2/3 complex. This observation has potential functional relevance since it suggests that SipA is likely to interfere with the binding of the Arp2/3 complex, thus potentially preventing the branching of SipA-stabilized actin filaments. However, more experiments will be required to test this hypothesis and to examine its potential functional significance.

It has been previously reported that SipA prevents the activity of the actin-severing proteins ADF/cofilin (25), resulting in further stabilization of SipA-bound filaments, enhanced membrane ruffles and bacterial internalization. Our structure provides a plausible mechanistic explanation for such activity. The actin-binding sites of SipA and ADF/Cofilin do not overlap (44). However, SipA-bound filaments exhibit a different helical symmetry from those bound by cofilin, which likely interferes with its binding. In fact, comparison of the structures of SipA- and cofilin-bound filaments predicts clashes between the D-loop region and the C-terminal loop (S348-F352) of actin in its SipA-bound configuration and cofilin potentially explaining the inhibitory activity.

Our structure of SipA bound to actin did not reveal the presence of non-globular N-terminal arms engaging actin subunits on the opposite strand of the actin filaments, as observed in a previous low-resolution cryo-EM structure (29). Thus, we could not confirm the "molecular staple" model. However, our structure did show an unstructured extension at the C-terminal end, which inserts deep into the actin filament. Furthermore, as previously shown(29), we found that this extension is required for the ability of SipA to increase the efficiency of bacterial invasion of cultured epithelial cells. This extension most likely contributes to bacterial invasion by facilitating the polymerization of G-actin, rather than by stabilizing F-actin, as removing this extension affected the former but not the latter.

In addition to its ability to modulate actin dynamics to mediate bacterial invasion, SipA has also been implicated in the stimulation of intestinal inflammation (26, 27), which is critical for *S*. Typhimurium virulence (51). Although the mechanisms by which SipA modulates the inflammatory response are poorly understood, this activity has been largely correlated with its amino-terminal domain (26, 27, 47). However, as the modulation of actin dynamics by bacterial pathogens has also been linked to the stimulation of innate immune responses involving inflammation (48), it has been unclear whether the carboxy-terminal, actin-binding domain of SipA contributes to this phenotype. Indeed, the lack of structural information for the SipA-actin interface has hampered previous efforts to obtain a mutant that would specifically lack this activity without altering the overall structure of this effector protein (28). This issue is complicated by the redundant nature of the SipA-actin interface. In fact, a previously reported mutant utilized in a study to attempt to address this issue (28) in our hands retains the ability to modulate actin dynamics (Fig. 6). Using structurally guided mutagenesis, we were able to identify a mutant that is specifically defective in actin binding without altering the overall conformation of SipA. As predicted by its significantly decreased ability to bind actin, an *S*. Typhimurium strain expressing this mutant stimulated more diffuse actin cytoskeleton rearrangements, less prominent ruffles, and was defective in bacterial invasion of cultured epithelial cells. In fact, these phenotypes were similar to those of a *S*. Typhimurium strain lacking *sipA*. To address the potential contribution of the actin-binding activity to the stimulation of intestinal inflammation, we tested the *S*. Typhimurium strain expressing the actin-binding defective SipA mutant for its ability to stimulate intestinal inflammation in a mouse model of infection. Surprisingly, we found that the *S*. Typhimurium strain expressing the SipA mutant was able to stimulate increased inflammation in comparison to the wild type. These results indicate that the actin-binding activity of SipA does not contribute to its ability to induce inflammation.

Taken together, our structure function studies have revealed a unique mechanism by which a bacterial effector protein modulates actin dynamics. Indeed, the ability of a protein to bind

four actin subunits with a single globular domain has not been previously reported for other actin-binding proteins. Furthermore, our studies have clarified the mechanism of action of this effector protein and provide unique insight into the *Salmonella* spp. virulence mechanisms.

Materials and Methods

Bacterial strains and growth conditions

The wild-type *Salmonella enterica* serovar Typhimurium SL1344 strain and its ∆*sipA* or ∆spiA (SPI-2 T3SS deficient) isogenic mutants have been previously described (20, 52). The *S.* Typhimurium SL1344 derivative encoding SipA[S537E/K636E] was constructed by allelic exchange as previously described (53). All bacterial strains were cultured in L-Broth containing 0.3 M NaCl to an OD600 of 0.9, conditions known to stimulate the expression of the SPI-1 T3SS (54).

Plasmid construction, protein expression, and purification.

DNA manipulations were carried out using the standard Gibson assembly protocol (55). *S.* Typhimurium SipA$_{447-685}$ was expressed in a modified pQE vector with GST and a 6xHis tag at its amino and carboxy-terminal ends, respectively. SipA$_{447-685}$, SipA$_{498-659}$, and SipA$_{513-659}$, as well as SipA$_{447-685}$ carrying specific point mutations, were expressed in *Escherichia coli* BL21 (DE3) and purified using affinity chromatography. Briefly, bacterial cells were cultured in LB medium at 37°C to an absorbance at OD$_{600}$ of 0.8 to 1.0, induced with 0.2 mM isopropyl β-d-1-thiogalactopyranoside (IPTG) and grown for an additional 16 to 20 h at 20°C. The bacteria were harvested by centrifugation and lysed using a One-Shot cell disrupter following the manufacturer's instructions (Constant Systems Ltd., Daventry, United Kingdom). Cell lysates were cleared by centrifugation, and the supernatant was loaded onto a Glutathione Sepharose 4B gravity-driven column. After elution, the GST tag was removed by digestion with HRV 3C protease. The digested protein mixture was further purified by passing through a second Glutathione Sepharose 4B and a Ni^{2+}–nitrilotriacetic acid column.

Actin was purified as previously reported (56). Briefly, actin was purified from chicken muscle acetone powder made from flash-frozen chicken (purchased from a local Trader Joe's store). After one cycle of polymerization and depolymerization, G-actin was isolated by gel

filtration through Sephacryl S-300 column in Ca-G-buffer (2 mM Tris·HCl, pH 8.0; 0.2 mM ATP; 0.1 mM $CaCl_2$; 1 mM NaN3; 0.5 mM DTT).

Cell culture, bacterial infections, and visualization of the actin cytoskeleton

All cell lines were routinely tested for the presence of mycoplasma by a standard PCR method, checked for their morphological features, growth characteristics and functionalities, but were not authenticated by short tandem repeat profiling. The ability of *S*. Typhimurium strains to invade cultured human intestinal epithelial Henle-407 cells (obtained from the Roy Curtiss III collection in 1987) using the gentamicin protection assay was carried out as described previously (57). Briefly, ~24 h after seeding onto tissue culture plates, cells were infected for 20 min with the indicated *S*. Typhimurium strains with an m.o.i. of 20. Infected cells were washed and incubated for 2 h in DMEM containing gentamicin (100 µg ml^{-1}) to kill extracellular bacteria. Alternatively, bacterial invasion was also measured using a differential staining protocol (inside-out staining) as described before (58). Briefly, cells were infected as indicated above and 20 min after infection, the cells were fixed in formalin and stained with rabbit anti-*Salmonella* lipopolysaccharide (LPS) antibody (generated at Poconos farms), followed by fluorescently labeled (Alexa Fluor 488) secondary anti-rabbit antibody. Cells were subsequently permeabilized with Triton X 100 (0.3 %) and stained with the anti-Salmonella LPS antibody followed by a fluorescently labeled (Alexa Fluor 594) anti-rabbit secondary antibody. In this protocol, extracellular bacteria are stained by both secondary antibodies, while intracellular bacteria are only stained by the Alexa Fluor 594-labeled secondary antibody. Internalized bacteria were quantified by capturing a minimum of 30 fields per condition (containing a minimum of 1000 cells each) on a Nikon Eclipse inverted microscope equipped with an Andor's Zyla 5.5 sCMOS camera.

To visualize actin cytoskeleton rearrangements due to bacterial invasion, cells were infected, fixed, and permeabilized as indicated above and then stained with Atto-488-labeled phalloidin (to visualize the actin cytoskeleton), an anti-*S.* Typhimurium LPS antibody (SIFIN,

catalog no.: TS 1624), and Alexa Fluor 594 fluorescently labeled secondary antibody to visualize bacteria. Cells were visualized under an Andor Dragonfly spinning disk fluorescence microscope (Oxford Scientific).

Mouse infections and measurement of intestinal inflammation

Groups of age- and sex-matched C57BL/6 mice were orally administered streptomycin (100 µl of 200 mg/ml solution) and 24 hrs after antibiotic treatment were orally infected by stomach gavage with 10^7 cfu in 100 µl of PBS with the indicated strains. Forty-eight hours after infection, mice were euthanized and the cecum was collected. Ceca were weighted as a proxy measurement of inflammation and then fixed in formalin and embedded in paraffin for sectioning and H&E staining. Embedding, sectioning, and staining were performed by the Yale Pathology Tissue Services. The sample sizes were empirically determined to optimize the numbers based on our previous experience with equivalent experiments. The mice were randomly assigned to the experimental groups, but experimenters were not blind to the assignment.

EM sample preparation, data acquisition, and image processing

After cation exchange from Ca^{2+} to Mg^{2+} using 0.1 volumes of 10×ME buffer (0.5 mM $MgCl_2$, 2 mM EGTA, pH7.5), Mg-ATP-actin monomers were first polymerized in KMEI buffer (50 mM KCl, 1 mM $MgCl_2$, 1 mM EGTA and 10 mM Imidazole pH 7.0) for 20 min, followed by the addition of SipA with a molar ratio of 1.7:1. The CapZ protein was added into the polymerized actin filaments with a molar ratio of 1:30 to avoid the formation of long actin filaments.

For negative staining, 3.5 µl of actin filaments (final concentration: 3 µM) or SipA (final concentration:5 µM)-decorated actin filaments (final concentration: 3µM) were applied onto glow-discharged carbon-coated grids and negatively stained with 0.8% (w/v) uranyl formate. Images were collected using a Talos L120C electron microscope (Thermo Fisher) operating at 120 kV.

For cryo-EM analysis, 3 μl of the sample was deposited onto glow-discharged Quantifoil R2/1 300 Mesh Gold Holey Carbon Grids (SPI supplies) and plunged into liquid ethane using a Vitrobot Mark IV (Thermo Fisher) at 10°C and 100% humidity. The freezing conditions were screened on a Glacios cryo-TEM (Thermo Fisher) operating at 200 kV. Cryo-EM data collection was performed using a Titan Krios electron microscope (Thermo Fisher) equipped with a K3 direct electron detector (Gatan) operating at 300 kV. Data were collected at a calibrated magnification of 81,000x (1.068 Å per pixel) in super-resolution mode, with defocus values between −2.5 μm and −1.2 μm. Images were recorded using the beam image shift implemented in SerialEM (59). Forty-four frames were taken at 0.05s per frame, with a dose rate of 29 e⁻ per pixel per second. In total, 4740 movies were recorded with an exposure time of 2.2 s. Motion correction and contrast transfer function (CTF) estimation were done using Relion 4.0 (60). Filaments were manually picked by using sxhelixboxer.py in SPARX (61). Layer-line images were calculated with SPARX (project and periodogram commands). In total, 278,740 filament segments were boxed out from 2,370 images and subjected to further 2D and 3D calculations in Relion 4.0. The cryo-EM map of ADP-actin (EMDB accession number: 7938) was used as the initial map and helical symmetry parameters (rise: 109.9 Å and twist: 53.3^{O}) were imposed on the reconstruction. 3D classification resulted in classes with different SipA occupancies. A class with most particles were selected and subjected to additional round of 3D classification. Three classes were combined (69,483 particles) for further 3D refinement. DeepEMhancer (33) was employed to improve map connectivity and interpretability. The final map has a global nominal resolution of 3.6 Å (based on the FSC using the 0.143 criterion calculated in Relion 4.0). The local resolution was calculated with ResMap in Relion 4.0. All the structural calculations were carried out on the Yale High-Performance Computing servers. The statistics for data collection and map reconstruction are summarized in Table S1. The data processing workflow is described in Supplementary Fig. 12.

Model Building and refinement.

The cryo-EM structure of Mg-ADP-actin filaments (PDB ID: 6DJO) (38) and the crystal structure of SipA (PDB ID: 1Q5Z) (29) were used as initial templates to dock into the EM maps using Chimera (35). The modeled structure was manually adjusted in Coot (62) and refined with Phenix (63). The SipA-actin interfaces were calculated with PDBePISA (64). Figures were generated using PyMOL and Chimera. The statistics for model refinement are listed in Table S1.

Actin co-sedimentation assay.

To test the binding affinity of the different SipA mutants, actin monomers were first polymerized into filaments in KMEI buffer (50 mM KCl, 1 mM MgCl2, 1 mM EGTA and 10 mM Imidazole pH 7.0) for 20 min. Then different concentrations (0, 1, 2, 3, 4, and 10 μM) of polymerized actin filaments were incubated with 4 μM of wild-type SipA or SipA mutants for 10 min at room temperature. Followed by ultracentrifugation at 120,000 x g in a TLA100 rotor (Beckman Instruments, Fullerton, CA) at 4°C for 30 min, an equal amount of the supernatant from each reaction was loaded onto SDS–PAGE Gel and stained with Coomassie blue.

To test the actin nucleation activity of $SipA_{447-685}$, $SipA_{498-659}$, and $SipA_{513-659}$, the SipA proteins were first dialyzed against a Tris buffer without salt (25 mM Tris, pH 7.5) three times before adding to G-actin. After 4 hs of incubation, the samples were ultracentrifuged at 120,000 x g in a TLA100 rotor (Beckman Instruments, Fullerton, CA) at 4°C for 30 min, then an equal amount of the supernatant and pellet from every reaction were loaded onto the SDS–PAGE gel and stained with Coomassie blue.

Data Availability.

The EM maps and atomic coordinates have been deposited in Electron Microscopy Data Bank (EMDB) under accession number EMD-43188 and Protein Data Bank (PDB) under accession code 8VFM.

Ethics statement

All animal experiments were conducted according to protocols approved by Yale University's Institutional Animal Care and Use Committee (IACUC) under protocol number 2022–07858. The IACUC is governed by applicable Federal and State regulations, including those of the Animal Welfare Act, Public Health Service and the United States Department of Agriculture and is guided by the US Government Principles for the Utilization and Care of Vertebrate Animals Used in Testing, Research and Training.

Acknowledgements

We thank Dr. Jianfeng Lin, Dr. Shenping Wu, Dr. Marc Llaguno, and Dr. Xinran Liu for managing the microscope resources, the Yale High Performance Computing support team for guidance and use of the computing infrastructure. We also thank Florian Schueder for assistance in the imaging of the actin cytoskeleton rearrangements induced by the different *S*. Typhimurium strains. This work was supported by NIH Grant R01AI055472 to J. E. G.

References

1. A. Voetsch *et al.*, FoodNet estimate of the burden of illness caused by nontyphoidal Salmonella infections in the United States. *Clin Infect Dis.* **38**, S127-134 (2004).
2. C. Parry, E. Threlfall, Antimicrobial resistance in typhoidal and nontyphoidal salmonellae. *Curr Opin Infect Dis.* **21**, 531-538 (2008).
3. J. Crump, E. Mintz, Global trends in typhoid and paratyphoid Fever. *Clin Infect Dis.* **50**, 241-246 (2010).
4. M. E. Ohl, S. I. Miller, Salmonella: a model for bacterial pathogenesis. *Annu. Rev. Med.* **52**, 259-274. (2001).
5. G. Grassl, B. Finlay, Pathogenesis of enteric Salmonella infections. *Curr Opin Gastroenterol.* **24**, 22-26 (2008).
6. C. Parry, T. T. Hien, G. Dougan, N. White, J. Farrar, Typhoid fever. *N Engl J Med.* **347**, 1770-1782 (2002).
7. M. Raffatellu, R. Wilson, S. Winter, A. Bäumler, Clinical pathogenesis of typhoid fever. *J Infect Dev Ctries* **2**, 260-266 (2008).
8. G. Dougan, S. Baker, Salmonella enterica serovar Typhi and the pathogenesis of typhoid fever. *Annu Rev Microbiol.* **68**, 317-336 (2014).
9. J. E. Galán, Salmonella interaction with host cells: Type III Secretion at Work. *Annu. Rev. Cell Dev. Biol.* **17**, 53-86 (2001).
10. D. LaRock, A. Chaudhary, S. Miller, Salmonellae interactions with host processes. *Nat Rev Microbiol.* **13**, 191-205 (2015).
11. P. Hume, V. Singh, A. Davidson, V. Koronakis, Swiss Army Pathogen: The Salmonella Entry Toolkit. *Front Cell Infect Microbiol.* **7**, 348-358 (2017).
12. S. Fattinger, M. Sellin, W. Hardt, Salmonella effector driven invasion of the gut epithelium: breaking in and setting the house on fire. *Curr Opin Microbiol.* **64**, 9-18 (2021).
13. W.-D. Hardt, L.-M. Chen, K. E. Schuebel, X. R. Bustelo, J. E. Galán, *Salmonella typhimurium* encodes an activator of Rho GTPases that induces membrane ruffling and nuclear responses in host cells. *Cell* **93**, 815-826 (1998).
14. C. Bakshi *et al.*, Identification of SopE2, a Salmonella secreted protein which is highly homologous to SopE and involved in bacterial invasion of epithelial cells. *J. Bacteriol.* **182**, 2341-2344 (2000).
15. J. C. Patel, J. E. Galan, Differential activation and function of Rho GTPases during Salmonella-host cell interactions. *J Cell Biol* **175**, 453-463 (2006).
16. D. Zhou, L. M. Chen, L. Hernandez, S. B. Shears, J. E. Galan, A Salmonella inositol polyphosphatase acts in conjunction with other bacterial effectors to promote host cell actin cytoskeleton rearrangements and bacterial internalization. *Mol Microbiol* **39**, 248-259 (2001).
17. D. Humphreys, A. Davidson, P. Hume, V. Koronakis, Salmonella virulence effector SopE and Host GEF ARNO cooperate to recruit and activate WAVE to trigger bacterial invasion. *Cell Host Microbe.* **11**, 129-139 (2012).

18. A. Criss, J. Casanova, Coordinate regulation of Salmonella enterica serovar Typhimurium invasion of epithelial cells by the Arp2/3 complex and Rho GTPases. *Infect Immun.* **71**, 2885-2891 (2003).

19. S. Stender *et al.*, Identification of SopE2 from Salmonella typhimurium, a conserved guanine nucleotide exchange factor for Cdc42 of the host cell. *Mol Microbiol.* **36**, 1206-1211. (2000).

20. D. Zhou, M. Mooseker, J. E. Galán, Role of the *S. typhimurium* actin-binding protein SipA in bacterial internalization. *Science* **283**, 2092-2095 (1999).

21. R. D. Hayward, V. Koronakis, Direct nucleation and bundling of actin by the SipC protein of invasive Salmonella. *EMBO J.* **18**, 4926-4934. (1999).

22. C. Collazo, J. E. Galán, The invasion-associated type III system of *Salmonella typhimurium* directs the translocation of Sip proteins into the host cell. *Mol. Microbiol.* **24**, 747-756 (1997).

23. J. Chang, J. Chen, D. Zhou, Delineation and characterization of the actin nucleation and effector translocation activities of Salmonella SipC. *Mol Microbiol.* **55(**1379-1389 (2005).

24. E. McGhie, R. Hayward, V. Koronakis, Cooperation between actin-binding proteins of invasive Salmonella: SipA potentiates SipC nucleation and bundling of actin. *EMBO J.* **20**, 2131-2139 (2001).

25. McGhie EJ, Hayward RD, K. V., Control of actin turnover by a salmonella invasion protein. *Mol Cell.* **13**, 497-510 (2004).

26. Wall DM *et al.*, Identification of the Salmonella enterica serotype typhimurium SipA domain responsible for inducing neutrophil recruitment across the intestinal epithelium. *Cell Microbiol.* **9**, 2299-2313 (2007).

27. S. Hapfelmeier *et al.*, Role of the Salmonella pathogenicity island 1 effector proteins SipA, SopB, SopE, and SopE2 in Salmonella enterica subspecies 1 serovar Typhimurium colitis in streptomycin-pretreated mice. *Infect Immun.* **72**, 795-809 (2004).

28. D. Li, X. Wang, L. Wang, D. Zhou, The actin-polymerizing activity of SipA is not essential for Salmonella enterica serovar Typhimurium-induced mucosal inflammation. *Infect Immun.* **81**, 1541-1549 (2013).

29. M. Lilic *et al.*, Salmonella SipA polymerizes actin by stapling filaments with nonglobular protein arms. *Science* **301**, 1918-1921. (2003).

30. V. E. Galkin *et al.*, The bacterial protein SipA polymerizes G-actin and mimics muscle nebulin. *Nat Struct. Biol.* **9**, 518-521. (2002).

31. J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589 (2021).

32. R. Evans *et al.*, Protein complex prediction with AlphaFold-Multimer. *bioRxiv* 10.1101/2021.10.04.463034, 2021.2010.2004.463034 (2021).

33. R. Sanchez-Garcia *et al.*, DeepEMhancer: a deep learning solution for cryo-EM volume post-processing. *Commun Biol.* **4**, 874 (2021).

34. S. Chou, T. Pollard, Mechanism of actin polymerization revealed by cryo-EM structures of actin filaments with three different bound nucleotides. *Proc Natl Acad Sci U S A.* **116**, 4265-4274 (2019).

35. T. Goddard *et al.*, UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein Sci.* **27**, 14-25 (2018).

36. P. Emsley, K. Cowtan, Coot: model-building tools for molecular graphics. *Acta Crystallogr.* **D 60**, 2126–2132 (2004).

37. P. Adams *et al.*, PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Cryst.* **D66**, 213-221 (2010).

38. S. Chou, T. Pollard, Cryo-electron microscopy structures of pyrene-labeled ADP-Pi- and ADP-actin filaments. *Nat Commun.* **11**, 5897 (2020).

39. M. Yuen, C. A. C. Ottenheijm, Nebulin: big protein with big responsibilities. *J Muscle Res Cell Motil* **41**, 103-124 (2020).

40. Z. Wang *et al.*, Structures from intact myofibrils reveal mechanism of thin filament regulation through nebulin. *Science* **375**, eabn1934 (2022).

41. Z. Wang *et al.*, Structures from intact myofibrils reveal mechanism of thin filament regulation through nebulin. *375* **375**, eabn1934 (2022).

42. P. Bieling, K. Rottner, From WRC to Arp2/3: Collective molecular mechanisms of branched actin network assembly. *Curr Opin Cell Biol.* **80**, 102156 (2023).

43. S. Chou, M. Chatterjee, T. Pollard, Mechanism of actin filament branch formation by Arp2/3 complex revealed by a high-resolution cryo-EM structureof the branch junction. *Proc Natl Acad Sci U S A.* **119**, e2206722119 (2022).

44. K. Tanaka *et al.*, Structural basis for cofilin binding and actin filament disassembly. *Nat Commun.* **9**, 1860 (2018).

45. J. Shaevitz, Z. Gitai, The structure and function of bacterial actin homologs. *Cold Spring Harb Perspect Biol.* **2**, a000364 (2010).

46. E. Ozyamak, J. Kollman, A. Komeili, Bacterial actins and their diversity. *Biochemistry* **52**, 6928-6939 (2013).

47. M. Raffatellu *et al.*, SipA, SopA, SopB, SopD, and SopE2 contribute to Salmonella enterica serotype typhimurium invasion of epithelial cells. *Infect Immun* **73**, 146-154 (2005).

48. S. Ahrens *et al.*, F-actin is an evolutionarily conserved damage-associated molecular pattern recognized by DNGR-1, a receptor for dead cells. *Immunity* **36**, 635-645 (2012).

49. C. E. Stebbins, J. E. Galan, Structural mimicry in bacterial virulence. *Nature* **412**, 701-705 (2001).

50. N. Elde, H. Malik, The evolutionary conundrum of pathogen mimicry. *Nat Rev Microbiol.* **7**, 787-797 (2009).

51. J. Galan, Salmonella Typhimurium and inflammation: a pathogen-centric affair. *Nat. Rev. Microbiol.* **19**, 716–725 (2021).

52. S. Spano, X. Liu, J. E. Galan, Proteolytic targeting of Rab29 by an effector protein distinguishes the intracellular compartments of human-adapted and broad-host Salmonella. *Proc Natl Acad Sci U S A* **108**, 18418-18423 (2011).

53. K. Kaniga, J. C. Bossio, J. E. Galan, The Salmonella typhimurium invasion genes invF and invG encode homologues of the AraC and PulD family of proteins. *Mol Microbiol* **13**, 555-568 (1994).

54. J. E. Galán, R. Curtiss III, Expression of *Salmonella typhimurium* genes required for invasion is regulated by changes in DNA supercoiling. *Infect. Immun.* **58**, 1879-1885 (1990).

55. D. G. Gibson *et al.*, Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Methods* **6**, 343-345 (2009).

56. S. Z. Chou, T. D. Pollard, Mechanism of actin polymerization revealed by cryo-EM structures of actin filaments with three different bound nucleotides. *Proc Natl Acad Sci U S A* **116**, 4265-4274 (2019).

57. J. E. Galán, R. Curtiss III, Cloning and molecular characterization of genes whose products allow *Salmonella typhimurium* to penetrate tissue culture cells. *Proc. Natl. Acad. Sc. USA* **86**, 6383-6387 (1989).

58. L. M. Chen, S. Hobbie, J. E. Galan, Requirement of CDC42 for Salmonella-induced cytoskeletal and nuclear responses. *Science* **274**, 2115-2118 (1996).

59. D. Mastronarde, Automated electron microscope tomography using robust prediction of specimen movements. *J Struct Biol.* **152**, 36-51 (2005).

60. D. Kimanius, L. Dong, G. Sharov, T. Nakane, S. H. W. Scheres, New tools for automated cryo-EM single-particle analysis in RELION-4.0. *Biochem J* **478**, 4169-4185 (2021).

61. M. Hohn *et al.*, SPARX, a new environment for Cryo-EM image processing. *J Struct Biol* **157**, 47-55 (2007).

62. P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486-501 (2010).

63. P. V. Afonine *et al.*, Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta Crystallogr D Struct Biol* **74**, 531-544 (2018).

64. E. Krissinel, K. Henrick, Inference of macromolecular assemblies from crystalline state. *J. Mol.Biol.* **372**, 774–797 (2007).

Figure 1. Overall structure of SipA$_{447-685}$-decorated actin filaments. Side views of SipA$_{447-685}$ decorated actin filaments. SipA$_{447-685}$ subunits are colored in magenta in one groove and yellow in the other groove, and actin strands are colored in different shades of greens.

Figure 2. Atomic model of SipA$_{447-685}$-decorated actin filament. (**A**) Ribbon diagram of SipA$_{447-685}$-decorated actin filaments. Each asymmetric subunit contains two SipA molecules (magenta and yellow) and four actin subunits (light and dark green). (**B**) Alignment of the globular domain of SipA associated with actin filaments (magenta) to its crystal structure (PDB ID: 1Q5Z) (palecyan). (**C**) Superimposition of the structure of an actin subunit in the SipA-decorated filament to the structure of the bare actin filament (PDB ID: 6DJO). The D-loop region is highlighted with a box. The four subdomains (SD1, SD2, SD3, and SD4), inner domain (ID) and outer domain (OD) are labeled. (**D**) Alignment of actin filament in our structure to the published ADP-actin filament structure (PDB ID: 6DJO).

Figure 3. Each SipA molecule interacts with four actin subunits. (**A**) Ribbon diagram showing that one SipA$_{447-685}$ subunit (magenta) interacts with four actin subunits (N, cyan; N+1, blue; N+2, orange; N+3, green; other subunits: gray). (**B**) SipA-binding interface on the four actin subunits. The actin residues on the actin-SipA interfaces are highlighted in magenta. SipA is hidden for clarity (**C**) Ribbon diagram showing that one SipA$_{447-685}$ subunit (magenta) interacts with four actin subunits. The structure is rotated 180° along the filament axis relative to (**A**) and (**B**). (**D**) Actin-binding interface on SipA$_{447-685}$. SipA$_{447-685}$ is in the same orientation as in (**C**). Secondary structure elements are labeled, and the interacting residues are colored according to the colors of interacting actin subunits in (**C**).

Figure 4. Comparison of the actin-binding interface of SipA with other actin binding molecules. (**A-C**) Superimposition of the structures of SipA and phalloidin (PDB ID: 7BTI) (**A**), jasplakinolide (PDB ID: 6T23) (**B**), or pyrene (PDB ID: 7K20) (**C**) bound to actin. (**D**) Structure of Nebulin (PDB ID: 7QIM) bound to actin filaments showing that nebulin and SipA bind to the same groove of the actin filament. (**E**) Structure of the Arp2/3 complex bound to actin filaments (PDB ID: 8E9B) depicting that SipA shares the same binding site. (**F**) Cofilin clashes with the actin D-loop region and the C-terminal loop (S348-F352) in the SipA-decorated actin filament. The inner domain (subdomains 3 and 4) of an actin subunit is used for structure superimposition. SipA, magenta; actin filament, light gray; phalloidin, green; jasplakinolide, yellow; nebulin, cyan; Arp2/3 complex, orange; cofilin, blue.

Figure 5. SipA binding sites are absent from the bacterial actin homologs MreB and ParM. (**A**-**B**) Alignment of MreB (tan) (**A**) or ParM (light purple) (**B**) to an actin subunit bound to SipA (white). Residues at different SipA-actin interfaces are highlighted following the same color scheme as in Figure 3A. The loops and helices of actin not present in ParM or MreB but involved in the SipA-actin interactions are highlighted. The four subdomains (SD1, SD2, SD3, and SD4), inner domain (ID) and outer domain (OD) of actin subunit are labeled.

Figure 6. Structure-function analysis of the SipA-actin interaction. (**A**) Determination of the binding affinity of different SipA constructs to actin filaments with a co-sedimentation assay. The SipA mutations are on their binding interfaces. The bars are the average dissociation constants calculated from three independent experiments, and the errors are the standard deviations. (**B**) Actin cytoskeletal rearrangements observed after infection with wild-type *S*. Typhimurium (left), a Δ*sipA* isogenic mutant (middle), and a strain expressing the SipA$^{S537E/K636E}$ mutant (right), which showed decreased binding affinity to actin. Actin filaments were stained with Atto 488-phalloidin

(green), bacteria were stained with an antibody directed to the *S*. Typhimurium LPS (red), and mammalian cell nuclear DNA was stained by 4',6-diamidino-2-phenylindole (DAPI) (blue). (**C-E**) Cultured epithelial cell invasion of wild type *S*. Typhimurium, its isogenic Δ*sipA* mutant or a mutant expressing SipA$^{S537E/K636E}$, which is defective for actin binding. The invasion ability of different strains was measured by the gentamicin protection (**C**) or differential (inside-out) staining (**D** and **E**) assays. Values in (**C**) are the ratio of the inoculum that survived antibiotic treatment due to internalization, normalized to wild type (w. t.), which was set to 1. The mean was calculated from six experiments (***: P<0.001; ****: P<0.0001, two-tailed Student *t*-test). w. t.: wild type. (**D**) Quantification of internalized bacteria by the differential staining assay. Values represent mean ± SD of the ratio of internalized vs total bacteria derived from quantifying a minimum of 1,000 cells per condition. (*: P<0.05; **: P<0.01, two tailed Student *t* test). (**E**) Representative fluorescence microscopy images showing stained nonpermeabilized (green) or all (nonpermeabilized and permeabilized, red) *S*. Typhimurium after infection. The mammalian cell nuclear DNA was stained by 4',6-diamidino-2-phenylindole (DAPI) (blue).

Figure 7. The pro-inflammatory activity of SipA is independent of actin-binding. Mice were infected with *S*. Typhimurium strains lacking *sipA*, expressing the actin-binding defective mutant SipA$^{S537E/K636E}$, or expressing wild-type SipA. To enhance the detection of SipA-dependent inflammation (27), a Δ*spiA* mutation was introduced in all strains, rendering them deficient for the SPI-2 T3SS. Gross pathology (**A**), weight (**B**), and histopathology (**C**) of ceca obtained from mice infected with the different strains or uninfected controls, as indicated, are shown. Values in (**B**) are the weight of each cecum in mg and represent the mean ± SD (*: P<0.05; ****: P<0.0001; ns: not signifcant; two tailed Student *t* test)

Pointed end

90°

90°

90°

Barbed end

Figure 1

A

B

1Q5Z

N

C

C

SD4

SD2

6DJO

ID

OD

SD3

SD1

SipA

D

6DJO

Figure 2

Figure 3

A **SipA** **Phalloidin**

B **Jasplakinolide**

C Actin-SipA Actin-Pyrene **Pyrene**
V45 H539
D-Loop

D **Nebulin**

E **Arp2/3 complex**

F **C-terminal loop** **D-Loop** Cofilin

Figure 4

Figure 5

Figure 6

A

uninfected

wild-type

$\Delta sipA$

$sipA^{S537E/K636E}$

B

<!-- chart: cecum weight (mg) vs uninfected, wild-type, ΔsipA, sipA^S537E/K636E; significance markers ****, *, ns, * -->

C

uninfected

wild type

$\Delta sipA$

$sipA^{S537E/K636E}$

Figure 7

Supplementary Figures for:

Cryo-EM structure of the bacterial effector protein SipA bound to F-actin reveals a unique mechanism for filament stabilization

Emily Z. Guo[1], Steven Z. Chou[2], Maria Lara-Tejero[1], and Jorge E. Galán[1]*

[1]Department of Microbial Pathogenesis, Yale University School of Medicine, New Haven, CT 06536, and [2]Department of Molecular Biology and Biophysics, UConn Health, Farmington, CT 06030

Supplementary Fig. 1. Diagram of SipA and AlphaFold prediction of its structure.

Supplementary Fig. 2. Actin-binding affinity of SipA447-685. as determined by a co-sedimentation assay.

Actin

SipA$_{447-685}$-Actin

Supplementary Fig. 3. Representative electron micrographs of negatively stained actin filaments alone and with SipA$_{447-685}$

Supplementary Fig. 4. Layer line images of actin filaments decorated with SipA. (**A**) Averaged power spectrum of helix segments boxed out from 1300 micrographs. (**B**) Power spectrum calculated from map projections. The Equator line (n = 0) and the first rise line (n = 0) are indicated with red and yellow lines, respectively. Rise = (pixel size) × (layer-line image size)/ (layer-line height) = 1.068 × 4,096/39.91 = 109.6 Å.

Supplementary Fig. 5. Lower electron map occupancy of $SipA_{446-684}$ in actin filaments. (**A**) representative 2D class averages of $SipA_{447-685}$ decorated actin filaments. Arrows show the density of $SipA_{447-685}$. (**B**) Initial 3D reconstruction of $SipA_{447-685}$ decorated actin filaments. Arrows show the density of $SipA_{447-685}$.

.

Supplementary Fig. 6. Global resolution estimation of the SipA$_{446-684}$-actin complex. Fourier shell correlation for maps constructed from two half data sets. A cutoff value of 0.143 is used.

Supplementary Fig. 7. Local resolution estimation for SipA-decorated F-actin with Resmap.

Supplemental Fig. 8. Cryo-EM map and visualization of the side chains of selective residues of actin near the ADP binding site of the model. The density for the fitted coordinates shows features consistent with a resolution of ~ 3.6 Å.

Supplemental Fig. 9. The globular domain (violet) and the C-terminal extension (magenta) of SipA. Actin filament is colored in gray. Ribbon diagram is shown on the right panel.

Supplementary Fig. 10. The unstructured arm regions are necessary for the G-actin nucleation function of SipA but are dispensable for its ability to stabilize F-actin. (**A** and **B**) G-actin nucleation activity of different SipA constructs as determined by a co-sedimentation assay. Values shown in (**A**) are the mean ± SD of three independent experiments (*: $P<0.05$; **: $P<0.01$; unpaired two-sided t-test). (**C** and **D**) F-actin binding activity of different SipA constructs as determined by a co-sedimentation assay. Values shown in (**C**) are the mean ± SD of three independent experiments (ns: differences not significant, unpaired two-sided t-test). (**E**) A SipA mutant lacking the unstructured C-terminal region exhibits reduced epithelial cell invasion ability. The invasive ability of the different strains was measured by the gentamicin protection assay as described in the methods. Values represent the percentage of the inoculum that survived antibiotic treatment due to internalization and are the mean ± SD of six independent determinations normalized to wild type, which was set to 1. (**: $P<0.01$; ***: $P<0.001$; unpaired two-sided t-test). WT: wild type.

Supplementary Fig. 11. Residues S537 and K636 of SipA interact with the D-loop and C-terminus of actin. Interfaces between SipA (magenta) and actin (green) are shown. Interface residues S537 and K635 of SipA, as well as the D loop and the C-terminal residue E364 of actin are indicated.

Supplementary Fig. 12. Data processing workflow.

Supplementary Table S1. Statistics for data collection, map reconstruction and model refinement

Cryo-EM data collection and model refinement	
Data collection and map reconstruction	
Microscope	Titan Krios
Voltage (kV)	300
Detector	K3
Magnification	81,000
Electron dose (e^-/$Å^2$)	68.1384
Defocus range (μm)	−1.2 to −2.5
Pixel size (Å)	1.068
Symmetry	C1; rise: 109.862 Å; twist: 53.3346 ^{O}
Initial filament segments (no.)	320,109
Final filament segments (no.)	81,799
Map resolution (Å)	3.6
FSC threshold	0.143
Model Refinement	
Initial model used (PDB IDs)	6DJO, 1Q5Z
Model resolution (Å)	3.78
FSC threshold	0.5
Model composition	8 actin subunits, 4 SipA subunits
Non-hydrogen atoms	28321
Protein residues	3616
Ligands	8 Mg^{2+}, 8 ADP.
B factors (Å2)	
Protein	112.29
Ligand	98.01
R.m.s. deviations	
Bond lengths (Å)	0.010
Bond angles (°)	1.317
Geometry validation	
MolProbity score	1.71
All-atom Clashscore	6.53
Rotamers outliers (%)	0.00
Ramachandran plot	
Favored (%)	94.90
Allowed (%)	5.10
Disallowed (%)	0.00

Preliminary Full wwPDB EM Validation Report (i)

Feb 21, 2024 – 11:44 PM EST

Deposition ID : D_1000280011

This is a Preliminary Full wwPDB EM Validation Report.

This report is produced by the wwPDB Deposition System
during initial deposition but before annotation of the structure.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev70
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	685	20% · 76%
1	B	685	19% · 76%
1	C	685	19% · 76%
1	I	685	19% · 76%
2	J	377	81% 17% ·
2	L	377	81% 18% ·
2	M	377	81% 18% ·
2	N	377	81% 18% ·

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	R	377	81% 17%
2	S	377	79% 20%
2	U	377	80% 18%
2	W	377	82% 16%

2 Entry composition ⓘ

There are 4 unique types of molecules in this entry. The entry contains 28316 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cell invasion protein SipA.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	B	162	Total 1223	C 773	N 216	O 230	S 4	0	0
1	A	162	Total 1223	C 773	N 216	O 230	S 4	0	0
1	C	162	Total 1223	C 773	N 216	O 230	S 4	0	0
1	I	162	Total 1223	C 773	N 216	O 230	S 4	0	0

- Molecule 2 is a protein called Actin, alpha skeletal muscle.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	J	371	Total 2900	C 1837	N 489	O 553	S 21	0	0
2	L	371	Total 2900	C 1837	N 489	O 553	S 21	0	0
2	M	371	Total 2900	C 1837	N 489	O 553	S 21	0	0
2	N	371	Total 2900	C 1837	N 489	O 553	S 21	0	0
2	R	371	Total 2900	C 1837	N 489	O 553	S 21	0	0
2	S	371	Total 2900	C 1837	N 489	O 553	S 21	0	0
2	U	371	Total 2900	C 1837	N 489	O 553	S 21	0	0
2	W	371	Total 2900	C 1837	N 489	O 553	S 21	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
3	J	1	Total 1	Mg 1	0
3	L	1	Total 1	Mg 1	0
3	M	1	Total 1	Mg 1	0
3	N	1	Total 1	Mg 1	0
3	R	1	Total 1	Mg 1	0
3	S	1	Total 1	Mg 1	0
3	U	1	Total 1	Mg 1	0
3	W	1	Total 1	Mg 1	0

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
4	J	1	27	10	5	10	2	0
4	L	1	Total 27	C 10	N 5	O 10	P 2	0
4	M	1	Total 27	C 10	N 5	O 10	P 2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
4	N	1	Total 27	C 10	N 5	O 10	P 2	0
4	R	1	Total 27	C 10	N 5	O 10	P 2	0
4	S	1	Total 27	C 10	N 5	O 10	P 2	0
4	U	1	Total 27	C 10	N 5	O 10	P 2	0
4	W	1	Total 27	C 10	N 5	O 10	P 2	0

3 Residue-property plots ⓘ

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cell invasion protein SipA

Chain B:

• Molecule 1: Cell invasion protein SipA

Chain A:

• Molecule 1: Cell invasion protein SipA

Chain C: 19% 76%

• Molecule 1: Cell invasion protein SipA

Chain I:

• Molecule 2: Actin, alpha skeletal muscle

Chain J:

- Molecule 2: Actin, alpha skeletal muscle

Chain L:

- Molecule 2: Actin, alpha skeletal muscle

Chain M:

- Molecule 2: Actin, alpha skeletal muscle

Chain N:

• Molecule 2: Actin, alpha skeletal muscle

Chain R:

• Molecule 2: Actin, alpha skeletal muscle

Chain S:

• Molecule 2: Actin, alpha skeletal muscle

Chain U:

• Molecule 2: Actin, alpha skeletal muscle

Chain W:

4 Experimental information ⓘ

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=53.2°, rise=109.6 Å, axial sym=C1	Depositor
Number of segments used	320109	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	68.1384	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.959	Depositor
Minimum map value	-0.068	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.037	Depositor
Recommended contour level	0.0434	Depositor
Map size (Å)	341.75998, 341.75998, 341.75998	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.068, 1.068, 1.068	Depositor

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: HIC, ADP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#|Z|>5$	RMSZ	$\#|Z|>5$
1	A	0.34	0/1247	0.55	0/1688
1	B	0.34	0/1247	0.55	0/1688
1	C	0.34	0/1247	0.55	0/1688
1	I	0.34	0/1247	0.55	0/1688
2	J	0.50	0/2950	0.59	0/3994
2	L	0.50	0/2950	0.59	0/3994
2	M	0.50	0/2950	0.59	0/3994
2	N	0.50	0/2950	0.59	0/3994
2	R	0.50	0/2950	0.59	0/3994
2	S	0.50	0/2950	0.59	0/3994
2	U	0.50	0/2950	0.59	0/3994
2	W	0.50	0/2950	0.58	0/3994
All	All	0.48	0/28588	0.58	0/38704

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1223	0	1204	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1223	0	1204	19	0
1	C	1223	0	1204	17	0
1	I	1223	0	1204	16	0
2	J	2900	0	2872	37	0
2	L	2900	0	2872	41	0
2	M	2900	0	2872	37	0
2	N	2900	0	2872	37	0
2	R	2900	0	2872	35	0
2	S	2900	0	2872	47	0
2	U	2900	0	2872	40	0
2	W	2900	0	2872	34	0
3	J	1	0	0	0	0
3	L	1	0	0	0	0
3	M	1	0	0	0	0
3	N	1	0	0	0	0
3	R	1	0	0	0	0
3	S	1	0	0	0	0
3	U	1	0	0	0	0
3	W	1	0	0	0	0
4	J	27	0	12	2	0
4	L	27	0	12	2	0
4	M	27	0	12	0	0
4	N	27	0	12	0	0
4	R	27	0	12	0	0
4	S	27	0	12	3	0
4	U	27	0	12	1	0
4	W	27	0	12	0	0
All	All	28316	0	27888	351	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (351) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:537:SER:HB3	1:I:540:PHE:HB2	1.71	0.72
1:C:537:SER:HB3	1:C:540:PHE:HB2	1.72	0.71
1:A:537:SER:HB3	1:A:540:PHE:HB2	1.71	0.71
1:B:537:SER:HB3	1:B:540:PHE:HB2	1.72	0.70
2:L:113:LYS:HE2	2:U:195:GLU:HB3	1.73	0.70
1:A:554:LEU:O	1:A:572:ARG:NH2	2.25	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:554:LEU:O	1:C:572:ARG:NH2	2.25	0.69
1:B:554:LEU:O	1:B:572:ARG:NH2	2.25	0.68
1:I:554:LEU:O	1:I:572:ARG:NH2	2.25	0.67
2:S:39:ARG:NH2	2:U:268:GLY:O	2.28	0.66
1:B:522:GLN:HE22	2:W:322:PRO:HA	1.65	0.61
2:U:18:LYS:NZ	4:U:802:ADP:O1B	2.34	0.61
2:J:207:GLU:OE2	2:J:210:ARG:NH2	2.36	0.59
2:L:207:GLU:OE2	2:L:210:ARG:NH2	2.36	0.59
2:R:207:GLU:OE2	2:R:210:ARG:NH2	2.36	0.59
2:S:207:GLU:OE2	2:S:210:ARG:NH2	2.36	0.59
2:M:207:GLU:OE2	2:M:210:ARG:NH2	2.36	0.59
2:N:207:GLU:OE2	2:N:210:ARG:NH2	2.36	0.59
2:U:207:GLU:OE2	2:U:210:ARG:NH2	2.36	0.59
2:W:207:GLU:OE2	2:W:210:ARG:NH2	2.36	0.59
1:I:522:GLN:HE22	2:R:322:PRO:HA	1.68	0.58
2:J:113:LYS:HE2	2:W:195:GLU:HB3	1.86	0.57
2:L:152:VAL:HG22	2:L:298:VAL:HB	1.87	0.57
2:S:152:VAL:HG22	2:S:298:VAL:HB	1.87	0.57
2:J:152:VAL:HG22	2:J:298:VAL:HB	1.87	0.56
2:R:152:VAL:HG22	2:R:298:VAL:HB	1.87	0.56
1:C:607:LYS:HD3	1:C:645:ASN:HB2	1.87	0.56
2:N:152:VAL:HG22	2:N:298:VAL:HB	1.87	0.56
2:U:152:VAL:HG22	2:U:298:VAL:HB	1.87	0.56
1:I:607:LYS:HD3	1:I:645:ASN:HB2	1.87	0.55
2:M:152:VAL:HG22	2:M:298:VAL:HB	1.87	0.55
1:B:607:LYS:HD3	1:B:645:ASN:HB2	1.87	0.55
1:A:607:LYS:HD3	1:A:645:ASN:HB2	1.87	0.55
2:R:133:TYR:HH	2:R:374:CYS:HG	1.52	0.55
2:W:152:VAL:HG22	2:W:298:VAL:HB	1.87	0.55
2:R:189:LEU:HD22	2:R:209:VAL:HG13	1.89	0.55
1:C:645:ASN:HA	1:C:648:ASP:HB2	1.89	0.54
2:J:12:ASN:HD21	2:J:105:LEU:HD22	1.72	0.54
2:S:12:ASN:HD21	2:S:105:LEU:HD22	1.72	0.54
2:L:12:ASN:HD21	2:L:105:LEU:HD22	1.72	0.54
2:N:12:ASN:HD21	2:N:105:LEU:HD22	1.72	0.54
2:U:12:ASN:HD21	2:U:105:LEU:HD22	1.72	0.54
2:U:189:LEU:HD22	2:U:209:VAL:HG13	1.89	0.54
2:W:274:ILE:HA	2:W:277:THR:HG22	1.90	0.54
2:J:274:ILE:HA	2:J:277:THR:HG22	1.90	0.54
2:R:274:ILE:HA	2:R:277:THR:HG22	1.90	0.54
2:L:189:LEU:HD22	2:L:209:VAL:HG13	1.89	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:S:274:ILE:HA	2:S:277:THR:HG22	1.89	0.54
2:W:189:LEU:HD22	2:W:209:VAL:HG13	1.89	0.54
2:J:189:LEU:HD22	2:J:209:VAL:HG13	1.89	0.54
2:M:189:LEU:HD22	2:M:209:VAL:HG13	1.89	0.54
2:N:189:LEU:HD22	2:N:209:VAL:HG13	1.89	0.54
2:N:274:ILE:HA	2:N:277:THR:HG22	1.90	0.54
2:M:274:ILE:HA	2:M:277:THR:HG22	1.90	0.54
2:J:197:GLY:HA2	2:M:112:PRO:HG3	1.90	0.54
2:U:274:ILE:HA	2:U:277:THR:HG22	1.90	0.54
2:M:12:ASN:HD21	2:M:105:LEU:HD22	1.72	0.54
2:S:189:LEU:HD22	2:S:209:VAL:HG13	1.89	0.54
2:W:12:ASN:HD21	2:W:105:LEU:HD22	1.72	0.54
2:R:12:ASN:HD21	2:R:105:LEU:HD22	1.72	0.53
2:J:230:ALA:O	2:J:233:SER:OG	2.26	0.53
2:L:274:ILE:HA	2:L:277:THR:HG22	1.90	0.53
1:I:645:ASN:HA	1:I:648:ASP:HB2	1.89	0.53
2:S:14:SER:O	2:S:183:ARG:NH2	2.42	0.53
1:A:645:ASN:O	1:A:649:GLY:N	2.42	0.53
2:L:14:SER:O	2:L:183:ARG:NH2	2.42	0.53
1:B:645:ASN:HA	1:B:648:ASP:HB2	1.89	0.53
2:L:270:GLU:HB2	2:U:39:ARG:CZ	2.39	0.53
2:M:14:SER:O	2:M:183:ARG:NH2	2.42	0.53
2:N:230:ALA:O	2:N:233:SER:OG	2.26	0.53
2:W:14:SER:O	2:W:183:ARG:NH2	2.42	0.53
1:B:645:ASN:O	1:B:649:GLY:N	2.42	0.52
1:A:645:ASN:HA	1:A:648:ASP:HB2	1.89	0.52
2:J:14:SER:O	2:J:183:ARG:NH2	2.42	0.52
2:U:14:SER:O	2:U:183:ARG:NH2	2.42	0.52
2:R:107:GLU:OE2	2:R:116:ARG:NE	2.42	0.52
2:W:147:ARG:HH12	2:W:330:ILE:HG13	1.75	0.52
2:M:147:ARG:HH12	2:M:330:ILE:HG13	1.75	0.52
2:N:14:SER:O	2:N:183:ARG:NH2	2.42	0.52
2:R:14:SER:O	2:R:183:ARG:NH2	2.42	0.52
1:I:645:ASN:O	1:I:649:GLY:N	2.42	0.52
2:L:332:PRO:O	2:L:335:ARG:NH1	2.43	0.52
2:L:147:ARG:HH12	2:L:330:ILE:HG13	1.75	0.52
2:M:86:TRP:HH2	2:M:119:MET:HG3	1.75	0.52
2:R:332:PRO:O	2:R:335:ARG:NH1	2.43	0.52
2:M:332:PRO:O	2:M:335:ARG:NH1	2.43	0.52
2:R:147:ARG:HH12	2:R:330:ILE:HG13	1.75	0.52
2:U:86:TRP:HH2	2:U:119:MET:HG3	1.75	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:U:332:PRO:O	2:U:335:ARG:NH1	2.43	0.52
2:W:86:TRP:HH2	2:W:119:MET:HG3	1.75	0.52
1:C:645:ASN:O	1:C:649:GLY:N	2.42	0.52
2:N:86:TRP:HH2	2:N:119:MET:HG3	1.75	0.52
2:R:86:TRP:HH2	2:R:119:MET:HG3	1.75	0.52
1:C:551:ARG:NH1	1:C:576:LEU:O	2.44	0.52
2:J:200:PHE:HB3	2:J:205:GLU:HB3	1.92	0.52
2:J:332:PRO:O	2:J:335:ARG:NH1	2.43	0.52
2:W:332:PRO:O	2:W:335:ARG:NH1	2.43	0.52
2:L:86:TRP:HH2	2:L:119:MET:HG3	1.75	0.51
2:N:200:PHE:HB3	2:N:205:GLU:HB3	1.92	0.51
2:J:86:TRP:HH2	2:J:119:MET:HG3	1.75	0.51
2:L:230:ALA:O	2:L:233:SER:OG	2.26	0.51
2:S:200:PHE:HB3	2:S:205:GLU:HB3	1.92	0.51
1:I:551:ARG:NH1	1:I:576:LEU:O	2.43	0.51
2:J:53:TYR:HB2	2:J:65:LEU:HD21	1.93	0.51
2:N:53:TYR:HB2	2:N:65:LEU:HD21	1.93	0.51
2:S:86:TRP:HH2	2:S:119:MET:HG3	1.75	0.51
2:U:200:PHE:HB3	2:U:205:GLU:HB3	1.92	0.51
2:W:107:GLU:OE2	2:W:116:ARG:NE	2.42	0.51
2:M:200:PHE:HB3	2:M:205:GLU:HB3	1.92	0.51
2:S:332:PRO:O	2:S:335:ARG:NH1	2.43	0.51
2:U:53:TYR:HB2	2:U:65:LEU:HD21	1.93	0.51
1:B:551:ARG:NH1	1:B:576:LEU:O	2.44	0.51
2:M:53:TYR:HB2	2:M:65:LEU:HD21	1.93	0.51
2:S:53:TYR:HB2	2:S:65:LEU:HD21	1.93	0.51
2:W:230:ALA:O	2:W:233:SER:OG	2.26	0.51
2:J:39:ARG:NH2	2:M:268:GLY:O	2.35	0.51
2:N:71:ILE:HG12	2:N:76:ILE:HG12	1.93	0.51
2:U:71:ILE:HG12	2:U:76:ILE:HG12	1.93	0.51
2:U:147:ARG:HH12	2:U:330:ILE:HG13	1.75	0.51
2:L:200:PHE:HB3	2:L:205:GLU:HB3	1.92	0.51
2:N:332:PRO:O	2:N:335:ARG:NH1	2.43	0.51
2:W:200:PHE:HB3	2:W:205:GLU:HB3	1.92	0.51
2:L:53:TYR:HB2	2:L:65:LEU:HD21	1.93	0.51
2:M:107:GLU:OE2	2:M:116:ARG:NE	2.42	0.51
2:W:71:ILE:HG12	2:W:76:ILE:HG12	1.93	0.51
1:A:551:ARG:NH1	1:A:576:LEU:O	2.44	0.50
2:J:71:ILE:HG12	2:J:76:ILE:HG12	1.93	0.50
2:M:78:ASN:ND2	2:M:81:ASP:OD2	2.44	0.50
2:S:78:ASN:ND2	2:S:81:ASP:OD2	2.45	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:78:ASN:ND2	2:J:81:ASP:OD2	2.45	0.50
2:N:147:ARG:HH12	2:N:330:ILE:HG13	1.75	0.50
2:W:53:TYR:HB2	2:W:65:LEU:HD21	1.93	0.50
2:S:147:ARG:HH12	2:S:330:ILE:HG13	1.75	0.50
2:U:107:GLU:OE2	2:U:116:ARG:NE	2.42	0.50
2:U:230:ALA:O	2:U:233:SER:OG	2.27	0.50
2:J:147:ARG:HH12	2:J:330:ILE:HG13	1.75	0.50
2:N:107:GLU:OE2	2:N:116:ARG:NE	2.42	0.50
2:L:78:ASN:ND2	2:L:81:ASP:OD2	2.44	0.50
2:R:200:PHE:HB3	2:R:205:GLU:HB3	1.92	0.50
2:R:53:TYR:HB2	2:R:65:LEU:HD21	1.93	0.50
2:N:169:TYR:HA	2:U:42:GLY:HA2	1.94	0.50
2:W:78:ASN:ND2	2:W:81:ASP:OD2	2.45	0.50
2:N:78:ASN:ND2	2:N:81:ASP:OD2	2.44	0.50
2:R:78:ASN:ND2	2:R:81:ASP:OD2	2.45	0.50
2:L:71:ILE:HG12	2:L:76:ILE:HG12	1.93	0.49
1:I:523:ASN:OD1	1:I:618:LYS:NZ	2.45	0.49
2:M:71:ILE:HG12	2:M:76:ILE:HG12	1.93	0.49
2:U:222:ASP:OD2	2:U:225:ASN:ND2	2.46	0.49
2:U:78:ASN:ND2	2:U:81:ASP:OD2	2.45	0.49
2:M:230:ALA:O	2:M:233:SER:OG	2.26	0.49
2:N:11:ASP:OD1	2:N:106:THR:OG1	2.31	0.49
2:S:230:ALA:O	2:S:233:SER:OG	2.27	0.49
2:J:222:ASP:OD2	2:J:225:ASN:ND2	2.46	0.49
2:L:18:LYS:NZ	4:L:802:ADP:O1B	2.46	0.49
2:L:186:THR:HG1	2:L:213:LYS:HZ3	1.58	0.49
2:M:62:ARG:HB2	2:M:67:LEU:HD11	1.95	0.49
2:R:71:ILE:HG12	2:R:76:ILE:HG12	1.93	0.49
1:B:523:ASN:OD1	1:B:618:LYS:NZ	2.45	0.49
2:S:71:ILE:HG12	2:S:76:ILE:HG12	1.93	0.49
2:W:62:ARG:HB2	2:W:67:LEU:HD11	1.95	0.49
1:C:523:ASN:OD1	1:C:618:LYS:NZ	2.45	0.49
2:U:11:ASP:OD1	2:U:106:THR:OG1	2.31	0.49
2:U:62:ARG:HB2	2:U:67:LEU:HD11	1.95	0.49
1:A:538:MET:SD	1:A:539:HIS:ND1	2.77	0.49
2:N:62:ARG:HB2	2:N:67:LEU:HD11	1.95	0.49
2:N:222:ASP:OD2	2:N:225:ASN:ND2	2.46	0.49
2:W:11:ASP:OD1	2:W:106:THR:OG1	2.31	0.48
2:M:272:ALA:HB1	2:M:276:GLU:HB3	1.95	0.48
2:U:272:ALA:HB1	2:U:276:GLU:HB3	1.95	0.48
2:N:272:ALA:HB1	2:N:276:GLU:HB3	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:W:222:ASP:OD2	2:W:225:ASN:ND2	2.46	0.48
2:N:9:VAL:HG12	2:N:104:LEU:HB3	1.95	0.48
2:U:9:VAL:HG12	2:U:104:LEU:HB3	1.95	0.48
1:B:604:GLU:OE2	1:B:641:TYR:OH	2.32	0.48
2:J:186:THR:HG1	2:J:213:LYS:HZ3	1.58	0.48
2:L:9:VAL:HG12	2:L:104:LEU:HB3	1.95	0.48
2:M:9:VAL:HG12	2:M:104:LEU:HB3	1.95	0.48
2:R:9:VAL:HG12	2:R:104:LEU:HB3	1.95	0.48
2:W:9:VAL:HG12	2:W:104:LEU:HB3	1.95	0.48
2:M:11:ASP:OD1	2:M:106:THR:OG1	2.31	0.48
2:W:272:ALA:HB1	2:W:276:GLU:HB3	1.95	0.48
1:A:523:ASN:OD1	1:A:618:LYS:NZ	2.45	0.48
2:N:195:GLU:HB3	2:R:113:LYS:HE2	1.96	0.48
2:S:9:VAL:HG12	2:S:104:LEU:HB3	1.96	0.48
2:S:180:LEU:HD11	2:S:260:THR:HG22	1.96	0.48
1:I:604:GLU:OE2	1:I:641:TYR:OH	2.32	0.48
2:J:9:VAL:HG12	2:J:104:LEU:HB3	1.95	0.48
2:S:62:ARG:HB2	2:S:67:LEU:HD11	1.95	0.48
1:A:604:GLU:OE2	1:A:641:TYR:OH	2.32	0.48
2:J:180:LEU:HD11	2:J:260:THR:HG22	1.96	0.48
2:R:180:LEU:HD11	2:R:260:THR:HG22	1.96	0.48
2:S:272:ALA:HB1	2:S:276:GLU:HB3	1.95	0.48
2:J:62:ARG:HB2	2:J:67:LEU:HD11	1.95	0.47
2:L:180:LEU:HD11	2:L:260:THR:HG22	1.96	0.47
2:R:62:ARG:HB2	2:R:67:LEU:HD11	1.95	0.47
1:I:628:ARG:NH2	1:I:629:GLU:OE2	2.47	0.47
2:J:272:ALA:HB1	2:J:276:GLU:HB3	1.95	0.47
2:R:230:ALA:O	2:R:233:SER:OG	2.26	0.47
2:W:180:LEU:HD11	2:W:260:THR:HG22	1.96	0.47
2:L:62:ARG:HB2	2:L:67:LEU:HD11	1.95	0.47
2:S:222:ASP:OD2	2:S:225:ASN:ND2	2.46	0.47
2:M:195:GLU:HB3	2:S:113:LYS:HE2	1.96	0.47
1:C:628:ARG:NH2	1:C:629:GLU:OE2	2.47	0.47
2:L:272:ALA:HB1	2:L:276:GLU:HB3	1.95	0.47
2:N:180:LEU:HD11	2:N:260:THR:HG22	1.96	0.47
2:L:11:ASP:OD1	2:L:106:THR:OG1	2.31	0.47
2:M:222:ASP:OD2	2:M:225:ASN:ND2	2.46	0.47
1:B:628:ARG:NH2	1:B:629:GLU:OE2	2.47	0.47
1:C:522:GLN:HE22	2:S:322:PRO:HA	1.80	0.47
1:C:604:GLU:OE2	1:C:641:TYR:OH	2.32	0.47
2:L:107:GLU:OE2	2:L:116:ARG:NE	2.42	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:222:ASP:OD2	2:L:225:ASN:ND2	2.46	0.47
2:U:180:LEU:HD11	2:U:260:THR:HG22	1.96	0.47
1:I:538:MET:SD	1:I:539:HIS:ND1	2.77	0.47
2:M:180:LEU:HD11	2:M:260:THR:HG22	1.96	0.47
1:A:628:ARG:NH2	1:A:629:GLU:OE2	2.47	0.47
2:M:197:GLY:HA2	2:S:112:PRO:HG3	1.95	0.47
2:S:11:ASP:OD1	2:S:106:THR:OG1	2.31	0.47
2:S:107:GLU:OE2	2:S:116:ARG:NE	2.42	0.47
2:L:168:GLY:HA2	2:S:44:MET:HB2	1.96	0.47
2:R:272:ALA:HB1	2:R:276:GLU:HB3	1.95	0.47
2:J:11:ASP:OD1	2:J:106:THR:OG1	2.31	0.46
2:S:157:ASP:OD1	4:S:802:ADP:O3'	2.32	0.46
2:R:11:ASP:OD1	2:R:106:THR:OG1	2.31	0.46
2:M:17:VAL:HG23	2:M:33:SER:HB2	1.98	0.46
2:J:107:GLU:OE2	2:J:116:ARG:NE	2.42	0.46
2:L:169:TYR:HA	2:S:42:GLY:H	1.81	0.46
2:W:17:VAL:HG23	2:W:33:SER:HB2	1.98	0.46
2:J:157:ASP:OD1	4:J:802:ADP:O3'	2.30	0.46
2:R:222:ASP:OD2	2:R:225:ASN:ND2	2.46	0.46
2:U:17:VAL:HG23	2:U:33:SER:HB2	1.98	0.46
1:A:575:ILE:HG22	1:A:576:LEU:HG	1.98	0.46
2:J:205:GLU:HA	2:J:208:ILE:HG12	1.98	0.46
1:B:575:ILE:HG22	1:B:576:LEU:HG	1.98	0.45
2:J:17:VAL:HG23	2:J:33:SER:HB2	1.98	0.45
2:S:197:GLY:HA2	2:U:112:PRO:HG3	1.98	0.45
2:N:17:VAL:HG23	2:N:33:SER:HB2	1.98	0.45
2:W:205:GLU:HA	2:W:208:ILE:HG12	1.98	0.45
1:B:538:MET:SD	1:B:539:HIS:ND1	2.77	0.45
2:L:17:VAL:HG23	2:L:33:SER:HB2	1.98	0.45
2:L:205:GLU:HA	2:L:208:ILE:HG12	1.98	0.45
2:S:17:VAL:HG23	2:S:33:SER:HB2	1.98	0.45
2:S:205:GLU:HA	2:S:208:ILE:HG12	1.98	0.45
1:C:575:ILE:HG22	1:C:576:LEU:HG	1.98	0.45
2:L:197:GLY:HA2	2:N:112:PRO:HG3	1.99	0.45
2:J:18:LYS:NZ	4:J:802:ADP:O1B	2.50	0.45
2:J:202:THR:OG1	2:J:203:THR:N	2.49	0.45
2:M:205:GLU:HA	2:M:208:ILE:HG12	1.98	0.45
2:N:205:GLU:HA	2:N:208:ILE:HG12	1.98	0.45
2:R:17:VAL:HG23	2:R:33:SER:HB2	1.98	0.45
2:R:205:GLU:HA	2:R:208:ILE:HG12	1.98	0.45
2:N:147:ARG:HH22	2:N:330:ILE:HG13	1.82	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:R:151:ILE:HD11	2:R:278:THR:HG23	1.99	0.45
2:U:205:GLU:HA	2:U:208:ILE:HG12	1.98	0.45
2:W:156:GLY:O	2:W:303:THR:OG1	2.35	0.45
2:M:147:ARG:HH22	2:M:330:ILE:HG13	1.82	0.45
2:S:14:SER:N	4:S:802:ADP:O3B	2.49	0.45
2:U:147:ARG:HH22	2:U:330:ILE:HG13	1.82	0.45
2:L:151:ILE:HD11	2:L:278:THR:HG23	1.99	0.45
2:L:157:ASP:OD1	4:L:802:ADP:O3'	2.33	0.45
1:B:532:VAL:HG11	2:W:266:PHE:HE1	1.82	0.44
1:I:575:ILE:HG22	1:I:576:LEU:HG	1.98	0.44
2:U:151:ILE:HD11	2:U:278:THR:HG23	1.99	0.44
2:S:151:ILE:HD11	2:S:278:THR:HG23	1.99	0.44
2:U:289:ILE:HG22	2:U:293:LEU:HG	2.00	0.44
1:C:538:MET:SD	1:C:539:HIS:ND1	2.77	0.44
1:C:602:LYS:O	1:C:605:THR:OG1	2.33	0.44
2:M:151:ILE:HD11	2:M:278:THR:HG23	1.99	0.44
2:S:195:GLU:HB3	2:U:113:LYS:HE2	2.00	0.44
1:I:602:LYS:O	1:I:605:THR:OG1	2.33	0.44
2:M:289:ILE:HG22	2:M:293:LEU:HG	2.00	0.44
2:W:147:ARG:HH22	2:W:330:ILE:HG13	1.82	0.44
2:W:151:ILE:HD11	2:W:278:THR:HG23	1.99	0.44
2:N:289:ILE:HG22	2:N:293:LEU:HG	2.00	0.44
1:B:602:LYS:O	1:B:605:THR:OG1	2.33	0.44
2:J:151:ILE:HD11	2:J:278:THR:HG23	1.99	0.44
2:N:151:ILE:HD11	2:N:278:THR:HG23	1.99	0.44
2:N:202:THR:OG1	2:N:203:THR:N	2.49	0.44
2:L:39:ARG:NH2	2:N:268:GLY:O	2.38	0.44
1:I:627:ALA:HA	1:I:632:LEU:HB3	2.00	0.44
2:L:147:ARG:HH22	2:L:330:ILE:HG13	1.82	0.44
2:R:147:ARG:HH22	2:R:330:ILE:HG13	1.82	0.44
1:B:627:ALA:HA	1:B:632:LEU:HB3	2.00	0.43
2:M:133:TYR:HH	2:M:374:CYS:HG	1.58	0.43
2:W:289:ILE:HG22	2:W:293:LEU:HG	2.00	0.43
1:A:627:ALA:HA	1:A:632:LEU:HB3	2.00	0.43
2:S:147:ARG:HH22	2:S:330:ILE:HG13	1.82	0.43
2:J:147:ARG:HH22	2:J:330:ILE:HG13	1.82	0.43
2:U:202:THR:OG1	2:U:203:THR:N	2.49	0.43
2:M:202:THR:OG1	2:M:203:THR:N	2.49	0.43
2:L:169:TYR:HA	2:S:42:GLY:HA2	2.01	0.43
2:L:289:ILE:HG22	2:L:293:LEU:HG	2.00	0.43
2:R:202:THR:OG1	2:R:203:THR:N	2.49	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:W:283:MET:O	2:W:290:ARG:NH2	2.46	0.43
2:L:166:TYR:HD2	2:S:64:ILE:HD13	1.84	0.43
1:C:627:ALA:HA	1:C:632:LEU:HB3	2.00	0.42
2:J:289:ILE:HG22	2:J:293:LEU:HG	2.00	0.42
2:R:156:GLY:O	2:R:303:THR:OG1	2.35	0.42
2:R:289:ILE:HG22	2:R:293:LEU:HG	2.00	0.42
2:S:289:ILE:HG22	2:S:293:LEU:HG	2.00	0.42
1:A:602:LYS:O	1:A:605:THR:OG1	2.33	0.42
2:L:202:THR:OG1	2:L:203:THR:N	2.49	0.42
2:N:283:MET:O	2:N:290:ARG:NH2	2.46	0.42
2:S:271:SER:OG	2:S:272:ALA:N	2.52	0.42
2:S:156:GLY:O	2:S:303:THR:OG1	2.35	0.42
2:L:271:SER:OG	2:L:272:ALA:N	2.53	0.42
2:S:202:THR:OG1	2:S:203:THR:N	2.49	0.42
2:W:36:GLY:HA3	2:W:65:LEU:HD23	2.02	0.42
2:J:156:GLY:O	2:J:303:THR:OG1	2.34	0.42
2:M:156:GLY:O	2:M:303:THR:OG1	2.35	0.42
2:M:271:SER:OG	2:M:272:ALA:N	2.52	0.42
2:N:36:GLY:HA3	2:N:65:LEU:HD23	2.02	0.42
2:R:271:SER:OG	2:R:272:ALA:N	2.52	0.42
2:U:36:GLY:HA3	2:U:65:LEU:HD23	2.02	0.42
2:U:283:MET:O	2:U:290:ARG:NH2	2.46	0.42
1:C:578:ASN:ND2	2:M:83:GLU:OE1	2.53	0.42
2:S:283:MET:O	2:S:290:ARG:NH2	2.46	0.42
2:M:36:GLY:HA3	2:M:65:LEU:HD23	2.02	0.41
2:W:271:SER:OG	2:W:272:ALA:N	2.52	0.41
2:N:156:GLY:O	2:N:303:THR:OG1	2.35	0.41
1:A:531:SER:HB2	2:U:265:SER:HB2	2.03	0.41
1:A:538:MET:H	1:A:538:MET:HG3	1.65	0.41
2:S:18:LYS:NZ	4:S:802:ADP:O1B	2.52	0.41
1:A:535:MET:HB2	1:A:630:ALA:HB2	2.03	0.41
2:J:271:SER:OG	2:J:272:ALA:N	2.52	0.41
2:S:36:GLY:HA3	2:S:65:LEU:HD23	2.02	0.41
1:A:554:LEU:HD11	1:A:610:ILE:HG22	2.03	0.41
1:I:619:LEU:O	1:I:623:ALA:N	2.53	0.41
2:L:156:GLY:O	2:L:303:THR:OG1	2.35	0.41
1:B:538:MET:H	1:B:538:MET:HG3	1.65	0.41
2:M:283:MET:O	2:M:290:ARG:NH2	2.46	0.41
2:S:357:ILE:HG12	2:S:373:LYS:HG3	2.03	0.41
1:I:535:MET:HB2	1:I:630:ALA:HB2	2.02	0.41
2:N:172:PRO:HG3	2:U:41:GLN:HE21	1.85	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:N:357:ILE:HG12	2:N:373:LYS:HG3	2.03	0.41
2:U:357:ILE:HG12	2:U:373:LYS:HG3	2.03	0.41
2:U:271:SER:OG	2:U:272:ALA:N	2.52	0.41
1:B:535:MET:HB2	1:B:630:ALA:HB2	2.03	0.40
1:B:554:LEU:HD11	1:B:610:ILE:HG22	2.03	0.40
2:W:357:ILE:HG12	2:W:373:LYS:HG3	2.03	0.40
2:N:271:SER:OG	2:N:272:ALA:N	2.52	0.40
2:R:357:ILE:HG12	2:R:373:LYS:HG3	2.03	0.40
1:C:535:MET:HB2	1:C:630:ALA:HB2	2.03	0.40
2:S:47:MET:N	2:S:47:MET:SD	2.95	0.40
2:J:36:GLY:HA3	2:J:65:LEU:HD23	2.02	0.40
2:L:169:TYR:HB3	2:S:40:HIS:HB2	2.02	0.40
1:B:522:GLN:HG2	1:B:615:GLN:NE2	2.37	0.40
1:C:619:LEU:O	1:C:623:ALA:N	2.53	0.40
2:J:53:TYR:CZ	2:J:61:LYS:HE2	2.57	0.40
2:R:36:GLY:HA3	2:R:65:LEU:HD23	2.02	0.40
2:R:53:TYR:CZ	2:R:61:LYS:HE2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/685 (23%)	144 (90%)	16 (10%)	0	100	100
1	B	160/685 (23%)	144 (90%)	16 (10%)	0	100	100
1	C	160/685 (23%)	144 (90%)	16 (10%)	0	100	100
1	I	160/685 (23%)	144 (90%)	16 (10%)	0	100	100
2	J	368/377 (98%)	353 (96%)	15 (4%)	0	100	100
2	L	368/377 (98%)	353 (96%)	15 (4%)	0	100	100
2	M	368/377 (98%)	354 (96%)	14 (4%)	0	100	100

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	N	368/377 (98%)	354 (96%)	14 (4%)	0	100	100
2	R	368/377 (98%)	353 (96%)	15 (4%)	0	100	100
2	S	368/377 (98%)	353 (96%)	15 (4%)	0	100	100
2	U	368/377 (98%)	353 (96%)	15 (4%)	0	100	100
2	W	368/377 (98%)	353 (96%)	15 (4%)	0	100	100
All	All	3584/5756 (62%)	3402 (95%)	182 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2   Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	128/580 (22%)	127 (99%)	1 (1%)	81	91
1	B	128/580 (22%)	127 (99%)	1 (1%)	81	91
1	C	128/580 (22%)	127 (99%)	1 (1%)	81	91
1	I	128/580 (22%)	127 (99%)	1 (1%)	81	91
2	J	313/319 (98%)	312 (100%)	1 (0%)	92	97
2	L	313/319 (98%)	312 (100%)	1 (0%)	92	97
2	M	313/319 (98%)	312 (100%)	1 (0%)	92	97
2	N	313/319 (98%)	312 (100%)	1 (0%)	92	97
2	R	313/319 (98%)	312 (100%)	1 (0%)	92	97
2	S	313/319 (98%)	312 (100%)	1 (0%)	92	97
2	U	313/319 (98%)	312 (100%)	1 (0%)	92	97
2	W	313/319 (98%)	312 (100%)	1 (0%)	92	97
All	All	3016/4872 (62%)	3004 (100%)	12 (0%)	91	97

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	566	ARG
1	A	566	ARG
1	C	566	ARG
1	I	566	ARG
2	J	355	MET
2	L	355	MET
2	M	355	MET
2	N	355	MET
2	R	355	MET
2	S	355	MET
2	U	355	MET
2	W	355	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	522	GLN
1	C	522	GLN
1	I	522	GLN
2	J	12	ASN
2	L	12	ASN
2	M	12	ASN
2	N	12	ASN
2	R	12	ASN
2	S	12	ASN
2	S	161	HIS
2	U	12	ASN
2	U	41	GLN
2	W	12	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	#$|Z| > 2$	Counts	RMSZ	#$|Z| > 2$
2	HIC	W	73	2	8,11,12	1.61	2 (25%)	6,14,16	1.56	1 (16%)
2	HIC	N	73	2	8,11,12	1.59	2 (25%)	6,14,16	1.55	1 (16%)
2	HIC	S	73	2	8,11,12	1.59	2 (25%)	6,14,16	1.54	1 (16%)
2	HIC	U	73	2	8,11,12	1.59	2 (25%)	6,14,16	1.55	1 (16%)
2	HIC	M	73	2	8,11,12	1.59	2 (25%)	6,14,16	1.55	1 (16%)
2	HIC	R	73	2	8,11,12	1.60	2 (25%)	6,14,16	1.53	1 (16%)
2	HIC	J	73	2	8,11,12	1.58	2 (25%)	6,14,16	1.54	1 (16%)
2	HIC	L	73	2	8,11,12	1.60	2 (25%)	6,14,16	1.56	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HIC	W	73	2	-	1/5/6/8	0/1/1/1
2	HIC	N	73	2	-	1/5/6/8	0/1/1/1
2	HIC	S	73	2	-	1/5/6/8	0/1/1/1
2	HIC	U	73	2	-	1/5/6/8	0/1/1/1
2	HIC	M	73	2	-	1/5/6/8	0/1/1/1
2	HIC	R	73	2	-	1/5/6/8	0/1/1/1
2	HIC	J	73	2	-	1/5/6/8	0/1/1/1
2	HIC	L	73	2	-	1/5/6/8	0/1/1/1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	W	73	HIC	CD2-CG	3.24	1.41	1.36
2	N	73	HIC	CD2-CG	3.23	1.41	1.36
2	U	73	HIC	CD2-CG	3.22	1.41	1.36
2	R	73	HIC	CD2-CG	3.21	1.41	1.36
2	S	73	HIC	CD2-CG	3.21	1.41	1.36
2	L	73	HIC	CD2-CG	3.21	1.41	1.36
2	M	73	HIC	CD2-CG	3.20	1.41	1.36
2	J	73	HIC	CD2-CG	3.16	1.40	1.36
2	S	73	HIC	CZ-NE2	-2.06	1.42	1.48
2	L	73	HIC	CZ-NE2	-2.04	1.42	1.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	M	73	HIC	CZ-NE2	-2.04	1.42	1.48
2	R	73	HIC	CZ-NE2	-2.04	1.42	1.48
2	W	73	HIC	CZ-NE2	-2.03	1.42	1.48
2	U	73	HIC	CZ-NE2	-2.03	1.42	1.48
2	J	73	HIC	CZ-NE2	-2.01	1.42	1.48
2	N	73	HIC	CZ-NE2	-2.00	1.42	1.48

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	M	73	HIC	CB-CA-C	-3.18	105.51	111.47
2	W	73	HIC	CB-CA-C	-3.17	105.52	111.47
2	L	73	HIC	CB-CA-C	-3.17	105.53	111.47
2	S	73	HIC	CB-CA-C	-3.16	105.53	111.47
2	J	73	HIC	CB-CA-C	-3.16	105.55	111.47
2	N	73	HIC	CB-CA-C	-3.16	105.55	111.47
2	U	73	HIC	CB-CA-C	-3.15	105.56	111.47
2	R	73	HIC	CB-CA-C	-3.14	105.58	111.47

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	J	73	HIC	CA-CB-CG-ND1
2	L	73	HIC	CA-CB-CG-ND1
2	M	73	HIC	CA-CB-CG-ND1
2	N	73	HIC	CA-CB-CG-ND1
2	R	73	HIC	CA-CB-CG-ND1
2	S	73	HIC	CA-CB-CG-ND1
2	U	73	HIC	CA-CB-CG-ND1
2	W	73	HIC	CA-CB-CG-ND1

There are no ring outliers.

No monomer is involved in short contacts.

5.5   Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	#$|Z| > 2$	Counts	RMSZ	#$|Z| > 2$
4	ADP	J	802	3	24,29,29	1.01	1 (4%)	29,45,45	1.56	3 (10%)
4	ADP	M	802	3	24,29,29	1.04	1 (4%)	29,45,45	1.41	3 (10%)
4	ADP	W	802	3	24,29,29	1.06	2 (8%)	29,45,45	1.47	3 (10%)
4	ADP	L	802	3	24,29,29	1.00	1 (4%)	29,45,45	1.48	3 (10%)
4	ADP	R	802	3	24,29,29	1.01	1 (4%)	29,45,45	1.48	4 (13%)
4	ADP	N	802	3	24,29,29	0.99	0	29,45,45	1.39	4 (13%)
4	ADP	S	802	3	24,29,29	1.00	1 (4%)	29,45,45	1.52	2 (6%)
4	ADP	U	802	3	24,29,29	1.11	1 (4%)	29,45,45	1.59	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	J	802	3	-	2/12/32/32	0/3/3/3
4	ADP	M	802	3	-	4/12/32/32	0/3/3/3
4	ADP	W	802	3	-	3/12/32/32	0/3/3/3
4	ADP	L	802	3	-	2/12/32/32	0/3/3/3
4	ADP	R	802	3	-	3/12/32/32	0/3/3/3
4	ADP	N	802	3	-	3/12/32/32	0/3/3/3
4	ADP	S	802	3	-	3/12/32/32	0/3/3/3
4	ADP	U	802	3	-	3/12/32/32	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	U	802	ADP	C2'-C1'	-2.54	1.49	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	R	802	ADP	C2'-C1'	-2.38	1.50	1.53
4	J	802	ADP	C2'-C1'	-2.37	1.50	1.53
4	W	802	ADP	C2'-C1'	-2.30	1.50	1.53
4	L	802	ADP	C2'-C1'	-2.22	1.50	1.53
4	M	802	ADP	C2'-C1'	-2.20	1.50	1.53
4	S	802	ADP	C2'-C1'	-2.09	1.50	1.53
4	W	802	ADP	C5-C4	2.00	1.46	1.40

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	S	802	ADP	PA-O3A-PB	-5.04	115.53	132.83
4	U	802	ADP	PA-O3A-PB	-4.65	116.88	132.83
4	J	802	ADP	PA-O3A-PB	-4.13	118.65	132.83
4	R	802	ADP	PA-O3A-PB	-3.99	119.14	132.83
4	W	802	ADP	PA-O3A-PB	-3.96	119.24	132.83
4	N	802	ADP	C3'-C2'-C1'	3.84	106.76	100.98
4	L	802	ADP	PA-O3A-PB	-3.84	119.65	132.83
4	M	802	ADP	C3'-C2'-C1'	3.76	106.63	100.98
4	J	802	ADP	C3'-C2'-C1'	3.65	106.48	100.98
4	L	802	ADP	C3'-C2'-C1'	3.65	106.48	100.98
4	M	802	ADP	PA-O3A-PB	-3.61	120.44	132.83
4	U	802	ADP	C3'-C2'-C1'	3.40	106.09	100.98
4	W	802	ADP	C3'-C2'-C1'	3.37	106.05	100.98
4	S	802	ADP	C3'-C2'-C1'	3.27	105.90	100.98
4	N	802	ADP	PA-O3A-PB	-3.17	121.95	132.83
4	R	802	ADP	C3'-C2'-C1'	3.12	105.68	100.98
4	J	802	ADP	N3-C2-N1	-2.87	124.19	128.68
4	R	802	ADP	N3-C2-N1	-2.66	124.51	128.68
4	L	802	ADP	N3-C2-N1	-2.44	124.86	128.68
4	N	802	ADP	N3-C2-N1	-2.37	124.97	128.68
4	M	802	ADP	N3-C2-N1	-2.23	125.19	128.68
4	R	802	ADP	C4-C5-N7	-2.22	107.09	109.40
4	W	802	ADP	N3-C2-N1	-2.17	125.29	128.68
4	N	802	ADP	C4-C5-N7	-2.13	107.18	109.40
4	U	802	ADP	N3-C2-N1	-2.12	125.37	128.68
4	U	802	ADP	C4-C5-N7	-2.05	107.26	109.40

There are no chirality outliers.

All (23) torsion outliers are listed below:

Preliminary Full wwPDB EM Validation Report
(*Not For Manuscript Review*)

Mol	Chain	Res	Type	Atoms
4	J	802	ADP	C5'-O5'-PA-O3A
4	L	802	ADP	C5'-O5'-PA-O3A
4	M	802	ADP	C5'-O5'-PA-O3A
4	M	802	ADP	C3'-C4'-C5'-O5'
4	N	802	ADP	C5'-O5'-PA-O2A
4	R	802	ADP	C5'-O5'-PA-O3A
4	S	802	ADP	C5'-O5'-PA-O1A
4	U	802	ADP	C5'-O5'-PA-O3A
4	W	802	ADP	C5'-O5'-PA-O1A
4	W	802	ADP	C5'-O5'-PA-O2A
4	M	802	ADP	O4'-C4'-C5'-O5'
4	R	802	ADP	C3'-C4'-C5'-O5'
4	N	802	ADP	C5'-O5'-PA-O3A
4	J	802	ADP	C5'-O5'-PA-O1A
4	L	802	ADP	C5'-O5'-PA-O1A
4	M	802	ADP	C5'-O5'-PA-O1A
4	N	802	ADP	C5'-O5'-PA-O1A
4	R	802	ADP	C5'-O5'-PA-O1A
4	U	802	ADP	C5'-O5'-PA-O1A
4	S	802	ADP	C3'-C4'-C5'-O5'
4	S	802	ADP	C5'-O5'-PA-O3A
4	W	802	ADP	C5'-O5'-PA-O3A
4	U	802	ADP	C3'-C4'-C5'-O5'

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	J	802	ADP	2	0
4	L	802	ADP	2	0
4	S	802	ADP	3	0
4	U	802	ADP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand ADP J 802

Bond lengths

Bond angles

Torsions

Rings

Ligand ADP M 802

Bond lengths

Bond angles

Torsions

Rings

Ligand ADP W 802

Bond lengths

Bond angles

Torsions

Rings

Ligand ADP L 802

Bond lengths

Bond angles

Torsions

Rings

Ligand ADP R 802

Bond lengths

Bond angles

Torsions

Rings

Ligand ADP N 802

Bond lengths

Bond angles

Torsions

Rings

Ligand ADP S 802

Bond lengths	Bond angles
Torsions	Rings

Ligand ADP U 802

Bond lengths	Bond angles
Torsions	Rings

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6   Map visualisation ⓘ

This section contains visualisations of the EMDB entry D_1000280011. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1   Orthogonal projections ⓘ

6.1.1   Primary map

X   Y   Z

6.1.2   Raw map

X   Y   Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices ⓘ

6.2.1 Primary map

X Index: 160	Y Index: 160	Z Index: 160

6.2.2 Raw map

X Index: 160	Y Index: 160	Z Index: 160

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices ⓘ

6.3.1 Primary map

X Index: 168 Y Index: 152 Z Index: 153

6.3.2 Raw map

X Index: 168 Y Index: 152 Z Index: 153

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) ⓘ

6.4.1 Primary map

X

Y

Z

6.4.2 Raw map

X

Y

Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views ⓘ

6.5.1 Primary map

X   Y   Z

The images above show the 3D surface view of the map at the recommended contour level 0.0434. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map

X   Y   Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation ⓘ

This section was not generated. No masks/segmentation were deposited.

7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)

The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate ⓘ

The volume at the recommended contour level is 345 nm^3; this corresponds to an approximate mass of 311 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3   Rotationally averaged power spectrum ⓘ

Rotationally averaged power spectrum

*Reported resolution corresponds to spatial frequency of 0.278 Å^{-1}

8   Fourier-Shell correlation ⓘ

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1   FSC ⓘ

FSC

(Plot of Correlation vs Spatial frequency (Å⁻¹))

Legend:
- Unmasked-calculated FSC
- 0.143
- 0.5
- Half-bit
- Reported resolution 3.60 Å*

*Reported resolution corresponds to spatial frequency of 0.278 $\mathring{A}^{-1}$

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.60	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.37	7.52	4.51

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.37 differs from the reported value 3.6 by more than 10 %

9 Map-model fit ⓘ

This section contains information regarding the fit between EMDB map D_1000280011 and PDB model D_1000280011. Per-residue inclusion information can be found in section 3 on page 7.

9.1 Map-model overlay ⓘ

X   Y   Z

The images above show the 3D surface view of the map at the recommended contour level 0.0434 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2   Q-score mapped to coordinate model ⓘ

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3   Atom inclusion mapped to coordinate model ⓘ

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0434).

9.4 Atom inclusion (i)

At the recommended contour level, 97% of all backbone atoms, 82% of all non-hydrogen atoms, are inside the map.

Preliminary Full wwPDB EM Validation Report

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0434) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8210	0.3280
A	0.7630	0.2680
B	0.6570	0.1810
C	0.6800	0.2270
I	0.6230	0.1910
J	0.8510	0.3520
L	0.8570	0.3610
M	0.8660	0.3760
N	0.8520	0.3580
R	0.8350	0.3270
S	0.8590	0.3620
U	0.8580	0.3610
W	0.8260	0.3150

1.0

0.0

<0.0