1 Title: **The human pupil and face encode sound affect and provide objective signatures of**
2 **tinnitus and auditory hypersensitivity disorders**
3
4

5 Authors: Samuel S. Smith[1,2,3]*, Kelly N. Jahn[1,2]*^, Jenna A. Sugai[1]*, Ken E. Hancock[1,2], Daniel
6 B. Polley[1,2,4]
7
8 1 – Eaton-Peabody Laboratories, Massachusetts Eye and Ear, Boston MA, 02114 USA
9 2 – Department of Otolaryngology – Head and Neck Surgery, Harvard Medical School, Boston
10 MA 02114 USA
11 3 – Lead contact
12 4 – Corresponding author
13
14 * Equal contribution
15 ^ Current address: Department of Speech, Language, and Hearing, The University of Texas at
16 Dallas, 1966 Inwood Road, Dallas, TX 75235

17
18
19

Summary

Central disinhibition works like an amplifier, boosting neural sensitivity to progressively weaker
peripheral inputs arising from degenerating sensory organs. The Excess Central Gain model
posits that central disinhibition also gives rise to cardinal features of sensory disorders
including sensory overload and phantom percepts. Here, we tested predictions of this model
with neural, autonomic, and behavioral approaches in participants with sound sensitivity and
tinnitus (phantom ringing). We confirmed enhanced auditory neural gain but found no
association with their sound aversion and anxiety. Instead, we hypothesized that symptom
severity was linked to affective sound encoding. In neurotypical controls, emotionally evocative
sounds elicited pupil dilations, electrodermal responses and facial reactions that scaled with
valence. Participants with disordered hearing exhibited disrupted pupil and facial reactivity that
accurately predicted their self-reported tinnitus and hyperacusis severity. These findings
highlight auditory-limbic dysregulation in tinnitus and sound sensitivity disorders and introduce
approaches for their objective measurement.

34
35
36

Introduction

Damage or degeneration of peripheral sensory organs causes reduced sensitivity to
environmental stimuli. It can also precipitate the opposite outcome: a hypersensitivity to environmental
stimuli along with the perception of phantom stimuli that have no physical source in the environment.
Sensory phantoms occur in all modalities but are more often heard than seen, tasted, felt, or smelled.
Approximately 12% of adults hear an indefatigable phantom sound every day of their waking life.
Tinnitus typically manifests as a continuous ringing, whooshing, roaring, or sizzling phantom sound that
is often accompanied by a generalized sensitivity and discomfort with moderately intense
environmental sounds[1,2]. Most evidence suggests that disinhibition in auditory processing centers of the
brain is a proximal cause of tinnitus and sound sensitivity[3–7]. Central auditory disinhibition related to
tinnitus and sound sensitivity can arise from normal aging[8,9], as a compensatory response to hearing
loss and auditory nerve degeneration[10–19], from traumatic injury of the brain or cervical ganglia[20–22], or

simply from the abrupt cessation of prescription GABA agonists[23,24]. Whatever the source, the Excess
Central Gain model posits that disinhibition has the knock-on effect of increasing the synchrony and
activity rate of local excitatory neurons in silence (the basis of the phantom sound) as well as
disproportionately strong responses to moderately intense sound (the basis of loudness
hypersensitivity)[25–29]. As expected, boosting levels of central inhibition through direct activation
protocols in animals or GABA-acting drugs in humans can mitigate tinnitus symptoms[30,31].
In some cases, persons with tinnitus and sound sensitivity also present with an intense
generalized aversion to sound, anxiety about sound exposure, social withdrawal, depression, and even
suicidal ideations[32]. It is unclear how central gain in low-level auditory processing is related to the
anxiety and mood disruptions that can accompany these disorders. One possibility is that the
connection between elevated central gain and affective dysregulation may literally reflect abnormally
strong coupling between auditory forebrain and limbic centers[33]. A recent animal study of auditory
threat learning has shown that a selective plasticity in auditory corticoamygdalar projection neurons
asymmetrically increased corticofugal spike-LFP coupling and sound-evoked activity in the lateral
amygdala[34], suggesting that increased neural gain in cortical projection neurons can be passed on to
postsynaptic brain regions that regulate affective evaluation of valence and arousal. To this point,
several neuroimaging studies in participants with tinnitus and sound sensitivity have identified
abnormally strong coupling between auditory cortex and the amygdala, insula, anterior cingulate cortex,
and medial prefrontal cortex[33,35–37]. Whereas psychoacoustic and low-level auditory assays were
generally not correlated with extra-auditory features of sound aversion and psychological burden[38–43],
neuroimaging assays of enhanced coupling with extra-auditory networks have shown stronger
correlations with individual differences in tinnitus and sound sensitivity severity[37,44].
There are no objective measurements for the severity of tinnitus and sound sensitivity disorders.
Instead, assessments rely on subjective self-report questionnaires, which can introduce vulnerabilities
for malingering and false disability claims[39] and offer less insight into the underlying causes and
potential treatments. Here, we hypothesized that involuntary autonomic and behavioral responses may
have untapped potential as non-invasive, objective markers of tinnitus and sound sensitivity severity
that are relatively easily to implement in laboratory and clinical settings. Autonomic responses (e.g.,
pupil dilation and skin conductance) and involuntary facial movements provide a wealth of information
into affective processing of valence and arousal in humans[45–48] as well as other animals[34,49]. Although
human studies have largely relied on visual stimuli, it is clear that speech, music and other sounds
provide a rich medium for conveying valence and arousal cues. This led us to ask whether emotionally
evocative sounds elicit autonomic and facial responses that are modulated by perceived valence. We
then evaluated whether these objective measures could identify a bias towards negative valence and
hyper-arousal in persons with tinnitus and sound sensitivity that might more accurately predict the
severity of sound aversion and psychological burden they report.
85
86
87
88   **Results**
Of 196 adults recruited to our study, we identified 71 eligible participants with normal hearing
thresholds who completed the full course of testing (**Figure S1A**). An experienced clinician assigned
participants with chronic tinnitus and/or auditory hypersensitivity to the disordered hearing (DH, N=35)
group and participants without tinnitus or auditory hypersensitivity to the neurotypical (NT, N=36) group.
The distribution of age and hearing thresholds were closely matched between DH and NT groups,
though, as expected, uncomfortable listening level thresholds in DH subjects were significantly lower

than NT subjects (**Figure S1B-C**, description of statistical testing described in the figure legends throughout).

Increased central gain is not correlated tinnitus or sound sensitivity burden

The Enhanced Central Gain model posits that disinhibition in the auditory cortex and subcortical auditory structures produces hypersynchronized and hyperactive population activity among excitatory neurons in silence and hyperresponsivity to sounds of increasing physical intensity (**Figure 1A**). Although we could not directly measure the rate or synchrony of spiking, we were able to directly measure the gain of the intensity versus neural response growth function through an approach that slowly increased and decreased the intensity of a 40 Hz amplitude modulated tone (**Figure 1B**, top). Scalp EEG recordings revealed a synchronized envelope following response (EFR) at 40 Hz that increased and decreased in amplitude with the change in sound intensity (**Figure 1B**, bottom). Gain (i.e., the output per unit step in input) could be explicitly measured as the slope of the neural growth function. As predicted by the Excess Central Gain model, neural activity grew significantly more steeply with increasing sound level in DH participants (**Figure 1C**).

Tinnitus and sound sensitivity are heterogenous co-morbid conditions. The cohort of DH participants presented with varying degrees of sound aversion, distress, anxiety and depression related to their disorder, as identified by standard clinical questionnaires for hyperacusis and tinnitus (HQ and THI, respectively, **Figure 1D**). This variability provided us with a valuable opportunity to study the relationship between excess central gain and psychological burden. Although central gain was significantly increased at a group level in DH participants (**Figure 1E**, top), it was not correlated with individual differences in hyperacusis or tinnitus severity (**Figures 1F and 1G**, respectively).

As illustrated in the cartoon model (**Figure 1A**), an extension of the Excess Central Gain model posits that hyperactive, hypersynchronized, and hyperresponsive projection neurons in the central auditory pathway drive abnormally strong functional connectivity and auditory recruitment in downstream limbic networks. Limbic hyperresponsivity, in turn, would elicit abnormal autonomic response for sounds that would not otherwise be encoded as unpleasant or highly arousing. In essence, a failure to homeostatically regulate neural activity in auditory brain networks disrupts allostatic processes that balance autonomic and behavioral responses to stressors. In this model, excess central gain is a distal, upstream precipitator of limbic-autonomic dysfunction that more directly determines whether an individual feels utterly debilitated versus slightly annoyed by their tinnitus. As such, central gain would not be expected to be highly correlated with the tinnitus or hyperacusis burden reported in surveys, but a more direct marker of limbic or autonomic dysfunction would.

Pupil-indexed affective processing is correlated with tinnitus and sound sensitivity burden

In constant light levels, pupil diameter provides an autonomic marker of brain-wide neuromodulator release[50,51] that, depending on task design, can index executive load[52], or affective valence and arousal[53]. To determine whether autonomic and behavioral evaluation of sound affect was biased towards negative valence and hyper-arousal in DH subjects, we measured pupil diameter and skin conductance while subjects listened to 6s audio clips from an established database of emotionally evocative sounds (**Figure 2A**)[54]. These sounds spanned a wide valence range, including sounds rated as highly unpleasant (vomiting, screaming) to pleasant (music) (**Figure 2B**). On average, selected sounds smoothly tiled the valence range (**Figure S2A**) but were idiosyncratic at the level of individual participants, presumably reflecting their individual hedonic sound associations (**Figure S2B**). When individually ordered by valence rating, we found that DH subjects rated sounds as less pleasant overall, particularly relatively pleasant or neutral sounds, suggesting a complementary metric to the generalized sound aversion captured by subjective self-report questionnaires[55] (**Figure 2C**).

142   These emotionally evocative sounds elicited pupil dilations within approximately 0.5s followed
143 by a slower increase in sound-evoked skin conductance (**Figure 2D; Figure S3**). Pupil dilation scaled
144 with self-reported sound valence across all participants but was significantly greater overall in DH
145 subjects, indicating a hypersensitized autonomic response to affective arousal (**Figure 2E**). Sound-
146 evoked changes in skin conductance also scaled with self-reported valence but did not differ between
147 groups (**Figure 2F**). Like the EEG measure of central gain (**Figure 1E**), sound-evoked pupil dilation
148 was also significantly elevated in DH participants (**Figure 2G**). Unlike central gain, the degree of sound-
149 evoked pupil was significantly correlated with self-reported hyperacusis burden (**Figure 2H**) and tinnitus
150 burden (**Figure 2I**). Average sound-evoked skin conductance did not differ between groups and was
151 not correlated with tinnitus or hyperacusis severity (**Figure 2J-L**).
152
153 **Pupil hyper-responsivity is specific to affective processing**
154   These findings support our hypothesis that affective sound encoding is disrupted in persons with
155 debilitating tinnitus and sound sensitivity, which can be measured through autonomic markers of
156 affective arousal such as pupil dilation. Two negative control experiments support the assertion that
157 pupil-indexed hyper-arousal was specific to affective processing. First, pupil dynamics in DH and NT
158 participants were indistinguishable when entrained to periodic changes in light intensity (**Figure S4A-
159 B**). Second, we measured pupil-indexed listening effort in a multi-talker speech intelligibility task
160 (**Figure S4C**) and observed reduced behavioral accuracy and increased pupil diameter at a more
161 challenging signal to noise ratio (**Figure S4D**). Importantly, behavioral accuracy and pupil-indexed
162 listening effort were equivalent between NT and DH participants, confirming that DH subjects do not
163 have widespread deficits in challenging listening tasks[56] or global autonomic regulation, but rather a
164 more specific behavioral and autonomic aversion to the affective quality of sound (**Figure S4E-F**).
165
166 **Subtle facial movements identify auditory valence and hearing disorder severity**
167   To identify additional windows into auditory affective processing that might complement or
168 extend upon pupil dynamics we turned our attention to subtle and rapid movements of the face. Facial
169 movements have a long history in human emotion research, though these studies have almost
170 exclusively relied on visual stimuli to convey affective cues[57]. To establish whether emotionally
171 evocative sounds would also produce facial movements related to affective valence and arousal, we
172 quantified high-resolution facial videography recordings that were performed in tandem with the pupil
173 and electrodermal recordings (**Figure 3A**). We developed an analysis pipeline to localize changes in
174 facial texture and observed that sounds do elicit subtle and rapid facial movements (**Figure 3B**). For
175 example, in one instance an unpleasant "buzz" sound elicited a tightening of the muscles around the
176 eyes and forehead (**Figure 3C**, left), whereas a more pleasant sound elicited a longer latency
177 movement of the mouth and jaw (**Figure 3C**, right). Amongst this spatiotemporal complexity, we found
178 stereotyped patterns that were associated with individual valence ratings. Pleasant sounds evoked
179 increased movements in the area of the zygomaticus major, a muscle involved in shaping the corners
180 of the lips (**Figure 3D**, left). Unpleasant sounds triggered increased movements globally but were
181 particularly robust around the eyes and brows (**Figure 3D**, middle).
182   Like the pupil, we hypothesized that sound-evoked facial expressions would convey
183 downstream limbic dysfunction in individuals with tinnitus and sound sensitivity. This prediction was
184 upheld but, interestingly, in the opposite direction of changes in pupil dilation. Whereas facial
185 movement amplitude scaled with valence in NT participants, DH participants exhibited a generally
186 blunted affect that did not change across self-reported valence (**Figure 3E-G**). Importantly, reduced
187 facial movements were significantly correlated with individual differences in sound sensitivity burden
188 (**Figure 3H**) and tinnitus burden (**Figure 3I**). Together, these results show that the face encodes not

189   only sound affect but can also provide a window into the intense aversion and lifestyle burden that can
190   accompany tinnitus and sound sensitivity.
191
192   **Combined measures of sound affect best determine psychological burden**
193   Here, we have identified neural (**Figure 1C**), autonomic (**Figure 2E**), voluntary behavioral
(**Figure 2C**), and involuntary behavioral (**Figure 3F**) measures that distinguish DH and NT subjects at a
group level. A major goal for research on sensory disorders is to identify and refine a set of objective
measurements – akin to what tumor imaging and biopsy data provide for oncologists or EEG measures
of epileptiform activity provide to neurologists – that inform future care providers about the subtype and
severity of the sensory disorder, the likelihood to benefit from a particular treatment, or whether they
have demonstrated an improvement subsequent to treatment. While some measurements have been
developed to identify whether an individual has tinnitus, they are insensitive to severity and lifestyle
burden[39,58,59]. Here, we found that pupil dilation and facial reactions both demonstrated a correlation
with symptom severity, though it was not clear whether these measures were redundant, nor how
accurately they could predict an individual's self-reported severity.
204   To this end, we incorporated all measures where DH and NT cohorts were significantly different
at a group level (pupil diameter, facial movement, behavioral valence rating, central gain) as well as
potential co-variates of lifestyle burden (age) in an elastic net regression analysis, developed to be
robust to superfluous predictor variables (see Methods). For the severity of self-reported sound
sensitivity, a cross-validated fitting procedure settled on an optimal model that combined measures of
affective responses – pupil dilation, facial movement, and behavioral valence rating (**Figure 4B**). This
model was significantly correlated with hyperacusis questionnaire scores and was moderately accurate
in predicting the hyperacusis index score compiled across NH and DT participants ($R = 0.6/R^2 = 0.36$,
Figure 4C). When the same approach was applied to tinnitus, age, we noted that central gain, and
behavioral valence rating were minimally weighted, a moderate weight was afforded to pupil dilation,
but facial reactivity was heavily weighted (**Figure 4D-E**). This model was highly correlated with self-
reported severity and exhibited fairly high accuracy in predicting individual tinnitus burden scores ($R =$
$0.78/ R^2 = 0.61$, **Figure 4D-F**). These findings demonstrate that pupil and facial reactivity to emotionally
evocative sounds capture distinct features of generalized aversion, distress and anxiety that can
accompany tinnitus and sound sensitivity. Although autonomic measures and affective processing have
rarely been considered as biomarkers for these conditions, predictions of hyperacusis and tinnitus
severity scores were significantly poorer when they were left out and the optimal model instead relied
on neural and voluntary behavioral measures (likelihood ratio test, $p<0.001$).
222
223
224
225   **Discussion**
226   **Extending upon the Excess Central Gain model of tinnitus and sound sensitivity**
227   Pioneering neuroimaging studies in human subjects with tinnitus and reduced sound level
tolerance identified auditory hyper-responsivity in the inferior colliculus and auditory cortex as the
proverbial "ghost in the machine"[60], i.e., a biological substrate for these perceptual disorders[42,61].
Subsequent findings in human subjects and animal models highlighted the juxtaposition of normal or
even hyper-responsive event-related potentials arising from later stages with reduced sound-evoked
potentials arising from the auditory nerve[12,15,62] Findings of this ilk laid the foundation for the Enhanced
Central Gain theory for tinnitus and sound sensitivity, wherein central auditory circuits leverage
disinhibition to compensate for reduced bottom-up input. In so doing, vulnerabilities for hyper-

synchrony, hyperactivity, and abnormally steep sound intensity growth functions are introduced that
generate phantom sounds and reduced tolerance of moderately intense sounds[25,26]. Here, we used
40Hz amplitude modulation to enhance the relative contribution of the auditory cortex to the EFR[63] and
derived the instantaneous amplitude as the sound level swept up and down across a 70 dB range to
provide a direct demonstration of enhanced central gain (i.e., increased neural output per unit step in
sound input) in participants with tinnitus and reduced sound level tolerance.
Disinhibition and enhanced central gain at later stages of the auditory pathway may generate
the sensory qualities of the phantom sound and disproportionate loudness but – in and of itself – is
unlikely to account for the attentional and affective sequelae of tinnitus and sound sensitivity. Central
gain does not directly account for why the phantom sound effortlessly recedes from conscious
awareness when not attended to for some but is irrepressible for others? Likewise, central gain does
not account for why internally and externally generated sounds a mild nuisance for some but are
irritating, overwhelming, and anxiety-producing in others? That central gain does not account for these
qualities does not mean that phenomenological models featuring auditory nerve degeneration and
enhanced central gain are wrong, but rather that they are incomplete[64].
Excitatory neurons from primary and higher-order auditory cortex project widely throughout
prefrontal cortex executive control networks and limbic brain centers in neocortex and amygdala[65–67].
While relatively complex multi-regional feedback loops[68], altered perceptual inference mechanisms[69], or
stochastic resonance[70] may account for the attentional and affective components of tinnitus and sound
sensitivity disorder, a more parsimonious explanation that directly builds upon the Excess Central Gain
model is simply that hyperactive auditory efferent projections into extra-auditory executive and limbic
networks produce functional hyperconnectivity and knock-on dysregulation in these networks, and this
dysregulation more directly accounts for the attentional and affective components of these disorders,
respectively. Validation for this model is already evidenced by demonstrations of enhanced functional
connectivity in animals and humans, though deeper insights into the nature of dysregulated
spontaneous and sound-evoked processing in these regions awaits direct circuit interrogation
approaches in animals and non-invasive imaging in humans. Importantly, indirect but objective indices
of affective and executive disruptions do not require specialized, costly, and low-throughput brain
imaging systems. Pupil diameter, eye position, and involuntary facial reactivity have a well-established
relationship with executive and affective processing and task designs to probe these relationships in
neurotypical and neurodivergent populations have been the subject of study for many decades[49,53,71].
Here, we adapted pupil and facial assays of affective processing to the auditory modality and
demonstrated their improved ability to account for the affective component of tinnitus and sound
sensitivity disorder. While pupil and behavioral measurements reported here and elsewhere did not
suggest deficits related to executive load or listening effort[56], the attentional component of tinnitus
disorder awaits more direct interrogation with paradigms that probe the persistence of attention to
invariant sensory features with varying implicit significance.
272
Limitations of the current study
One general limitation of observational research on human sensory disorders relates to
uncertainty about the underlying precipitators of tinnitus and sound sensitivity in our subjects. Hearing
loss, age, and certain medications can directly impact central gain and tinnitus[10-24], motivating us to
strictly screen and match these variables in NT and DH cohorts to control for a confounding influence.
Nevertheless, the underlying cause and duration of tinnitus and sound sensitivity in our DH cohort is
unknown and may contribute to the heterogeneity reported here. Another limitation is that the EEG,
pupil, and facial measures described here provide little insight into their underlying generators in the
central and autonomic nervous systems. We leveraged the excellent temporal resolution of EEG to

derive measures of central gain that would be impossible with measures like fMRI but acknowledge that
non-invasive imaging approaches could significantly extend and enrich the observations reported here.
Another consideration is that while optimized multi-variate models featuring these objective markers of
affective processing captured a sizeable fraction of the variance in clinical outcome measures and
clearly outstripped other predictors studied here, these linear models fall short of providing the
sensitivity and selectivity required of a diagnostic measure. One important point to consider here is that
predictive accuracy cannot exceed the internal noise (e.g., test re-test reliability) and internal validity of
the outcome measure. Whether because tinnitus and sound sensitivity are inherently dynamic disorders
that ebb and flow over time or because the structure of the questionnaires elicit variable answers, the
upper bound of the predictive accuracy reported here is capped by the reliability of the outcome
measures themselves[72]. One of the underlying motivations for our study was to develop alternatives to
self-report questionnaires, not only because of their implicit subjectivity but also because their validity
as an instrument to probe the psychological burden arising from their tinnitus and hyperacusis
independent of their broader mental state is not certain[73,74]. We propose that the predictive accuracy
will be further improved by distilling the affective sound features used in the test battery, and combining
these refined stimulus sets with ecological momentary assessment approaches that embrace the
inherent dynamics of these disorders instead of reducing these subjects to a single measure of central
tendency.
300
Broader implications for clinical research on sensory disorders
The affective biomarkers described here were non-invasive and could be measured without
specialized equipment, suggesting a promising approach for complementing subjective self-report
questionnaires in clinical settings. These measures could prove to be useful biomarkers for other
neurological disorders where auditory aversion is a prominent clinical feature, particularly for
neurodevelopmental disorders such as autism spectrum disorder, where participant age or language
impairment may preclude objective assessments based on questionnaires[75,76]. Objective physiological
or neuroimaging biomarkers have been essential for the development of effective treatments for
epilepsy, stroke, and other neurological conditions. Better established objective biomarkers for sensory
overload will accelerate the pace of progress for clinical research on therapies for this debilitating
condition[77].

Supplemental Information
A supplementary document consisting of four supplemental figures is available to download.

Acknowledgements
This research was supported by research awards from the National Institute on Deafness and Other Communication Disorders to D.P. (P50DC015857) and K.J. (K01DC019647). Additional support for K.J. was provided by an ASHFoundation New Investigators Research Grant.

Author contributions
K.J. and D.P. conceived the project and designed the experiments. J.S. collected the data using software programmed by K.H and supported by K.J. and S.S. Data analysis was led by S.S. with contributions from K.J. Figure preparation and manuscript writing was performed by S.S. and D.P. with input from all authors.

Declaration of Interests
The authors declare no competing interests.

Figure 1. Increased central gain is not correlated tinnitus or sound sensitivity burden

A) Cartoon denotes a stage of central auditory processing (e.g., the auditory cortex) with excitatory projection neurons (red) and inhibitory interneurons (cool colors). In this model, disinhibition of excitatory neurons promotes elevated, hypersynchronous firing in silence (the purported generator of the phantom sound) and a steeper growth in spiking with sounds of increasing intensity (i.e., excess central gain, the purported generator of loudness hyperacusis). Hyperactive auditory projection neurons feed into downstream centers of limbic processing and autonomic regulation but, as distal upstream precipitator, excess central gain is less predictive of individual differences in psychoaffective burden than autonomic affective markers.

B) *Top:* Cartoon denotes the 64-channel array of scalp EEG electrodes and activity from a central electrode corresponding to the increasing intensity of a 40Hz amplitude modulated tone. Note that EEG amplitude is synchronized to the amplitude modulation rate.

Bottom: Spectrogram plots the amplitude of synchronized EEG activity across frequencies and time as the amplitude modulated 2kHz tone slowly increases and decreases across a 70 dB range. Note the rise and fall of the 40Hz envelope following response (EFR) amplitude as a function of time/sound intensity.

C) EFR growth as a function of sound intensity relative to the 2kHz audibility threshold measured for each participant (i.e., the sensation level, SL). NT and DH are neurotypical and disordered hearing participants, respectively. Central gain was measured as the change in neural response over a 25 dB change in sound level.

D) Hyperacusis and tinnitus severity for all participants based on Hyperacusis questionnaire (HQ) and Tinnitus Handicap Index (THI) scores, respectively (N = 35/35 NT/DH). Circles denote individual participants. Marginal distributions for each group are shown as normalized density functions. All participants can provide a meaningful HQ score but only participants with tinnitus can provide a meaningful THI value.

E) Central gain is significantly elevated in DH participants (two-sample t-test, p = 0.009, N = 36/33 NT/DH). Density functions display the central gain measure for each participant (individual circle) and sample mean (vertical lines).

357   **F)** Central gain is not correlated with hyperacusis severity (Pearson R = 0.16, p = 0.19, N = 68).

358   Shaded region denotes the 95% confidence interval. Solid line denotes linear fit. Circles denote

359   individual participants.

360   **G)** Central gain is not correlated with tinnitus severity (Pearson R = 0.13, p = 0.52, N = 22). Plotting

361   conventions as per *e.* Note that THI values are limited to participants with tinnitus.

Figure 2. Pupil-indexed affective processing is correlated with tinnitus and sound sensitivity burden

A) Schematic of trial design and experimental setup for combined autonomic and behavioral evaluation of sound valence. Alongside are spectrograms of six representative sounds filtered with a gammatone filter bank.

B) Mean behavioral valence rating of 60 sounds drawn from the IADS affective sound library (low/high scores indicate pleasant/unpleasant). Sounds are shown rank ordered and color-coded by labeled category.

C) Mean individually rank-ordered valence ratings demonstrate a significant bias overall towards unpleasant ratings in DH participants (Wilcoxon rank sum, p = 0.005, N = 36/35 NT/DH). When discretized into pleasant, neutral, and unpleasant categories (blue/gray/red, respectively), valence ratings were significantly elevated for pleasant and neutral sounds (asterisks, Wilcoxon rank sum, p = 0.026 and p = 0.048, respectively).

D) Mean ± SEM pupil diameter and skin conductance for all sounds during the 6s stimulus period. Responses are grouped by NT and DH participants.

E) Sound-evoked pupil dilations were larger overall in DH participants (two-sample t-test, p = 0.03, N=32/35 NT/DH). Pupil dilations were significantly increased for pleasant and unpleasant sounds (two-sample t-tests, p = 0.01 and 0.03, respectively) in DH compared to NT subjects.

F) Skin conductance did not differ overall (two-sample t-test, p = 0.59, N = 24/30 NT/DH) or as a
function of valence (two-sample t-tests, p > 0.5 for each valence category) in DH compared to NT
subjects.
G) Density functions display the sound-evoked pupil response averaged for each participant (individual
circle) and sample mean (vertical lines).
H) Pupil size is correlated with HQ score (Pearson R = 0.33, p = 0.006, N = 66). Shaded region
denotes the 95% confidence interval. Solid line denotes linear fit. Circles denote individual participants.
I) Pupil size is correlated with THI score (Pearson R = 0.56, p = 0.005, N = 24).
J) Density functions display the skin conductance response averaged for each participant (individual
circle) and sample mean (vertical lines).
K) Skin conductance is not significantly correlated with HQ score (Pearson R = 0.07, p = 0.63, N = 53).
Shaded region denotes the 95% confidence interval. Solid line denotes linear fit. Circles denote
individual participants.
L) Skin conductance is not significantly correlated with THI score (Pearson R = -0.03, p = 0.92, N = 20).

Figure 3. Subtle facial movements identify auditory valence and hearing disorder severity

A) Facial videography experimental setup.

B) Videos were processed with a custom analysis pipeline incorporating a face-mesh mapping and subsequent pixel quantification (see Methods).

C) Single trial facial movement amplitudes in two participants illustrate differences in amplitude and time course for an unpleasant (*left*) and pleasant (*right*) sound.

D) T-score contrast maps of differences in facial movements for unpleasant vs neutral (*left*), pleasant vs. neutral (*middle*), and for NT vs. DT participants (*right*) during the 6s stimulus period.

E) Mean ± SEM facial movement for all sounds during the 6s stimulus period. Responses are grouped into NT/DH.

F) Facial reactions were reduced overall in DH participants (two-sample t-test, p = 0.046, N = 36/33 NT/DH) and were significantly reduced for unpleasant sounds (two-sample t-test, p = 0.003) in DH compared to NT subjects.

G) Density functions display sound-evoked facial movements averaged for each participant (individual circle) and sample mean (vertical lines).

H) Facial movement is negatively correlated with HQ score (Pearson R = -0.31, p = 0.0096, N = 68). Shaded region denotes the 95% confidence interval. Solid line denotes linear fit. Circles denote individual participants.

I) Facial movement is negatively correlated with THI score (Pearson R = -0.42, p = 0.042, N = 23).

415
Figure 4. Combined measures of sound affect best determine psychological burden

A) An elastic net regression model was fit to individual hyperacusis severity scores (HQ scores). A tuning parameter, λ, controls the extent to which the coefficients contributing least to predictive accuracy are suppressed.

B) A radar plot displays the coefficient weight (i.e., |β|) in the optimal model of HQ score (filled shape), The optimal model with minimal cross-validated error retained non-zero coefficients for pupil, facial movement, and valence ratings.

C) The HQ scores predicted via elastic-net regression versus participants' actual scores.

D) Plotting conventions as per *A*, for tinnitus burden (THI) scores.

E) Plotting conventions as per *B*, for THI scores. Predictors from the optimal elastic-net featured physiological measures of sound affect (face, pupil) as the highest weighted coefficients.

F) Plotting conventions as per *C*, for THI scores.

Methods

428

429

430 **Experimental model and subject details.**

431 All procedures were approved by the Mass General Brigham Institutional Review Board.

432 *Participants.* Data are from 71 adults between 19 and 60 years of age (mean age = 33.9 years, 42
433 females) that were fluent in English. Participants were recruited as part of a larger study through flyers,
434 word of mouth, and by posting to the Mass General Brigham participant recruitment website. As
435 detailed in **Figure S1A**, screening and grouping of the 196 potentially eligible participants were
436 performed by licensed clinicians. Participants were required to have normal hearing (unremarkable
437 otoscopic evaluation and air conduction thresholds for tones 0.25 to 8 kHz ≤ 25 dB HL, 75 participants
438 excluded). Participants were required to have normal cognitive function (telephone Montreal Cognitive
439 Assessment, t-MOCA ≥18). We assessed mental health status and an ability to tolerate sound stimuli
440 used in our experiments based on their response to question 24 in Beck's Depression Inventory,
441 question 23 of the Tinnitus Reactivity Questionnaire, and question 23 of the Sound Reactivity
442 Questionnaire (10 participants excluded). Of the remaining participants, 16 were lost to follow-up after
443 the initial orientation and four were excluded for not completing the full course of testing.
444 Participants were assigned to neurotypical or disordered hearing groups by experienced
445 clinicians based on their responses to an open-ended questionnaire and clinical evaluation. Participants
446 were excluded if they had catastrophic tinnitus (> 77 on Tinnitus Handicap Index, THI, 1 participant
447 excluded), intermittent tinnitus (14 participants excluded), or inconsistently reported having tinnitus
448 across questionnaires (4 participants excluded). Of the remaining participants, those who did not report
449 sound sensitivity, nor intermittent or chronic tinnitus were classified as neurotypical (N=36, 25 females).
450 The remaining subjects were assigned to the disordered hearing (DH) group (N=35, 17 females) based
451 on their clinical evaluation of chronic tinnitus and/or abnormal sound sensitivity. Among these
452 participants, we found that EEG data (N=2) and skin conductance recordings (N=17) were not usable
453 on account of poor electrode contact. We excluded pupil data (N = 4) and facial movement (N = 2) due
454 to inordinately high rates of blinking or failed tracking.

455

456 **Method details**

457 Participants performed psychophysical and questionnaire assessments remotely, between two in-lab
458 sessions (approximately 3 hours each).

459 *Patient reported outcome measures.* The psychoaffective burden of tinnitus and hyperacusis
460 symptoms was assessed with the Hyperacusis Questionnaire (HQ)[78] and the Tinnitus Handicap
461 Inventory (THI)[79]. Both NT and DH participants can provide meaningful questionnaire responses to the
462 HQ, which focuses on discomfort and aversion experienced with environmental sounds. By contrast,
463 only participants who experience tinnitus can provide meaningful responses to the THI, which focuses
464 on the lifestyle burden associated with phantom auditory percepts.

465 *Pure tone audiometry and uncomfortable loudness level testing.* Prior to EEG recordings, air
466 conduction thresholds were measured for each ear with insert earphones (EarTone-3A) for pure tones
467 ranging from 0.25 – 8 kHz in octave intervals using the modified Hughson-Westlake procedure.
468 Uncomfortable loudness level (ULL) assessment was also performed for each ear just prior to the EEG
469 session as well as during the remote tablet-based testing. Prior to the EEG session, ULL was
470 determined with the Contour Test of Loudness Perception[80], which presented three amplitude
471 modulated 2kHz tones (200ms duration, 5ms raised cosine onset and offset ramps) beginning 5-dB
472 below their 2 kHz hearing threshold and ascended in 5-dB steps until the participant rated the sound as

"Uncomfortably Loud". Participants completed four runs and the median intensity level across the four
runs determined the demarcation of all seven loudness categories. Tablet-based testing was performed
with a calibrated tablet computer and circumaural headphones (Bose AE2). For tablet-based testing,
participants dragged a virtual slider to adjust pure tone sound intensity to the point where sound was
judged to be uncomfortably loud, as described previously[72]. The ULL was the average across three
repetitions.

Electroencephalography (EEG) measurements of central gain. Participants were seated in a
reclining chair within a sound-treated booth and watched a movie of their choice with the volume muted
and subtitles on. Continuous audio and EEG monitoring ensured that participants remained in an
awake, restful state for the duration of testing. EEG recordings were performed with a 64-channel array
of scalp electrodes and insert earphones (EarTone 3A) connected to an electrically isolated digitizer
and signal processor (BioSemi ActiveTwo, Cortech Solutions Inc.). Envelope following responses
(EFRs) were measured in response to auditory stimuli delivered to the left ear via insert headphones
(EarTone-3A). Stimuli were sinusoidal amplitude-modulated tones with a carrier frequency of 2 kHz, a
modulation rate of 40-Hz, and 100% modulation depth. Stimulus delivery (sampling rate: 100 kHz) and
data acquisition (sampling rate: 8192 Hz) were coordinated and aligned through custom LabVIEW
applications. The stimulus was repeated 160 times with alternating polarity of the carrier signal.
Stimulus intensity was continuously ramped up (8 seconds) and down (8 seconds) over a 70 dB range,
corresponding to a rate of intensity change of 8.75 dB/second. The upper limit of the intensity range
was initially set to the median sound rated 6, "Loud, but comfortable" with additional downward
adjustments made upon subject request.

Affective sound evaluations. Participants listened to 60 emotionally evocative environmental sounds
from the original International Affective Digitized Sounds (IADS) corpus and the extended IADS-E
corpus[54,81]. The IADS corpus contains 167 naturally occurring sounds that have been extensively
validated for quantifying differences in affective reactions. For the present study, a subset of sixty IADS
stimuli were chosen to span previously established valence categories[53,81,82]. Stimuli were presented
binaurally through calibrated circumaural headphones (Bose AE2) in random order at a root-mean-
square level of 75 dB SPL. A single trial consisted of a 5-second pre-stimulus baseline period, a 6-
second stimulus, a 5-second silent post-stimulus period, and 30 seconds for the participant to self-
report their behavioral response to the stimulus and for the physiological responses to return to
baseline. Behavioral evaluations asked participants to rate the valence, arousal, and loudness of the
preceding sound with a nine-point Self-Assessment Manikin scale[83]. Participants' heads were stabilized
throughout the session with a padded head support frame with an adjustable chin and forehead rest
(SR Research Ltd.).

Pupil, skin conductance, and facial recordings. Changes in pupil size, skin conductance, and facial
expressions were recorded during each of the 60 trials of sound tokens selected from the IADS corpus.
Participants sat in isoluminant conditions and fixated on a cross in the center of a front-facing monitor.
Sound-evoked changes in pupil size were recorded using the EyeLink 1000 Plus (SR Research Ltd.) at
a sampling rate of 1 kHz. Prior to testing, the participant's pupil size dynamic range was characterized
by presenting alternating bright and dark screens on the computer monitor. Skin conductance
responses (SCRs) were recorded using the skin conductance module of the BioSemi ActiveTwo
system (Cortech Limited Inc.) with two electrodes placed on the hand. Videos of the participant's facial
expressions were recorded at a sampling rate of 120 Hz with a Genie Nano-M2020 camera (Teledyne
DALSA) fitted with a 16mm IP/CCTV lens (TAMRON).

Mulit-talker speech intelligibility task. The details of the task procedure are similar to previous
work[84]. Briefly, participants were required to report four digits that were spoken by a male target

speaker (F_0=115 Hz) masked by two additional speakers (F_0=90 Hz, F_0=175 Hz) who were also
vocalizing digits. Digits (1-9, excluding the bisyllabic 'se-ven') were pseudorandomly selected for each
speaker such that each speaker produced a distinct digit at any given time. Stimuli were presented
diotically through calibrated circumaural headphones (Bose AE2). After familiarization with the task,
participants performed randomly interleaved blocks where four blocks had a signal-to-noise ratio of 9
dB SNR, and four blocks at 0 dB SNR. A block consisted of 10 trials (each a 4-digit sequence), where
the first two trials were adaptive in difficulty, designed to re-familiarize the participant with the target,
and were excluded from analyses. In each trial, digits were spaced with 0.68 s between onsets, and a
virtual keypad appeared 1 s following the fourth digit to allow participants to report the target digits.
Feedback was not provided during testing. Pupil size was simultaneously recorded throughout, as
described above.

Quantification and statistical analysis

Ranked valence rating. Each participant's rank-ordered valence ratings were used to order their
individual autonomic responses (e.g., in **Figure 2E**). As multiple ratings could have the same integer
value, and hence their relative rank-order was arbitrary, all responses for a given integer valence rating
were represented by their mean.

Token spectrograms. Sounds were filtered via a 64 channel gammatone filterbank with center
frequencies spaced between .1 and 10 kHz on an ERB scale. Energy was calculated using 50 ms
windows with 25 ms overlap. Values were converted to dB prior to display. Spectrogram plots omit
sound frequencies and include additional modifications not present in the source material to honor the
terms of use associated with the IADS and IADS-E stimulus batteries.

EEG data analysis. Analysis of EEG data was performed in MATLAB (MathWorks, Natick, MA) using
FieldTrip software[85]. Data were down-sampled and filtered using zero-phase Butterworth bandpass
filters. Eye movement artifacts were identified and removed using independent component analysis[86,87].
The 160 sweeps were averaged together and principal component analysis was used to identify the
optimal subset of EEG channels across which to analyze the EFR response[88–90]. After pre-processing,
amplitude-intensity functions were quantified using spectrograms, wherein multiple short-term Fourier
transforms were computed on consecutive overlapping 1-second intervals using a rectangular
window[63]. The second half of the sweep was time-reversed and vector-averaged with the first half of
the sweep, combining data from the upward and downward sweeps into a single upward function. A
first order polynomial was fit (least-squares) to each growth function between 40 and 65 dB SL. Central
gain was quantified as the slope of this fit.

Pupillometry data analysis. Analysis of pupillometry data was performed in MATLAB (MathWorks,
Natick, MA). To avoid including periods with blinks or missing data, a custom script thresholded
absolute pupil size and pupil derivative, padding flagged periods with 100 samples (**Figure S3B**).
Thresholding was verified by visual inspection. For each participant, z-score normalization was
performed using the mean and standard deviation pooled from traces in the 3 seconds prior to all 60
trials (**Figure S3C**). The covariate of baseline pupil-size was regressed out linearly
(evoked=0.58×baseline+0.39) (**Figure S3D**). Missing data resulting from blink extraction were replaced
through linear interpolation. Trials missing more than 50% of data were excluded from analysis
Otherwise, flagged missing data were linearly interpolated (**Figure S3E**). Evoked pupil responses were
summarized as the mean response between 2 and 5 seconds re. stimulus onset. For the dynamic light
stimulus and multi-talker speech intelligibility task, pupillometry analysis procedures matched those
described for the responses to IADS stimuli. Trial responses in the multi-talker speech intelligibility task

were summarized as the mean pupil diameter between 0.5 and 3.5 second after the onset of the first
digit.

Skin conductance data analysis. SCR data were pre-processed in MATLAB (MathWorks, Natick,
MA) using the Ledalab Version 3.2.2 toolbox[91,92]. A non-negative deconvolution approach was used to
separate the skin conductance data into continuous tonic signals (i.e., slow-varying skin conductance
level) and phasic signals (i.e., fast-varying SCRs)[91]. For each stimulus trial, the integrated SCR (iSCR)
was calculated by taking the time integral of the phasic signal during the eleven seconds following
sound onset. The trial average iSCR was calculated for each participant for each sound token.

Facial videography data analysis (Figure 3B). We identified and mapped 478 3-dimensional facial
landmarks using the Python MediaPipe toolbox at a downsampled rate of 20 Hz[93]. Face-mesh
landmark positions were linearly interpolated across blinks. Landmark positions were temporally
smoothed with a Gaussian kernel (sd of 5 samples). Contrast limited adaptive histogram equalization
was applied to each video frame. Histogram of oriented gradients analyses (resolution of 8 orientations,
local normalization, 2x2 blocks) were performed on each frame, on windows anchored to 77 uniformly
spaced landmark. Windows were 20% the height and width of the fitted face-mesh. Euclidean distance
relative to features in the preceding/proceeding second was derived and smoothed with a Gaussian
kernel (sd of 10 samples). For each participant, Euclidean distances were numerically standardized
(i.e., mode and sd calculated with kernel density estimates) across all 60 sound token responses. Trials
that deviated by more than five times the average deviation were removed. Facial landmarks were
clustered via kernel k-means[94], where kernels were formed for each participant by calculating the
Euclidian distance between facial region responses and applying a Gaussian kernel, before averaging
the matrix across participants, and running k-means with 6 clusters (the maximum number of clusters
that retained facial symmetry). Trial responses were summarized as the mean across all 77 facial
regions between 0 and 6 seconds re. stimulus onset.

Elastic net regression. Elastic net regression[95] was used to investigate the relationship between
predictor variables (pupil, face, valence rating, age, and EFR slope), and severity scores (THI, HQ).
Predictor variables were first standardized such that fitted coefficients approximated relative predictor
strength. Elastic net regression incorporates both regularization and feature selection. The predictive
accuracy of the model is penalized for any non-zero coefficients within the model. To regulate
correlated predictor variables, the coefficient penalty is a weighted sum of the L1 and L2 norms (here,
equally weighted). The value λ scales the size of the coefficient penalty where for larger values of λ any
coefficients that are not predictive of the outcome variable are suppressed (**Figure 4A,D**). Fivefold-
validation, repeated for 500 random initializations, was used to derive λ that minimized out-of-sample
mean square error (MSE). With the predictor variables selected via the elastic net, we refit a linear
regression model on all available data (including individuals not in the original elastic net regression
due to missing values when including other predictor variables) and contrasted against a linear
regression model without the selected autonomic measures (pupil, face) with a likelihood ratio test.

Statistical analysis. Statistical analyses were performed with MATLAB (Mathworks, Natick, MA). Non-
parametric tests were used when assumptions of parametric tests did not apply (i.e., for behavioral
interval data). To assess statistical significance, we used a p-value criterion of $p < 0.05$ (symbolized with
an asterisk, when appropriate $p < 0.01$ was also symbolized with two asterisks). Specific statistical
details can be found in the corresponding figure legends.

References

1. Jarach, C.M., Lugo, A., Scala, M., van den Brandt, P.A., Cederroth, C.R., Odone, A., Garavello, W., Schlee, W., Langguth, B., and Gallus, S. (2022). Global prevalence and incidence of tinnitus: A systematic review and meta-analysis. JAMA Neurol.
2. Paulin, J., Andersson, L., and Nordin, S. (2016). Characteristics of hyperacusis in the general population. Noise Health *18*, 178.
3. Miyakawa, A., Wang, W., Cho, S.-J., Li, D., Yang, S., and Bao, S. (2019). Tinnitus correlates with downregulation of cortical glutamate decarboxylase 65 expression but not auditory cortical map reorganization. J. Neurosci. *39*, 9989–10001.
4. Sedley, W., Parikh, J., Edden, R.A.E., Tait, V., Blamire, A., and Griffiths, T.D. (2015). Human auditory cortex neurochemistry reflects the presence and severity of tinnitus. J. Neurosci. *35*, 14822–14828.
5. Yang, S., Weiner, B.D., Zhang, L.S., Cho, S.-J.J., and Bao, S. (2011). Homeostatic plasticity drives tinnitus perception in an animal model. Proc. Natl. Acad. Sci. *108*, 14974–14979. 10.1073/pnas.1107998108.
6. Eggermont, J.J., and Roberts, L.E. (2004). The neuroscience of tinnitus. Trends Neurosci. *27*, 676–682.
7. Roberts, L.E., and Salvi, R. (2019). Overview: Hearing loss, tinnitus, hyperacusis, and the role of central gain at Elsevier.
8. Harris, K.C., Dias, J.W., McClaskey, C.M., Rumschlag, J., Prisciandaro, J., and Dubno, J.R. (2022). Afferent loss, GABA, and Central Gain in older adults: Associations with speech recognition in noise. J. Neurosci. *42*, 7201–7212.
9. Cisneros-Franco, J.M., Ouellet, L., Kamal, B., and de Villers-Sidani, E. (2018). A brain without brakes: reduced inhibition is associated with enhanced but dysregulated plasticity in the aged rat auditory cortex. Eneuro *5*.
10. Resnik, J., and Polley, D.B. (2017). Fast-spiking GABA circuit dynamics in the auditory cortex predict recovery of sensory processing following peripheral nerve damage. Elife *6*, e21452.
11. Resnik, J., and Polley, D.B. (2021). Cochlear neural degeneration disrupts hearing in background noise by increasing auditory cortex internal noise. Neuron *109*, 984–996.
12. Vasilkov, V., Caswell-Midwinter, B., Zhao, Y., de Gruttola, V., Jung, D.H., Liberman, M.C., and Maison, S.F. (2023). Evidence of cochlear neural degeneration in normal-hearing subjects with tinnitus. Sci. Rep. *13*, 19870.
13. Kotak, V.C., Takesian, A.E., MacKenzie, P.C., and Sanes, D.H. (2013). Rescue of inhibitory synapse strength following developmental hearing loss. PLoS One *8*, e53438.
14. Kumar, M., Handy, G., Kouvaros, S., Zhao, Y., Brinson, L.L., Wei, E., Bizup, B., Doiron, B., and Tzounopoulos, T. (2023). Cell-type-specific plasticity of inhibitory interneurons in the rehabilitation of auditory cortex after peripheral damage. Nat. Commun. *14*, 4170.
15. Gu, J.W., Herrmann, B.S., Levine, R.A., and Melcher, J.R. (2012). Brainstem auditory evoked potentials suggest a role for the ventral cochlear nucleus in tinnitus. J. Assoc. Res. Otolaryngol. *13*, 819–833.
16. Bramhall, N.F., Konrad-Martin, D., and McMillan, G.P. (2018). Tinnitus and auditory perception after a history of noise exposure: Relationship to auditory brainstem response measures. Ear Hear. *39*, 881.
17. Balaram, P., Hackett, T.A., and Polley, D.B. (2019). Synergistic transcriptional changes in AMPA and GABAA receptor genes support compensatory plasticity following unilateral hearing loss. Neuroscience *407*, 108–119.
18. Schaette, R., and McAlpine, D. (2011). Tinnitus with a normal audiogram: physiological evidence for hidden hearing loss and computational model. J. Neurosci. *31*, 13452–13457.
19. Schaette, R., Turtle, C., and Munro, K.J. (2012). Reversible induction of phantom auditory sensations through simulated unilateral hearing loss. PLoS One *7*, e35238. 10.1371/journal.pone.0035238.
20. Kurioka, T., Mizutari, K., Satoh, Y., Kobayashi, Y., and Shiotani, A. (2023). Blast-induced central auditory neurodegeneration affects tinnitus development regardless of peripheral cochlear

damage. J. Neurotrauma.

21.   Harris, J.L., Yeh, H.-W., Choi, I.-Y., Lee, P., Berman, N.E., Swerdlow, R.H., Craciunas, S.C., and
Brooks, W.M. (2012). Altered neurochemical profile after traumatic brain injury: 1H-MRS
biomarkers of pathological mechanisms. J. Cereb. Blood Flow Metab. *32*, 2122–2134.

22.   Benedetti, B., Weidenhammer, A., Reisinger, M., and Couillard-Despres, S. (2022). Spinal cord
injury and loss of cortical inhibition. Int. J. Mol. Sci. *23*, 5622.

23.   Laskey, C., and Opitz, B. (2020). Tinnitus associated with benzodiazepine withdrawal syndrome:
A case report and literature review. Ment. Heal. Clin. *10*, 100–103.

24.   Busto, U., Sellers, E.M., Naranjo, C.A., Cappell, H., Sanchez-Craig, M., and Sykora, K. (1986).
Withdrawal reaction after long-term therapeutic use of benzodiazepines. N. Engl. J. Med. *315*,
854–859.

25.   Herrmann, B., and Butler, B.E. (2021). Hearing loss and brain plasticity: the hyperactivity
phenomenon. Brain Struct. Funct. *226*, 2019–2039.

26.   Noreña, A.J. (2011). An integrative model of tinnitus based on a central gain controlling neural
sensitivity. Neurosci. Biobehav. Rev. *35*, 1089–1109.

27.   McGill, M., Hight, A.E., Watanabe, Y.L., Parthasarathy, A., Cai, D., Clayton, K., Hancock, K.E.,
Takesian, A., Kujawa, S.G., and Polley, D.B. (2022). Neural signatures of auditory
hypersensitivity following acoustic trauma. Elife *11*, e80015.

28.   Zeng, F.-G. (2020). Tinnitus and hyperacusis: central noise, gain and variance. Curr. Opin.
Physiol. *18*, 123–129.

29.   Koops, E.A., and van Dijk, P. (2021). Hyperacusis in tinnitus patients relates to enlarged
subcortical and cortical responses to sound except at the tinnitus frequency. Hear. Res. *401*,
108158.

30.   Witkin, J.M., Lippa, A., Smith, J.L., Cook, J.M., and others (2022). Can GABAkines quiet the
noise? The GABAA receptor neurobiology and pharmacology of tinnitus. Biochem. Pharmacol.
201, 115067.

31.   Masri, S., Chan, N., Marsh, T., Zinsmaier, A., Schaub, D., Zhang, L., Wang, W., and Bao, S.
(2021). Chemogenetic activation of cortical parvalbumin-positive interneurons reverses noise-
induced impairments in gap detection. J. Neurosci. *41*, 8848–8857.

32.   De Ridder, D., Schlee, W., Vanneste, S., Londero, A., Weisz, N., Kleinjung, T., Shekhawat, G.S.,
Elgoyhen, A.B., Song, J.-J., and Andersson, G. (2021). Tinnitus and tinnitus disorder: Theoretical
and operational definitions (an international multidisciplinary proposal). Prog. Brain Res. *260*, 1–
25.

33.   Elgoyhen, A.B., Langguth, B., De Ridder, D., and Vanneste, S. (2015). Tinnitus: perspectives
from human neuroimaging. Nat. Rev. Neurosci. *16*, 632–642.

34.   Asokan, M.M., Watanabe, Y., Kimchi, E.Y., and Polley, D.B. (2023). Potentiation of cholinergic
and corticofugal inputs to the lateral amygdala in threat learning. Cell Rep. *42*.

35.   Leaver, A.M., Seydell-Greenwald, A., and Rauschecker, J.P. (2016). Auditory–limbic interactions
in chronic tinnitus: Challenges for neuroimaging research. Hear. Res. *334*, 49–57.

36.   Chen, Y.-C., Li, X., Liu, L., Wang, J., Lu, C.-Q., Yang, M., Jiao, Y., Zang, F.-C., Radziwon, K.,
and Chen, G.-D. (2015). Tinnitus and hyperacusis involve hyperactivity and enhanced
connectivity in auditory-limbic-arousal-cerebellar network. Elife *4*, e06576.

37.   Husain, F.T., and Khan, R.A. (2023). Review and Perspective on Brain Bases of Tinnitus. J.
Assoc. Res. Otolaryngol., 1–14.

38.   Knudson, I.M., and Melcher, J.R. (2016). Elevated acoustic startle responses in humans:
relationship to reduced loudness discomfort level, but not self-report of hyperacusis. J. Assoc.
Res. Otolaryngol. *17*, 223–235.

39.   Smalt, C.J., Sugai, J.A., Koops, E.A., Jahn, K.N., Hancock, K.E., and Polley, D.B. (2022).
Automatic identification of tinnitus malingering based on overt and covert behavioral responses
during psychoacoustic testing. NPJ Digit. Med. *5*, 127.

40.   Liu, Z., Yao, L., Wang, X., Monaghan, J.J.M., Schaette, R., He, Z., and McAlpine, D. (2021).
Generalizable sample-efficient siamese autoencoder for tinnitus diagnosis in listeners with
subjective tinnitus. IEEE Trans. Neural Syst. Rehabil. Eng. *29*, 1452–1461.

41.   Yakunina, N., and Nam, E.-C. (2021). What makes tinnitus loud? Otol. Neurotol. *42*, 235–241.

42. Gu, J.W., Halpin, C.F., Nam, E.-C., Levine, R.A., and Melcher, J.R. (2010). Tinnitus, diminished
sound-level tolerance, and elevated auditory activity in humans with clinically normal hearing
sensitivity. J. Neurophysiol. *104*, 3361–3370.

43. Munro, K.J., Turtle, C., and Schaette, R. (2014). Plasticity and modified loudness following short-
term unilateral deprivation: evidence of multiple gain mechanisms within the auditory system. J.
Acoust. Soc. Am. *135*, 315–322.

44. Vanneste, S., Plazier, M., Van Der Loo, E., Van de Heyning, P., Congedo, M., and De Ridder, D.
(2010). The neural correlates of tinnitus-related distress. Neuroimage *52*, 470–480.

45. Kreibig, S.D. (2010). Autonomic nervous system activity in emotion: A review. Biol. Psychol. *84*,
394–421.

46. Damasio, A., and Carvalho, G.B. (2013). The nature of feelings: evolutionary and neurobiological
origins. Nat. Rev. Neurosci. *14*, 143–152.

47. Critchley, H.D., Rotshtein, P., Nagai, Y., O'Doherty, J., Mathias, C.J., and Dolan, R.J. (2005).
Activity in the human brain predicting differential heart rate responses to emotional facial
expressions. Neuroimage *24*, 751–762.

48. Quadt, L., Critchley, H., and Nagai, Y. (2022). Cognition, emotion, and the central autonomic
network. Auton. Neurosci. *238*, 102948.

49. Dolensek, N.N., Gehrlach, D.A., Klein, A.S., and Gogolla, N. (2020). Facial expressions of
emotion states and their neuronal correlates in mice. Science (80-.). *368*, 89–94.
10.1126/SCIENCE.AAZ9468/SUPPL_FILE/AAZ9468S3.MP4.

50. Reimer, J., McGinley, M.J., Liu, Y., Rodenkirch, C., Wang, Q., McCormick, D.A., and Tolias, A.S.
(2016). Pupil fluctuations track rapid changes in adrenergic and cholinergic activity in cortex. Nat.
Commun. *7*, 13289.

51. Joshi, S., and Gold, J.I. (2020). Pupil size as a window on neural substrates of cognition. Trends
Cogn. Sci. *24*, 466–480. 10.1016/J.TICS.2020.03.005.

52. Koelewijn, T., de Kluiver, H., Shinn-Cunningham, B.G., Zekveld, A.A., and Kramer, S.E. (2015).
The pupil response reveals increased listening effort when it is difficult to focus attention. Hear.
Res. *323*, 81–90.

53. Partala, T., and Surakka, V. (2003). Pupil size variation as an indication of affective processing.
Int. J. Hum. Comput. Stud. *59*, 185–198.

54. Yang, W., Makita, K., Nakao, T., Kanayama, N., Machizawa, M.G., Sasaoka, T., Sugata, A.,
Kobayashi, R., Hiramoto, R., and Yamawaki, S. (2018). Affective auditory stimulus database: An
expanded version of the International Affective Digitized Sounds (IADS-E). Behav. Res. Methods
50, 1415–1429.

55. Enzler, F., Fournier, P., and Norena, A.J. (2021). A psychoacoustic test for diagnosing
hyperacusis based on ratings of natural sounds. Hear. Res. *400*, 108124.

56. Zeng, F.-G., Richardson, M., and Turner, K. (2020). Tinnitus does not interfere with auditory and
speech perception. J. Neurosci. *40*, 6007–6017.

57. Ekman, P. (1993). Facial expression and emotion. Am. Psychol. *48*, 384.

58. Jackson, R., Vijendren, A., and Phillips, J. (2019). Objective measures of tinnitus: a systematic
review. Otol. \& Neurotol. *40*, 154–163.

59. Haider, H.F., Hoare, D.J., Ribeiro, S.F., Ribeiro, D., Caria, H., Trigueiros, N., Borrego, L.M.,
Szczepek, A.J., Papoila, A.L., Elarbed, A., et al. (2021). Evidence for biological markers of
tinnitus: A systematic review. Prog. Brain Res. *262*, 345–398.

60. Ryle, G. (1949). The Concept of Mind (London: Hutchinson's University Library). Cited on *25*.

61. Melcher, J.R., Sigalovsky, I.S., Guinan Jr, J.J., and Levine, R.A. (2000). Lateralized tinnitus
studied with functional magnetic resonance imaging: abnormal inferior colliculus activation. J.
Neurophysiol. *83*, 1058–1072.

62. Bramhall, N.F., McMillan, G.P., Gallun, F.J., and Konrad-Martin, D. (2019). Auditory brainstem
response demonstrates that reduced peripheral auditory input is associated with self-report of
tinnitus. J. Acoust. Soc. Am. *146*, 3849–3862.

63. Picton, T.W., Van Roon, P., and John, M.S. (2007). Human auditory steady-state responses
during sweeps of intensity. Ear Hear. *28*, 542–557. 10.1097/AUD.0B013E31806DC2A7.

64. Sedley, W. (2019). Tinnitus: does gain explain? Neuroscience *407*, 213–228.

65. Chen, X., Sun, Y.-C., Zhan, H., Kebschull, J.M., Fischer, S., Matho, K., Huang, Z.J., Gillis, J.,
and Zador, A.M. (2019). High-throughput mapping of long-range neuronal projection using in situ
sequencing. Cell *179*, 772–786.
66. Romanski, L.M., and LeDoux, J.E. (1993). Information cascade from primary auditory cortex to
the amygdala: corticocortical and corticoamygdaloid projections of temporal cortex in the rat.
Cereb. Cortex *3*, 515–532.
67. Romanski, L.M., Bates, J.F., and Goldman-Rakic, P.S. (1999). Auditory belt and parabelt
projections to the prefrontal cortex in the rhesus monkey. J. Comp. Neurol. *403*, 141–157.
68. Leaver, A.M., Renier, L., Chevillet, M.A., Morgan, S., Kim, H.J., and Rauschecker, J.P. (2011).
Dysregulation of limbic and auditory networks in tinnitus. Neuron *69*, 33–43.
69. Sedley, W., Friston, K.J., Gander, P.E., Kumar, S., and Griffiths, T.D. (2016). An integrative
tinnitus model based on sensory precision. Trends Neurosci. *39*, 799–812.
10.1016/j.tins.2016.10.004.
70. Krauss, P., Tziridis, K., Metzner, C., Schilling, A., Hoppe, U., and Schulze, H. (2016). Stochastic
resonance controlled upregulation of internal noise after hearing loss as a putative cause of
tinnitus-related neuronal hyperactivity. Front. Neurosci. *10*, 597.
71. Schall, J.D., and Thompson, K.G. (1999). Neural selection and control of visually guided eye
movements. Annu. Rev. Neurosci. *22*, 241–259.
72. Chen, J.X., Whitton, J.P., Parthasarathy, A., Hancock, K.E., and Polley, D.B. (2020). Fluctuations
in subjective tinnitus ratings over time: Implications for clinical research. Otol. Neurotol. Off. Publ.
Am. Otol. Soc. Am. Neurotol. Soc. [and] Eur. Acad. Otol. Neurotol. *41*, e1167.
73. Aazh, H., and Moore, B.C.J. (2017). Factors associated with depression in patients with tinnitus
and hyperacusis. Am. J. Audiol. *26*, 562–569.
74. Smith, S.S., Kitterick, P.T., Scutt, P., Baguley, D.M., and Pierzycki, R.H. (2021). An exploration
of psychological symptom-based phenotyping of adult cochlear implant users with and without
tinnitus using a machine learning approach. Prog. Brain Res. *260*, 283–300.
75. Tavassoli, T., Miller, L.J., Schoen, S.A., Brout, J.J., Sullivan, J., and Baron-Cohen, S. (2018).
Sensory reactivity, empathizing and systemizing in autism spectrum conditions and sensory
processing disorder. Dev. Cogn. Neurosci. *29*, 72–77.
76. Williams, Z.J., He, J.L., Cascio, C.J., and Woynaroski, T.G. (2021). A review of decreased sound
tolerance in autism: Definitions, phenomenology, and potential mechanisms. Neurosci.
Biobehav. Rev. *121*, 1–17.
77. Polley, D.B., and Schiller, D. (2022). The promise of low-tech intervention in a high-tech era:
Remodeling pathological brain circuits using behavioral reverse engineering. Neurosci.
Biobehav. Rev. *137*, 104652.
78. Khalfa, S., Dubal, S., Veuillet, E., Perez-Diaz, F., Jouvent, R., and Collet, L. (2002).
Psychometric normalization of a hyperacusis questionnaire. Orl *64*, 436–442.
79. Newman, C.W., Jacobson, G.P., and Spitzer, J.B. (1996). Development of the tinnitus handicap
inventory. Arch. Otolaryngol. Neck Surg. *122*, 143–148.
80. Cox, R.M., Alexander, G.C., Taylor, I.M., and Gray, G.A. (1997). The contour test of loudness
perception. Ear Hear. *18*, 388–400.
81. Bradley, M.M., and Lang, P.J. (2007). The International Affective Digitized Sounds (; IADS-2):
Affective ratings of sounds and instruction manual (Technical report B-3. University of Florida,
Gainesville, Fl).
82. Szibor, A., Lehtimäki, J., Ylikoski, J., Aarnisalo, A.A., Mäkitie, A., and Hyvärinen, P. (2018).
Attenuation of positive valence in ratings of affective sounds by tinnitus patients. Trends Hear.
22, 2331216518816215.
83. Bradley, M.M., and Lang, P.J. (1994). Measuring emotion: the self-assessment manikin and the
semantic differential. J. Behav. Ther. Exp. Psychiatry *25*, 49–59.
84. Parthasarathy, A., Hancock, K.E., Bennett, K., DeGruttola, V., and Polley, D.B. (2020). Bottom-
up and top-down neural signatures of disordered multi-talker speech perception in adults with
normal hearing. Elife *9*, e51419.
85. Oostenveld, R., Fries, P., Maris, E., and Schoffelen, J.-M. (2011). FieldTrip: open source
software for advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput.

Intell. Neurosci. *2011*, 1–9.

86. Jung, T.-P., Makeig, S., Humphries, C., Lee, T.-W., Mckeown, M.J., Iragui, V., and Sejnowski,
T.J. (2000). Removing electroencephalographic artifacts by blind source separation.
Psychophysiology *37*, 163–178.

87. Vigário, R.N. (1997). Extraction of ocular artefacts from EEG using independent component
analysis. Electroencephalogr. Clin. Neurophysiol. *103*, 395–404.

88. Bharadwaj, H.M., Masud, S., Mehraei, G., Verhulst, S., and Shinn-Cunningham, B.G. (2015).
Individual differences reveal correlates of hidden hearing deficits. J. Neurosci. *35*, 2161–2172.

89. Bharadwaj, H.M., and Shinn-Cunningham, B.G. (2014). Rapid acquisition of auditory subcortical
steady state responses using multichannel recordings. Clin. Neurophysiol. *125*, 1878–1888.

90. Lu, H., Mehta, A.H., Bharadwaj, H.M., Shinn-Cunningham, B.G., and Oxenham, A.J. (2020).
Comment on 'Rapid acquisition of auditory subcortical steady state responses using
multichannel recordings.' Clin. Neurophysiol. Off. J. Int. Fed. Clin. Neurophysiol. *131*, 1833.

91. Benedek, M., and Kaernbach, C. (2010). Decomposition of skin conductance data by means of
nonnegative deconvolution. Psychophysiology *47*, 647–658. 10.1111/J.1469-
8986.2009.00972.X.

92. Benedek, M., and Kaernbach, C. (2010). A continuous measure of phasic electrodermal activity.
J. Neurosci. Methods *190*, 80–91. 10.1016/J.JNEUMETH.2010.04.028.

93. Lugaresi, C., Tang, J., Nash, H., McClanahan, C., Uboweja, E., Hays, M., Zhang, F., Chang, C.-
L., Yong, M.G., and Lee, J. (2019). Mediapipe: A framework for building perception pipelines.
arXiv Prepr. arXiv1906.08172.

94. Zhou, F., Torre, F.D. La, and Hodgins, J.K. (2013). Hierarchical aligned cluster analysis for
temporal clustering of human motion. IEEE Trans. Pattern Anal. Mach. Intell. *35*, 582–596.
10.1109/TPAMI.2012.137.

95. Zou, H., and Hastie, T. (2005). Regularization and variable selection via the elastic net. J. R.
Stat. Soc. Ser. B Stat. Methodol. *67*, 301–320.

848