Reliability of task-based fMRI in the dorsal horn of the human spinal cord

Alice Dabbagh[1], Ulrike Horn[1], Merve Kaptan[1,2], Toralf Mildner[3], Roland Müller[3], Jöran Lepsien[3], Nikolaus Weiskopf[4,5,6], Jonathan C.W. Brooks[7], Jürgen Finsterbusch[8], Falk Eippert[1]

Max Planck Research Group Pain Perception, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

Department of Anesthesiology, Perioperative and Pain Medicine, Stanford University, CA, USA

Methods & Development Group Nuclear Magnetic Resonance, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

Department of Neurophysics, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

Felix Bloch Institute for Solid State Physics, Faculty of Physics and Earth Sciences, University of Leipzig, Leipzig, Germany

Wellcome Centre for Human Neuroimaging, Institute of Neurology, University College London, London, UK

School of Psychology, University of East Anglia Wellcome Wolfson Brain Imaging Centre (UWWBIC), Norwich, United Kingdom

Department of Systems Neuroscience, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

Address for correspondence: Alice Dabbagh & Falk Eippert; Max Planck Research Group Pain Perception, Max Planck Institute for Human Cognitive and Brain Sciences, Stephanstraße 1a, 04103 Leipzig, Germany.

Number of pages: 48

Number of figures / tables: 7 / 1

Number of supplementary pages: 7

Number of supplementary figures / tables: 3 / 4

Abstract: The application of functional magnetic resonance imaging (fMRI) to the human spinal cord is still a relatively small field of research and faces many challenges. Here we aimed to probe the limitations of task-based spinal fMRI at 3T by investigating the reliability of spinal cord blood oxygen level dependent (BOLD) responses to repeated nociceptive stimulation across two consecutive days in 40 healthy volunteers. We assessed the test-retest reliability of subjective ratings, autonomic responses, and spinal cord BOLD responses to short heat pain stimuli (1s duration) using the intraclass correlation coefficient (ICC). At the group level, we observed robust autonomic responses as well as spatially specific spinal cord BOLD responses at the expected location, but no spatial overlap in BOLD response patterns across days. While autonomic indicators of pain processing showed good-to-excellent reliability, both β-estimates and z-scores of task-related BOLD responses showed poor reliability across days in the target region (gray matter of the ipsilateral dorsal horn). When taking into account the sensitivity of gradient-echo echo planar imaging (GE-EPI) to draining vein signals by including the venous plexus in the analysis, we observed BOLD responses with fair reliability across days. Taken together, these results demonstrate that heat pain stimuli as short as one second are able to evoke a robust and spatially specific BOLD response, which is however strongly variable within participants across time, resulting in low reliability in the dorsal horn gray matter. Further improvements in data acquisition and analysis techniques are thus necessary before event-related spinal cord fMRI as used here can be reliably employed in longitudinal designs or clinical settings.

Keywords: spinal cord; fMRI; heat pain; reliability; spatial specificity; human

1. Introduction

Functional magnetic resonance imaging (fMRI) is a non-invasive method routinely used for brain imaging, with its first application in the human spinal cord about 30 years ago (Yoshizawa et al., 1996). Compared to the brain, the spinal cord is a more challenging environment for fMRI (Bosma & Stroman, 2014; Cohen-Adad, 2017; Eippert, Kong, Jenkinson, et al., 2017; Giove et al., 2004; Kinany, Pirondini, Micera, et al., 2022) and the number of studies using this technique has increased only slowly at first. However, the continued development and improvement of scanner hardware (Cohen-Adad et al., 2011; Lopez-Rios et al., 2023; Topfer et al., 2016), image acquisition protocols for spinal cord fMRI (Barry et al., 2021; Finsterbusch et al., 2013; Kinany,et al., 2022), shimming procedures (Finsterbusch et al., 2012; Islam et al., 2019; Tsivaka et al., 2023), denoising strategies (Brooks et al., 2008; Kong et al., 2012; Vannesjo et al., 2019) and software tools tailored to preprocessing and analyzing spinal cord data (De Leener et al., 2017, 2018; Rangaprakash & Barry, 2022) have made spinal fMRI more robust, sensitive and accessible, and accordingly has met with growing numbers of spinal fMRI studies more recently (Kinany et al., 2022; Landelle et al., 2021; Powers et al., 2018; Tinnermann et al., 2021).

Apart from a few notable exceptions (Conrad et al., 2018; Martucci et al., 2019, 2021; Rowald et al., 2022; Stroman, 2002; Stroman et al., 2004), most spinal cord fMRI studies have focused on cross-sectional designs in healthy volunteers, i.e. have not employed longitudinal designs or looked at the diagnostic or prognostic potential of spinal fMRI in clinical settings. This is different to brain imaging, the use of which in longitudinal settings and for biomarker development in the clinical context has been extensively discussed (Cole & Franke, 2017; Elliott et al., 2020; Khalili-Mahani et al., 2017; Kragel et al., 2021; Woo & Wager, 2016). Considering that the spinal cord is affected in a large range of neurological conditions – such as multiple sclerosis (Filippi & Rocca, 2013), neuropathic pain (Colloca et al., 2017) or spinal cord injury (Ahuja et al., 2017) – spinal cord fMRI could also be a valuable tool in clinical contexts, e.g., for tracking or predicting disease progression and treatment. However, the successful application of spinal cord fMRI in longitudinal settings and for diagnostic or prognostic purposes requires – at a minimum – achieving a high reliability of the method, with reliability being the extent to which measurement outcomes are consistent across different contexts. Test-retest reliability, for instance, describes the stability of a measure over time, i.e. it quantifies the precision of a method, or in other words, the expected variation over time, given that the underlying process of interest remains the same (Brandmaier et al., 2018; Lavrakas, 2008; Noble et al., 2021).

Studies assessing the test-retest reliability of spinal cord fMRI have mostly focused on resting-
state signals (Barry et al., 2016; Hu et al., 2018; Kaptan et al., 2023; Kong et al., 2014; Kowalczyk
et al., 2023; Liu et al., 2016; San Emeterio Nateras et al., 2016). Only three studies have examined
task-related signal changes, with two of those using motor tasks (Bouwman et al., 2008; Weber
et al., 2016b) and one using a pain task (Weber et al., 2016a). While these task-based studies
provided important initial insights into the reliability of spinal cord fMRI, they had modest sample
sizes (with at most N = 12) and mostly assessed reliability within a single scan session, thus
circumventing some of the challenges inherent to longitudinal studies, such as repositioning of
participants, and day-to-day variations in physiological state and mood (note that Bouwman and
colleagues looked at a time-interval of 10 weeks, but only acquired data from three participants).

Here, we set out to provide a comprehensive assessment of the reliability of task-based spinal
fMRI by investigating heat-pain evoked spinal cord BOLD responses. We chose the domain of
pain for this endeavor for two reasons: on the one hand, changes in spinal cord processing are
assumed to contribute to chronic pain (D'Mello & Dickenson, 2008; Kuner & Flor, 2017; Prescott
et al., 2014) and on the other hand the development of pain biomarkers is currently a topic of
intense interest (Davis et al., 2020; Leone et al., 2022; Mouraux & Iannetti, 2018; Sluka et al.,
2023; Tracey, 2021), making spinal fMRI a prime candidate for inclusion in such clinical
developments. In contrast to previous studies, we acquired data on two consecutive days using
an identical experimental set up and a relatively large sample of 40 participants, as specified in
an accompanying preregistration. We first analyzed the spatial distribution of the response across
multiple spinal segments as well as its spread into the venous plexus surrounding the spinal cord.
We then quantified the spatial consistency of the response patterns (using the Dice coefficient)
and assessed test-retest reliability of BOLD responses in multiple ways (using the intraclass-
correlation coefficient). Importantly, we simultaneously collected peripheral physiological data and
compared their reliability to that of the BOLD data as only this allows for disambiguating between
different causes for possibly low reliability in spinal cord BOLD responses, i.e. either poor data
quality of spinal cord fMRI or variability in the underlying process (nociceptive processing).

2. Methods

2.1 Participants

40 healthy participants (20 female, mean age: 27.3 years, range: 18 – 35 years) participated in this study. This sample size was based on a preregistered power calculation (G*Power, Faul et al., 2007, version 3.1.9.7.), where we estimated that a sample of 36 participants would be necessary to detect a medium-sized effect (d = 0.5) with 90% power at an alpha-level of 0.05 when using a one-sample t-test against the baseline. All participants had normal or corrected-to-normal vision and a BMI < 24, were right-handed and gave written informed consent. The study was approved by the Ethics Committee of the Medical Faculty of the University of Leipzig.

2.2 Thermal stimulation

We employed phasic painful heat stimuli (duration: 1s, including 0.23s of ramp-up and ramp-down each, temperature: 48°C, baseline: 32°C), which were applied to the inner left forearm of the participants via an MRI compatible thermode with a ramp-speed of 70°C/s (surface of 9cm^2, PATHWAY CHEPS; Medoc, Ramat Yishai, Israel). The stimuli were applied on five different areas of the inner left forearm (Figure 1), in order to minimize possible sensitization and habituation over runs.

2.3 Experimental procedure

This study was part of a larger methodological project aimed at investigating the relationship between spinal cord BOLD responses and employed echo times. While we describe the entire data acquisition and experiment for the sake of transparency, here we solely focus on the data relevant for the issue of reliability – the echo-time dependence is the focus of an upcoming report.

At the beginning of the experiment, the participants were informed about the study and any remaining questions were discussed. We then outlined the five stimulation areas on the arm (most likely corresponding to dermatome C6 see Lee et al., 2008), using ink that remained visible over both days of the experiment. Before the main experiment started, the participants were familiarized with the heat stimulus by administering it twice to the right forearm, and then twice to each of the 5 possible stimulation areas on the left forearm. This served to minimize orienting / novelty responses, which could lead to detrimental movement at the beginning of a run.

After this familiarization, we prepared the participants for the placement in the scanner. We attached a breathing belt to measure respiration, as well as three electrodes to record the

electrocardiogram (ECG; one electrode was placed on the left parasternal line at the level of the
1^{st} / 2^{nd} rib, another electrode on the left medioclavicular line at the level of the 9^{th} or 10^{th} rib, and
the ground electrode on the left side of the chest, one hand-width below the armpit). Two
electrodes were placed on the right hand to record skin conductance responses (SCR, one
electrode on the thenar eminence, one electrode on the hypothenar eminence). The thermode
was placed on the left arm. A custom-built MR-compatible extension mechanism (Mueller et al.,
2024) was attached to the thermode that allowed for an easy repositioning of the thermode
between scans from outside the scanner bore, without moving the participants and without
changing the thermode pressure on the skin. After lying down on the scanner table, the
participants were asked to tilt their head slightly towards their chest in order to minimize cervical
lordosis (Cohen-Adad et al., 2021) and the isocenter was set approximately to the participants'
larynx. Before the experiment started, the eye tracker was calibrated to measure the pupil
diameter throughout the experiment (eye tracker settings were validated or, if necessary, re-
calibrated before the beginning of each run). The participants were instructed to avoid moving,
avoid excessive swallowing, and to breathe normally (see Cohen-Adad et al., 2021) as well as to
look at a fixation cross on a screen for the entire duration of each run while avoiding excessive
blinking.

We split the experiment up across two consecutive days and measured five runs in each MRI
session (Day 1 and Day 2). One run consisted of 20 trials, with one trial lasting between 11 and
13 seconds (1s stimulations and jittered inter-trial interval of 10-12s) and the duration of one run
being four minutes and 48 seconds. We measured each run with a different echo time (TE; 7 runs:
TE = 25ms, 30ms, 35ms, 40ms, 45ms, 50ms, 55ms) and a repetition time (TR) of 1800ms; two
runs were additionally measured with the shortest possible TR (TE = 40ms & TR = 1560ms; TE =
25ms & TR = 1320ms). Splitting up the experiment across two days gave us the opportunity to
assess the reliability of task-based spinal fMRI data. For this reason, we measured one run with
a TE of 40ms (approximating T_2^* in the spinal cord, Barry & Smith, 2019) and a TR of 1800ms on
each of the two measurement days (Fig. 1), which is the main focus of this study (from now
referred to as the Reliability Run). In total, we therefore measured ten runs per participant, (nine
TE&TR combinations and one additional Reliability Run), with five runs per day. Run order and
targeted skin patch were pseudo-randomized and counter-balanced across participants, but kept
identical across both days for the Reliability Run (see Fig. 1).

Figure 1. Experimental design. *Top*: 40 participants were measured on two consecutive days (sessions). On each
day, we acquired five runs (each rectangle represents one run, also indicating the employed echo time [TE]). The runs
were randomized across the two measurements days, with the exception that the Reliability Run (TE of 40ms, yellow
rectangle) was measured on both days and is the focus of this report. Runs with a TR of 1800ms (medium grey
rectangles) were combined with the Reliability Run for the Combined Runs average. Runs with a short TR (dark grey
rectangles) were not included in this study. *Bottom*: Left: Before the measurements, we divided the area of the forearm
into 5 equally sized patches, adapted to the individual proportions of the participant's forearm. We drew the stimulation
areas onto the left arm with a pen, to be able to target them easily when changing the thermode position in the scanner
and to stimulate the same areas on the second day of the experiment. During each run, the participants received 20
heat stimuli with a duration of 1s and a temperature of 48°C. Across both days, we targeted the same skin patch for the
Reliability Run. Middle: We measured heart rate, respiration, skin conductance and pupil dilation for each run and at
the end of each run, participants were asked to rate the overall stimulus intensity using a numerical rating scale from 0-
100 (0: no percept, 50: pain threshold, 100: unbearable pain). Right: Sagittal view of an example participant on the left,
with the yellow rectangle indicating the rostro-caudal extent of the EPI slice stack (covering spinal cord segments C5 to
C8, which translates to vertebrae C4 to C7). An axial example EPI slice (top right) demonstrates the data quality
obtained with an in-plane resolution of 1x1mm (slice thickness: 5mm) and the masks employed are shown in the lower
right, with the region of interest being the left dorsal horn in spinal cord segment C6 (depicted in green).

2.4 Data acquisition

The MRI data were acquired on a Siemens PRISMA FIT 3 Tesla scanner (Siemens, Erlangen, Germany), equipped with a whole-body radio-frequency (RF) transmit coil. We used a 64-channel RF head-and-neck coil, of which head coil element 7 and neck coil element groups 1 and 2 were utilized (all receive-only). We started the data acquisition with a localizer scan, followed by positioning the EPI slice stack and adjust volume (60 x 60 x 100 mm). A single EPI volume was then acquired, initializing the scanner's "Advanced shim mode", with the resulting shim applied to all following EPI acquisitions. The angle as well as centering of the adjust volume was identical to that of the EPI acquisitions, but it was slightly larger in superior-inferior-direction. We then acquired a z-shim reference scan that allowed for the automatic determination of the optimal z-shim moment for each slice of the EPI slice stack (Finsterbusch et al., 2012; Kaptan et al., 2022). A sagittal field map was obtained to estimate the static B0 field distribution. After this we measured a high-resolution T2-weighted structural scan for registration purposes, followed by two T2*-weighted ME-GRE scans to map T2* with two different resolutions. Finally, we measured the five functional runs. Prior to each functional run we acquired ten functional volumes with posterior-to-anterior phase encoding. In the following paragraph, we provide details on all protocols, except for the acquired field map, T2*-weighted ME-GRE and functional volumes measured with posterior-to-anterior phase encoding, since these were not utilized in the course of this study and will be described elsewhere (as will the EPI runs with shortened TR).

The EPI z-shim reference scan (TE: 40ms, total acquisition time: 55 seconds) consisted of 21 volumes with equidistant z-shim moments compensating for field inhomogeneities between +0.21 and -0.21 mT/m (in steps of 0.021 mT/m). The fMRI runs were measured via a single-shot 2D gradient-echo EPI sequence with 16 slices, covering the spinal cord from the 4th cervical vertebra to the 1st thoracic vertebra, with a resolution of 1 x 1 x 5mm (slice orientation: oblique axial; TR: 1.8s, TE different between runs: 25ms | 30m | 35ms | 40ms | 45ms | 50ms | 55ms, readout flip angle (FA): 75°, field of view (FOV): 128 x 128mm^2, FOV position: centered rostro-caudally at level of 4th spinal disc, GRAPPA acceleration factor: 2, partial Fourier factor: 6/8, phase-encoding direction: AP, echo-spacing: 0.47ms, bandwidth per pixel: 1220 Hz/Pixel, slice angulation: perpendicular to each participant's spinal cord, fat saturation and anterior and posterior saturation bands). All EPI acquisitions were performed with automatic slice-wise z-shimming (Finsterbusch et al., 2012; Kaptan et al., 2022). Three initial dummy shots were performed before the first functional image was acquired to achieve steady-state conditions. With the employed repetition time and flip angle, this approach brought all MR images to within 0.12% of the steady-state signal for gray matter, allowing us to include all images in the analysis. We

also acquired a high-resolution T2-weighted structural scan via a SPACE sequence with a
resolution of 0.8 x 0.8 x 0.8mm (slice orientation: sagittal, repetition time (TR): 1.5s, TE: 0.12s,
FA: 120°, number of slices: 64, field-of-view (FOV): 256 x 256 mm^2, GRAPPA acceleration factor:
3, bandwidth per pixel: 625 Hz/pixel; Cohen-Adad et al., 2021).

In addition to the MRI data, we acquired peripheral physiological data (respiration, heart rate, skin
conductance and pupil diameter) throughout the entire experiment on both measurement days.
Respiration, heartbeat, and skin conductance responses were recorded using a BrainAmp ExG
system (Brain Products, Gilching, Germany) and pupil diameter was assessed via the Eyelink
1000 Plus system (SR research, Ottawa, Canada). Furthermore, after every run, the participants
were asked to verbally rate the average intensity of the stimuli on a numerical rating scale (NRS)
ranging from 0 to 100, where 0 translated to "no percept", 50 marked the pain threshold and 100
translated to "unbearable pain".

2.5 Peripheral physiological data analysis
2.5.1   Heart period responses (HPR)
The ECG data were preprocessed using EEGLAB (Delorme & Makeig, 2004) and the FMRIB plug-
in for EEGLAB, provided by the University of Oxford Centre for Functional MRI of the Brain
(FMRIB) to remove MR-artifacts from the data traces recorded during functional runs (Niazy et al.,
2005). Using in-house Matlab scripts, R-peaks were automatically detected, and manually
corrected, if necessary. To obtain heart period time series, each inter-beat interval (IBI) was
assigned to its following R-peak and the resulting IBI time series was linearly interpolated to
achieve a sampling rate of 10Hz. Additionally, we filtered the IBI time series using a second-order
Butterworth band-pass filter with cut-off frequencies at 0.01 Hz and 0.5 Hz (Paulus et al., 2016).
The IBI traces were subdivided into event-epochs of -1 to 10s relative to stimulus onset and
baseline-corrected to the average IBI within 5s before until stimulus onset. We then extracted the
minimum of the IBI trace in an interval of 0 - 8s after stimulus onset of each trial and averaged the
resulting 20 HPR values of the Reliability Run per day and participant. To test for differences in
the HPRs between both days, we entered the average HPR of each participant and day into a
pair-wise two-sided t-test.

2.5.2   Skin conductance responses (SCR)
In two participants, SCR could not be recorded due to technical issues, leading to a sample size
of 38 participants for SCR analyses. SCR data were down-sampled to 100Hz and low-pass filtered
with a cut-off frequency of 1 Hz. The SCR traces were subdivided into event-epochs of -1 to 10s

relative to stimulus onset and baseline-corrected to stimulus onset. We then extracted the peak of
the skin conductance trace in an interval of 0 - 8s after stimulus onset of each trial and averaged
the resulting 20 SCR values of the Reliability Run to acquire one average peak value per day and
participant. To test for differences in the SCRs between both days, we entered the average SCR
of each participant and day into a pair-wise two-sided t-test.

2.5.3   Pupil dilation responses (PDR)

In six participants, eye tracking data of sufficient quality could not be recorded, leading to a sample
size of 34 participants for pupil dilation analyses. Eyeblinks that were automatically detected by
the EyeLink software were removed within a period of ± 100ms surrounding each blink. After the
automatic blink detection, we manually corrected for any additional blinks or artifacts in the data
trace by interpolating across the affected data segments. Blinking periods or otherwise missing
data were interpolated linearly and the data were down-sampled to 100Hz and low-pass filtered
with a cut-off frequency of 4 Hz. The pupil data traces were subdivided into event-epochs of -1 to
10s relative to stimulus onset and baseline-corrected to stimulus onset. We then extracted the
peak of the pupil dilation trace in an interval of 0 – 4s after stimulus onset of each epoch and
averaged the resulting 20 PDR values of the Reliability Run to acquire one average peak value
per day and participant. To test for differences in the PDRs between both days, we entered the
average PDR of each participant and day into a pair-wise two-sided t-test.

2.6 fMRI data analysis

Preprocessing and statistical analyses were carried out using FSL (version 6.0.3), SCT (Version
5.5) as well as custom MATLAB (version 2021a) and Python (version 3.9.13) scripts. The following
procedures were carried out separately for the Reliability Run of each measurement day and
participant.

2.6.1   Preprocessing

Correction for thermal noise. As a first step, we applied non-local Marchenko-Pastur principal
component analysis (MP-PCA, https://github.com/NYU-DiffusionMRI/mppca_denoise, Veraart et
al., 2016) on the unprocessed EPI data of the Reliability Run to reduce thermal noise (Ades-Aron
et al., 2021; Diao et al., 2021; Kaptan et al., 2023). The application of MP-PCA resulted in a
substantial spinal cord tSNR increase (63.6%; from 11.69 before MP-PCA to 19.12 after MP-PCA),
but only a marginal increase in spatial smoothness in the spinal cord (5.7%; from 1.23 before MP-
PCA   to   1.30   after   MP-PCA;   estimated   via   AFNI's   3dFWHMx   tool:
https://afni.nimh.nih.gov/pub/dist/doc/program_help/3dFWHMx.html)

Motion correction. Motion correction was carried out in two steps. We first created a mean image

of the 160 EPI volumes (after thermal noise correction). This mean image was used as a target

image for the motion correction as well as to automatically segment the spinal cord. Based on the

segmentation, we created a cylindrical mask, which was used to prevent adverse effects of non-

spinal movement on the motion parameter estimation. Motion correction was then carried out

slice-wise (allowing for x- and y-translations), using spline interpolation and a 2^{nd} degree

polynomial function for regularization along the z-direction (De Leener et al., 2017). As a second

step, we repeated the motion correction of the original time series of the Reliability Run, now using

the mean image of the initially motion-corrected time-series as the target image.

Segmentation. In order to obtain a high-resolution segmentation of the spinal cord, we used the

T2-weighted acquisition with 0.8mm isotropic voxels. For this purpose, we first applied the

ANTs N4 bias field correction algorithm on the raw structural data to correct for intensity

inhomogeneities due to RF coil imperfections (Tustison & Gee, 2010). As a next step, we denoised

the structural data via Adaptive Optimized Nonlocal Means (AONLM) filtering (Manjón et al., 2010)

to account for spatially varying noise levels and increase the SNR. To improve the robustness and

quality of the final segmentation, we used an iterative procedure: the data were initially segmented

using the SCT DeepSeg algorithm (Gros et al., 2019), smoothed along the z-direction using an

anisotropic Gaussian kernel with 6mm sigma (in straightened space), and again segmented via

the DeepSeg algorithm. To obtain a spinal cord segmentation of the EPI data, we used the mean

image of the motion-corrected time series as input for SCT's DeepSeg algorithm.

Registration to template space. While the statistical analyses on the individual level took place in

each participant's native space, group-level analyses were carried out in a common anatomical

space defined by the PAM50 template of the spinal cord (De Leener et al., 2018). For the individual

transformations from native to template space, we utilized the denoised and segmented structural

T2-weighted image. In line with SCT's recommended registration procedure (De Leener et al.,

2017), the vertebral levels were identified and labeled, and the spinal cord was straightened. Using

an iterative, slice-wise non-linear registration procedure based on segmentations, the structural

image was then registered to the template. The resulting inverse warping field served to initialize

the registration of the PAM50 template to the motion-corrected mean functional image via SCT's

multi-step non-rigid registration (De Leener et al., 2017). Based on this registration, we obtained

a warping field to move the native-space mean functional image of each participant to template

space using spline interpolation.

Correction for physiological noise. We employed several steps to reduce physiological noise. First,

we identified volumes with excessive motion whose effects we aimed to remove during general

linear model estimation. For this purpose, we calculated the root mean square difference between

successive volumes (dVARS) as well as the root mean square intensity difference of each volume

to a reference volume (refRMS) using FSL's fsl_motion_outliers algorithm. Volumes presenting

with dVARS or refRMS values three standard deviations above the time series mean were defined

as outliers and individually modelled as regressors of no interest in subsequent analyses (on

average 2% [range: 0.6% - 5.6%] of the 160 volumes per run were regarded as outliers). Second,

the respiratory and preprocessed cardiac signals (see section 2.4.1 electrocardiogram) were used

to create a physiological noise model (PNM), which approximates to what extent fMRI signal

changes can be explained by respiratory and cardiac activity (Brooks et al., 2008). The

approximation is based on the estimated cardiac and respiratory cycle phase during which the

slices were obtained. The approach is derived from the retrospective image correction procedure

(RETROICOR; Glover et al., 2000) and has been adapted for spinal cord fMRI to obtain slice-wise

physiological regressors for subsequent analyses (Brooks et al., 2008; Kong et al., 2012). We

extracted 32 noise regressors to estimate cardiac, respiratory and interaction effects as well as

an additional regressor to model the cerebrospinal fluid (CSF) signal, which was derived from

voxels in the CSF and spinal cord space that exhibited high levels of signal variance.

2.6.2　Statistical analysis

General linear model. The statistical analysis of the fMRI data was based on the general linear

model (GLM) approach implemented in FSL's FEAT (FMRI Expert Analysis Tool;

http://fsl.fmrib.ox.ac.uk/fsl/fslwiki/FEAT; Woolrich et al., 2001) and included spatial smoothing via

FSL's Susan tool with an isotropic 2mm (full width half maximum) Gaussian kernel as well as high-

pass filtering at 100s. The first-level design matrix included a regressor for the heat stimulus

onsets convolved with a double-gamma hemodynamic response function (HRF) as well as a

temporal derivative. The following regressors of no interest were added to the design matrix for

robust denoising: 33 slice-wise PNM regressors (describing cardiac, respiratory and CSF effects),

two slice-wise motion regressors (describing movement along x and y; calculated during motion

correction), and one regressor for each volume with excessive motion. From the first-level analysis

we obtained a β-estimate map for the main effect of heat for each participant and day, which we

registered to the template using the previously estimated warping fields.

Masks. For the subsequent analyses, we used multiple masks. In template space, we first derived

z-coordinates to divide the spinal cord according to the spinal segmental levels C5 to C8 (which

are the levels covered by our EPI slice-stack). The coordinates for each segment were obtained

from SCT (version 6.1, De Leener et al., 2018) and are based on findings from Frostell et al.,

(2016). Segmental masks for the four gray matter horns of the spinal cord were derived by

cropping the unthresholded probabilistic gray matter masks of each gray matter horn according to the same segmental coordinates. Additionally, we utilized segmental masks for the four quadrants of the cord, encompassing both gray matter and white matter, using the same segmental coordinates. The mask of the left dorsal horn in segment C6 (number of isotropic 0.5mm voxels: 502) was used to investigate the main effect of heat, as well as the reliability between the days, along with the mask of the right ventral horn of segment C6 (number of isotropic 0.5mm voxels: 719), which was defined as a control region. To assess BOLD activity patterns beyond the target region, we used a cord mask of spinal segment C6 dilated by 6 voxels (i.e., including an area occupied by draining veins), which we then subdivided into 4 quadrants (left dorsal, number of isotropic 0.5mm voxels: 5888; left ventral, number of isotropic 0.5mm voxels: 6671; right dorsal, number of isotropic 0.5mm voxels: 5888, right ventral, number of isotropic 0.5mm voxels: 6671).

Average and day-wise BOLD responses. To first investigate whether phasic heat stimulation as employed here evokes a significant BOLD response at all, we averaged the normalized β-maps over both days within each participant and then submitted the resulting 40 β-maps to a one-sample t-test. Correction for multiple comparison was carried out via voxel-wise non-parametric permutation testing as implemented in FSL's randomise algorithm (Winkler et al., 2014) in an anatomically informed target region (left dorsal horn, segment C6) at a threshold of $p_{FWE} < 0.05$ (family-wise error corrected). Second, we also aimed to test for significant responses on each of the two days using exactly the same statistical procedure, but now using the 40 β-maps from each day as input. Finally, we aimed to test for an overlap of significant responses across both days (i.e., a conjunction) and therefore created a binary mask of the significant voxels of each day, which we subsequently multiplied with each other to determine significant voxels overlapping across both days.

Spatial specificity. We also aimed to describe the spatial organization of the BOLD responses in the part of the spinal cord covered by our slice-stack. For this purpose, we performed a one-sample t-test (again within a permutation-testing framework) using the averaged β-estimates over both days within a cord mask including spinal segments C5 to C8. From the resulting uncorrected group-level p-maps, we assessed the number of voxels surviving liberal thresholding at $p < 0.001$ uncorrected in each spinal segment. We then calculated what percentage of the total number of active voxels in the respective cord segment were located in the dorsal left, dorsal right, ventral left and ventral right cord quadrant (including both gray and white matter parts of the cord). In order to supplement this analysis with more fine-grained information regarding gray matter responses, we additionally assessed the number of supra-threshold voxels within the four gray matter horns.

Finally, we aimed to assess to what degree BOLD responses also occur 'outside' the spinal cord

proper. While our target tissue of interest is the gray matter of the dorsal horn, this is drained

through a hierarchy of veins: small veins coalesce into radially-oriented intramedullary veins,

which further drain into the circumferential spinal veins of the superficial pial venous plexus, a

structure itself permeated by longitudinal veins. From here, the blood drains into the internal

vertebral plexus before progressing to the external vertebral plexus, ultimately joining the systemic

circulation (Thron, 2016). Taking into account this network of draining veins outside the gray

matter is crucial for spinal cord fMRI, as these venous pathways can influence the spatial

specificity of the BOLD response, potentially diluting the signal across a larger area than the region

of neuronal activity. To better understand the resulting signal spread, we performed an additional

group analysis within an extended region encompassing spinal segment C6, dilated by 4 voxels

(i.e., 2mm, covering parts of the venous drainage system, such as the pial venous plexus) on a

slice-by-slice basis.

2.7 Reliability

2.7.1   Contextual differences

Before quantifying the reliability of the response measures, we tested for differences between the

scanning days that could explain changes in the BOLD fMRI results, such as differences in the

general physiological state of the participants and in the fMRI data quality. To assess differences

in the physiological state of participants, we calculated three metrics: heart rate, heart rate

variability and spontaneous fluctuations of electrodermal activity. All three metrics served to

describe underlying differences in tonic autonomous nervous system activity, e.g. due to stress or

emotional arousal (Bach et al., 2010; Berntson et al., 2017; Dawson et al., 2017). Heart rate was

quantified as beats per minute (bpm) and heart rate variability (HRV) was calculated as the root

mean square of successive peak-to-peak interval differences between normal heartbeats

(RMSSD) in milliseconds. Spontaneous fluctuations in electrodermal activity were calculated by i)

setting up a GLM where stimulus onsets were convolved with a canonical skin conductance

response function (implemented in PsPM: https://bachlab.github.io/PsPM/ ; Bach et al., 2009,

2013) and ii) then using the residual activity (i.e. after removal of modelled stimulus-evoked

responses) to calculate the area under the curve of the remaining skin conductance traces (Bach

et al., 2010). To describe the overall quality of the fMRI data, we i) estimated motion by calculating

the root mean square intensity differences of each volume to a reference volume and ii) calculated

the temporal signal-to-noise ratio (tSNR) of the motion-corrected EPI data.

2.7.2   Intra-class correlation coefficient

To assess the reliability of responses to painful heat stimulation across two days, we calculated

the intra-class correlation coefficient according to Shrout & Fleiss (1979), a widely used statistical

measure to assess the reliability of repeated measurements. Specifically, ICC(3,1) serves to

assess the consistency of measurements across different occasions or days, and is defined as

the ratio of the between-participant variance and the total variance (Caceres et al., 2009). ICC(3,1)

was calculated using the following formula:

$$ICC(3,1) = \frac{BMS - EMS}{BMS + (k-1)EMS}$$

The formula includes three components: BMS (Between Participants Mean Squares), which

represents the variance between different participants; EMS (Error Mean Squares), representing

the residual variability, which includes inconsistencies in repeated measurements for the same

participant; and k, indicating the number of measurements (Caceres et al., 2009). The numerator

(BMS - EMS) reflects the variability between participants after accounting for measurement errors,

while the denominator (BMS + (k - 1) * EMS) combines the total variability between participants

with measurement errors adjusted by the number of measurements. This ratio quantifies the

consistency of the measurements and indicates the degree to which participants maintain their

relative ranking across days (Caceres et al., 2009; Liljequist et al., 2019). The calculation of the

reliability coefficients for all of the measures described below was implemented via the Python

package Pingouin (version 0.5.3, Vallat, 2018). The interpretation of the resulting reliability

estimates followed the conventions by Cicchetti (1994), where ICC values smaller than 0.4

indicate poor reliability, values from 0.4 - 0.59 imply fair reliability, values from 0.6 - 0.74 represent

good reliability and values from 0.75 - 1.0 are defined as excellent reliability.

2.7.3   Subjective and peripheral physiological responses

We calculated the test-retest reliability for the verbal ratings of pain intensity (data from two

participants were missing due to technical issues, resulting in N = 38) as well as for SCR, PDR,

and HPR. For verbal ratings, we employed the single rating obtained after all trials and for the

peripheral physiological measures, we employed the peak response value (averaged across trials)

of each participant obtained on each of the two days (see sections 2.5.1., 2.5.2, 2.5.3 for closer

description of peak value extraction).

2.7.4   BOLD responses

To assess the test-retest reliability of spinal cord heat-evoked BOLD responses across two

consecutive days, we extracted β-estimates for the main effect of heat for each participant and

session. We assessed the reliability through the application of different anatomical masks (see

section 2.6.2, under *Masks*). Two masks covered the area of interest, one of them limited to the

gray matter horn, while the other mask incorporated draining vein territory, and two further masks

captured the gray matter in a control region as well the draining vein territory adjacent to the control

region. The first mask was the left dorsal horn mask on spinal segment C6, the second mask was

the left dorsal quadrant of the dilated cord mask on the level of C6, the third mask was the right

ventral horn on C6 and the fourth mask was the right ventral quadrant of the dilated cord mask of

C6. The calculation of reliability in the left dorsal horn, was based on three metrics, namely the

mean β over the entire region of interest, the peak β estimate in the ROI regardless of its location

within the ROI, and finally the average of the top 10% β values in the ROI. We also calculated the

reliability for the ROI mean, peak value and average of the top10% using the z-scores from the

z-maps, since the z-scores scale the parameter estimate (β maps) by the standard error of each

voxel, thereby considering the underlying variation within runs.

The reliability assessment described so far aimed to quantify the similarity of the response

amplitudes, quantified via the β estimates or z-scores, over both days. However, in the context of

fMRI, not only the response amplitude holds importance but also the spatial patterns of the

response – specifically, we wanted to know whether the BOLD response on Day 1 occurred in the

same location as the BOLD response on Day 2. To compare the spatial patterns of the BOLD

responses between days, we calculated Dice coefficients, which quantified the amount of overlap

of the active voxels in the left dorsal horn in spinal segment C6 (Rombouts et al., 1999; Wilson et

al., 2017).

$$Dice\ coefficient = \frac{2\ x\ V_{overlap}}{(V_1 + V_2)}$$

V_1 and V_2 define the number of active, i.e., above-threshold voxels on each day, and $V_{overlap}$ is the

number of voxels that overlap. We calculated Dice coefficients on the group and individual level

using binarized statistical maps. On the group level we binarized the uncorrected p-maps at the

thresholds 0.001, 0.01 and 0.5, and on the individual level we binarized the z statistic image for

the main effect of heat, thresholded at +/- 1.96 (i.e., p < 0.05 uncorrected).

2.8 Post-hoc analyses

As reported in the Results section, test-retest reliability was low for the peak activation in the left

dorsal horn on C6. Additionally, in the same ROI there was no spatial overlap between the group-

level results of both days, accompanied by a low Dice coefficient for even rather liberal thresholds.

To investigate possible reasons for this surprising lack of response consistency, we carried out

three further sets of analyses, which we had not specified in the preregistration.

2.8.1 Increasing the number of runs

The Reliability Run was optimized for assessing reliability in terms of keeping the measurement parameters and stimulation position on the arm identical across days. However, since one run consisted only of 20 trials and our stimulus duration of 1s was relatively short, this data is likely noisier than data of spinal fMRI paradigms with more trials or more prolonged stimuli, which might explain the low reliability. For this reason, we also investigated the fMRI activation maps and reliability metrics using the average over multiple runs per day, resulting in a four-fold increase of trials, since we included all runs with a TR of 1800ms (including the Reliability Run), i.e. four runs per day (the two additional runs with shorter TRs were excluded from this analysis due to the different TR and flip angle employed in those acquisitions).

The preprocessing followed the steps described above and was done separately for each day. To bring all individual runs into a common space, motion correction was carried out exactly as described in section 2.6.1, however, instead of correcting a single run we concatenated all suitable runs of the respective day and motion-corrected the entire concatenated time series. Registration to template space followed the identical procedures as above, only now the mean image of the concatenated and motion-corrected time series served as the destination image. For all subsequent analyses we used the same procedures as described above (section 2.6.2), with the difference that the β maps of the individual runs were averaged across runs and only then entered the group-level analysis. For the rating and all peripheral physiological data, we also combined the data of the four runs per day and calculated the reliability coefficients accordingly. The results of this analysis are referred to as "Combined Runs".

2.8.2 Accounting for spontaneous activations

Another cause of the low reliability of task-evoked BOLD responses could be spontaneous fluctuations in the BOLD signal, which were not accounted for in the GLM, and which might increase trial-to-trial variability. A study by Fox and colleagues (2006) showed that trial-to-trial BOLD response variability in the left somatomotor cortex could be reduced by regressing out the BOLD signal of the right somatomotor cortex. The authors argue that the spontaneous fluctuations of both regions correlated due to the interhemispheric connectivity between the regions. Regressing out the signal of the opposite hemisphere mainly decreased noise, whereas the accompanied reduction of the task-relevant signal was non-significant. Since previous spinal fMRI studies have found evidence for resting-state functional connectivity between the left and right dorsal horn (Barry et al., 2014, 2016; Eippert, Kong, Winkler, et al., 2017; Harita & Stroman, 2017; Kaptan et al., 2023; Kinany et al., 2020; Kong et al., 2014; Vahdat et al., 2020), we aimed to test if a similar analysis strategy could decrease noise due to spontaneous fluctuations, and improve

reliability (though we are aware that this could also be negatively affected e.g. due to pain-induced
responses in contralateral dorsal horns (Fitzgerald, 1982). For this purpose, we extracted the time
series of the contralateral (right) dorsal horn of each slice, and used it as an additional slice-wise
regressor in the GLM to regress out spontaneous fluctuations in the ipsilateral (left) dorsal horn.
Otherwise, the analysis followed the procedure outlined above.

2.8.3   Within-run reliability

Given the low reliability across days, we wanted to assess if reliability would be equally low within
runs, as such comparisons would not involve the potentially detrimental impact of repositioning
the participants in the scanner as well as possibly imperfect matches of the normalized parameter
estimate maps in template space. For this purpose, we adopted a split-half approach and divided
the Reliability Run into two subsets: odd and even trials for the odd-even reliability analysis, or the
first and second half for the early-late reliability analysis, with the corresponding trial regressor
entered in the general linear model (GLM). Subsequently, we obtained two β maps from both the
odd-even and early-late GLMs, representing the respective trial selections. These β maps were
then subjected to the spatial normalization procedure described in section 2.6.1. Reliability
coefficients (ROI mean, ROI peak, average of top 10% β estimates extracted for each participant)
were calculated between the respective trial selections of both approaches, separately for each
day and the resulting ICC scores were averaged across days. We also calculated both within-run
reliability measures for SCR, PDR and HPR, calculating the respective response peaks for each
set of trials and averaging ICC values across days.

2.8.4   Correlations between BOLD and non-BOLD response measures

In an additional exploratory analysis inspired by a reviewer's comment, we assessed the across-
participant correlations between peripheral physiological as well as subjective responses and
BOLD responses (results are reported in Supplementary Table 3). For this purpose, for every
participant we averaged the top 10% β estimates and z-scores in the left dorsal horn of C6, along
with SCR, PDR, HPR and subjective ratings across days. We then calculated Pearson's r for each
of the eight correlations (only including participants with responses in both variables): β estimates
with SCR, HPR, PDR, and rating as well as z-scores with SCR, HPR, PDR, and rating. Since we
assumed that higher BOLD responses would go along with stronger peripheral physiological and
subjective responses (positive associations expected for all responses but HPR, as here a
negative-going response indicates cardiac acceleration as is typical in response to nociceptive
input), we base our results on one-tailed p-values as indicators of statistical significance of the
correlation strength.

2.8.5   Correlations between BOLD parameter estimates and indicators of data quality

Inspired by a reviewer's comments, we calculated correlations between changes in data quality

metrics and BOLD parameter estimates across days, as this should allow for insights into possible

data quality contributions to across-day reliability of BOLD responses. For every participant we

calculated the difference from Day 1 to Day 2 for i) motion estimates, ii) normalization quality

estimates and iii) indicators of participant positioning. Motion estimates were obtained via root

mean square intensity differences of each motion-corrected volume to reference volume (see

refRMS, section 2.6.1). Normalization quality was estimated via computing Dice coefficients

between the segmentation of the normalized EPI and the PAM50 cord mask (see section 2.7.4 for

Dice coefficient). Participant positioning estimates were obtained by calculating the angulation of

the slice stack relative to the direction of the B0 field, since the slice stack was always positioned

to be orthogonal to the longitudinal axis of the spinal cord (see Fig. 1 for example). The angle

between the normal vector of the slice package extracted from the DICOM header and the

scanner's z-axis (0,0,1) therefore serves as a proxy for the positioning of the neck – and thus the

orientation of the draining veins – relative to B0. For each of these measures, we correlated the

absolute difference across days with the absolute difference of BOLD parameter estimates across

days, quantified as the top 10% β estimates and z-scores of the left dorsal horn in spinal cord

segment C6. Since we expected a positive correlation (i.e., greater differences across days would

be associated with greater variation of BOLD responses), we report one-tailed p-values alongside

the correlations in Supplementary Table 4.

2.9 Open Science

This study was preregistered before the start of data acquisition and the preregistration is openly

available on the Open Science Framework (https://osf.io/a58h9); differences between the

analyses suggested in the preregistration and the analyses carried out here (as well as the

reasons behind these changes) are listed in the Supplementary Material. The underlying data and

code are currently only accessible to reviewers, but will be made openly available upon publication

via OpenNeuro and GitHub, respectively. The intended data-sharing via OpenNeuro was

mentioned in the Informed Consent Form signed by the participants and approved by the Ethics

Committee at the Medical Faculty of the University of Leipzig.

 3. Results

3.1 Behavioral and physiological responses

Across both days, the participants reported an average stimulus intensity of 71.7 (Fig. 2, left, n = 38, SD = 12.1), indicating that the employed heat stimuli were perceived as clearly painful (responses greater than 50 indicated pain). This subjective percept was accompanied by robust physiological changes (Fig. 2), as evidenced in skin conductance responses (SCR), pupil dilation responses (PDR) and heart period responses (HPR). All measures showed rather similar responses when compared across days ($t_{rating}(37) = 0.15$, $p_{rating} = 0.88$; $t_{SCR}(37) = 0.86$, $p_{SCR} = 0.39$; $t_{PDR}(33) = 0.67$, $p_{PDR} = 0.51$; $t_{HPR}(39) = 0.48$, $p_{HPR} = 0.63$; Fig. 2).

Figure 2. Subjective and peripheral physiological responses. Left: verbal ratings of stimulus intensity on a numerical rating scale with 50 indicating the pain threshold; half-violins and boxplots depict the distribution over participants and grey dots show the raw values (jittered slightly for visualization purposes). Right: Averaged traces of the skin conductance response (SCR), pupil dilation response (PDR) and heart period response (HPR) in response to the stimulus, with error bands reflecting the standard error of the mean across the group and the gray rectangle representing the stimulus duration.

3.2 BOLD responses: amplitudes

Averaged across days, we observed a significant response in the ipsilateral dorsal horn in spinal segment C6 ($t(39) = 4.51$, $p_{corr} = 0.002$, 61 supra-threshold voxels; Fig. 3). Using the same

analysis parameters, we also observed significant responses for each day separately
(t_{day1}(39) = 3.58, p_{corr} = 0.035, 2 supra-threshold voxels; t_{day2}(39) = 4.50, p_{corr} = 0.0018, 48 supra-
threshold voxels). When comparing the spatial pattern of active voxels (at a threshold of p < 0.05
corrected) for Day 1 and Day 2, there was no overlap, with the active voxels of Day 1 being located
consistently more caudal in segment C6 compared to the active voxels of Day 2 (Fig. 3, sagittal
view), despite the heat stimulation occurring at the identical location on the forearm.

Figure 3. BOLD responses. Axial view of group-level results in the left dorsal horn in spinal segment C6 thresholded
at p_{FWE} < 0.05, with the average across both days depicted on the very left, followed by Day 1 and Day 2. The rightmost
plot shows a sagittal view of the activation maps of both days, with purple voxels belong to Day 1 while red voxels
belong to Day 2; the green outline marks spinal cord segment C6 for visualization purposes. Data are overlaid on the
T2*-weighted PAM50 template in axial views, and on the T2-weighted PAM50 template in the sagittal view.

3.3 BOLD responses: spatial specificity

3.3.1 Entire spinal cord

To assess the spatial specificity of BOLD responses, we used the group-level results of the across-
day average within the cord mask from spinal segment C5 to C8. We counted the number of active
voxels in each segment using a liberal threshold (p < 0.001 uncorrected) and assessed what
percentage of those voxels was located in each of the 4 cord quadrants: left dorsal, right dorsal,
left ventral and right ventral (Fig. 4; for exact percentages and day-wise results see Supplementary
Table 1). The highest number of voxels that survived thresholding was located in spinal cord
segment C5, followed by C6 and C7, with C8 holding the lowest number of supra-threshold voxels.
As can be seen in the percentages, segments C6 demonstrated the highest level of spatial
(i.e., neuroanatomical) specificity, followed closely by C5. In both segments, the majority of active
voxels were concentrated in the left dorsal quadrant, and a smaller number of active voxels were
found in the right dorsal quadrant, with a relatively small percentage of active voxels observed in
the ventral region. Conversely, the above-threshold voxels in C7 were mostly located in the right
ventral quadrant, and in spinal segment C8 in the right dorsal part.

Figure 4. Spatial specificity of BOLD responses across cord quadrants. Number of supra-threshold voxels across the four cord quadrants of all spinal segments from C5 to C8. The total on top of each bar shows the overall number of active voxels in the entire cord mask of each level. Colors indicate the number of active voxels across the dorsal left (DL) and right (DR) as well as ventral left (VL) and right (VR) quadrants. All results shown here are based on the group-level across-day average (uncorrected p < 0.001).

3.3.2 Gray matter

In order to obtain a more detailed understanding of spatial specificity in the spinal cord gray matter (instead of the cord quadrants, as reported above), we also projected all active voxels in the four gray matter horns onto an exemplary spinal cord slice, either from all segments (grey dots in Fig. 5A,) or only from target segment C6 (red dots in Fig. 5A) and furthermore visualized their distribution along the left-right and dorsal ventral axis. Across all segments, the highest number of active voxels was located in the left dorsal horn, but a substantial number of active voxels was also present in the other horns. Conversely, for our target segment C6, the clear majority of voxels is located in the ipsilateral dorsal horn, with a lesser number of voxels in the contralateral dorsal horn and no active voxels in the ventral horns.

3.3.3 Surrounding tissue

To investigate the impact of draining veins on the location of BOLD responses, we also assessed the spatial pattern of active voxels (p < 0.001 uncorrected, across-day group-level average) in a mask of spinal segment C6 that also included the venous plexus (Fig. 5B). Several aspects are worth noting here. First, in line with the previously presented data, there is almost no ventral horn activation and thus also no BOLD responses in the venous plexus on the anterior surface of the cord. Second, gray matter responses are consistently present throughout segment C6 in the ipsilateral dorsal horn and with lesser prominence also in the contralateral dorsal horn. Most importantly though, the strongest BOLD responses are actually observed at the dorsal surface of the cord in the region of the veins draining the dorsal cord: these responses are evident both ipsi- and contralaterally, at times spanning both the left and right dorsal surface.

Figure 5. Spatial specificity of BOLD responses. A: Positions of supra-threshold voxels in the gray matter horns of all spinal segments from C5 to C8, collapsed over z (grey), vs. only in spinal segment C6 (red), with jitter added for visualization purposes. The lines on outside of plot show the distribution of the voxels across hemicords (lines on the left average over the left and right horns, lines on the bottom average over the dorsal and ventral horns); colors indicate the employed mask (red: only spinal segment C6, grey: collapsed over z). The mask used for visualization here was obtained by combining a slice of the cord and all four GM horn masks in C6. B: Six example slices across segment C6 showing supra-threshold voxels within a dilated cord mask to allow for observing draining vein responses. All results shown here are based on the group-level across-day average (uncorrected p < 0.001).

3.4 Reliability

3.4.1   Physiological state and data quality across days

To test for differences in the participants' general physiological state across days, we calculated run-wise heart rate, heart rate variability, and spontaneous fluctuations of the electrodermal activity (Fig. 6A-C). While heart-rate showed a slight increase from Day 1 to Day 2 (t(39) = -2.15, p = 0.04), neither heart rate variability (t(39) = 1.23, p = 0.22) nor spontaneous fluctuations in electrodermal activity (t(37)=0.04, p = 0.97) showed significant differences across days. To assess fMRI data quality across days, we investigated motion effects (quantified as root mean square intensity difference to a reference volume for each run per participant and day; Fig. 6D) and temporal signal-to-noise ratio (tSNR; Fig. 5E) after motion correction. Neither motion effects (t(39) = 1.1, p = 0.28) nor tSNR (F(1,38) = 0.3, p = 0.59) showed significant differences across days; for tSNR this pattern held across all slices.

Figure 6. Physiological state and data quality across days. A – D show the average physiological state or fMRI quality indicators of each participant and day, visualized via box-plots, half-violin plots and grey lines that indicate participant-wise changes across days. A) Heart-rate quantified as beats per minute. B) Heart-rate variability quantified as root mean square of successive differences between normal heartbeats in ms. C) Spontaneous fluctuations in skin conductance (SCF) quantified as area under the curve. D) fMRI motion quantified as root mean square intensity differences of each volume to reference volume. E) fMRI signal quality quantified as temporal signal-to-noise ratio (tSNR) within a cord mask of each slice.

3.4.2 Behavioral and peripheral physiological test-retest reliability

As a positive control analysis, we first assessed the reliability of the behavioral and peripheral physiological measures in order to ascertain that responses to noxious thermal stimulation can in principle be reliably assessed (Fig. 7, Table 1; Supplementary Figure 1). Subjective ratings (ICC = 0.72), skin conductance (ICC_{SCR} = 0.77) and heart period (ICC_{HPR} = 0.77) exhibited good-to-excellent test-retest reliability, whereas pupil dilation (ICC_{PDR} = 0.34) showed poor test-retest reliability.

3.4.3 fMRI test-retest reliability

We calculated the reliability of the BOLD response amplitudes using four different masks: i) the left dorsal horn of C6 (ROI; grey matter area of interest), ii) an enlarged mask of the dorsal left

cord quadrant of C6 (dilated ROI; including the venous plexus containing veins draining the dorsal

horn), iii) the right ventral horn of C6 (control; grey matter area of no interest), and iv) an enlarged

mask of the right ventral quadrant of C6 (dilated control; including the venous plexus with veins

draining the ventral horn). To our surprise, all investigated metrics (β estimates of i) peak voxel,

ii) top 10% of voxels, and iii) average across all voxels) showed poor reliability (all ICC < 0.4) in

our target region, i.e., the left dorsal horn. This pattern did not change when we also took into

account noise at the individual level, by repeating these analyses on z-values instead of β-

estimates (with the latter only reflecting response amplitudes without taking into account residual

noise; see Table 1). When assessing a larger region however (including draining vein territory),

reliability was in the poor to fair range (ICC between 0.23 and 0.59). For the control areas, reliability

in the gray matter region was consistently in the poor range (all ICC < 0.4), and in the poor to fair

range (ICC between 0.41 and 0.53) in the dilated control region (as shown in Fig. 7).

Figure 7. Test-retest reliability across both days for subjective ratings, peripheral physiological data and BOLD

response amplitudes. Reliability is indicated via ICCs (plotted as dots with 95% CI represented as a line). ICCs are

reported for (from left to right) verbal ratings, skin conductance response amplitude (SCR), pupil dilation response

amplitude (PDR), heart period response amplitude (HPR), top 10% β-estimate in the left dorsal horn of C6 (ROI), in the

dilated left dorsal quadrant of C6 (dilated ROI), in the right ventral horn of C6 (control), and in the dilated right ventral

quadrant of C6 (dilated control). Colors indicate ICC interpretation according to Cicchetti (1994): dark red: ICC <

0.4, poor; medium red: ICC 0.4 - 0.59, fair; orange: ICC 0.6 - 0.74, good; yellow: ICC 0.75 - 1.0, excellent. Individual

values underlying the ICC calculation are shown in Supplementary Figure 1.

Table 1

Intraclass correlation coefficient (ICC) and 95% confidence interval for subjective ratings, peripheral physiological data and BOLD response amplitudes.

Measures				ICC (95% CI)
		Ratings		0.72 (0.53–0.85)
		SCR		0.77 (0.61–0.88)
		PDR		0.34 (0–0.60)
		HPR		0.77 (0.55–0.85)
fMRI ROI	Dorsal horn left	β	peak	0.20 (-0.12–0.48)
			top 10%	0.20 (-0.11–0.48)
			avg	0.03 (-0.28–0.33)
		z-score	peak	0.24 (-0.08–0.51)
			top 10%	0.17 (-0.14–0.46)
			avg	0.10 (-0.22–0.39)
	Dilated left dorsal quadrant	β	peak	0.53 (0.26–0.72)
			top 10%	0.59 (0.35–0.76)
			avg	0.23 (-0.08–0.51)
fMRI Control	Ventral horn right	β	peak	0.18 (-0.13–0.46)
			top 10%	0.17 (-0.14–0.46)
			avg	0.17 (-0.15–0.45)
	Dilated right ventral quadrant	β	peak	0.53 (0.02–0.58)
			top 10%	0.45 (0.14–0.46)
			avg	0.41 (0.12–0.64)

3.4.4 fMRI spatial consistency

To also compare the spatial patterns of the BOLD responses between days, we calculated Dice

coefficients (DC). On the group level, the p-maps in the left dorsal horn of C6

(at p < 0.05 corrected) did not show any overlap across days (DC = 0). Using a more liberal

thresholding increased the DC slightly (uncorrected p < 0.01: DC = 0.009, uncorrected p < 0.05:

DC = 0.26). On the individual level (using an uncorrected p < 0.05), z-maps of 35 participants held

supra-threshold voxels in the left dorsal horn (C6), but in only 5 participants these overlapped

across days (mean DC across all 35 participants with suprathreshold voxels in ROI on both days:

0.05, mean DC of 5 participants with overlap: 0.33, range 0.02 – 0.65).

3.5 Post-hoc analyses

3.5.1   Increased number of runs

In order to assess whether an increase in stimulus numbers would lead to a higher reliability, we

used all runs with a TR of 1800ms (four runs per day instead of just one). When first assessing

response amplitude, we noticed that this led to a clear enhancement in the strength of the group-

level BOLD response, not only in the average across days ($t(39) = 6.64$, $p_{corr} < 0.001$, 331 supra-

threshold voxels), but also for Day 1 ($t(39) = 4.77$, $p_{corr} = 0.002$, 94 supra-threshold voxels) and

for Day 2 separately ($t(39) = 5.55$, $p_{corr} < 0.001$, 285 supra-threshold voxels; Supplementary

Figure 2). Most importantly, with the increased stimulus numbers we now observed an overlap of

activation across both days (94 supra-threshold voxels at $p < 0.05$ corrected), leading to an

improved dice coefficient (DC = 0.80 for group-level p-maps at $p_{corr} < 0.05$), meaning that spatial

consistency improved substantially. Contrary to our expectations, increasing the number of trials

did not lead to improvements in the reliability of the non-BOLD data or the BOLD response

amplitude in the target region (Supplementary Fig. 3 and Supplementary Table 2).

3.5.2   Accounting for spontaneous activity

When accounting for spontaneous fluctuations of BOLD activity in the left dorsal horn by adding

the time-course of the right dorsal horn as a slice-wise regressor to the GLM, we did not observe

an increase in test-retest reliability (Supplementary Fig. 3 and Supplementary Table 2).

3.5.3   Within-run reliability

Interestingly, the within-run reliability depended on the selection of trials. Between odd and even

trials, reliability in the target region (left dorsal horn, C6) reached the level "good" (ICC = 0.69,

Supplementary Fig. 3 and Supplementary Table 2) for the top 10% β estimates, and still "fair" for

the ROI average (ICC = 0.52), whereas comparing the first half of trials to the second resulted in

poor reliability (ICC = 0.36, top 10% β). Odd-even reliability was excellent for all three peripheral

measures (Supplementary Fig. 3 and Supplementary Table 2), and slightly lower, but still fair to

excellent, for the first vs. second half of trials.

                                                    4. Discussion

In this study we aimed to probe the limitations of task-based spinal fMRI – a young field facing

many challenges – by investigating the robustness of spinal cord BOLD responses to repeated

nociceptive stimulation across two consecutive days. For this purpose, we first examined if BOLD

activation patterns occurred in the expected region of the spinal cord and assessed the spatial

specificity of the response across a larger area. In our main investigation, we focused on the test-

retest reliability of the BOLD response amplitude as well as the consistency of its spatial pattern.

To disambiguate between effects on reliability of either data quality or variability in the underlying

process of nociception, we also assessed the reliability of several simultaneously-recorded

peripheral physiological measures.

4.1 Heat-pain evoked responses

In order to ascertain that the chosen stimulation parameters (contact heat stimuli at 48°C for 1s)

would elicit a robust response, we recorded subjective ratings as well as peripheral physiological

responses. Our results are in line with the general observation that painful stimulation activates

the autonomic nervous system (Boucsein, 2012; Cowen et al., 2015; Kyle & McNeil, 2014),

exemplified here by increased skin conductance, pupil dilation and heart rate in response to the

stimulus. Although these responses are not specific for pain per se, as they can generally indicate

increased arousal or salience (Lee et al., 2020), along with perceptual ratings being clearly above

pain threshold, they suggest that a robust pain response was evoked by our brief nociceptive

stimulation.

Our data furthermore showed that a brief contact heat stimulus of 1s can already evoke a

measurable group-level BOLD response in the dorsal horn of the spinal cord. This response was

observed in the expected segmental level and survived strict permutation-based correction for

multiple comparisons. In humans, there is ample evidence that heat pain stimulation leads to

activation in the ipsilateral DH of the spinal gray matter at the expected rostrocaudal location

(Bosma & Stroman, 2015; Brooks et al., 2012; Eippert et al., 2009; Geuter & Buchel, 2013; Nash

et al., 2013; Oliva et al., 2022; Seifert et al., 2023; Sprenger, Eichler, et al., 2018; Summers et al.,

2010; Weber et al., 2016b, see Kolesar et al., 2015 for review), though these studies have

consistently used longer stimulus durations. While results from one spinal fMRI study in the motor

domain suggested that short stimuli may elicit weaker BOLD responses than expected (Giulietti

et al., 2008), the application of short stimuli in event-related designs allows for a larger number of

trials and may therefore boost power, as well as enable greater variability in the timing of the

stimulus presentation (D'Esposito et al., 1999). Such features could be helpful for investigating

the cognitive modulation of pain (Atlas & Wager, 2012; Villemure & Bushnell, 2002; Wiech, 2016)

at the spinal level with more sophisticated paradigms than currently employed.

4.2 Spatial specificity

After confirming that BOLD responses indeed occurred in the ipsilateral dorsal horn of the

expected segment, we next investigated the response pattern beyond this target area, to allow

insights into the spatial specificity of BOLD responses. In the target segment of the spinal cord

(C6), the ipsilateral dorsal horn indeed showed the highest number of active voxels across all four

gray matter horns, though in all horns smaller numbers of active voxels were observed. Activation

beyond the ipsilateral dorsal horn has been reported in previous spinal cord fMRI studies (Cahill

& Stroman, 2011; Geuter & Buchel, 2013; Summers et al., 2010; Yang et al., 2015; see Kolesar

et al., 2015 for review) and one factor contributing to this could be spatial inaccuracies, e.g. due

to distortions during image acquisition, suboptimal registration to template space and spatial

smoothing (Bosma & Stroman, 2015; Eippert, Kong, Jenkinson, et al., 2017; Hoggarth et al.,

2022). On the other hand, activations in the ventral horns as well as the contralateral dorsal horn

could also be indicative of neural processing in these areas: not only is there evidence for

functional connectivity between the dorsal horns in the human spinal cord (for review, Harrison et

al., 2021), but also evidence from animal models for dorsal commissural interneurons (Bannatyne,

2006; Petkó & Antal, 2000) and primary afferents that project contralaterally (Culberson et al.,

1979; Light & Perl, 1979). Furthermore, autoradiographic rat data also show widespread

responses to noxious heat stimuli (Coghill et al., 1991, 1993) and a painful heat stimulus might

trigger motor responses as well as the active inhibition of thereof (Pierrot-Deseilligny & Burke,

2012; Purves et al., 2019). Taking these aspects into account, we would argue that activations

outside of the ipsilateral dorsal horn likely reflect more than noise.

Apart from activation in spinal cord segment C6, we also observed active voxels in segments C5,

C7 and C8 (when using a liberal uncorrected threshold), a pattern that has been observed

previously in human data (Geuter & Buchel, 2013; Rempe et al., 2015; Seifert et al., 2023; Weber

et al., 2016a). In addition, Shekhtmeyster et al. (2023) provide evidence for cross-segmental spinal

cord activation of glial cells of mice in response to nociceptive stimulation, hinting at similarly

widespread spinal processing mechanisms. On the one hand, a part of this large rostrocaudal

extent could be due to dermatomal variability between participants and the fact that adjacent spinal

roots can innervate overlapping areas of skin (Lee et al., 2008). However, this aspect seems to

be more relevant for tactile as opposed to pain and temperature dermatomes (Lorenz et al., 1996;

Sherrington, 1898, as cited in Lee et al., 2008) and may therefore not explain the activity patterns

we observed. Interestingly however, it has also been suggested that the size of dermatomes

depends on central communication via spinal levels (Denny-Brown et al., 1973; Denny-Brown &

Kirk, 1968; Kirk & Denny-Brown, 1970), emphasizing a dynamic view of cutaneous innervation.

Building on this, the large rostrocaudal patterns may also be explained by inter-segmental

nociceptive processing via e.g. propriospinal neurons (Flynn et al., 2011; Pierrot-Deseilligny &

Burke, 2012) or inter-segmental projection patterns of primary afferents (Kato et al., 2004; Pinto

et al., 2010).

Beyond the gray matter of the dorsal horn, the activation seemed to bleed into the subarachnoid

space, with strong peaks just outside of the spinal cord, where the large veins that drain the spinal

cord are located (Thron, 2016). Typically, the evaluation of BOLD effects in the spinal cord is

restricted to either a cord mask or a gray matter mask, meaning that the extent to which such

activation patterns are prevalent in the literature is uncertain. However, spinal fMRI studies

employing a hypercapnia challenge reported stronger signal changes at the edge of the cord and

in the CSF compared to inside the spinal cord (Barry et al., 2021; Cohen-Adad et al., 2010) and

several spinal fMRI studies employing painful heat stimuli also reported activations on the outer

edge of the cord or even overlapping into the CSF (Geuter & Buchel, 2013; Nash et al., 2013;

Oliva et al., 2022; Rempe et al., 2015; Sprenger et al., 2015; Sprenger, Stenmans, et al., 2018).

The bias towards draining veins is a well-known drawback of gradient-echo EPI and – considering

our results – it might be advisable in the future to try disentangling micro- and macrovascular

contributions to the spinal cord BOLD response, for instance by modeling the respective time

courses (Kay et al., 2020), leveraging differences in TE- (Markuerkiaga et al., 2021; Uludağ et al.,

2009) and phase dependencies (Stanley et al., 2021) to remove signal contributions from large

veins, or suppressing draining vein signal during data acquisition (Li et al., 2022). A first proof-of-

principle step might however be to obtain individual vasculature maps and investigate the

relationship between vascular anatomy and BOLD activation patterns in the spinal cord, a concept

that aligns with the initial approach of Cohen-Adad et al. (2010), whose findings highlight the

importance of vascular dynamics.

4.3 Test-retest reliability across consecutive days

The main objective of this study was to investigate the test-retest reliability of task-based spinal

cord BOLD responses across two consecutive days. Test-retest reliability describes to what extent

repeated measurements yield similar results, given that the underlying true value has not changed

(Lavrakas, 2008) and this also applies to inter-individual variation, with consistent differences

between participants indicating good reliability. Considering that many factors contribute to the

processing of pain (Bushnell et al., 2013; Heinricher & Fields, 2013), we also collected verbal

ratings as well as peripheral physiological responses in response to painful stimulation – apart

from offering a different window on the reliability of pain processing, these measures also served

as controls against which we could compare the reliability of the BOLD responses. We observed

that verbal ratings exhibited high reliability, a finding that has also been reported in previous

studies (Letzen, 2014; Quiton & Greenspan, 2008; Upadhyay, 2015), though it is unclear to what

extent this reflects the stability of actual perceptual differences or might be driven by biases in

reporting pain or differences in interpreting the rating scale. Peripheral physiological measures

also mostly showed high reliability, providing complementary evidence that participants can

indeed be distinguished reliably based on their peripheral physiological response to pain (the low

reliability of pupil dilation is likely due to noisy data on account of the non-ideal setup for eye-

tracking with the 64-channel coil employed here). Together these data provide a solid foundation

for investigating the reliability of spinal cord BOLD responses, as they indicate a generally high

reliability of supra-spinal measures of pain processing.

When looking at the test-retest reliability of spinal cord BOLD responses across days, we observed

that the reliability of the response amplitudes in the region of interest (left dorsal horn of segment

C6) was consistently in the poor range, similar to results obtained by Weber and colleagues when

investigating within-day test-retest reliability of spinal cord BOLD responses to heat-pain

stimulation (Weber et al., 2016a). One could argue that two of our chosen metrics (ROI mean and

peak) are suboptimal for assessing reliability, as the former included many non-responsive voxels

and the latter may merely constitute an outlier (given the tSNR of the data). However, the more

constrained approach of using the top 10% resulted in very similar reliability and higher reliability

values have been reported when investigating BOLD responses to painful stimulation using similar

approaches in the brain (Bi, 2021; Gay et al., 2015; Letzen, 2014; Quiton et al., 2014; Upadhyay,

2015). While the majority of these studies used stimuli longer than 10s (Gay et al., 2015; Quiton

et al., 2014; Upadhyay, 2015), Letzen et al. (2014) reported good test-retest reliability for 4s

contact heat stimuli and lower trial as well as participant numbers, similarly to Bi et al. (2021), who

observed fair to moderate test-retest reliability when using 4ms radiant heat-laser stimuli on

roughly half the number of participants compared to our study, albeit with slightly higher trial

numbers. Partly, these differences may be explained by the larger tSNR typically achieved in brain

compared to spinal cord fMRI as well as the lower spatial resolution employed in these studies,

which further increases tSNR.

Interestingly, an extended mask that covered the venous plexus surrounding the spinal cord yielded good reliability for the top 10% of the parameter estimates. One possible interpretation of this finding is that spinal cord BOLD response amplitudes could indeed be a reliable measure, but with the employed gradient-echo EPI acquisition's sensitivity to macrovascular responses (Bandettini et al., 1994; Duong et al., 2003; Gati et al., 1997; Uludağ et al., 2009), the actual response peak might be shifted from the gray matter towards the draining veins – in brain fMRI such differences would not be immediately noticeable, considering the typically lower spatial resolution, potentially causing signals from veins and gray matter to blend within individual voxels. Furthermore, the brain's anatomical structure, where large draining vessels often lie directly on top of the cortical gray matter, contrasts with the spinal cord's architecture, in which these larger draining vessels are located outside the white matter, surrounding the cord (Duvernoy, 1999; Gray, 2021).

In addition to response amplitudes, we also investigated the spatial consistency of the response pattern, i.e., if active voxels overlapped across days. The group level results showed a significant response in the target region for each session separately, but, the activation patterns did not overlap between days – while the location on the dorsal-ventral dimension remained similar, the patterns differed rostro-caudally within the same segment. This was paralleled on the individual level, where only few participants had overlapping responses in the area of interest, leading to very low dice coefficients at either level. It is noteworthy that a higher-than-expected amount of spatial variability in the z-direction within participants upon thermal stimulation has also been reported in a recent within-day design by Seifert and colleagues (2023). It is currently unclear if this is an indicator of large variability in spinal nociceptive processing, or the result of low tSNR (due to residual noise), an insufficient number of trials or a low stimulus intensity (Upadhyay, 2015). A partial answer was given by a post-hoc analysis where we increased the amount of data by averaging over multiple runs per session: here, the spatial overlap on the group level improved substantially, yet the reliability of the response amplitude did not improve.

There are several factors specific to across-day set-ups that could have negatively impacted the reliability of both response amplitudes and spatial patterns. One such factor is the quality of the spatial normalization, since differences thereof between days could have adverse effects on reliability (especially for a small structure such as the DH), yet this did not receive support by our analyses based on EPI-template Dice coefficients. Further inconsistencies between sessions may have been caused for example by differences in the positioning of the participants in the scanner, resulting in a different tilt of the head and neck, which is supported by the moderate correlation between differences in BOLD parameter estimates and the angulation of the slice stack relative

to the static magnetic field (as a proxy for neck positioning). Due to the anatomical organization of

the draining veins, the resulting different curvature / orientation of the neck and thus spinal cord

across days may have impacted the relative contribution of the longitudinal and radial veins to the

overall signal (Giove et al., 2004; Viessmann et al., 2019). It is also possible that the general

physiological state of the participants produced some variation the fMRI responses (Dubois &

Adolphs, 2016), but except for a slight increase in heart rate from day one to day two, all other

markers of physiological arousal remained the same. In one post-hoc analysis, we tried to partially

address these limitations by computing the within-run reliability using odd/even trials as well as

the first vs. second half of trials, where no re-positioning in the scanner occurred and spatial

normalization was equal. Interestingly, odd-even reliability of the BOLD fMRI results was in the

good range, while early-late reliability was still poor. The main difference between these two

assessments is the temporal distance between the trials, which was greater for early-late

reliability. This indicates that the heat-pain evoked activations display considerable variability

within a single run, potentially due to mechanisms physiological mechanisms such as adaptation,

habituation and sensitization (Greffrath et al., 2007; Hollins et al., 2011; Latremoliere & Woolf,

2009) as well as potential technical issues such as scanner drift. While it is unlikely that the same

mechanisms account for across-day variability, it indicates that factors beyond positioning and

spatial normalization can contribute to low reliability.

It is important to point out that a systematic comparison of our results with reliability estimates

obtained by resting-state spinal cord fMRI studies is unfortunately not possible, as these studies

not only varied vastly in their sample sizes (N = 1 to N = 45) but consistently used a within-day

design (Barry et al., 2016; Hu et al., 2018; Kaptan et al., 2023; Kong et al., 2014; Liu et al., 2016;

San Emeterio Nateras et al., 2016), thus not encountering the issues of across-day measurements

that might bring about low reliability. A notable exception is a recent study by Kowalczyk and

colleagues (2024), where a between-day design also resulted in mostly 'poor' voxelwise ICCs,

though the spatial patterns of connectivity showed near-perfect agreement.

4.4 General considerations on reliability

To put our observation of mostly low reliability of spinal cord BOLD responses into a larger context,

it is important to mention that a recent meta-analysis investigating univariate BOLD responses in

the brain to several common tasks from various domains also observed generally low reliability

(Elliott et al., 2020). Several ways to improve the reliability of fMRI have been discussed (Elliott et

al., 2021; Kragel et al., 2021), such as multivariate analysis (Gianaros et al., 2020; Han et al.,

2022), modeling stable variability, and the aggregation of more data (Elliott et al., 2021), all of
which might be applicable in the context of spinal cord fMRI as well.

A further aspect deserving discussion is our quantification of reliability, which was carried out using
the intra-class correlation coefficient ICC(3,1) (Shrout & Fleiss, 1979), a common measure for
test-retest reliability of fMRI data (Caceres et al., 2009; Elliott et al., 2020; Noble et al., 2021). The
ICC is a useful metric to investigate inter-individual differences, since it quantifies to what extent
participants can be "re-identified" across repeated measurements by means of the stability of their
rating in relation to that of other participants (Brandmaier et al., 2018; Hedge et al., 2018; Liljequist
et al., 2019). In order to obtain a high ICC, the variation between participants should be large, and
the variation within participants as well as the general measurement error should be small.
However, traditional univariate analyses of BOLD responses via the GLM – as also employed here
– are designed to minimize between-participant variability in order to gain a robust group-level
response (Fröhner et al., 2019; Hedge et al., 2018). Given the possible sources of noise discussed
in this study and elsewhere (Eippert, Kong, Jenkinson, et al., 2017; Kinany, Pirondini, Micera, et
al., 2022; Summers et al., 2014), minimizing the measurement noise holds the potential to both
improve reliability and optimize main effects on the group level.

1002

4.5 Limitations
There are several limitations of this work that need to be considered. First, the BOLD responses
elicited by 1s contact heat stimuli may exhibit lower reliability compared to those from longer
stimulus durations in a block design, which are typically more effective in detecting effects and
could yield more robust activation patterns (Bennett & Miller, 2013). The limited number of trials
further constrains our assessment of test-retest reliability, potentially making it more restrictive
than studies using more powerful experimental designs; here it is also important to mention the
low degrees of freedom of our time-series (considering that only 160 volumes were acquired per
run and that extensive denoising was carried out). Second, an assessment of the test-retest
reliability of tSNR values – as possible for example via a short resting-state acquisition – in the left
dorsal horn across different days would have provided valuable insights into the consistency of
the fMRI signal quality and should be considered in future studies. Third, one might argue that
instead of delivering stimuli with the same temperature on both days, we could have instead
matched stimulus intensity across days based on subjectively perceived intensity (to account for
confounds that might differ across days). Fourth, while the use of MP-PCA resulted in a substantial
tSNR increase (without a strong spatial smoothness penalty), future studies might look in more
detail at potential violations of underlying assumptions as well as artificial activation spreading,

which has recently been observed under certain conditions (Fernandes et al., 2023). Finally, we

might have achieved an increased across-day reliability by minimizing variability in participant

position (and thus also spatial inaccuracies), for example by using personalized casts (Power et

al., 2019).

1024

1025   4.6 Conclusion

We observed that heat pain stimuli as short as 1s can evoke a robust BOLD response in the

ipsilateral dorsal horn of the relevant spinal cord segment, making such stimuli suitable for use in

cognitive neuroscience experiments that require variable trial designs and large numbers of trials.

Although autonomic and subjective indicators of pain processing showed mostly good-to-excellent

reliability, BOLD response patterns varied strongly within participants, resulting in poor test-retest

reliability in the target region. Interestingly, using an extended analysis region including the

draining veins improved reliability across days, suggesting that future studies should aim to

disentangle macro- and microvascular contributions to the spatial response profile. Our results

indicate that further improvements in data acquisition and analysis techniques are necessary

before event-related spinal cord fMRI can be reliably employed in longitudinal designs or clinical

settings. To facilitate such endeavours, all data and code of this study are publicly available, thus

allowing others to develop and improve pre-processing and analysis strategies to overcome

current limitations.

1039

1040

1041   **Data and code availability**: The underlying data and code are currently only accessible to

1042   reviewers, but will be made openly available upon publication via OpenNeuro and GitHub,

1043   respectively.

1044

1045   **Ethics:** All participants gave written informed consent. The study was approved by the Ethics

1046   Committee of the Medical Faculty of the University of Leipzig.

1047

Author contributions: Author contributions are listed alphabetically according to CRediT

taxonomy (https://credit.niso.org).

Conceptualization: AD, FE

Data curation: AD

Formal analysis: AD, FE, UH

Funding acquisition: FE

Investigation: AD

Methodology: AD, FE, JL, RM

Project administration: AD, FE

Resources: JF, JL, RM

Software: AD, FE, UH, MK

Supervision: FE, TM, NW

Visualization: AD

Writing – original draft: AD, FE

Writing – review & editing: JB, AD, FE, JF, UH, MK, JL, RM, TM, NW

1063

1064   **Acknowledgements**:   We would like to thank all volunteers who participated in this study.
1065   Additionally, we want to thank everyone who assisted in data acquisition as well as Lisa-Marie
1066   Pohle for her help in the randomization of trials and conditions. This manuscript will be part of a
1067   doctoral thesis.

1068

Funding information: FE received funding from the Max Planck Society and the European
Research Council (under the European Union's Horizon 2020 research and Innovation Program;
grant agreement No 758974). MK was supported by a grant from the National Institute of Health
(Grant Number R01NS109450). JB received funding from the UK Medical Research Council
(MR/N026969/1). NW received funding from the European Research Council under the European
Union's Seventh Framework Programme (FP7/2007- 2013, ERC grant agreement No 616905),
the European Union's Horizon 2020 research and innovation program (under the grant agreement
No 681094) and the BMBF (01EW1711A & B) in the framework of ERA-NET NEURON.

1077

1078   **Competing interest**: The Max Planck Institute for Human Cognitive and Brain Sciences has an
1079   institutional research agreement with Siemens Healthcare. Nikolaus Weiskopf holds a patent on

1080   acquisition of MRI data during spoiler gradients (US 10,401,453 B2). Nikolaus Weiskopf was a

1081   speaker at an event organized by Siemens Healthcare and was reimbursed for the travel

1082   expenses.

1083

References

Ades-Aron, B., Lemberskiy, G., Veraart, J., Golfinos, J., Fieremans, E., Novikov, D. S., & Shepherd, T. (2021). Improved Task-based Functional MRI Language Mapping in Patients with Brain Tumors through Marchenko-Pastur Principal Component Analysis Denoising. *Radiology*, *298*(2), 365–373. https://doi.org/10.1148/radiol.2020200822

Ahuja, C. S., Wilson, J. R., Nori, S., Kotter, M. R. N., Druschel, C., Curt, A., & Fehlings, M. G. (2017). Traumatic spinal cord injury. *Nature Reviews Disease Primers*, *3*(1), 17018. https://doi.org/10.1038/nrdp.2017.18

Atlas, L. Y., & Wager, T. D. (2012). How expectations shape pain. *Neuroscience Letters*, *520*(2), 140–148. https://doi.org/10.1016/j.neulet.2012.03.039

Bach, D. R., Flandin, G., Friston, K. J., & Dolan, R. J. (2009). Time-series analysis for rapid event-related skin conductance responses. *Journal of Neuroscience Methods*, *184*(2), 224–234. https://doi.org/10.1016/j.jneumeth.2009.08.005

Bach, D. R., Friston, K. J., & Dolan, R. J. (2010). Analytic measures for quantification of arousal from spontaneous skin conductance fluctuations. *International Journal of Psychophysiology*, *76*(1), 52–55. https://doi.org/10.1016/j.ijpsycho.2010.01.011

Bach, D. R., Friston, K. J., & Dolan, R. J. (2013). An improved algorithm for model-based analysis of evoked skin conductance responses. *Biological Psychology*, *94*(3), 490–497. https://doi.org/10.1016/j.biopsycho.2013.09.010

Bandettini, P. A., Wong, E. C., Jesmanowicz, A., Hinks, R. S., & Hyde, J. S. (1994). Spin-echo and gradient-echo epi of human brain activation using bold contrast: A comparative study at 1.5 T. *NMR in Biomedicine*, *7*(1–2), 12–20. https://doi.org/10.1002/nbm.1940070104

Bannatyne, B. A. (2006). Differential Projections of Excitatory and Inhibitory Dorsal Horn Interneurons Relaying Information from Group II Muscle Afferents in the Cat Spinal Cord. *Journal of Neuroscience*, *26*(11), 2871–2880. https://doi.org/10.1523/JNEUROSCI.5172-05.2006

Barry, R. L., Conrad, B. N., Maki, S., Watchmaker, J. M., McKeithan, L. J., Box, B. A., Weinberg, Q. R., Smith, S. A., & Gore, J. C. (2021a). Multi-shot acquisitions for stimulus-evoked spinal cord BOLD fMRI. *Magnetic Resonance in Medicine*, *85*(4), 2016–2026. https://doi.org/10.1002/mrm.28570

Barry, R. L., Conrad, B. N., Maki, S., Watchmaker, J. M., McKeithan, L. J., Box, B. A., Weinberg, Q. R., Smith, S. A., & Gore, J. C. (2021b). Multi-shot acquisitions for stimulus-evoked spinal cord BOLD fMRI. *Magnetic Resonance in Medicine*, *85*(4), 2016–2026. https://doi.org/10.1002/mrm.28570

Barry, R. L., Rogers, B. P., Conrad, B. N., Smith, S. A., & Gore, J. C. (2016). Reproducibility of resting state spinal cord networks in healthy volunteers at 7 Tesla. *NeuroImage*, *133*, 31–40. https://doi.org/10.1016/j.neuroimage.2016.02.058

Barry, R. L., & Smith, S. A. (2019). Measurement of T_2* in the human spinal cord at 3T. *Magnetic Resonance in Medicine*, *82*(2), 743–748. https://doi.org/10.1002/mrm.27755

Barry, R. L., Smith, S. A., Dula, A. N., & Gore, J. C. (2014). Resting state functional connectivity in the human spinal cord. *ELife*, *3*. https://doi.org/10.7554/eLife.02812

Bennett, C. M., & Miller, M. B. (2013). fMRI reliability: Influences of task and experimental design. *Cognitive, Affective, & Behavioral Neuroscience*, *13*(4), 690–702. https://doi.org/10.3758/s13415-013-0195-1

Bi, Y. (2021). Test–retest reliability of laser evoked pain perception and fMRI BOLD responses. *Scientific Reports*, 9.

Bi, Y., Hou, X., Zhong, J., & Hu, L. (2021). Test–retest reliability of laser evoked pain perception and fMRI BOLD responses. *Scientific Reports*, *11*(1), 1322. https://doi.org/10.1038/s41598-020-79196-z

Bosma, R. L., & Stroman, P. W. (2014). Assessment of data acquisition parameters, and analysis techniques for noise reduction in spinal cord fMRI data. *Magnetic Resonance Imaging*, *32*(5), 473–481. https://doi.org/10.1016/j.mri.2014.01.007

Bosma, R. L., & Stroman, P. W. (2015). Spinal cord response to stepwise and block presentation of thermal stimuli: A functional MRI study: Assessment of Spinal fMRI Paradigms. *Journal of Magnetic Resonance Imaging*, *41*(5), 1318–1325. https://doi.org/10.1002/jmri.24656

Boucsein, W. (2012). *Electrodermal activity* (2nd ed). Springer.

Bouwman, C. J. C., Wilmink, J. T., Mess, W. H., & Backes, W. H. (2008). Spinal cord functional MRI at 3 T: Gradient echo echo-planar imaging versus turbo spin echo. *NeuroImage*, *43*(2), 288–296. https://doi.org/10.1016/j.neuroimage.2008.07.024

Brandmaier, A. M., Wenger, E., Bodammer, N. C., Kühn, S., Raz, N., & Lindenberger, U. (2018). Assessing reliability in neuroimaging research through intra-class effect decomposition (ICED). *ELife*, *7*, e35718. https://doi.org/10.7554/eLife.35718

Brooks, J. C. W., Beckmann, C. F., Miller, K. L., Wise, R. G., Porro, C. A., Tracey, I., & Jenkinson, M. (2008). Physiological noise modelling for spinal functional magnetic resonance imaging studies. *NeuroImage*, *39*(2), 680–692. https://doi.org/10.1016/j.neuroimage.2007.09.018

Brooks, J. C. W., Kong, Y., Lee, M. C., Warnaby, C. E., Wanigasekera, V., Jenkinson, M., & Tracey, I. (2012). Stimulus Site and Modality Dependence of Functional Activity within the Human Spinal Cord. *Journal of Neuroscience*, *32*(18), 6231–6239. https://doi.org/10.1523/JNEUROSCI.2543-11.2012

Bushnell, M. C., Čeko, M., & Low, L. A. (2013). Cognitive and emotional control of pain and its disruption in chronic pain. *Nature Reviews Neuroscience*, *14*(7), 502–511. https://doi.org/10.1038/nrn3516

Caceres, A., Hall, D. L., Zelaya, F. O., Williams, S. C. R., & Mehta, M. A. (2009a). Measuring fMRI reliability with the intra-class correlation coefficient. *NeuroImage*, *45*(3), 758–768. https://doi.org/10.1016/j.neuroimage.2008.12.035

Caceres, A., Hall, D. L., Zelaya, F. O., Williams, S. C. R., & Mehta, M. A. (2009b). Measuring fMRI reliability with the intra-class correlation coefficient. *NeuroImage*, *45*(3), 758–768. https://doi.org/10.1016/j.neuroimage.2008.12.035

Cadotte, D. W., Cadotte, A., Cohen-Adad, J., Fleet, D., Livne, M., Wilson, J. R., Mikulis, D., Nugaeva, N., & Fehlings, M. G. (2015). Characterizing the Location of Spinal and Vertebral Levels in the Human Cervical Spinal Cord. *American Journal of Neuroradiology*, *36*(4), 803–810. https://doi.org/10.3174/ajnr.A4192

Cahill, C. M., & Stroman, P. W. (2011). Mapping of neural activity produced by thermal pain in the healthy human spinal cord and brain stem: A functional magnetic resonance imaging study. *Magnetic Resonance Imaging*, *29*(3), 342–352. https://doi.org/10.1016/j.mri.2010.10.007

Cicchetti, D. V. (1994). Guidelines, criteria, and rules of thumb for evaluating normed and standardized assessment instruments in psychology. *Psychological Assessment*, *6*(4), 284–290. https://doi.org/10.1037/1040-3590.6.4.284

Coghill, R. C., Mayer, D. J., & Price, D. D. (1993). The roles of spatial recruitment and discharge frequency in spinal cord coding of pain: A combined electrophysiological and imaging investigation. *Pain*, *53*(3), 295–309. https://doi.org/10.1016/0304-3959(93)90226-F

Coghill, R. C., Price, D. D., Hayes, R. L., & Mayer, D. J. (1991). Spatial distribution of nociceptive processing in the rat spinal cord. *Journal of Neurophysiology*, *65*(1), 133–140. https://doi.org/10.1152/jn.1991.65.1.133

Cohen-Adad, J. (2017). Functional Magnetic Resonance Imaging of the Spinal Cord: Current Status and Future Developments. *Seminars in Ultrasound, CT and MRI*, *38*(2), 176–186. https://doi.org/10.1053/j.sult.2016.07.007

Cohen-Adad, J., Alonso-Ortiz, E., Abramovic, M., Arneitz, C., Atcheson, N., Barlow, L., Barry, R. L., Barth, M., Battiston, M., Büchel, C., Budde, M., Callot, V., Combes, A. J. E., De Leener, B., Descoteaux, M., de Sousa, P. L., Dostál, M., Doyon, J., Dvorak, A., … Xu, J. (2021). Generic acquisition protocol for quantitative MRI of the spinal cord. *Nature Protocols*, *16*(10), 4611–4632. https://doi.org/10.1038/s41596-021-00588-0

Cohen-Adad, J., Gauthier, C. J., Brooks, J. C. W., Slessarev, M., Han, J., Fisher, J. A., Rossignol, S., & Hoge, R. D. (2010). BOLD signal responses to controlled hypercapnia in human spinal cord. *NeuroImage*, *50*(3), 1074–1084. https://doi.org/10.1016/j.neuroimage.2009.12.122

Cohen-Adad, J., Mareyam, A., Keil, B., Polimeni, J. R., & Wald, L. L. (2011). 32-Channel RF coil optimized for brain and cervical spinal cord at 3 T: 32ch Head/c-Spine Coil at 3 T. *Magnetic Resonance in Medicine*, *66*(4), 1198–1208. https://doi.org/10.1002/mrm.22906

Cole, J. H., & Franke, K. (2017). Predicting Age Using Neuroimaging: Innovative Brain Ageing Biomarkers. *Trends in Neurosciences*, *40*(12), 681–690. https://doi.org/10.1016/j.tins.2017.10.001

Colloca, L., Ludman, T., Bouhassira, D., Baron, R., Dickenson, A. H., Yarnitsky, D., Freeman, R., Truini, A., Attal, N., Finnerup, N. B., Eccleston, C., Kalso, E., Bennett, D. L., Dworkin, R. H., & Raja, S. N. (2017). Neuropathic pain. *Nature Reviews Disease Primers*, *3*(1), 17002. https://doi.org/10.1038/nrdp.2017.2

Conrad, B. N., Barry, R. L., Rogers, B. P., Maki, S., Mishra, A., Thukral, S., Sriram, S., Bhatia, A., Pawate, S., Gore, J. C., & Smith, S. A. (2018). Multiple sclerosis lesions affect intrinsic functional connectivity of the spinal cord. *Brain*, *141*(6), 1650–1664. https://doi.org/10.1093/brain/awy083

Cowen, R., Stasiowska, M. K., Laycock, H., & Bantel, C. (2015). Assessing pain objectively: The use of physiological markers. *Anaesthesia*, *70*(7), 828–847. https://doi.org/10.1111/anae.13018

Culberson, J. L., Haines, D. E., Kimmel, D. L., & Brown, P. B. (1979). Contralateral projection of primary afferent fibers to mammalian spinal cord. *Experimental Neurology*, *64*(1), 83–97. https://doi.org/10.1016/0014-4886(79)90007-4

Davis, K. D., Aghaeepour, N., Ahn, A. H., Angst, M. S., Borsook, D., Brenton, A., Burczynski, M. E., Crean, C., Edwards, R., Gaudilliere, B., Hergenroeder, G. W., Iadarola, M. J., Iyengar, S., Jiang, Y., Kong, J.-T., Mackey, S., Saab, C. Y., Sang, C. N., Scholz, J., … Pelleymounter, M. A. (2020). Discovery and validation of biomarkers to aid the development of safe and effective pain therapeutics: Challenges and opportunities. *Nature Reviews Neurology*, *16*(7), 381–400. https://doi.org/10.1038/s41582-020-0362-2

De Leener, B., Fonov, V. S., Collins, D. L., Callot, V., Stikov, N., & Cohen-Adad, J. (2018). PAM50: Unbiased multimodal template of the brainstem and spinal cord aligned with the ICBM152 space. *NeuroImage*, *165*, 170–179. https://doi.org/10.1016/j.neuroimage.2017.10.041

De Leener, B., Lévy, S., Dupont, S. M., Fonov, V. S., Stikov, N., Louis Collins, D., Callot, V., & Cohen-Adad, J. (2017). SCT: Spinal Cord Toolbox, an open-source software for processing spinal cord MRI data. *NeuroImage*, *145*, 24–43. https://doi.org/10.1016/j.neuroimage.2016.10.009

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, *134*(1), 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Denny-Brown, D., & Kirk, E. (1968). Hyperesthesia from spinal and root lesions. *Transactions of the American Neurological Association*, *93*, 116–120.

Denny-Brown, D., Kirk, E. J., & Yanagisawa, N. (1973). The tract of Lissauer in relation to sensory transmission in the dorsal horn of spinal cord in the macaque monkey. *The Journal of Comparative Neurology*, *151*(2), 175–199. https://doi.org/10.1002/cne.901510206

D'Esposito, M., Zarahn, E., & Aguirre, G. K. (1999). Event-related functional MRI: Implications for cognitive psychology. *Psychological Bulletin*, *125*(1), 155–164. https://doi.org/10.1037/0033-2909.125.1.155

Diao, Y., Yin, T., Gruetter, R., & Jelescu, I. O. (2021). PIRACY: An Optimized Pipeline for Functional Connectivity Analysis in the Rat Brain. *Frontiers in Neuroscience*, *15*, 602170. https://doi.org/10.3389/fnins.2021.602170

D'Mello, R., & Dickenson, A. H. (2008). Spinal cord mechanisms of pain. *British Journal of Anaesthesia*, *101*(1), 8–16. https://doi.org/10.1093/bja/aen088

Dubois, J., & Adolphs, R. (2016). Building a Science of Individual Differences from fMRI. *Trends in Cognitive Sciences*, *20*(6), 425–443. https://doi.org/10.1016/j.tics.2016.03.014

Duong, T. Q., Yacoub, E., Adriany, G., Hu, X., U??urbil, K., & Kim, S.-G. (2003). Microvascular BOLD contribution at 4 and 7 T in the human brain: Gradient-echo and spin-echo fMRI with suppression of blood effects. *Magnetic Resonance in Medicine*, *49*(6), 1019–1027. https://doi.org/10.1002/mrm.10472

Duvernoy, H. M. (1999). *The Human Brain: Surface, Three-Dimensional Sectional Anatomy with MRI, and Blood Supply* (2nd ed.). Springer Vienna.

Eippert, F., Finsterbusch, J., Bingel, U., & Buchel, C. (2009). Direct Evidence for Spinal Cord Involvement
in Placebo Analgesia. *Science*, *326*(5951), 404–404. https://doi.org/10.1126/science.1180142

Eippert, F., Kong, Y., Jenkinson, M., Tracey, I., & Brooks, J. C. W. (2017). Denoising spinal cord fMRI data:
Approaches to acquisition and analysis. *NeuroImage*, *154*, 255–266.
https://doi.org/10.1016/j.neuroimage.2016.09.065

Eippert, F., Kong, Y., Winkler, A. M., Andersson, J. L., Finsterbusch, J., Büchel, C., Brooks, J. C. W., &
Tracey, I. (2017). Investigating resting-state functional connectivity in the cervical spinal cord at 3 T.
NeuroImage, *147*, 589–601. https://doi.org/10.1016/j.neuroimage.2016.12.072

Elliott, M. L., Knodt, A. R., & Hariri, A. R. (2021). Striving toward translation: Strategies for reliable fMRI
measurement. *Trends in Cognitive Sciences*, *25*(9), 776–787. https://doi.org/10.1016/j.tics.2021.05.008

Elliott, M. L., Knodt, A. R., Ireland, D., Morris, M. L., Poulton, R., Ramrakha, S., Sison, M. L., Moffitt, T. E.,
Caspi, A., & Hariri, A. R. (2020). What Is the Test-Retest Reliability of Common Task-Functional MRI
Measures? New Empirical Evidence and a Meta-Analysis. *Psychological Science*, *31*(7), 792–806.
https://doi.org/10.1177/0956797620916786

Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: A flexible statistical power analysis
program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, *39*(2), 175–191.
https://doi.org/10.3758/BF03193146

Fernandes, F. F., Olesen, J. L., Jespersen, S. N., & Shemesh, N. (2023). MP-PCA denoising of fMRI time-
series data can lead to artificial activation "spreading". NeuroImage, 273, 120118.
https://doi.org/10.1016/j.neuroimage.2023.120118

Filippi, M., & Rocca, M. A. (2013). Present and future of fMRI in multiple sclerosis. *Expert Review of
Neurotherapeutics*, *13*(sup2), 27–31. https://doi.org/10.1586/14737175.2013.865871

Finsterbusch, J., Eippert, F., & Büchel, C. (2012). Single, slice-specific z-shim gradient pulses improve T2*-
weighted imaging of the spinal cord. *NeuroImage*, *59*(3), 2307–2315.
https://doi.org/10.1016/j.neuroimage.2011.09.038

Finsterbusch, J., Sprenger, C., & Büchel, C. (2013). Combined T2*-weighted measurements of the human
brain and cervical spinal cord with a dynamic shim update. *NeuroImage*, *79*, 153–161.
https://doi.org/10.1016/j.neuroimage.2013.04.021

Fitzgerald, M. (1982). The contralateral input to the dorsal horn of the spinal cord in the decerebrate spinal
rat. *Brain Research*, *236*(2), 275–287. https://doi.org/10.1016/0006-8993(82)90714-4

Flynn, J. R., Graham, B. A., Galea, M. P., & Callister, R. J. (2011). The role of propriospinal interneurons in
recovery from spinal cord injury. *Neuropharmacology*, *60*(5), 809–822.
https://doi.org/10.1016/j.neuropharm.2011.01.016

Fox, M. D., Snyder, A. Z., Zacks, J. M., & Raichle, M. E. (2006). Coherent spontaneous activity accounts
for trial-to-trial variability in human evoked brain responses. *Nature Neuroscience*, *9*(1), 23–25.
https://doi.org/10.1038/nn1616

Fröhner, J. H., Teckentrup, V., Smolka, M. N., & Kroemer, N. B. (2019). Addressing the reliability fallacy in
fMRI: Similar group effects may arise from unreliable individual effects. *NeuroImage*, *195*, 174–189.
https://doi.org/10.1016/j.neuroimage.2019.03.053

Frostell, A., Hakim, R., Thelin, E. P., Mattsson, P., & Svensson, M. (2016). A review of the segmental
diameter of the healthy human spinal cord. *Frontiers in neurology, 7*, 230582.

Gati, J. S., Menon, R. S., U??urbil, K., & Rutt, B. K. (1997). Experimental determination of the BOLD field
strength dependence in vessels and tissue. *Magnetic Resonance in Medicine*, *38*(2), 296–302.
https://doi.org/10.1002/mrm.1910380220

Gay, C. W., Papuga, M. O., Bishop, M. D., & Dougherty, P. (2015). The frequency and reliability of cortical
activity using a novel strategy to present pressure pain stimulus over the lumbar spine. *Journal of
Neuroscience Methods*, *239*, 108–113. https://doi.org/10.1016/j.jneumeth.2014.10.010

Geuter, S., & Buchel, C. (2013). Facilitation of Pain in the Human Spinal Cord by Nocebo Treatment. *Journal
of Neuroscience*, *33*(34), 13784–13790. https://doi.org/10.1523/JNEUROSCI.2191-13.2013

Gianaros, P. J., Kraynak, T. E., Kuan, D. C.-H., Gross, J. J., McRae, K., Hariri, A. R., Manuck, S. B., Rasero, J., & Verstynen, T. D. (2020). Affective brain patterns as multivariate neural correlates of cardiovascular disease risk. *Social Cognitive and Affective Neuroscience*, *15*(10), 1034–1045. https://doi.org/10.1093/scan/nsaa050

Giove, F., Garreffa, G., Giulietti, G., Mangia, S., Colonnese, C., & Maraviglia, B. (2004). Issues about the fMRI of the human spinal cord. *Magnetic Resonance Imaging*, *22*(10), 1505–1516. https://doi.org/10.1016/j.mri.2004.10.015

Giulietti, G., Giove, F., Garreffa, G., Colonnese, C., Mangia, S., & Maraviglia, B. (2008). Characterization of the functional response in the human spinal cord: Impulse-response function and linearity. *NeuroImage*, *42*(2), 626–634. https://doi.org/10.1016/j.neuroimage.2008.05.006

Gray, H. (2021). *Gray's anatomy: The anatomical basis of clinical practice* (S. Standring, N. Anand, & R. Tunstall, Eds.; Forty-second edition). Elsevier.

Greffrath, W., Baumgärtner, U., & Treede, R.-D. (2007). Peripheral and central components of habituation of heat pain perception and evoked potentials in humans. *Pain*, *132*(3), 301–311. https://doi.org/10.1016/j.pain.2007.04.026

Gros, C., De Leener, B., Badji, A., Maranzano, J., Eden, D., Dupont, S. M., Talbott, J., Zhuoquiong, R., Liu, Y., Granberg, T., Ouellette, R., Tachibana, Y., Hori, M., Kamiya, K., Chougar, L., Stawiarz, L., Hillert, J., Bannier, E., Kerbrat, A., … Cohen-Adad, J. (2019). Automatic segmentation of the spinal cord and intramedullary multiple sclerosis lesions with convolutional neural networks. *NeuroImage*, *184*, 901–915. https://doi.org/10.1016/j.neuroimage.2018.09.081

Han, X., Ashar, Y. K., Kragel, P., Petre, B., Schelkun, V., Atlas, L. Y., Chang, L. J., Jepma, M., Koban, L., Losin, E. A. R., Roy, M., Woo, C.-W., & Wager, T. D. (2022). Effect sizes and test-retest reliability of the fMRI-based neurologic pain signature. *NeuroImage*, *247*, 118844. https://doi.org/10.1016/j.neuroimage.2021.118844

Harita, S., & Stroman, P. W. (2017). Confirmation of resting-state BOLD fluctuations in the human brainstem and spinal cord after identification and removal of physiological noise: Resting-State BOLD fMRI in the Human Brainstem and Spinal Cord. *Magnetic Resonance in Medicine*, *78*(6), 2149–2156. https://doi.org/10.1002/mrm.26606

Harrison, O. K., Guell, X., Klein-Flügge, M. C., & Barry, R. L. (2021). Structural and resting state functional connectivity beyond the cortex. *NeuroImage*, *240*, 118379. https://doi.org/10.1016/j.neuroimage.2021.118379

Hedge, C., Powell, G., & Sumner, P. (2018). The reliability paradox: Why robust cognitive tasks do not produce reliable individual differences. *Behavior Research Methods*, *50*(3), 1166–1186. https://doi.org/10.3758/s13428-017-0935-1

Heinricher, M. M., & Fields, H. L. (2013). Central nervous system mechanisms of pain modulation. In *Wall and Melzack's textbook of pain* (6th ed.).

Hoggarth, M. A., Wang, M. C., Hemmerling, K. J., Vigotsky, A. D., Smith, Z. A., Parrish, T. B., Weber, K. A., & Bright, M. G. (2022). Effects of variability in manually contoured spinal cord masks on fMRI co-registration and interpretation. *Frontiers in Neurology*, *13*, 907581. https://doi.org/10.3389/fneur.2022.907581

Hollins, M., Harper, D., & Maixner, W. (2011). Changes in pain from a repetitive thermal stimulus: The roles of adaptation and sensitization: *Pain*, *152*(7), 1583–1590. https://doi.org/10.1016/j.pain.2011.02.049

Hu, Y., Jin, R., Li, G., Luk, K. D., & Wu, Ed. X. (2018). Robust spinal cord resting-state fMRI using independent component analysis-based nuisance regression noise reduction. *Journal of Magnetic Resonance Imaging*, *48*(5), 1421–1431. https://doi.org/10.1002/jmri.26048

Islam, H., Law, C. S. W., Weber, K. A., Mackey, S. C., & Glover, G. H. (2019). Dynamic per slice shimming for simultaneous brain and spinal cord fMRI. *Magnetic Resonance in Medicine*, *81*(2), 825–838. https://doi.org/10.1002/mrm.27388

Kaptan, M., Horn, U., Vannesjo, S. J., Mildner, T., Weiskopf, N., Finsterbusch, J., Brooks, J. C. W., & Eippert, F. (2023). Reliability of resting-state functional connectivity in the human spinal cord: Assessing the impact of distinct noise sources. *NeuroImage*, *275*, 120152. https://doi.org/10.1016/j.neuroimage.2023.120152

Kaptan, M., Vannesjo, S. J., Mildner, T., Horn, U., Hartley-Davies, R., Oliva, V., Brooks, J. C. W., Weiskopf, N., Finsterbusch, J., & Eippert, F. (2022). Automated slice-specific z-shimming for functional magnetic resonance imaging of the human spinal cord. *Human Brain Mapping*, hbm.26018. https://doi.org/10.1002/hbm.26018

Kato, G., Furue, H., Katafuchi, T., Yasaka, T., Iwamoto, Y., & Yoshimura, M. (2004). Electrophysiological mapping of the nociceptive inputs to the substantia gelatinosa in rat horizontal spinal cord slices: Mapping of nociceptive inputs in rat substantia gelatinosa. *The Journal of Physiology*, *560*(1), 303–315. https://doi.org/10.1113/jphysiol.2004.068700

Kay, K., Jamison, K. W., Zhang, R.-Y., & Uğurbil, K. (2020). A temporal decomposition method for identifying venous effects in task-based fMRI. *Nature Methods*, *17*(10), 1033–1039. https://doi.org/10.1038/s41592-020-0941-6

Khalili-Mahani, N., Rombouts, S. A. R. B., van Osch, M. J. P., Duff, E. P., Carbonell, F., Nickerson, L. D., Becerra, L., Dahan, A., Evans, A. C., Soucy, J.-P., Wise, R., Zijdenbos, A. P., & van Gerven, J. M. (2017). Biomarkers, designs, and interpretations of resting-state fMRI in translational pharmacological research: A review of state-of-the-Art, challenges, and opportunities for studying brain chemistry. *Human Brain Mapping*, *38*(4), 2276–2325. https://doi.org/10.1002/hbm.23516

Kinany, N., Pirondini, E., Mattera, L., Martuzzi, R., Micera, S., & Van De Ville, D. (2022). Towards reliable spinal cord fMRI: Assessment of common imaging protocols. *NeuroImage*, *250*, 118964. https://doi.org/10.1016/j.neuroimage.2022.118964

Kinany, N., Pirondini, E., Micera, S., & Van De Ville, D. (2020). Dynamic Functional Connectivity of Resting-State Spinal Cord fMRI Reveals Fine-Grained Intrinsic Architecture. *Neuron*, *108*(3), 424-435.e4. https://doi.org/10.1016/j.neuron.2020.07.024

Kinany, N., Pirondini, E., Micera, S., & Van De Ville, D. (2022). Spinal Cord fMRI: A New Window into the Central Nervous System. *The Neuroscientist*, 107385842211018. https://doi.org/10.1177/10738584221101827

Kirk, E. J., & Denny-Brown, D. (1970). Functional variation in dermatomes in the macaque monkey following dorsal root lesions. *The Journal of Comparative Neurology*, *139*(3), 307–320. https://doi.org/10.1002/cne.901390304

Kolesar, T. A., Fiest, K. M., Smith, S. D., & Kornelsen, J. (2015). Assessing Nociception by Fmri of the Human Spinal Cord: A Systematic Review. *Magnetic Resonance Insights*, *8s1*, MRI.S23556. https://doi.org/10.4137/MRI.S23556

Kong, Y., Eippert, F., Beckmann, C. F., Andersson, J., Finsterbusch, J., Büchel, C., Tracey, I., & Brooks, J. C. W. (2014). Intrinsically organized resting state networks in the human spinal cord. *Proceedings of the National Academy of Sciences*, *111*(50), 18067–18072. https://doi.org/10.1073/pnas.1414293111

Kong, Y., Jenkinson, M., Andersson, J., Tracey, I., & Brooks, J. C. W. (2012). Assessment of physiological noise modelling methods for functional imaging of the spinal cord. *NeuroImage*, *60*(2), 1538–1549. https://doi.org/10.1016/j.neuroimage.2011.11.077

Kowalczyk, O. S., Medina, S., Tsivaka, D., McMahon, S. B., Williams, S. C. R., Brooks, J. C. W., Lythgoe, D. J., & Howard, M. A. (2024). Spinal fMRI demonstrates segmental organisation of functionally connected networks in the cervical spinal cord: A test–retest reliability study. Human Brain Mapping, 45(2), e26600. https://doi.org/10.1002/hbm.26600

Kragel, P. A., Han, X., Kraynak, T. E., Gianaros, P. J., & Wager, T. D. (2021). Functional MRI Can Be Highly Reliable, but It Depends on What You Measure: A Commentary on Elliott et al. (2020). *Psychological Science*, *32*(4), 622–626. https://doi.org/10.1177/0956797621989730

Kuner, R., & Flor, H. (2017). Structural plasticity and reorganisation in chronic pain. *Nature Reviews Neuroscience*, *18*(1), 20–30. https://doi.org/10.1038/nrn.2016.162

Kyle, B. N., & McNeil, D. W. (2014). Autonomic Arousal And Experimentally Induced Pain: A Critical Review of the Literature. *Pain Research and Management*, *19*(3), 159–167. https://doi.org/10.1155/2014/536859

Landelle, C., Lungu, O., Vahdat, S., Kavounoudias, A., Marchand-Pauvert, V., De Leener, B., & Doyon, J. (2021). Investigating the human spinal sensorimotor pathways through functional magnetic resonance imaging. *NeuroImage*, *245*, 118684. https://doi.org/10.1016/j.neuroimage.2021.118684

Latremoliere, A., & Woolf, C. J. (2009). Central Sensitization: A Generator of Pain Hypersensitivity by Central Neural Plasticity. *The Journal of Pain*, *10*(9), 895–926. https://doi.org/10.1016/j.jpain.2009.06.012

Lavrakas, P. J. (2008). *Encyclopedia of Survey Research Methods*. SAGE Publications.

Lee, I.-S., Necka, E. A., & Atlas, L. Y. (2020). Distinguishing pain from nociception, salience, and arousal: How autonomic nervous system activity can improve neuroimaging tests of specificity. *NeuroImage*, *204*, 116254. https://doi.org/10.1016/j.neuroimage.2019.116254

Lee, M. W. L., McPhee, R. W., & Stringer, M. D. (2008). An evidence-based approach to human dermatomes. *Clinical Anatomy*, *21*(5), 363–373. https://doi.org/10.1002/ca.20636

Leone, C., Di Stefano, G., Di Pietro, G., Bloms-Funke, P., Boesl, I., Caspani, O., Chapman, S. C., Finnerup, N. B., Garcia-Larrea, L., Li, T., Goetz, M., Mouraux, A., Pelz, B., Pogatzki-Zahn, E., Schilder, A., Schnetter, E., Schubart, K., Tracey, I., Troconiz, I. F., … Treede, R.-D. (2022). IMI2-PainCare-BioPain-RCT2 protocol: A randomized, double-blind, placebo-controlled, crossover, multicenter trial in healthy subjects to investigate the effects of lacosamide, pregabalin, and tapentadol on biomarkers of pain processing observed by non-invasive neurophysiological measurements of human spinal cord and brainstem activity. *Trials*, *23*(1), 739. https://doi.org/10.1186/s13063-022-06431-5

Letzen, J. E. (2014). *Test-Retest Reliability of Pain-Related Brain Activity in Healthy Controls Undergoing Experimental Thermal Pain*. 7.

Letzen, J. E., Sevel, L. S., Gay, C. W., O'Shea, A. M., Craggs, J. G., Price, D. D., & Robinson, M. E. (2014). Test-Retest Reliability of Pain-Related Brain Activity in Healthy Controls Undergoing Experimental Thermal Pain. *The Journal of Pain*, *15*(10), 1008–1014. https://doi.org/10.1016/j.jpain.2014.06.011

Li, L., Law, C., Marrett, S., Chai, Y., Huber, L., Jezzard, P., & Bandettini, P. (2022). Quantification of cerebral blood volume changes caused by visual stimulation at 3 T using DANTE-prepared dual-echo EPI. *Magnetic Resonance in Medicine*, *87*(4), 1846–1862. https://doi.org/10.1002/mrm.29099

Light, A. R., & Perl, E. R. (1979). Reexamination of the dorsal root projection to the spinal dorsal horn including observations on the differential termination of coarse and fine fibers. *Journal of Comparative Neurology*, *186*(2), 117–131. https://doi.org/10.1002/cne.901860202

Liljequist, D., Elfving, B., & Skavberg Roaldsen, K. (2019). Intraclass correlation – A discussion and demonstration of basic features. *PLOS ONE*, *14*(7), e0219854. https://doi.org/10.1371/journal.pone.0219854

Liu, X., Zhou, F., Li, X., Qian, W., Cui, J., Zhou, I. Y., Luk, K. D. K., Wu, Ed. X., & Hu, Y. (2016). Organization of the intrinsic functional network in the cervical spinal cord: A resting state functional MRI study. *Neuroscience*, *336*, 30–38. https://doi.org/10.1016/j.neuroscience.2016.08.042

Lopez-Rios, N., Gilbert, K. M., Papp, D., Cereza, G., Foias, A., Rangaprakash, D., May, M. W., Guerin, B., Wald, L. L., Keil, B., Stockmann, J. P., Barry, R. L., & Cohen-Adad, J. (2023). An 8-channel Tx dipole and 20-channel Rx loop coil array for MRI of the cervical spinal cord at 7 Tesla. *NMR in Biomedicine*, *36*(11), e5002. https://doi.org/10.1002/nbm.5002

Lorenz, J., Hansen, H. C., Kunze, K., & Bromm, B. (1996). Sensory deficits of a nerve root lesion can be objectively documented by somatosensory evoked potentials elicited by painful infrared laser stimulations: A case study. *Journal of Neurology, Neurosurgery, and Psychiatry*, *61*(1), 107–110. https://doi.org/10.1136/jnnp.61.1.107

Manjón, J. V., Coupé, P., Martí-Bonmatí, L., Collins, D. L., & Robles, M. (2010). Adaptive non-local means denoising of MR images with spatially varying noise levels: Spatially Adaptive Nonlocal Denoising. *Journal of Magnetic Resonance Imaging*, *31*(1), 192–203. https://doi.org/10.1002/jmri.22003

Markuerkiaga, I., Marques, J. P., Gallagher, T. E., & Norris, D. G. (2021). Estimation of laminar BOLD activation profiles using deconvolution with a physiological point spread function. *Journal of Neuroscience Methods*, *353*, 109095. https://doi.org/10.1016/j.jneumeth.2021.109095

Martucci, K. T., Weber, K. A., & Mackey, S. C. (2019). Altered Cervical Spinal Cord Resting-State Activity in Fibromyalgia. *Arthritis & Rheumatology*, *71*(3), 441–450. https://doi.org/10.1002/art.40746

Martucci, K. T., Weber, K. A., & Mackey, S. C. (2021). Spinal Cord Resting State Activity in Individuals With Fibromyalgia Who Take Opioids. *Frontiers in Neurology*, *12*. https://www.frontiersin.org/articles/10.3389/fneur.2021.694271

Mouraux, A., & Iannetti, G. D. (2018). The search for pain biomarkers in the human brain. *Brain*, *141*(12), 3290–3307. https://doi.org/10.1093/brain/awy281

Mueller, R., Dabbagh, A., & Eippert, F. (2024). ThermoSlide: an MR-compatible thermode positioning device (v1.0). Zenodo. https://doi.org/10.5281/zenodo.10475479

Nash, P., Wiley, K., Brown, J., Shinaman, R., Ludlow, D., Sawyer, A.-M., Glover, G., & Mackey, S. (2013). Functional magnetic resonance imaging identifies somatotopic organization of nociception in the human spinal cord: *Pain*, *154*(6), 776–781. https://doi.org/10.1016/j.pain.2012.11.008

Niazy, R. K., Beckmann, C. F., Iannetti, G. D., Brady, J. M., & Smith, S. M. (2005). Removal of FMRI environment artifacts from EEG data using optimal basis sets. *NeuroImage*, *28*(3), 720–737. https://doi.org/10.1016/j.neuroimage.2005.06.067

Noble, S., Scheinost, D., & Constable, R. T. (2021). A guide to the measurement and interpretation of fMRI test-retest reliability. *Current Opinion in Behavioral Sciences*, *40*, 27–32. https://doi.org/10.1016/j.cobeha.2020.12.012

Oliva, V., Hartley-Davies, R., Moran, R., Pickering, A. E., & Brooks, J. C. (2022). Simultaneous brain, brainstem and spinal cord pharmacological-fMRI reveals involvement of an endogenous opioid network in attentional analgesia. *ELife*, *11*, e71877. https://doi.org/10.7554/eLife.71877

Paulus, P. C., Castegnetti, G., & Bach, D. R. (2016). Modeling event-related heart period responses. *Psychophysiology*, *53*(6), 837–846. https://doi.org/10.1111/psyp.12622

Petkó, M., & Antal, M. (2000). Propriospinal afferent and efferent connections of the lateral and medial areas of the dorsal horn (laminae I-IV) in the rat lumbar spinal cord. *Journal of Comparative Neurology*, *422*(2), 312–325. https://doi.org/10.1002/(SICI)1096-9861(20000626)422:2<312::AID-CNE11>3.0.CO;2-A

Pierrot-Deseilligny, E., & Burke, D. (2012). *The Circuitry of the Human Spinal Cord: Spinal and Corticospinal Mechanisms of Movement*. Cambridge University Press.

Pinto, V., Szucs, P., Lima, D., & Safronov, B. V. (2010). Multisegmental Aδ- and C-Fiber Input to Neurons in Lamina I and the Lateral Spinal Nucleus. *The Journal of Neuroscience*, *30*(6), 2384–2395. https://doi.org/10.1523/JNEUROSCI.3445-09.2010

Power, J. D., Silver, B. M., Silverman, M. R., Ajodan, E. L., Bos, D. J., & Jones, R. M. (2019). Customized head molds reduce motion during resting state fMRI scans. *NeuroImage*, *189*, 141–149. https://doi.org/10.1016/j.neuroimage.2019.01.016

Powers, J., Ioachim, G., & Stroman, P. (2018). Ten Key Insights into the Use of Spinal Cord fMRI. *Brain Sciences*, *8*(9), 173. https://doi.org/10.3390/brainsci8090173

Prescott, S. A., Ma, Q., & De Koninck, Y. (2014). Normal and abnormal coding of somatosensory stimuli causing pain. *Nature Neuroscience*, *17*(2), 183–191. https://doi.org/10.1038/nn.3629

Purves, D., Augustine, G. J., Fitzpatrick, D., Hall, W., LaMantia, A.-S., & White, L. (2019). *Neurosciences*. De Boeck Superieur.

Quiton, R. L., & Greenspan, J. D. (2008). Across- and within-session variability of ratings of painful contact heat stimuli: *Pain*, *137*(2), 245–256. https://doi.org/10.1016/j.pain.2007.08.034

Quiton, R. L., Keaser, M. L., Zhuo, J., Gullapalli, R. P., & Greenspan, J. D. (2014). Intersession reliability of fMRI activation for heat pain and motor tasks. *NeuroImage: Clinical*, *5*, 309–321. https://doi.org/10.1016/j.nicl.2014.07.005

Rangaprakash, D., & Barry, R. L. (2022). Neptune: A toolbox for spinal cord functional MRI data processing and quality assurance. *Proceedings 30th Scientific Meeting*. International Society for Magnetic Resonance in Medicine. https://archive.ismrm.org/2022/0396.html

Rempe, T., Wolff, S., Riedel, C., Baron, R., Stroman, P. W., Jansen, O., & Gierthmühlen, J. (2015). Spinal and supraspinal processing of thermal stimuli: An fMRI study: Processing of Thermal Stimuli. *Journal of Magnetic Resonance Imaging*, *41*(4), 1046–1055. https://doi.org/10.1002/jmri.24627

Reyes del Paso, G. A., Montoro, C., Muñóz Ladrón de Guevara, C., Duschek, S., & Jennings, J. R. (2014). The effect of baroreceptor stimulation on pain perception depends on the elicitation of the reflex cardiovascular response: Evidence of the interplay between the two branches of the baroreceptor system. *Biological Psychology*, *101*, 82–90. https://doi.org/10.1016/j.biopsycho.2014.07.004

Rombouts, S. A., Barkhof, F., Hoogenraad, F. G., Sprenger, M., Valk, J., & Scheltens, P. (1999). Test-Retest Analysis With Functional MR of the Activated Area in the Human Visual Cortex: *Journal of Neuro-Ophthalmology*, *19*(2), 112. https://doi.org/10.1097/00041327-199906000-00012

Rowald, A., Komi, S., Demesmaeker, R., Baaklini, E., Hernandez-Charpak, S. D., Paoles, E., Montanaro, H., Cassara, A., Becce, F., Lloyd, B., Newton, T., Ravier, J., Kinany, N., D'Ercole, M., Paley, A., Hankov, N., Varescon, C., McCracken, L., Vat, M., … Courtine, G. (2022). Activity-dependent spinal cord neuromodulation rapidly restores trunk and leg motor functions after complete paralysis. *Nature Medicine*, *28*(2), 260–271. https://doi.org/10.1038/s41591-021-01663-5

San Emeterio Nateras, O., Yu, F., Muir, E. R., Bazan, C., Franklin, C. G., Li, W., Li, J., Lancaster, J. L., & Duong, T. Q. (2016). Intrinsic Resting-State Functional Connectivity in the Human Spinal Cord at 3.0 T. *Radiology*, *279*(1), 262–268. https://doi.org/10.1148/radiol.2015150768

Seifert, A. C., Xu, J., Kong, Y., Eippert, F. C., Miller, K. L., Tracey, I., & Vannesjo, S. J. (2023). *Thermal Stimulus Task fMRI in the Cervical Spinal Cord at 7 Tesla* [Preprint]. Bioengineering. https://doi.org/10.1101/2023.01.31.526451

Shekhtmeyster, P., Duarte, D., Carey, E. M., Ngo, A., Gao, G., Olmstead, J. A., Nelson, N. A., & Nimmerjahn, A. (2023). Trans-segmental imaging in the spinal cord of behaving mice. *Nature Biotechnology*. https://doi.org/10.1038/s41587-023-01700-3

Sherrington, C. S. (1898). Experiments in Examination of the Peripheral Distribution of the Fibres of the Posterior Roots of Some Spinal Nerves. Part II. *Philosophical Transactions of the Royal Society of London. Series B, Containing Papers of a Biological Character*, *190*, 45–186.

Shrout, P. E., & Fleiss, J. L. (1979). Intraclass Correlations: Uses in Assessing Rater Reliability. *Psychological Bulletin*, *86*(2), 420.

Sluka, K. A., Wager, T. D., Sutherland, S. P., Labosky, P. A., Balach, T., Bayman, E. O., Berardi, G., Brummett, C. M., Burns, J., Buvanendran, A., Caffo, B., Calhoun, V. D., Clauw, D., Chang, A., Coffey, C. S., Dailey, D. L., Ecklund, D., Fiehn, O., Fisch, K. M., … the A2CPS Consortium. (2023). Predicting chronic postsurgical pain: Current evidence and a novel program to develop predictive biomarker signatures. *Pain*, *164*(9), 1912–1926. https://doi.org/10.1097/j.pain.0000000000002938

Sprenger, C., Eichler, I.-C., Eichler, L., Zöllner, C., & Büchel, C. (2018). Altered Signaling in the Descending Pain-modulatory System after Short-Term Infusion of the μ-Opioid Agonist Remifentanil. *The Journal of Neuroscience*, *38*(10), 2454–2470. https://doi.org/10.1523/JNEUROSCI.2496-17.2018

Sprenger, C., Finsterbusch, J., & Buchel, C. (2015). *Spinal Cord–Midbrain Functional Connectivity Is Related to Perceived Pain Intensity: A Combined Spino-Cortical fMRI Study*. 10.

Sprenger, C., Stenmans, P., Tinnermann, A., & Büchel, C. (2018). Evidence for a spinal involvement in temporal pain contrast enhancement. *NeuroImage*, *183*, 788–799. https://doi.org/10.1016/j.neuroimage.2018.09.003

Stanley, O. W., Kuurstra, A. B., Klassen, L. M., Menon, R. S., & Gati, J. S. (2021). Effects of phase regression on high-resolution functional MRI of the primary visual cortex. *NeuroImage*, *227*, 117631. https://doi.org/10.1016/j.neuroimage.2020.117631

Stroman, P. (2002). Mapping of Neuronal Function in the Healthy and Injured Human Spinal Cord with Spinal fMRI. *NeuroImage*, *17*(4), 1854–1860. https://doi.org/10.1006/nimg.2002.1305

Stroman, P. W., Kornelsen, J., Bergman, A., Krause, V., Ethans, K., Malisza, K. L., & Tomanek, B. (2004). Noninvasive assessment of the injured human spinal cord by means of functional magnetic resonance imaging. *Spinal Cord*, *42*(2), 59–66. https://doi.org/10.1038/sj.sc.3101559

Summers, P. E., Brooks, J. C. W., & Cohen-Adad, J. (2014). Spinal Cord fMRI. In *Quantitative MRI of the Spinal Cord* (pp. 221–239). Elsevier. https://doi.org/10.1016/B978-0-12-396973-6.00015-0

Summers, P. E., Ferraro, D., Duzzi, D., Lui, F., Iannetti, G. D., & Porro, C. A. (2010). A quantitative comparison of BOLD fMRI responses to noxious and innocuous stimuli in the human spinal cord. *NeuroImage*, *50*(4), 1408–1415. https://doi.org/10.1016/j.neuroimage.2010.01.043

Thron, A. K. (2016). *Vascular Anatomy of the Spinal Cord: Radioanatomy as the Key to Diagnosis and Treatment*. Springer.

Tinnermann, A., Büchel, C., & Cohen-Adad, J. (2021). Cortico-spinal imaging to study pain. *NeuroImage*, *224*, 117439. https://doi.org/10.1016/j.neuroimage.2020.117439

Topfer, R., Starewicz, P., Lo, K.-M., Metzemaekers, K., Jette, D., Hetherington, H. P., Stikov, N., & Cohen-Adad, J. (2016). A 24-channel shim array for the human spinal cord: Design, evaluation, and application. *Magnetic Resonance in Medicine*, *76*(5), 1604–1611. https://doi.org/10.1002/mrm.26354

Tracey, I. (2021). Neuroimaging enters the pain biomarker arena. *Science Translational Medicine*, *13*(619), eabj7358. https://doi.org/10.1126/scitranslmed.abj7358

Tsivaka, D., Williams, S. C. R., Medina, S., Kowalczyk, O. S., Brooks, J. C. W., Howard, M. A., Lythgoe, D. J., & Tsougos, I. (2023). A second-order and slice-specific linear shimming technique to improve spinal cord fMRI. *Magnetic Resonance Imaging*, *102*, 151–163. https://doi.org/10.1016/j.mri.2023.06.012

Tustison, N. J., & Gee, J. (2010). N4ITK: Nick's N3 ITK Implementation For MRI Bias Field Correction. *The Insight Journal*. https://doi.org/10.54294/jculxw

Uludağ, K., Müller-Bierl, B., & Uğurbil, K. (2009). An integrative model for neuronal activity-induced signal changes for gradient and spin echo functional imaging. *NeuroImage*, *48*(1), 150–165. https://doi.org/10.1016/j.neuroimage.2009.05.051

Upadhyay, J. (2015). Test–retest reliability of evoked heat stimulation BOLD fMRI. *Journal of Neuroscience Methods*, 9.

Vahdat, S., Khatibi, A., Lungu, O., Finsterbusch, J., Büchel, C., Cohen-Adad, J., Marchand-Pauvert, V., & Doyon, J. (2020). Resting-state brain and spinal cord networks in humans are functionally integrated. *PLOS Biology*, *18*(7), e3000789. https://doi.org/10.1371/journal.pbio.3000789

Vallat, R. (2018). Pingouin: Statistics in Python. *Journal of Open Source Software*, *3*(31), 1026. https://doi.org/10.21105/joss.01026

Vannesjo, S. J., Clare, S., Kasper, L., Tracey, I., & Miller, K. L. (2019). A method for correcting breathing-induced field fluctuations in T2*-weighted spinal cord imaging using a respiratory trace. *Magnetic Resonance in Medicine*, *81*(6), 3745–3753. https://doi.org/10.1002/mrm.27664

Veraart, J., Novikov, D. S., Christiaens, D., Ades-aron, B., Sijbers, J., & Fieremans, E. (2016). Denoising of diffusion MRI using random matrix theory. *NeuroImage*, *142*, 394–406. https://doi.org/10.1016/j.neuroimage.2016.08.016

Villemure, C., & Bushnell, M. C. (2002). Cognitive modulation of pain: How do attention and emotion influence pain processing? *Pain*, *95*(3), 195–199. https://doi.org/10.1016/S0304-3959(02)00007-6

Weber, K. A., Chen, Y., Wang, X., Kahnt, T., & Parrish, T. B. (2016a). Functional magnetic resonance imaging of the cervical spinal cord during thermal stimulation across consecutive runs. *NeuroImage*, *143*, 267–279. https://doi.org/10.1016/j.neuroimage.2016.09.015

Weber, K. A., Chen, Y., Wang, X., Kahnt, T., & Parrish, T. B. (2016b). Lateralization of cervical spinal cord activity during an isometric upper extremity motor task with functional magnetic resonance imaging. *NeuroImage*, *125*, 233–243. https://doi.org/10.1016/j.neuroimage.2015.10.014

Wiech, K. (2016). Deconstructing the sensation of pain: The influence of cognitive processes on pain perception. *Science*, *354*(6312), 584–587. https://doi.org/10.1126/science.aaf8934

Wilson, S. M., Bautista, A., Yen, M., Lauderdale, S., & Eriksson, D. K. (2017). Validity and reliability of four language mapping paradigms. *NeuroImage: Clinical*, *16*, 399–408. https://doi.org/10.1016/j.nicl.2016.03.015

Winkler, A. M., Ridgway, G. R., Webster, M. A., Smith, S. M., & Nichols, T. E. (2014). Permutation inference
for the general linear model. *NeuroImage*, *92*, 381–397. https://doi.org/10.1016/j.neuroimage.2014.01.060

Woo, C.-W., & Wager, T. D. (2016). What reliability can and cannot tell us about pain report and pain
neuroimaging: *PAIN*, *157*(3), 511–513. https://doi.org/10.1097/j.pain.0000000000000442

Woolrich, M. W., Ripley, B. D., Brady, M., & Smith, S. M. (2001). Temporal Autocorrelation in Univariate
Linear Modeling of FMRI Data. *NeuroImage*, *14*(6), 1370–1386. https://doi.org/10.1006/nimg.2001.0931

Viessmann, O., Scheffler, K., Bianciardi, M., Wald, L. L., & Polimeni, J. R. (2019). Dependence of resting-
state fMRI fluctuation amplitudes on cerebral cortical orientation relative to the direction of B0 and
anatomical axes. *Neuroimage, 196*, 337-350.

Yang, P.-F., Wang, F., & Chen, L. M. (2015). Differential fMRI Activation Patterns to Noxious Heat and
Tactile Stimuli in the Primate Spinal Cord. *Journal of Neuroscience*, *35*(29), 10493–10502.
https://doi.org/10.1523/JNEUROSCI.0583-15.2015

Yoshizawa, T., Nose, T., Moore, G. J., & Sillerud, L. O. (1996). Functional Magnetic Resonance Imaging of
Motor Activation in the Human Cervical Spinal Cord. *NeuroImage*, *4*(3), 174–182.
https://doi.org/10.1006/nimg.1996.0068

1585

1586

1587

1588

1589

1590

1591 **Supplementary Material**

1592

1593

1594 **Reliability of task-based fMRI in the dorsal horn of the human spinal cord**

1595

1596 Alice Dabbagh, Ulrike Horn, Merve Kaptan, Toralf Mildner, Roland Müller, Jöran Lepsien,

1597 Nikolaus Weiskopf, Jonathan C.W. Brooks, Jürgen Finsterbusch, Falk Eippert

1598

1599   **Supplementary Figure 1. Individual values underlying ICC calculation.** Participant-wise data of Day 1 and Day 2

1600   peak values for skin conductance responses (SCR), pupil dilation responses (PDR) and heart period responses

1601   (HPR), as well as average top 10% β values of the left dorsal horn (ROI) and the dilated left dorsal quadrant (Dilated

1602   ROI) in spinal cord segment C6.

1603

Supplementary Figure 2. Group-level fMRI results. Results are shown for the Reliability Run (i.e., one run per day;

top row; same image as in main manuscript used for comparison purposes here) and Combined Runs (i.e. average of

four runs per day; bottom row). Only voxels surviving a threshold of p < 0.05 (corrected for multiple comparisons via a

permutation test in a mask of the left dorsal horn in spinal cord segment C6) are displayed on top of a T2*-weighted

spinal cord template (PAM50) in axial view, and on top of a T2-weighted spinal cord template (PAM50) in the sagittal

view. In contrast to the Reliability Run, the Combined Runs demonstrated substantial overlap across both days. Spinal

cord segment C6 is outlined in green in the sagittal images.

1611

Supplementary Figure 3. Test-retest reliability across both days for subjective ratings, peripheral physiological

data and BOLD response amplitudes. Reliability is indicated via ICCs (plotted as dots with 95% CI represented as a

line). *Combined Runs*: ICCs are reported for (from left to right) verbal ratings, SCR, PDR, HPR and top 10% β-estimate

in the left dorsal horn of C6 (ROI). *Denoising DHR*: ICC of the top 10% β-estimate in the left dorsal horn of C6 (ROI) of

only the Reliability Run, obtained from a GLM with an additional regressor of right dorsal horn activity. *Within-run*

reliability: ICCs obtained from comparing either odd and even trials numbers (odd-even), or the first and second half of

a run (early-late). Colors indicate ICC interpretation according to Cicchetti (1994): dark red: ICC < 0.4, poor; medium

red: ICC 0.4 - 0.59, fair; orange: ICC 0.6 - 0.74, good; yellow: ICC 0.75 - 1.0, excellent.

Supplementary Table 1

Percent suprathreshold voxels in the four cord quadrants of each spinal segment.

Spinal cord segment	Total number of active voxels (p < 0.001)	DL	DR	VL	VR
Average					
C5	1140	50.1%	34.5%	5.6%	9.8%
C6	674	61.3%	38.4%	0.3%	0%
C7	604	14.6%	26.7%	13.4%	45.4%
C8	62	22.6%	56.5%	8.1%	12.9%
Day 1					
C5	438	79.0%	14.4%	4.6%	2.1%
C6	65	83.1%	10.8%	6.2%	0%
C7	59	37.3%	25.4%	5.1%	32.2%
C8	3	0%	100%	0%	0%
Day 2					
C5	302	28.1%	49.7%	4.0%	18.2%
C6	332	66.3%	27.7%	5.1%	0.9%
C7	235	9.8%	11.9%	48.9%	29.4%
C8	17	0%	82.4%	0%	17.6%

Notes. Results are based on the group-level results of each day's Reliability Run. ROI names refer to spinal cord quadrants in the respective segment. *Abbreviations*: dorsal left (DL), dorsal right (DR), ventral left (VL), ventral right (VR). This analysis was carried out using the masks of the four cord quadrants separately for each spinal segment.

1620

1621

Supplementary Table 2

Intraclass correlation coefficient and 95% confidence interval for subjective ratings, peripheral physiological data and BOLD response amplitudes of post-hoc analyses.

Change in analysis pipeline	Measures			ICC (95% CI)
	Ratings			0.71 (0.51–0.83)
	SCR			0.61 (0.36–0.78)
	PDR			0.19 (-0.16–0.81)
	HPR			0.73 (0.54–0.85)
Combined runs average	DH left	β	peak	0.20 (-0.12–0.48)
			Top 10%	0.09 (-0.22–0.39)
			avg	-0.08 (-0.38–0.23)
		z-score	peak	0.07 (-0.24–0.37)
			Top 10%	-0.01 (-0.31–0.30)
			avg	0.03 (-0.28–0.34)
DHR regressor	DH left	β	peak	0.11 (-0.21–0.40)
			Top 10%	0.08 (-0.25–0.37)
			avg	-0.07 (-0.37–0.24)
		z-score	peak	0.11 (-0.2–0.41)
			Top 10%	0.06 (-0.25–0.36)
			avg	0.025 (-0.29–0.33)
Within-run reliability Odd-even	SCR			0.81 (0.67–0.90)
	PDR			0.82 (0.66–0.90)
	HPR			0.86 (0.76–0.93)
	DH left	β	peak	0.63 (0.41–0.79)
			Top 10%	0.69 (0.49–0.83)
			avg	0.52 (0.27–0.71)
Early-late	SCR			0.56 (0.31–0.74)
	PDR			0.61 (0.35–0.78)
	HPR			0.79 (0.64–0.89)
	DH left	β	peak	0.31 (0.02–0.56)
			Top 10%	0.36 (0.07–0.59)
			avg	-0.003 (-0.31–0.31)

Abbreviations: skin conductance response (SCR), pupil dilation response (PDR), heart period response (HPR), dorsal horn (DH), right dorsal horn (DHR) in spinal cord segment C6.

Supplementary Table 3

Correlations between response measures.

Peripheral / subjective measure	BOLD parameter estimate (top 10%)	Pearson's r	p-value One-tailed
Ratings	β	-0.18	0.855
	z-score	-0.30	0.965
SCR	β	0.34	0.017 *
	z-score	0.36	0.014 *
HPR	β	-0.28	0.038 *
	z-score	-0.28	0.039 *
PDR	β	-0.11	0.722
	z-score	-0.05	0.611

Notes. Results are based on individual BOLD responses and peripheral physiological responses and subjective ratings of the reliability run. We correlated responses averaged across days for each measure. BOLD responses were quantified as the top 10 % of β estimates and z-scores extracted from the left dorsal horn in spinal cord segment C6. *Abbreviations*: skin conductance response (SCR), pupil dilation response (PDR), heart period response (HPR). For further information see section 2.8.4. * p < 0.05

1622

Supplementary Table 4

Correlations between BOLD parameter estimates and indicators of data quality.

Data quality estimate	BOLD parameter estimate (top 10%)	Pearson's r	p-value One-tailed
Motion	β	0.16	0.159
	z-score	0.19	0.118
Normalization quality	β	-0.25	0.939
	z-score	-0.14	0.8
Angulation relative to B0	β	0.41	0.004 **
	z-score	0.40	0.005 **

Notes. Results are based on individual BOLD responses and data quality indicators of the reliability run. We correlated absolute differences across days for each measure. BOLD responses were quantified as the top 10 % of β estimates and z-scores extracted from the left dorsal horn in spinal cord segment C6. Motion was quantified as root mean square intensity differences of each volume to reference volume. Normalization quality was quantified as the Dice coefficient between the segmentation of the normalized mean functional image and the PAM50 cord mask across the same z-range. The angulation relative to B0 describes the angle between the scanner's z-axis (aligned with B0) and the z direction in the slice-stack. For further information see section 2.8.5. * p < 0.05, ** p < 0.01

1623

Deviations from preregistration

In the preregistration, we stated that in addition to ICC(3,1) we would report the Pearson
correlation coefficient and ICC(2,1) as indicators of reliability. However, for the sake of brevity we
ultimately decided to report only ICC(3,1), as all indicators were found to be highly similar.

In the preregistration, we stated that we aimed to calculate voxel-wise ICC maps. However, given
that we observed almost no overlap of activation across days in the analysis reported here (thus
making a voxel-wise assessment pointless), we decided to focus solely on the ROI assessments.

In the preregistration, we stated that we would investigate spatial aspects of reliability via x-, y-,
and z-coordinates. However, for the sake of brevity we decided to instead employ Dice coefficients
(i.e. a measure of spatial overlap), as we deemed them a more succinct and comprehensive
representation of our data, also considering the complexity of the manuscript.

In the preregistration, we stated that we would assess reliability only in the ipsilateral dorsal horn
of spinal cord segment C6. However, after observing robust activation extending to areas outside
the gray matter, we chose to also investigate a larger region encompassing the draining vein
territory.

Any other analyses carried out here, but not included in the preregistration, are clearly indicated
as post-hoc analyses in the manuscript (see section 2.8).

1641