EryDB: a transcriptomic profile database for erythropoiesis and erythroid-related diseases

Guangmin Zheng[1,2,#], Song Wu[1,2,3,#], Zhaojun Zhang[1,2,4,#], Zijuan Xin[1,2], Lijuan Zhang[1,2], Siqi Zhao[1,2], Jing Wu[1,2], Yanxia Liu[1,4], Meng Li[1,4], Xiuyan Ruan[1,4], Yiming Bao[1,2,3,*], Hongzhu Qu[1,2,4,*], Xiangdong Fang[1,2,4,*]

1 CAS Key Laboratory of Genome Sciences and Information, Beijing Institute of Genomics, Chinese Academy of Sciences / China National Center for Bioinformation, Beijing 100101, China

2 University of Chinese Academy of Sciences, Beijing 100049, China

3 National Genomics Data Center, Beijing Institute of Genomics, Chinese Academy of Sciences / China National Center for Bioinformation, Beijing 100101, China

4 Beijing Key Laboratory of Genome and Precision Medicine Technologies, Beijing Institute of Genomics, Chinese Academy of Sciences / China National Center for Bioinformation, Beijing 100101, China

#Contributed equally.

*Corresponding authors.

E-mail: baoym@big.ac.cn (Bao Y), quhongzhu@big.ac.cn (Qu H), fangxd@big.ac.cn (Fang X).

Running title: *Guangmin Z et al* / A transcriptomic database of erythropoiesis and erythroid-related diseases

Total number of words: 3162

Total Figure number: 4

Total Table number: 1

Total supplementary tables: 3

Abstract

Erythropoiesis is a finely regulated and complex process that involves multiple transformations from hematopoietic stem cells to mature red blood cells at hematopoietic sites from the embryonic to the adult stages. Investigations into its molecular mechanisms have generated a wealth of expression data, including bulk and single-cell RNA sequencing data. A comprehensively integrated and well-curated erythropoiesis-specific database will greatly facilitate the mining of gene expression data and enable large-scale research of erythropoiesis and erythroid related diseases. Here, we present EryDB (https://ngdc.cncb.ac.cn/EryDB/home), an open-access and comprehensive database dedicated to the collection, integration, analysis, and visualization of transcriptomic data for erythropoiesis and erythroid-related diseases. Currently, the database includes expertly curated quality-assured data of 3,803 samples and 1,187,119 single cells derived from 107 public studies of three species (*Homo sapiens*, *Mus musculus*, and *Danio rerio*), nine tissue types, and five diseases. EryDB provides users with the ability to not only browse the molecular features of erythropoiesis between tissues and species, but also perform computational analyses of single-cell and bulk RNA sequencing data, thus serving as a convenient platform for customized queries and analyses. EryDB v1.0 is freely accessible at https://ngdc.cncb.ac.cn/EryDB/home.

KEYWORDS: Erythropoiesis; Database; Data analysis; Transcriptome; scRNA-seq; Data integration

Introduction

Erythrocytes (red blood cells), the most abundant of the blood cells, are responsible for transporting oxygen from lungs to other tissues [1]. Erythrocyte impairment is attributed to acquired/hereditary erythropoietic diseases in which abnormalities occur in causative genes, including genes that encode erythrocyte membrane proteins, globins, and enzymes, as well as genes that are required for erythropoiesis [2]. Erythropoiesis is a finely regulated and complicated process that involves multiple transformations from hematopoietic stem cells to mature red blood cells at hematopoietic sites from the embryonic to adult stages [3]. Moreover, the current understanding of erythropoiesis and the pathophysiology of erythropoietic defects is limited. The underlying mechanisms need to be better defined to improve *in vitro* regeneration of red blood cells [4, 5] for the development of clinical treatments [3, 6]. Large amounts of multi-omics data are now available [7-13], providing unprecedented insights into erythroid biology and generating many novel hypotheses. However, most of the omics data are available as raw data deposited in public repositories, such as the Genome Sequence Archive (GSA) [14] and Sequence Read Archive (SRA) [15], making it challenging for clinicians and researchers to reanalyze the data and gain biological insights [16]. Although some research groups have made their erythroid profile data web-accessible [9, 11, 17], no erythropoiesis-specific database that can

facilitate the reuse of omics data is available to the erythroid biology community. An
authoritative and integrated omics database that contains the molecules, pathways, and
events involved in erythropoiesis will help to mitigate these shortcomings and
accelerate research in this area.
Here, we present EryDB (https://ngdc.cncb.ac.cn/EryDB/home), a database that
contains curated, quality-assured, and pre-analyzed transcriptome data of
erythropoiesis and erythroid-related diseases. The EryDB platform allows users to
search, browse, and analyze gene expression profiles or related data among various
tissues or experimental types. Furthermore, gene expression changes under various
pathological conditions can be analyzed towards developing diagnostic and prognostic
tools for various erythropoietic diseases.
In this paper, we describe the breadth of the transcriptomic data available in EryDB,
which includes bulk and single-cell RNA sequencing (RNA-seq) data. The datasets
are classified into four major erythropoiesis-related categories, namely erythroid
differentiation, genes, compounds, and diseases that users can mine to obtain
erythroid-specific information. Customized filtering of the data can be performed
under any of the four categories. These tools allow users to easily retrieve erythroid-
specific data of interest to them. Moreover, EryDB provides a functional module
named the Erythroid Atlas that can be used to integrate and compare erythropoiesis-
related datasets from different tissues, namely bone marrow, peripheral blood, cord
blood, and embryos. The rapid data query and analysis capabilities provided by
EryDB enable efficient data mining for basic and clinical research.
107
Data collection and processing

Data collection and classification
The process that was used to construct EryDB is outlined in **Figure 1**. Transcriptome
data of erythroid cells were obtained from the Gene Expression Omnibus (GEO) [18],
Sequence Read Archive (SRA) [15], and Genome Sequence Archive (GSA) [14]
databases using the keywords "erythroid" or "erythrocyte". We also searched PubMed
to identify recently published single-cell transcriptome studies by the keywords
"blood mononuclear cell" and by the keywords "single cell" and "erythroid". We
screened the titles, abstracts, and sample information to identify studies that
specifically performed gene expression profiling of erythroid cells. For the single-cell
RNA-seq (scRNA-seq) data, only erythroid-related studies with >300 cells were
included. We also included published [19-22] and unpublished data from our
laboratory. Finally, for each dataset, we have defined a custom ID for the Erythroid
Database for user-friendly searching, and we also keep the source as one of the
metadata of the dataset.
123

Metadata curation
Metadata were downloaded from GEO, SRA, GSA and manually curated by experts in
erythropoiesis (Figure 1). Repetitive and redundant information was removed. Each
dataset was manually annotated with biologically relevant information, including

model organism, sample tissue, experiment type, and omics type. A description of
each study, GEO or GSA identifiers, and references to the original publications were
also included in the metadata. All the datasets are labeled with species, tissue, and
experiment type (*in vivo*/*in vitro*).
132

Pre-processing and quality control of bulk transcriptomic data

For the RNA-seq datasets, we downloaded raw data, but included pre-processed data
or raw count files when raw data were not available. We use the identical processing
pipeline for the data sets that can obtain the raw sequencing data .The adaptor and
low-quality sequences were removed using *Trimmomatic* (v.0.36) [23]. Gene
expression was quantified using *Salmon* (v.1.4.0) [24] (Figure 1). Trimmed mean of
M-values and Voom normalization were performed using the *EdgeR* (v.3.42.4) [25]
and *limma* (v.3.48.3) [26] packages in R (v.4.3.0), respectively. For the integrated
datasets in erythroid atlas in EryDB, we have ensured that all sample use the same
analysis pipeline and then remove batch before proceeding to downstream analysis.
We obtained gene annotation for Homo sapiens (GRCh38), Mus musculus (GRCm38),
and Danio rerio (Zv9) from Ensemble (release-91).
145

Pre-processing and quality control of single-cell transcriptomic data

For the scRNA-seq dataset, we downloaded the processed expression matrices when
no raw data were available, the raw data were reprocessed using the method used in
the original publication, including quality control and normalization. Low-quality
cells (expressing <300 genes) and mitochondrial genes (>20% of the reads mapped to
mitochondrial genes) were excluded. Library size normalization was performed using
the *Seurat* (v.4.3.0)[27] package. The batch effects of combined expression data were
removed using *Seurat* CCA or Harmony (v.0.1.1) [28]. For the integrated single-cell
datasets in in erythroid atlas in EryDB, we also ensured that comparative analyses
were performed after using the identical processing pipeline.
156

Implementation

158 EryDB is an interactive web application built using the Python/Flask (1.0.2) web
159 framework. We used *Vue*, *Plotly*, *Encharts*, and *Highcharts* with Python for data
160 visualization, and *iView.Table* to construct searchable tables.
161

Database content and usage

163 The EryDB database is a free public biological resource for erythropoiesis-related
164 research. Users can browse datasets, examine the expression of erythropoiesis-related
165 genes, and compare transcriptome differences and cell variations in the context of
166 erythropoiesis and erythroid-related diseases.
167

Data records

In the current release, EryDB contains high-quality analytical results of 118 datasets

and 3,803 samples from 107 studies of three species, *Homo sapiens*, *Mus musculus*,

and *Danio rerio*. Fifteen of these studies were scRNA-seq studies involving 1,187,119

single cells; the others were bulk RNA-seq studies. Among the datasets, 60 (50.85%),

49 (41.53%), and 9 (7.63%) were for *Homo sapiens, Mus musculus*, and *Danio rerio*,

respectively. The 118 datasets were associated with 11 tissue types: bone marrow, cord

blood, peripheral blood, spleen, embryo, fetal tissue, heart, kidney, induced

pluripotent stem cells (iPSC), cell line, and other (unknown). Data statistics for

EryDB are shown in **Table 1**. The accession numbers for all the erythroid-related

datasets are listed in **Supplementary Table 1**.

User-friendly search modules for data queries

EryDB provides three methods for datasets searches. First, a quick-search box on the

home page that allows real-time queries by specifying genes, tissues, dataset ID,

organism name or PubMed Unique Identifier (PMID). Second, category-specific

searches by erythroid differentiation, genes, compounds, or diseases that are also

available on the home page: Erythroid Differentiation—the process of erythroid

differentiation is subdivided into 12 cell types (**Figure 2A**), and datasets that contain a

cell type of interest are listed by clicking of the cell type; Genes—important genes

related to erythroid differentiation and development are summarized as a keyword

cloud and a sorted table, and datasets associated with a gene of interest can be

accessed by clicking on the gene name (**Figure 2B**); Compounds—molecular

compounds related to erythroid differentiation and development are listed (**Figure 2C**),

and datasets associated with a compound of interest can be accessed by clicking on the

compound name; and Diseases—the five main diseases related to erythropoiesis are

available in 14 datasets (**Figure 2D**), and datasets associated with a disease of interest

can be accessed by clicking on the disease name. The query results from these four

categories are presented as tables that contains the dataset ID, species, tissue, and

experiment type (**Figure 2E**). The table heading in each column is a drop-down box

that enables further filtering of the datasets according to specific conditions. Third,

target datasets can be selected by clicking on "Search" at the top of the home page and

choosing one or more of the options on the navigation bar on the left-hand side of the

page, namely cell type, reported gene, compound type, disease type, species, tissue,

experiment type, and omics type (**Figure 2F**). Detailed information about the content

of the datasets that can be searched under these conditions is listed in **Supplementary**

Table 2. The results of such queries are listed in tables that are similar to those

obtained by category-specific searches.

Clicking on any word in the results tables from all three types of searches expands the

overview and the gene expression profiles of the dataset. The overview includes a

research summary, research organization, sequencing platform information, specific

sample information, data processing method, overall design introduction, data

contributors, and contact information for the dataset (**Figure 2G**). The sample

accession numbers are linked to the database source of the original data. Because of

the different technologies used for bulk RNA-seq and scRNA-seq, the gene expression

profile datasets from the two technologies are presented separately.

Comprehensive multidimensional online data exploration

Bulk RNA-seq data exploration

EryDB provides four analysis modules for the bulk transcriptome datasets on the
overview page: Expression Profile—analysis of expression changes for a gene, sample
group, and dataset specified by users (**Figure 3A**). The detailed results are shown
when the sample bar is selected; Principal Component Analysis—a two-dimensional
(2D) plot of the principal component analysis results is obtained, where individual
sample's group are color-coded (**Figure 3B**); Differential Analysis—a volcano plot of
differentially expressed genes (DEGs) between two user-specified groups and a
browsable table below the plot are obtained (**Figure 3C**); and Enrichment Analysis—
enrichment analysis results of DEGs between user-specified groups, including Gene
Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) functional
and pathway annotations (**Figure 3D**).

To demonstrate the database functions, we applied EryDB to the GEO dataset
GSE128268, which is a bulk RNA-seq dataset of an erythroid cell culture of human
umbilical cord-derived CD34+ hematopoietic stem and progenitor cells [29]. We
found that the hemoglobin subunit gene *HBA1* was highly expressed at the colony-
forming unit erythroid (CFU-E) stage (Figure 3A). Furthermore, the burst forming
unit-erythroid (BFU-E) and CFU-E cells had transcriptome profiles that were clearly
distinct, and both profiles were distinct from those of umbilical cord-derived CD34+
hematopoietic stem and progenitor cells (Figure 3B). To determine which pathways
were most upregulated in CFU-E cells, we compared the gene expression signatures
between CFU-E and BFU-E cells and found that gas and oxygen transport were
upregulated in CFU-E cells (Figure 3D). Analysis of the DEGs showed that
hemoglobin markers (e.g., *HBA1*, *HBA2*, and *HBG2*) were among the top 25 most
upregulated genes at the CFU-E stage (Figure 3C). Together, these results suggest that
CFU-E cells undergo a switch that results in a hemoglobin-based oxygen-carrier
phenotype.

The Erythroid Atlas page in EryDB allows comparisons to be made between
transcriptome datasets associated with erythroid development. Users can explore the
gene expression profiles of different differentiation systems in humans, mice, and
zebrafish (**Figure 3E**). For example, changes in the expression of the transcription
factor gene *RUNX2* at different erythroid differentiation stages in humans (*in vivo*) or
changes induced *in vitro* can be browsed (**Figure 3F**). The results were obtained by
the integration of 11 datasets that contain a total of 207 samples. All the data are
analyzed and normalized in a standard pipeline to eliminate the batch effect of the data
sources. Moreover, DEGs can be identified from volcano plots and tables between any
two stages of the erythroid differentiation process by selecting corresponding tags in
the drop-down box, which enables users to identify genes of interest for further study.

scRNA-seq data exploration

EryDB provides six analysis modules for scRNA-seq data: Visualization & Feature—

enables dimension reduction analysis and signature expression queries among cells.
Gene expression among single cells is displayed as interactive t-distributed stochastic
neighbor embedding (t-SNE) or uniform maximal approximation projection (UMAP)
plots (**Figure 4A** and **B**), and the expression of specific genes among cells can be
queried in an extra t-SNE plot (**Figure 4C**); Marker & Enrichment—provides marker
gene expression of cell types as a heatmap (**Figure 4D**) and enables functional
enrichment analysis of genes in cell populations. The KEGG pathway or GO
enrichment results for each stage are presented as bar graphs (**Figure 4E**); Difference
& Enrichment—the expression of DEGs in the same cell type from different group is
presented in a volcano plot and the functional enrichment results of these DEGs are
presented in a bar plot; Differentiation Trajectory—pseudotime analysis of cells where
pseudotime trajectories of cell differentiation are visualized in 2D space (**Figure 4F**),
and cell differentiation trajectories can be displayed by cell type (**Figure 4G**); Cell–
Cell Interaction—the interaction of ligands and receptors between different cell types
can be viewed as a circus plot, where the thickness of the lines indicates the
interaction strength. The communication types include growth factors, cytokines,
checkpoints, and others, which can be queried by selection in the box provided
(**Figure 4H**); and Cell–Cell Communication—cellular communication is visualized as
a river diagram in which each pattern represents different levels of communication
(**Figure 4I**). User can view the interaction of a signaling pathway of interest (**Figure
4J**) and the contribution of specific ligand–receptor pairs in this pathway (**Figure 4K**).
Together, these analyses cover most of the statistical bases of single-cell
transcriptomes.
To demonstrate the database functions, we applied EryDB to the GSA dataset
CRA002443, which is a single-cell transcriptome dataset of human 5–19 week post-
coitum fetal tissue (W5–19) [7], and explored the erythroid lineage properties. A total
of 12 cell clusters were identified in the W9 sample, and the number of erythroid cell
lineages in this sample was higher than the numbers in the other samples (Figure 4A
and B). Expression of the cyclin gene *CCNB1* was upregulated in erythroblasts
(Figure 4C). Markers for each cell population are plotted as a heatmap (Figure 4D)
that can be enlarged for convenient browsing of gene expression. The hemoglobin
gene *HBB* was highly expressed in erythroid cell lineages in the W9 sample, and
genes expressed in erythroblasts were enriched in several biological processes,
including response to UV and regulation of deoxyribonuclease activity (Figure 4E). In
the W9 group, endoderm-derived cell lineages had the strongest cell-cell interactions
among the top 40 pairs of ligands and receptors, but they showed much less
interactions in early erythroblasts (Figure 4H). Four communication patterns were
identified among different cell types (Figure 4I). The transforming growth factor
TGFβ signaling pathway in two of the patterns (patterns 1 and 2) was communicated
between most of the cells by multiple ligands and receptors (Figure 4J and K).
Similarly, the Erythroid Atlas page provides integrative analysis via six modules
across differentiated systems from the scRNA-seq datasets.
In summary, these examples demonstrate the power of EryDB to efficiently test novel

hypotheses using publicly available data of genes that regulate erythroid
differentiation and development. All visualizations can be downloaded as SVG or
PNG files, related result data tables can be downloaded in comma-separated value
(CSV) format or Excel sheets, and integrated expression matrix and metadata can be
downloaded for advancer users to do other personalized analysis. Besides, the analysis
results can be filtered with adjustable parameters. The fully functional design of
EryDB is intended to facilitate the widespread reuse of publicly available datasets to
address novel and unresolved questions and allow comparative analyses of similar
studies to resolve study-specific biases.
308
Discussion
A survey of erythroid-related omics databases confirmed that the currently available
databases are important resources that can be used to interpret gene regulation or
mutations related to erythropoiesis and erythroid-related diseases. Some erythroid-
related databases are no longer accessible (e.g., EpoDB [30], Hembase [31], and
ErythronDB [32]). Accessible databases (e.g., dbRBC [33], RBCmembrane [34], and
BloodSpot [35, 36]) that contain specific omics data associated with red blood cells
(RBCs) (**Supplmentary Table 1**) tend to lack diversity in their data sources, species
covered, or functional analysis tools, and do not contain single-cell sequencing data.
EryDB was designed to integrate most of the accessible public transcriptome datasets
related to erythropoiesis and enable the investigation of gene expression dynamics in
erythroid differentiation and erythropoiesis-related diseases. Compared with the other
RBC-related databases, EryDB has more user-friendly data search options, has more
extensive data sources, and more comprehensive analysis capabilities. In the sub-
module search of the datasets, users can select the search that meets their research
interests by selecting the relevant module. For example, in the disease module,
researchers can easily examine the gene expression changes under certain pathological
conditions for blood disease diagnosis [37-40] and in the compounds module, users
can obtain information about the small molecules that promote or inhibit erythroid
differentiation. The *in vitro* RBC regeneration data from our laboratory are unique in
EryDB and can be integrated with *in vivo*-derived erythroid cell data. This allows
users to compare the differences and mechanisms underlying erythropoiesis from
different global sources. EryDB also provides comprehensive functional analysis tools.
For example, cell–cell interaction analysis based on scRNA-seq datasets generated a
broad spectrum of information on the interactions between erythroid cells and the
other cell types. We found that *in vivo* erythroid cells had stronger cellular interactions
with the other cell types than *in vitro* erythroid cells had.
EryDB will be useful not only for computational biologists but also for bench
clinicians interested in erythropoietic disorders and for researchers involved in basic
erythropoiesis-related research. In future releases of EryDB, the transcriptomic data
will be continuously updated, and other types of omics data will be added. We will
also develop new analytical feature for further exploration of the multi-omics data.
341

Availability

EryDB v1.0 is freely accessible at https://ngdc.cncb.ac.cn/EryDB/home.

Author's contributions

Xiangdong Fang: Supervision, Project administration, Funding acquisition, Resources, Conceptualization.

Yiming Bao: Supervision, Project administration, Funding acquisition, Resources, Conceptualization.

Guangmin Zheng: Software, Visualization, Writing - Original Draft, Writing - Review & Editing.

Song Wu: Software, Visualization.

Zhaojun Zhang: Data curation, Resources, Investigation, Validation, Writing - Review & Editing.

Hongzhu Qu: Validation, Writing - Review & Editing.

Zijuan Xin: Data curation, Resources, Investigation.

Lijuan Zhang: Data curation, Resources, Investigation.

Meng Li: Validation.

Siqi Zhao: Validation.

Jing Wu: Validation.

Yanxia Liu: Validation.

Xiuyan Ruan: Validation.

All authors reviewed and approved the final manuscript.

Competing interests

The authors have declared no competing interests.

Acknowledgements

The authors acknowledged the helps from all the research groups for their RNA data. This research was supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (grant no. XDA16010602), National Natural Science Foundation of China (Grant Nos. 81870097, 82070114, 82270126), the National Key Research and Development Program of China (2022YFC2406803). The authors would like to thank Liwen (www.liwenbianji.cn) and Editage (www.editage.cn) for the English language review.

ORCID: 0000-0002-3781-170X.   Guangmin Zheng

ORCID: 0000-0002-6628-8620.   Xiangdong Fang

References

[1] Palis J. Primitive and definitive erythropoiesis in mammals. Front Physiol 2014; 5:3.

[2] Di Pierro E, Brancaleoni V, Granata F. Advances in understanding the pathogenesis of congenital erythropoietic porphyria. Br J Haematol 2016;

173(3):365-379.

[3] Caulier AL, Sankaran VG. Molecular and cellular mechanisms that regulate human
erythropoiesis. Blood 2022; 139(16):2450-2459.

[4] Batta K, Menegatti S, Garcia-Alegria E, Florkowska M, Lacaud G, Kouskoff V.
Concise Review: Recent Advances in the In Vitro Derivation of Blood Cell
Populations. Stem Cells Transl Med 2016; 5(10):1330-1337.

[5] Christaki EE, Politou M, Antonelou M, Athanasopoulos A, Simantirakis E,
Seghatchian J *et al*. Ex vivo generation of transfusable red blood cells from
various stem cell sources: A concise revisit of where we are now. Transfus Apher
Sci 2019; 58(1):108-112.

[6] Weiss MJ, dos Santos CO. Chaperoning erythropoiesis. Blood 2009;
113(10):2136-2144.

[7] Wang X, Yang L, Wang YC, Xu ZR, Feng Y, Zhang J *et al*. Comparative analysis
of cell lineage differentiation during hepatogenesis in humans and mice at the
single-cell transcriptome level. Cell Res 2020; 30(12):1109-1126.

[8] Yang D, Jang I, Choi J, Kim MS, Lee AJ, Kim H *et al*. 3DIV: A 3D-genome
Interaction Viewer and database. Nucleic Acids Res 2018; 46(D1):D52-D57.

[9] Tusi BK, Wolock SL, Weinreb C, Hwang Y, Hidalgo D, Zilionis R *et al*.
Population snapshots predict early haematopoietic and erythroid hierarchies.
Nature 2018; 555(7694):54-60.

[10] Weinreb C, Wolock S, Tusi BK, Socolovsky M, Klein AM. Fundamental limits
on dynamic inference from single-cell snapshots. Proc Natl Acad Sci U S A 2018;
115(10):E2467-E2476.

[11] Popescu DM, Botting RA, Stephenson E, Green K, Webb S, Jardine L *et al*.
Decoding human fetal liver haematopoiesis. Nature 2019; 574(7778):365-371.

[12] Cao J, O'Day DR, Pliner HA, Kingsley PD, Deng M, Daza RM *et al*. A human
cell atlas of fetal gene expression. Science 2020; 370(6518).

[13] Vanuytsel K, Matte T, Leung A, Naing ZH, Morrison T, Chui DHK *et al*. Induced
pluripotent stem cell-based mapping of beta-globin expression throughout human
erythropoietic development. Blood Adv 2018; 2(15):1998-2011.

[14] Members C-N, Partners. Database Resources of the National Genomics Data
Center, China National Center for Bioinformation in 2022. Nucleic Acids Res
2022; 50(D1):D27-D38.

[15] Leinonen R, Sugawara H, Shumway M, International Nucleotide Sequence
Database C. The sequence read archive. Nucleic Acids Res 2011; 39(Database
issue):D19-21.

[16] Rung J, Brazma A. Reuse of public genome-wide gene expression data. Nat Rev
Genet 2013; 14(2):89-99.

[17] Reyes M, Filbin MR, Bhattacharyya RP, Billman K, Eisenhaure T, Hung DT *et al*.
An immune-cell signature of bacterial sepsis. Nat Med 2020; 26(3):333-340.

[18] Barrett T, Troup DB, Wilhite SE, Ledoux P, Rudnev D, Evangelista C *et al*. NCBI
GEO: mining tens of millions of expression profiles--database and tools update.
Nucleic Acids Res 2007; 35(Database issue):D760-765.

[19] Xin Z, Zhang W, Gong S, Zhu J, Li Y, Zhang Z *et al*. Mapping Human
Pluripotent Stem Cell-derived Erythroid Differentiation by Single-cell
Transcriptome Analysis. Genomics Proteomics Bioinformatics 2021; 19(3):358-
376.

[20] Ren Y, Zhu J, Han Y, Li P, Wu J, Qu H *et al*. Regulatory association of long
noncoding RNAs and chromatin accessibility facilitates erythroid differentiation.
Blood Adv 2021; 5(23):5396-5409.

[21] Ding N, Xi J, Li Y, Xie X, Shi J, Zhang Z *et al*. Global transcriptome analysis for
identification of interactions between coding and noncoding RNAs during human
erythroid differentiation. Front Med 2016; 10(3):297-310.

[22] Yang Y, Wang H, Chang KH, Qu H, Zhang Z, Xiong Q *et al*. Transcriptome
dynamics during human erythroid differentiation and development. Genomics
2013; 102(5-6):431-441.

[23] Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 2014; 30(15):2114-2120.

[24] Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and
bias-aware quantification of transcript expression. Nat Methods 2017; 14(4):417-
419.

[25] Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics
2010; 26(1):139-140.

[26] Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W *et al*. limma powers
differential expression analyses for RNA-sequencing and microarray studies.
Nucleic Acids Res 2015; 43(7):e47.

[27] Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd *et al*.
Comprehensive Integration of Single-Cell Data. Cell 2019; 177(7):1888-1902
e1821.

[28] Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K *et al*. Fast,
sensitive and accurate integration of single-cell data with Harmony. Nat Methods
2019; 16(12):1289-1296.

[29] Schulz VP, Yan H, Lezon-Geyda K, An X, Hale J, Hillyer CD *et al*. A Unique
Epigenomic Landscape Defines Human Erythropoiesis. Cell Rep 2019;
28(11):2996-3009 e2997.

[30] Stoeckert CJ, Jr., Salas F, Brunk B, Overton GC. EpoDB: a prototype database
for the analysis of genes expressed during vertebrate erythropoiesis. Nucleic Acids
Res 1999; 27(1):200-203.

[31] Goh SH, Lee YT, Bouffard GG, Miller JL. Hembase: browser and genome portal
for hematology and erythroid biology. Nucleic Acids Res 2004; 32(Database
issue):D572-574.

[32] Kingsley PD, Greenfest-Allen E, Frame JM, Bushnell TP, Malik J, McGrath KE
et al. Ontogeny of erythroid gene expression. Blood 2013; 121(6):e5-e13.

[33] Patnaik SK, Helmberg W, Blumenfeld OO. BGMUT: NCBI dbRBC database of
allelic variations of genes encoding antigens of blood group systems. Nucleic

471   Acids Res 2012; 40(Database issue):D1023-1029.

472 [34] An X, Mohandas N. Disorders of red cell membrane. Br J Haematol 2008;
473   141(3):367-375.

474 [35] Bagger FO, Sasivarevic D, Sohi SH, Laursen LG, Pundhir S, Sønderby CK *et al*.
475   BloodSpot: a database of gene expression profiles and transcriptional programs for
476   healthy and malignant haematopoiesis. Nucleic Acids Research 2016;
477   44(D1):D917-D924.

478 [36] Bagger FO, Kinalis S, Rapin N. BloodSpot: a database of healthy and malignant
479   haematopoiesis updated with purified and single cell mRNA sequencing profiles.
480   Nucleic Acids Research 2019; 47(D1):D881-D885.

481 [37] Masuda T, Wang X, Maeda M, Canver MC, Sher F, Funnell AP *et al*.
482   Transcription factors LRF and BCL11A independently repress expression of fetal
483   hemoglobin. Science 2016; 351(6270):285-289.

484 [38] Lee HY, Gao X, Barrasa MI, Li H, Elmes RR, Peters LL *et al*. PPAR-alpha and
485   glucocorticoid receptor synergize to promote erythroid progenitor self-renewal.
486   Nature 2015; 522(7557):474-477.

487 [39] Wan Y, Zhang Q, Zhang Z, Song B, Wang X, Zhang Y *et al*. Transcriptome
488   analysis reveals a ribosome constituents disorder involved in the RPL5
489   downregulated zebrafish model of Diamond-Blackfan anemia. BMC Med
490   Genomics 2016; 9:13.

491 [40] Deena Iskander GW, Guanlin Wang, Guanlin Wang. Single-cell profiling of
492   human bone marrow progenitors reveals mechanisms of failing erythropoiesis in
493   Diamond-Blackfan anemia. Science translational medicine 2021;
494   13(610):eabf0113.

496 **Figure legends**

497 **Figure 1 Construction process of EryDB**

498 Transcriptome data for erythroid cells were obtained from Gene Expression Omnibus
499 (GEO), Sequence Read Archive (SRA), and Genome Sequence Archive (GSA).
500 Metadata were downloaded from GEO, SRA, GSA and manually curated by experts in
501 erythropoiesis.

503 **Figure 2 Dataset query strategies in EryDB**

504 **A.** The Erythroid Differentiation category allows datasets to be filtered by cell type. **B.**
505 The Genes category allows datasets to be filtered by selecting a gene of interest. **C.**
506 The Compounds category allows datasets under the action of a certain molecular
507 compound to be obtained. **D.** The Diseases category allows datasets to be obtained by
508 selecting a specific disease. **E.** Query results are presented as a table. **F.** Navigation
509 bar on the left-hand side of the "Search" page lists the conditions for selecting target
510 datasets. **G.** Detailed meta information included in the dataset overview page.

512 **Figure 3 Statistical analysis function for bulk RNA-seq data**

513 **A.** Expression profile of hemoglobin subunit gene *HBA1* in dataset GSE128268 as an

example to demonstrate the functions of EryDB. **B.** Principal component analysis of samples in the GSE128268 dataset. **C.** Volcano plot of differentially expressed genes (DEGs) between samples from the CFU-E and BFU-E groups in GSE128268. **D.** Enrichment analysis (biological process terms of Gene Ontology) of DEGs between samples from the CFU-E and BFU-E groups in GSE128268. **E.** Screenshot of the Erythroid Atlas page. Datasets for different differentiation stages can be explored on this page. **F.** Expression of the transcription factor gene *RUNX2* among different developmental stages across multiple datasets of bone marrow tissue as an example. The tissue of interest can be selected from the drop-down box on the left-hand side.

Figure 4 Statistical analysis function for scRNA-seq data

A. Interactive two-dimensional (2D) t-SNE plot of the single cells in the W9 group in dataset CRA002443 as an example to demonstrate the functions of EryDB; cell clusters are color-coded. **B.** Interactive three-dimensional UMAP plot of the single cells in the W9 group; cell clusters are color-coded. **C.** Expression of the cyclin gene *CCNB1* among erythroblasts as an example. Panels A, B, and C illustrate the usage of the Visualization & Feature module. **D.** Expression of marker genes at different cell stages. **E.** Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) functional and pathway enrichment for genes in the erythroblasts cluster. The cluster and enriched type can be selected in the drop-down boxes in the Marker & Enrichment module. **F.** Pseudotime trajectories of cell differentiation visualized in 2D. **G.** Cell differentiation trajectories displayed by cell type. Panels F and G show the usage of the Differentiation Trajectory module. **H.** Interaction of ligands and receptors between different cell types. Interaction of the top 40 ligands and receptors pairs is shown. The communication type can be defined in the drop-down boxes in the Cell–Cell Interaction module. **I.** Cell–Cell communication is visualized as a river diagram. **J.** Interaction of the TGFβ signaling pathway between cells as an example. **K.** Contribution of specific ligand–receptor pairs in the TGFβ signaling pathway. Panels I, J, and K illustrate the usage of the Cell–Cell Communication module.

Tables
Table 1. Statistics of the contents of EryDB

Supplementary material
Supplementary Table 1. All erythroid-related datasets in EryDB
Supplementary Table 2. Content of the datasets that can be searched by users
Supplementary Table 3. Other publicly available erythropoiesis-related databases

A

HSC — MPP — CMP — MEP — BFU-E — CFU-E — Pro-E — Baso-E — Poly-E — Ortho-E — Retic — RBC

B

GATA1 KLF1 WTAP BCL11A TFAM ALAS2 METTL14 UCA1 ALAS1 PHB2 LINCRED1 PTBP1 ERYTHA TAL1 PI4KAP2 METTL3 ATF4 REDRUM HMGA2 ZBTB7A GATA1 TMCC2 BCL11A BLOODLINC RUNX1C GRB10 CD169 ... (word cloud of gene names)

C

- TRβ agonists
- ERBB inhibitors
- Iron
- IDH2 inhibitors
- BET bromodomain inhibitors
- PPAR-α agonists
- HbF regulators
- Corticosteroids
- Erythropoietin

D

Number of datasets

Disease	0	1	2	3	4	5	6
Aplastic anemia				3			
Fanconi anemia		1					
Hemoglobinopathies				3			
Epo-resistant anemia		1					
Diamond-Blackfan anemia							6

E

Dataset	Species	Tissue	Experiment type	Title	Omics type	Sample count	Cell count
Input Dataset 🔍	Choose Species ▾	Choose Tissue ▾	Choose··· ▾	Input Title 🔍	Choose··· ▾	Choose··· ▾	Choose··· ▾
PMID34644394	*Homo sapiens*	Cord Blood (CB)	*In vivo*	RNA regulation during erythroid differentiation of hematopoietic stem cells	Bulk RNA-seq	30	
PMID31597962	*Homo sapiens*	Embryonic Fetal Liver (EMFL)	*In vivo*	Decoding human fetal liver haematopoiesis	scRNA-Seq	17	113, 063

F

Choose cell type:
Please select cell type ▾

Choose reported gene:
Please select reported gene ▾

Choose compound type:
Please select compound type ▾

Choose disease type:
Please select disease type ▾

Choose species:
Please select species ▾

Choose tissue:
Please select tissue ▾

Choose experiment type:
Please select experiment type ▾

Choose omics type:
Please select sequence type ▾

G

Direct targets of pSTAT5 signalling in erythropoiesis

Dataset: GSE94301
Organism: *Mus musculus*
Source: Cell line (J2E)
Experiment type: *In vitro*
Omics type: Expression profiling by high throughput sequencing; Genome binding/occupancy profiling by high throughput sequencing
Status: Public on Sep 06, 2017
Summary: Identification of direct transcriptional targets of EPO-JAK2-STAT5 signalling in erythroid cells
Organization name: Monash University
Platforms: GPL18635 Ion Torrent Proton (Mus musculus)
Citation(s): 28732065 30127368
Samples (4) and group:

Accession No ⇅	Sample name ⇅	Organism ⇅	Tissue ⇅	Cell type ⇅	Source ⇅	Growth mode ⇅	Group ⇅
GSM2472756	J2E 4sU-RNA-seq plus epo rep1	*Mus musculus*	Cell line (J2E)		Treatment: treated with erythropoietin	*In vitro*	plus_epo
GSM2472757	J2E 4sU-RNA-seq plus epo rep2	*Mus musculus*	Cell line (J2E)		Treatment: treated with erythropoietin	*In vitro*	plus_epo
GSM2472758	J2E 4sU-RNA-seq minus epo rep1	*Mus musculus*	Cell line (J2E)		Treatment: untreated	*In vitro*	minus_epo
GSM2472759	J2E 4sU-RNA-seq minus epo rep2	*Mus musculus*	Cell line (J2E)		Treatment: untreated	*In vitro*	minus_epo

Total 4 items   ‹ 1 ›   10/page ▾   Goto 1

Table 1 Statistics of the contents of EryDB

Species	Tissue	Omics type	No. of studies	No. of datasets	No. of samples	No. of single cells
Homo sapiens	Fetal tissue (e.g., liver, spleen, kidney)	scRNA-seq	3	3	98	434,156
		Bulk RNA-seq	5	5	80	/
	Bone marrow	scRNA-seq	6	7	130	267,566
		Bulk RNA-seq	3	6	88	/
	Peripheral blood	scRNA-seq	1	1	2	26,740
		Bulk RNA-seq	10	7	474	/
	Induced pluripotent stem cells	scRNA-seq	2	2	731	27,666
		Bulk RNA-seq	1	1	5	/
	Cord blood	scRNA-seq	2	2	7	29,920
		Bulk RNA-seq	13	13	142	/
	Cell line	Bulk RNA-seq	6	6	82	/
	Other (unknown)	Bulk RNA-seq	1	1	6	/
Mus musculus	Fetal tissue (e.g., liver, yolk sac, caudal half)	scRNA-seq	1	1	6	353,528
		Bulk RNA-seq	18	22	542	/
	Bone marrow	scRNA-seq	2	1	9	28,224
		Bulk RNA-seq	10	15	1,162	/
	Cell line	Bulk RNA-seq	4	6	46	/
	Spleen	Bulk RNA-seq	4	6	43	/
	Embryo	Bulk RNA-seq	3	3	24	/
	Heart	Bulk RNA-seq	2	1	6	/
Danio rerio	Fetal tissue	scRNA-seq	1	1	1	19,319
	Embryo	Bulk RNA-seq	7	6	93	/
	Kidney	Bulk RNA-seq	2	2	24	/
Sum			107	118	3,803	1,187,119