Group size modulates kinship dynamics and selection on social traits

Peng He[1,7], **Michael N. Weiss**[2,3], **Samuel Ellis**[2], **Daniel W. Franks**[4,5],

Michael A. Cant[6], **Darren P. Croft**[2,3] and **Rufus A. Johnstone**[1,7]

[1]Department of Zoology, University of Cambridge, Cambridge, CB2 3EJ, UK

[2]Centre for Research in Animal Behaviour, University of Exeter, Exeter, EX4 4QG, UK

[3]Center for Whale Research, Friday Harbour, Washington, 98250, USA

[4]Department of Biology, University of York, York, YO10 5DD, UK

[5]Department of Computer Science, University of York, York, YO10 5GH, UK

[6]Centre for Ecology and Conservation, University of Exeter, Cornwall, TR10 9FE, UK

[7]Correspondence: peng.he@outlook.de (PH), raj1003@cam.ac.uk (RAJ)

ABSTRACT

An individual's relatedness to its group may change with age due to demographic processes, and such kinship dynamics can shape age-linked social behaviours. Existing models, however, have focused almost exclusively on dispersal and mating, leaving the consequences of group size local variation unexplored. Here, we extend these models to incorporate such variation within a single, genetically connected meta-population. We then contrast predicted kinship dynamics and age-linked selection for helping/harming in coexisting smaller and larger groups in social mammal contexts, followed by exploring predictions under broader demographic contexts to clarify how demographic conditions, including local group size, influence patterns of female and male kinship dynamics. We show an individual's age-specific relatedness to others is higher, and changes faster (especially when younger), in smaller social mammal groups; consequently, smaller groups favour more extreme helping/harming (especially when younger) — in the context of bi-sexual philopatry with non-local mating (e.g., whales) that favours female

shift from harming to helping with age (which explains the evolution of menopause and post-reproductive helping), such shifts are earlier in smaller groups. Further explorations suggest that under group size local variation, kinship dynamics in a group reflect not only its own demographic conditions, but also those in coexisting groups of different size. Collectively, our study generates new insights into how female and male kinship dynamics emerge and vary under within-population group-size heterogeneity, and how such locally dynamic, varying kinship environments may be linked to observed variations in age-linked social traits, such as the timing of menopause in social mammals.

KEYWORDS: Group size variation | Inclusive fitness | Life history evolution | Menopause | Population asymmetric subdivision | Social evolution

INTRODUCTION

There is clear evidence that the mean relatedness of an individual to others in its social group may change systematically with age [1, 2]. Such kinship dynamics, which are sensitive to a species' demography, are predicted to influence sex- and age-specific social behaviours such as helping and harming [1-4]. For example, among social mammals, only humans and some toothed whales [5] exhibit a life history pattern of early reproductive cessation (compared to males), followed by an extended period of post-reproductive life during which they assist kin [3, 4, 6, 7]. This unusual life history pattern is associated with unusual kinship dynamics: females in these species typically experience an increase in their relatedness to others in their social groups as they age, a trend that has been invoked to help explain the evolution of menopause and post-reproductive helping [2-4, 6].

Despite growing interest in kinship dynamics, existing models have focused chiefly on how these dynamics are shaped by typical rates of female and male philopatry versus dispersal [3, 8], and of local versus non-local mating [2, 4]. It remains unknown how local variation in demographic factors, such as social group size, modulates local kinship dynamics following population asymmetric subdivision, and the sex- and age-linked trends in social behaviours expected under

such locally dynamic yet varying kinship environments. There is abundant evidence that group size varies within natural populations [9-18], and that it is often correlated with social behaviours such as the amount of assistance provided by non-breeding helpers in cooperative breeders [19-23]. But the impacts of group size local variation on kinship dynamics and predicted changes in social behaviour with sex and age remain unknown.

Here, we extend existing models of kinship dynamics [2, 4] to incorporate group size local deterministic variation within a genetically connected meta-population. We then use it to examine individuals' age-specific relatedness to their groupmates in coexisting smaller and larger groups, and explore how such emerging patterns of kinship dynamics influence age-linked selection for helping/harming (with a focus on social mammals), followed by discussing key challenges and uncertainties involved in empirically testing predictions on these age-linked social behaviours (using the killer whales *Orcinus orca* as an example). Finally, we explore our model in a broader range of demographic contexts to clarify the mechanisms through which demographic conditions, including group size local variation, influences emergent patterns of kinship dynamics females and males experience as they age. Together, our analyses provide new insights into how female and male kinship dynamics emerge and vary in coexisting local groups with deterministic differences in their sizes, and demonstrates a group-size-related mechanism that might be invoked to explain widespread variations in social traits in animal populations, such as the timing of reproductive cessation in social mammals [24, 25].

METHODS

Population demographic assumptions

Building on the baseline model of Johnstone and Cant [4] and Ellis et al [2], we consider an infinite, diploid, bisexual population, divided into discrete social groups [a.k.a. islands, demes or patches, 26, 27], a proportion u of which are of α-type consisting of $n_{f\alpha}$ female and $n_{m\alpha}$ male breeders, while the remaining fraction $1 - u$ are of β-type consisting of $n_{f\beta}$ female and $n_{m\beta}$ male breeders (the baseline model assumes all groups have the same number of breeders). That is, grouptype is defined solely by the total number of breeders in a group, with $n_{f\alpha} + n_{m\alpha} <$

$n_{f\beta} + n_{m\beta}$, such that α-type groups are persistently smaller while β-type groups are persistently larger. Such differences reflect deterministic (or structural) variation in the fixed numbers of breeders across coexisting grouptypes in a population, rather than their stochastic fluctuations [28] or statistical distributions. Also note that under the infinite island population model, inter-individual relatedness is generated within local groups (as it is consistently zero between groups). As in the baseline model, time proceeds in discrete steps, in each of which every female produces some very large number, p, of offspring (regardless of group size), a fraction x of which are females while the rest are males (fecundity and primary sex ratio do not influence predictions but are introduced for clarity — see appendix for details; the assumption of a very large number of offspring in each sex makes sure that available breeding vacancies in each group at each timestep are always fully claimed by successful offspring — see below, such that group sizes remain consistent over time). At each timestep, the probability that an offspring born in a group is sired by a male from the same group is m in the baseline model (i.e., a measure of a local male's mating success — with probability $1 - m$ the offspring is sired by a non-local male); in the context of group size local variation, this local siring probability is conditioned on grouptype in which the offspring is born, to account for differences in male numbers (see appendix for derivations). After reproduction, each female or male offspring disperses from its natal group (to a random group in the population) with probability d_f or d_m, respectively. Then, offspring in each group (both those born locally that did not disperse, together with immigrants from the rest of the population) compete in a fair lottery manner for the available breeding vacancies created by the deaths in their own sex in that group — at each timestep each female or male breeder dies with probability μ_f or μ_m, respectively; in the presence of group size local variation, the probability that a native offspring wins a breeding vacancy (i.e., being recruited to its natal group a new breeder) is also conditioned on grouptype in which it is born (see derivations in the appendix). Offspring that fail to obtain any breeding vacancy die, and the cycle then repeats.

Modelling kinship dynamics

Given above demographic assumptions, we derive (in the appendix) expressions for the expected relatedness of males and females of varying ages to others in the same social group — distinguishing between smaller (α-type) and larger (β-type) groups. In the results section, we

explore the predicted patterns of an individual's sex-age-specific average relatedness to its groupmates in populations in which the proportion of α-type groups is either low ($u = 0.1$), medium ($u = 0.5$) or high ($u = 0.9$) — when $u = 0$ or $u = 1$ group-size heterogeneity disappears and our model reduces to the baseline model in which all groups are uniformly larger or smaller, respectively. For simplicity, we assume that $n_{f\alpha} = n_{m\alpha} = 5$ and $n_{f\beta} = n_{m\beta} = 20$ throughout (i.e., a smaller group contains 10 individuals, while a larger group contains 40 individuals, with the same number of males and females in either grouptype). The mortality rate of female and male breeders at each timestep across their lifespan in both grouptypes is $\mu_f = \mu_m = 0.1$. To make our explorations biologically relevant, we consider the three typical dispersal and mating patterns of social mammals addressed by Johnstone and Cant [4] and Ellis et al [2], represented by (I) whales: *bisexual philopatry with non-local mating*, where $d_f = 0.15$, $d_m = 0.15$, and $m = 0$, (II) apes: *female-biased dispersal with local mating*, where $d_f = 0.85$, $d_m = 0.15$, and $m = .82$, and (III) typical mammals: *male-biased dispersal with local mating*, where $d_f = 0.15$, $d_m = 0.85$, and $m = .82$. To reveal the effects of group size variation on kinship dynamics, under each of these social mammal scenarios, we also show in the appendix for comparison the kinship dynamics in distinct, group-size homogeneous populations consisting of either all α-type or all β-type groups — as defined in the group-size 'heterogeneous' population, while keeping all other parameter values consistent. Besides, we also explore our model under other demographic contexts (including group size) to clarify how female and male kinship dynamics emerge and vary under group size local variation (see appendix for details).

Evaluating selective pressures

We use the derived sex-age-explicit relatedness to quantify the strength of selection on helping/harming in both grouptypes, using the neighbor-modulated inclusive fitness approach [2, 29]. We assume helping/harming is genetically determined by a mutant allele, and the analysis proceeds in three steps (full expressions are given in the appendix). We first derive sex-grouptype-specific reproductive values under our demographic assumptions; then, we construct fitness functions for a focal individual in each sex-grouptype class, incorporating its own fecundity and mortality as well as those of others; next, we combine these fitness functions with the sex-age-explicit relatedness to exmine the inclusive-fitness effect of the behaviour expressed

in given age, sex, and grouptype, provided it alters the fecundity or mortality of the focal individual and its groupmates (see fully details in the appendix).

More specifically, for the inclusive fitness calculations, we assume the behaviour confers a total marginal fecundity or mortality benefits (or costs) of magnitude b (say $0 < |b| \ll 1$, where $0 > b$ for helping while $b < 0$ for harming) on the local recipients (either sex), distributed equally among them, and confers a constant, marginal fecundity or mortality cost c on the actor ($0 < c \ll 1$; see details in the appendix). Then, the critical b^* at which the net inclusive fitness effect is zero can be derived, and we then use the $\frac{c}{b^*}$ as a measure of sex-age-specific selective pressure on the behaviour — the larger the positive value of $\frac{c}{b^*}$, the stronger the selection for helping, whereas the smaller the negative value of $\frac{c}{b^*}$, the stronger the selection for harming. To reveal how the prevalence of smaller groups in a population modulates selection, we explore populations with either low ($u = 0.1$), medium ($u = 0.5$) or high ($u = 0.9$) proportions of smaller groups. For comparison, we also show (in the appendix) the corresponding selective pressures in the counterpart, group-size homogeneous populations [i.e., as in 2] consisting entirely of smaller or entirely of larger groups (all other parameter values held constant).

RESULTS

Local variation in kinship dynamics

Across the whale, typical mammal, and the ape cases characterized by distinct patterns of female and male dispersal and prevalence of local mating, individuals' average relatedness to their group is generally higher in smaller than larger groups. Of greater interest is the prediction that the rate of change in relatedness with age is typically higher in smaller than larger social groups — and that such differences are more pronounced at younger than older ages (Fig. 1). Across these cases, intra-population variation in group size does not alter the general, qualitative patterns of kinship dynamics experienced by individuals of either sex [2], nor does the prevalence of smaller groups in the population. With given prevalence of smaller groups in the population, the differences in the patterns of kinship dynamics experienced by both sexes in

smaller versus larger groups are most pronounced when both sexes are philopatric and mate non-locally (i.e., the whale case). With given patterns of mating and dispersal, the more prevalent the smaller groups are in the population (i.e., a higher u), the larger the differences in the patterns of kinship dynamics experienced by both sexes in smaller versus larger groups (Fig. 1). The predicted patterns and variations of female and male kinship dynamics under broader demographic contexts (i.e., varying magnitude of group size contrast, sex ratio, mortality, the prevalence of local mating, dispersal, and the proportion of smaller groups in the population) are enclosed in the appendix (Fig. A4-A7).

Fig. 1 Predicted mean relatedness of an individual female (red) or male(blue) to others in its social group (smaller: 5 females and 5 males, solid lines; larger: 20 females and 20 males, dashed lines), as a function of its age (scaled relative to the mean lifespan under constant mortality $\mu_f = \mu_m = 0.1$ at each timestep). The columns in (A) or (B) show the three typical patterns of dispersal and mating that characterize social mammal systems [2, 4], referred to as: the 'whale' case (*bisexual philopatry with non-local mating*), the typical mammal case (*male-biased dispersal with local mating*), and the 'ape' case (*female-biased dispersal with local mating*), as detailed in methods. The rows show three different proportions of smaller groups in the population: low ($u = 10\%$), medium ($u = 50\%$), or high ($u = 90\%$).

Local variation in selection

We find that, across the whale, typical mammal, and the ape cases, selection for helping/harming is generally higher in smaller than larger groups, and the rate of change in selection with age is higher in smaller than larger social groups, especially at earlier than later life history stages (Fig. 2). Across these social mammal contexts, the presence of group size variation does not qualitatively alter the predicted patterns of selective pressures on the social behaviour [2], neither does the prevalence (u) of smaller groups in the population. In the whale and ape cases, we find that females in both smaller and larger groups experience a significant shift in selection — from being more harmful when younger to being more helpful when older, when the behaviour affects fecundity, and that such shifts occur earlier in smaller groups (Fig. 2-A). With given

prevalence of smaller groups in the population, the differences in the critical ages of these shifts

between small and large groups are less significant in the ape than in the whale case; however,

the impacts of the prevalence of smaller groups differs between the cases: in the whale case,

such differences in critical age are more marked when smaller groups are more prevalent, while

in the ape case such differences are most pronounced at low but least pronounced at

intermediate prevalence of smaller groups in the population (Fig. 2-B).

Fig. 2 Predicted strength of selection for helping (positive values) or harming (negative values) as a function of sex (female: red; male: blue), age (scaled relative to the mean lifespan under constant mortality $\mu_f = \mu_m = 0.1$ at each timestep), and grouptype (solid lines: smaller groups; dashed lines: larger groups), given predicted patterns of sex-grouptype-specific kinship dynamics, provided the behaviour affect the fecundity (A) or mortality (B) of an actor and the recipients. Dotted horizontal lines mark selectively neutral condition ($\frac{c}{b^*} = 0$) and dotted vertical lines in (A) mark the critical age (dots at curve intersections) at which selection on females shifts from favouring more harmful to favouring more helpful in smaller (5 females and 5 males) and larger (20 females and 20 males) groups. The columns in (A) or (B) show the three typical patterns of dispersal and mating that characterize social mammal systems [2, 4], referred to as: the 'whale' case (*bisexual philopatry with non-local mating*), the typical mammal case (*male-biased dispersal with local mating*), and the 'ape' case (*female-biased dispersal with local mating*), as detailed in methods. The rows show three different proportions of smaller groups in the population: low ($u = 10\%$), medium ($u = 50\%$), or high ($u = 90\%$).

DISCUSSION

Group size is a key component of social evolution [9, 11, 13, 15, 17, 30-33]. Building on existing models, we generated new insights into how patterns of female and male kinship dynamics emerge and vary in local groups following population asymmetric subdivision, and subsequently, how selection for age-linked social traits varies in such locally dynamic kinship environments.

Comparing distinct, group-size homogeneous populations across the whale, typical mammal, and ape contexts shows the consistency (Fig. A1) that individuals are more related to others in populations characterised by smaller groups [see also 31, 34]. Our results show that this expectation of higher relatedness in smaller social mammal groups extends to group-size heterogeneous populations (so does the expectation of stronger selection for helping/harming in smaller groups, Fig. A2). However, we argue that this correspondence is conditional rather than universal: the strength and even direction of local group size effects depend on sex, life history stage, sex ratio, and social systems collectively realized by patterns of dispersal and local mating. For example, in the whale and ape contexts, an older male's average relatedness to its groupmates becomes increasingly insensitive to group size (as differences between smaller and larger groups diminish with age); under unbiased, moderate dispersal and mortality between the sexes, combined with moderate local mating, males even exhibit consistently higher relatedness in larger groups (Fig. A4), a pattern contrary to the seemingly general expectation of higher relatedness in smaller social mammal groups. More fundamentally, we find that an individual's kinship ties to others within a local group reflects not only its own size, but also the demographic conditions experienced by the other grouptype (e.g., altering the demographies in one grouptype changes patterns of kinship dynamics in both grouptypes, Fig. A4-A5). Such population-scale coupling means that predictions of individuals' age-specific relatedness in populations with group size local variation cannot be obtained by interpolating those by existing models assuming no group size local variation (e.g., Fig. A3), because the underlying processes by which such coupling is played out differ between group-size homogeneous versus heterogeneous populations (e.g., Fig. A1-A2, A6-A7). In summary, our incorporation of group size local variation does more than extend previous group-size homogeneous models — it shows that within a genetically connected meta-population the effects of sex, age, dispersal, and mating on an

individual's average relatedness to its groupmates cannot be understood by examining each local grouptype in isolation whilst coexisting.

The qualitative patterns of kinship dynamics for both sexes across the social mammal cases by our model, and the way they relate to the underlying demographic processes, are the same for both smaller and larger groups. Differences in local kinship dynamics nevertheless result from the asymmetric impact of these demographic processes between smaller and larger groups. In the presence of group size local variation, a native offspring (either sex) in a larger group is relatively more likely to be sired by a local male and recruited as a new breeder in its natal group (see proofs in appendix); while higher probabilities of local recruiting and siring strengthen local relatedness (by increasing the chance of individuals sharing local ancestry), we show that an individual's relatedness to others is consistently lower in larger groups across the mammal cases. This apparent paradox arises because two opposing mechanisms act simultaneously: while lower local recruiting and siring means introducing relatively more unrelated lineages into smaller groups, reducing the probability of identity by descent (i.e., diluting local relatedness), smaller groups also contain fewer breeders, so two individuals are much more likely to trace their ancestry back to the same parents (i.e., stronger genetic drift, as the probability of lineage coalescence scales inversely with the numbers of breeders — see recursions in the appendix). Our results across social mammal contexts thus suggest that, when these opposing mechanisms are combined, the relatively stronger coalescence in smaller groups outweighs local relatedness dilution by relatively higher gene inflow (i.e., lower local recruiting and/or siring probabilities), yielding higher relatedness therein. Interestingly, such outweighing is consistent across these social mammal contexts, despite they are characterized by distinct patterns dispersal and local mating, whereas the opposite can be true under other demographic regimes (e.g., under moderate local mating and dispersal of the sexes, males in smaller groups are consistently less related to their groups, Fig. A4).

Our study also provide new insights into the theory of social life history evolution, by showing how locally varying kinship environments can generate and explain variations in age-linked trends in social behaviours, such as the timing of shifts from harming to helping (i.e., social life history traits such as menopause in social mammals [24, 25]). However, here we also highlight the empirical challenges of testing these evolutionary predictions in natural populations.

Particularly, a common yet key complication that comes into play is group size variation itself: natural group sizes often fluctuate across ecological and evolutionary timescales, making it difficult to identify consistently smaller from larger groups (especially in fission-fusion societies where group size and membership change frequently and dramatically), whereas with the scope focused on the qualitative, expected effects of smaller versus larger groups on local kinship dynamics and selection gradients for helping/harming, our model assumes persistent, deterministic variation in local group size (i.e., group size is treated as a local demographic state variable instead of a random variable). While the expected effects of group size on kinship dynamics may be detectable in clearly demographically and socially structured populations, inferring the more profound evolutionary causal link between group size and age-related trends in social traits is more challenging. We suggest that a key step is to identify the social organizational scales at which groups can be biologically meaningfully classified as qualitatively smaller or larger. For example, at broader scales, one might contrast smaller and larger subpopulations, whereas at finer scales within these, one could contrast smaller and larger social groups.

To illustrate such challenges, we draw on facts revealed by long-term census datasets from three killer whale populations. At the broadest social scale the data permits, comparisons among the northern resident, the southern resident, and the transient populations [24] shows that 1) resident populations typically form larger pods than the transients, and 2) transient females achieve 95% of their lifetime reproductive outputs earlier than females in the resident populations [24]. This pattern is qualitatively consistent with our prediction that smaller social groups should be associated with earlier selective shifts from harming to helping; however, inference at this scale is constrained not only by the very small sample size (i.e., 3 populations), which precludes strong statistical test, but also by the clear ecological differences between the resident and transient ecotypes, which confound any causal interpretation of group size effects. Thus, beyond small sample size, this example demonstrates the challenge of disentangling group size from correlated ecological and evolutionary factors. At a finer social scale, matrilines form the core social units within each population, while individuals from different matrilines associate into stable social clusters as pods [2, 35]. Using long-term, individually-explicit census data available from the Southern resident population comprising three stable pods [36], we examined associations between pod size, age class, and female reproductive strategy (see appendix for

details). Estimated age-class- and pod-specific differences in annual birth probability were small, with slightly higher predicted birth probabilities for younger than older mothers, particularly in smaller pods (Fig. A8); while this pattern is again qualitatively consistent with our prediction (smaller groups favour earlier shifts from female harming to helping), inference at this scale remains constrained by only three pods and restricted variation in pod size in a single population. Consequently, substantial uncertainty remains on how much these observations reflect underlying effects of age and pod size on reproductive strategies (though the dataset provides a foundation for future rigorous tests of our predictions).

In conclusion, while empirically testing the predicted patterns of age-linked trends in social behaviours in relation to group size remains challenging, we suggest that group size local variation is a fundamental component underlying the diverse kinship environments individuals of both sexes experience as they age in natural populations, and such variations may therefore help to explain widespread variations in social traits, including the evolution and timing of menopause in social mammals [24, 25].

DATA AVAILABILITY

The expressions for calculating sex- and age-specific relatedness and selective pressures on social behaviours are given in the appendix. All the code and data used are available at https://github.com/ecopeng/GroupSize_KinshipDynamics_Selection.

AUTHORS' CONTRIBUTIONS

PH and RAJ led the conceptualization, modelling, and writing for this study with supports and inputs from other co-authors. MNW, SE and DPC led the data curation, analysis, and interpretation. DWF, MAC, DPC and RAJ contributed to acquiring funding for this study. All authors gave final approval for the publication of this study.

COMPETING INTERESTS

We do not have conflicts of interest to disclose.

ACKNOWLEDGEMENTS

We thank Andy Gardner and two anonymous reviewers for their constructive comments, and the Natural Environmental Research Council for granting us the funding (No. NE/S010327/1) to conduct this study.

REFERENCES

1. Croft D.P., Weiss M.N., Nielsen M.L.K., Grimes C., Cant M.A., Ellis S., Franks D.W., Johnstone R.A. 2021 Kinship dynamics: patterns and consequences of changes in local relatedness. *P Roy Soc B-Biol Sci* **288**(1957). (doi:10.1098/rspb.2021.1129).

2. Ellis S., Johnstone R.A., Cant M.A., Franks D.W., Weiss M.N., Alberts S.C., Balcomb K.C., Benton C.H., Brent L.J.N., Crockford C., et al. 2022 Patterns and consequences of age-linked change in local relatedness in animal societies. *Nat Ecol Evol* **6**(11), 1766-1776. (doi:10.1038/s41559-022-01872-2).

3. Johnstone R.A., Cant M.A. 2008 Sex differences in dispersal and the evolution of helping and harming. *Am Nat* **172**(3), 318-330. (doi:10.1086/589899).

4. Johnstone R.A., Cant M.A. 2010 The evolution of menopause in cetaceans and humans: the role of demography. *Proc Biol Sci* **277**(1701), 3765-3771. (doi:10.1098/rspb.2010.0988).

5. Ellis S., Franks D.W., Nattrass S., Cant M.A., Bradley D.L., Giles D., Balcomb K.C., Croft D.P. 2018 Postreproductive lifespans are rare in mammals. *Ecol Evol* **8**(5), 2482-2494. (doi:10.1002/ece3.3856).

6. Croft D.P., Johnstone R.A., Ellis S., Nattrass S., Franks D.W., Brent L.J.N., Mazzi S., Balcomb K.C., Ford J.K.B., Cant M.A. 2017 Reproductive conflict and the evolution of menopause in killer whales. *Curr Biol* **27**(2), 298-304. (doi:10.1016/j.cub.2016.12.015).

7. Johnstone R.A., Cant M.A. 2019 Evolution of menopause. *Curr Biol* **29**(4), R112-R115. (doi:10.1016/j.cub.2018.12.048).

8. Cant M.A., Johnstone R.A. 2008 Reproductive conflict and the separation of reproductive generations in humans. *P Natl Acad Sci USA* **105**(14), 5332-5336. (doi:10.1073/pnas.0711911105).

9. Majolo B., Vizioli A.D., Schino G. 2008 Costs and benefits of group living in primates: group size effects on behaviour and demography. *Anim Behav* **76**, 1235-1247. (doi:10.1016/j.anbehav.2008.06.008).

10. Lehmann J., Korstjens A.H., Dunbar R.I.M. 2007 Group size, grooming and social cohesion in primates. *Anim Behav* **74**, 1617-1629. (doi:10.1016/j.anbehav.2006.10.025).

11. Brown C.R., Brown M.B. 1996 *Coloniality in the cliff swallow: the effect of group size on social behavior*, University of Chicago Press.

12. Crockett C.M., Eisenberg J.F. 1987 Howlers: variations in group size and demography. In *Primate Societies* (eds. Smuts B.B., Cheney D.L., Seyfarth R.M., Wrangham R.W., Struhsaker T.T.), pp. 54-68, Chicago University Press.

13. Krause J., Ruxton G.D. 2002 *Living in groups*. New York, Oxford University Press.

14. Alexander R.D. 1974 The evolution of social behavior. *Annu Rev* **5**(1), 325-383. (doi:10.1146/annurev.es.05.110174.001545).

15. Brown C.R., Brown M.B., Roche E.A., O'Brien V.A., Page C.E. 2016 Fluctuating survival selection explains variation in avian group size. *P Natl Acad Sci USA* **113**(18), 5113-5118. (doi:10.1073/pnas.1600218113).

16. Safran R.J. 2004 Adaptive site selection rules and variation in group size of barn swallows: individual decisions predict population patterns. *Am Nat* **164**(2), 121-131. (doi:10.1086/422198).

17. Dornhaus A., Powell S., Bengston S. 2012 Group size and its effects on collective organization. *Annu Rev Entomol* **57**, 123-141. (doi:10.1146/annurev-ento-120710-100604).

18. Dunbar R.I.M., Mac Carron P., Shultz S. 2018 Primate social group sizes exhibit a regular scaling pattern with natural attractors. *Biol Letters* **14**(1). (doi:10.1098/rsbl.2017.0490).

19. Clutton-Brock T.H., Russell A.F., Sharpe L.L. 2004 Behavioural tactics of breeders in cooperative meerkats. *Anim Behav* **68**, 1029-1040. (doi:10.1016/j.anbehav.2003.10.024).

20. Garcia-Ruiz I., Taborsky M. 2022 Group augmentation on trial: helpers in small groups enhance antipredator defence of eggs. *Biol Lett* **18**(10), 20220170. (doi:10.1098/rsbl.2022.0170).

21. Josi D., Heg D., Takeyama T., Bonfils D., Konovalov D.A., Frommen J.G., Kohda M., Taborsky M. 2021 Age- and sex-dependent variation in relatedness corresponds to reproductive skew, territory inheritance, and workload in cooperatively breeding cichlids. *Evolution* **75**(11), 2881-2897. (doi:10.1111/evo.14348).

22. Farabaugh S.M., Brown E.D., Hughes J.M. 1992 Cooperative territorial defense in the Australian Magpie, *Gymnorhina Tibicen* (Passeriformes, Cracticidae), a group-living songbird. *Ethology* **92**(4), 283-292. (doi:10.1111/j.1439-0310.1992.tb00966.x).

23. Pike K.N., Ashton B.J., Morgan K.V., Ridley A.R. 2019 Social and Individual Factors Influence Variation in Offspring Care in the Cooperatively Breeding Western Australian Magpie. *Front Ecol Evol* **7**. (doi:10.3389/fevo.2019.00092).

24. Nielsen M.L.K., Ellis S., Towers J.R., Doniol-Valcroze T., Franks D.W., Cant M.A., Weiss M.N., Johnstone R.A., Balcomb K.C., Ellifrit D.K., et al. 2021 A long postreproductive life span is a shared trait among genetically distinct killer whale populations. *Ecol Evol* **11**(13), 9123-9136. (doi:10.1002/ece3.7756).

25. Thomas F., Renaud F., Benefice E., De Meeus T., Guegan J.F. 2001 International variability of ages at menarche and menopause: patterns and main determinants. *Hum Biol* **73**(2), 271-290. (doi:10.1353/hub.2001.0029).

26. Rousset F. 2004 *Genetic structure and selection in subdivided populations*, Princeton University Press.

27. Wright S. 1931 Evolution in Mendelian populations. *Genetics* **16**(2), 0097-0159. (doi:10.1093/genetics/16.2.97).

28. Rousset F., Ronce O. 2004 Inclusive fitness for traits affecting metapopulation demography. *Theor Popul Biol* **65**(2), 127-141. (doi:10.1016/j.tpb.2003.09.003).

29. Taylor P.D., Frank S.A. 1996 How to make a kin selection model. *J Theor Biol* **180**(1), 27-37. (doi:10.1006/jtbi.1996.0075).

30. Kaufhold S.P., van Leeuwen E.J.C. 2019 Why intergroup variation matters for understanding behaviour. *Biol Letters* **15**(11). (doi:10.1098/rsbl.2019.0695).

31. Powers S.T., Lehmann L. 2017 When is bigger better? The effects of group size on the evolution of helping behaviours. *Biol Rev* **92**(2), 902-920. (doi:10.1111/brv.12260).

32. Bose A.P.H., Koch L., Dabernig-Heinz J., Grimm J., Sefc K.M., Jordan A. 2022 Patterns of sex-biased dispersal are consistent with social and ecological constraints in a group-living cichlid fish. *BMC Ecol Evol* **22**(1). (doi:10.1186/s12862-022-01980-4).

33. Peña J., Nöldeke G. 2018 Group size effects in social evolution. *J Theor Biol* **457**, 211-220. (doi:10.1016/j.jtbi.2018.08.004).

34. Dyble M., Clutton-Brock T.H. 2020 Contrasts in kinship structure in mammalian societies. *Behav Ecol* **31**(4), 971-977. (doi:10.1093/beheco/araa043).

35. Bigg M., Olesiuk P., Ellis G.M., Ford J., Balcomb K.C. 1990 Social organization and genealogy of resident killer whales (*Orcinus orca*) in the coastal waters of British Columbia and Washington State. *Rep Int Whal Comm* **12**, 383-405.

36. Wiles G.J. 2016 Periodic status review for the killer whale in Washington. Washington Department of Fish and Wildlife, Olympia, Washington, USA.