Full Title: **Cyfip2 controls the acoustic startle threshold through FMRP, actin**

polymerization, and GABA$_B$ receptor function.

Short Title: **Cyfip2 controls startle sensitivity via FMRP, Rac1 and GABA$_B$ receptors.**

Jacob C. Deslauriers[1], Rohit P. Ghotkar[1,3], Lindsey A. Russ[1,4], Jordan A. Jarman[1,5], Rubia M.

Martin[1,6], Rachel G. Tippett[1], Sureni H. Sumathipala[1], Derek F. Burton[1,7], D. Chris Cole[1], Kurt C.

Marsden*[1,2]

1. Department of Biological Sciences, North Carolina State University, Raleigh, North

9   Carolina, USA

2. Center for Human Health and the Environment (CHHE), North Carolina State University,

Raleigh, North Carolina, USA

3. Current address: Putnam Associates, Boston, Massachusetts, USA

4. Current address: Department of Pharmacology & Physiology, Georgetown University,

Washington D.C., USA

5. Current address: Department of Physiology and Biophysics, Boston University, Boston,

MA, USA

6. Current address: U.S. Environmental Protection Agency, Raleigh-Durham-Chapel Hill,

North Carolina, USA

7. Current address: Biogen, Durham, North Carolina, USA

*Corresponding author: kcmarsde@ncsu.edu

Author contributions: Conceptualization: JCD and KCM; Data curation: JCD, RAM; Formal

analysis: JCD, RAM, RGT, SHS, DCC, KCM; Funding acquisition: KCM; Investigation: JCD,

RPG, LAR, JAJ, RAM, RGT, SHS, DFB, DCC, KCM; Methodology: JCD, RAM, KCM; Project

administration: KCM; Supervision: JCD, DFB, DCC, KCM; Visualization: JCD, RAM, KCM;

Writing – original draft: JCD; Writing – review and editing: JCD, KCM

<u>Abstract</u>

Animals process a constant stream of sensory input, and to survive they must detect and

respond to dangerous stimuli while ignoring innocuous or irrelevant ones. Behavioral responses

are elicited when certain properties of a stimulus such as its intensity or size reach a critical

value, and such behavioral thresholds can be a simple and effective mechanism to filter sensory

information. For example, the acoustic startle response is a conserved and stereotyped

defensive behavior induced by sudden loud sounds, but dysregulation of the threshold to initiate

this behavior can result in startle hypersensitivity that is associated with sensory processing

disorders including schizophrenia and autism. Through a previous forward genetic screen for

regulators of the startle threshold a nonsense mutation in *Cytoplasmic Fragile X Messenger*

Ribonucleoprotein (FMRP)-interacting protein 2 (*cyfip2*) was found that causes startle

hypersensitivity in zebrafish larvae, but the molecular mechanisms by which Cyfip2 establishes

the acoustic startle threshold are unknown. Here we used conditional transgenic rescue and

CRISPR/Cas9 to determine that Cyfip2 acts though both Rac1 and FMRP pathways, but not the

closely related FXR1 or FXR2, to establish the acoustic startle threshold during early

neurodevelopment. To identify proteins and pathways that may be downstream effectors of

Rac1 and FMRP, we performed a candidate-based drug screen that indicated that Cyfip2 can

also act acutely to maintain the startle threshold branched actin polymerization and N-methyl D-

aspartate receptors (NMDARs). To complement this approach, we used unbiased discovery

proteomics to determine that loss of Cyfip2 alters cytoskeletal and extracellular matrix

components while also disrupting oxidative phosphorylation and GABA receptor signaling.

Finally, we functionally validated our proteomics findings by showing that activating GABA$_B$

receptors, which like NMDARs are also FMRP targets, restores normal startle sensitivity in

cyfip2 mutants. Together, these data reveal multiple mechanisms by which Cyfip2 regulates

excitatory/inhibitory balance in the startle circuit to control the processing of acoustic

information.

Introduction

To navigate their environments to find food and avoid predation, animals must be able to filter out extraneous stimuli but respond appropriately to salient ones, a process known as sensorimotor gating. Specific attributes of a stimulus can trigger a response; for visual stimuli the luminance, size, and speed of the stimulus determine if escape and reorientation responses are initiated [1–3]. Similarly, the intensity and frequency of acoustic stimuli determine whether a response is made [4,5]. One way in which animals can control their responses to sensory stimuli is by establishing a behavioral threshold such that when one or more of these stimulus attributes reaches a critical value a specific behavioral response is initiated. Behavioral thresholds are a fundamental mechanism of sensorimotor gating used across the animal kingdom to regulate a wide range of behavioral responses including both collective responses, such as fanning behaviors for hive climate regulation in bees [6,7] and shoaling behavior in fish [8,9], as well as individual responses to odors [10–13], tactile stimuli [14–17], changes in luminance and contrast of visual stimuli [1,18–21], and sound frequency and intensity in mammals and fish [5,22–24]. That behavioral over-responsiveness to visual, tactile, and acoustic stimuli is observed across a number of neuropsychiatric conditions including autism, anxiety, and schizophrenia [4,22,24–28] highlights the importance of setting such behavioral thresholds at an appropriate level. Yet our knowledge of the molecular mechanisms that both establish and maintain behavioral thresholds is limited.

Previously, to identify genes that regulate the threshold for initiating the acoustic startle response, a highly conserved behavior initiated following sudden loud sounds that may indicate danger [5,29–31], we conducted a standard 3-generation, ENU-based forward genetic screen in larval zebrafish [32]. We identified a set of five mutant lines that display acoustic startle hypersensitivity, and through whole-genome sequencing of the *triggerhappy* mutant line, we identified a causal, nonsense mutation in *cytoplasmic Fragile X Messenger Ribonucleoprotein (FMRP)-Interacting protein 2 (cyfip2).* Cyfip2 was first identified as an interactor of FMRP and

the elongation initiation factor 4E (eIF4E), through which it participates in translational

repression of many target transcripts [33,34]. Cyfip2, but not the closely related Cyfip1, can also

bind the Fragile X-related proteins FXR1 and FXR2, but the function of these interactions is

unknown [33]. Additionally, Cyfip2 interacts with the activated form of the small Rho GTPase

Rac1, and it is a member of the WAVE Regulatory Complex (WRC) in which it helps regulate

Arp2/3 activation and branched actin polymerization [35–41]. Cyfip2 is vital for proper neuronal

migration and cell movement, axonal growth and guidance, as well as synapse formation and

function in flies, mice, and zebrafish [32,33,37,42–47]. Homozygous *cyfip2* mutations are

embryonically lethal in mammals and fatal after 7-8 days post-fertilization (dpf) in zebrafish

[32,42,44,48]. Despite its key role in multiple aspects of neurodevelopment, the links between

how Cyfip2 regulates RNA translation, actin polymerization, and behavior have not been

defined. Our previous work demonstrated that loss of Cyfip2 causes acoustic startle

hypersensitivity that is reversible upon transgenic expression of GFP-tagged Cyfip2, alters the

morphology but not the electrophysiological properties of the startle command-like Mauthner

cells (M-cells), and causes hyperexcitability of the spiral fiber neurons (SFNs), a set of hindbrain

excitatory interneurons that project to the M-cell axon hillock [32]. It is unclear, however,

whether Cyfip2 acts via Rac1-mediated actin polymerization or through FMRP-mediated

translational repression to control the startle threshold. Furthermore, the downstream molecular

changes that directly modulate the function of the startle circuit have not been identified.

In this study we used an inducible rescue approach in *cyfip2* mutant zebrafish larvae to

demonstrate that both Cyfip2's Rac1 and FMRP interactions are required for establishing the

acoustic startle threshold during early neurodevelopment. Using CRISPR-Cas9 gene

knockdown we find that FXR1 and FXR2 are dispensable for startle regulation and that Cyfip2

acts through FMRP. Furthermore, with a candidate-based pharmacological approach we show

that Cyfip2 mediates Arp2/3-induced branched actin polymerization and may modulate N-

methyl-D-aspartate receptors (NMDARs) to alter neuronal function in the acoustic startle circuit.

Finally, we performed discovery proteomics to define molecular pathways disrupted by loss of

Cyfip2 *in vivo*. Our results indicate roles for Cyfip2 in mitochondrial function, oxidative

phosphorylation, and inhibitory Gamma-Aminobutyric Acid (GABA) receptor signaling. We

confirmed the functional importance of this last finding using the $GABA_B$ receptor agonist

baclofen, which rescues *cyfip2* mutants' hypersensitivity. Together these data establish a novel

pathway that links Cyfip2, actin dynamics, RNA translation, and excitatory/inhibitory balance in

the control of acoustic responsiveness.

Results

<u>Cyfip2 establishes the acoustic startle threshold through both Rac1 and FMRP during early</u>

<u>neurodevelopment.</u>

Cyfip2 has four known protein-interaction domains [49] (Fig. 1A), and it can act through

Rac1 to promote actin polymerization (Fig. 1B) and FMRP to regulate RNA translation (Fig. 1C).

cyfip2^{p400} mutants have a single base pair transversion (nt1024; T to A) resulting in a premature

stop codon at amino acid position 343 (Fig. 1A) [32]. *cyfip2^{p400}* mutant zebrafish larvae (5 dpf)

were previously shown to display acoustic startle hypersensitivity that could be rescued by

expressing Cyfip2 at 30 hours post fertilization (hpf) using a stable heatshock-inducible

transgenic line, *Tg(hsp70:cyfip2-EGFP)* [32]. We replicate those findings here, using a 60-

stimulus assay consisting of 10 trials at each of 6 intensities with a 20 second (s) interstimulus

interval (ISI) to measure startle sensitivity (Fig. 1D). A 40-min heatshock at 30 hpf restores

normal sensitivity in transgenic (*Tg+*) but not in non-transgenic (*Tg-*) *cyfip2* mutants (Fig. 1D,E).

Previous studies have established that C179R and K723E amino acid substitutions prevent

Cyfip2 from binding with Rac1 and FMRP, respectively [34,39,45]. To determine if Cyfip2

engages Rac1-mediated actin regulation and/or FMRP/eIF4E-mediated translational repression

pathways to establish the acoustic startle threshold, we induced C179R (Δ*Rac1*) and K723E

(Δ*FMRP*) point mutations in the *Tg(hsp70:cyfip2-EGFP)* construct and created stable transgenic

lines for each (Fig. 1A). We expressed either wildtype or mutant (Δ*Rac1*; Δ*FMRP)* versions of

Cyfip2 in mutants at 30 hpf with a 40-minute heatshock at 38°C, followed by acoustic startle

testing at 5 dpf. While expression of wildtype *Tg(hsp70:cyfip2-EGFP)* at 30 hpf rescues mutant

hypersensitivity, expression of Δ*Rac1* or Δ*FMRP* versions of Cyfip2 in mutants was insufficient

to rescue mutant hypersensitivity (Fig. 1E).

cyfip2 mutants also display a number of kinematic defects in their performance of the

startle response [32]. To determine if Rac1 and FMRP binding are also required for Cyfip2 to

regulate startle kinematics, we analyzed startle latency, duration, head turn angle (C1 angle),

and total distance traveled during the response in *Tg-* and *Tg+* fish after the same 40-min

heatshock protocol at 30 hpf (Fig. S1). Heatshock induction of wildtype Cyfip2 restored normal

latency, duration, and C1 angle, but not distance traveled (Fig. S1). All kinematic defects

remained in *cyfip2* mutant larvae expressing Δ*Rac1*-Cyfip2 (Fig. S1), but expression of Δ*FMRP-*

Cyfip2 was sufficient to rescue startle duration and induced a trend toward rescue of startle

latency and C1 turn angle (Fig. S1). These data suggest that Rac1 binding is required for all

aspects of Cyfip2-mediated startle regulation, while FMRP binding is required to regulate the

startle threshold but is largely dispensable for startle kinematics.

147
Figure 1. Cyfip2 establishes the acoustic startle threshold through Rac1 and FMRP. (A) Cyfip2
protein interacting domain diagram of wildtype (top) and mutant (bottom) Cyfip2 proteins. Black
arrowheads indicate the positions of induced mutations in Cyfip2, eliminating the Rac1- (C179R) or
FMRP/eIF4E (K723E)-binding capacity of Cyfip2. (B) Cyfip2 actin regulatory pathway wherein Cyfip2
(orange) upon stimulation by Rac1-GTP triggers WAVE1 activation, Arp2/3-complex initiation and
branched actin nucleation. (C) Cyfip2 translational repression pathway in which Cyfip2, eIF4E (teal), and
FMRP (pink) along with the poly-A binding protein (PABP; gray), sequester neurodevelopmentally
important mRNAs from being translated. (D) Average startle frequency (%) after 10 trials at 13.6, 25.7,
29.2, 35.5, 39.6 and 53.6 dB for 5 dpf *cyfip2* siblings (+/) and mutant (-/-) larvae heatshocked at 30 hpf for
40 minutes at 38˚C. The average startle frequency curve for *cyfip2* siblings (+/; open circles, dashed line),
cyfip2 mutants (-/-; closed circles, solid line) and *cyfip2* mutants harboring the *Tg(hsp70:cyfip2-EGFP)*+
transgene (-/-; Tg+; closed circles, solid green line). (E) Sensitivity indices, calculated as the area under

the startle frequency curves, for 5 dpf *cyfip2* siblings and mutants, following a 40-minute heatshock at 30
hpf to express either wildtype (*Tg+*; green), Rac1- (ΔRac1+; blue) or FMRP/eIF4E- (ΔFMRP+; pink)
binding deficient versions of Cyfip2-EGFP. Comparisons were made to both non-transgenic (*Tg-*) and
non-heatshocked controls. All indices (mean ± SD) compared using a Kruskal-Wallis test with Dunn's
multiple comparisons correction; $p^{****} < 0.0001$. (F) Sensitivity indices for 5 dpf *cyfip2* sibling (+/) and
mutant (-/-) larvae following 1-cell stage injection with CRISPR-Cas9 and a single, *scrambled* guide RNA
(gRNA) or dual gRNA cocktails targeting *fmr1*, *fxr1,* or *fxr2*. *scrambled* gRNA injected (white bar, closed
circles); *fmr1* gRNA injected (dark gray bar closed circles); *fxr1* gRNA injected (medium gray bar; closed
circles); *fxr2* gRNA injected (light gray bar, closed circles). Comparisons were made both within genotype
and between genotypes by condition. All indices (mean ± SD) compared using an Ordinary one-way
ANOVA with Sidak's multiple comparisons correction; $p^* < 0.05$; $p^{**} < 0.01$; $p^{****} < 0.0001$.
171

One possible explanation for these results is that expression levels may differ between

the three heatshock transgenic lines. We therefore measured expression levels of each

transgenic Cyfip2-GFP protein 6 hours after a 40-min heatshock by fluorescence intensity (Fig.

S2A). The ΔRac1 and ΔFMRP lines displayed GFP expression that was not significantly

different than the wildtype *Tg+* line but which trended lower. To induce expression of wildtype

Cyfip2-GFP at levels more comparable to the ΔRac1 and ΔFMRP lines after 40-minute

heatshock, we delivered a 15-min heatshock at 30 hpf in the wildtype *Tg+* line. The 15-min

heatshock reduced peak Cyfip2-GFP expression to levels comparable to or below that of the

ΔRac1 and ΔFMRP lines, and this level of expression was also sufficient to rescue acoustic

startle sensitivity in *cyfip2* mutants (Fig. S2A-C). Thus, the level of transgene expression cannot

account for the failure of the ΔRac1 and ΔFMRP constructs to rescue startle phenotypes, and

these findings support our conclusion that Cyfip2 utilizes both Rac1- and FMRP-mediated

pathways to establish the acoustic startle threshold.

<u>Cyfip2 acts through FMRP but not FXR1/2 to establish the acoustic startle threshold.</u>

We previously found that FMRP is not required to establish a normal startle threshold, as

fmr1^{hu2787} mutants' startle sensitivity is unaffected [32]. Having established that the K723 residue

that Cyfip2 uses to bind FMRP is required for normal startle sensitivity, however, we

hypothesized that Cyfip2 may instead interact with other members of the Fragile X protein

family, Fragile X-related proteins 1 and 2 (FXR1/2), to establish the acoustic startle threshold

[33]. We designed pairs of CRISPR guide RNAs (gRNAs) targeting each of the *fmr1, fxr1* and

fxr2 genes. We injected *fmr1-, fxr1-* or *fxr2*-specific CRISPR-Cas9 cocktails into 1-cell stage

cyfip2 sibling and mutant embryos and measured acoustic startle sensitivity in these crispant

larvae at 5 dpf (Fig. 1F). Highly efficient CRISPR-induced mutagenesis was observed for all 3

genes, with 4 of the 6 gRNAs inducing edits as confirmed by PCR amplification and Sanger

sequencing (Fig. S3A-F). Quantitative PCR confirmed that mRNA expression of all three genes

was strongly reduced by CRISPR/Cas9 injection (Fig. S3G). FMRP crispants had significantly

increased startle sensitivity compared to larvae injected with a scrambled gRNA plus Cas9, and

sensitivity of FMRP crispants was even further increased in the *cyfip2* mutant background (Fig.

1F). However, startle sensitivity was unaltered in FXR1 or FXR2 crispants in either *cyfip2*

siblings or mutants (Fig. 1F). These data indicate that FMRP, but not FXR1/2, regulates the

startle threshold. The discrepancy between our hypersensitive *fmr1* crispants and prior analysis

of non-hypersensitive *fmr1^{hu2787}* mutants could be due to genomic adaptation that has been

reported in the *fmr1^{hu2787}* mutant line that may partially compensate for the loss of FMRP [50]. In

support of this, an independently created CRISPR-generated *fmr1* mutant line displayed

additional behavioral and developmental phenotypes not originally seen in the *fmr1^{hu2787}* line

[51]. *fmr1^{hu2787}* mutant larvae have been found to express other autism-related phenotypes,

however, such as altered social behavior and preference for reduced sensory stimulation [52],

as well as increased brain activity in response to acoustic stimulation [53]. Together our

conditional rescue and *fmr1* crispant data provide clear evidence that Cyfip2 acts in part through

FMRP to establish the acoustic startle threshold.

<u>Cyfip2 can acutely maintain the acoustic startle threshold through Rac1 and FMRP pathways.</u>

Our previous study found that Cyfip2 is important for both establishing the acoustic

startle threshold during early neurodevelopment, and for actively modulating the threshold later

in development between 4 and 6 dpf [32]. Here we sought to define a critical window for Cyfip2

expression in regulating the startle threshold using our heatshock transgenic lines. Our results

show that while a 40-minute heatshock to induce expression of wildtype Cyfip2 (*Tg+*) at 30 hpf

is sufficient for behavioral rescue in 5 dpf *cyfip2* mutant larvae (Fig. 1D,E), a 40-min heatshock

at 2, 3, or 4 dpf fails to rescue *cyfip2* mutant hypersensitivity (Fig. S4). Heatshock-induced

expression at 5 dpf, 4 hours prior to testing, resulted in a trend toward rescue and a bi-modal

distribution with some *cyfip2* mutants remaining hypersensitive and a second population

showing restoration of normal sensitivity (Fig. 2A). These results are similar to our prior findings

[32], and they suggest that Cyfip2 can not only function during the development of the startle

circuit but can also actively maintain the circuit's threshold after it has formed. Next, to

determine if Cyfip2 employs both of its canonical pathways in maintaining the startle threshold

after 4 dpf, we expressed either $\Delta Rac1$ or $\Delta FMRP$ versions of Cyfip2 in *cyfip2* mutants at 5 dpf

with a 40-min heatshock followed by acoustic startle behavior testing 6 hours later. Similar to

what we observed with the developmental heatshock (Fig. 1E), neither expression of the $\Delta Rac1$

nor the $\Delta FMRP$ version of Cyfip2 at 5 dpf rescued acoustic hypersensitivity of *cyfip2* mutants

(Fig 2A). These findings suggest that Cyfip2 uses both its Rac1- and FMRP-mediated pathways

to both establish and dynamically maintain the acoustic startle threshold throughout

neurodevelopment.

Figure 2. Cyfip2 acutely regulates branched actin polymerization and NMDARs to establish the acoustic startle threshold. (A) Sensitivity indices for 5 dpf *cyfip2* sibling (+/) and mutant (-/-) larvae, following a 40-minute heatshock at 120 hpf (5 dpf) to express either wildtype (*Tg+*; green), Rac1- (Δ*Rac1+*; blue) or FMRP/eIF4E- (Δ*FMRP+*; pink) binding deficient versions of Cyfip2-EGFP. Comparisons were made to non-transgenic (*Tg-*), heatshocked sibling (+/) and mutant (-/-) controls. All indices (mean ± SD) compared using a Kruskal-Wallis test with Dunn's multiple comparisons correction; p-values listed; $p** < 0.01$, $p**** < 0.0001$. (B) Sensitivity indices for 5 dpf *cyfip2* wildtype (+/+; white bar) and heterozygous (+/-; gray bar) larvae, treated for 30 minutes on d5 with 5, 20 or 50 µM CK-869. Comparisons were made both within genotype and within condition. All indices (mean ± SD) compared using a Kruskal-Wallis test with Dunn's multiple comparisons correction; $p* < 0.05$; $p**** < 0.0001$. (C) Sensitivity indices for 5 dpf Tüpfel longfin (TLF) larvae treated for 30 minutes on d5 with the highest, non-lethal doses the formin antagonist (SMIFH2; 5 µM), PAK3 antagonist (IPA-3; 50 µM) and ROCK antagonist (GSK429286; 100 µM). Comparisons were made between respective treatments and the DMSO controls. All indices (mean ± SD) were compared using a Kruskal-Wallis test with Dunn's multiple comparisons correction; All comparisons made were non-significant (n.s.). (D) Sensitivity indices for 5 dpf *cyfip2* sibling (+/) and mutant (-/-) larvae, treated for 30 minutes on d5 with 100 or 500 µM MK-801. Comparisons were made both between genotypes within condition and between conditions by genotype. All indices (mean ± SD) were compared using an Ordinary one-way ANOVA with Tukey's multiple comparisons correction. $p** < 0.01$; $p*** < 0.001$; $p**** < 0.0001$.

255
256

<u>Cyfip2 maintains the acoustic startle threshold through branched actin polymerization.</u>

Regulation of the actin cytoskeleton is a vital cellular process that within the context of the nervous system is critical for cell migration and movement, synapse formation, function and plasticity, receptor anchoring and trafficking, as well as axon growth and guidance [54]. Given our findings that Cyfip2 can act through the Rac1-WAVE pathway to establish and modulate the acoustic startle threshold, we hypothesized that Cyfip2-mediated branched actin polymerization specifically modulates the startle threshold. To test this hypothesis, we incubated 5 dpf *cyfip2* heterozygous and wildtype larvae in the Arp2/3 antagonist CK-869 at 5, 20, or 50 µM for 30 minutes, followed by acoustic startle testing. In control conditions, *cyfip2* heterozygotes display startle sensitivity equal to that of wildtypes (Fig. 2B), and incubation at 5 µM did not significantly alter startle sensitivity of either heterozygotes or wildtypes (Fig 2B). At 20 µM, wildtype larvae are unaffected by CK-869, but *cyfip2* heterozygotes have significantly increased startle sensitivity compared to controls (Fig 2B). 50 µM CK-869 caused both *cyfip2* wildtype and heterozygous larvae to become significantly hypersensitive compared to controls, phenocopying *cyfip2* mutants. Thus, Arp2/3-mediated, branched actin polymerization is necessary for acutely maintaining the acoustic startle threshold. That *cyfip2* heterozygotes display hypersensitivity at 20 µM but wildtypes do not indicates that a single functional copy of *cyfip2* is insufficient to maintain normal startle circuit function when branched actin polymerization is limited by a moderate dose of CK-869. We found similar results when exposing larvae to another Arp2/3 antagonist, CK-666, with both *cyfip2* heterozygotes and wildtypes phenocopying mutant hypersensitivity at a 50 µM concentration (Table S2). These findings support the conclusion that Cyfip2-dependent Arp2/3-mediated branched actin polymerization is necessary to maintain the acoustic startle threshold.

The actin cytoskeleton is dynamic and requires the action of both branched and

unbranched actin regulatory pathways to maintain cellular structure and function. Unbranched

filamentous (F) actin is polymerized from globular actin monomers by dimeric complexes of

formin proteins, which bind at the barbed ends of new filaments and promote their elongation

[55–58]. Another form of actin regulation involves the action of cofilin, an actin severing protein

that cleaves existing filaments to create new barbed ends and increase the rate of actin turnover

within the cell [59]. To determine whether unbranched actin and severing pathways play a role

in regulating the acoustic startle threshold we incubated 5 dpf wildtype larvae in a formin

antagonist, SMIFH2, or the cofilin disinhibitors, IPA-3 and GSK429286, for 30 minutes followed

by acoustic startle testing. Treatment with 5 µM SMIFH2 or with 50 µM IPA-3 or 100 µM

GSK429286 did not significantly affect startle sensitivity in wildtype larvae (Fig 2C). We also

tested these drugs at higher concentrations, which were lethal after a 30-minute exposure, as

well as longer exposures at lower concentrations, which had no effect on startle sensitivity

(Table S1). Our data with the formin inhibitor SMIFH2 are in contrast to a recent finding showing

that Formin 2B morpholino knockdown caused a decrease in Mauthner cell-mediated fast startle

responses in zebrafish larvae [60]. That we did not observe any change in startle frequency

could be due to the acute nature of our pharmacological approach as opposed to the

morpholino-mediated developmental knockdown of Formin 2B. Our findings suggest that acute

perturbations to unbranched actin filaments and actin turnover do not play a significant role in

regulating the acoustic startle threshold. Altogether, these data further support our conclusion

that Cyfip2-mediated, branched actin polymerization is a key pathway for acutely maintaining

the acoustic startle threshold.

Cyfip2 may regulate NMDA receptors to modulate the acoustic startle threshold.

While we have established that Cyfip2 mediates the establishment and maintenance of

the acoustic startle threshold through both branched actin and FMRP regulatory pathways, it is

unclear what molecular mechanisms directly modulate the excitability of the startle circuit. To

identify molecules that may be downstream effectors of Cyfip2 in modulating activity of the

startle circuit, we conducted a candidate-based small-molecule screen with compounds

previously shown to alter startle sensitivity in wildtype zebrafish larvae (Table 1) [61]. In this

screen we incubated *cyfip2* wildtype, heterozygous, and mutant larvae in each compound for 30

min prior to and during acoustic startle testing. Consistent with previous findings, *N-*

phenylanthrinilic acid (NPAA; Cl^- channel antagonist), Meclofenamic acid (MA; K^+ channel and

gap junction antagonist), Phenoxybenzamine (POBA; alpha-adrenergic receptor and calmodulin

antagonist), Etazolate (ETAZ; phosphodiesterase 4 (PDE4) inhibitor), and MK-801 (N-methyl-D-

aspartate receptor (NMDAR) antagonist) all increased startle sensitivity in a dose-dependent

manner (Table 1). BMS204352, a different K^+ channel antagonist, did not alter acoustic startle

sensitivity at either 10 or 50 μM concentrations, and NSC-23766, a Rac1 antagonist, reduced

sensitivity in siblings at 100 μM, but not *cyfip2* mutants.

Compound	Concentration (μM)	Effect By Genotype		
		cyfip2(+/+)	*cyfip2(+/-)*	*cyfip2(-/-)*
NPAA (Cl-channel antagonist)	1	93.21% of Control, p > 0.99	110.21% of Control, p < 0.99	113.57% of Control, p > 0.99
	5	140.11% of Control, p = 0.9785	137.32% of Control, p = 0.1385	140.29% of Control, p > 0.2787
	10	**234.14% of Control, p** = 0.0038**	**201.29% of Control, p**** < 0.0001**	**155.06% of Control, p* = 0.0189**
MA (K+ channel/gap jxn. antagonist)	1	131.6% of Control, p > 0.99	11.87% of Control, p > 0.99	132.29% of Control, p = 0.5732
	5	137.24% of Control, p > 0.99	138.46% of Control, p = 0.0622	112.29% of Control, p > 0.99
	10	214.13% of Control, p = 0.0833	**169.81% of Control, p* = 0.0188**	**155.76% of Control, p* = 0.0288**
POBA (AAR/calmodulin antagonist)	1	102.9% of Control, p > 0.99	107.08% of Control, p > 0.99	101.04% of Control, p > 0.99
	10	136.17% of Control, p > 0.99	106.58% of Control, p > 0.99	99.01% of Control, p > 0.99
	50	**175.8% of Control, p* = 0.0123**		99.76% of Control, p > 0.99
ETAZ (PDE4 inhibitor)	1	141.79% of Control, p > 0.99	126.36% of Control, p > 0.99	121.74% of Control, p > 0.99
	10	168.6% of Control, p = 0.5472	**172.26% of Control, p* = 0.0137**	152.73% of Control, p = 0.3977

	50	**236.64% of Control, p** = 0.0029**	**179.99% of Control, p** = 0.0015**	**149.75% of Control, p* = 0.0105**
MK-801 (NMDAR antagonist)	100	120.19% of Control, p > 0.99		**145.93% of Control, p** = 0.0024**
	500	**212.69% of Control, p**** < 0.0001**		**163.26% of Control, p*** = 0.0002**
BMS-204352 (K+ channel antagonist)	10	79.48% of Control, p = 0.4269		89.55% of Control, p > 0.99
	50	76.08% of Control, p = 0.6759		108.74% of Control, p > 0.99
NSC-23766 (Rac1 antagonist)	100	**49.14% of Control, p*** = 0.0002**		81.60% of Control, p > 0.99
	200	86.26% of Control, p > 0.99		105.28% of Control, p > 0.99

320
Table 1. Cyfip2 may regulate NMDARs to control acoustic startle sensitivity. Mean startle index
comparisons, listed as percentage (%) of the mean startle index of vehicle-treated controls by *cyfip2*
genotype and drug concentration, for larvae treated with compounds targeting the indicated pathways
[61] to increase acoustic startle sensitivity. All significant differences (p < 0.05) are listed (**bold**) for
comparisons using a Kruskal-Wallis test and Dunn's multiple comparisons correction. NPAA (*N*-
phenylanthranilic acid); MA (meclofenamic acid); POBA (phenoxybenzamine); ETAZ (etazolate).

To determine whether any of the targeted pathways may be downstream of Cyfip2, we looked

for conditions in which there was a clear *cyfip2* genotype-specific effect on sensitivity. The

NMDA receptor blocker MK-801 showed the clearest such effect, with a low dose (100 µM)

elevating startle sensitivity only in *cyfip2* mutants but not siblings (Table 1). Therefore, Cyfip2-

mediated cytoskeletal and/or translational regulation may impact the expression and/or function

of NMDA receptors within the startle circuit to modulate the acoustic startle threshold.

Proteomic analysis reveals that Cyfip2-mediated regulation of GABA$_B$ receptors is critical for

startle sensitivity.

To complement our candidate drug screen with an unbiased approach to identify

proteins and molecular pathways regulated by Cyfip2, likely through its role in translational

regulation, we conducted a proteomic analysis of *cyfip2* wildtype, heterozygous, and mutant

larvae at 5 dpf. All larvae used were siblings and were genotyped by PCR and Sanger

sequencing and then pooled in groups of 30 per genotype and snap frozen with liquid nitrogen.

Three independent pools of 30 larvae were analyzed for each genotype. Protein lysates were

submitted to the Molecular Education, Technology and Research Innovation Center (METRIC)

at NC State University for protein digestion and LC-MS. Raw LC-MS files were processed and

quantified using MaxQuant (Max Planck Institute of Biochemistry) and imported into Perseus

software for transformation and identification of Differentially Expressed Proteins (DEPs) for

subsequent Ingenuity Pathway Analysis (IPA).

Comparative analysis of *cyfip2* heterozygous and mutant versus wildtype proteomes

identified a total of 221 differentially expressed proteins (DEPs) in heterozygotes and 127 DEPs

in mutants (Fig. S5A-B; Tables S7,S8). Cyfip2 was the most strongly downregulated protein in

mutants, providing a key validation of our unbiased approach (Fig. 3A,S5A; Tables S7,S8).

Cyfip2 was slightly but significantly downregulated in heterozygotes as well (Fig. S5B), providing

a basis for the sensitization of Cyfip2 heterozygotes to the actin inhibitor CK-869 (Fig. 2B).

Cyfip1 expression was not significantly altered in either genotype, indicating that it likely does

not act to compensate for the loss of Cyfip2 (Fig. S5A-B). 66 DEPs were shared between *cyfip2*

heterozygotes and mutants, while 155 and 61 DEPs were specific to each group, respectively

(Fig. S5C). The top 5 upregulated proteins identified in *cyfip2^{p400}* heterozygotes in descending

order included: microtubule actin cross-linking factor 1 (MACF1), acyl-CoA dehydrogenase

family member 11 (ACAD11), cullin 2 (CUL2), calcium channel, voltage dependent, L-type alpha

1S (CACNA1S) and ubiquitin specific peptidase 24 (USP24) (Fig. 3A; top, green). The top 5

downregulated proteins identified in *cyfip2^{p400}* heterozygotes in descending order included:

tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein theta or 14-3-3

protein theta (YWHAQ), SEC61 translocon subunit alpha 1 (SEC61A1), ribosomal protein S5

(RPS5), enolase 2 (ENO2), and ribosomal protein L18A (RPL18A) (Fig 3A; top, red). The top 5

upregulated proteins identified in *cyfip2^{p400}* mutants in descending order included: ACAD11,

mitochondrial NADH dehydrogenase 4 (MT-ND4), NIPBL cohesion loading factor b (NIPBL),

troponin C2 fast skeletal type (TNNC2), and USP24 (Fig 3A; bottom, green). The top 5

downregulated proteins identified in *cyfip2^{p400}* mutants in descending order included:

cytoplasmic FMR1-interacting protein 2 (CYFIP2), collagen type VI alpha 3 (COL6A3), stress-

induced phosphoprotein 1 (STIP1), thymosin beta (TMSB10/TMSB4X) and collagen type 1

alpha 1 (COL1A1) (Fig 3A; bottom, green). These changes highlight the diverse set of roles that

Cyfip2 plays, impacting translational machinery, metabolism, and the extracellular matrix.

IPA analysis of DEPs in *cyfip2* heterozygotes and mutants revealed disruption of

multiple pathways common to both genotypes. Notable disrupted pathways shared between

cyfip2 heterozygotes and mutants include oxidative phosphorylation, mitochondrial dysfunction,

EIF2 signaling, sirtuin signaling, eIF4/p70S6K signaling, 14-3-3-mediated signaling, and GABA

receptor signaling (Fig. 3B). Analysis of the most affected diseases and functions for each

genotype supports a role for Cyfip2-mediated regulation in neuromuscular disease, disorder of

the basal ganglia, dyskinesia, Huntington disease, breast cancer, familial encephalopathy, and

neurological disorders (Fig. 3C). These data provide further evidence of the central importance

of Cyfip2 for neural function.

Finally, we sought to confirm that these pathways play a functional role in regulating

startle sensitivity, and so we focused on GABA receptor signaling as the most likely to

contribute to the hyperexcitability of *cyfip2* mutant larvae. We applied agonists of both ionotropic

$GABA_A$ receptors (muscimol) and metabotropic $GABA_B$ receptors (baclofen) for 1 hour prior to

startle testing. Muscimol did not have a consistent impact on mutant hypersensitivity at 1, 10, or

100 µM (Fig. 3D), but baclofen induced a clear restoration of normal sensitivity in *cyfip2* mutants

at 10 µM (Fig. 3E). Importantly, baclofen application did not cause sedation at this concentration

as mutants were equally responsive as untreated siblings, and the same dose of baclofen did

not affect startle sensitivity in sibling larvae. Thus, Cyfip2 maintains a normal startle threshold by

promoting GABAergic inhibition through the activity of $GABA_B$ receptors. Together our findings

establish a set of molecular pathways downstream of Cyfip2 that enable the proper processing

of acoustic stimuli to limit sensory over-responsiveness.

Figure 3. Loss of Cyfip2 causes widespread proteomic changes and GABA_B receptor signaling is critical for startle sensitivity. (A) Bubble plots reporting the level of significance of the top 10 dysregulated proteins for both *cyfip2* heterozygous (top) and mutant (bottom) groups compared to wildtype controls. The size of the dot is proportional to the significance of the results while the color code represents the log₂ fold change; top five upregulated (green), and top five downregulated (red) proteins

are shown. (B-C) Heat maps displaying the impacted canonical pathways (B) and diseases and biological
functions (C) from IPA functional analysis. The blue-colored gradient indicates the degree of enrichment
for the listed pathways or diseases/functions, as represented by the – log of the P value for each
pathway, disease and/or function. (D-E) Sensitivity indices for 5 dpf *cyfip2* sibling (+/) and mutant (-/-)
larvae, treated for 60 minutes prior to testing with muscimol (D) or (E) baclofen. All indices (mean ± SD)
were compared using a one-way ANOVA with Sidak's multiple comparisons correction. p* < 0.05, p** <
0.01; p*** < 0.001; p**** < 0.0001.

Discussion

Through its ability to modulate both the actin cytoskeleton and protein translation, Cyfip2

is well-positioned to be a critical factor for many processes in neurodevelopment [33,34,37–

39,45,49]. Further highlighting its importance, Cyfip2 has been implicated in an array of

neuropsychiatric and other conditions, including schizophrenia [62], autism [49,63–65], binge

eating [66–68], obesity [69], amyotrophic lateral sclerosis (ALS) [70,71], Alzheimer's disease

[72], epilepsy [65,73–76], and cancer [77–82]. Here we focused on Cyfip2's role in a common

endophenotype of schizophrenia and autism, increased acoustic startle responsiveness

[4,24,27,28]. By combining conditional transgenesis, CRISPR/Cas9 gene editing,

pharmacology, and discovery proteomics in an *in vivo*, vertebrate model system, we found that

both actin and translational regulation pathways are required for Cyfip2 to establish and

maintain a normal startle threshold. Our data indicate that through these pathways Cyfip2

modulates both excitatory (NMDA receptor) and inhibitory (GABA$_B$ receptor) function to

establish and maintain proper sensory responsiveness.

Cyfip2 acts through both Rac1 and FMRP to establish the acoustic startle threshold.

The actin regulating function of Cyfip2-Rac1 interactions is essential for many

neurodevelopmental processes including neuronal outgrowth and maturation, synapse

formation and function, axon guidance and cell migration [38,44,45,83]. The functional

importance of Cyfip2-FMRP interactions, though less well characterized than Cyfip1-FMRP

interactions, are thought to similarly impact the expression of many key neurodevelopmental

proteins that are directly involved in axon growth, synapse maturation, and synaptic plasticity

[34,45,84–86]. Here we used a heatshock-inducible expression system to reveal that Cyfip2

requires the ability to interact with both Rac1 and FMRP to establish the acoustic startle

threshold early in neurodevelopment (Fig. 1E). The ability of wildtype Cyfip2 to restore normal

startle sensitivity in mutant larvae when expression is induced at 30 hpf (Fig. 1D,E), but not at

48, 72, or 96 hpf (Fig. S4) reveals some potential ways that it may affect the underlying neural

circuits. Prior to the rescue window, by 8-15 hpf the command-like Mauthner cells have been

specified [87] and begun extending their axons (17-18 hpf) and lateral dendrites (22-23 hpf)

[88,89]. During the rescue window from ~30-48 hpf, other neurons in the startle circuit continue

to migrate to their final positions in the ventral hindbrain, and the synaptic contacts within the

circuit begin to form and mature, including those between the auditory nerve, Mauthner cells,

and excitatory Spiral Fiber Neurons (SFNs) [89–91]. Actin dynamics would be required during

this time to facilitate neuronal migration, axon and dendrite growth, and synapse formation.

These processes would also require precisely regulated RNA translation through complexes like

Cyfip2-FMRP-eIF4E to produce the many proteins needed to establish these connections.

When we induced Cyfip2 expression at 5 dpf we observed a clear bimodal distribution with

some mutants remaining hypersensitive and a second population with normal sensitivity (Fig.

2A). This pattern was not observed when Rac1 or FMRP binding was abolished, suggesting that

both pathways are also needed for Cyfip2 to modulate the startle circuit in this acute context,

which would likely occur through changes in neuronal and/or synaptic function rather than

altered connectivity. That Cyfip2 expression between 48-96 hpf did not restore the startle

threshold in mutants could be due to insufficient levels, but together our conditional expression

experiments indicate that Cyfip2 is able to most reliably function when the circuit is in a less

mature state.

Our findings also show that Cyfip2-Rac1 but not Cyfip2-FMRP binding is required for the

performance of the startle response, as kinematic parameters including latency, turn angle, and

duration were largely restored to normal by wildtype and ΔFMRP versions of Cyfip2 but

remained altered in ΔRac1-Cyfip2 expressing cyfip2 mutants (Fig. S1). These data demonstrate

that the actin regulation and translation regulation functions of Cyfip2, while both are required in

some contexts, also have some non-overlapping roles. This is consistent with findings in

zebrafish larvae showing that Cyfip2's interaction with FMRP is dispensable but that its

interaction with the Wave Regulatory Complex (WRC), which like the Rac1-Cyfip2 interaction

regulates actin polymerization (Fig. 1B), is required for retinal ganglion cell (RGC) axons to

properly navigate to their targets in the contralateral optic tectum [45]. In the startle context, our

data showing that ΔFMRP-Cyfip2 drives a stronger rescue for turn angle and duration than for

latency suggest that the Cyfip2-FMRP translational regulation pathway contributes more to the

initial sensory processing of acoustic stimuli than the regulation of motor output in the spinal

cord. The actin regulatory function of Cyfip2, however, appears to be critical for all of Cyfip2's

known roles in the startle circuit.

Cyfip2-dependent startle threshold regulation requires FMRP but not FXR1/2.

Previously, we observed that mutants from the fmr1^hu2787 line have normal startle

sensitivity, suggesting that FMRP plays no role in regulating the startle threshold [32]. So here

we tested whether Cyfip2, which in contrast to the closely related Cyfip1 has the capacity to also

bind with the Fragile X-related proteins FXR1 and FXR2 [33], may instead rely on these binding

partners to modulate the startle threshold. Like FMRP, both FXR1 and FXR2 regulate RNA

translation [92,93] and are expressed in the brain during early vertebrate neurodevelopment,

though divergent expression patterns emerge for the FXR1/2 proteins in later development and

adulthood in most vertebrates [94–96]. By analyzing FMRP, FXR1, and FXR2 crispants,

however, we found that FXR1 and FXR2 are dispensable but that FMRP is required for normal

startle sensitivity (Fig. 1F). This data is consistent with the distinct expression patterns of the

FMRP, FXR1 and FXR2 proteins, as well as the inability of FXR1/2 to functionally compensate

for the loss of FMRP [94–96]. Similarly, in *Drosophila*, which have only one Fragile X protein

family member (dFMR1), re-expressing human FMRP (hFMR1), but not human FXR1 or FXR2,

in dFMR1 mutants is sufficient to specifically rescue aberrant neuronal phenotypes [97]. As

discussed above, the fact that *fmr1* crispants but not *fmr1^{hu2787}* mutants show startle

hypersensitivity may be due to genomic adaptation in the ENU-induced *fmr1^{hu2787}* line [50]. That

we observed that *fmr1* crispants show heightened startle sensitivity in the *cyfip2* mutant

background compared to *cyfip2* mutants alone (Fig. 1F) further strengthens our conclusion that

Cyfip2 and FMRP work cooperatively to regulate the startle threshold.

Cyfip2-dependent branched actin dynamics are required for maintaining the acoustic startle

threshold.

Our rescue experiments indicate that Cyfip2's actin regulatory function through its

binding with Rac1 is required during startle circuit development (Fig. 1E) and that this pathway

may also facilitate a more acute role for Cyfip2 in maintaining the startle threshold (Fig. 2A).

This conclusion is bolstered by our finding that inhibition of Arp2/3 with 20 µM CK-869 for 30

minutes prior to testing uncovers startle hypersensitivity in *cyfip2* heterozygotes but not

wildtypes (Fig. 2B). Thus in wildtypes, Cyfip2 must act acutely to facilitate actin polymerization

to maintain the startle threshold in the face of this challenge. Arp2/3-mediated F-actin nucleation

creates branched actin filaments, while formin-mediated nucleation produces unbranched

filaments [56–58,98]. Our data show that only branched actin nucleation is required for acute

maintenance of the startle threshold, while both formin activity and cofilin-mediated actin

filament severing play no acute role in regulating the startle threshold (Fig. 2C). Formin 2B has

been shown with morpholinos to be required developmentally for normal startle responsiveness,

however, and it appears to play a role in the growth of Spiral Fiber Neuron (SFN) axons [60].

SFNs are excitatory interneurons that receive input from the contralateral auditory nerve and

project their axons across the midline to the contralateral Mauthner cell, providing a key driving

force to initiate the startle response [99]. We previously found that SFNs, but not Mauthner cells,

have heightened excitability in *cyfip2* mutants [32], making them a likely place for Cyfip2 to

regulate the startle threshold. Functioning acutely, Cyfip2 may impact synaptic input onto SFNs,

and it is possible that inhibitory and/or excitatory synapses on SFNs may be modulated by

Cyfip2 to maintain the startle threshold. Cyfip1 and Cyfip2 are both enriched at excitatory

synapses and regulate dendritic complexity and spine maturation in mouse cortical neurons

[84,85]. Both Cyfip1 and Cyfip2 are also found at inhibitory postsynaptic sites in mouse

hippocampal neurons, and overexpression of either protein disrupts excitatory/inhibitory (E/I)

synaptic balance [86]. It is likely that Cyfip2 functions similarly in the zebrafish startle circuit to

regulate neuronal excitability, as our data implicate both excitatory (NMDA receptors; Fig. 2D)

and inhibitory (GABA receptors; Fig. 3B,E) pathways.

Cyfip2 may control sensory processing and other disease-related functions by regulating

neurotransmitter receptors, mitochondrial function, and/or cytoskeletal remodeling.

Our candidate drug screen to identify potential downstream effectors of Cyfip2 in

regulating the startle threshold builds on previous work showing that NMDA receptor function is

required for normal startle sensitivity (Fig 2D; Table 1) [61]. While our screen confirmed the

known roles of Cl$^-$ and K$^+$ channels, gap junctions, calmodulin, and PDE4 in regulating startle

sensitivity, only the NMDA receptor blocker MK-801 produced a *cyfip2* genotype-specific

response, indicating that Cyfip2 may impact NMDA receptor expression and/or function in the

startle circuit. The mRNAs of three critical NMDA receptor subunit genes – GRIN1, GRIN2A,

and GRIN2B – are all targets of FMRP-mediated translational regulation [100], providing a

potential mechanism for the enhanced sensitivity of *cyfip2* mutants to the NMDAR inhibitor MK-

801. Our unbiased proteomic analysis of *cyfip2* heterozygotes and mutants did not uncover

dysregulation in excitatory synaptic pathways compared to wildtypes, although this may be

because we analyzed protein lysates from whole larvae and thus may have diluted out any

changes in NMDA receptor expression in specific neuronal subpopulations. It could also be the

case that Cyfip2 modulates the membrane localization of NMDA receptors through actin-

mediated trafficking rather than impacting total expression levels.

Ingenuity Pathway Analysis (IPA) of our proteomic data revealed that inhibitory GABA

receptor signaling is significantly disrupted in *cyfip2* heterozygotes and mutants (Fig. 3B). We

confirmed that Cyfip2-mediated regulation of GABA receptor function plays a key role in the

startle threshold, showing that activation of $GABA_B$ but not $GABA_A$ receptors is sufficient to

rescue the startle hypersensitivity phenotype in *cyfip2* mutants (Fig. 3D,E). Cyfip2 most likely

modulates $GABA_B$ receptors through translational regulation via FMRP, as both $GABA_B$ receptor

transcripts in mouse (GABRAB1 and GABRAB2) are targets of FMRP [100]. Further

experiments are needed, though, to determine where and how Cyfip2 affects $GABA_B$ receptor

expression in the startle circuit. Our data are consistent with recent findings using rats in which

baclofen-mediated activation of $GABA_B$ receptors restored normal auditory processing in

Cntnap2 knockout animals [101]. $GABA_B$ receptors function both pre-synaptically to regulate

neurotransmitter release and post-synaptically to activate inward-rectifying K^+ channels that

cause hyperpolarization [102,103]. $GABA_B$ receptors are also expressed at high levels

throughout the auditory system [104], and baclofen treatment has been shown to improve social

avoidance in some individuals with autism [105–107]. It is currently unknown whether baclofen

affects sensory processing in clinical populations, however. Our data fit with a growing body of

evidence that $GABA_B$ receptors are important modulators of auditory function with direct clinical

applications.

The most significantly disrupted pathways in our proteomic analysis of *cyfip2*

heterozygotes and mutants were oxidative phosphorylation and mitochondrial dysfunction (Fig.

3B). It is unclear if these metabolic functions influence the activity of the acoustic startle circuit,

although these pathways are essential within neurons for neuronal development and plasticity,

cell death, axon extension and branching, and synaptogenesis [108,109], and so they may also

contribute to the many disease associations for Cyfip2 listed above. Our data also show that in

the absence of Cyfip2, mitochondrial proteins (ACAD11, MT-ND4) increase in abundance, and

cytoskeletal (TMSB10/TMSB4X) and extracellular matrix (ECM) proteins (COL6A3, COL1A1)

decrease in abundance (Fig 3A; Table S8). TMSB4X and TMSB10 both suppress actin

polymerization [110,111], so their downregulation in *cyfip2* mutants may reflect an attempt to

compensate for the loss of Cyfip2- and WRC-mediated actin polymerization. ECM collagens like

COL6A3 and COL1A1 are important for multiple aspects of neural development including axon

guidance [112,113], and so this may reflect another potential mechanism for Cyfip2's

developmental role in regulating the startle circuit. Our analysis of diseases and functions

impacted by the loss of Cyfip2 include multiple neurological and neuromuscular conditions (Fig.

3C). These findings reinforce the known associations between *cyfip2* and ALS [70,71] and

Alzheimer's disease [72], further underscoring the importance of Cyfip2 for neural function

beyond the startle circuit. Further work on the links between Cyfip2, its molecular effectors, and

the development, function, and maintenance of neural circuits will improve our understanding of

and ability to treat these varied conditions.

<u>Materials & Methods</u>

<u>Zebrafish Husbandry and Maintenance</u>

All animal use and procedures were approved by the North Carolina State University

Institutional Animal Care and Use Committee (IACUC). Zebrafish embryos were obtained from

the Zebrafish International Resource Center (ZIRC), the University of Pennsylvania, or

generated at North Carolina State University and raised in a recirculating housing system.

Animals were fed and housed at a 5 zebrafish/L density under a 14h:10h light-dark cycle at

28°C.

Embryos were generated by a male and female pair placed in a mating box

(Aquaneering) containing system water and artificial grass. The following morning, 2-3 hours

into the light cycle, embryos were collected in petri dishes containing E3 embryo media (5 mM

NaCl, 0.17 mM KCl, 0.33 mM CaCl2·2H2O, 0.33 mM MgSO4 in water). Embryos were

examined under a brightfield compound microscope for fertilization and proper development and

were kept in groups ≤ 65. Embryos were placed in a 29°C incubator on a 14h:10h light-dark

cycle. 50% of E3 media changed daily, and any embryos with gross morphological defects were

removed and euthanized.

DNA Extraction & Genotyping

Fin biopsies were obtained from adult fish anesthetized in 0.02% Tricaine (MS-222;

Fisher) in system water. Fin clips were taken using a razor blade to remove ~2-3 mm of tissue

from the tail fin and samples were immediately fixed in 100% MeOH. Larval samples were

individually fixed in 100% MeOH following behavioral testing. DNA was extracted using the

HotShot DNA lysis method which consisted of a tissue lysis with base solution (25mM NaOH,

0.2mM EDTA), sample incubation at 95°C for 30 minutes, and sample neutralization with

neutralizing solution (40mM Tris-HCl). *cyfip2*^{p400} fish were genotyped using either dCAPS PCR

and restriction digest with ApoI-HF [32] or the rhAmp SNP Genotyping System (IDT). rhAmp

SNP genotyping was carried out using *cyfip2* locus and allele specific primers (Table S3)

targeting the wildtype and *cyfip2*^{p400} alleles. Genotyping for *Tg(hsp70:cyfip2-GFP),*

Tg(hsp70:cyfip2-(C179R)-GFP) and *Tg(hsp70:cyfip2-(K723E)-GFP)* was accomplished by PCR

amplification with primers specific to GFP (Table S4) followed by agarose gel electrophoresis.

Molecular Cloning

Alternative *cyfip2* rescue constructs (ΔRac1; ΔFMRP) were generated from a pENTR

cyfip2-EGFP plasmid [32] using custom primers and the Q5 Site Directed Mutagenesis Kit

(NEB) to induce the desired C179R (ΔRac1) and K723E (ΔFMRP) mutations. Mutagenesis was

confirmed using restriction digest and Sanger sequencing (Table S3). LR Gateway Cloning

(ThermoFisher) was used to insert the altered *cyfip2*-EGFPs into the pDEST I-SceI hsp70

destination vector. Transgenic lines were created by microinjection into 1-cell stage embryos

with a transgenesis mix containing phenol red, I-SceI enzyme, and the pDEST I-SceI

hsp70:*cyfip2*-(C179R)-EGFP or pDEST I-SceI hsp70:*cyfip2*-(K723E)-EGFP plasmid.

Inducible Heatshock Rescue & Imaging

Inducible expression of *cyfip2-EGFP,* as well as C179R and K723E variants, was

initiated at 30 hpf by placing dechorionated larvae into 96-well plates and incubating at 38°C for

15 or 40 minutes [32]. Following heatshock, larvae were returned to Petri dishes, and given 4

days of recovery at 29°C. GFP fluorescence was confirmed between 4-6 hours post-heatshock

for startle experiments using a Nikon SMZ25 stereo microscope with a GFP bandpass filter and

Lumen 200 fluorescence illumination system. For day 5 heatshock rescue experiments, larvae

were given 4 hours of recovery at 29°C prior to startle sensitivity testing.

For imaging experiments, larvae were treated as above for transgene expression at 30

hpf and at 29°C for 1 hour recovered in petri dishes in groups ≤ 65. After 1 hour of recovery

fluorescence was verified, and larvae were visualized using the stereo microscope system

described above and larval images were captured at 1-, 3-, 6-, 18-, 24-, 30-, and 42-hours post-

heatshock using a Nikon DS-Qi2 monochrome microscope camera. Image analysis was

conducted using FIJI (ImageJ) analysis software to manually define ROIs encompassing the

entire larval body, excluding the eye and auto fluorescent yolk sac. Fluorescence intensity

values reflect the mean gray values recorded for respective ROIs.

Chemical Exposures

For all exposures, groups of 10-20 larvae (5 dpf) were incubated for specified periods of

between 30 minutes and 16 hours within 35 mm Petri dishes in 2 mL of each drug solution.

Drug solutions remained on larvae during startle testing for 30 minutes to 1-hour exposures. For

16-hour incubations larvae first received fresh E3 prior to testing. Following incubation, larvae

were placed on the 6x6 acrylic testing grid and run through the acoustic startle assay. CK-869,

CK-666, MK-801, *N*-phenylanthranilic acid (NPAA), meclofenamic acid (MA),

phenoxybenzamine (POBA), etazolate (ETAZ), BMS 204352, muscimol, and baclofen were

obtained from Sigma-Aldrich. SMIFH2, IPA-3, GSK429286 and NSC23766 were acquired from

Tocris through Fisher Scientific.

<u>Behavioral Assays</u>

All larvae were tested at 5 days post-fertilization (dpf) unless otherwise stated. On the

day of testing, embryos were thoroughly screened for developmental defects, and those with

gross morphological defects were removed prior to behavior testing. *cyfip2^{p400}* larvae without

inflated swim bladders were not discarded, as *cyfip2* mutant larvae fail to inflate their swim

bladders [32]. Larvae were adapted to the testing arena lighting and temperature conditions for

30 minutes prior to testing. As previously described, the behavioral testing system consists of a

36-well acrylic grid attached to an acoustic-vibrational shaker (Bruel-Kjaer), a photron mini UX-

50 camera, LED lighting, InfraRed illuminator, and an acrylic IR diffuser [5,32,61].

To test the acoustic startle response, 5 dpf larvae were presented with 60 total stimuli

with a 20 second interstimulus interval (ISI), with 10 pseudo-randomized trials at each of the

following 6 stimulus intensities: 13.6, 25.7, 29.2, 35.5, 39.6 and 53.6 dB. All stimuli were

calibrated using a PCB Piezotronics accelerometer (#355B04) and signal conditioner

(#482A21), and voltage outputs were converted to dB using the formula dB = 20 log (V/0.775)

[32].

<u>Behavioral Analysis</u>

All responses in the acoustic startle assay were tracked using FLOTE analysis software

[5,32,61]. Short latency C-bends (SLCs) were identified by FLOTE using defined kinematic

parameters (latency, turn angle, duration, and maximum angular velocity). Startle sensitivity was

calculated by measuring the SLC frequency at each of the six stimulus intensities during the 60-

stimulus startle assay. Sensitivity indices were defined as the area under the startle frequency

vs. stimulus intensity curve calculated using Prism software (GraphPad).

<u>Larval Sample Preparation for Proteomics.</u>

Larvae from incrosses of *cyfip2*^{+/p400} carriers were raised as described above. At 3 dpf, DNA was

extracted from larvae using the Zebrafish Embryonic Genotyping (ZEG) apparatus (DanioLab).

ZEG samples were used for PCR amplification and then submitted for Sanger sequencing to

determine the genotype at the *cyfip2*^{p400} locus. At 5 dpf larvae were sorted into pools of 30

larvae for each genotype: homozygous wild type, *cyfip2*^{p400} heterozygous, and homozygous

mutant. Samples were snap frozen in liquid nitrogen and stored at -80˚C. They were then

resuspended and lysed in 100 µL ammonium bicarbonate (ABC; pH 8) containing 1% sodium

deoxycholate (SDC) using a Branson SLPe Sonicator (40:0.15;4C) delivering two 20 second

pulses at 20% amplitude intensity separated by 10 seconds between pulses. Lysates were

centrifuged at 10,000 rpm for 5 minutes at 4˚C, and the supernatants were retained and

quantified using Pierce BCA protein quantification (ThermoFisher; Cat #: 23225) and an

IMPLEN NP80 nanophotometer. Lysates were submitted the same-day to the Molecular

Education, Technology and Research Innovation Center (METRIC) at NC State University for

protein digestion and LC-MS.

<u>Protein Digestion and LC-MS</u>

Each lysed sample was normalized to 200 µg of protein in 200 µL of ABC/SDC solution.

Disulfide reduction was conducted by adding 15 µL 50 mM DTT and incubating at 56˚C for 30

minutes. 200 µL of 8M urea in 0.1 M Tris buffer (pH 8) was added and samples were transferred

to Vivicon 30kD Molecular Weight Cut-off (MWCO) filters. Samples were centrifuged at 12,000 x

g for 10 minutes at 21˚C. 200 µL of 8 M urea in 0.1 M Tris buffer (pH 8) was added to the top of

each filter, as well as 64 µL 55 mM iodoacetamide (IAA) solution and samples were incubated

for 1 hour in the dark at room temperature. Samples were centrifuged at 12,000 x g for 20

minutes. 100 µL of 2 M urea, 10 mM $CaCl_2$ in 0.1 M Tris buffer (pH 8) was added and samples

were centrifuged at 12,000 x g for 20 minutes. The previous step was repeated twice. 100 µL of

0.1 M Tris buffer (pH 7.5) was added and samples were centrifuged at 12,000 x g for 20-45

minutes. This step was repeated twice, with a 1-hour centrifuge period on the final spin. 200 µL

of 0.02 µg/mL trypsin in 0.1 M Tris buffer (pH 7.5) was added, and samples incubated overnight

at 37˚C. Following protein digestion with trypsin, samples were placed into fresh microcentrifuge

reservoirs, and 50 µL of quench solution (0.001% zwittergent3-16 in water, 1% formic acid) was

added and samples centrifuged at 12,000 x g for 1 hour. 450 µL quench solution was applied to

each filter and samples were centrifuged at 14,000 x g for 1 hour. Solutions were dried using a

speedvac concentrator and samples were stored dry until LC-MS. Samples were reconstituted

in 100 µL of mobile phase A (98% water, 2% acetonitrile, 0.1% formic acid) and peptides were

quantified via Pierce BCA assay. Samples were normalized to the lowest peptide concentration

for every sample and nanoLC-MS was conducted using a Thermo Orbitrap Exploris 480. This

work was performed in part by METRIC at NC State University, which is supported by the State

of North Carolina.

<u>Proteomics data analysis</u>

Shotgun proteomics raw files were processed and quantified with MaxQuant (version 2.2.0.0).

Briefly, the built-in Andromeda search engine scored MS2 spectra against fragment masses of

tryptic peptides derived from a *Danio rerio* reference proteome containing 93,351 entries

including isoforms (UniProt, accessed March 22, 2019). Our database search required variable

modifications (methionine oxidation and N-terminal acetylation) and a fixed modification

(cysteine carbamido-methylation) along with a minimum peptide length of 7 amino acids and

limited the search space to a maximum peptide mass of 4600 Da and a maximum of two missed

cleavages. The false discovery rate was controlled with a target-decoy approach at less than

1% for peptide spectrum matches and protein group identifications.

<u>Bioinformatics</u>

Label-Free quantification (LFQ) intensities from MaxQuant were imported into Perseus software

(version 2.0.7.0) and transformed to logarithmic scale with base two. Missing values were

replaced with values from the normal distribution, reducing the distributions to a factor of "0.3″

(width) and down-shifting by "1.8″ standard deviations while simulating random values to

replace the missing values. This protein quantification was used to measure the fold-enrichment

between *cyfip2^{p400}* heterozygous/homozygous and *cyfip2* wildtype groups. Statistical

significance was calculated using a two-way Student t-test and FPR ($p<0.05$). Differentially

expressed proteins (DEPs) were submitted to ingenuity pathway analysis (IPA) to identify their

function, specific processes, and related enriched pathways/diseases.

Statistical Methods

All statistical analyses were performed using Prism (GraphPad). All data sets were tested for

normality in Prism. Subsequent parametric or non-parametric tests and post-hoc analyses were

performed using Prism, and significance values ($p < 0.05$) were reported.

Acknowledgements

We would like to thank Leonard Collins and Taufika Williams from METRIC at NC State

University for their assistance with proteomic analysis. A special thank you to Kimberly Scofield

and Kara Carlson for feedback on the manuscript.

Funding Disclosure

We are grateful for financial support from the National Institute for Neurological Disease and

Stroke (R01-NS116354 to K.C.M.) and for seed funds provided by North Carolina State

University's Center for Human Health and the Environment (CHHE) through a National Institute

of Environmental Health Science center grant (P30 ES025128).

Competing Interests

The authors declare that no competing interests exist.

References

1. Burgess HA, Granato M. Modulation of locomotor activity in larval zebrafish during light
adaptation. J Exp Biol. 2007;210: 2526–2539. doi:10.1242/jeb.003939

2. Vagnoni E, Lourenco SF, Longo MR. Threat modulates perception of looming visual
stimuli. Curr Biol. 2012;22: R826-7. doi:10.1016/j.cub.2012.07.053

3. Randlett O, Haesemeyer M, Forkin G, Shoenhard H, Schier AF, Engert F, et al. Distributed
Plasticity Drives Visual Habituation Learning in Larval Zebrafish. Curr Biol. 2019;29: 1337-
1345.e4. doi:10.1016/j.cub.2019.02.039

4. Geyer MA, Swerdlow NR, Mansbach RS, Braff DL. Startle response models of
sensorimotor gating and habituation deficits in schizophrenia. Brain Res Bull. 1990;25:
485–498. doi:10.1016/0361-9230(90)90241-q

5. Burgess HA, Granato M. Sensorimotor gating in larval zebrafish. J Neurosci. 2007;27:
4984–4994. doi:27/18/4984 [pii]

6. Weidenmüller A, Kleineidam C, Tautz J. Collective control of nest climate parameters in
bumblebee colonies. Anim Behav. 2002;63: 1065–1071. doi:10.1006/anbe.2002.3020

7. Weidenmuller A. The control of nest climate in bumblebee (Bombus terrestris) colonies:
interindividual variability and self reinforcement in fanning response. Behav Ecol. 2004;15:
120–128. doi:10.1093/beheco/arg101

8. Ward AJW, Sumpter DJT, Couzin ID, Hart PJB, Krause J. Quorum decision-making
facilitates information transfer in fish shoals. Proc Natl Acad Sci U S A. 2008;105: 6948–
6953. doi:10.1073/pnas.0710344105

9. Cresci A, De Rosa R, Agnisola C. Assessing the Influence of Personality on Sensitivity to
Magnetic Fields in Zebrafish. J Vis Exp. 2019. doi:10.3791/59229

10. Michel WC, Sanderson MJ, Olson JK, Lipschitz DL. Evidence of a novel transduction
pathway mediating detection of polyamines by the zebrafish olfactory system. J Exp Biol.
2003;206: 1697–1706. doi:10.1242/jeb.00339

11. Dudova I, Vodicka J, Havlovicova M, Sedlacek Z, Urbanek T, Hrdlicka M. Odor detection
threshold, but not odor identification, is impaired in children with autism. Eur Child Adolesc
Psychiatry. 2011;20: 333–340. doi:10.1007/s00787-011-0177-1

12. Tavassoli T, Baron-Cohen S. Olfactory detection thresholds and adaptation in adults with
autism spectrum condition. J Autism Dev Disord. 2012;42: 905–909. doi:10.1007/s10803-
011-1321-y

13.   Takahashi H, Tsuboi A. Olfactory avoidance test (mouse). Bio Protoc. 2017;7: e2153.
772   doi:10.21769/BioProtoc.2153

773   14.   Blakemore S-J, Tavassoli T, Calò S, Thomas RM, Catmur C, Frith U, et al. Tactile
774   sensitivity in Asperger syndrome. Brain Cogn. 2006;61: 5–13.
775   doi:10.1016/j.bandc.2005.12.013

776   15.   Kohashi T, Oda Y. Initiation of Mauthner- or non-Mauthner-mediated fast escape evoked
777   by different modes of sensory input. J Neurosci. 2008;28: 10641–10653.
778   doi:10.1523/JNEUROSCI.1435-08.2008

779   16.   DeLorey TM, Sahbaie P, Hashemi E, Li W-W, Salehi A, Clark DJ. Somatosensory and
780   sensorimotor consequences associated with the heterozygous disruption of the autism
781   candidate gene, Gabrb3. Behav Brain Res. 2011;216: 36–45.
782   doi:10.1016/j.bbr.2010.06.032

783   17.   Puts NAJ, Wodka EL, Tommerdahl M, Mostofsky SH, Edden RAE. Impaired tactile
784   processing in children with autism spectrum disorder. J Neurophysiol. 2014;111: 1803–
785   1811. doi:10.1152/jn.00890.2013

786   18.   Armington JC. The electroretinogram, the visual evoked potential, and the area-luminance
787   relation. Vision Res. 1968;8: 263–276. doi:10.1016/0042-6989(68)90014-x

788   19.   Campbell FW, Maffei L. The influence of spatial frequency and contrast on the perception
789   of moving patterns. Vision Res. 1981;21: 713–721. doi:10.1016/0042-6989(81)90080-8

790   20.   Yilmaz M, Meister M. Rapid innate defensive responses of mice to looming visual stimuli.
791   Curr Biol. 2013;23: 2011–2015. doi:10.1016/j.cub.2013.08.015

792   21.   Wang Y, Wu W, Zhang X, Hu X, Li Y, Lou S, et al. A Mouse Model of Visual Perceptual
793   Learning Reveals Alterations in Neuronal Coding and Dendritic Spine Density in the Visual
794   Cortex. Front Behav Neurosci. 2016;10: 42. doi:10.3389/fnbeh.2016.00042

795   22.   Bakker MJ, Tijssen MA, van der Meer JN, Koelman JH, Boer F. Increased whole-body
796   auditory startle reflex and autonomic reactivity in children with anxiety disorders. J
797   Psychiatry Neurosci. 2009;34: 314–322. Available:
798   https://www.ncbi.nlm.nih.gov/pubmed/19568483

799   23.   Han K, Chen H, Gennarino VA, Richman R, Lu H-C, Zoghbi HY. Fragile X-like behaviors
800   and abnormal cortical dendritic spines in cytoplasmic FMR1-interacting protein 2-mutant
801   mice. Hum Mol Genet. 2015;24: 1813–1823. doi:10.1093/hmg/ddu595

802   24.   Takahashi H, Nakahachi T, Stickley A, Ishitobi M, Kamio Y. Relationship between
803   physiological and parent-observed auditory over-responsiveness in children with typical
804   development and those with autism spectrum disorders. Autism. 2018;22: 291–298.
805   doi:10.1177/1362361316680497 [doi]

806   25.   Grillon C, Ameli R, Goddard A, Woods SW, Davis M. Baseline and fear-potentiated startle
807   in panic disorder patients. Biol Psychiatry. 1994;35: 431–439. doi:10.1016/0006-
808   3223(94)90040-x

26.   Grillon C, Davis M. Acoustic startle and anticipatory anxiety in humans: effects of monaural
right and left ear stimulation. Psychophysiology. 1995;32: 155–161. doi:10.1111/j.1469-
8986.1995.tb03307.x [doi]

27.   Chamberlain PD, Rodgers J, Crowley MJ, White SE, Freeston MH, South M. A potentiated
startle study of uncertainty and contextual anxiety in adolescents diagnosed with autism
spectrum disorder. Mol Autism. 2013;4: 31. doi:10.1186/2040-2392-4-31

28.   Kohl S, Heekeren K, Klosterkötter J, Kuhn J. Prepulse inhibition in psychiatric disorders--
apart from schizophrenia. J Psychiatr Res. 2013;47: 445–452.
doi:10.1016/j.jpsychires.2012.11.018

29.   Eaton RC, Bombardieri RA, Meyer DL. The Mauthner-initiated startle response in teleost
fish. J Exp Biol. 1977;66: 65–81. doi:10.1242/jeb.66.1.65

30.   Kimmel CB, Patterson J, Kimmel RO. The development and behavioral characteristics of
the startle response in the zebra fish. Dev Psychobiol. 1974;7: 47–60.
doi:10.1002/dev.420070109

31.   Davis M, Gendelman DS, Tischler MD, Gendelman PM. A primary acoustic startle circuit:
lesion and stimulation studies. J Neurosci. 1982;2: 791–805. doi:10.1523/JNEUROSCI.02-
06-00791.1982

32.   Marsden KC, Jain RA, Wolman MA, Echeverry FA, Nelson JC, Hayer KE, et al. A Cyfip2-
Dependent Excitatory Interneuron Pathway Establishes the Innate Startle Threshold. Cell
Rep. 2018;23: 878–887. doi:S2211-1247(18)30471-6 [pii]

33.   Schenck A, Bardoni B, Moro A, Bagni C, Mandel JL. A highly conserved protein family
interacting with the fragile X mental retardation protein (FMRP) and displaying selective
interactions with FMRP-related proteins FXR1P and FXR2P. Proc Natl Acad Sci U S A.
2001;98: 8844–8849. doi:10.1073/pnas.151231598 [doi]

34.   Napoli I, Mercaldo V, Boyl PP, Eleuteri B, Zalfa F, Rubeis SD, et al. The fragile X syndrome
protein represses activity-dependent translation through CYFIP1, a new 4E-BP. Cell.
2008;134: 1042–1054. doi:10.1016/j.cell.2008.07.031 [doi]

35.   Kobayashi K, Kuroda S, Fukata M, Nakamura T, Nagase T, Nomura N, et al. p140Sra-1
(specifically Rac1-associated protein) is a novel specific target for Rac1 small GTPase. J
Biol Chem. 1998;273: 291–295. doi:10.1074/jbc.273.1.291 [doi]

36.   Eden S, Rohatgi R, Podtelejnikov AV, Mann M, Kirschner MW. Mechanism of regulation of
WAVE1-induced actin nucleation by Rac1 and Nck. Nature. 2002;418: 790–793.
doi:10.1038/nature00859

37.   Schenck A, Bardoni B, Langmann C, Harden N, Mandel JL, Giangrande A. CYFIP/Sra-1
controls neuronal connectivity in Drosophila and links the Rac1 GTPase pathway to the
fragile X protein. Neuron. 2003;38: 887–898. doi:S0896627303003544 [pii]

38.   Bogdan S, Grewe O, Strunk M, Mertens A, Klämbt C. Sra-1 interacts with Kette and Wasp
and is required for neuronal and bristle development in Drosophila. Development.
2004;131: 3981–3989. doi:10.1242/dev.01274

39.   Chen Z, Borek D, Padrick SB, Gomez TS, Metlagel Z, Ismail AM, et al. Structure and
849   control of the actin regulatory WAVE complex. Nature. 2010;468: 533–538.
850   doi:10.1038/nature09623 [doi]

851   40.   Chen B, Padrick SB, Henry L, Rosen MK. Biochemical reconstitution of the WAVE
852   regulatory complex. Methods Enzymol. 2014;540: 55–72. doi:10.1016/B978-0-12-397924-
853   7.00004-2

854   41.   Chen B, Chou HT, Brautigam CA, Xing W, Yang S, Henry L, et al. Rac1 GTPase activates
855   the WAVE regulatory complex through two distinct binding sites. Elife. 2017;6:
856   10.7554/eLife.29795. doi:10.7554/eLife.29795 [doi]

857   42.   Trowe T, Klostermann S, Baier H, Granato M, Crawford AD, Grunewald B, et al. Mutations
858   disrupting the ordering and topographic mapping of axons in the retinotectal projection of
859   the zebrafish, Danio rerio. Development. 1996;123: 439–450. doi:10.1242/dev.123.1.439

860   43.   Schenck A, Qurashi A, Carrera P, Bardoni B, Diebold C, Schejter E, et al. WAVE/SCAR, a
861   multifunctional complex coordinating different aspects of neuronal connectivity. Dev Biol.
862   2004;274: 260–270. doi:10.1016/j.ydbio.2004.07.009

863   44.   Pittman AJ, Gaynes JA, Chien CB. Nev (Cyfip2) is Required for Retinal Lamination and
864   Axon Guidance in the Zebrafish Retinotectal System. Dev Biol. 2010;344: 784–794.
865   doi:10.1016/j.ydbio.2010.05.512 [doi]

866   45.   Cioni JM, Wong HH, Bressan D, Kodama L, Harris WA, Holt CE. Axon-Axon Interactions
867   Regulate Topographic Optic Tract Sorting via CYFIP2-Dependent WAVE Complex
868   Function. Neuron. 2018;97: 1078-1093.e6. doi:S0896-6273(18)30052-7 [pii]

869   46.   Lee Y, Kim D, Ryu JR, Zhang Y, Kim S, Kim Y, et al. Phosphorylation of CYFIP2, a
870   component of the WAVE-regulatory complex, regulates dendritic spine density and neurite
871   outgrowth in cultured hippocampal neurons potentially by affecting the complex assembly.
872   Neuroreport. 2017;28: 749–754. doi:10.1097/WNR.0000000000000838

873   47.   Zhang Y, Kang Hyae R, Lee S-H, Kim Y, Ma R, Jin C, et al. Enhanced Prefrontal Neuronal
874   Activity and Social Dominance Behavior in Postnatal Forebrain Excitatory Neuron-Specific
875   Cyfip2 Knock-Out Mice. Front Mol Neurosci. 2020;13: 574947.
876   doi:10.3389/fnmol.2020.574947

877   48.   Kumar V, Kim K, Joseph C, Kourrich S, Yoo SH, Huang HC, et al. C57BL/6N mutation in
878   cytoplasmic FMRP interacting protein 2 regulates cocaine response. Science. 2013;342:
879   1508–1512. doi:10.1126/science.1245503 [doi]

880   49.   Abekhoukh S, Bardoni B. CYFIP family proteins between autism and intellectual disability:
881   links with Fragile X syndrome. Front Cell Neurosci. 2014;8: 81.
882   doi:10.3389/fncel.2014.00081 [doi]

883   50.   Barthelson K, Baer L, Dong Y, Hand M, Pujic Z, Newman M, et al. Zebrafish chromosome
884   14 gene differential expression in the fmr1 h u2787 model of Fragile X syndrome. Front
885   Genet. 2021;12: 625466. doi:10.3389/fgene.2021.625466

51. Hu J, Chen L, Yin J, Yin H, Huang Y, Tian J. Hyperactivity, memory defects, and
craniofacial abnormalities in zebrafish fmr1 mutant larvae. Behav Genet. 2020;50: 152–
160. doi:10.1007/s10519-020-09995-7

52. Zhu SI, McCullough MH, Pujic Z, Sibberas J, Sun B, Darveniza T, et al. fmr1 Mutation
Alters the Early Development of Sensory Coding and Hunting and Social Behaviors in
Larval Zebrafish. J Neurosci. 2023;43: 1211–1224. doi:10.1523/JNEUROSCI.1721-
22.2022

53. Constantin L, Poulsen RE, Scholz LA, Favre-Bulle IA, Taylor MA, Sun B, et al. Altered
brain-wide auditory networks in a zebrafish model of fragile X syndrome. BMC Biol.
2020;18: 125. doi:10.1186/s12915-020-00857-6

54. Spence EF, Soderling SH. Actin Out: Regulation of the Synaptic Cytoskeleton. J Biol
Chem. 2015;290: 28613–28622. doi:10.1074/jbc.R115.655118 [doi]

55. Chalkia D, Nikolaidis N, Makalowski W, Klein J, Nei M. Origins and evolution of the formin
multigene family that is involved in the formation of actin filaments. Mol Biol Evol. 2008;25:
2717–2733. doi:10.1093/molbev/msn215

56. Pellegrin S, Mellor H. The Rho family GTPase Rif induces filopodia through mDia2. Curr
Biol. 2005;15: 129–133. doi:10.1016/j.cub.2005.01.011

57. Schirenbeck A, Bretschneider T, Arasada R, Schleicher M, Faix J. The Diaphanous-related
formin dDia2 is required for the formation and maintenance of filopodia. Nat Cell Biol.
2005;7: 619–625. doi:10.1038/ncb1266

58. Peng J, Wallar BJ, Flanders A, Swiatek PJ, Alberts AS. Disruption of the Diaphanous-
related formin Drf1 gene encoding mDia1 reveals a role for Drf3 as an effector for Cdc42.
Curr Biol. 2003;13: 534–545. doi:10.1016/s0960-9822(03)00170-2

59. Maciver SK, Hussey PJ. The ADF/cofilin family: actin-remodeling proteins. Genome Biol.
2002;3: reviews3007. doi:10.1186/gb-2002-3-5-reviews3007

60. Nagar D, James TK, Mishra R, Guha S, Burgess SM, Ghose A. The Formin Fmn2b Is
Required for the Development of an Excitatory Interneuron Module in the Zebrafish
Acoustic Startle Circuit. eNeuro. 2021;8. doi:10.1523/ENEURO.0329-20.2021

61. Wolman MA, Jain RA, Liss L, Granato M. Chemical modulation of memory formation in
larval zebrafish. Proc Natl Acad Sci U S A. 2011;108: 15468–15473.
doi:10.1073/pnas.1107156108 [doi]

62. Focking M, Lopez LM, English JA, Dicker P, Wolff A, Brindley E, et al. Proteomic and
genomic evidence implicates the postsynaptic density in schizophrenia. Mol Psychiatry.
2015;20: 424–432. doi:10.1038/mp.2014.63 [doi]

63. Hoeffer CA, Sanchez E, Hagerman RJ, Mu Y, Nguyen DV, Wong H, et al. Altered mTOR
signaling and enhanced CYFIP2 expression levels in subjects with fragile X syndrome.
Genes Brain Behav. 2012;11: 332–341. doi:10.1111/j.1601-183X.2012.00768.x [doi]

64.   Noroozi R, Omrani MD, Sayad A, Taheri M, Ghafouri-Fard S. Cytoplasmic FMRP
interacting protein 1/2 (CYFIP1/2) expression analysis in autism. Metab Brain Dis. 2018;33:
1353–1358. doi:10.1007/s11011-018-0249-8 [doi]

65.   Zweier M, Begemann A, McWalter K, Cho MT, Abela L, Banka S, et al. Spatially clustering
de novo variants in CYFIP2, encoding the cytoplasmic FMRP interacting protein 2, cause
intellectual disability and seizures. Eur J Hum Genet. 2019;27: 747–759.
doi:10.1038/s41431-018-0331-z

66.   Kirkpatrick SL, Goldberg LR, Yazdani N, Babbs RK, Wu J, Reed ER, et al. Cytoplasmic
FMR1-Interacting Protein 2 Is a Major Genetic Factor Underlying Binge Eating. Biol
Psychiatry. 2017;81: 757–769. doi:10.1016/j.biopsych.2016.10.021

67.   Babbs RK, Beierle JA, Ruan QT, Kelliher JC, Chen MM, Feng AX, et al. Cyfip1
Haploinsufficiency Increases Compulsive-Like Behavior and Modulates Palatable Food
Intake in Mice: Dependence on Cyfip2 Genetic Background, Parent-of Origin, and Sex. G3
. 2019;9: 3009–3022. doi:10.1534/g3.119.400470

68.   Babbs RK, Beierle JA, Yao EJ, Kelliher JC, Medeiros AR, Anandakumar J, et al. The effect
of the demyelinating agent cuprizone on binge-like eating of sweetened palatable food in
female and male C57BL/6 substrains. Appetite. 2020;150: 104678.
doi:10.1016/j.appet.2020.104678

69.   Manigandan S, Yun JW. Loss of cytoplasmic FMR1-interacting protein 2 (CYFIP2) induces
browning in 3T3-L1 adipocytes via repression of GABA-BR and activation of mTORC1. J
Cell Biochem. 2022;123: 863–877. doi:10.1002/jcb.30231

70.   Nachmany H, Wald S, Abekasis M, Bulvik S, Weil M. Two potential biomarkers identified in
mesenchymal stem cells and leukocytes of patients with sporadic amyotrophic lateral
sclerosis. Dis Markers. 2012;32: 211–220. doi:10.3233/DMA-2011-0885

71.   Lilo E, Wald-Altman S, Solmesky LJ, Ben Yaakov K, Gershoni-Emek N, Bulvik S, et al.
Characterization of human sporadic ALS biomarkers in the familial ALS transgenic
mSOD1(G93A) mouse model. Hum Mol Genet. 2013;22: 4720–4725.
doi:10.1093/hmg/ddt325

72.   Tiwari SS, Mizuno K, Ghosh A, Aziz W, Troakes C, Daoud J, et al. Alzheimer-related
decrease in CYFIP2 links amyloid production to tau hyperphosphorylation and memory
loss. Brain. 2016;139: 2751–2765. doi:10.1093/brain/aww205

73.   Nakashima M, Kato M, Aoto K, Shiina M, Belal H, Mukaida S, et al. De novo hotspot
variants in CYFIP2 cause early-onset epileptic encephalopathy. Ann Neurol. 2018;83: 794–
806. doi:10.1002/ana.25208 [doi]

74.   Begemann A, Sticht H, Begtrup A, Vitobello A, Faivre L, Banka S, et al. New insights into
the clinical and molecular spectrum of the novel CYFIP2-related neurodevelopmental
disorder and impairment of the WRC-mediated actin dynamics. Genet Med. 2021;23: 543–
554. doi:10.1038/s41436-020-01011-x

75.   Biembengut ÍV, Silva ILZ, Souza T de ACB de, Shigunov P. Cytoplasmic FMR1 interacting
protein (CYFIP) family members and their function in neural development and disorders.
Mol Biol Rep. 2021;48: 6131–6143. doi:10.1007/s11033-021-06585-6

76.   Biembengut ÍV, Shigunov P, Frota NF, Lourenzoni MR, de Souza TACB. Molecular
Dynamics of CYFIP2 Protein and Its R87C Variant Related to Early Infantile Epileptic
Encephalopathy. Int J Mol Sci. 2022;23. doi:10.3390/ijms23158708

77.   Takahashi H, Nemoto T, Yoshida T, Honda H, Hasegawa T. Cancer diagnosis marker
extraction for soft tissue sarcomas based on gene expression profiling data by using
projective adaptive resonance theory (PART) filtering method. BMC Bioinformatics. 2006;7:
399. doi:10.1186/1471-2105-7-399

78.   Mongroo PS, Noubissi FK, Cuatrecasas M, Kalabis J, King CE, Johnstone CN, et al. IMP-1
displays cross-talk with K-Ras and modulates colon cancer cell survival through the novel
proapoptotic protein CYFIP2. Cancer Res. 2011;71: 2172–2182. doi:10.1158/0008-
5472.CAN-10-3295

79.   Jiao S, Li N, Cai S, Guo H, Wen Y. Inhibition of CYFIP2 promotes gastric cancer cell
proliferation and chemoresistance to 5-fluorouracil through activation of the Akt signaling
pathway. Oncol Lett. 2017;13: 2133–2140. doi:10.3892/ol.2017.5743

80.   Vandamme T, Beyens M, Boons G, Schepers A, Kamp K, Biermann K, et al. Hotspot
DAXX, PTCH2 and CYFIP2 mutations in pancreatic neuroendocrine neoplasms. Endocr
Relat Cancer. 2019;26: 1–12. doi:10.1530/ERC-18-0120

81.   Liu Y, Liu H, Bian Q. Identification of Potential Biomarkers Associated with Basal Cell
Carcinoma. Biomed Res Int. 2020;2020: 2073690. doi:10.1155/2020/2073690

82.   Li Y, Song X, Liu L, Yue L. NUAK2 silencing inhibits the proliferation, migration and
epithelial□to□mesenchymal transition of cervical cancer cells via upregulating CYFIP2. Mol
Med Rep. 2021;24. doi:10.3892/mmr.2021.12457

83.   Steffen A, Rottner K, Ehinger J, Innocenti M, Scita G, Wehland J, et al. Sra-1 and Nap1 link
Rac to actin assembly driving lamellipodia formation. EMBO J. 2004;23: 749–759.
doi:10.1038/sj.emboj.7600084

84.   Rubeis SD, Pasciuto E, Li KW, Fernandez E, Marino DD, Buzzi A, et al. CYFIP1
coordinates mRNA translation and cytoskeleton remodeling to ensure proper dendritic
spine formation. Neuron. 2013;79: 1169–1182. doi:10.1016/j.neuron.2013.06.039 [doi]

85.   Pathania M, Davenport EC, Muir J, Sheehan DF, Lopez-Domenech G, Kittler JT. The
autism and schizophrenia associated gene CYFIP1 is critical for the maintenance of
dendritic complexity and the stabilization of mature spines. Transl Psychiatry. 2014;4:
e374. doi:10.1038/tp.2014.16 [doi]

86.   Davenport EC, Szulc BR, Drew J, Taylor J, Morgan T, Higgs NF, et al. Autism and
Schizophrenia-Associated CYFIP1 Regulates the Balance of Synaptic Excitation and
Inhibition. Cell Rep. 2019;26: 2037-2051.e6. doi:S2211-1247(19)30129-9 [pii]

87.　Hanneman E, Trevarrow B, Metcalfe WK, Kimmel CB, Westerfield M. Segmental pattern of
development of the hindbrain and spinal cord of the zebrafish embryo. Development.
1988;103: 49–58. doi:10.1242/dev.103.1.49

88.　Eaton RC, Farley RD. Development of the mauthner neurons in embryos and larvae of the
zebrafish, Brachydanio rerio. Copeia. 1973;1973: 673. doi:10.2307/1443067

89.　Kimmel CB, Hatta K, Metcalfe WK. Early axonal contacts during development of an
identified dendrite in the brain of the zebrafish. Neuron. 1990;4: 535–545.
doi:10.1016/0896-6273(90)90111-r

90.　Kimmel CB, Sessions SK, Kimmel RJ. Morphogenesis and synaptogenesis of the zebrafish
Mauthner neuron. J Comp Neurol. 1981;198: 101–120. doi:10.1002/cne.901980110

91.　Tanimoto M, Ota Y, Horikawa K, Oda Y. Auditory input to CNS is acquired coincidentally
with development of inner ear after formation of functional afferent pathway in zebrafish. J
Neurosci. 2009;29: 2762–2767. doi:10.1523/JNEUROSCI.5530-08.2009

92.　Siomi MC, Siomi H, Sauer WH, Srinivasan S, Nussbaum RL, Dreyfuss G. FXR1, an
autosomal homolog of the fragile X mental retardation gene. EMBO J. 1995;14: 2401–
2408. doi:10.1002/j.1460-2075.1995.tb07237.x

93.　Zhang Y, O'Connor JP, Siomi MC, Srinivasan S, Dutra A, Nussbaum RL, et al. The fragile
X mental retardation syndrome protein interacts with novel homologs FXR1 and FXR2.
EMBO J. 1995;14: 5358–5366. doi:10.1002/j.1460-2075.1995.tb00220.x

94.　Bakker CE, de Diego Otero Y, Bontekoe C, Raghoe P, Luteijn T, Hoogeveen AT, et al.
Immunocytochemical and biochemical characterization of FMRP, FXR1P, and FXR2P in
the mouse. Exp Cell Res. 2000;258: 162–170. doi:10.1006/excr.2000.4932

95.　Agulhon C, Blanchet P, Kobetz A, Marchant D, Faucon N, Sarda P, et al. Expression of
FMR1, FXR1, and FXR2 genes in human prenatal tissues. J Neuropathol Exp Neurol.
1999;58: 867–880. doi:10.1097/00005072-199908000-00009

96.　Tucker B, Richards R, Lardelli M. Expression of three zebrafish orthologs of human FMR1-
related genes and their phylogenetic relationships. Dev Genes Evol. 2004;214: 567–574.
doi:10.1007/s00427-004-0438-9

97.　Coffee RL Jr, Tessier CR, Woodruff EA 3rd, Broadie K. Fragile X mental retardation protein
has a unique, evolutionarily conserved neuronal function not shared with FXR1P or
FXR2P. Dis Model Mech. 2010;3: 471–485. doi:10.1242/dmm.004598

98.　Boczkowska M, Rebowski G, Kast DJ, Dominguez R. Structural analysis of the transitional
state of Arp2/3 complex activation by two actin-bound WCAs. Nat Commun. 2014;5: 3308.
doi:10.1038/ncomms4308

99.　Lacoste AM, Schoppik D, Robson DN, Haesemeyer M, Portugues R, Li JM, et al. A
convergent and essential interneuron pathway for Mauthner-cell-mediated escapes. Curr
Biol. 2015;25: 1526–1534. doi:10.1016/j.cub.2015.04.025 [doi]

100.   Darnell JC, Van Driesche SJ, Zhang C, Hung KYS, Mele A, Fraser CE, et al. FMRP
1037   stalls ribosomal translocation on mRNAs linked to synaptic function and autism. Cell.
1038   2011;146: 247–261. doi:10.1016/j.cell.2011.06.013

1039   101.   Möhrle D, Wang W, Whitehead SN, Schmid S. GABAB Receptor Agonist R-Baclofen
1040   Reverses Altered Auditory Reactivity and Filtering in the Cntnap2 Knock-Out Rat. Front
1041   Integr Neurosci. 2021;15. doi:10.3389/fnint.2021.710593

1042   102.   Waldmeier PC, Kaupmann K, Urwyler S. Roles of GABAB receptor subtypes in
1043   presynaptic auto- and heteroreceptor function regulating GABA and glutamate release. J
1044   Neural Transm. 2008;115: 1401–1411. doi:10.1007/s00702-008-0095-7

1045   103.   Wu C, Sun D. GABA receptors in brain development, function, and injury. Metab Brain
1046   Dis. 2015;30: 367–379. doi:10.1007/s11011-014-9560-1

1047   104.   Tureček R, Melichar A, Králíková M, Hrušková B. The role of GABAB receptors in the
1048   subcortical pathways of the mammalian auditory system. Front Endocrinol . 2023;14:
1049   1195038. doi:10.3389/fendo.2023.1195038

1050   105.   Berry-Kravis E, Hagerman R, Visootsak J, Budimirovic D, Kaufmann WE, Cherubini M,
1051   et al. Arbaclofen in fragile X syndrome: results of phase 3 trials. J Neurodev Disord.
1052   2017;9: 3. doi:10.1186/s11689-016-9181-6

1053   106.   Veenstra-VanderWeele J, Cook EH, King BH, Zarevics P, Cherubini M, Walton-Bowen
1054   K, et al. Arbaclofen in Children and Adolescents with Autism Spectrum Disorder: A
1055   Randomized, Controlled, Phase 2 Trial. Neuropsychopharmacology. 2017;42: 1390–1398.
1056   doi:10.1038/npp.2016.237

1057   107.   Erickson CA, Veenstra-Vanderweele JM, Melmed RD, McCracken JT, Ginsberg LD,
1058   Sikich L, et al. STX209 (arbaclofen) for autism spectrum disorders: an 8-week open-label
1059   study. J Autism Dev Disord. 2014;44: 958–964. doi:10.1007/s10803-013-1963-z

1060   108.   Mattson MP, Gleichmann M, Cheng A. Mitochondria in neuroplasticity and neurological
1061   disorders. Neuron. 2008;60: 748–766. doi:10.1016/j.neuron.2008.10.010

1062   109.   Smith GM, Gallo G. The role of mitochondria in axon development and regeneration.
1063   Dev Neurobiol. 2018;78: 221–237. doi:10.1002/dneu.22546

1064   110.   Safer D, Elzinga M, Nachmias VT. Thymosin beta 4 and Fx, an actin-sequestering
1065   peptide, are indistinguishable. J Biol Chem. 1991;266: 4029–4032. Available:
1066   https://www.ncbi.nlm.nih.gov/pubmed/1999398

1067   111.   Yu FX, Lin SC, Morrison-Bogorad M, Atkinson MA, Yin HL. Thymosin beta 10 and
1068   thymosin beta 4 are both actin monomer sequestering proteins. J Biol Chem. 1993;268:
1069   502–509. doi:10.1016/s0021-9258(18)54179-x

1070   112.   Cheng IH, Lin Y-C, Hwang E, Huang H-T, Chang W-H, Liu Y-L, et al. Collagen VI
1071   protects against neuronal apoptosis elicited by ultraviolet irradiation via an
1072   Akt/phosphatidylinositol 3-kinase signaling pathway. Neuroscience. 2011;183: 178–188.
1073   doi:10.1016/j.neuroscience.2011.03.057

1074   113.   Fox MA. Novel roles for collagens in wiring the vertebrate nervous system. Curr Opin
1075   Cell Biol. 2008;20: 508–513. doi:10.1016/j.ceb.2008.05.003

1076

1077

<u>Figure Legends</u>

<u>Figure 1. Cyfip2 establishes the acoustic startle threshold through Rac1 and FMRP.</u>

(A) Cyfip2 protein interacting domain diagram of wildtype (top) and mutant (bottom) Cyfip2

proteins. Black arrowheads indicate the positions of induced mutations in Cyfip2, eliminating the

Rac1- (C179R) or FMRP/eIF4E (K723E)-binding capacity of Cyfip2. (B) Cyfip2 actin regulatory

pathway wherein Cyfip2 (orange) upon stimulation by Rac1-GTP triggers WAVE1 activation,

Arp2/3-complex initiation and branched actin nucleation. (C) Cyfip2 translational repression

pathway in which Cyfip2, eIF4E (teal), and FMRP (pink) along with the poly-A binding protein

(PABP; gray), sequester neurodevelopmentally important mRNAs from being translated. (D)

Average startle frequency (%) after 10 trials at 13.6, 25.7, 29.2, 35.5, 39.6 and 53.6 dB for 5 dpf

cyfip2 siblings (+/) and mutant (-/-) larvae heatshocked at 30 hpf for 40 minutes at 38°C. The

average startle frequency curve for cyfip2 siblings (+/; open circles, dashed line), cyfip2 mutants

(-/-; closed circles, solid line) and cyfip2 mutants harboring the *Tg(hsp70:cyfip2-EGFP)*+

transgene (-/-; Tg+; closed circles, solid green line). (E) Sensitivity indices, calculated as the

area under the startle frequency curves, for 5 dpf cyfip2 siblings and mutants, following a 40-

minute heatshock at 30 hpf to express either wildtype (Tg+; green), Rac1- (ΔRac1+; blue) or

FMRP/eIF4E- (ΔFMRP+; pink) binding deficient versions of Cyfip2-EGFP. Comparisons were

made to both non-transgenic (Tg-) and non-heatshocked controls. All indices (mean ± SD)

compared using a Kruskal-Wallis test with Dunn's multiple comparisons correction; $p^{****} <$

0.0001. (F) Sensitivity indices for 5 dpf *cyfip2* sibling (+/) and mutant (-/-) larvae following 1-cell

stage injection with CRISPR-Cas9 and a single, scrambled guide RNA (gRNA) or dual gRNA

cocktails targeting *fmr1*, *fxr1*, or *fxr2*. scrambled gRNA injected (white bar, closed circles); fmr1

gRNA injected (dark gray bar closed circles); *fxr1* gRNA injected (medium gray bar; closed

circles); *fxr2* gRNA injected (light gray bar, closed circles). Comparisons were made both within

genotype and between genotypes by condition. All indices (mean ± SD) compared using an

Ordinary one-way ANOVA with Sidak's multiple comparisons correction; $p^* < 0.05$; $p^{**} < 0.01$;

$p^{****} < 0.0001$.

1105

Figure 2. Cyfip2 acutely regulates branched actin polymerization and NMDARs to

establish the acoustic startle threshold.

(A) Sensitivity indices for 5 dpf *cyfip2* sibling (+/) and mutant (-/-) larvae, following a 40-minute

heatshock at 120 hpf (5 dpf) to express either wildtype (Tg+; green), Rac1- (ΔRac1+; blue) or

FMRP/eIF4E- (ΔFMRP+; pink) binding deficient versions of Cyfip2-EGFP. Comparisons were

made to non-transgenic (Tg-), heatshocked sibling (+/) and mutant (-/-) controls. All indices

(mean ± SD) compared using a Kruskal-Wallis test with Dunn's multiple comparisons correction;

p-values listed; $p^{**} < 0.01$, $p^{****} < 0.0001$. (B) Sensitivity indices for 5 dpf *cyfip2* wildtype (+/+;

white bar) and heterozygous (+/-; gray bar) larvae, treated for 30 minutes on d5 with 5, 20 or 50

µM CK-869. Comparisons were made both within genotype and within condition. All indices

(mean ± SD) compared using a Kruskal-Wallis test with Dunn's multiple comparisons correction;

$p^* < 0.05$; $p^{****} < 0.0001$. (C) Sensitivity indices for 5 dpf Tüpfel longfin (TL) larvae treated for

30 minutes on d5 with the highest, non-lethal doses the formin antagonist (SMIFH2; 5 µM),

PAK3 antagonist (IPA-3; 50 µM) and ROCK antagonist (GSK429286; 100 µM). Comparisons

were made between respective treatments and the DMSO controls. All indices (mean ± SD)

were compared using a Kruskal-Wallis test with Dunn's multiple comparisons correction; All

comparisons made were non-significant (n.s.). (D) Sensitivity indices for 5 dpf *cyfip2* sibling (+/)

and mutant (-/-) larvae, treated for 30 minutes on d5 with 100 or 500 µM MK-801. Comparisons

were made both between genotypes within condition and between conditions by genotype. All

indices (mean ± SD) were compared using an Ordinary one-way ANOVA with Tukey's multiple

comparisons correction. $p^{**} < 0.01$; $p^{***} < 0.001$; $p^{****} < 0.0001$.

1127

Figure 3. Loss of Cyfip2 causes widespread proteomic changes and GABAB receptor

signaling is critical for startle sensitivity.

(A) Bubble plots reporting the level of significance of the top 10 dysregulated proteins for both

cyfip2 heterozygous (top) and mutant (bottom) groups compared to wildtype controls. The size

of the dot is proportional to the significance of the results while the color code represents the

log2 fold change; top five upregulated (green), and top five downregulated (red) proteins are

shown. (B-C) Heat maps displaying the impacted canonical pathways (B) and diseases and

biological functions (C) from IPA functional analysis. The blue-colored gradient indicates the

degree of enrichment for the listed pathways or diseases/functions, as represented by the – log

of the P value for each pathway, disease and/or function. (D-E) Sensitivity indices for 5 dpf

cyfip2 sibling (+/) and mutant (-/-) larvae, treated for 60 minutes prior to testing with muscimol

(D) or (E) baclofen. All indices (mean ± SD) were compared using a one-way ANOVA with

Sidak's multiple comparisons correction. $p^* < 0.05$, $p^{**} < 0.01$; $p^{***} < 0.001$; $p^{****} < 0.0001$.

1141

1142   **Tables**

Compound	Concentration (µM)	Effect By Genotype		
		cyfip2(+/+)	*cyfip2(+/-)*	*cyfip2(-/-)*
NPAA (Cl-channel	1	93.21% of Control, p > 0.99	110.21% of Control, p < 0.99	113.57% of Control, p > 0.99

antagonist)	5	140.11% of Control, p = 0.9785	137.32% of Control, p = 0.1385	140.29% of Control, p > 0.2787
	10	**234.14% of Control, p** = 0.0038**	**201.29% of Control, p**** < 0.0001**	**155.06% of Control, p* = 0.0189**
MA (K+ channel/gap jxn. antagonist)	1	131.6% of Control, p > 0.99	11.87% of Control, p > 0.99	132.29% of Control, p = 0.5732
	5	137.24% of Control, p > 0.99	138.46% of Control, p = 0.0622	112.29% of Control, p > 0.99
	10	214.13% of Control, p = 0.0833	**169.81% of Control, p* = 0.0188**	**155.76% of Control, p* = 0.0288**
POBA (AAR/calmodulin antagonist)	1	102.9% of Control, p > 0.99	107.08% of Control, p > 0.99	101.04% of Control, p > 0.99
	10	136.17% of Control, p > 0.99	106.58% of Control, p > 0.99	99.01% of Control, p > 0.99
	50	**175.8% of Control, p* = 0.0123**		99.76% of Control, p > 0.99
ETAZ (PDE4 inhibitor)	1	141.79% of Control, p > 0.99	126.36% of Control, p > 0.99	121.74% of Control, p > 0.99
	10	168.6% of Control, p = 0.5472	**172.26% of Control, p* = 0.0137**	152.73% of Control, p = 0.3977
	50	**236.64% of Control, p** = 0.0029**	**179.99% of Control, p** = 0.0015**	**149.75% of Control, p* = 0.0105**
MK-801 (NMDAR antagonist)	100	120.19% of Control, p > 0.99		**145.93% of Control, p** = 0.0024**
	500	**212.69% of Control, p**** < 0.0001**		**163.26% of Control, p*** = 0.0002**
BMS-204352 (K+ channel antagonist)	10	79.48% of Control, p = 0.4269		89.55% of Control, p > 0.99
	50	76.08% of Control, p = 0.6759		108.74% of Control, p > 0.99
NSC-23766 (Rac1 antagonist)	100	**49.14% of Control, p*** = 0.0002**		81.60% of Control, p > 0.99
	200	86.26% of Control, p > 0.99		105.28% of Control, p > 0.99

1143

Table 1. Cyfip2 may regulate NMDARs to control acoustic startle sensitivity. Mean startle index

comparisons, listed as percentage (%) of the mean startle index of vehicle-treated controls by *cyfip2*

genotype and drug concentration, for larvae treated with compounds targeting the indicated pathways

[61] to increase acoustic startle sensitivity. All significant differences (p < 0.05) are listed (**bold**) for

comparisons using a Kruskal-Wallis test and Dunn's multiple comparisons correction. NPAA (*N*-

phenylanthranilic acid); MA (meclofenamic acid); POBA (phenoxybenzamine); ETAZ (etazolate).

1150

1151

Figure 1 Deslauriers et al. 2023

Figure 2 Deslauriers et al. 2023

Figure 3 Deslauriers et al. 2023