CEP41, a ciliopathy-linked centrosomal protein, regulates microtubule

assembly and cell division

Running title: CEP41 regulates microtubule assembly and cell division

Shweta Shyam Prassanawar[1], Tuhin Sarkar[1], and Dulal Panda[1,2]*

[1]Department of Biosciences and Bioengineering, Indian Institute of Technology Bombay, Mumbai,

India

[2]National Institute of Pharmaceutical Education and Research, SAS Nagar, Punjab, India

*Corresponding author: Dulal Panda

Email: panda@iitb.ac.in, panda@niper.ac.in

Keywords: Centrosomal proteins, CEP41, microtubules, microtubule stabilization, ciliopathy,

rhodanese-like domain.

Summary

A novel role of CEP41 as a microtubule-associated protein that promotes microtubule assembly

and regulates cell cycle progression could offer insights into the development of ciliopathies.

Abstract

Centrosomal proteins play pivotal roles in orchestrating microtubule dynamics, and their dysregulation leads to various disorders, including cancer and ciliopathies. Understanding the multifaceted roles of centrosomal proteins is vital to comprehend their involvement in disease development. Here, we report novel cellular functions of CEP41, a centrosomal and ciliary protein implicated in Joubert syndrome, a neurodevelopmental ciliopathy. We show that CEP41 is an essential microtubule-associated protein with microtubule-stabilizing activity. *In vitro*, purified CEP41 binds to preformed microtubules, promotes microtubule nucleation, and suppresses microtubule disassembly. When overexpressed in cultured cells, CEP41 localizes to microtubules and promotes formation of stable microtubule bundles. Conversely, shRNA-mediated knockdown of CEP41 causes disassembly of the interphase microtubule network, emphasizing its role in microtubule organization. Using GFP-tagged deletion constructs, we demonstrate that CEP41's association with microtubules relies on its conserved rhodanese homology domain and the N-terminal region. Furthermore, CEP41 depletion inhibits cell proliferation and induces G1 arrest, suggesting its potential role in cell cycle regulation. These insights into the cellular functions of CEP41 hold promise for unraveling the impact of its mutations in ciliopathies.

Introduction

Several cellular processes essential for cell growth and survival rely on the dynamic remodeling of the microtubule network. In animal cells, the centrosome is the key organelle responsible for the controlled assembly and organization of microtubules (Conduit et al., 2015). Due to its pivotal role in microtubule regulation, centrosomes influence several microtubule-dependent cellular processes, including cell division, cell migration, organelle positioning, and maintaining cell shape and polarity. Particularly noteworthy is the centrosome's role during mitosis, which ensures the formation of a dynamic bipolar spindle for precise chromosome segregation (Doxsey, 2001). Furthermore, the centrosome is a major signaling hub due to its indispensable role in ciliogenesis, involving several centrosomal proteins integral for proper cilia functioning (Bettencourt-Dias et al., 2011). Given these critical functions, defects in centrosome functioning adversely affect cell growth and development, resulting in a spectrum of diseases from cancer to developmental anomalies (Bettencourt-Dias et al., 2011; Nigg and Raff, 2009). Specifically, mutations in centrosomal proteins functioning at the cilium-centrosome interface have been linked to diverse genetic disorders broadly classified as ciliopathies (Vertii et al., 2015). These include developmental disorders such as Joubert syndrome, Bardet-Biedl syndrome, Meckel-Gruber syndrome, and polycystic kidney disease (Reiter and Leroux, 2017). Understanding the cellular functions of ciliopathy-associated centrosomal proteins is crucial for comprehending their role in disease development and devising targeted therapeutic strategies.

CEP41 is a centrosomal protein mutated in neurodevelopmental disorders, Joubert syndrome, and autism spectrum disorder (Korvatska et al., 2011; Lee et al., 2012; Patowary et al., 2019). CEP41 is essential for cilia assembly and indispensable in maintaining ciliary structure and motility by transporting ciliary tubulin glutamylase (Lee et al., 2012). CEP41 has also been reported to be necessary for cilia reabsorption, with its involvement in activating the Aurora kinase A-dependent cilia disassembly pathway (Ki et al., 2020). Knockdown of CEP41 in zebrafish embryos severely hampers development due to disrupted ciliary functions and vascular impairment (Ki et al., 2020; Lee et al., 2012). Although CEP41's ciliary functions have been well-studied, its role at the centrosome remains unexplored. In this regard, it is of significant interest to gain further information about the cellular functions of CEP41 to comprehend the impact of its mutations in ciliopathies.

Since CEP41 is associated with microtubule-based structures in cells (Gache et al., 2010; Lee et al., 2012), we aimed to explore its microtubule-associated functions. We provide evidence that CEP41 is a novel microtubule-associated protein (MAP) with microtubule-stabilizing activity. We used purified proteins to show that CEP41 binds to microtubules, promotes microtubule assembly, and stabilizes microtubules against disassembly. In cultured cells, CEP41 stabilizes microtubules and is essential for microtubule organization and cell proliferation. We

72 also show that CEP41 interacts with the microtubules through its rhodanese-like domain and the
73 coiled-coil motifs. Further, our study demonstrates CEP41's involvement in mitosis, hinting at its
74 yet unexplored role in cell cycle regulation. The findings advance our understanding of CEP41 as
75 a microtubule-associated protein potentially involved in maintaining the microtubule network in
76 cells.

Results

CEP41 interacts with tubulin and binds to microtubules

Based on previously reported localization of CEP41 at different microtubule-based structures,
such as the centrosome, cilia, and basal bodies (Andersen et al., 2003; Jakobsen et al., 2011;
Lee et al., 2012), we hypothesized that CEP41 could be a microtubule-interacting protein with
microtubule-associated functions. We performed a co-immunoprecipitation assay to test whether
CEP41 interacts with tubulin. CEP41 antibody, when incubated with the whole cell lysate of
NIH3T3 cells, co-immunoprecipitated tubulin, indicating a physical interaction between the two
proteins (Fig. 1A). Likewise, CEP41 was detected in the immunoprecipitants of α-tubulin,
confirming that it is a novel binding partner of tubulin (Fig. 1B). As negative controls, rabbit IgG,
mouse IgG, and Protein A agarose beads did not precipitate tubulin or CEP41. The positive
interaction of CEP41 with tubulin prompted us to probe its interaction with microtubules further.
We first checked if CEP41 is a part of the MAP family of proteins. MAPs are microtubule-binding
proteins capable of modulating microtubule dynamics by imparting stabilizing or destabilizing
effects (Bodakuntla et al., 2019). Brain tissue is a rich source of tubulin and MAPs, which can be
fractionated by repeated cycles of tubulin polymerization and depolymerization, yielding tubulin
rich with MAPs (Vallee, 1986). CEP41 was present in the crude brain tissue extract and MAP-rich
fractions (Fig. 1C). The presence of CEP41 in both the first and second cycles of MAP-rich
fractions indicates that CEP41 can bind and co-sediment with microtubules. It also suggests that
CEP41 cycles with other MAPs and can be a novel member of the microtubule-associated protein
family.

Next, we sought to establish a direct interaction between CEP41 and microtubules *in*
vitro. To this end, we cloned and expressed histidine-tagged CEP41 in bacteria and affinity
purified the protein (Fig. 1D) and validated it by immunoblotting with appropriate antibodies (Fig.
S1A). Using purified proteins, we assessed the physical interaction of CEP41 with brain
microtubules in a microtubule pelleting assay (Fig. 1E). Preformed taxol-stabilized microtubules
were incubated with CEP41 and then sedimented by ultracentrifugation. CEP41 pelleted with the
microtubules and was detected in the polymer fraction, whereas without microtubules, it did not
sediment (Fig. 1E). The band intensity of the pelleted CEP41 increased significantly in a
concentration-dependent manner (Fig. 1F), indicating that it was physically bound to microtubules

and sedimented along with microtubules (Fig. 1F). To confirm that the sedimentation of CEP41
with microtubules was attributed to their interaction and not its aggregation, CEP41 was
sedimented in similar conditions without microtubules and, as expected, remained in the soluble
fraction (Fig. S1B).

CEP41 promotes microtubule nucleation *in vitro*

The interaction of CEP41 with microtubules prompted us to check whether CEP41 regulates
microtubule assembly. Spontaneous polymerization of microtubules *in vitro* occurs at higher
tubulin concentrations as microtubule nucleation is a kinetically unfavorable process and requires
the formation of stable tubulin oligomers (Roostalu and Surrey, 2017). Microtubule stabilizers,
such as taxol and certain MAPs, enhance tubulin-tubulin interactions and promote microtubule
assembly (Bre and Karsenti, 1990; Kumar, 1981; Tovey and Conduit, 2018). At first, we checked
if CEP41 could induce spontaneous tubulin polymerization without any polymerization promoter.
Interestingly, CEP41 polymerized tubulin in the presence of GTP, as observed in a turbidity-
based assay (Fig. 2A). CEP41-induced tubulin polymerization showed a classic sigmoidal curve
with an initial lag phase followed by an elongation phase that saturated over time (Fig. 2A). With
increasing CEP41 concentrations, the lag time decreased, and the microtubule mass increased.
Tubulin alone at a similar concentration did not polymerize. Also, bovine serum albumin and
alkaline phosphatase as negative controls did not induce microtubule assembly. Compared to the
control, where tubulin did not polymerize, in the presence of 1 μM CEP41, there was a 10-fold
increase in the assembly rate (Fig. 2B). Furthermore, the rate increased by 13, 39, and 59-fold in
the presence of 2, 3, and 5 μM CEP41, respectively (Fig. 2B). Next, we sedimented the
microtubules formed in the presence of CEP41 and observed a distinct increase in the polymer
mass with increasing concentrations of CEP41 (Fig. 2C). In the absence of CEP41, tubulin did
not polymerize, as evident by the less sedimented mass. The band intensities of the pelleted
microtubules increased significantly with increasing CEP41. Likewise, increasing concentrations
of CEP41 reduced the mass of unpolymerized soluble tubulin, indicating tubulin heterodimers
polymerizing into microtubules (Fig. 2C). The polymer/soluble ratio increased by 1.8, 2.3, 3, and
3.9-fold in the presence of 1, 2, 3, and 5 μM CEP41, respectively (Fig. 2D).

We also visualized the microtubules formed in the presence of CEP41 using
transmission electron microscopy (Fig. 2E). Polymers were not observed in the absence of
CEP41 when tubulin was incubated only with GTP (Fig. 2E, top left panel). Similarly, CEP41
without tubulin showed no polymer formation (Fig. 2E, top right panel). In the presence of CEP41,
microtubules were seen with a dense mesh-like network of polymers in most fields (Fig. 2E,
bottom panel). Also, in some areas, microtubule branching was visible (black arrows), which
might have contributed to the formation of the mesh-like network (Fig. 2E, bottom right panel),
suggesting that in addition to nucleating microtubules, CEP41 may promote new filament growth

from an existing microtubule. We also visualized the CEP41-induced tubulin polymers by confocal

microscopy by immunostaining the microtubules using α-tubulin antibody (Fig. 2F). As observed

previously, here, too, in the absence of CEP41, there were no polymers and only small tubulin

aggregates (Fig. 2F, left panel). In the presence of CEP41, long filaments in a mesh-like network

were seen along with thick microtubule bundles (Fig. 2F, right panel). The results demonstrate

that CEP41 can promote microtubule nucleation and polymerization *in vitro*.

CEP41 enhances the rate of taxol-induced microtubule assembly

Next, we examined the effect of CEP41 on microtubule assembly in the presence of taxol, which

promotes tubulin polymerization at lower tubulin concentrations. CEP41 promoted tubulin

assembly in the presence of taxol (Fig. 3A). Compared to the control, the extent of polymerization

increased by 1.3, 1.4, 1.5, and 1.7-fold in the presence of 1, 2, 3, and 5 μM CEP41, respectively,

as evident by an increase in light scattering. Also, CEP41 significantly increased the rate of

tubulin polymerization, resulting in faster and more polymer formation (Fig. 3B). In comparison to

the rate of polymerization in the presence of taxol, the rate increased by 1.4, 1.8, 2.1, and 2.3-fold

in the presence of 1, 2, 3, and 5 μM CEP41, respectively, suggesting that CEP41 may assist in

the nucleation phase of microtubule assembly. Similarly, in a sedimentation assay, compared to

only taxol, more polymeric tubulin was sedimented in the presence of CEP41 (Fig. 3C). The band

intensity of pelleted microtubules increased significantly with increasing CEP41 concentration.

The polymer/soluble ratio increased by 1.3, 1.4, 1.7, and 1.9-fold in the presence of 1, 2, 3, and 5

μM CEP41, respectively (Fig. 3D). The data suggest that CEP41 is a promoter of microtubule

assembly.

CEP41 prevents dilution-induced disassembly of microtubules *in vitro*

Furthermore, we assessed the effect of CEP41 on dilution-induced microtubule

disassembly. Microtubules undergo depolymerization upon dilution, as the concentration of

tubulin subunits falls below the critical concentration required to maintain the assembly rate (Karr

and Purich, 1979). We polymerized microtubules in the presence of DMSO and then diluted them

10-fold in both the absence and presence of CEP41. Microtubules resisted depolymerization

when incubated with CEP41, as evidenced by more polymer mass sedimented compared to the

control (Fig. 3E). In the presence of 1, 3, and 5 μM CEP41, the polymer/soluble tubulin ratio

increased by 5, 10, and 16-fold, respectively, suggesting inhibition of microtubule

depolymerization (Fig. 3F). Additionally, the presence of CEP41 in the polymer fractions

complements the earlier microtubule-binding assay result, indicating that CEP41 interacts with

the microtubules and sediments along with them (Fig. 3E). Together, the results suggest that

CEP41 binds to microtubules and imparts a stabilizing effect, promoting assembly and preventing

disassembly.

Exogenously expressed CEP41 colocalizes with interphase microtubules and associates with the spindle poles and midbody during mitosis

Previous studies have reported the localization of CEP41 at the cilia and basal bodies and established its essential role in cilia functioning (Ki et al., 2020; Lee et al., 2012). We wanted to investigate the cilia-unrelated functions of CEP41 and characterize its localization during cell division. Since the polyclonal antibody available for CEP41 was unsuitable for immunofluorescence analysis, we exogenously expressed GFP-tagged CEP41 in human cervical cancer (HeLa) and mouse fibroblast (NIH3T3) cells. In HeLa cells, CEP41-GFP, when expressed in low amounts, showed centrosomal localization, whereas, in high-expressing cells, it localized to microtubules and showed a dense fiber-like network spread throughout the cell (Fig. 4A; Fig. S2). Similar CEP41 localization was observed in NIH3T3 cells as well (Fig. S3A). A distinct overlap between CEP41-GFP and α-tubulin fluorescence was visible, indicating colocalization (Fig. 4A). In contrast, GFP did not show such colocalization and exhibited diffused staining in the cytoplasm and nucleus in both cell lines (Fig. 4A; Fig. S2; Fig. S3A). CEP41-GFP localizing on microtubules further strengthens our findings that it binds and interacts with microtubules in cells. To measure the extent of colocalization, we determined Pearson's correlation coefficient (r), which measures the level of correlation between the two fluorophores. An r value close to 1 indicates a strong correlation, whereas a value close to 0 or negative indicates no correlation. The coefficient value r for CEP41 and microtubules was estimated to be 0.8 ± 0.01, indicating a strong correlation between the fluorophores (Fig. 4B). The r value for GFP control and microtubules was -0.25 ± 0.05, showing no correlation.

We next determined the localization of CEP41 during different cell cycle stages. In HeLa cells, CEP41-GFP localized to the spindle poles and microtubule-based structures throughout mitosis (Fig. 4C). In interphase and prophase, it localized to the centrosome and cytoplasmic microtubules and, upon nuclear envelope breakdown, was prominently seen at both the spindle poles and kinetochore microtubules, specifically near the poles. During telophase, CEP41 accumulated at the intermediate region of the cleavage furrow and, throughout cytokinesis, remained associated with the midbody. Similar localization of CEP41 at the midbody was observed in NIH3T3 cells (Fig. S3B), and GFP as negative control did not localize to the mitotic spindle (Fig. 4C). Our result is the first evidence to show the involvement of CEP41 in cell division, suggesting its potential functions in mitosis and cytokinesis.

CEP41 interacts with microtubules through its rhodanese-like domain and the N-terminal region

As CEP41 lacked any known microtubule-binding domain, we sought to identify its microtubule-interacting region. Prediction of its secondary and tertiary structure revealed that CEP41

comprises two coiled-coil motifs at the N-terminal region and a rhodanese-like domain (RHOD) at the intermediate region. In contrast, the C-terminal segment of the protein appeared to be primarily disordered (Fig. 5A; Fig. S4). To investigate this further, we generated several deletion constructs of CEP41 fused with GFP at the C-terminal and expressed the truncated proteins in HeLa cells through transient transfection (Fig. 5B-D). In the case of the N-terminal deletion construct (CEP41$^{\Delta N}$), the protein did not localize to microtubules and instead exhibited diffused localization in the cytoplasm, similar to the GFP control (Fig. 5C). For the construct with the rhodanese domain deleted (CEP41$^{\Delta RHOD}$), we noticed the formation of substantial aggregates in the cytoplasm, likely stemming from protein misfolding following the deletion of a conserved domain. Interestingly, the CEP41 construct with the C-terminal region deleted displayed a localization pattern similar to that of the full-length protein and exhibited colocalization with microtubules. This suggests that the C-terminal disordered region is dispensable for CEP41-microtubule interaction. Moreover, when only the RHOD domain was expressed without the N and C-terminal regions, the protein displayed a diffused distribution in the cytoplasm (Fig. 5C). These observations suggest that the coiled-coil motifs and the RHOD domain are essential for CEP41's interaction with microtubules. Similar results were obtained in dividing cells, where only the CEP41 construct with the C-terminal region deleted localized to spindle poles and the midbody, underscoring the significance of the RHOD domain and the N-terminal region in CEP41's functioning (Fig. 5D; Fig. S5). Remarkably, both these regions, the coiled-coil motifs, and the RHOD domain, are highly conserved among CEP41 orthologues, implying their essential role in CEP41's cellular functions (Fig. S6).

Overexpression of CEP41 induces formation of stable microtubule bundles

We also observed noticeable microtubule bundling in HeLa cells overexpressing CEP41 (Fig. 6A). Most CEP41-overexpressing cells exhibited thick bundles of straight microtubules (Fig. 6A, i), while in some cases, the microtubules concentrated in a circular ring around the nucleus (Fig. 6A, ii). No significant microtubule bundling was observed in GFP control cells, as they showed a well-spread interphase arrangement of microtubules (Fig. 6A). We counted the number of cells with prominent microtubule bundles, and 72 ± 12% of the cells expressing CEP41-GFP showed microtubule bundling (Fig. 6B). In contrast, only 15 ± 2% of GFP-expressing cells showed distinct bundles. Due to significant bundling, we expected a concomitant increase in the microtubule intensity, so we quantified the mean microtubule intensity per cell. The fluorescence intensity increased by 42 ± 4% in CEP41-overexpressing cells compared to non-transfected control cells, with no significant change in GFP-control cells (Fig. 6C). Next, we assessed the stability of these microtubule bundles. CEP41-induced microtubule bundles were positively stained with acetylated tubulin antibody (Fig. 6D). Acetylation of the Lys40 residue is a post-translational modification of α-tubulin and a marker for stable, long-lived microtubules (Fig. 6D) (Piperno et al., 1987). Non-

transfected control and GFP-control cells showed no significant microtubule bundling and a basal
level of acetylated tubulin staining, whereas the acetylated tubulin intensity increased by 81 ±
18% in CEP41-overexpressing cells (Fig. 6D). Further, we examined the stability of CEP41-
induced microtubule bundles against the depolymerizing action of nocodazole. The interphase
microtubule network depolymerized entirely in control and GFP-expressing cells (Fig. 6E).
However, in CEP41-GFP expressing cells, prominent microtubule bundles were visible, which
were resistant against nocodazole-induced depolymerization. We quantified the mean
microtubule intensity per cell, which was increased by 75 ± 12% in CEP41-overexpressing cells
compared to the control (Fig. 6E). The formation of stable microtubule bundles and localization of
CEP41 to these bundles hints toward a possible role of CEP41 in microtubule cross-linking.
Together with the *in vitro* findings, the results suggest that CEP41 is a microtubule-stabilizing
protein involved in microtubule nucleation and organization.

Depletion of CEP41 depolymerizes the interphase microtubule network

We further elucidated the role of CEP41 in microtubule regulation via loss of function experiments
where we depleted the endogenous levels of CEP41 using shRNA. Given that CEP41 localization
was similar in HeLa and NIH3T3 cells, we chose the NIH3T3 cell line for the depletion
experiments as it is a non-cancerous fibroblast cell line capable of undergoing ciliogenesis.
CEP41 is an essential ciliary protein reported to be crucial for ciliogenesis and cilia functioning
and is expressed in more amounts in NIH3T3 than in HeLa cells (Graser et al., 2007; Ki et al.,
2020; Lee et al., 2012). Immunoblotting of NIH3T3 cell lysate, 48 h post-transfection, showed a
74 ± 2% reduction in the protein level compared to untransfected and scrambled shRNA-
transfected cells (Fig. 7A,B). Further, immunofluorescence analysis of CEP41-depleted cells
revealed that CEP41 depletion depolymerized microtubules and disrupted the interphase
microtubule network (Fig. 7C). Control and scrambled control cells exhibited a well-spread
microtubule network with thin hair-like filaments throughout the cell. However, CEP41-depleted
cells lacked a well-spread microtubule network and showed fewer microtubules (Fig. 7C). We
quantified the fluorescence intensity of microtubules per cell, which was reduced by 47 ±1.2% in
CEP41-depleted cells compared to the control cells (Fig. 7D). CEP41-depleted cells also
exhibited a drastic change in cell morphology as they were significantly smaller than control cells
(Fig. 7C). The cytoplasmic area of CEP41-depleted cells was reduced by 55 ± 7% compared to
the control cells (Fig. 7E). The results indicate that CEP41 is vital for maintaining the interphase
microtubule network and cell shape.

CEP41 is essential for cell proliferation and cell cycle progression

During the immunofluorescence analysis of CEP41-depleted cells, we observed that the total
number of cells per field was lesser than the control and scrambled control. To investigate this

further, we studied the effect of CEP41 depletion on cell proliferation. After 48 hours of
transfection with shRNAs, cells were allowed to grow for one cycle. CEP41-depleted cells did not
double in number in comparison to control untransfected cells. The percentage inhibition of cell
proliferation was calculated to be 72 ± 17% for CEP41-depleted cells, whereas it was 20 ± 9% for
scrambled control cells (Fig. 8A). Further, we checked the effect of CEP41 depletion on the cell
cycle progression to determine if cell proliferation is inhibited due to a cell cycle block (Fig. 8B).
CEP41-depleted and scrambled control cells were grown in puromycin-containing media and then
subjected to a double thymidine block to synchronize cells at the G1/S boundary. Synchronized
cells were released from the thymidine block and allowed to grow for 8 h to progress into mitosis.
Control untransfected and scrambled control cells progressed into mitosis with 47 and 48.5%
cells in the G2/M stage, respectively (Fig. 8B,C). However, CEP41-depleted cells did not
progress to mitosis as efficiently as control cells, with only 28% of cells in the G2/M phase.
Interestingly, 51.5% of CEP41-depleted cells remained in the G1 state post-thymidine release,
compared to only 38% of control cells (Fig. 8B,C). Failure of CEP41-depleted cells to move from
G1 to G2/M suggests a potential involvement of CEP41 in the G1/S transition. Together, the
results indicate that CEP41 plays an essential role in cell cycle regulation, possibly via its
microtubule-associated functions.

Discussion

The key finding of our study is the identification of CEP41 as a microtubule-associated protein
that promotes microtubule assembly and stabilization. We provide evidence that CEP41 is vital
for maintaining the cellular microtubule network with indispensable roles in cell cycle progression
and proliferation. We also show that the coiled-coil motifs and the rhodanese-like domain of
CEP41 are essential for its interaction with the microtubules.

CEP41 is a microtubule-interacting protein

CEP41 has been suggested to be a microtubule-interacting protein (Andersen et al., 2003; Gache
et al., 2010); however, its direct binding and association with microtubules have not been shown.
Here, we demonstrate CEP41's binding to microtubules in cells and *in vitro*. In a co-sedimentation
assay, purified CEP41 was sedimented with preformed microtubules, suggesting an apparent
interaction between the proteins. Further, in prior studies, endogenous CEP41 was reported to
localize only at the centrosome and ciliary axoneme. In our investigation, CEP41 colocalized with
the microtubule lattice upon overexpression in HeLa and NIH3T3 cells. This observation
resembles end-binding proteins like EB1 and CLIP-170, which, at endogenous concentrations,
localize exclusively to growing microtubule tips yet bind throughout the microtubule lattice *in vitro*
(Ligon et al., 2006). CEP41's interaction with tubulin was also evident in co-immunoprecipitation
and *in vitro* microtubule polymerization assays. Thus, the results suggest that, at endogenous

levels, CEP41 could potentially be involved in microtubule anchoring and organization due to its
direct binding to the lattice or soluble tubulin. Furthermore, CEP41 was detected in the MAP-
enriched fraction of tubulin extracted from goat brain tissue, implying that it is a part of the MAP
family of proteins, which are well known to bind to microtubules and regulate their assembly.
MAPs can also modulate microtubule dynamics by influencing the interaction of other proteins
with microtubules (Bodakuntla et al., 2019). CEP41 is involved in the recruitment of tubulin
glutamylase TTLL6 to the ciliary axoneme; however, how it is involved in this regulation is unclear
(Lee et al., 2012). We propose that it could be due to CEP41's direct interaction with
microtubules, where it could function as an intermediary and promote localization of TTLL6 to the
ciliary microtubules. Similarly, cilia and spindle-associated protein (CSAP) have been reported to
bind to TTLL5 and regulate its interaction with microtubules (Bompard et al., 2018). Another
axoneme-associated protein, CEP162, is shown to mediate the interaction of transition zone
components with ciliary microtubule (Wang et al., 2013). Establishing CEP41's direct interaction
with tubulin and microtubules provides valuable insights into its role in microtubule-dependent
cellular functions. Furthermore, we demonstrated that the interaction between CEP41 and
microtubules is mediated by its rhodanese-like domain and coiled-coil motifs. Coiled-coil motifs in
centrosomal proteins are common, where they play an essential role in sub-cellular localization
and contribute to protein-protein interactions (Salisbury, 2003). However, the rhodanese
homology domain is primarily found in phosphatases and yeast ubiquitin-specific proteases
(Bordo and Bork, 2002). Although catalytically inactive, the RHOD domain in CEP41 might be
responsible for its stability and interactions with other proteins. Individuals affected by Joubert
syndrome carry heterozygous mutations in the conserved coiled-coil region and the RHOD
domain (Lee et al., 2012). We propose that these mutations might impede CEP41's interaction
with microtubules, consequently disrupting its microtubule-associated functions.

CEP41 is a novel regulator of microtubule organization and bundling

Regulation of microtubule dynamics in cells is a complex process involving the coordinated
activity of several proteins, including MAPs, centrosomal proteins, motor proteins, and several
protein kinases that control the activation and recruitment of the nucleating factors (Bodakuntla et
al., 2019; Kumari and Panda, 2018; Prassanawar and Panda, 2019; Tovey and Conduit, 2018).
We propose CEP41 as another member of this regulatory network with newly identified
microtubule-stabilizing activity. Upon CEP41 overexpression, we observed extensive microtubule
bundling and an increase in the overall microtubule content in HeLa cells. Similarly, in CEP41-
expressing NIH3T3 cells, a dense network of microtubules emanating from the centrosome was
seen, suggesting that CEP41 positively regulates microtubule assembly. Further, purified CEP41
promoted *in vitro* microtubule assembly and prevented dilution-induced microtubule disassembly,
again hinting at its function as a microtubule stabilizer. Microtubule nucleation is a cooperative

process and requires the initial assembly of stable microtubule seeds that grow into polymers
(Kuchnir Fygenson et al., 1995; Voter and Erickson, 1984). CEP41 might interact with tubulin and
stabilize the initial oligomeric tubulin assemblies. TPX2, a microtubule-stabilizing protein, is
thought to nucleate microtubules similarly by stabilizing oligomers (Schatz et al., 2003). The
microtubule bundling observed in cells could also be due to an increase in the rate of microtubule
assembly mediated by CEP41 overexpression. Given CEP41's direct interaction with
microtubules, it could also act as a cross-linker and enhance lateral interactions among the
microtubules, prompting bundling. Another centrosomal and ciliary protein, ENKD1, has been
shown to exhibit similar microtubule bundling and stabilization in cells and *in vitro* (Tiryaki et al.,
2022). Moreover, in NIH3T3 cells, CEP41 depletion disrupted the interphase microtubule
network, emphasizing its importance in organizing and maintaining the cellular microtubule array.
Since CEP41 is a component of the centrosome and we showed its capability to nucleate and
stabilize microtubules, we propose its possible involvement in regulating microtubule nucleation
at the centrosome. Although a direct interaction between CEP41 and γ-tubulin wasn't detected in
a co-immunoprecipitation assay (Fig. S7), CEP41 may interact with other γ-TuRC components.

CEP41 is essential for cell proliferation

During cell division, the centrosome orchestrates a complete reorganization of the interphase
microtubule network into a dynamic mitotic spindle (Doxsey et al., 2005). This highly dynamic
transition requires several centrosomal and microtubule-associated proteins that constantly
modulate microtubule dynamics. Owing to the microtubule nucleating and stabilizing functions of
CEP41, we hypothesized its involvement in the organization of spindle microtubules.
Interestingly, CEP41 localized at the spindle poles and the midbody in dividing cells. In addition,
CEP41 depletion inhibited cell proliferation and hampered cell cycle progression by arresting cells
at the G1 phase, validating our hypothesis that CEP41 is an essential protein with critical
functions during cell division. Cell cycle arrest at the G1 phase also hints toward faulty
centrosome duplication, suggesting a possible role of CEP41 in centriole biogenesis or
maturation, although this needs further assessment (Nigg, 2007). Earlier, the defective
phenotypes of CEP41 knockdown were attributed only to ciliary dysfunctions (Ki et al., 2020; Lee
et al., 2012). However, our findings suggest possible disruption of other cellular processes,
specifically microtubule-dependent processes. For instance, retarded cell migration and impaired
vascular development were prominent in CEP41-deficient zebrafish embryos (Ki et al., 2020), and
our results suggest that it could also be due to CEP41's role in microtubule organization.
Similarly, CEP41 was reported to be essential for ciliogenesis and cilia disassembly (Graser et
al., 2007; Lee et al., 2012), and here, we report its involvement in cell cycle regulation. Cilia
disassembly and the onset of the cell cycle are tightly coordinated processes, with many
centrosomal and ciliary proteins as key players functioning in this crosstalk (Izawa et al., 2015;

Jackson, 2011). IFT88 is one example of an essential ciliary protein that is reported to regulate
the G1/S transition in non-ciliated cells (Robert et al., 2007). Similarly, CEP164, a centrosomal
protein, regulates the G2/M transition and is vital for primary cilia formation (Graser et al., 2007).
CEP41 could also be a key player in this crosstalk that warrants further investigation. The
presence of a conserved RHOD domain in CEP41, the function of which remains unexplored,
strongly hints at the possibility of CEP41's involvement in cell cycle regulation. The RHOD
domain is found in the catalytic domain of Cdc25 phosphatase and the non-catalytic domain of
MAPK-phosphatases (Bordo and Bork, 2002). Both of these classes of proteins play roles in
regulating various signaling pathways associated with cell growth and survival. Interestingly,
CEP41 was shown to be essential for angiogenesis through its role in activating Aurora kinase A
and HIF1α (hypoxia-inducible factor 1α), which regulates the cilia disassembly pathway (Ki et al.,
2020). Therefore, investigating CEP41's role in such regulatory pathways will be an exciting
endeavor.

Overall, the results identify a potential role of CEP41 as a centrosomal MAP involved in
maintaining stable microtubule structures in cells and provide a background to explore its
functions as a cell cycle regulator.

Materials and methods

Construction of expression vectors

Full-length cDNA encoding human CEP41 (GenBank accession no. NM_018718.3) cloned in
pcDNA3.1 vector with a C-terminal eGFP tag was procured from Origene (USA). The CEP41
cDNA was amplified and subcloned into a bacterial expression vector pET28a(+) at Nde1 and
SalI restriction sites with an N-terminal hexahistidine tag. Deletion constructs with deleted N-
terminal (ΔN: residues deleted 33-137), rhodanese-like domain (ΔRHOD: 169-266), C-terminal
(ΔC: 273-373), and both N- and C- terminal (ΔNC: 33-137 and 273-373) regions were generated
by PCR using full-length CEP41 in pcDNA3.1 as a template. All constructs were purified using
endotoxin-free plasmid extraction kits (Qiagen) and sequenced to verify intact reading frames.

Cell culture and transfection

Mouse embryo-derived fibroblast (NIH3T3) and human cervical carcinoma (HeLa) cells were
obtained from the cell repository of the National Centre for Cell Sciences (India). The cells were
cultured in Dulbecco's Modified Eagle's Medium (HiMedia) supplied with 10% (v/v) fetal bovine
serum (Invitrogen) and 1% (v/v) antibiotic-antimycotic solution (HiMedia). The cells were free from
microbial or fungal contamination and were maintained at 37°C in a humidified cell culture
incubator (Sanyo) with 5% CO_2. Cells at 60-70% confluency were transiently transfected with

plasmids or shRNAs using Lipofectamine 3000 according to the manufacturer's protocol
(Invitrogen).

Cell lysis and immunoblotting

Immunoblotting analysis was performed as described earlier (Kumari et al., 2021; Srivastava and
Panda, 2018). Briefly, cells were trypsinized, lysed in a cold lysis buffer, and centrifuged, and the
protein concentration of the resulting supernatant was determined using Bradford's reagent
(Bradford, 1976). Equal concentrations of the samples were resolved on SDS-PAGE gel and
electroblotted onto PVDF membranes (Merck Millipore, USA) (Srivastava and Panda, 2018). The
blots were blocked at room temperature (RT, 25 ℃) for 1 h with 5% skim milk in TBST (Tris-
buffered saline pH 7.5 with 0.1% Tween-20), followed by incubation with primary antibody at 4 ℃
overnight, and peroxidase-conjugated secondary antibody at RT for 1 h. The blots were
developed using the chemiluminescence of luminol (Thermo Fisher Scientific), transferred on X-
ray film, and band intensities were quantified using ImageJ (National Institutes of Health,
Bethesda, USA).

Primary antibodies used were mouse anti-α-tubulin (T9026, Sigma-Aldrich, at 1:1000),
rabbit anti-CEP41 (17566-1-AP, Proteintech, at 1:800), mouse anti-β-actin (A2228, Sigma-
Aldrich, at 1:2000), mouse anti-GAPDH (sc-47724, Santa Cruz Biotech, at 1:500), and mouse
anti-polyhistidine (H1029, Sigma-Aldrich, at 1:1000). Secondary antibodies used were
horseradish peroxidase (HRP)-conjugated anti-mouse IgG (7076S, Cell Signaling Technology)
and anti-rabbit IgG (1706615, Bio-Rad) at 1:2000.

Co-immunoprecipitation assay

NIH3T3 cells seeded in T-75 flasks were harvested at 80-90% confluency and lysed in a cold
lysis buffer (50 mM Tris, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 0.1% Triton X, 500 μM sodium
orthovanadate, 2 mM PMSF, PIC, pH 7.4). The lysate was subjected to co-immunoprecipitation
with desired antibodies (Kumari et al., 2021). The cell lysate was pre-cleared with Protein A
agarose beads (Millipore, Sigma), and 700 μg of the pre-cleared lysate was incubated with 2 μg
of either anti-CEP41, anti-α-tubulin, rabbit IgG or mouse IgG at 4°C on a rocker overnight,
followed by incubation with pre-washed Protein A agarose beads for 2 h. The beads were
collected by centrifugation, washed, and the resin-bound immune complex was resuspended in
SDS loading buffer and boiled for 5 min. The samples were analyzed by SDS-PAGE followed by
western blotting. Rabbit IgG, mouse IgG, and Protein A agarose beads incubated with pre-
cleared lysate, without any antibody, were used as negative controls. The experiment was
performed twice.

Protein expression and purification

Recombinant full-length CEP41 was synthesized in *E.coli* BL21 (DE3) pLysE cells using the pET vector expression system with induction conditions of 30°C for 8 h with 0.5 mM IPTG. Post-induction, the cells were collected by centrifugation, lysed, and sonicated in ice-cold lysis buffer (50 mM Tris, 300 mM NaCl, 1mg/ml lysozyme, 2 mM PMSF, 0.1% Triton X-100, 0.01% β-ME, 10% glycerol, pH 8). Subsequently, the clarified cell lysate was incubated with pre-equilibrated Ni-NTA resin (Takara) for 2 h at 4°C in slow shaking conditions. The his-tagged protein was eluted with an elution buffer (50 mM Tris, 300 mM NaCl, and 250 mM imidazole, pH 8), concentrated, and dialyzed in a storage buffer (25 mM Tris, 50 mM NaCl, 10% glycerol, and pH 7). The dialyzed protein was aliquoted, snap-frozen, and stored at -80°C. Before every experiment, the histidine tag was cleaved using Thrombin CleanCleave Kit (Thermo Fisher Scientific), per the manufacturer's protocol. The thrombin-bound beads were separated by centrifugation, and the supernatant passed through the Ni-NTA column. The resulting cleaved CEP41 was concentrated and analyzed by SDS-PAGE and immunoblotted using anti-CEP41 and anti-polyhistidine antibodies.

Tubulin was purified from the goat brain tissue, as described previously (Hamel and Lin, 1981; Panda et al., 2005). Briefly, the goat brain tissue was homogenized in PEM buffer (50 mM PIPES, 3 mM $MgCl_2$, 1 mM EGTA, pH 6.8) containing 2 mM PMSF and 0.1% β-Me and centrifuged at 33,000 g and 4°C. The clear cell extract was polymerized at 37°C with 1 M glutamate, 10% DMSO, and 0.5 mM GTP (Sigma-Aldrich), centrifuged, and then depolymerized on ice. Two cycles of such polymerization and depolymerization were performed, and the resulting pure tubulin was snap-frozen and stored at -80°C. The purity of the protein was assessed by SDS-PAGE analysis.

Similarly, microtubule-associated proteins were isolated from the goat brain tissue employing two cycles of microtubule assembly and disassembly, as described earlier (Panda et al., 2005). After centrifugation, the clarified brain tissue extract was polymerized with 4 M glycerol and 0.5 mM GTP, followed by centrifugation and cold depolymerization. The presence of CEP41 in the MAP-rich fractions was detected by resolving 150 µg of the brain tissue extract and supernatant from the 1[st] and 2[nd] depolymerization steps on SDS-PAGE gel and immunoblotted with anti-CEP41 antibody. The experiment was performed twice.

Microtubule binding assay

The binding of CEP41 with microtubules was determined, as described earlier (Asthana et al., 2012). Tubulin (30 µM) was polymerized with taxol (20 µM) and 1 mM GTP in PEM buffer. 3 µM of the polymerized microtubules were incubated with CEP41 (1, 2, 3, and 5 µM) for 20 min at

37°C in PEM buffer containing taxol (5 μM) and 1 mM GTP. The reaction mixture was centrifuged
at 60,000 g for 20 min at 30°C in an ultracentrifuge Optima MAX-XP (Beckman Coulter), and the
resulting microtubule pellet was gently washed with warm PEM buffer containing 10% DMSO.
The pellet was dissolved in SDS running buffer, and equal volumes of the soluble and polymer
fractions were resolved on 12% SDS-PAGE gel and immunoblotted with anti-CEP41 antibody. As
a control, CEP41 (1, 2, 3, and 5 μM), without microtubules, was incubated and centrifuged in the
same conditions and analyzed by SDS-PAGE.

Light scattering assay

The effect of CEP41 on *in vitro* microtubule assembly was studied, as mentioned before
(Chaudhary et al., 2016). Tubulin (30 μM) was incubated with CEP41 (1, 2, 3, and 5 μM) for 20
min on ice in PEM buffer, followed by the addition of GTP (1 mM). The reaction kinetics was
monitored at 37°C by measuring light scattering at 350 nm using Spectramax M2e (Molecular
Devices). Bovine serum albumin (BSA) and alkaline phosphatase (10 μM) were used as negative
controls. Similarly, the effect of CEP41 on taxol-induced tubulin polymerization was determined.
Tubulin (14 μM) was incubated with CEP41 (1, 2, 3, and 5 μM), followed by the addition of taxol
(4 μM) and GTP (1 mM), and the reaction kinetics were measured. The rate of tubulin
polymerization was calculated from the slope of the linear part (elongation phase) of the
polymerization curve.

Sedimentation assay

Tubulin (30 μM) was polymerized with CEP41 (1, 2, 3, and 5 μM) in PEM buffer and GTP (1 mM)
at 37°C for 30 min. The reaction mixture was ultracentrifuged at 50,000 g at 30°C for 30 min. The
supernatant was separated, and the pellet was washed with warm PEM buffer with 10% DMSO.
The pellet was dissolved in SDS running buffer, and equal volumes of supernatant and pellet
were resolved on 15% SDS-PAGE. Similarly, the effect of CEP41 on taxol-induced tubulin
polymerization was determined. Tubulin (14 μM) was incubated with CEP41 (1, 2, 3, and 5 μM)
for 20 min on ice in PEM buffer, followed by the addition of taxol (4 μM) and GTP (1 mM). The
reaction mixture was polymerized for 30 min at 37°C, followed by centrifugation and SDS-PAGE
analysis.

Transmission electron microscopy

Tubulin (30 μM) was polymerized in the presence of CEP41 (5 μM) and GTP (1 mM) in PEM
buffer for 30 min at 37°C, and the microtubules formed were imaged in an electron microscope as
described before (Asthana et al., 2012). Briefly, the polymerized mixture was spotted on carbon-
formvar coated copper grids, air-dried, washed with double distilled water, and stained with

0.75% uranyl acetate. The grids were imaged in an electron microscope, JEM 2100 ultra HRTEM
instrument at 200 kV (JEOL). The experiment was performed three times.

Dilution-induced disassembly assay

The effect of CEP41 on dilution-induced microtubule depolymerization was determined, as
described earlier (Asthana et al., 2012). Tubulin (20 μM) was polymerized with 10% DMSO and 1
mM GTP in PEM buffer. The polymerized microtubules were diluted 10-fold in PEM buffer
containing 1 mM GTP and CEP41 (1, 3, and 5 μM) and incubated for 10 min at 37°C. The
reaction mixture was then ultracentrifuged at 60,000 g for 20 min at 30°C. The supernatant was
separated, and the pellet was gently washed with warm PEM buffer containing 10% DMSO. The
pellet was dissolved in SDS running buffer, and equal volumes of the soluble and polymer
fractions were resolved on 12% SDS-PAGE gel, followed by immunoblotting using anti-α-tubulin
and anti-CEP41 antibodies.

Immunofluorescence and microscopy

Cells were seeded at 50,000 cells/mL density on clean glass coverslips in 24-well plates and
incubated overnight. After incubation, cells were transfected with desired plasmids and incubated
for 24 h, followed by fixation and immunostaining, as described earlier (Asthana et al., 2013;
Kumari and Panda, 2022). Briefly, the cells were fixed with 3.7% formaldehyde, followed by
permeabilization with 0.1% Triton-X 100 and blocking with 4% BSA, and then incubated with
primary antibodies in BSA for 3 h at RT, followed by incubation with secondary antibodies for 1 h
at RT. DNA was stained with Hoechst (10 μg/ml) for 15 min at RT, and following three washes,
the coverslips were mounted and cured overnight. Primary antibodies used were mouse anti-α-
tubulin (T9026, Sigma-Aldrich) and rabbit anti-acetyl-alpha-tubulin (Lys40) (SAB5600134, Sigma-
Aldrich) at 1:400. Secondary antibodies were anti-mouse IgG (Invitrogen, A11005) and anti-rabbit
IgG (Invitrogen, A11012) coupled with Alexa Fluor 594, at 1:400. Images were acquired in z-
stacks in a spinning-disc CSU-X1 (Yokogawa) or a laser scanning LSM 780 (Zeiss) confocal
microscope using a 63x oil immersion lens. The exposure and gain settings were kept identical
among the control and treated groups to quantify fluorescence intensities. Images were
processed in Zen Blue software, and intensities were quantified using ImageJ. Colocalization
analyses of GFP-expressing cells were done using the Coloc 2 plugin available in ImageJ Fiji
software. Measurement of cell area was done in Zen Blue software by drawing cell boundaries
based on microtubule staining. All experiments were performed thrice, with a minimum of 50 cells
quantified for each group.

For the nocodazole-induced disassembly assay, HeLa cells transiently transfected with
CEP41-GFP and GFP constructs were incubated for 24 h and then treated with 500 nM

nocodazole (Sigma-Aldrich) for 1 h and subjected to immunostaining as described above. For

imaging CEP41-expressing mitotic cells, HeLa cells were transfected with CEP41-GFP and GFP

constructs and then synchronized by double thymidine block using 2 mM thymidine (Sigma-

Aldrich) (Ma and Poon, 2011). Post 8 h of thymidine release, cells were fixed and immunostained.

In vitro microtubule staining

The microtubules formed in the presence of CEP41 were visualized by immunostaining (Asthana

et al., 2012; Girotra et al., 2017). Tubulin (30 µM) was polymerized with CEP41 (5 µM) and GTP

(1 mM) in PEM buffer for 30 min at 37°C, and the polymerized solution was loaded on clean poly-

L-lysine coated coverslips, incubated for 20 min, and blot dried. The coverslips were fixed with

3.7% formaldehyde and immunostained using an α-tubulin antibody. The experiment was

performed three times.

CEP41 depletion using shRNA

The endogenous level of CEP41 in NIH3T3 cells was depleted using shRNA, as described before

(Kumari et al., 2023). Cells were seeded in 60 mm dishes at 50-60% confluency and transiently

transfected with CEP41 shRNA (5'-GCTTACAGTTACCCAATTGCA-3', TRCN0000143499,

Sigma-Aldrich) or scramble shRNA (SHC002, Sigma-Aldrich) using Lipofectamine 3000. Post-

transfection, the cells were incubated for 48 h, harvested, and lysed. The levels of CEP41 were

determined by western blotting using anti-CEP41 and anti-β-actin (loading control) antibodies. For

immunostaining of depleted cells, NIH3T3 cells were grown on coverslips in 24-well plates,

transfected with CEP41 and scrambled shRNA, and 48 h post-transfection subjected to

immunostaining as described above.

Cell proliferation assay

Sulforhodamine B (SRB) assay was performed to determine the effect of CEP41 depletion on cell

proliferation. NIH3T3 cells were seeded in 96-well plates (8000 cells/well) and transfected with

CEP41 and scramble shRNA, as mentioned above. After 48 h, the cells were fixed with chilled

50% trichloroacetic acid and processed for the SRB assay as described before (Ray et al., 2007;

Skehan et al., 1990). The percentage inhibition of cell proliferation was determined, and the

experiment was performed four times.

Cell cycle analysis

587 The effect of CEP41 depletion on cell cycle progression was determined by flow cytometric

588 analysis (Gajula et al., 2013; Mohan and Panda, 2008). CEP41-depleted and scrambled control

589 cells were grown in puromycin (4 µg/ml)-containing medium for 2 weeks to enrich shRNA-

590 expressing cells. The cells were then synchronized by a double thymidine block and 8 h post-

thymidine release, trypsinized, fixed with 70% ethanol, and stained with propidium iodide (30 µg/ml). Using a BD FACSAria flow cytometer (Becton Dickinson), 2000 cells were analyzed in each group, and the experiment was performed twice. The data were processed in the FlowJo software (Tree Star).

In silico analysis

The secondary and tertiary structures of CEP41 were predicted using PSI-PRED (McGuffin et al., 2000) and AlphaFold (Jumper et al., 2021; Varadi et al., 2022), respectively. The image was created using PyMoL (DeLano, 2002). Coiled-coil regions in CEP41 were predicted using Ncoils in the Waggawagga prediction software (Simm et al., 2015). Sequence alignment of CEP41 orthologues was carried out using Clustal Omega (Sievers et al., 2011).

Statistics

Data were analyzed and plotted using GraphPad Prism version 8.0.2 and represented as mean ± standard deviation with error bars denoting the standard deviation. Experiments were performed as three independent sets unless mentioned otherwise. An unpaired, two-tailed *t*-test was used to determine statistical significance between the mean of two groups, and for significance between three or more groups, a one-way analysis of variance (ANOVA) was applied, followed by Tukey's multiple-comparison test. $P < 0.05$ was considered statistically significant.

Acknowledgments

The authors thank IIT Bombay for the central facilities of spinning disk and laser scanning confocal microscopes, FACS, and cryo-HRTEM. We thank Prof. Subba Rao Gangi Shetty (IISc, Bangalore) for providing the shRNA clones of CEP41. S.S.P. thanks the Council of Scientific and Industrial Research, Government of India for her doctoral fellowship.

Competing interests

The authors declare no competing or financial interests.

Funding

This work was supported by the JC Bose National Fellowship (JCB/2019/000016) awarded to D.P. from the Department of Science and Technology, Government of India.

Data availability

All relevant data can be found within the article and its supplementary information.

References

Andersen, J. S., Wilkinson, C. J., Mayor, T., Mortensen, P., Nigg, E. A. and Mann, M. (2003).
Proteomic characterization of the human centrosome by protein correlation profiling. *Nature*
426, 570-574.

Asthana, J., Kuchibhatla, A., Jana, S. C., Ray, K. and Panda, D. (2012). Dynein light chain 1
(LC8) association enhances microtubule stability and promotes microtubule bundling.
Journal of Biological Chemistry **287**, 40796-40805.

Asthana, J., Kapoor, S., Mohan, R. and Panda, D. (2013). Inhibition of HDAC6 deacetylase
activity increases its binding with microtubules and suppresses microtubule dynamic
instability in MCF-7 cells. *Journal of Biological Chemistry* **288**, 22516-22526.

Bettencourt-Dias, M., Hildebrandt, F., Pellman, D., Woods, G. and Godinho, S. A. (2011).
Centrosomes and cilia in human disease. *Trends in Genetics* **27**, 307-315.

Bodakuntla, S., Jijumon, A. S., Villablanca, C., Gonzalez-Billault, C. and Janke, C. (2019).
Microtubule-Associated Proteins: Structuring the Cytoskeleton. *Trends Cell Biol* **29**, 804-
819.

Bompard, G., van Dijk, J., Cau, J., Lannay, Y., Marcellin, G., Lawera, A., van der Laan, S.
and Rogowski, K. (2018). CSAP Acts as a Regulator of TTLL-Mediated Microtubule
Glutamylation. *Cell Rep* **25**, 2866-2877.

Bordo, D. and Bork, P. (2002). The rhodanese/Cdc25 phosphatase superfamily. Sequence-
structure-function relations. *EMBO Rep* **3**, 741-746.

Bradford, M. (1976). A Rapid and Sensitive Method for the Quantitation of Microgram Quantities
of Protein Utilizing the Principle of Protein-Dye Binding. *Anal Biochem* **72**, 248-254.

Bre, M. H. and Karsenti, E. (1990). Effects of brain microtubule-associated proteins on
microtubule dynamics and the nucleating activity of centrosomes. *Cell Motil Cytoskeleton*
15, 88-98.

Chaudhary, V., Venghateri, J. B., Dhaked, H. P. S., Bhoyar, A. S., Guchhait, S. K. and
Panda, D. (2016). Novel Combretastatin-2-aminoimidazole Analogues as Potent Tubulin
Assembly Inhibitors: Exploration of Unique Pharmacophoric Impact of Bridging Skeleton and
Aryl Moiety. *J Med Chem* **59**, 3439-3451.

Conduit, P. T., Wainman, A. and Raff, J. W. (2015). Centrosome function and assembly in
animal cells. *Nat Rev Mol Cell Biol* **16**, 611-624.

DeLano, W. L. (2002). The PyMOL Molecular Graphics System, Version 1.1. Schrodinger LLC.

Doxsey, S. (2001). Re-evaluating centrosome function. *Nat Rev Mol Cell Biol* **2**, 688-698.

Doxsey, S., Zimmerman, W. and Mikule, K. (2005). Centrosome control of the cell cycle.
Trends Cell Biol **15**, 303-311.

Gache, V., Waridel, P., Winter, C., Juhem, A., Schroeder, M., Shevchenko, A. and Popov, A.
V. (2010). Xenopus meiotic microtubule-associated interactome. *PLoS One* **5**, e9248.

Gajula, P. K., Asthana, J., Panda, D. and Chakraborty, T. K. (2013). A synthetic dolastatin 10
analogue suppresses microtubule dynamics, inhibits cell proliferation, and induces apoptotic
cell death. *J Med Chem* **56**, 2235-2245.

Girotra, M., Srivastava, S., Kulkarni, A., Barbora, A., Bobra, K., Ghosal, D., Devan, P., Aher,
A., Jain, A., Panda, D., et al. (2017). The C-terminal tails of heterotrimeric kinesin-2 motor
subunits directly bind to α-tubulin1: Possible implications for cilia-specific tubulin entry.
Traffic **18**, 123-133.

Graser, S., Stierhof, Y. D., Lavoie, S. B., Gassner, O. S., Lamla, S., Le Clech, M. and Nigg, E.
A. (2007). Cep164, a novel centriole appendage protein required for primary cilium
formation. *Journal of Cell Biology* **179**, 321-330.

Hamel, E. and Lin, C. M. (1981). Glutamate-Induced Polymerization of Tubulin - Characteristics
of the Reaction and Application to the Large-Scale Purification of Tubulin. *Arch Biochem*
Biophys **209**, 29-40.

Izawa, I., Goto, H., Kasahara, K. and Inagaki, M. (2015). Current topics of functional links
between primary cilia and cell cycle. *Cilia* **4**, 1-13.

Jackson, P. K. (2011). Do cilia put brakes on the cell cycle? *Nat Cell Biol* **13**, 340-342.

Jakobsen, L., Vanselow, K., Skogs, M., Toyoda, Y., Lundberg, E., Poser, I., Falkenby, L. G.,
Bennetzen, M., Westendorf, J., Nigg, E. A., et al. (2011). Novel asymmetrically localizing
components of human centrosomes identified by complementary proteomics methods.
EMBO Journal **30**,.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly accurate
protein structure prediction with AlphaFold. *Nature* **596**, 1520-1535.

Karr, T. L. and Purich, D. L. (1979). A microtubule assembly/disassembly model based on drug
effects and depolymerization kinetics after rapid dilution. *Journal of Biological Chemistry*
254, 10885-10888.

Ki, S. M., Kim, J. H., Won, S. Y., Oh, S. J., Lee, I. Y., Bae, Y., Chung, K. W., Choi, B., Park, B.,
Choi, E., et al. (2020). CEP41-mediated ciliary tubulin glutamylation drives angiogenesis
through AURKA -dependent deciliation. *EMBO Rep* **21**, e48290.

Korvatska, O., Estes, A., Munson, J., Dawson, G., Bekris, L. M., Kohen, R., Yu, C. E.,
Schellenberg, G. D. and Raskind, W. H. (2011). Mutations in the TSGA14 gene in families
with autism spectrum disorders. *American Journal of Medical Genetics, Part B:*
Neuropsychiatric Genetics **156**, 303-311.

Kuchnir Fygenson, D., Flyvbjerg, H., Sneppen, K., Libchaber, A. and Leibler, S. (1995).
Spontaneous nucleation of microtubules. *Phys Rev E* **51**, 5058.

Kumar, N. (1981). Taxol-induced polymerization of purified tubulin. Mechanism of action. *Journal*
of Biological Chemistry **256**, 10435-10441.

Kumari, A. and Panda, D. (2018). Regulation of microtubule stability by centrosomal proteins.
IUBMB Life **70**, 602-611.

Kumari, A. and Panda, D. (2022). Monitoring the Disruptive Effects of Tubulin-Binding Agents on
Cellular Microtubules. In *Microtubules: Methods and Protocols* (ed. H. Inaba), pp. 431-448.
Humana, New York, NY: Springer.

Kumari, A., Shriwas, O., Sisodiya, S., Santra, M. K., Guchhait, S. K., Dash, R. and Panda, D.
(2021). Microtubule-targeting agents impair kinesin-2-dependent nuclear transport of β-
catenin: Evidence of inhibition of Wnt/β-catenin signaling as an important antitumor
mechanism of microtubule-targeting agents. *FASEB Journal* **35**, e21539.

Kumari, A., Prassanawar, S. S. and Panda, D. (2023). β-III Tubulin Levels Determine the
Neurotoxicity Induced by Colchicine-Site Binding Agent Indibulin. *ACS Chem Neurosci* **14**,
19-34.

Lee, J. E., Silhavy, J. L., Zaki, M. S., Schroth, J., Bielas, S. L., Marsh, S. E., Olvera, J.,
Brancati, F., Iannicelli, M., Ikegami, K., et al. (2012). CEP41 is mutated in Joubert
syndrome and is required for tubulin glutamylation at the cilium. *Nat Genet* **44**, 193-199.

Ligon, L. A., Shelly, S. S., Tokito, M. K. and Holzbaur, E. L. F. (2006). Microtubule binding
proteins CLIP-170, EB1, and p150Glued form distinct plus-end complexes. *FEBS Lett* **580**,
1327-1332.

Ma, H. T. and Poon, R. Y. C. (2011). Synchronization of HeLa Cells. In *Methods in Molecular*
Biology (ed. G. Banfalvi), pp 151-161. Humana, New York, NY: Springer

McGuffin, L. J., Bryson, K. and Jones, D. T. (2000). The PSIPRED protein structure prediction
server. *Bioinformatics* **16**, 404-405.

Mohan, R. and Panda, D. (2008). Kinetic stabilization of microtubule dynamics by estramustine
is associated with tubulin acetylation, spindle abnormalities, and mitotic arrest. *Cancer Res*
68, 6181-6189.

Nigg, E. A. (2007). Centrosome duplication: of rules and licenses. *Trends Cell Biol* **17**, 215-221.

Nigg, E. A. and Raff, J. W. (2009). Centrioles, Centrosomes, and Cilia in Health and Disease.
Cell **139**, 663-678.

Panda, D., Rathinasamy, K., Santra, M. K. and Wilson, L. (2005). Kinetic suppression of
microtubule dynamic instability by griseofulvin: implications for its possible use in the
treatment of cancer. *Proceedings of the National Academy of Sciences* **102**, 9878-9883.

Patowary, A., Won, S. Y., Oh, S. J., Nesbitt, R. R., Archer, M., Nickerson, D., Raskind, W. H.,
Bernier, R., Lee, J. E. and Brkanac, Z. (2019). Family-based exome sequencing and case-
control analysis implicate CEP41 as an ASD gene. *Transl Psychiatry* **9**, 4.

Piperno, G., LeDizet, M. and Chang, X. J. (1987). Microtubules containing acetylated alpha-
tubulin in mammalian cells in culture. *J Cell Biol* **104**, 289-302.

Prassanawar, S. S. and Panda, D. (2019). Tubulin heterogeneity regulates functions and
dynamics of microtubules and plays a role in the development of drug resistance in cancer.
Biochemical Journal **476**, 1359-1376.

Ray, S., Mohan, R., Singh, J. K., Samantaray, M. K., Shaikh, M. M., Panda, D. and Ghosh, P.
(2007). Anticancer and antimicrobial metallopharmaceutical agents based on palladium,
gold, and silver N-heterocyclic carbene complexes. *J Am Chem Soc* **129**, 15042-15053.

Reiter, J. F. and Leroux, M. R. (2017). Genes and molecular pathways underpinning
ciliopathies. *Nat Rev Mol Cell Biol* **18**, 533-547.

Robert, A., Margall-Ducos, G., Guidotti, J. E., Brégerie, O., Celati, C., Bréchot, C. and
Desdouets, C. (2007). The intraflagellar transport component IFT88/polaris is a
centrosomal protein regulating G1-S transition in non-ciliated cells. *J Cell Sci* **120**, 628-637.

Roostalu, J. and Surrey, T. (2017). Microtubule nucleation: Beyond the template. *Nat Rev Mol*
Cell Biol **18**, 702-710.

Salisbury, J. L. (2003). Centrosomes: Coiled-coils organize the cell center. *Current Biology* **13**,
R88-R90.

Schatz, C. A., Santarella, R., Hoenger, A., Karsenti, E., Mattaj, I. W., Gruss, O. J. and
Carazo-Salas, R. E. (2003). Importin α-regulated nucleation of microtubules by TPX2.
EMBO Journal **22**,.

Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., Lopez, R., McWilliam, H.,
Remmert, M., Söding, J., et al. (2011). Fast, scalable generation of high-quality protein
multiple sequence alignments using Clustal Omega. *Mol Syst Biol* **7**, 2060-2070.

Simm, D., Hatje, K. and Kollmar, M. (2015). Waggawagga: Comparative visualization of coiled-
coil predictions and detection of stable single α-helices (SAH domains). *Bioinformatics* **31**,
767-769.

Skehan, P., Storeng, R., Scudiero, D., Monks, A., Mcmahon, J., Vistica, D., Warren, J. T.,
Bokesch, H., Kenney, S. and Boyd, M. R. (1990). New colorimetric cytotoxicity assay for
anticancer-drug screening. *J Natl Cancer Inst* **82**, 1107-1112.

Srivastava, S. and Panda, D. (2018). A centrosomal protein STARD9 promotes microtubule
stability and regulates spindle microtubule dynamics. *Cell Cycle* **17**, 2052-2068.

Tiryaki, F., Deretic, J. and Firat-Karalar, E. N. (2022). ENKD1 is a centrosomal and ciliary
microtubule-associated protein important for primary cilium content regulation. *FEBS*
Journal **289**, 3789-3812.

Tovey, C. A. and Conduit, P. T. (2018). Microtubule nucleation by γ-tubulin complexes and
beyond. *Essays Biochem* **62**, 765-780.

Vallee, R. B. (1986). Reversible assembly purification of microtubules without assembly-
promoting agents and further purification of tubulin, microtubule-associated proteins, and
MAP fragments. *Methods Enzymol* **134**, 89-104.

Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova, G., Yuan, D.,
Stroe, O., Wood, G., Laydon, A., et al. (2022). AlphaFold Protein Structure Database:
Massively expanding the structural coverage of protein-sequence space with high-accuracy
models. *Nucleic Acids Res* **50**, D439-D444.

Vertii, A., Bright, A., Delaval, B., Hehnly, H. and Doxsey, S. (2015). New frontiers: discovering
cilia-independent functions of cilia proteins. *EMBO Rep* **16**, 1275-1287.

Voter, W. A. and Erickson, H. P. (1984). The kinetics of microtubule assembly. Evidence for a
two-stage nucleation mechanism. *Journal of Biological Chemistry* **259**, 10430-10438.

Wang, W. J., Tay, H. G., Soni, R., Perumal, G. S., Goll, M. G., MacAluso, F. P., Asara, J. M.,
Amack, J. D. and Bryan Tsou, M. F. (2013). CEP162 is an axoneme-recognition protein
promoting ciliary transition zone assembly at the cilia base. *Nat Cell Biol* **15**, 591-601.

 Figure legends

Fig 1. CEP41 interacts with tubulin and binds to microtubules. Co-immunoprecipitation of

CEP41 and α-tubulin from NIH3T3 cells. The whole cell lysate was incubated with **(A)** CEP41 IgG

or **(B)** α-tubulin IgG and immunoprecipitated using Protein A agarose beads. The inputs and

eluates (IP) were immunoblotted with indicated antibodies. The experiments were performed

twice, and representative blots are shown. **(C)** CEP41 was present in the MAP-rich tubulin

fraction isolated from goat brain tissue. MAP-rich tubulin was isolated from a goat brain by two

cycles of polymerization and depolymerization using 4 M glycerol. An equal concentration of

fractions was loaded on SDS-PAGE gel and immunoblotted with CEP41 antibody. The

experiment was performed twice, and a representative blot is shown. **(D)** Coomassie Brilliant Blue

stained SDS-PAGE gel image of purified tubulin, his-CEP41, and his-tag cleaved CEP41. **(E)**

Representative SDS-PAGE gel image (top) and immunoblot (bottom) of co-sedimentation assay

performed with CEP41 and preformed microtubules. Taxol-stabilized microtubules (3 μM) were

incubated with indicated concentrations of CEP41 at 37°C. The polymers were sedimented by

ultra-centrifugation, and an equal volume of the pellet fraction was resolved on SDS-PAGE gel

and immunoblotted with anti-CEP41 antibody (bottom). **(F)** Band intensities of CEP41 in the pellet

fraction were quantified using ImageJ. The experiment was performed three times, and error bars

denote the standard deviation. Statistical significance was determined by Student's t-test (*$P <$

0.05; **$P < 0.01$; ***$P < 0.001$).

Fig 2. CEP41 promoted tubulin assembly in the presence of GTP. (A) *In vitro* polymerization

of tubulin (30 μM) without (●) or with 1 (■), 2 (▲), 3 (▼), and 5 (♦) μM CEP41 and 1 mM GTP

was monitored at 37°C. Bovine serum albumin (●) and alkaline phosphatase (■) were negative

controls. The experiment was performed thrice, and a representative plot is shown. **(B)** The rate

of tubulin polymerization was calculated from the slope of the elongation phase (**$P < 0.01$; ***P

< 0.001). Error bars represent the standard deviation from three independent sets of experiments.

(C) Tubulin (30 μM) was polymerized in the absence or presence of 1, 2, 3, and 5 μM CEP41 for

45 min at 37°C, and the soluble and polymer fractions were separated by ultra-centrifugation. An

equal volume of the fractions was resolved on SDS-PAGE gel; a representative gel image is

shown. **(D)** Band intensities of tubulin in the soluble and the polymer fractions were quantified

using ImageJ, and the polymer/soluble ratio was determined (*$P < 0.05$; ***$P < 0.001$; ****$P <$

0.0001). Error bars represent the standard deviation from three independent sets of experiments.

(E) Transmission electron microscopy images of microtubules formed in the presence of CEP41.

Tubulin (30 μM) was polymerized in the presence of 5 μM CEP41 for 45 min at 37°C. The scale

bar is 1 μm. Black arrows indicate microtubule branching. **(F)** Tubulin was polymerized in the

same manner mentioned above. The reaction mixture was loaded on poly-L-lysine coated

coverslips, immunostained with α-tubulin antibody, and imaged in a spinning-disk confocal

microscope at 63x magnification. The scale bar is 10 μm. The experiments were performed three
times, and representative images are shown. Statistical significance was determined by Student's
t-test.

Fig 3. CEP41 enhanced the rate of taxol-induced tubulin polymerization and prevented
dilution-induced microtubule disassembly. (A) The effect of CEP41 on taxol-induced tubulin
polymerization was monitored at 37°C. Tubulin (15 μM) was incubated with different
concentrations of CEP41 (0 (●), 1 (■), 2 (▲), 3 (▼), 5 (♦) μM) followed by taxol (4 μM) and GTP
(1 mM). The experiment was performed thrice, and a representative plot is shown. **(B)** The rate of
tubulin polymerization was calculated from the slope of the elongation phase (log phase of the
polymerization curve in **A** (*$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; ****$P < 0.0001$). Error bars
represent the standard deviation from three independent sets of experiments. **(C)** Tubulin (15 μM)
was polymerized in the presence of taxol (4 μM), and different concentrations (0, 1, 2, 3, and 5
μM) of CEP41 for 30 min at 37°C and the soluble and polymer fractions were separated by ultra-
centrifugation. An equal volume of soluble and polymer fractions was resolved on SDS-PAGE
gel, and a representative gel image is shown. **(D)** Band intensities of tubulin in the soluble and the
polymer fraction were quantified using ImageJ, and the polymer/soluble ratio was calculated (*$P <$
0.05; **$P < 0.01$; ***$P < 0.001$). Error bars represent the standard deviation from three
independent sets of experiments. **(E)** Representative immunoblot of dilution-induced disassembly
assay performed with CEP41 and preformed microtubules. Microtubules were polymerized in the
presence of DMSO and diluted 10-fold, followed by incubation with indicated concentrations of
CEP41 at 37°C. The polymers were sedimented by ultra-centrifugation, and equal volumes of the
supernatant and pellet fractions were resolved on SDS-PAGE gel and immunoblotted with anti-α-
tubulin (top) and anti-CEP41 antibody (bottom). **(F)** Band intensities of tubulin in the soluble and
the polymer fractions were quantified using ImageJ, and the polymer/soluble ratio was calculated.
The experiment was performed three times, and error bars denote the standard deviation.
Statistical significance was determined by Student's *t*-test (**$P < 0.01$; ***$P < 0.001$; ****$P <$
0.0001).

Fig 4. Exogeneously expressed CEP41 colocalized with microtubules and associated with
spindle poles and midbody in HeLa cells. (A) Fluorescence microscopy images of HeLa cells
transfected with CEP41-GFP or GFP constructs. After 24 h of transfection, cells were fixed with
3.7% paraformaldehyde and immunostained for α-tubulin (red). DNA was stained with Hoechst
(blue). The scale bar is 10 μm. **(B)** Pearson's correlation coefficient (*r*) of GFP-expressing cells
was determined using the Coloc2 plugin in ImageJ. Data represent the mean ± standard
deviation of three independent experiments with 170 cells analyzed in each group. **(C)**
Localization of CEP41-GFP at different stages of the cell cycle. A double thymidine block was
used to synchronize HeLa cells transfected with either CEP41-GFP or GFP constructs to enrich

mitotic cells. The cells were fixed and stained for α-tubulin (red) and DNA (blue). Images were

captured in a laser scanning confocal microscope at 63x magnification. Scale bar: 5 μm, and the

insets show magnification of the midbody. The bottom last panel is for GFP control showing no

GFP localization at the spindle poles.

Fig 5. CEP41 interacted with microtubules via its RHOD domain and the N-terminal region.

(A) AlphaFold-predicted structure of CEP41. Color code, N-terminal region: pink, coiled-coil

motifs: dark blue, RHOD domain: orange, C-terminal region: cyan. The image was created using

PyMoL. (B) Deletion constructs of CEP41. Cyan box (filled) indicates the predicted nuclear exit

sequence, and grey boxes indicate the predicted coiled-coil motifs. **(C)** Fluorescence microscopy

images of HeLa cells transfected with GFP-tagged full-length and truncated CEP41 constructs

(green). Transfected cells were immunostained for α-tubulin (red) 24 h post-transfection. DNA

was stained with Hoechst (blue). Images were captured in a spinning-disk confocal microscope at

63x magnification. Scale bar: 10 μm. **(D)** Summarized results of the localization of GFP-tagged

truncated CEP41. The experiment was performed twice, with 100 cells analyzed in each group.

Fig 6. Overexpression of CEP41 induced the formation of stable microtubule bundles. (A)

Fluorescence microscopy images of HeLa cells transfected with CEP41-GFP or GFP constructs

and immunostained for α-tubulin (red). DNA was stained with Hoechst (blue). The scale bar is 10

μm, and white arrows point to microtubule bundles. Panels, i and ii, show different forms of

microtubule bundles formed in HeLa cells upon CEP41 overexpression. **(B)** Percentage of cells

with microtubule bundles plotted for GFP and CEP41-GFP expressing cells. The data are an

average of three independent experiments with 300 cells counted in each case (**$P < 0.01$). **(C)**

Microtubule intensity of 250 cells in each case was quantified using ImageJ (****$P < 0.0001$).

Data represent the mean ± standard deviation of three independent experiments. **(D)** CEP41

overexpression enhanced the level of tubulin acetylation. HeLa cells transfected with CEP41-GFP

and GFP constructs were stained for acetylated tubulin (red) and DNA (blue), scale bar: 10 μm.

The intensity of acetylated tubulin per cell was quantified for 150 cells in each case (****$P <$

0.0001). **(E)** Exogenous expression of CEP41 stabilized microtubules against nocodazole-

induced depolymerization. HeLa cells transfected with GFP, or CEP41-GFP constructs, were

treated with nocodazole (500 nM) for 1 h and stained for α-tubulin (red) and DNA (blue), scale

bar: 10 μm. The white arrows mark microtubule bundles. Microtubule intensity of 150 cells in each

case was quantified using ImageJ (****$P < 0.0001$). The experiments were performed three times,

and error bars denote standard deviation. Statistical significance was determined by Student's *t*-

test.

Fig 7. CEP41 depletion disrupted the interphase microtubule organization in NIH3T3 cells.

(A) Representative immunoblot showing CEP41 levels in NIH3T3 cells transfected with

scrambled or CEP41 shRNA. β-actin was used as a loading control. **(B)** Band intensities were

quantified using ImageJ, and the CEP41/β-actin intensity ratios were plotted (ns $P > 0.05$; **$P <$

0.01). Error bars represent the standard deviation from three independent sets of experiments.

(C) Fluorescence microscopy images of NIH3T3 cells transfected with scrambled or CEP41

shRNA. Transfected cells were fixed and stained for α-tubulin (red) and DNA (blue). Images were

captured in a spinning-disk confocal microscope at 63x magnification. Scale bar: 10 μm. **(D)**

Microtubule intensity and **(E)** Cell area of 250 cells in each case were quantified using ImageJ (ns

$P > 0.05$; *$P < 0.05$; **$P < 0.01$; ****$P < 0.0001$). Data represent the mean ± standard deviation of

three independent sets. Statistical significance was determined by Student's t-test.

Fig 8. CEP41 depletion inhibited proliferation and cell cycle progression in NIH3T3 cells.

(A) CEP41 depletion inhibited cell proliferation. NIH3T3 cells transfected with scrambled or

CEP41 shRNA. Percentage inhibition of cell proliferation was determined with respect to control.

The data were an average of four independent sets represented as mean ± standard deviation

(**$P < 0.01$). Statistical significance was determined by Student's t-test. **(B)** CEP41-depleted cells

failed to progress to the G2/M phase. NIH3T3 cells transfected with scrambled or CEP41 shRNA

were synchronized by a double thymidine block and, 8 h post-thymidine release, were stained

with propidium iodide and subjected to flow cytometry analysis. Representative cytograms of

control, scrambled control, and CEP41-depleted cells are shown. Control (0 h) denotes cells

synchronized at the G1 stage. **(C)** Percentage of cells at different cell cycle stages. The data

represent mean ± SEM of two independent sets with 2000 cells analyzed in each case (****$P <$

0.0001). Statistical significance was determined by one-way ANOVA test.

D

	Localization		
	Microtubules	**Spindle poles**	**Midbody**
FL	+	+	+
ΔN	-	-	-
ΔRHOD	-	-	-
ΔC	+	+	+
ΔNC	-	-	-