Steroid signaling controls sex-specific development in an invertebrate

Lydia Grmai[1,♦], Erin Jimenez[1], Ellen Baxter[1], Mark Van Doren[1,*]

[1]Department of Biology, Johns Hopkins University, Baltimore, Maryland, USA

[♦]Current address: Department of Cell Biology, University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania, USA

*author correspondence: vandoren@jhu.edu

Summary

In vertebrate sexual development, two important steroid hormones – testosterone and estrogen – regulate the sex-specific development of many tissues. In contrast, invertebrates utilize a single steroid hormone, ecdysone, to regulate developmental timing in both sexes. Here, we show that in the fruit fly *Drosophila melanogaster*, sex-specific ecdysone activity controls important aspects of gonad sexual dimorphism. Rather than being regulated at the level of hormone production, hormone activity is regulated cell-autonomously through sex-specific hormone reception. We found that *Ecdysone receptor (EcR)* is regulated downstream of the sex determination factor Doublesex (Dsx), the founding member of the Dsx/Mab3 Related Transcription Factor (DMRT) family that regulates gonad development in all animals. EcR is restricted to the developing ovary and repressed in the testis at a time when ecdysone initiates ovary morphogenesis. EcR activity drives ovary development and antagonizes development of the testis stem cell niche. Interestingly, de-repression of the ecdysone response in the testis led to rapid feminization, which is known to cause infertility. This work demonstrates that invertebrates can also use steroid hormone signaling to control sex-specific development. Further, it may help explain recent work showing that vertebrate sexual development is surprisingly cell-autonomous. For example, in birds that have a mixture of cells with male and female genotypes, the male cells develop as male and the female cells develop as female despite exposure to the same circulating hormones. Sex-specific regulation of steroid hormone response in vertebrates may well underly such cellular sexual fate choices.

Main

Sexual dimorphism is a hallmark feature of nearly all metazoans and encompasses all discernible differences between males and females of the same species. Sex-specific gonad development is instructed by members of the Doublesex/Mab-3 Related Transcription Factor (DMRT) family in all animals for which a role has been characterized, including flies, nematodes, chickens, mice, and humans. Loss-of-function studies in both flies and mice have underscored the importance of Dsx and DMRTs in sex-specific gonad development. Loss of *Drosophila doublesex (dsx)* impairs sexual development of the gonad[1,2] and loss of sexual identity in the adult gonad causes infertility[3]. In mice, DMRT1 is expressed in the developing testis and deletion of the *DMRT1* locus causes defects in spermatogenesis and germline loss[4]. Additionally, altering *DMRT1* expression in the adult mouse gonad causes sex reversal and gametogenesis defects[5–7].

The *Drosophila dsx* locus encodes two sex-specific protein isoforms - Dsx^M and Dsx^F - that bear identical DNA binding domains and unique C-termini that are thought to recruit distinct interacting partners[8]. As a result, they each drive unique transcriptional programs that instruct most known aspects of sexual differentiation, including abdominal pigmentation, genital development, some aspects of courtship behavior, and gonadogenesis[9]. The somatic sex determination cascade in *Drosophila* initiates with the expression and activity of female-specific RNA-binding proteins Sex-lethal (Sxl) and Transformer (Tra). Tra binds newly transcribed *dsx* RNA to elicit a female splicing pattern that yields the female-specific protein isoform, Dsx^F. In the absence of Sxl/Tra, default *dsx* splicing yields the male isoform, Dsx^M. Dsx is required for proper sexual differentiation of the gonad stem cell niche[2], which in both sexes is established by somatic cells at the apical end of the gonad that anchor self-renewing germline stem cells (GSCs). An important component of the testis GSC niche, the "hub," consists of a tight cluster of somatic cells surrounded by GSCs. In contrast, in adult ovaries each of the 16-20 ovarioles contain GSC niches, important components of which are "cap cells (CCs)" contacting the GSCs and apically localized stacks of 6-7 "terminal filament (TF)" cells that are flatter in morphology. Finally, the timing of gonad stem cell niche establishment is sexually dimorphic; the testis hub is established by the end of embryogenesis (24 hrs after fertilization, AF)[10], while female TFs and CCs form several days later in the third larval instar (L3, 4.5 days AF)[11].

Mammalian gonads are the primary site of sex hormone synthesis and are key effectors of sexual development[12,13]. Testosterone and estrogen, produced in male and female gonads respectively, instruct many secondary sex characteristics throughout the organism[14]. Defects in steroid hormone synthesis and/or signaling are associated with common Differences of Sex Development such as congenital adrenal hyperplasia and androgen insensitivity syndrome[15]. Ecdysone is the dominant steroid hormone in insects like *Drosophila* and regulates larval molting, gut regeneration, and adult female reproduction[16–19]. Pulses of ecdysone

hormone are produced from the prothoracic gland (PG) and secreted into the hemolymph in the hours preceding each larval molt. The Ecdysone receptor (EcR) is a nuclear hormone receptor, similar to the mammalian testosterone and estrogen receptors, that is activated by the ecdysone derivative, 20-hydroxyecdysone (20E). EcR-dependent transcriptional activation requires a number of co-factors, including the mammalian RXR ortholog Ultraspiracle (Usp) and the SRC-3 homolog Taiman (Tai)[20]. Blocking ecdysone biosynthesis or ecdysone uptake or reducing EcR levels leads to developmental arrest and eventually death[21]. Studies on the role of ecdysone signaling in the gonad have largely focused on the adult ovary, where EcR is required for female GSC maintenance[17], border cell migration[22], and male somatic stem cell maintenance[23]. EcR target gene activation has been implicated in the establishment of the female gonad niche during development[11], although its role in regulating sex-specific gonadogenesis has not been examined.

Dsx regulates gonadal ecdysone signaling

We examined transcriptional activation via EcR in developing gonads using a transgenic reporter (*EcRE-lacZ*)[24,25,26] and protein detection of the EcR target Broad (Br)[11]. In late larval ovaries, *EcRE-lacZ* produced high levels of β-gal throughout the apical cap, including TF cells (**Fig. 1a**). Similarly, Br was expressed strongly in all somatic cells of the late larval ovary, including the developing TFs[11] (**Extended Data Fig. 1a**). In the larval testis, however, little to no *EcRE-lacZ* activity or Br expression was detected, including in the testis niche (**Fig. 1b, Extended Data Fig. 1b**). However, Br was detected in the testis terminal epithelium cells (**Extended Data Fig. 1c**), indicating that the testis environment receives 20E and supports ecdysone signaling in the cells that can respond. The Z1 isoform of Br is the most ecdysone-responsive isoform in the developing ovary[11] (**Extended Data Fig. 1d**), and Br-Z1 was consistently absent from the male stem cell niche (**Extended Data Fig. 1e**).

In mammals, sex-biased hormone activity is regulated at the level of hormone production: females produce higher levels of estrogen, while males produce higher levels of testosterone. In contrast, we found that a female bias in hormone levels could not alone account for the observed dimorphism in ecdysone activity: supplying exogenous 20E or a membrane-permeable, non-steroidal EcR agonist (chromafenozide) did not activate ecdysone signaling at a stage in which EcR transcriptional activity was inducible in ovaries (**Extended Data Fig. 2a-f**). Ecdysone is converted to its active form, 20E, via monooxygenation by the P450 enzyme Shade[27] (Shd). We found that depletion of *shd* from the ovary did not reproduce female niche defects seen upon blocking EcR activity[11] (**Extended Data Fig. 2g-h**), suggesting that our observed female bias in ecdysone activity is likely not due to differential 20E availability by local conversion in the larval ovary. This is consistent with our observation that cells of the testis terminal epithelium can respond to ecdysone signaling (**Extended Data Fig. 1c**).

We previously conducted a collaborative genome-wide search for putative Dsx targets using genomic and computational approaches[8]. Using a position-weight matrix, we identified 41 sequences across the *EcR* locus with high similarity to the Dsx consensus binding sequence, many of which were conserved in other *Drosophila* species[8,28]. Several Dsx occupancy regions overlapping with putative Dsx binding sites were also mapped to *EcR* by DamID-seq[8]. We therefore explored whether sex differences in *EcR* expression might account for sexually dimorphic ecdysone activity in the gonad. EcR was expressed at high levels in the female somatic gonad (**Fig. 1c**) and at very low levels in the larval testis (**Fig. 1d**). We also saw expression of EcR in terminal epithelium cells of the testis (**Extended Data Fig. 1h, k**). An EcR co-factor, Taiman (Tai), that mediates ecdysone signaling in the adult ovary[20,22] was expressed in the somatic gonad of both larval ovaries and testes **(Fig. 1e-f)**.

To test whether EcR expression and activity are regulated downstream of Dsx, we assessed EcR and Br levels in *dsx*-mutant gonads. We generated female animals heterozygous for the dsx^D allele[29], which carries a mutation in the Tra binding site and cannot be spliced into dsx^F. Thus, XX dsx^D/+ animals express both Dsx^M and Dsx^F and mimic *dsx* loss-of-function[29]. In the gonad, this leads to stochastic establishment of hubs or TFs regardless of chromosomal sex[2]. XX dsx^D/+ gonads exhibited significantly lower levels of EcR protein (**Fig. 2a-d**) and *EcR* mRNA (**Extended Data Fig. 3a-c**) compared with control ovaries. While Br-Z1 was predictably female-specific in control gonads **(Fig. 2e-f)**, in dsx^D/+ gonads Br-Z1 expression in the niche correlated with niche identity rather than chromosomal sex. In other words, dsx^D/+ gonads with a hub expressed very low levels of Br-Z1 **(Fig. 2g)** while dsx^D/+ gonads with TFs expressed higher levels of Br-Z1 **(Fig. 2h)**, supporting the idea that high ecdysone signaling promotes female niche development.

To evaluate whether Dsx is a transcriptional regulator of *EcR*, we generated a transgenic reporter, EcR^{6kb}-*gfp*, that expresses GFP under control of a 6.3-kb intronic enhancer element in the *EcR* locus. This enhancer contains two predicted Dsx binding sites that are highly conserved across *Drosophila* species. EcR^{6kb}-*gfp* was expressed in somatic cells of the larval ovary, while very low levels were observed in the larval testis **(Fig. 2i, j)**. Interestingly, EcR^{6kb}-*gfp* activity was lower in XX intersex gonads with perturbed Dsx function (dsx^D/+) compared with control XX gonads **(Fig. 2k-m; Extended Data Fig. 3d-f)**.

EcR antagonizes testis development

Since EcR activity was female-biased during gonad development, we investigated whether the requirement for EcR in gonad niche development is sexually dimorphic. EcR activation by 20E is required for development of the female gonad stem cell niche[11,30]. To investigate the role of EcR in gonad niche development, we blocked EcR activation in the somatic gonad by over-expressing a dominant-negative allele of *EcR-A*[31] (EcR^{DN}) using the somatic gonad driver *traffic jam (tj)-GAL4* (*tj-GAL4; UAS-EcR^{DN}* –

abbreviated as *tj>EcR^{DN}* throughout). Importantly, since EcR acts as a repressor in the absence of hormone[32], *EcR^{DN}* mis-expression specifically blocks the activator function of EcR. As previously observed, we found that *EcR^{DN}* expression in the female somatic gonad prevented the differentiation of Engrailed-expressing TF cells[11] **(Fig. 3a, b)**. Interestingly, transcriptional activation via EcR was not required for establishment of the male gonad niche, as testes expressing *EcR^{DN}* did not contain fewer hub cells than control testes. In fact, *tj>EcR^{DN}* testes showed a slight increase in hub cell number and hub volume compared with controls **(Extended Data Fig. 4)**.

To test whether activating EcR in the developing testis is deleterious to male niche development, we first identified the rate-limiting components for ecdysone signaling in the larval testis. Neither over-expression of EcR nor supplementing larval food with 20E were sufficient to activate signaling, as determined by quantifying Br expression **(Extended Data Fig. 2, Extended Data Fig. 5a-e)**. We hypothesized that at least one essential EcR co-factor must also be over-expressed to yield EcR activation. Tai is expressed in TF cells and intermingling somatic cells **(Fig. 1e)** and at low levels in hub cells (**Fig. 1f**). Tai protein was also detected in differentiating cyst cells **(Fig. 1f)**. Since Tai was also identified as a putative Dsx target[8], we tested requirement for Tai in EcR-dependent transcriptional activation. Co-expression of *tai* and *EcR*, but not expression of either alone, in the male somatic gonad led to a significant increase in Br expression **(Extended Data Fig. 5a-e)**. Activation of ecdysone signaling in the testis by co-expression of *EcR* and *tai* (*tj>EcR,tai*) led to a significant decrease in larval hub cell number compared with control testes **(Fig. 3c-e)**. Hub cell loss was apparent in as early as L1 larval gonads upon EcR activation **(Fig. 3e and Extended Data Fig. 6a-c)**. Interestingly, hub cell loss was progressive beyond larval development. Loss of all hub cells renders testes unable to support germline and somatic differentiation and results in acellular testes[33,34]. In some *tj>EcR,tai* pupal testes, no cells were detected within the gonad proper **(Extended Data Fig. 6d-e)**, suggesting that these testes have lost all hub cells before adulthood. While EcR activation caused a loss of male niche identity, we did not find acquisition of female niche identity in testes with EcR and Tai over-expression. Expression of the TF marker Engrailed increased in cyst cells upon EcR activation **(Extended Data Fig. 7**, but the testis appeared morphologically normal with no evidence of TF formation.

EcR activity feminizes the adult testis

While EcR activation was not sufficient to sex-transform the gonad niche, we observed a striking male-to-female transformation of somatic support cells in the testis upon co-expression of EcR and Tai. In the adult testis, cyst cells surround the developing spermatogenic cysts while in the ovary, somatic support cells include escort cells that nurture germ cells during the earliest stages of differentiation and epithelial

follicle cells that support maturing oocytes. The transcription factor Chronologically inappropriate morphogenesis (Chinmo) is normally expressed in hub cells and early cyst cells of L3 and adult testes, while in females it is expressed in germ cells but is absent from the somatic gonad at both stages[3,35,36] (**Fig. 4a, b** and **Extended Data Fig. 8a-b**). Co-expression of EcR and Tai caused a loss of Chinmo expression in cyst cells but not hub cells **(Fig. 4c).** Interestingly, this was accompanied by a gain in expression of the follicle cell markers Fas3 and Castor **(Fig. 4d-f)**, which also occurs upon Chinmo depletion from the testis[3,36] (**Fig. 4g).** Repression of *chinmo* and activation of *Br* by ecdysone signaling has also been observed in developing neuroblasts and in larval wing imaginal discs[37,38] and has been proposed to regulate many aspects of development[39]. To further investigate the mutually repressive relationship between EcR and *chinmo* in the somatic gonad, we examined whether transcriptional activation via EcR was required for the male-to-female transformation observed upon loss of *chinmo*. Indeed, interfering with EcR action by expressing *EcR^{DN}* blocked follicle-like transformation observed with loss of *chinmo* (**Fig. 4g-h**). Follicle-like cells were present in 92.3% of *tj>chinmo^{RNAi}* testes, but they appeared in only 7.3% of *tj>EcR^{DN}; >chinmo^{RNAi}* testes (**Fig. 4i**). In addition, depletion of *chinmo* in the testis led to induction of the EcR target Br in the transformed follicle-like cells (**Fig. 4j-l**). These data are consistent with a model where EcR and *chinmo* exhibit mutually repressive interactions in the somatic gonad.

Discussion

Taken together, our data demonstrate that ecdysone signaling can be used to control sex-specific development in an invertebrate model. This does not require sex-specific ecdysone production and, indeed, sufficient ecdysone is produced in both sexes to activate signaling in cells competent to respond to the hormone (**Extended Data Fig. 1a, c**). Here, we demonstrate that at the stage of establishment of the female gonadal stem cell niche, ecdysone signaling is repressed in most somatic cells of the larval testis, including the hub and somatic support cells. Further, our work shows that this repression is necessary for proper testis development: activation of ecdysone signaling causes larval and adult hub defects as well as adult somatic sex transformation of toward a female follicle-like cell identity. Our data suggest that repression of ecdysone signaling in the testis is at least partly due to Dsx-dependent regulation of *EcR* expression. Therefore, invertebrates like *Drosophila* that utilize the steroid hormone ecdysone to regulate development in both sexes can also employ this same hormone to effect sexual differentiation via cell autonomous regulation of its receptor.

We show that somatic gonad development and maintenance feature mutually exclusive expression of the transcription factors Chinmo and Br. Repression of *chinmo* by EcR and Br has been observed in other developmental contexts[37–39], presenting this as a common regulatory cassette. Thus, Chinmo/EcR mutual repression may well be used to control sexual dimorphism in other tissues. It is notable that upon EcR

hyperactivation in the adult testis, cyst cells appear transformed to a female-like identity, whereas hub cells are lost but do not undergo a similar transformation toward a female niche (TF/CC) identity. Such a transformation is observed in *dsx* mutants[2], indicating that there must be additional regulatory mechanisms downstream of Dsx, in addition to EcR, that promote sex-specific niche development and maintenance. Candidates for this include *fruitless,* which encodes a transcription factor involved in hub maintenance during larval stages[40], and *bric a brac 1* and *2 (bab1/2)*, which are important for TF cell specification[41]. Interestingly, Fruitless, Bab1/2, Br, and Chinmo all encode transcription factors with BTB (Bric a brac, Tramtrack, Broad) domains that promote homotypic and heterotypic protein interactions. Further, they are all either predicted or known to be Dsx target genes[2,8,40,41]. Thus, the regulation of these factors by Dsx and EcR, in addition to their potential physical associations, could allow for a particularly rich network of regulatory interactions controlling sexual dimorphism in the gonad and other tissues.

The autonomous regulation of ecdysone signaling we report in *Drosophila* gonads may provide a model for how cell-autonomous sexual identity is regulated in vertebrates. Birds, like mammals, utilize estrogen and testosterone to control development of sexually dimorphic characteristics. Indeed, manipulating the activity of aromatase, which converts testosterone to estrogen, can induce sex-reversal during chicken development[42]. However, in birds with a mixture of cells with male (ZZ) and female (ZW) sex chromosome genotypes, the cells with a male genotype develop as male and the cells with a female genotype develop as female, despite the fact that these animals have a single circulating concentration of estrogen and testosterone[43,44]. Further, while male (ZZ) chickens with one mutant allele of *DMRT1* still develop testes, all other tissues analyzed are phenotypically female[45]. These data clearly indicate that even when estrogen and testosterone are used to control sex-specific development, the responding tissues can cell-autonomously differ in how they respond to these hormones. The simplest explanation for this is that these tissues autonomously regulate their response to these steroid hormones, similarly to how *Drosophila* cells regulate sex-specific response to ecdysone signaling in the gonads.

Taken together, our data present a model for sexual differentiation via steroid hormone signaling whereby Dsx establishes sex-specific transcriptional programs in the gonad via EcR/Chinmo expression. Thus, once sexual identity is specified within the gonad it is self-sustaining, and continuous mutual repression of EcR and Chinmo ensure maintenance of gametogenic potential in the testis and ovary, respectively. In this way, tissues such as the developing gonad can achieve local sex-specific steroid hormone signaling via transcriptional regulation of its receptor. Our work uncovers a novel mode of sex hormone regulation that could extend to other *Drosophila* tissues and may also be an important feature of hormone-dependent sexual differentiation across many species.

Acknowledgments

We thank the Bloomington *Drosophila* Stock Center and the Developmental Systems Hybridoma Bank for making available fly stocks and antibodies. We are also grateful to Natalie Murphy, Jiaxin Lin, Harrison Curnutte, and Dr. Shyama Nandakumar for project and manuscript feedback. This work was funded by NIH R01GM113001 (MVD).

Author contributions

Conceptualization: L.G., E.J. and M.V.D.; Methodology: L.G., E.J., E.B., and M.V.D.; Investigation: L.G., E.J., and E.B.; Formal analysis: L.G.; Validation: L.G.; Resources: M.V.D.; Writing – original draft: L.G.; Writing – review and editing: L.G. and M.V.D.; Supervision: L.G. and M.V.D.

Methods

Fly stocks and husbandry

The following flies were used and are described in FlyBase: $Oregon^R$, dsx^1, $Df(3R)dsx^3$, dsx^D, *msl3-gfp*, *tj-gal4 (P{GawB}NP1624)*, *UAS-lacZ*, *UAS-EcR.C*, *UAS-Tai*, $UAS\text{-}EcR\text{-}A^{W650A}$ (referred to as $UAS\text{-}EcR^{DN}$ in text). *EcR6kb-gfp* enhancer line was generated in this study; see below for cloning details.

Antibodies

The following antibodies were used: rabbit anti-Vasa (1:10,000; gift of Ruth Lehmann, Skirball Institute/NYU School of Medicine, NY, USA), guinea pig anti-Tj (1:1000), mouse anti-Fas3 (1:100; DSHB), rat anti-N-cadherin (1:20; DSHB), mouse anti-EcR (Ag10.2; 1:20; DSHB), mouse anti-Broad-core (25E9.D7; 1:40; DSHB), mouse anti-Broad-Z1 (1:50; DSHB), mouse anti-β-galactosidase (40-1a; 1:40; DSHB). Cross-adsorbed secondary antisera (Invitrogen) were raised in goat and diluted to 4 μg/mL for staining.

Larval hormone feeding experiments

Larval 20-hydroxyecdysone (20E) and chromafenozide (CF) feeding experiments were performed as follows. First, embryos were collected for 4 hours on apple juice agar petri dishes supplemented with regular yeast paste. 21 hours after the start of embryo collection, larvae that hatched prematurely were removed. First instar larvae that hatched over the following 4 hours were then transferred to a fresh apple juice plate supplemented with regular yeast paste. Newly ecdysed third instar larvae were transferred to a fresh apple juice plate supplemented with either control yeast paste (dry yeast mixed with H_2O lacking hormone), yeast paste plus 20E (dry yeast mixed with 1 mM 20E in H_2O), or yeast paste plus 1 mM CF (dry yeast mixed with 1 mM CF in H_2O). Wandering third instar larvae were dissected 48 hours later for gonad staining and imaging.

Immunofluorescence

Testes and ovaries were dissected in PBS and fixed in 5% formaldehyde (diluted in PBS) for 20 minutes at room temperature (RT). Fixed tissue was washed twice for 10 minutes each in PBTx (PBS + 0.1% Triton X-100). Blocking was performed for 30 minutes or overnight in BBTx (PBS + 0.1% Triton X-100 + 1% BSA). Primary antibodies were diluted in BBTx and incubated overnight at 4° C. Primary antibodies were washed off twice for 10 minutes each in BBTx. Secondary antibodies were diluted in BBTx, incubated for 2-3 hours at RT in the dark, and washed off in PBTx. Samples were then incubated in 1 μg/mL DAPI for 10 minutes at RT in the dark, and finally washed twice in PBTx. HCR-FISH was performed according to manufacturer's protocol (Molecular Instruments, Inc.) and *EcR* hybridization probes were generated commercially. Tissue mounting medium was supplemented with DABCO anti-fade reagent (*company*) prior to confocal analysis. Confocal images were captured using a Zeiss LSM 710 confocal microscope using 20x, 40x, or 63x objective lens.

ImageJ quantifications

For quantification of *EcR^{6kb}-gfp* gonad expression shown in Fig. 2x, total GFP intensity from a representative Z-slice was quantified for each gonad, then normalized to total DAPI intensity within the same area.

S-phase (EdU incorporation) assay

EdU incorporation was performed using the Click-iT EdU Cell Proliferation kit (Invitrogen). Gonads were dissected in Schneider's medium for 10-15 minutes, then incubated in Schneider's medium containing 10 μM EdU for 30 minutes at room temperature. Tissues were then fixed, washed, and stained as normal (see "immunofluorescence" methods above). Click-iT reaction was performed following secondary antibody incubation. according to manufacturer's guidelines.

Generating EcR^{6kb}-GFP constructs

Cloning was performed using a modified version of pStinger[46] that replaces P-elements with *attB* site to facilitate ΦC31-mediated, site-directed genome integration. The MCS from *pStinger* (Barolo et al. 2000) was excised and ligated into *pARE-GFP^{nls}* (see Chatterjee & Bohmann 2012 for vector map) to generate *pStinger-attB*. The *hsp70* minimal promoter sequence was removed (since the 6.3kb putative enhancer contains a core promoter) to generate *pStinger-attB*. This vector was used as the starting point to generate *EcR^{6kb}-GFP*. The 6.3-kb putative enhancer region was amplified from a BAC construct containing part of the *EcR* locus (*BACR08A11*) using the primers listed below, then digested with SphI/AgeI and ligated into

pStinger-attB using the NEBuilder HiFi DNA cloning assembly kit (New England Biosciences, Inc.). Positive clones were screened by PCR and sequence integrity was verified using Sanger sequencing. Plasmid was injected into *Drosophila* embryos and stable lines were generated via BestGene, Inc.

Primers:

EcR6kb_sphI_ageI_fwd: tagtgctactgcatagca

EcR6kb+sph1_ageI_rev: ctatgcagccgccatata

Figure Legends – main body

Figure 1. Ecdysone signaling activity is female-biased in the developing gonad.

(a-b) Beta-galactosidase expression (green) from a transgenic *EcRE-lacZ*[25] reporter is higher in the ovarian somatic gonad (a) than in the testicular somatic gonad (b). (c-d) EcR protein expression is higher in the ovary (c) than in the testis (d). (e-f) Expression of Taiman-GFP from a genomic duplication is detected in several somatic cell types of an L3 ovary (e), including the apical cap, terminal filaments, and intermingled cells. In an L3 testis (d), Tai is detected at low levels in the hub and CySCs and at higher levels in differentiating cyst cells. Vasa (red) labels the germline, Traffic jam (Tj, blue) labels somatic cells. Scale bars = 25 μm. Yellow brackets indicate TF cells; yellow dotted outline indicates the hub.

Figure 2. EcR activity is regulated by Dsx in the developing somatic gonad.

(a-c) EcR protein levels are higher in an L3 ovary (a, green) than in an L3 testis (b). EcR expression is diminished in XX *dsx*-mutant gonads (*dsx^D^/+*, c-d). DAPI (blue) labels nuclei. (e-h) Br-Z1, an EcR target, is expressed in TFs of a control ovary (e) and is absent from hub cells in a control testis (f). Br-Z1 expression is high in an L3 XX *dsx^D^/+* TF structure (g) and low in a hub structure (h). N-cad labels the gonad niche (TF or hub cells) (i-j) Expression of a 6.3-kb intronic EcR enhancer (*EcR^6kb^-gfp*) in larval gonads. Gonads shown in (i) and (j) contain two copies of enhancer construct (homozygous). *EcR^6kb^-gfp* is expressed throughout the somatic gonad of an LL3 ovary (h, arrowheads) and is expressed at very low levels in the hub (arrowhead) and CySCs of an L3 testis (j). (k-l) *EcR^6kb^-gfp* expression is greatly reduced in *dsx*-mutant (*EcR^6kb^-gfp/+; dsx^D^/+*) gonads (l) compared with control (*EcR^6kb^-gfp/+; +*) ovaries (k). Heterozygous expression of *EcR^6kb^-gfp* in a control testis is shown in (m). In k-m, DAPI (blue) labels nuclei and Tj (red) labels somatic cells. In **i-m**, arrowheads indicate TF cells and arrows indicate the hub. Scale bars = 25 μm.

Figure 3. EcR signaling is deleterious to male somatic gonad establishment.

(a) TFs are established in a control LL3 ovary (*tj>lacZ*). Engrailed (green) labels TF cells. (b) Blocking EcR activation in the somatic gonad (*tj>EcR^DN^*) prevents the establishment of terminal filaments in an L3 ovary. (c) In a control L3 testis, Fas3 (green) labels intact hub cells. (d) Activation of EcR by co-expressing EcR and Tai (*tj>EcR,tai*) in the male somatic gonad produces smaller hubs with visibly fewer cells. Fas3 (green) labels hub cells. (g) Quantification of hub cells per LL3 (blue dots) or L1 (yellow dots) testis in various genotypes. For statistical significance here and in subsequent figures, asterisks indicate statistical significance by Student's t-test as follows: *$p<0.05$; **$p<0.01$; ***$p<0.001$; ****$p<0.0001$. Scale bars = 25 μm.

Figure 4. EcR activation in the testis causes adult somatic sex transformation.

(a-c) Chinmo expression in adult gonads and with EcR activation in the testis. In an adult ovary (a), Chinmo protein (green) is absent from the female somatic gonad (arrowheads). Chinmo is expressed in early female germ cells as previously observed[3]. (b) In an adult testis, Chinmo is detectable in the hub and CySCs. (c) Chinmo expression is dramatically reduced upon EcR activation in the adult testis. (d-h) Expression of follicle cell markers Fas3 (red) and Castor (green) in adult gonads. Adult ovaries (d) show expression of Fas3 (red) and Castor (green) in early follicle cells, while in a control adult testis (e) Castor and Fas3 are absent from the cyst cell lineage. In adult testes, as seen in previous figures, Fas3 is expressed in hub cells. EcR activation in the testis (f) causes somatic feminization in the adult testis, characterized by the presence of Fas3- and Castor-positive somatic aggregates. Depletion of *chinmo* in the male somatic gonad (g) also causes somatic feminization as previously observed[3]. Blocking EcR activation by over-expressing EcR^{DN} (h) suppresses somatic feminization in the absence of Chinmo. (i) Quantification of feminization rescue in $tj>chinmo^{RNAi}$ testes by blocking EcR activation. An abbreviated description of phenotypic categories: 0=no feminization or germline defects; 1=germline defects but no feminization; 2=mild feminization; 3=moderate feminization; 4=severe feminization; 5=acellular testes. Feminization and acellularity are indicated on the graph in blue. More complete descriptions and visual examples of each phenotypic category can be found in **Extended Data Fig. 9**. (j-l) Expression of Br-C (green) in the adult testis upon *chinmo* depletion. Br-C is expressed in follicle cells during mid-oogenesis in the adult ovary (j). Br-C is expressed at moderate levels in early cyst cells but is absent from differentiating cyst cells in an adult testis (k). In a $tj>chinmo^{RNAi}$ testis (l), Br-C is expressed in later cyst cells distal from the hub (hub and testis apex are out of frame in l and l'). Scale bars = 50 μm.

<u>**Figure Legends – Extended Data**</u>

Extended Data Figure 1. Expression of EcR and Br isoforms in the developing gonad.

(a-b) Expression of the EcR target Br-C in an LL3 ovary (a) and testis (b). Br-C expression in terminal epithelium cells of a larval testis is shown in (c). Vasa (green) labels the germline and Tj (blue) labels somatic cells. (d-e) Expression of Br-Z1 isoform in an LL3 ovary (d) and testis (e). Vasa (magenta) labels the germline. (f-k) Expression of EcR-A (f-h) and EcR-B2 isoforms in the LL3 ovary (f, i), testis (g, j) and testicular terminal epithelium cells (h, k). Vasa (green) labels the germline and Tj labels somatic cells. Scale bars = 25 μm.

Extended Data Figure 2.

(a-f) Expression of *EcRE-lacZ* reporter in L3 ovaries (a-c) or L3 testes (d-f) upon feeding with 10% DMSO (vehicle, a and d), 20-hydroxyecdysone (20E, b and e), or chromafenozide (CF, c and f). (g-h) Depletion of *shd* in the larval ovary does not visibly impair ovary development. Note that *dcg-gal4* used in this experiment is expressed in both the ovary and surrounding fat body. Vasa (green) labels the germline, N-cad (red) labels TF cells, and Tj (blue) labels intermingled somatic cells. Scale bars = 25 μm.

Extended Data Figure 3.

(a-c) Detection of *EcR* mRNA (red) in LL3 gonads by HCR-FISH. A *dsx*-mutant ($dsx^D/+$) gonad (c) shows lower *EcR* mRNA abundance than control ovaries (a). *EcR* mRNA in an LL3 testis is shown in (b). (d-f) Detection of *GFP* mRNA (red) in LL3 gonads by HCR-FISH. $dsx^D/+$ gonad (f) contains higher *GFP* mRNA abundance than control ovaries (d). Scale bars = 25 μm.

Extended Data Figure 4.

(a-b) Neither hub specification nor hub maintenance are compromised by blocking EcR activity (*tj>EcR-^{DN}*). Fas3 (green) labels hub cells, Vasa (red) labels the germline, and Tj (blue) labels somatic cells. (c-e) Quantification of hub cell number (c), hub volume (d), and germline stem cell (GSC) number (e) upon blocking EcR activation in the somatic gonad. Scale bars = 10μm.

Extended Data Figure 5.

(a-d) Br-C expression in LL3 control testes upon somatic over-expression of EcR (b), Tai (c), or EcR and Tai simultaneously (d). (e) Quantification of relative Br-C intensity from a-d. Scale bars = 25 μm.

Extended Data Figure 6.

(a-c) Representative images of early L1 (24-28h after egg lay, AEL) gonads, identified by the presence of a Fas3-positive hub (green). Ovaries lack a hub (a) while testes contain a Fas3-positive hub (green) by this stage (b). III staining. Co-expression of *EcR* and *tai* in the early somatic gonad (c) leads to a visibly smaller hub that contains significantly fewer hub cells than control L1 testes (see **Fig. 3e** for quantification). Vasa (red) labels the germline, Tj (blue) labels somatic gonadal precursor cells, which give rise to the somatic gonad in both sexes. (d-e) Pupal testes upon EcR/Tai co-expression. In a control pupal testis (d), the germline and somatic stem cells are maintained by the presence of a hub (Fas3, green) and thus differentiating germ cells (Vasa, red) are observed in the testis. In some pupal testes, following EcR/Tai co-expression in the somatic gonad (e), no Fas3-positive hub cells are present and the absence of Vasa and DAPI staining indicates that both germline and somatic cell lineages have been lost. DAPI-positive muscle sheath cells can be found at the periphery of the testis. Scale bars = 10 μm.

Extended Data Figure 7.

(a-c) Expression of the TF marker Engrailed (En) in LL3 gonads. En (green) labels TF cells in a control ovary (a) but is not detected in the niche of a control testis (b). En is not detected in hub cell nuclei upon EcR/Tai co-expression in the somatic gonad (c). En induction is observed in the cyst cell lineage upon EcR activation (c). In b and c, the hub is outlined with a yellow dotted line. Scale bars = 25 μm.

Extended Data Figure 8.

Mutually exclusive expression of Chinmo (red) and Br-Z1 (green) in somatic cells of LL3 gonads. In an LL3 ovary (a), Chinmo is absent from somatic cells (a', inset arrowheads) while Br-Z1 is expressed broadly in the female somatic gonad. We note that Chinmo is expressed in female larval progenitor germ cells (PGCs) (a'). In an LL3 testis (b), Chinmo is expressed in the hub, CySCs/cyst cells, and GSCs/early germ cells. Br-Z1 is not detectable in the hub or early cyst cells. Vasa (blue) labels the germline. Scale bars = 25 μm.

Extended Data Figure 9.

Phenotypic category descriptions and representative images for rescue quantification in Figure 4i. (a) Category 0: no evidence of feminization (accumulation of Fas3/Castor-expressing somatic aggregates), early/late spermatogonia and large-nuclei spermatocytes present. (b) Category 1: No evidence of feminization, but germline defects seen (shortening of mitotic region and/or absence of large-nuclei spermatocytes). (c) Category 2: mild feminization (a few cells express Fas3 and/or Castor, but aggregates have not yet formed). (d) Category 3: moderate feminization (follicle-like aggregates present clearly expressing Fas3 and Castor, few germ cells are present and mostly seem arrested in spermatogonial stage). (e) Category 4:

severe feminization (large follicle-like aggregates present, few/no germ cells remain). (f) Category 5: acellular. Vasa (blue) labels the germline. Scale bars = 50 μm.

References

1.   Le Bras, S. & Van Doren, M. Development of the male germline stem cell niche in Drosophila. *Dev. Biol.* **294**, 92–103 (2006).

2.   Camara, N., Whitworth, C., Dove, A. & Van Doren, M. Doublesex controls specification and maintenance of the gonad stem cell niches in Drosophila. *Development* **146**, (2019).

3.   Grmai, L., Hudry, B., Miguel-Aliaga, I. & Bach, E. A. Chinmo prevents transformer alternative splicing to maintain male sex identity. *PLoS Genet.* **14**, e1007203 (2018).

4.   Raymond, C. S., Murphy, M. W., O'Sullivan, M. G., Bardwell, V. J. & Zarkower, D. Dmrt1, a gene related to worm and fly sexual regulators, is required for mammalian testis differentiation. *Genes Dev.* **14**, 2587–2595 (2000).

5.   Matson, C. K. *et al.* DMRT1 prevents female reprogramming in the postnatal mammalian testis. *Nature* **476**, 101–104 (2011).

6.   Minkina, A. *et al.* DMRT1 protects male gonadal cells from retinoid-dependent sexual transdifferentiation. *Dev. Cell* **29**, 511–520 (2014).

7.   Lindeman, R. E. *et al.* Sexual cell-fate reprogramming in the ovary by DMRT1. *Curr. Biol.* **25**, 764–771 (2015).

8.   Clough, E. *et al.* Sex- and tissue-specific functions of Drosophila doublesex transcription factor target genes. *Dev. Cell* **31**, 761–773 (2014).

9.   Zarkower, D. DMRT genes in vertebrate gametogenesis. *Curr. Top. Dev. Biol.* **102**, 327–356 (2013).

10.   Wingert, L. & DiNardo, S. Traffic jam functions in a branched pathway from Notch activation to niche cell fate. *Development* **142**, 2268–2277 (2015).

11.   Gancz, D., Lengil, T. & Gilboa, L. Coordinated regulation of niche and stem cell precursors by hormonal signaling. *PLoS Biol.* **9**, e1001202 (2011).

12.   Reinisch, J. M., Ziemba-Davis, M. & Sanders, S. A. Hormonal contributions to sexually dimorphic behavioral development in humans. *Psychoneuroendocrinology* **16**, 213–278 (1991).

13.   Angelopoulou, R., Lavranos, G. & Manolakou, P. Establishing sexual dimorphism in humans. *Coll. Antropol.* **30**, 653–658 (2006).

14.   Wilson, C. A. & Davies, D. C. The control of sexual differentiation of the reproductive system and brain. *Reproduction* **133**, 331–359 (2007).

15.   Gomez-Lobo, V., Oelschlager, A.-M. A. & North American Society for Pediatric and Adolescent Gynecology. Gynecological challenges in the diagnosis and care of patients with DSD: The role of the obstetrician gynecologist in the multidisciplinary approach to the patient. *Am. J. Med. Genet. C Semin. Med. Genet.* **175**, 300–303 (2017).

16.   Li, T. & Bender, M. A conditional rescue system reveals essential functions for the ecdysone receptor (EcR) gene during molting and metamorphosis in Drosophila. *Development* **127**, 2897–2905 (2000).

17.   Ables, E. T. & Drummond-Barbosa, D. The steroid hormone ecdysone functions with intrinsic chromatin remodeling factors to control female germline stem cells in Drosophila. *Cell Stem Cell* **7**, 581–592 (2010).

18.   Knapp, E. & Sun, J. Steroid signaling in mature follicles is important for Drosophila ovulation. *Proc Natl Acad Sci USA* **114**, 699–704 (2017).

19.   Zipper, L., Jassmann, D., Burgmer, S., Görlich, B. & Reiff, T. Ecdysone steroid hormone remote controls intestinal stem cell fate decisions via the PPARγ-homolog Eip75B in Drosophila. *eLife* **9**, (2020).

20.   Bai, J., Uehara, Y. & Montell, D. J. Regulation of invasive cell behavior by taiman, a Drosophila protein related to AIB1, a steroid receptor coactivator amplified in breast cancer. *Cell* **103**, 1047–1058 (2000).

21.   Okamoto, N. *et al.* A membrane transporter is required for steroid hormone uptake in drosophila. *Dev. Cell* **47**, 294-305.e7 (2018).

22.   Manning, L. *et al.* A hormonal cue promotes timely follicle cell migration by modulating transcription profiles. *Mech. Dev.* **148**, 56–68 (2017).

23.   Li, Y., Ma, Q., Cherry, C. M. & Matunis, E. L. Steroid signaling promotes stem cell maintenance in the Drosophila testis. *Dev. Biol.* **394**, 129–141 (2014).

24.   Koelle, M. R. *et al.* The Drosophila EcR gene encodes an ecdysone receptor, a new member of the steroid receptor superfamily. *Cell* **67**, 59–77 (1991).

25.   White, K. P., Hurban, P., Watanabe, T. & Hogness, D. S. Coordination of Drosophila metamorphosis by two ecdysone-induced nuclear receptors. *Science* **276**, 114–117 (1997).

26.   Riddihough, G. & Pelham, H. R. An ecdysone response element in the Drosophila hsp27 promoter. *EMBO J.* **6**, 3729–3734 (1987).

27.   Petryk, A. *et al.* Shade is the Drosophila P450 enzyme that mediates the hydroxylation of ecdysone to the steroid insect molting hormone 20-hydroxyecdysone. *Proc Natl Acad Sci USA* **100**, 13773–13778 (2003).

28.   Yi, W. & Zarkower, D. Similarity of DNA binding and transcriptional regulation by Caenorhabditis elegans MAB-3 and Drosophila melanogaster DSX suggests conservation of sex determining mechanisms. *Development* **126**, 873–881 (1999).

29.   Nöthiger, R. *et al.* Genetic and developmental analysis of the sex-determining gene 'double sex' (*dsx*) of *Drosophila melanogaster. Genet. Res.* **50**, 113–123 (1987).

30.   Hodin, J. & Riddiford, L. M. The ecdysone receptor and ultraspiracle regulate the timing and progression of ovarian morphogenesis during Drosophila metamorphosis. *Dev. Genes Evol.* **208**, 304–317 (1998).

31.   Cherbas, L., Hu, X., Zhimulev, I., Belyaeva, E. & Cherbas, P. EcR isoforms in Drosophila: testing tissue-specific requirements by targeted blockade and rescue. *Development* **130**, 271–284 (2003).

32.   Dobens, L., Rudolph, K. & Berger, E. M. Ecdysterone regulatory elements function as both transcriptional activators and repressors. *Mol. Cell. Biol.* **11**, 1846–1853 (1991).

33.   Voog, J. *et al.* Escargot restricts niche cell to stem cell conversion in the Drosophila testis. *Cell Rep.* **7**, 722–734 (2014).

34.   Herrera, S. C. *et al.* Proliferative stem cells maintain quiescence of their niche by secreting the Activin inhibitor Follistatin. *Dev. Cell* **56**, 2284-2294.e6 (2021).

35.   Flaherty, M. S. *et al.* chinmo is a functional effector of the JAK/STAT pathway that regulates eye development, tumor formation, and stem cell self-renewal in Drosophila. *Dev. Cell* **18**, 556–568 (2010).

36.   Ma, Q., Wawersik, M. & Matunis, E. L. The Jak-STAT target Chinmo prevents sex transformation of adult stem cells in the Drosophila testis niche. *Dev. Cell* **31**, 474–486 (2014).

37.   Syed, M. H., Mark, B. & Doe, C. Q. Steroid hormone induction of temporal gene expression in Drosophila brain neuroblasts generates neuronal and glial diversity. *eLife* **6**, (2017).

38.   Narbonne-Reveau, K. & Maurange, C. Developmental regulation of regenerative potential in Drosophila by ecdysone through a bistable loop of ZBTB transcription factors. *PLoS Biol.* **17**, e3000149 (2019).

39.   Truman, J. W. & Riddiford, L. M. Chinmo is the larval member of the molecular trinity that directs Drosophila metamorphosis. *Proc Natl Acad Sci USA* **119**, e2201071119 (2022).

40.   Zhou, H., Whitworth, C., Pozmanter, C., Neville, M. C. & Van Doren, M. Doublesex regulates fruitless expression to promote sexual dimorphism of the gonad stem cell niche. *PLoS Genet.* **17**, e1009468 (2021).

41.   Williams, T. M. *et al.* The regulation and evolution of a genetic switch controlling sexually dimorphic traits in Drosophila. *Cell* **134**, 610–623 (2008).

42.   Estermann, M. A., Major, A. T. & Smith, C. A. Genetic regulation of avian testis development. *Genes (Basel)* **12**, (2021).

43.   Agate, R. J. *et al.* Neural, not gonadal, origin of brain sex differences in a gynandromorphic finch. *Proc Natl Acad Sci USA* **100**, 4873–4878 (2003).

44. Zhao, D. *et al.* Somatic sex identity is cell autonomous in the chicken. *Nature* **464**, 237–242 (2010).

45. Ioannidis, J. *et al.* Primary sex determination in birds depends on DMRT1 dosage, but gonadal sex does not determine adult secondary sex characteristics. *Proc Natl Acad Sci USA* **118**, (2021).

46. Barolo, S., Carver, L. A. & Posakony, J. W. GFP and beta-galactosidase transformation vectors for promoter/enhancer analysis in Drosophila. *BioTechniques* **29**, 726, 728, 730, 732 (2000). Sciwheel inserting bibliography...

Figure 1

Figure 2

Figure 3

Figure 4

Extended Data Figure 1

Extended Data Figure 2

EcRE-lacZ ovaries (XX)

β-gal   Tj   Vasa   β-gal

10% DMSO

a   a'

20E feeding

b   b'

CF feeding

c   c'

EcRE-lacZ testes (XY)

β-gal   Tj   Vasa   β-gal

d   d'

e   e'

f   f'

dcg>mCherry^{RNAi}   *dcg>shd^{RNAi}*

β-gal
Tj
N-cad
Vasa

g   h

N-cad

g'   h'

Extended Data Figure 3

Extended Data Figure 4

Extended Data Figure 5

Extended Data Figure 6

Extended Data Figure 7

Extended Data Figure 8

Extended Data Figure 9