RESEARCH ARTICLES

Phosphorylated transcription factor PuHB40 is involved in

ROS-dependent anthocyanin biosynthesis in pear exposed to high-light

stress

Lu Zhang[1], Lu Wang[1], Yuhao Gao[1], Shulin Yang[1], Jun Su[2], Junbei Ni[1], Yuanwen

Teng[1,3] and Songling Bai[1,*]

[1]College of Agriculture and Biotechnology, Zhejiang University, Hangzhou 310058,

Zhejiang, China

[2]Institute of Horticulture, Yunnan Academy of Agricultural Sciences, Kunming 650205,

China

[3]Hainan Institute of Zhejiang University, Sanya, Hainan 572000, China.

Short title: PuHB40 promotes ROS-dependent anthocyanin biosynthesis

*Corresponding author: Songling Bai (songlingbai@zju.edu.cn)

24   (songlingbai@zju.edu.cn).

ABSTRACT

In the context of global warming and climate change, as sessile organisms, plants are more vulnerable to environmental stresses than before. Stress-induced reactive oxygen species (ROS) accumulation results in plant cell damages and even cell death. Anthocyanins are important antioxidants that scavenge ROS to maintain ROS homeostasis. However, the mechanism underlying ROS-induced anthocyanin accumulation is unclear. In this study, we determined that the HD-Zip I family member PuHB40 mediates ROS-dependent anthocyanin biosynthesis under high-light stress. Specifically, PuHB40 is a transcription factor that induces PuMYB123-like/PubHLH3 complex for anthocyanin biosynthesis. The transcriptional activation by PuHB40 depends on its phosphorylation level, which is regulated by protein phosphatase 2A (PP2A). High ROS contents maintain the phosphorylation of PuHB40 at a high level, while also enhancing PuHB40-induced *PuMYB123-like* transcription by decreasing *PuPP2AA2* expression at the transcriptional and translational levels, ultimately leading to increased anthocyanin biosynthesis. Our study revealed the pathway regulating ROS-induced anthocyanin biosynthesis in pear, further clarifying the mechanism underlying abiotic stress-induced anthocyanin biosynthesis, which may have implications for improving plant stress tolerance.

Keywords: high-light stress, ROS, anthocyanin biosynthesis, PuHB40, PuMYB123-like,

PP2A phosphatase, dephosphorylation

IN A NUTSHELL

Background: Various abiotic stresses, including high light, promotes the accumulation of

anthocyanins in plants. Anthocyanins attenuate the negative effects of on photosynthesis

and physiological metabolism of plants under high-light stress by absorbing excess light

energy or by acting as an antioxidant to reduce ROS accumulation. Several reports have

showed that ROS can stimulate anthocyanin accumulation, but whether the high-light

induced anthocyanin accumulation depends on ROS is still under debate. Nevertheless,

the underlying mechanism of ROS-dependent anthocyanin biosynthesis is still not clear.

Question: What is the underlying molecular mechanism of high light stress-induced

anthocyanin biosynthesis?

Findings: We found that various stresses-induced anthocyanin accumulation in pear plant

is dependent on ROS. PuHB40 was the core TF involved in ROS-dependent anthocyanin

biosynthesis under high-light stress, while PuMYB13-like is the key MYB TF for

anthocyanin biosynthesis in ROS pathway. PuHB40 transcriptionally activates

PuMYB123-like/PubHLH3 complex for promoting anthocyanin biosynthesis. The

transcriptional activation by PuHB40 depends on its phosphorylation level, which is

regulated by protein phosphatase 2A (PP2A). High ROS contents maintain the

phosphorylation of PuHB40 at a high level, while also enhancing PuHB40-induced

PuMYB123-like transcription by decreasing *PuPP2AA2* expression at the transcriptional

and translational levels, ultimately leading to increased anthocyanin biosynthesis.

Next steps: An important task for next step is to explore whether PuHB40 regulation

pathway is applicable in other stresses and provide insight into theoretical basis of abiotic

stress-induced anthocyanin biosynthesis. In addition, investigate other post-translational

modifications involved in ROS-dependent anthocyanin biosynthesis. Moreover, we will

explore the effects of different stress levels on anthocyanin accumulation in pear fruits,

and try to regulate pear fruit coloration by appropriate stress.

INTRODUCTION

Because of global warming and climate change, plants are increasingly becoming more

susceptible to a variety of environmental stresses (e.g., drought, salinity, cold, and

high-light) during growth and development, which leads to changes in physiological and

biochemical metabolism, including the dramatic accumulation of reactive oxygen species

(ROS) (Hipsch et al. 2021; Mittler et al. 2022). Light is one of the most indispensable

environmental factors for plants. It provides energy for photosynthesis to transfer carbon

dioxide (CO_2) and water into energy-rich organic matter to sustain plant physiological

metabolism (Croce and van Amerongen 2014). Under high-light stress, the flux of

photons exceeds the energy that plant need to fix CO_2, O_2^- and $_1O^2$ would primarily

produce in the chloroplasts by photosystems i and ii, respectively, and H_2O_2 will also be

generated in peroxisomes if photorespiration is activated (Galvez-Valdivieso et al. 2009;

Kerchev et al. 2016; Wang et al. 2020; Mittler et al. 2022). Excessive ROS accumulation

would disrupt the biological machinery surrounding them, including membranes, proteins

and DNA, and then trigger a physiological or programmed pathway for cell death (Lee

and Gould 2002; Mittler 2017). Anthocyanins are important secondary metabolites that

can function as non-enzymatic antioxidants that maintain ROS homeostasis and limit the

damage due to stress-induced ROS. Specifically, anthocyanins decrease ROS production

by absorbing light energy at wavelengths that are not readily absorbed by pigments (e.g.,

green and blue wavelengths) and limiting the photoinhibitory effects of abiotic stresses

(Feild et al. 2001; Hoch et al. 2001; Steyn et al. 2002). They are also antioxidants that

serve as ROS scavengers. Notably, their antioxidant capacity is greater than that of other

flavonoids or even ascorbic acid and vitamin E (Gould et al. 2002; Bi et al. 2014; Acero

et al. 2019). Considering various abiotic stresses promote anthocyanin accumulation in

plants (Catala et al. 2011; Xie et al. 2012; Jaakola 2013; Nakabayashi et al. 2014) and

ROS can induce anthocyanin production (Xu et al. 2017; Shi et al. 2018), ROS may

influence abiotic stress-induced anthocyanin biosynthesis; this hypothesis has been tested

in a few studies (Qu et al. 2018).

Anthocyanins are synthesized in the phenylpropanoid pathway by various enzymes

encoded by structural genes. Examples include phenylalanine ammonia-lyase (PAL),

chalcone synthase (CHS), chalcone isomerase (CHI), flavanone 3-hydroxylase (F3H),

flavonoid 3′-hydroxylase (F3′H), dihydroflavonol 4-reductase (DFR), anthocyanin

synthase (ANS), and UDP-glucose: flavonoid 3-glucosyltransferase (UFGT)

(Winkel-Shirley 2001; Koes et al. 2005; Xu et al. 2015). The R2R3-MYB and basic

helix–loop–helix (bHLH) transcription factors (TFs) combine with WD-repeat proteins to

form MBW complexes in which MYBs function as major regulators of structural gene

expression (Koes et al. 2005; Jaakola 2013; Xu et al. 2015). The R2R3-MYB TFs are

subdivided into 25 subgroups, among which subgroup 6 contains TFs that are mainly

responsible for promoting anthocyanin biosynthesis, including PAP1 (AtMYB75), PAP2

(AtMYB90), and AtMYB113 (Dubos et al. 2010). The expression of MYB TFs is

regulated by various factors, including light (Takos et al. 2006; Li et al. 2012; Bai et al.

2019a), low/high temperatures (Choi et al. 2009; Lin-Wang et al. 2011; Xie et al. 2012),

and phytohormones (Shan et al. 2009; Das et al. 2012; Ni et al. 2023). The encoded MYB

TFs control the expression of anthocyanin biosynthetic structural genes. To date, the

molecular basis of ROS-induced anthocyanin biosynthesis remains relatively

uncharacterized.

ROS accumulation in cells has an effect on redox state and the post-translational

modification of proteins, especially on protein phosphorylation. ROS signal is transduced

by Ca^{2+}, Ca^{2+} binding protein, phosphatidic acid (PA) and phospholipase D (PLD) and

activates OXIDATIVE SIGNAL-INDUCIBLE 1 (OXI1), which subsequently activates

MAPK cascade pathway, which in turn promotes the phosphorylation of downstream

transcription factors, such as WRKY proteins and AP2/ethylene-response factors (ERFs)

(Kovtun et al. 2000; Jalmi and Sinha 2015; Byrne et al. 2020; Mittler et al. 2022).

Furthermore, the phosphorylation of transcription factors are also regulated by

phosphatase, such as PTP1, SAL1 and PP2C, whose activity are regulated by ROS levels

(Gupta and Luan 2003), oxidative stresses (Chan et al. 2016) or phytohormones (Mine et

al. 2017). The PP2A holoenzyme, which is a serine/threonine-specific phosphatase, is a

heterotrimeric complex comprising a catalytic subunit C, a regulatory subunit B, and a

structural subunit A. The A subunit has a conserved structure consisting of 15 tandem

repeats of a 39 amino acid sequence (i.e., HEAT motif) (Janssens and Goris 2001). In *A.*

thaliana, three A subunits of PP2A have been identified, namely PP2AA1 (RCN1),

PP2AA2, and PP2AA3. The phosphatase activity of PP2A plays an important role in

stress response (DeLong 2006; Uhrig et al. 2013; Lillo et al. 2014; Durian et al. 2016).

However, it is not clear whether PP2A phosphatases are involved in ROS regulation of

protein phosphorylation under stress conditions.

In this study, we determined that abiotic stress-induced anthocyanin biosynthesis is

at least partially dependent on ROS in pear. The HD-Zip I family member PuHB40 was

identified as the key TF involved in ROS-dependent anthocyanin biosynthesis. Notably,

PuHB40 functions as the upstream positive regulator of the PuMYB123-like/PubHLH3

complex during anthocyanin biosynthesis. The dephosphorylation of PuHB40, which is

catalyzed by protein phosphatase 2A (PP2A), inhibits its function related to ROS-induced

anthocyanin biosynthesis. Under abiotic stress conditions, ROS inhibit the transcription

and translation of *PuPP2AA2*. This study revealed the pathway regulating ROS-induced

anthocyanin biosynthesis in pear under high-light stress, providing important insights into

the mechanism mediating abiotic stress-induced anthocyanin production.

RESULTS

Stress induced ROS promoted anthocyanin biosynthesis in pear seedling

To test whether abiotic stress-induced anthocyanin biosynthesis depends on ROS

production, pear seedlings (*P. ussuriensis*) grown under continuous light were exposed to

abiotic stresses, including salt, drought, and high-light, and the ROS scavenger

N,N′-dimethylthiourea (DMTU) was used to decrease the ROS content under abiotic

stress conditions. The various abiotic stresses caused anthocyanins to accumulate in pear

seedlings after 7 days, but anthocyanin production decreased significantly following the

DMTU treatment (Supplemental Fig. S1A), indicating that ROS are potentially involved

in stress-induced anthocyanin production. Furthermore, ROS accumulation and

anthocyanin biosynthesis were examined at 0, 6, 12, 24, 48, 72, and 96 h after plants

underwent the high-light stress treatment (Fig. 1A). Anthocyanin accumulation peaked at

the 48-h time-point and then decreased, but this accumulation was significantly

suppressed in the seedlings that were also treated with DMTU (Fig. 1B). The NBT

staining results indicated that the superoxide radical (O^{2-}) accumulated in leaves only

after the 6-h time-point, but the accumulation was significantly inhibited by DMTU (Fig.

1C). The DAB staining and quantitative analysis revealed that the accumulation of

hydrogen peroxide (H_2O_2) peaked at 48 h and then decreased, but the DMTU treatment

suppressed this accumulation (Fig. 1, D and E). Moreover, in response to the different

stress treatments (e.g. high-light stress, drought stress, and salt stress), the anthocyanin

and H_2O_2 contents increased in the same leaf regions (Supplemental Fig. S1B).

According to these results, abiotic stress-induced anthocyanin accumulation depends on

ROS production. Hence, the ROS scavenger DMTU can inhibit ROS accumulation and

indirectly decrease anthocyanin production.

To identify the key genes responsible for abiotic stress-induced anthocyanin

biosynthesis, the expression of anthocyanin-related structural and regulatory genes was

analyzed. The *PuCHI*, *PuCHS*, *PuDFR*, *PuF3H*, and *PuANS* expression levels increased

significantly and peaked at 48 h after the high-light stress treatment, which was in

accordance with the changes in anthocyanin accumulation. The expression of these genes

was significantly inhibited by DMTU (Fig. 1, F-J). In addition, high-light responsive

genes *PuELIP1* and *PuELIP2* were significantly up-regulated in high-light stress

treatment and high-light stress plus DMTU treatment compared with the control group,

indicating that both groups suffered from high-light stress (Supplemental Fig. S1, C and

D). Interestingly, *PuMYB10*, which encodes a critical TF that regulates light-induced

anthocyanin biosynthesis in pears, was similarly expressed following the high-light stress

and DMTU treatments (Fig. 1K), even though these two treatments resulted in significant

differences in anthocyanin accumulation and the expression of downstream structural

genes. These findings suggest that other factors influencing transcription may be involved

in regulating ROS-dependent anthocyanin biosynthesis.

Identification of PuMYB123-like and its interaction with PubHLH3

To determine which TFs participate in ROS-dependent anthocyanin biosynthesis, the

high-light stress-treated pear seedlings underwent an RNA-seq analysis. Since MYBs

TFs play an important role in anthocyanin biosynthesis, we first screened for TFs from

the MYB family with expression patterns consistent with anthocyanin biosynthesis

structural genes. Finally, we identified a MYB transcription factor, *PuMYB123-like*,

whose expression increased rapidly in response to high-light stress and peaked at 48 h,

but the DMTU treatment had inhibitory effects on *PuMYB123-like* expression (Fig. 2A).

A phylogenetic analysis indicated that *PuMYB123-like* is a homolog of AT5G35550

(AtMYB123) in subgroup 5 (Supplemental Fig. S2A). Moreover, *PuMYB123-like* is most

closely related to *MdMYB6* from apple (*Malus domestica*) and *FaMYB9* from strawberry

(*Fragaria × ananassa*), which are reportedly involved in anthocyanin and

proanthocyanidin biosynthesis, respectively (Schaart et al. 2012; Xu et al. 2020) (Fig. 2B).

The multiple sequence alignment showed PuMYB123-like contains a typical R2R3

repeat in the N-terminal, with a bHLH motif in the R3 repeat region that can interact with

bHLH TFs (Supplemental Fig. S2B). The subcellular localization analysis showed that

the fluorescence of GFP-tagged PuMYB123-like was predominantly distributed in the

nucleus, consistent with the characteristics of TFs (Fig. 2C). The results of the

trans-activation assay suggested PuMYB123-like is a transcriptional activator (Fig. 2D).

Considered together, these results indicate that PuMYB123-like is a transcriptional

activator that may contribute to ROS-dependent anthocyanin biosynthesis.

Considering MYB TFs, bHLHs, and WD-repeat proteins combine to form MBW

complexes that control the expression of anthocyanin-related structural genes, we

assessed whether PuMYB123-like can interact with three candidate bHLH TFs

(PubHLH3, PubHLH33, and PubHLH64). According to the yeast two-hybrid (Y2H)

assay results, PuMYB123-like can bind to PubHLH3 (Fig. 2E). The interaction was

confirmed in pull-down and luciferase complementation imaging (LCI) assays, which

revealed PuMYB123-like can interact with PubHLH3 both *in vivo* and *in vitro* (Fig. 2, F

and G). Because of the self-activation of PubHLH33 and PubHLH64, the corresponding

genes were fragmented to assess potential interactions. The results showed that

PuMYB123-like can interact with the truncated PubHLH64 (encoded by 747–1,086 bp)

(Supplemental Fig. S3A), and weakly interact with the truncated PubHLH33 (encoded by

588–1,326 bp) (Supplemental Fig. S3B). The dual-luciferase assay demonstrated that

PuMYB123-like can significantly activate the promoters of the anthocyanin-related

structural genes *PuCHS*, *PuANS*, *PuDFR*, and *PuUFGT* (Fig. 2H). Moreover, PubHLH3

significantly enhanced the activation of the *PuCHS* and *PuANS* promoters by

PuMYB123-like, but PubHLH33 and PubHLH64 had no significant effect (Fig. 2I and

Supplemental Fig. S3C). On the basis of these observations, we speculated that

PuMYB123-like, which belongs to subgroup 2 of the R2R3-MYB family, promotes

anthocyanin-related structural genes expression as a component of the MBW complex

that includes PubHLH3.

PuMYB123-like promotes anthocyanin biosynthesis in pears

To confirm PuMYB123-like affects anthocyanin biosynthesis, *PuMYB123-like* was

overexpressed in pear calli. After a 3-day exposure to light, the anthocyanin contents and

anthocyanin-related structural gene expression levels were significantly higher in the calli

overexpressing *PuMYB123-like* than in the wild-type calli. In addition, the silencing of

PuMYB123-like in pear calli via RNA interference (RNAi) inhibited anthocyanin

biosynthesis and decreased the expression of anthocyanin-related structural genes (Fig.

3A-C). Considering PuMYB123-like belongs to subgroup 2 of the R2R3-MYB family,

which reportedly regulates proanthocyanin biosynthesis, the proanthocyanin content in

PuMYB123-like-overexpressing (*PuMYB123-like*-OE) pear calli was also examined.

There was no significant difference in the proanthocyanin contents of the wild-type and

transgenic pear calli (Supplemental Fig. S4A). Additionally, *PuMYB123-like* was

ectopically expressed in *Arabidopsis thaliana* to verify the function of the encoded TF.

Compared with the wild-type control, the two independent transgenic lines accumulated

more anthocyanin (Fig. 3D-F) and had shorter hypocotyls (Fig. 3G). The expression of

the anthocyanin-related structural genes *AtCHS*, *AtCHI*, *AtDFR*, *AtF3H*, *AtLODX*, and

AtPAP1 increased significantly in the *PuMYB123-like*-expressing lines (Fig. 3H). To

examine whether PuMYB123-like mediates anthocyanin biosynthesis in fruits,

PuMYB123-like was transiently overexpressed in the peel of 'Zaosu' pear fruits that were

bagged in advance. The comparison with the control peels (empty vector) after a 3-day

exposure to light indicated the *PuMYB123-like*-OE peels had significantly higher

anthocyanin contents (Supplemental Fig. S4, B and C), and anthocyanin-related structural

gene expression levels (i.e., *PpCHS, PpDFR, PpF3H, PpANS, PpUFGT, PpMYB10*, and

PpMYB114) (Supplemental Fig. S4D). Similarly, 'Hongzaosu' red pear fruit peels in

which *PuMYB123-like* was transiently silenced via virus-induced gene silencing (VIGS)

had lower anthocyanin contents and related structural gene expression levels than the

control peels (TRV) (Supplemental Fig. S4, E-G). Taken together, these results

demonstrate that PuMYB123-like positively regulate the ROS-dependent anthocyanin

biosynthesis in pears.

Identification of PuHB40 and its response to ROS

To further investigate the upstream regulator of *PuMYB123-like* in pear, the cis-elements

in its promoter were analyzed. Four CAATNATTG motifs (i.e., the binding site of

HD-Zip proteins) were identified (Fig. 4A). After screening for HD-Zip-encoding genes

with expression patterns similar to that of *PuMYB123-like*, *PuHB40* was identified as a

candidate gene. In response to the high-light stress treatment, the *PuHB40* expression

level gradually increased after 24 h, peaked at 48 h, and then decreased, but the DMTU

treatment significantly inhibited *PuHB40* expression (Fig. 4B). Additionally, the PuHB40

content changes in response to ROS were analyzed in an immunoblot involving an

affinity-purified anti-PuHB40 polyclonal antibody generated in this study (Supplemental

Fig. S5A). The immunoblot results showed that the PuHB40 protein content increased

significantly after 6 h of the high-light stress treatment, after which it gradually decreased,

but was maintained at a relatively high level. In contrast, the PuHB40 protein level

remained low after the DMTU treatment (Fig. 4C). Hence, PuHB40 was affected by ROS

at both the transcriptional and translational levels after the abiotic stress treatment.

Interestingly, the response of PuHB40 was much faster at the translational level than at

the transcriptional level in pear seedlings. To explore whether PuHB40 is responsive to

ROS directly, we treated pear calli with 4 mM H_2O_2, which resulted in significantly

increased anthocyanin biosynthesis (Fig. 4, D and E). Moreover, the expression of

PcHB40 as well as the anthocyanin-structural genes *PcCHI*, *PcDFR*, *PcF3H*, and *PcANS*

also increased (Fig. 4F). Following the H_2O_2 treatment, the *PcHB40* expression level

increased after 3 h (Fig. 4G), suggestive of a direct response to H_2O_2.

A phylogenetic analysis showed that PuHB40 is a homolog of AT4G36740.1

(ATHB40), while a multiple sequence alignment revealed a conserved homeodomain and

leucine zipper in PuHB40 (Supplemental Fig. S6, A and B). The trans-activation assay

confirmed that PuHB40-BD is a transcriptional activator (Fig. 4H). The subcellular

localization analysis detected PuHB40 in the nucleus (Fig. 4I). Accordingly, PuHB40 is a

transcriptional activator localized in the nucleus.

PuHB40 promotes *PuMYB123-like* transcript level

On the basis of the presence of four possible PuHB40-binding sites in the *PuMYB123-like* promoter (binding sites 1–4), we hypothesized that PuHB40 may modulate *PuMYB123-like* expression by binding to its promoter. To test this hypothesis, we replaced the possible binding sites with TTTTTTTTT. The dual-luciferase assay detected a significant decrease in the activation of the *PuMYB123-like* promoter by PuHB40 when binding site 4 was mutated (Fig. 5A). The electrophoretic mobility shift assay (EMSA) showed that PuHB40 can directly bind to the *PuMYB123-like* promoter fragment containing binding site 4 (Fig. 5B). The chromatin immunoprecipitation (ChIP) assay revealed the specific enrichment of PuHB40 in the *PuMYB123-like* promoter region harboring binding site 4 (Fig. 5C). Thus, the binding of PuHB40 to site 4 may activate *PuMYB123-like* expression.

To further verify the relationship between PuHB40 and *PuMYB123-like* expression, we transiently overexpressed *PuHB40* and silenced *PuMYB123-like* in pear seedlings. After a 3-day light treatment, anthocyanin accumulation was significantly greater in the *PuHB40*-OE seedlings than in the control seedlings, but the positive effect of PuHB40 on anthocyanin accumulation was significantly inhibited if *PuMYB123-like* was silenced (Fig. 5, D and E); a similar trend was detected in the expression of anthocyanin-related structural genes (Fig. 5F). These results confirmed that PuHB40 promotes anthocyanin biosynthesis in pear by regulating *PuMYB123-like* expression.

PuHB40 positively regulates anthocyanin biosynthesis

To analyze the PuHB40 function related to anthocyanin biosynthesis in pear, *PuHB40*-overexpressing (*PuHB40*-OE) transgenic pear calli and *PuHB40*-RNAi pear calli were generated. After a 2-day exposure to white light, the anthocyanin content was higher in the *PuHB40*-OE transgenic pear calli than in the control calli. Furthermore, anthocyanin accumulation was barely detectable in the *PuHB40*-RNAi pear calli (Fig. 6A-C). A qPCR analysis indicated that the anthocyanin-related genes *PcPAL*, *PcCHI*, *PcDFR*, *PcF3H*, *PcANS*, *PcUFGT*, *PcMYB10* and *PcMYB123-like* were highly expressed in the *PuHB40*-OE pear calli, whereas they were expressed at low levels in the *PuHB40*-RNAi pear calli (Fig. 6D). In addition, *PuHB40* was ectopically expressed in *A. thaliana*. Compared with the wild-type control, three independent transgenic lines ectopically overexpressing *PuHB40* had higher anthocyanin contents (Fig. 6E-G) and shorter hypocotyls (Fig. 6H). Moreover, the ectopic expression of *PuHB40* significantly increased the expression of anthocyanin-related structural genes (Fig. 6I).

To further analyze the *PuHB40* function in pear fruits, bagged mature 'Zaosu' and 'Hongzaosu' pear fruits were used for transient overexpression and VIGS experiments, respectively. The transient overexpression of *PuHB40* significantly promoted anthocyanin biosynthesis in the pear peel (Supplemental Fig. S7, A and B), while also significantly increasing the expression of anthocyanin-related structural genes and *PpMYB123-like* (Supplemental Fig. S7C). However, compared with the control fruit peel,

the *PuHB40*-TRV fruit peel had a lower anthocyanin content (Supplemental Fig. S7, D

and E) and lower anthocyanin-related structural gene and *PpMYB123-like* expression

levels (Supplemental Fig. S7F). Accordingly, PuHB40 may be involved in promoting

anthocyanin biosynthesis. Considered together, these results imply PuHB40 functions as

a positive transcriptional regulator governing anthocyanin production in pear.

Identification of PuPP2AA2 and its interaction with PuHB40

Because the increase in PuHB40 protein abundance occurred much earlier than the

increase in the *PuHB40* transcript level, we analyzed the possible post-translational

modifications of PuHB40. The PuHB40-interacting proteins in pear seedlings were

identified via a GST pull-down assay followed by a mass spectrometry analysis. Among

the 36 identified proteins, we focused on the A subunit of PP2A (Supplemental Table S4).

According to a phylogenetic analysis, the PP2A A subunit analyzed in the current study is

most closely related to the A2 subunit of PP2A in *A. thaliana.* Thus, it was named

PuPP2AA2 (Fig. 7A). During the subcellular localization analysis, PuPP2AA2 was

detected in the nucleus and cytoplasm (Fig. 7B). The *PuPP2AA2* expression level was

significantly lower after the high-light stress treatment than after the DMTU treatment

(Fig. 7C). To investigate the response of PuPP2AA2 to ROS at the translational level, we

transiently overexpressed the fusion construct encoding MYC-tagged PuPP2AA2

(PuPP2AA2-MYC) in pear seedlings and then exposed the plants to high-light stress

conditions with or without a DMTU treatment. The PuPP2AA2 protein abundance

decreased sharply under high-light conditions, but the decrease was significantly

attenuated after the DMTU treatment, which presumably decreased the ROS level (Fig.

7D). Thus, PuPP2AA2 is inhibited by high-light stress-induced ROS at both the

transcriptional and translational levels.

The GST pull-down assay showed that PuHB40 can interact with PuPP2AA2 *in*

vitro (Fig. 7E), whereas the LCI and bimolecular fluorescence complementation (BiFC)

assays reflected the interaction between PuHB40 and PuPP2AA2 *in vivo* (Fig. 7, F and

G). Taken together, these data confirmed that PuHB40 interacts with the A2 subunit of

PP2A, PuPP2AA2, *in vitro* and *in vivo*.

PuPP2AA2 inhibits PuHB40 trans-activation to *PuMYB123-like* by

dephosphorylating

To clarify whether the interaction between PuPP2AA2 and PuHB40 affects the

phosphorylation status of PuHB40, we transiently overexpressed the fusion construct

encoding GFP-tagged PuHB40 (PuHB40-GFP) in *Nicotiana benthamiana* leaves, which

were treated with cantharidin (i.e., a widely used PP2A inhibitor) (Honkanen 1993; Ren

et al. 2022). We also extracted proteins from the *N. benthamiana* leaves and separated

them in a Phos-tag gel and detected the fusion protein using an anti-GFP antibody. One of

the two bands detected in the gel corresponded to the phosphorylated form (P+) of

PuHB40-GFP, which migrated more slowly than the non-phosphorylated form (P−) of

PuHB40-GFP. Moreover, the P+:P− ratio increased after the cantharidin treatment (Fig.

8A), indicating that the phosphorylation level increased after PP2A activity was inhibited.

To further investigate whether PP2A can directly dephosphorylate PuHB40, we purified

the MYC, PuHB40-GFP, and PP2A complex from pear seedlings that were transiently

overexpressing the constructs encoding MYC, PuPP2AA2-MYC, and PuHB40-GFP

using MYC-Trap and GFP-Trap agarose beads. Because PuPP2AA2 is a structural

subunit that is tightly associated with the B and C subunits of the PP2A holoenzyme,

other PP2A subunits may be co-purified in the PuPP2AA2-MYC immunoprecipitation

complex (Zhou et al. 2004; Ren et al. 2022). Thus, the PP2A complex co-purified by

PuPP2AA2-MYC may contain all of the subunits and have phosphatase activity (Ren et

al. 2022). We incubated PuHB40-GFP with the MYC and PP2A complex, after which

different PuHB40-GFP isoforms were detected using the anti-PuHB40 antibody for the

Phos-tag analysis. The incubation with the PP2A complex led to the dephosphorylation of

PuHB40 and a decrease in the P+:P− ratio from 0.57 to 0.27 (Fig. 8B), implying PP2A

can directly dephosphorylate PuHB40. To confirm that the phosphatase activity of PP2A

was responsible for the decrease in the phosphorylation of PuHB40, we deleted the last

two tandem repeats (i.e., 14 and 15) of the HEAT motif in the C-terminal of PuPP2AA2.

The mutated sequence, which was named PuPP2AA2D, was unable to bind to the B and

C subunits of PP2A, thereby preventing the formation of a functional holoenzyme

(Ruediger et al. 1994). The subcellular localization analysis showed that PuPP2AA2D

was distributed in the nucleus as well as the cytoplasm, similar to the distribution of

PuPP2AA2 (Fig. 8C). The LCI and BiFC assays demonstrated that the deletion of tandem

repeats 14 and 15 did not affect the interaction between PuPP2AA2D and PuHB40 (Fig.

8, D and E).

We transiently coexpressed the fusion constructs encoding PuHB40-GFP and

PuPP2AA2-MYC as well as the fusion constructs encoding PuHB40-GFP and

MYC-tagged PuPP2AA2D (PuPP2AA2D-MYC) in *N. benthamiana* leaves. Different

isoforms of PuHB40-GFP were detected by the Phos-tag analysis involving the anti-GFP

antibody. Furthermore, the extent of the phosphorylation of PuHB40 decreased

significantly after the overexpression of *PuPP2AA2* (i.e., significant decrease in the

P+:P− ratio from 6.20 to 1.68), but there was no significant change in the

phosphorylation of PuHB40 after the overexpression of *PuPP2AA2D* (i.e.,

non-significant decrease in the P+:P− ratio from 6.20 to 5.83) (Fig. 8F). These results

indicate PP2A can dephosphorylate PuHB40 *in planta*. Hence, PP2A can

dephosphorylate PuHB40 *in vivo* and *in vitro*. Finally, to determine whether PuHB40

activity is influenced by its phosphorylation status, the PuHB40-induced transcription of

PuMYB123-like was determined in a dual-luciferase assay. The presence of PuPP2AA2

abolished the ability of PuHB40 to activate *PuMYB123-like* transcription. However, the

lack of PP2A phosphatase activity had no significant effect on PuHB40-activated

PuMYB123-like transcription (Fig. 8G). Therefore, PuHB40 may interact with

PuPP2AA2, with the resulting dephosphorylation by PP2A affecting whether PuHB40

can activate the transcription of *PuMYB123-like*.

PP2A attenuates PuHB40-induced anthocyanin biosynthesis

To elucidate the contribution of PuPP2AA2 to anthocyanin biosynthesis in pear, we

generated two *PuPP2AA2*-overexpressing transgenic pear calli lines (*PuPP2AA2*-OE #5

and #8), which were subsequently exposed to strong white light for 3 days. The

anthocyanin contents were much lower in *PuPP2AA2*-OE #5 and #8 than in the wild-type

pear calli (Fig. 9, A and B). The qPCR analysis indicated that the overexpression of

PuPP2AA2 downregulated the expression of anthocyanin-related genes (Fig. 9C). We

also transiently silenced and overexpressed *PuPP2AA2* in pear fruit peels. At the

injection site of the *PuPP2AA2*-OE fruit peels, there was almost a complete lack of red

coloration and the anthocyanin content was lower than that in the control fruit peels

(Supplemental Fig. S8, A and B). Moreover, the expression levels of anthocyanin-related

genes were significantly lower in the *PuPP2AA2*-OE pear fruit peels than in the control

fruit peels (Supplemental Fig. S8C). In contrast, the transient silencing of *PuPP2AA2*

significantly promoted anthocyanin accumulation (Supplemental Fig. S8, D and E) and

increased the expression of the related structural genes (Supplemental Fig. S8F)

suggestive of the negative regulatory effects of PuPP2AA2 on anthocyanin biosynthesis

and the red coloration of pear fruits.

To assess whether the phosphorylation status of PuHB40 affects its function

associated with ROS-induced anthocyanin biosynthesis in pear, we transiently

overexpressed *PuHB40* alone as well as with *PuPP2AA2* in pear seedlings, which were

then subjected to a 2-day light treatment. Anthocyanin accumulation and the expression

of anthocyanin-related genes as well as *PuMYB123-like* were greater in the pear seedlings

overexpressing *PuHB40* alone than in the control seedlings. In contrast, there were no

obvious differences in anthocyanin accumulation and the expression of the related genes

between the control pear seedlings and the seedlings coexpressing *PuHB40* and

PuPP2AA2 (Fig. 9, D-F). Additionally, bagged mature 'Hongzaosu' pear fruits were used

for the transient overexpression of *PuHB40* alone and with *PuPP2AA2*. Compared with

the corresponding control levels, the anthocyanin contents and the expression levels of

anthocyanin-related genes increased significantly in the pear fruit peels overexpressing

PuHB40 alone, but the presence of PuPP2AA2 weakened the effects of PuHB40 on

anthocyanin biosynthesis and the expression of the related genes (Fig. 9, G-I). Thus, the

dephosphorylation of PuHB40 by PP2A adversely affects anthocyanin biosynthesis.

DISCUSSION

Plants growing under natural conditions over long periods are increasingly becoming

more susceptible to different stresses because of the variability in environmental

conditions due to global warming and climate change (Lesk et al. 2016). In plant cells,

stress can induce the rapid accumulation of ROS, which are important signaling

molecules that activate plant mechanisms related to defense and acclimatization (Mittler

et al. 2022). However, excessive amounts of ROS can damage plant cells and may lead to

cell death (Mittler 2017) In response to a wide range of stresses and to maintain ROS

homeostasis, plants have evolved a series of mechanisms, including those mediating

anthocyanin accumulation.

In plants, anthocyanins are crucial antioxidants that function as ROS scavengers

(Gould et al. 2002; Neill et al. 2002; Bi et al. 2014; Jiang et al. 2023). Many studies

showed that anthocyanin accumulation is strongly induced by various stresses, such as

high-light stress, ultraviolet radiation, and low temperatures (Steyn et al. 2002; Catala et

al. 2011; Xie et al. 2012; Nakabayashi et al. 2014; Landi et al. 2015; Xu et al. 2015). An

earlier investigation involving raman spectroscopy confirmed that abiotic stress

conditions increase the ROS and anthocyanin contents in the leaves of coleus plants

(Altangerel et al. 2017). However, the link between ROS and anthocyanins remains

unclear. Although treating $A.$ $thaliana$ seedlings with H_2O_2 and ROS activators promotes

anthocyanin accumulation (Xu et al. 2017; Shi et al. 2018), evidence of the relationship

between ROS and anthocyanin accumulation is lacking. In this study, the ROS scavenger

DMTU was applied to eliminate ROS induced by various stresses, thereby serving as a

reliable negative control. Stress-induced anthocyanin accumulation in pear seedlings was

attenuated by DMTU (Supplemental Fig. S1A), indicating that the increase in the

anthocyanin content depends on ROS, which likely function as signaling molecules

linking the exposure to different stresses to changes in anthocyanin contents.

Anthocyanin biosynthesis is regulated by the MBW complex, with MYB TFs

playing a major role and responding to various environmental and internal factors (Xu et

al. 2015). R2R3-MYB proteins of subgroup 6 is responsible for anthocyanin biosynthesis

(Dubos et al. 2010). In rosaceous plants, MYB10 TFs, which belong to subgroup 6, are

essential promotors of light-induced anthocyanin biosynthesis (Takos et al. 2006; Espley

et al. 2007; Feng et al. 2010). Although most of the MYB TFs promoting anthocyanin

biosynthesis belong to subgroup 6, a few members of other subgroups are also involved.

For example, the subgroup 2 members MdMYB9 and MdMYB11 promote jasmonic

acid-induced anthocyanin biosynthesis by increasing the expression of

anthocyanin-related structural genes (An et al. 2015), whereas MdMYB308L, which is a

member of subgroup 4, promotes low-temperature-induced anthocyanin biosynthesis by

interacting with MdbHLH33 (An et al. 2020). In the current study, we determined that

PuMYB123-like, which belongs to subgroup 2, is the key MYB TF mediating

anthocyanin biosynthesis via the ROS-responsive pathway. Traditionally, PuMYB10 has

been considered the main MYB TF responsible for anthocyanin biosynthesis in pear

(Feng et al. 2010; Bai et al. 2019a), however, its expression is similar following the

high-light stress and DMTU treatments (Fig. 1K). The MYB subgroup 2 members are

primarily involved in proanthocyanidin biosynthesis, but in the present study, the

overexpression of *PuMYB123-like* did not affect proanthocyanidin accumulation in pear

calli (Supplemental Fig. S4A). Instead, PuMYB123-like promoted anthocyanin

biosynthesis by interacting with PubHLH3 (Fig. 3). Our study findings suggest that

members of other MYB subgroups may also have important regulatory effects on

anthocyanin biosynthesis, further clarifying the involvement of MYB TFs in

ROS-associated anthocyanin biosynthesis.

According to earlier research, HD-Zip I family members are crucial regulators of

plant responses to stress, especially drought and osmotic stress (Ariel et al. 2007; Harris

et al. 2011). But their role in other stresses is less well known. In this study, The HD-Zip I

family member PuHB40 was identified as a key TF involved in ROS-regulated

anthocyanin biosynthesis under high-light stress (Fig. 4). Considering the important role

of HD-Zip I family members in various stresses, PuHB40 regulation pathway for

high-light stress-induced anthocyanin biosynthesis might be universal in different stresses.

The response of HD-Zip I TFs to drought stress depends on abscisic acid (ABA)

(Soderman et al. 1996; Söderman et al. 1999; Olsson et al. 2004), but the relationship

with ROS remains unknown. In the current study, PuHB40 responded to high-light stress

in an ROS-dependent manner (Fig. 4B), mplying HD-Zip I TFs can respond to other

abiotic stresses (i.e., in addition to drought) through multiple signaling pathways. The

stability and activity of HD-Zip proteins are regulated by the phosphorylation. In apple,

MdHB1 and MdHB2 are phosphorylated by SnRK2 kinase, which enhances their protein

stability and activity, thereby activates *MdACO1* expression and promotes ethylene

biosynthesis (Jia et al. 2022). In Arabidopsis, ATHB6 is dephosphorylated by the PP2C

phosphatase ABI1 by interacting with it directly, thus functions as a negative regulator in

the ABA signaling pathway (Himmelbach et al. 2002). The phosphorylation status of

HD-Zip I TFs is mainly associated with phytohormone signaling pathway, especially with

ABA. In the present study, the phosphorylation of PuHB40 was regulated by ROS via

PP2A (Fig. 8). The decrease in the extent of the phosphorylation of PuHB40 decreased

the PuHB40-activated transcription of *PuMYB123-like*, ultimately decreasing

ROS-induced anthocyanin biosynthesis (Fig. 9, D and G). In plants, PP2A is a ubiquitous

and highly conserved serine/threonine-specific phosphatase (Farkas et al. 2007) that

catalyzes dephosphorylations with diverse regulatory effects on plant stress responses

(Lillo et al. 2014). In general, under salt and osmotic stress conditions, PP2A is inhibited,

resulting in phosphorylated target proteins that help protect against various stresses (Bian

et al. 2020; Li et al. 2020). However, the effects of ROS on PP2A phosphatase activity

under abiotic stress conditions are unclear. In this study, we observed that the structural

subunit of PP2A was repressed by ROS at the transcriptional and translational levels in

pear (Fig. 7, C and D), suggesting that the regulatory effects of PP2A on plant stress

responses may involve ROS. The A subunit of PP2A is the structural subunit that

provides a scaffold that binds to the B and C subunits (Kamibayashi et al. 1992; Ruediger

et al. 1994; Groves et al. 1999; Janssens and Goris 2001). In *A. thaliana*, three A subunits

of PP2A have been identified (RCN1, PP2AA2, and PP2AA3). A mutation to RCN1

results in severe phenotypic changes, including stunted growth (dwarfism), abnormal

embryogenesis, and sterility. This is in contrast to the mostly normal phenotypes of the

pp2aa2 and *pp2aa3* mutants (Zhou et al. 2004). Thus, it is widely believed that RCN1 is

critical for the overall regulation of PP2A, which may help to explain why there have

been relatively few studies on PP2AA2 and PP2AA3 (Lillo et al. 2014). In the current

study, we revealed that PuPP2AA2 affects ROS-induced anthocyanin biosynthesis by

dephosphorylating PuHB40 (Fig. 10), which clarifies the role of the A subunit of PP2A in

plants and highlights the importance of the other A subunits.

In summary, PuHB40 is the key TF regulating ROS-induced anthocyanin

biosynthesis. Specifically, it is a transcriptional activator that induces

PuMYB123-like/PubHLH3 complex for anthocyanin biosynthesis. The ability of

PuHB40 to activate target gene transcription depends on its phosphorylation level, which

is regulated by PP2A. Furthermore, ROS accumulation leads to the extensive

phosphorylation of PuHB40 by suppressing PuPP2AA2 production at the transcriptional

and translational levels, thereby enhancing the PuHB40-activated transcription of

PuMYB123-like and promoting anthocyanin biosynthesis (Fig. 10). Our study findings

have elucidated the molecular mechanisms by which abiotic stresses regulate anthocyanin

biosynthesis in pear, providing insights into how ROS modulate anthocyanin

biosynthesis.

MATERIALS AND METHODS

Plant materials and treatment

Pear seedlings 'Qingzhen No. 1' (*Pyrus ussuriensis*: genes from it are prefixed with 'Pu')

were kindly provided by Professor Wang Ran of Qingdao Agricultural University. The

pear seedlings were growth at 25 °C under a 16 h light/ 8 h dark photoperiod (photon flux

density 50 $\mu mol \cdot s^{-1} \cdot m^{-2}$). For high-light stress treatment, pear seedlings were exposed to

continuous LED white light (photon flux density 160 $\mu mol \cdot s^{-1} \cdot m^{-2}$) in a growth chamber.

For salt and drought stress, pear seedlings were growth on solid MS medium

supplemented with 30 mM sodium chloride and 2.5% polyethylene Glycol 8000 (PEG

8000) respectively and were exposed to continuous LED white light (photon flux density

50 $\mu mol \cdot s^{-1} \cdot m^{-2}$). For N, N'-dimethylthiourea (DMTU) treatment, seedlings were growth

on solid MS medium supplemented with 5 mM DMTU and other condition is same as

stress treatment. After treatment, the seedlings were collected, immediately frozen in

liquid nitrogen, and stored at -80 °C until used.

Pear calli were generated from the flesh of 'Clapp's Favorite' (*Pyrus communis*:

genes from it are prefixed with 'Pc') fruitlets as previously described (Bai et al. 2019b).

For treatment, the pear calli on good condition were subcultured to the MS solid medium

supplemented with 4 mM H_2O_2 for overnight firstly and then were exposed to continuous

white light. The pear calli were collected after treatment and stored at -80 °C until used.

Measurement of the total anthocyanin content

The measurement of anthocyanin content is as previously described (Zhang et al. 2022).

Briefly, 0.1 g powder of seedlings, frozen pear calli or peels were soaked in 500 μL

methanol: hydrochloric acid (99:1, v/v) solution and incubated overnight at 4 °C in

darkness. For pear seedlings, 250 μL chloroform and 250 μL double distilled water were

added to the extracting solution for mixing and centrifuged. The supernatant was

measured at 530, 657 nm using the DU800 spectrophotometer (Beckman Coulter, Brea,

CA, USA).

For pear calli and peels, the extracting solution centrifuged at 4 °C, 12000 rpm, for

10 minutes. And then, the absorbance of supernatant was measured at 530, 620, and 650

nm. The anthocyanin content was calculated as the following formula:

$([A_{530}-A_{650}]-0.2\times[A_{650}-A_{620}])/0.1$ for pear calli as well as peels and $(A_{530}-0.25*A_{657})/0.1$

for seedlings.

Histological staining detection of ROS

H_2O_2 and O^{2-} were detected as previously described with minor modification (Xu et al.

2017). Briefly, the leaves of pear seedlings were dipped in DAB staining solution (for

H_2O_2 detection) containing 1 mg ml^{-1} DAB and NBT staining solution (for O^{2-} detection)

containing 1 mg ml^{-1} NBT. The leaves were incubated for overnight in the dark with

gentle shaking (25 °C, 220 rpm min^{-1}). Then, the chlorophyll of leaves was removed with

bleaching solution (ethanol: acetic acid: glycerol=3:1:1) by incubating at 85 °C for 30

min. Finally, the leaves were kept in preservative solution (ethanol: glycerol = 4: 1) until

images were taken.

RNA isolation and reverse transcription quantitative PCR

Total RNA was extracted with a modified CTAB method (Zhang et al. 2012). Genomic

DNA removal and first-strand cDNA synthesis from 1 μg RNA were performed with

HiScript II First Strand cDNA Synthesis Kit (+gDNA wiper; Vazyme Biotech, China).

qPCR was conducted using ChamQ SYBR qPCR Master Mix (Vazyme, China) on CFX

Connect real-time PCR system (Bio-Rad, https://www. bio-rad.com/). The relative

expression level of genes was calculated using the $2^{-\triangle\triangle Ct}$ method (Livak and Schmittgen

2001) with the pear *PpActin* gene (GenBank accession no. JN684184) to normalize the

expression values, and the analyses were carried out with three biological replicates. The

primers for RT-qPCR are listed in Supplemental Data Set S2.

H_2O_2 content quantification

The powder of pear seedlings leaves was soaked in 1 ml of 20 mM potassium phosphate

buffer (pH 6.5) on ice and centrifuged, then the supernatant was transferred and

additional potassium phosphate buffer was added to 1.5 ml. A 100 μL plant extract was

used in each reaction and the H_2O_2 concentration was measured with the Amplex Red

hydrogen peroxide/peroxidase assay kit (Thermo Fisher Scientific, Cat. No. A22188).

Subcellular localization analysis

Subcellular localization analysis was performed as previously described (Ni et al. 2023).

the full-length complete coding sequences (CDS) (without termination codons) of the

target genes (*PuMYB123-like*, *PuHB40*, *PuPP2AA2* and *PuPP2AA2D*) were inserted into

pCAMBIA1300 vector to produce *Pro35S: PuMYB123-like-GFP*, *Pro35S: PuHB40-GFP*,

Pro35S: PuPP2AA2-GFP, and *Pro35S: PuPP2AA2D-GFP*, respectively. *Agrobacterium*

strain GV3101 harboring the plasmids were infiltrated into the leaves of *N. benthamiana*

transgenic lines containing red fluorescent protein in the nuclei. The GFP and mCherry

fluorescence signals were observed by confocal laser scanning microscopy (A1, Nikon,

Tokyo, Japan). The primers used for vector construction are listed in Supplemental Data

Set S1.

Trans-activation activity assay

A trans-activation activity assay was performed as previously described (Zhang et al.

2022). Briefly, the full-length CDS of *PuMYB123-like* and *PuHB40* was cloned to the

pGBKT7 (BD) vector and transformed into yeast *(Saccharomyces cerevisiae)* strain

AH109 cells independently. The yeast only containing BD vector is as negative control

and yeast containing BD vector with the VP16 fragment (VP16-BD) is as positive control.

The yeast transformants were screened on a selective synthetic dextrose medium lacking

tryptophan and histidine (SD/-Trp/-His) supplemented with X-α-gal. The primers used for

vector construction are listed in Supplemental Data Set S1.

Transient transformation assay in pear fruit and seedlings

'Hongzaosu' (also called 'Red Zaosu') pear fruits (*Pyrus pyrifolia* × *Pyrus communis*:

genes from it are prefixed with 'Pp') were collected from the orchard at the Institute of

Horticulture, Henan Academy of Agricultural Sciences, Henan, China. Fruits were

covered with lightproof double-layered paper bags during the growing season and were

harvested about 140 d after full bloom, then immediately transported to the laboratory.

Transient transformation assay in pear fruit was performed as previously described (Ni et

al. 2023). For the transient overexpression, the full-length CDS of *PuMYB123-like*,

PuHB40 and *PuPP2AA2* was cloned to the pGreenII 0029 62-SK vector and their

expression was driven by the 35S promoter. Pear fruits were infiltrated with *A.*

tumefaciens GV3101 cells containing the plasmid by a needle syringe. For VIGS assay,

the gene fragment (~200 to 300 bp) was inserted into the pTRV2 vector (gene-pTRV2)

and transformed into *A. tumefaciens* GV3101. Gene-pTRV2 or empty pTRV2 (negative

control) was mixed with pTRV1 in equal proportions and co-infiltrated into pear fruits by

a needle syringe. After infiltration, pear fruits were exposed to continuous white light (60

$\mu mol \cdot m^{-2} \cdot s^{-1}$) and UV-B light (1.1 $\mu mol \cdot m^{-2} \cdot s^{-1}$). The fruit peel surrounding the injection

sites was collected and stored at -80 °C until use.

For transient transformation assay in pear seedlings, the full-length CDS of *PuHB40*,

PuPP2AA2 and reverse complementary gene fragment (~200 to 300 bp) of

PuMYB123-like (*PuMYB123-like-RNAi*) were cloned to pCAMBIA1301, and then the

plasmids transformed into *A. tumefaciens* GV3101. *A. tumefaciens* GV3101 with

PuHB40 was mixed with pCAMBIA1301, *PuMYB123-like-RNAi*, *PuPP2AA2* in equal

proportions and co-infiltrated into seedlings by vacuum (0.8 kg·cm^{-2}, 5 min). The

seedlings were then incubated in darkness for 1-2 d and then transferred to a growth

chamber for treatment under continuous white light. Finally, the seedlings were collected,

frozen in liquid nitrogen immediately, and stored at -80°C until used. The primers used

for vector construction are listed in Supplemental Data Set S1.

Genetic transformation of pear calli

Transgenic pear calli were generated as previously described (Bai et al. 2019b). Briefly,

pear calli were soaked with *A. tumefaciens* EHA105 containing different vectors

(PuHB40-pCAMBIA1301, PuMYB123-like-pCAMBIA1301,

PuPP2AA2-pCAMBIA1301 for overexpression and PuHB40-RNAi-pCAMBIA1301,

PuMYB123-like-RNAi-pCAMBIA1301 for silencing) for 12-15 min and then

co-cultured on MS solid medium for 2 d in darkness. Positive pear calli were screened on

solid MS medium supplemented with antibiotics (10 mg/L hygromycin B) under

continuous darkness at 24 °C for 1-3 months and subcultured every 3 weeks. For light

treatment, pear calli in good condition were exposed to continuous white light. The

primers used for vector construction are listed in Supplemental Data Set S1.

Yeast hybrid assays (Y1H and Y2H)

Yeast one-hybrid assays were performed using the Matchmaker Gold Yeast One-Hybrid

System Kit (TaKaRa) according to the manufacturer's protocol. Briefly, the CDS of

PuHB40 was cloned into the pGADT7 vector, while the promoter fragment of

PuMYB123-like was ligated into the pAbAi vector. The yeast (*S. cerevisiae*) strain Y1H

Gold cells containing both of *PuHB40*-AD and pAbAi-bait were screened on SD/-Leu

medium containing an appropriate AbA concentration. The Y2H assay was carried out

based with the manufacturer's protocol of Matchmaker Gold Yeast Two-Hybrid System

Kit (Takara). The primers used for vector construction are listed in Supplemental Data

Set S1.

Pull down screening coupled with LC-MS/MS

The GST-tagged PuHB40 protein (PuHB40-GST) bound to GST-Trap agarose beads were

incubated with total proteins of pear seedlings at 4°C for overnight, the GST protein is as

negative control. The PuHB40-GST and GST proteins bound to beads were eluted by

glutathione (GSH) with gravity column and subjected to immunoblotting analysis with

anti-GST antibody. The eluent solution contained PuHB40-GST and GST proteins as well

as proteins interacting with PuHB40 in pear seedlings, and then these proteins were

further identified by mass spectrometry.

For pull-down assay, His-tagged PuMYB123-like were incubated with GST and

GST-tagged PubHLH3 respectively supplemented with GST-Trap agarose beads for

overnight, similarly, His-tagged PuHB40 were incubated with GST and GST-tagged

PuPP2AA2 respectively. Then they were eluted by GSH and subjected to immunoblotting

analysis with anti-GST antibody and anti-His antibody. The primers used for vector

construction are listed in Supplemental Data Set S1.

Luciferase complementation imaging assays

The coding sequences (without termination codons) of PuHB40, PubHLH3 was inserted

into pCAMBIA1300-nLUC vectors, and the CDS of PuMYB123-like, PuPP2AA2 and

PuPP2AA2D was cloned to pCAMBIA1300-cLUC vectors. The vectors were introduced

into *Agrobacterium* strain GV3101 individually and *A. tumefaciens* harboring

gene-nLUC were mixed with an equal volume of *Agrobacteria* containing gene-cLUC

and then co-infiltrated into *N. benthamiana* leaves. Five minutes before detection, 10 mM

beetle luciferin (Promega) was injected into the same position as the agrobacteria and

then captured by ChemiDoc™XRS+ (Bio-Rad). The *N. benthamiana* leaves co-infiltrated

with empty vectors expressing cLUC and nLUC as a negative control. The primers used

for vector construction are listed in Supplemental Data Set S1.

Bimolecular fluorescence complementation assay

The full-length PuHB40 was inserted into p2YC and p2YN, while PuPP2AA2 and

PuPP2AA2D were cloned to p2YN and p2YN respectively. Then, *A. tumefaciens*

GV3101 cells harboring gene-p2YN and gene-p2YC mixed in equal proportions and

co-infiltrated into *N. benthamiana* leaves by injection, Leaves were co-infiltrated with

gene-2YN and 2YC or 2YN and 2YC-gene as the negative control. The YFP signal was

detected by a confocal laser scanning microscope (Nikon, Japan) after 48 h-72 h. The

primers used for vector construction are listed in Supplemental Data Set S1.

Dual-luciferase assay

Dual-luciferase assays were performed as previously reported (Bai et al. 2019a). The

full-length CDS of PuHB40, PuPP2AA2 and PuPP2AA2D were cloned into pGreenII

0029 62-SK vector (gene-SK) and the fragments of *PuMYB123-like* promoter were

ligated into pGreenII 0800-LUC vector (promoter-LUC). All vectors were transformed

into *A. tumefaciens* strain GV3101 cells harboring the pSoup vector. *A. tumefaciens* strain

GV3101 contaning gene-SK and promoter-LUC were injected into *N. benthamiana*

leaves at a ratio of 1:10. The firefly luciferase and Renilla luciferase activities were

analyzed at 48 h after the infiltration by the Dual-Luciferase Reporter Assay System

(Promega) and the GloMax96 Microplate Luminometer (Promega). The primers used for

vector construction are listed in Supplemental Data Set S1.

Electrophoretic mobility shift assay

The 3' end biotin-labeled double-stranded DNA probes were prepared by annealing 3'

end biotin-labeled oligonucleotides to the corresponding complementary strands (95 °C

for 5 min and 70 °C for 20 min). The EMSA were performed using the Light-Shift[TM]

Chemiluminescent EMSA Kit (Thermo Scientific) based with manufacturer's protocol.

Briefly, recombinant PuHB40-His protein was incubated with labeled or nonlabeled

double-stranded probes at room temperature for 20 min in binding buffer. The samples

were then separated by 6.5% (w/v) native polyacrylamide gels at 4 °C and the

biotin-labeled probes was captured with ChemiDoc™XRS+ (Bio-Rad). The

oligonucleotide probes used in this study are listed in Supplemental Data Set S1.

ChIP assay

The ChIP assays were performed as previously described (Ni et al. 2023). The pear calli

transformed with the recombinant PuHB40-MYC construct and the empty construct

(MYC alone) were prepared for ChIP assays. The pear calli were treated in 1% (w/v)

formaldehyde under vacuum for 10 min for the cross-linking and then the chromatin was

extracted via sucrose gradient centrifugation. The chromatin DNA was sonication for 30

min on ice (30 s with 30 s intervals) with the Bioruptor Plus device (Diagenode) to shear

the chromatin to an average size of 200 bp to 500 bp. The sheared chromatin was

incubated with Anti-MYC (Sigma-Aldrich, 05-724) at 4 °C overnight and the amount of

immunoprecipitated chromatin was determined by qPCR. Each ChIP assay was repeated

3 times, and the enriched DNA fragments in each ChIP sample served as 1 biological

replicate for qPCR. The primers used for the ChIP-qPCR analysis are listed in

Supplemental Data Set S3.

Mn^{2+}-Phos-tag SDS-PAGE

Prior to Phos-tag analysis, samples need to be pretreated by TCA precipitation and then

protein samples were detected by 10% regular SDS-PAGE supplemented with 60 μM

Phos-tag (TM) Acrylamide AAL-107 (Wako) and 120 μM MnCl$_2$. Wash the gels by

gently agitating it 10 minutes for 3 times using general transfer buffer supplemented with

10 mmol/L EDTA after electrophoresis for 1.5 h and then washed by general transfer

buffer for 10 min. Finally, the gels were used for immunoblotting analysis.

Protein dephosphorylation assays *in vitro* and *vivo*

For dephosphorylation assays in vitro, *A. tumefaciens* GV3101 with PuHB40-GFP,

PuPP2AA2-MYC and MYC infiltrated into pear seedlings by vacuum (0.8 kg·cm^{-2}, 5

min). Then, PuHB40-GFP, PuPP2AA2-MYC and MYC were purified by

immunoprecipitation from pear seedlings with GFP-Trap and MYC-Trap agarose beads.

Purified PuPP2AA2-MYC was used as PP2A complex, since other subunits of PP2A

could co-immunoprecipitate by PuPP2AA2. Equal volume of purified PuHB40-GFP

mixed with PuPP2AA2-MYC and MYC respectively and then incubated an oscillator

(1200 g, 30 °C) for 2 h, which was terminated by adding protein loading buffer and

boiled at 98 °C for 10 min. Finally, the mixtures were used for immunoblotting analysis.

For dephosphorylation assays *in vivo*, *A. tumefaciens* GV3101 with PuHB40-GFP

was mixed with MYC, PuPP2AA2-MYC, PuPP2AA2D-MYC in equal proportions and

co-infiltrated into *N. benthamiana* leaves by injection. The total proteins were extracted

from *N. benthamiana* leaves after 48 h-72 h and subjected to TCA precipitation,

immediately followed by Phos-tag analyses. For cantharidin treatment, *A. tumefaciens*

GV3101 with PuHB40-GFP infiltrated into *N. benthamiana* leaves by injection and 50

μM cantharidin was injected into the same position as the agrobacteria after 24 h, and

then treated for 24 h. Finally, the total proteins were extracted and then used for Phos-tag

analyses.

Statistical analysis

All data in this study were analyzed using two-sided Student's *t*-test or one-way analysis

of variance (ANOVA) to determine statistical significance with GraphPad Prism v.9.0.

Detailed statistical analysis data are shown in Supplemental Data Set S5.

ACCESSION NUMBERS

Sequence data from this article can be found in an online pear genome database

(https://ngdc.cncb.ac.cn/gwh/Assembly/18534/show): PuHB40 (EVM0020592.1),

PuMYB123-like (EVM0034726.1), PuPP2AA2 (EVM0003533.1), PubHLH3

(EVM0031731.1), PubHLH33 (EVM0038434.1), PubHLH64 (EVM0003136.1),

PuELIP1 (EVM0019834.1), PuELIP2 (EVM0035306.1), PuCHS (EVM0036585.1),

PuCHI (EVM0024725.1), PuF3H (EVM0021098.1), PuDFR (EVM0012887.1), PuANS

(EVM0001840.1), PuUFGT (EVM0026501.1), PuMYB10 (EVM0039871.1),

PuMYB114 (EVM0020507.1).

SUPPLEMENTAL DATA

Supplemental Figure S1. Stresses induced anthocyanin biosynthesis is ROS-dependent.

Supplemental Figure S2. Identification of PuMYB123-like.

Supplemental Figure S3. The interaction between PuMYB123-like and PubHLHs.

Supplemental Figure S4. Functional analysis of PuMYB123-like in 'Hongzaosu' and

'Zaosu' pear fruit.

Supplemental Figure S5. Detection of specific anti-PuHB40 polyclonal antibody.

Supplemental Figure S6. Identification of PuHB40.

Supplemental Figure S7. Functional analysis of PuHB40 in 'Hongzaosu' and 'Zaosu'

pear fruit.

Supplemental Figure S8. Functional analysis of PuPP2AA2 in 'Hongzaosu' pear fruit.

Supplemental Data Set S1. Primers used for constructing vectors or EMSA in this study.

Supplemental Data Set S2. Primers for qPCR analysis in this study.

Supplemental Data Set S3. Primers for ChIP-qPCR analysis in this study.

Supplemental Data Set S4. PuHB40 possible interacting proteins were identified with

GST pull-down combined with mass spectrometry.

Supplemental File S1. Protein sequences used to generate phylogenetic tree in Fig. 2B.

Supplemental File S2. Phylogenetic tree of Fig. 2B in Newick format.

Supplemental File S3. Protein sequences used to generate phylogenetic tree in Fig. 7A.

Supplemental File S4. Phylogenetic tree of Fig. 7A in Newick format.

Supplemental File S5. Protein sequences used to generate phylogenetic tree in Fig. S2A.

Supplemental File S6. Phylogenetic tree of Fig. S2A in Newick format.

Supplemental File S7. Protein sequences used to generate phylogenetic tree in Fig. S6A.

Supplemental File S8. Phylogenetic tree of Fig. S6A in Newick format.

ACKNOWLEDGMENTS

We thank R. Wang at Qingdao Agricultural University for providing pear seedlings of

'Qingzhen No. 1' (*Pyrus ussuriensis*). We also thank Kunfeng Li for technical help and

providing plant materials.

FUNDING INFORMATION

This work was supported by the Zhejiang Provincial Natural Science Foundation of

China under Grant No. LR22C150001, The National Natural Science Foundation of

China (32072545 to YT, 32272639 to SB, 32269745 to JS), and the China Agriculture

Research System of MOF and MARA.

AUTHOR CONTRIBUTIONS

Y.T and S.B. planned the work. S.B. and L.Z. conceived and designed the experiments.

L.Z. performed most experiments. L.W., S.Y. helped conduct some of the experiments.

Y.G. and J.S. helped to analyze the data. L.Z. and S.B wrote the manuscript and, Y.T., J.N.

helped to revise the manuscript. All authors read and approved the final paper.

DECLARATION OF INTERESTS

837   The authors declare that they have no competing interests.

FIGURE LEGENDS

Figure 1. High light stress-induced anthocyanin biosynthesis is ROS-dependent.

A-K), Pear seedlings were treated with high light stress treatment (photon flux density:

approximately 160 μmol·s^{-1}·m^{-2}) or high light stress supplemented with N,

N'-dimethylthiourea (DMTU) treatment. DMTU was applied as ROS scavenger, and pear

seedlings under appropriate light condition were used as Mock (photon flux density:

approximately 50 μmol·s^{-1}·m^{-2}). Pear seedlings were sampled at 0 h, 6 h, 12 h, 24 h, 48 h,

72 h, 96 h. **A)**, Phenotype of pear seedlings (scale bar, 1 cm). **B)**, The anthocyanin

content for pear seedlings. **C and D)**, NBT staining for superoxide and DAB staining for

H_2O_2 in pear seedlings leaves, respectively (scale bar, 0.5 mm). **E)**, H_2O_2 quantification of

pear seedlings. **F-K)**, The expression levels of *PuCHI*, *PuCHS*, *PuDFR*, *PuF3H*, *PuANS*,

PuMYB10 in pear seedlings under different treatments. *PuACTIN* was amplified as an

internal control. Data are presented as means ± s.d. of 3 biological replicates. Asterisks

indicate significant differences compared with Mock group (two-tailed Student's *t-test*,

*$P < 0.05$, **$P < 0.01$), all exact P values are shown in Supplemental Data Set S5.

Figure 2. Identification of PuMYB123-like and its interaction with PubHLH3.

A) The expression levels of *PuMYB123-like* in pear seedlings under high-light stress

treatment or high-light stress supplemented with DMTU treatment, and pear seedlings

under appropriate light condition were used as Mock. *PuACTIN* was amplified as an

internal control. **B)** Phylogenetic analysis of PuMYB123-like. The phylogenetic tree was

generated using the Maximum-likelihood method of MEGA (version 10.1.8). Bootstrap

values from 1000 replicates for each branch are shown. The protein sequences from pear

(Pu, *Pyrus pyrifolia*, highlighted with blue dots), apple (Md, *Malus × domestica*,

highlighted with green dots), Arabidopsis (At, *Arabidopsis thaliana*, highlighted with

rose red dots) and strawberry (Fa, *Fragaria × ananassa*, highlighted with purple dots).

PuMYB123-like (EVM0034726.1) is bolded and enlarged. **C)** Subcellular localization of

PuMYB123-like in *N. benthamiana* leaf cells (scale bar, 25 μm). **D)** Trans-acting activity

of PuMYB123-like in yeast cells. **E)** Interaction between PuMYB123-like and PubHLH3

in a yeast two-hybrid assay, with pGADT7-T and pGBKT7-53 as positive controls. DDO,

SD/-Trp/-Leu medium; QDO/X/A, SD/-Trp/-Leu/-Ade/-His medium with X-α-gal and

Aureobasidin. **F)** The interaction between PuMYB123-like and PubHLH3 was detected

by luciferase complementation imaging assays. The pseudocolour scale bar indicates the

range of luminescence intensity. **G)** The interaction between PuMYB123-like and

PubHLH3 analyzed using a pull-down assay. PuMYB123-like-His was incubated with

GST or PubHLH3-GST. The interaction proteins were precipitated by GST-Trap agarose

beads and analysed by immunoblotting with an anti-GST or anti-His antibody. **H)** A

dual-luciferase assay demonstrated that PuMYB123-like promotes the promoter activity

of anthocyanin-related genes (*PuCHS*, *PuANS*, *PuDFR*, *PuUFGT*). The promoter of

anthocyanin-related genes was cloned into the pGreenII 0800-LUC (firefly luciferase)

vector, and the full-length CDS of *PuMYB123-like* was cloned into the pGreenII 0029

62-SK vector. The empty vector of pGreenII 0029 62-SK was used as control. **I)**

Dual-luciferase assay showed that interaction between PuMYB123-like and PubHLH3

significantly enhanced activation of PuMYB123-like on *PuCHS* and *PuANS* promoters.

Data are presented as means ± s.d. of 3 biological replicates **A)** and 6 biological replicates

H) and **I)**. Asterisks indicate significant differences compared with Mock **A)**, empty

vector **H)** and PuMYB123-like **I)** (two-tailed Student's *t-test*, $*P < 0.05$, $**P < 0.01$), all

exact *P* values are shown in Supplemental Data Set S5.

Figure 3. PuMYB123-like promotes anthocyanin biosynthesis.

A) Phenotype of pear calli overexpressing and silencing *PuMYB123-like* (scale bar, 1 cm). Wild type pear calli were used as the negative control. Pear calli were incubated under continuous white light at 17°C for 3 d. **B)** Expression levels of anthocyanin-related genes and *PuMYB123-like* in *PuMYB123-like*-OE *and PuMYB123-like*-RNAi pear calli after light treatment. *PcACTIN* was amplified as an internal control. **C)** Anthocyanin contents in *PuMYB123-like*-OE and *PuMYB123-like*-RNAi pear calli after light treatment. **D)** Phenotype of Arabidopsis seedlings overexpressing *PuMYB123-like* (scale bar, 2 mm). Wild-type 'Columbia 0' was used as the negative control. Arabidopsis seedlings were grown on 1/2 MS medium for 5 days. **E)** Expression levels of *PuMYB123-like* in transgenic Arabidopsi*s* lines. *AtPP2A* was amplified as an internal control. **F)** Anthocyanin contents in transgenic Arabidopsis lines. **G)** The hypocotyl length of transgenic Arabidopsis seedlings. **H)** Expression levels of anthocyanin-related genes in transgenic Arabidopsis lines. Data are presented as means ± s.d. of 3 biological replicates **B)**, **C)**, **E)**, **F)** and **H)** and 15 biological replicates **G)**. Asterisks indicate significant differences compared with wild type pear calli **B)** and **C)**, wild type Arabidopsis **E-H)**, (two-tailed Student's *t-test*, $*P < 0.05$, $**P < 0.01$), all exact P values are shown in Supplemental Data Set S5.

Figure 4. Identification of PuHB40 and its response to ROS.

A) Distribution of potential binding sites of PuHB40 in *PuMYB123-like* promoter region. Vertical red lines indicate the position of site1 to site4. **B)** The expression levels of *PuHB40* in pear seedlings under high-light stress treatment or high-light stress supplemented with DMTU treatment, and pear seedlings under appropriate light condition were used as Mock. *PuACTIN* was amplified as an internal control. **C)** Changes in the PuHB40 abundance in pear seedlings under high-light stress treatment or high-light stress supplemented with DMTU treatment for 0 h, 6 h, 24 h, 48 h. The PuHB40 protein was detected by immunoblotting with an affinity-purified specific anti-PuHB40 polyclonal antibody. **D)** Phenotype of pear calli after 4 mM H_2O_2 treatment for 2 days (scale bar, 1 cm). Pear calli treated with 0 mM H_2O_2 were used as control. **E)**

Anthocyanin contents in pear calli treated with 0 mM or 4 mM H_2O_2 for 2 days. **F)**

Expression levels of *PcHB40*, *PcMYB123-like* and anthocyanin-related genes in pear calli

treated with 0 mM or 4 mM H_2O_2 for 2 days. *PcACTIN* was amplified as an internal

control. **G)** Expression levels of *PcHB40* in pear calli after 0 mM or 4 mM H_2O_2

treatment for 0 h, 3 h, 6 h, 12 h, 24 h, 72 h. *PcACTIN* was amplified as an internal control.

H) Trans-acting activity of PuHB40 in yeast cells. **I)** Subcellular localization of PuHB40

in *N. benthamiana* leaf cells (scale bar, 25 μm). Data are presented as means ± s.d. of 3

biological replicates. Asterisks indicate significant differences compared with Mock **B)**, 4

mM H_2O_2 **E-F)** (two-tailed Student's *t-test*, *$P < 0.05$, **$P < 0.01$), all exact P values are

shown in Supplemental Data Set S5.

Figure 5. PuHB40 positively regulates anthocyanin biosynthesis by promoting

***PuMYB123-like* transcription.**

A) Effect of potential binding sites mutation on the activation of *PuMYB123-like* by

PuHB40. Four potential binding sites (site1 to site4) were mutated into TTTTTTTT,

respectively. The promoters of *PuMYB123-like* with or without potential binding site

mutation were cloned into the pGreenII 0800-LUC (firefly luciferase) vector, and the

full-length CDS of *PuHB40* was cloned into the pGreenII 0029 62-SK vector. The empty

vector of pGreenII 0029 62-SK was used as control. **B)** An EMSA confirmed that

PuHB40 binds directly to the site4 in the *PuMYB123-like* promoter. The

ProPuMYB123-like probe was the biotin-labeled fragment of *PuMYB123-like* promoter

containing site4 (AAATTATTG), whereas the cold probe was an unlabeled probe (50-,

100-, and 200-fold molar excess). *ProPuMYB123-like*-mut probe was the same as the

ProPuMYB123-like probe but with site4 (AAATTATTG) mutated to TTTTTTTT. The

His-tagged PuHB40 protein was purified. **C)** ChIP-qPCR assay results reflected the in

vivo binding of PuHB40 to the *PuMYB123-like* promoter. Cross-linked chromatin

samples were extracted from *PuHB40*-MYC-overexpressing pear calli with 3 biological

replicates and precipitated with an anti-MYC antibody. Eluted DNA was used to amplify

PuMYB123-like promoter fragments and eight regions (P1 to P8) were examined, among

which P6 containing site4. Pear calli overexpressing the MYC sequence were used as a

negative control. Three biological replicates of pear calli were used in the ChIP assay. **D)**

Transient overexpression of *PuHB40* with or without transient silencing of

PuMYB123-like in pear seedlings. The full-length CDS of *PuHB40* and reverse

complementary gene fragment (~200 to 300 bp) of *PuMYB123-like* were inserted into the

pCAMBIA1301 vector for the subsequent expression under the control of the 35S

promoter. The plasmids were transformed into *A. tumefaciens* GV3101 and infiltrated

into pear seedlings by vacuum. The infiltrated pear seedlings were firstly incubated in

darkness for 2 days and then exposed to continuous white light for 3 days. Pear seedlings

infiltrated with an empty pCAMBIA1301 vector were used as a control. **E)** Anthocyanin

content in pear seedlings. **F)** Expression levels of *PuHB40*, *PuMYB123-like* and

anthocyanin-related genes in pear seedlings. Data are presented as means ± s.d. of 3

biological replicates **C), E)**, **F)** and 6 biological replicates **A)**. Asterisks indicate

significant differences compared with empty vector **A)** and control **C), E)**, **F)** (two-tailed

Student's *t-test*, *$P < 0.05$, **$P < 0.01$; ns, no significance, $P > 0.05$), all exact P values

are shown in Supplemental Data Set S5.

Figure 6. PuHB40 promotes anthocyanin biosynthesis.

A) Phenotype of pear calli overexpressing and silencing *PuHB40* (scale bar, 1 cm). Wild

type pear calli were used as the negative control. Pear calli were incubated under

continuous white light at 17°C for 3 d. **B) a**nd **D)** Expression levels of *PuHB40*,

PpMYB123-like and anthocyanin-related genes in *PuHB40*-OE and *PuHB40*-RNAi pear

calli after light treatment. *PcACTIN* was amplified as an internal control. **C)** Anthocyanin

contents in *PuHB40*-OE and *PuHB40*-RNAi pear calli after light treatment. **E)** Phenotype

of Arabidopsis seedlings overexpressing *PuHB40* (scale bar, 2 mm). Wild-type

'Columbia 0' was used as the negative control. Arabidopsis seedlings were grown on 1/2

MS medium for 5 days. **F)** Expression levels of *PuHB40* in transgenic Arabidopsis lines.

AtPP2A was amplified as an internal control. **G)** Anthocyanin contents in transgenic

Arabidopsis lines. **H)** The hypocotyl length of transgenic Arabidopsis seedlings. **I)**

Expression levels of anthocyanin-related genes in transgenic Arabidopsis lines. Data are

presented as means ± s.d. of 3 biological replicates **B-D)**, **F), G), I)** and 15 biological

replicates **H)**. Asterisks indicate significant differences compared with wild type pear
calli **B-D)** and wild type Arabidopsis **F-I)** (two-tailed Student's *t-test*, **P* < 0.05, ***P* <
0.01), all exact *P* values are shown in Supplemental Data Set S5.

Figure 7. Identification of PuPP2AA2 and its interaction with PuHB40.
A) Phylogenetic analysis of PuPP2AA2. The phylogenetic tree was generated using the
Maximum-likelihood method of MEGA (version 10.1.8). Bootstrap values from 1000
replicates for each branch are shown. The protein sequences from pear (Pu, *Pyrus*
pyrifolia, highlighted with blue dots) and Arabidopsis (*Arabidopsis thaliana*, highlighted
with rose red dots). PuPP2AA2 (EVM0003533.1) is bolded and enlarged. **B)** Subcellular
localization of PuPP2AA2 in *N. benthamiana* leaf cells (scale bar, 25 μm). **C)** The
expression levels of *PuPP2AA2* in pear seedlings under high-light stress treatment or
high-light stress supplemented with DMTU treatment. *PuACTIN* was amplified as an
internal control. **D)** Changes in the PuPP2AA2 abundance in pear seedlings. Pear
seedlings transiently overexpressed MYC-tagged *PuPP2AA2* and were exposed to
high-light stress treatment or high-light stress supplemented with DMTU treatment for 0
h, 3 h, 6 h, 12 h. The PuPP2AA2 protein was detected by immunoblotting with anti-MYC
monoclonal antibody. **E)** The interaction between PuPP2AA2 and PuHB40 using a
pull-down assay. PuHB40-His was incubated with GST or PuPP2AA2-GST. The
interaction proteins were precipitated by GST-Trap agarose beads and analysed by
immunoblotting with an anti-GST or anti-His antibody. **F)** The interaction between
PuPP2AA2 and PuHB40 was detected by luciferase complementation imaging assays.
The pseudocolour scale bar indicates the range of luminescence intensity. **G)** Physical
association between PuPP2AA2 and PuHB40 confirmed by a BiFC assay. Fluorescence
indicates positive interactions (scale bar, 25 μm). Data are presented as means ± s.d. of 3
biological replicates. Asterisks indicate significant differences compared with high-light
stress supplemented with DMTU treatment (two-tailed Student's *t-test*, **P* < 0.05), all
exact *P* values are shown in Supplemental Data Set S5.

1006

Figure 8. PuHB40 was dephosphorylated by PP2A phosphatase.

A) Phos-tag analysis of phosphorylation of PuHB40 after cantharidin treatment. *A. tumefaciens* GV3101 containing PuHB40-GFP plasmid infiltrated into *N. benthamiana* leaves by injection and 50 μM cantharidin was injected into the same position as the agrobacteria after 24 h. *N. benthamiana* leaves injected corresponding concentration of DMSO were used as negative control. Protein samples were extracted from *N. benthamiana* leaves after 24 h and were separated in a Phos-tag gel (upper panel), and PuHB40-GFP with different phosphorylated status was detected using the anti-GFP antibody. Total PuHB40-GFP proteins were analyzed by SDS-PAGE without Phos-tag (indicated as "Regular"), and were detected using anti-GFP antibody. P+ and P- indicate phosphorylated and non-phosphorylated form of PuHB40, respectively. The intensity of the bands was measured with the Image J software and was calculated as the ratio of the P+/P- form. **B)** Immunoblotting analysis showing that the PP2A complex directly dephosphorylates PuHB40 *in vitro*. MYC, PP2A complex, and PuHB40-GFP were immunopurified from pear seedlings transiently overexpressing MYC, PuPP2AA2-MYC, and PuHB40-GFP using MYC-Trap or GFP-Trap, respectively. After incubation of PP2A complexes with PuHB40-GFP (MYC with PuHB40-GFP was set as control), protein samples in the reaction system were separated in Phos-tag gel (upper panel) and detected using the anti-PuHB40 antibody. The intensity of the bands was measured with the Image J software and calculated as the ratio of the P+/P-. Total proteins in the reaction system were analyzed by SDS-PAGE without Phos-tag (indicated as "Regular"). MYC, PuPP2AA2-MYC, and PuHB40-GFP were set as the loading controls and detected by the anti-MYC or anti-PuHB40 antibody, respectively. **C)** Subcellular localization of PuPP2AA2D in *N. benthamiana* leaf cells (scale bar, 25 μm). PuPP2AA2D is PuPP2AA2 with heat repeats 14 and 15 in the C-terminal deleted. **D)** The interaction between PuPP2AA2D and PuHB40 confirmed by a BiFC assay. Fluorescence indicates positive interactions (scale bar, 25 μm). **E)** The interaction between PuPP2AA2D and PuHB40 was detected by luciferase complementation imaging assays. The pseudocolour scale bar indicates the range of luminescence intensity. **F)** Immunoblotting analysis showing that PP2A dephosphorylates PuHB40 *in vivo*. Protein extracts from *N. benthamiana* leaves

co-infiltrated PuHB40-GFP with MYC, PuPP2AA2-MYC, or PuPP2AA2D-MYC were separated in Phos-tag gel (upper panel) and detected using the anti-GFP antibody (MYC with PuHB40-GFP was set as control). The intensity of the bands was measured with the Image J software and calculated the ratio of P+/P−. Total PuHB40-GFP, MYC, PuPP2AA2-MYC and PuPP2AA2D-MYC proteins were analyzed by SDS-PAGE without Phos-tag (indicated as "Regular"). PuHB40-GFP, MYC, PuPP2AA2-MYC and PuPP2AA2D-MYC were set as the loading controls and detected by the anti-GFP or anti-MYC antibody, respectively. **G)** Dual-luciferase assay showed that dephosphorylation of PuHB40 by PP2A significantly inhibited the activation of *PuMYB123-like* promoter by PuHB40. The *PuMYB123-like* promoter was cloned into the pGreenII 0800-LUC (firefly luciferase) vector, and the full-length CDS of *PuHB40*, PuPP2AA2 and PuPP2AA2D was cloned into the pGreenII 0029 62-SK vector. The PuHB40 was used as control. Data are presented as means ± s.d. of 6 biological replicates **G)**. Asterisks indicate significant differences compared PuHB40 **G)** (two-tailed Student's *t-test*, **$P < 0.01$; ns, no significance, $P > 0.05$), all exact P values are shown in Supplemental Data Set S5.

Figure 9. Attenuation of PuHB40-induced anthocyanin biosynthesis by PuPP2AA2.
A) Phenotype of pear calli overexpressing *PuPP2AA2* (scale bar, 1 cm). Wild type pear calli were used as the negative control. Pear calli were incubated under continuous white light at 17°C for 3 d. **B)** Anthocyanin contents in *PuPP2AA2*-OE after light treatment. **C)** Expression levels of *PuPP2AA2* and anthocyanin-related genes in *PuPP2AA2*-OE pear calli after light treatment. *PcACTIN* was amplified as an internal control. **D)** Transient overexpression of *PuHB40* with or without transient overexpressing of *PuPP2AA2* in pear seedlings. The full-length CDS of *PuHB40* and *PuPP2AA2* were inserted into the pCAMBIA1301 vector for the subsequent expression under the control of the 35S promoter. The plasmids were transformed into *A. tumefaciens* GV3101 and infiltrated into pear seedlings by vacuum. The infiltrated pear seedlings were firstly incubated in darkness for 2 days and then exposed to continuous white light for 3 days. Pear seedlings infiltrated with an empty pCAMBIA1301 vector were used as a control. **E)** Anthocyanin

content in pear seedlings. **F)** Expression levels of *PuHB40*, *PuPP2AA2*, *PuMYB123-like*

and anthocyanin-related genes in pear seedlings. **G)** Transient overexpression of *PuHB40*

with or without transient overexpressing of *PuPP2AA2* in fruit (scale bar, 1 cm). The

full-length CDS of *PuHB40* and *PuPP2AA2* were cloned to the pGreenII 0029 62-SK

vector for the subsequent expression under the control of the 35S promoter. Pear fruits

were infiltrated with *A. tumefaciens* GV3101 cells containing the recombinant plasmid

using a needle-free syringe. Fruits infiltrated with an empty pGreenII 0029 62-SK vector

were used as a control. The phenotypes were examined after a 1 day dark and 4 days light

treatment. **H)** Total anthocyanin content in pear fruits. **I)** Expression levels of *PuHB40*,

PuPP2AA2 and anthocyanin-related genes in pear fruits. Data are presented as means ±

s.d. of 3 biological replicates. Asterisks indicate significant differences compared with

wild type pear calli **B)** and **C)**, empty vector **E), F), H), I)** (two-tailed Student's *t-test*, *P

< 0.05, **P < 0.01; ns, no significance, $P > 0.05$), all exact P values are shown in

Supplemental Data Set S5.

1081

Figure 10. Proposed model for PuHB40-mediated ROS-dependent anthocyanin

biosynthesis in pears.

PuHB40 is the key transcription factor in ROS-induced anthocyanin biosynthesis. ROS

promote PuHB40 expression and then positively regulate its transcriptional activity on

PuMYB123-like/PubHLH3 complex to induce anthocyanin biosynthesis. The

transcriptional activity of PuHB40 depends on its phosphorylation level which is

regulated by PP2A. ROS maintain phosphorylation of PuHB40 at a high level, and

enhance the transcriptional activity of PuHB40 on *PuMYB123-like* by reducing the

transcriptional and translational levels of PuPP2AA2 under high-light stress, and thus

promote anthocyanin biosynthesis. The lightning indicates the intensity of transcriptional

regulation of *PuMYB123-like* by PuHB40.

1093

1094   **REFERENCES**

1095   **Acero N, Gradillas A, Beltran M, Garcia A, Munoz Mingarro D.** Comparison of

phenolic compounds profile and antioxidant properties of different sweet cherry

(*Prunus* *avium* L.) varieties. Food Chem. 2019:**279**:260-271.

https://doi.org/10.1016/j.foodchem.2018.12.008

Altangerel N, Ariunbold GO, Gorman C, Alkahtani MH, Borrego EJ, Bohlmeyer D,

Hemmer P, Kolomiets MV, Yuan JS, Scully MO. In vivo diagnostics of early abiotic

plant stress response via Raman spectroscopy. Proc Natl Acad Sci U S A.

2017:**114**(13):3393-3396. https://doi.org/10.1073/pnas.1701328114

An JP, Wang XF, Zhang XW, Xu HF, Bi SQ, You CX, Hao YJ. An apple MYB

transcription factor regulates cold tolerance and anthocyanin accumulation and

undergoes MIEL1-mediated degradation. Plant Biotechnol J. 2020:**18**(2):337-353.

https://doi.org/10.1111/pbi.13201

An XH, Tian Y, Chen KQ, Liu XJ, Liu DD, Xie XB, Cheng CG, Cong PH, Hao YJ.

MdMYB9 and MdMYB11 are involved in the regulation of the JA-induced

biosynthesis of anthocyanin and proanthocyanidin in apples. Plant Cell Physiol.

2015:**56**(4):650-662. https://doi.org/10.1093/pcp/pcu205

Ariel FD, Manavella PA, Dezar CA, Chan RL. The true story of the HD-Zip family.

Trends Plant Sci. 2007:**12**(9):419-426. https://doi.org/10.1016/j.tplants.2007.08.003

Bai S, Tao R, Yin L, Ni J, Yang Q, Yan X, Yang F, Guo X, Li H, Teng Y. Two B-box

proteins, PpBBX18 and PpBBX21, antagonistically regulate anthocyanin biosynthesis

via competitive association with *Pyrus pyrifolia* ELONGATED HYPOCOTYL 5 in the

peel of pear fruit. Plant J. 2019a:**100**(6):1208-1223. https://doi.org/10.1111/tpj.14510

Bai S, Tao R, Tang Y, Yin L, Ma Y, Ni J, Yan X, Yang Q, Wu Z, Zeng Y, et al. BBX16, a B-box protein, positively regulates light-induced anthocyanin accumulation by activating *MYB10* in red pear. Plant Biotechnol J. 2019b:**17**(10):1985-1997. https://doi.org/10.1111/pbi.13114

Bi X, Zhang J, Chen C, Zhang D, Li P, Ma F. Anthocyanin contributes more to hydrogen peroxide scavenging than other phenolics in apple peel. Food Chem. 2014:**152**:205-209. https://doi.org/10.1016/j.foodchem.2013.11.088

Bian C, Guo X, Zhang Y, Wang L, Xu T, DeLong A, Dong J. Protein phosphatase 2A promotes stomatal development by stabilizing SPEECHLESS in *Arabidopsis*. Proc Natl Acad Sci U S A. 2020:**117**(23):13127-13137. https://doi.org/10.1073/pnas.1912075117

Byrne DP, Shrestha S, Galler M, Cao M, Daly LA, Campbell AE, Eyers CE, Veal EA, Kannan N, Eyers PA. Aurora A regulation by reversible cysteine oxidation reveals evolutionarily conserved redox control of Ser/Thr protein kinase activity. Sci Signal. 2020:**13**(613). https://doi.org/10.1126/scisignal.aax2713

Catala R, Medina J, Salinas J. Integration of low temperature and light signaling during cold acclimation response in *Arabidopsis*. Proc Natl Acad Sci U S A. 2011:**108**(39):16475-16480. https://doi.org/10.1073/pnas.1107161108

Chan KX, Mabbitt PD, Phua SY, Mueller JW, Nisar N, Gigolashvili T, Stroeher E,

Grassl J, Arlt W, Estavillo GM, et al. Sensing and signaling of oxidative stress in

chloroplasts by inactivation of the SAL1 phosphoadenosine phosphatase. Proc Natl

Acad Sci U S A. 2016:**113**(31):E4567-4576. https://doi.org/10.1073/pnas.1604936113

Choi S, Kwon YR, Hossain MA, Hong S-W, Lee B-h, Lee H. A mutation in *ELA1*, an

age-dependent negative regulator of *PAP1/MYB75*, causes UV- and cold

stress-tolerance in *Arabidopsis thaliana* seedlings. Plant Sci. 2009:**176**(5):678-686.

https://doi.org/10.1016/j.plantsci.2009.02.010

Croce R, van Amerongen H. Natural strategies for photosynthetic light harvesting. Nat

Chem Biol. 2014:**10**(7):492-501. https://doi.org/10.1038/nchembio.1555

Das PK, Shin DH, Choi SB, Yoo SD, Choi G, Park YI. Cytokinins enhance

sugar-induced anthocyanin biosynthesis in Arabidopsis. Mol Cells. 2012:**34**(1):93-101.

https://doi.org/10.1007/s10059-012-0114-2

DeLong A. Switching the flip: protein phosphatase roles in signaling pathways. Curr

Opin Plant Biol. 2006:**9**(5):470-477. https://doi.org/10.1016/j.pbi.2006.07.015

Dubos C, Stracke R, Grotewold E, Weisshaar B, Martin C, Lepiniec L. MYB

transcription factors in *Arabidopsis*. Trends Plant Sci. 2010:**15**(10):573-581.

https://doi.org/10.1016/j.tplants.2010.06.005

Durian G, Rahikainen M, Alegre S, Brosche M, Kangasjarvi S. Protein Phosphatase

2A in the Regulatory Network Underlying Biotic Stress Resistance in Plants. Front

Plant Sci. 2016:**7**:812. https://doi.org/10.3389/fpls.2016.00812

Espley RV, Hellens RP, Putterill J, Stevenson DE, Kutty-Amma S, Allan AC. Red

colouration in apple fruit is due to the activity of the MYB transcription factor,

MdMYB10. Plant J. 2007:**49**(3):414-427.

https://doi.org/10.1111/j.1365-313X.2006.02964.x

Farkas I, Dombradi V, Miskei M, Szabados L, Koncz C. *Arabidopsis* PPP family of

serine/threonine phosphatases. Trends Plant Sci. 2007:**12**(4):169-176.

https://doi.org/10.1016/j.tplants.2007.03.003

Feild TS, Lee DW, Holbrook NM. Why Leaves Turn Red in Autumn. The Role of

Anthocyanins in Senescing Leaves of Red-Osier Dogwood. Plant Physiol.

2001:**127**(2):566-574. https://doi.org/10.1104/pp.010063

Feng S, Wang Y, Yang S, Xu Y, Chen X. Anthocyanin biosynthesis in pears is regulated

by a R2R3-MYB transcription factor PyMYB10. Planta. 2010:**232**(1):245-255.

https://doi.org/10.1007/s00425-010-1170-5

Galvez-Valdivieso G, Fryer MJ, Lawson T, Slattery K, Truman W, Smirnoff N,

Asami T, Davies WJ, Jones AM, Baker NR, et al. The high light response in

Arabidopsis involves ABA signaling between vascular and bundle sheath cells. Plant

Cell. 2009:**21**(7):2143-2162. https://doi.org/10.1105/tpc.108.061507

Gould KS, McKelvie J, Markham KR. Do anthocyanins function as antioxidants in

leaves? Imaging of H_2O_2 in red and green leaves after mechanical injury. Plant Cell

Environ. 2002:**25**(10):1261-1269. https://doi.org/10.1046/j.1365-3040.2002.00905.x

Groves MR, Hanlon N, Turowski P, Hemmings BA, Barford D. The structure of the

protein phosphatase 2A PR65/A subunit reveals the conformation of its 15 tandemly

repeated HEAT motifs. Cell. 1999:**96**(1):99-110.

https://doi.org/10.1016/S0092-8674(00)80963-0

Gupta R, Luan S. Redox control of protein tyrosine phosphatases and mitogen-activated

protein kinases in plants. Plant Physiol. 2003:**132**(3):1149-1152.

https://doi.org/10.1104/pp.103.020792

Harris JC, Hrmova M, Lopato S, Langridge P. Modulation of plant growth by HD-Zip

class I and II transcription factors in response to environmental stimuli. New Phytol.

2011:**190**(4):823-837. https://doi.org/10.1111/j.1469-8137.2011.03733.x

Himmelbach A, Hoffmann T, Leube M, Hohener B, Grill E. Homeodomain protein

ATHB6 is a target of the protein phosphatase ABI1 and regulates hormone responses in

Arabidopsis. Embo J. 2002:**21**(12):3029-3038. https://doi.org/10.1093/emboj/cdf316

Hipsch M, Lampl N, Zelinger E, Barda O, Waiger D, Rosenwasser S. Sensing stress

responses in potato with whole-plant redox imaging. Plant Physiol.

2021:**187**(2):618-631. https://doi.org/10.1093/plphys/kiab159

Hoch WA, Zeldin EL, McCown BH. Physiological significance of anthocyanins during

autumnal leaf senescence. Tree Physiol. 2001:**21**(1):1-8.

https://doi.org/10.1093/treephys/21.1.1

Honkanen RE. Cantharidin, another natural toxin that inhibits the activity of

serine/threonine protein phosphatases types 1 and 2A. FEBS Lett. 1993:**330**(3):283-286. https://doi.org/10.1016/0014-5793(93)80889-3

Jaakola L. New insights into the regulation of anthocyanin biosynthesis in fruits. Trends Plant Sci. 2013:**18**(9):477-483. https://doi.org/10.1016/j.tplants.2013.06.003

Jalmi SK, Sinha AK. ROS mediated MAPK signaling in abiotic and biotic stress-striking similarities and differences. Front Plant Sci. 2015:**6**:769. https://doi.org/10.3389/fpls.2015.00769

Janssens V, Goris J. Protein phosphatase 2A: a highly regulated family of serine/threonine phosphatases implicated in cell growth and signalling. Biochem J. 2001:**353**(3):417-439. https://doi.org/10.1042/0264-6021:3530417

Jia M, Li X, Wang W, Li T, Dai Z, Chen Y, Zhang K, Zhu H, Mao W, Feng Q, et al. SnRK2 subfamily I protein kinases regulate ethylene biosynthesis by phosphorylating HB transcription factors to induce *ACO1* expression in apple. New Phytol. 2022:**234**(4):1262-1277. https://doi.org/10.1111/nph.18040

Jiang T, He Y, Wu Z, Cui Y, Wang X, Huang H, Fan Y, Han M, Wang J, Wang S, et al. Enhancing stimulation of cyaniding, *GhLDOX3* activates reactive oxygen species to regulate tolerance of alkalinity negatively in cotton. Ecotox Environ Safe. 2023:**267**:115655. https://doi.org/10.1016/j.ecoenv.2023.115655

Kamibayashi C, Lickteig RL, Estes R, Walter G, Mumby MC. Expression of the A subunit of protein phosphatase 2A and characterization of its interactions with the

catalytic and regulatory subunits. J Biol Chem. 1992:**267**(30):21864-21872.

https://doi.org/10.1016/s0021-9258(19)36692-x

Kerchev P, Waszczak C, Lewandowska A, Willems P, Shapiguzov A, Li Z, Alseekh S,

Muhlenbock P, Hoeberichts FA, Huang J, et al. Lack of GLYCOLATE OXIDASE1,

but Not GLYCOLATE OXIDASE2, Attenuates the Photorespiratory Phenotype of

CATALASE2-Deficient Arabidopsis. Plant Physiol. 2016:**171**(3):1704-1719.

https://doi.org/10.1104/pp.16.00359

Koes R, Verweij W, Quattrocchio F. Flavonoids: a colorful model for the regulation and

evolution of biochemical pathways. Trends Plant Sci. 2005:**10**(5):236-242.

https://doi.org/10.1016/j.tplants.2005.03.002

Kovtun Y, Chiu WL, Tena G, Sheen J. Functional analysis of oxidative stress-activated

mitogen-activated protein kinase cascade in plants. Proc Natl Acad Sci U S A.

2000:**97**(6):2940-2945. https://doi.org/10.1073/pnas.97.6.2940

Landi M, Tattini M, Gould KS. Multiple functional roles of anthocyanins in

plant-environment interactions. Environ Exp Bot. 2015:**119**:4-17.

https://doi.org/10.1016/j.envexpbot.2015.05.012

Lee DW, Gould KS. Why leaves turn red: pigments called anthocyanins probably protect

leaves from light damage by direct shielding and by scavenging free radicals. AM SCI.

2002:**90**(6):524-531. https://doi.org/10.1511/2002.6.524

Lesk C, Rowhani P, Ramankutty N. Influence of extreme weather disasters on global

crop production. Nature. 2016:**529**(7584):84-87. https://doi.org/10.1038/nature16467

Li Y, Wang Y, Tan S, Li Z, Yuan Z, Glanc M, Domjan D, Wang K, Xuan W, Guo Y,

et al. Root Growth Adaptation is Mediated by PYLs ABA Receptor-PP2A Protein

Phosphatase Complex. Adv Sci. 2020:**7**(3):1901455.

https://doi.org/10.1002/advs.201901455

Li YY, Mao K, Zhao C, Zhao XY, Zhang HL, Shu HR, Hao YJ. MdCOP1 ubiquitin

E3 ligases interact with MdMYB1 to regulate light-induced anthocyanin biosynthesis

and red fruit coloration in apple. Plant Physiol. 2012:**160**(2):1011-1022.

https://doi.org/10.1104/pp.112.199703

Lillo C, Kataya AR, Heidari B, Creighton MT, Nemie-Feyissa D, Ginbot Z, Jonassen

EM. Protein phosphatases PP2A, PP4 and PP6: mediators and regulators in

development and responses to environmental cues. Plant Cell Environ.

2014:**37**(12):2631-2648. https://doi.org/10.1111/pce.12364

Lin-Wang K, Micheletti D, Palmer J, Volz R, Lozano L, Espley R, Hellens RP,

Chagne D, Rowan DD, Troggio M, et al. High temperature reduces apple fruit colour

via modulation of the anthocyanin regulatory complex. Plant Cell Environ.

2011:**34**(7):1176-1190. https://doi.org/10.1111/j.1365-3040.2011.02316.x

Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time

quantitative PCR and the $2^{-\triangle\triangle C_{T}}$ method. Methods. 2001:**25**(4):402-408.

https://doi.org/10.1006/meth.2001.1262

Mine A, Berens ML, Nobori T, Anver S, Fukumoto K, Winkelmuller TM, Takeda A,

Becker D, Tsuda K. Pathogen exploitation of an abscisic acid- and

jasmonate-inducible MAPK phosphatase and its interception by *Arabidopsis* immunity.

Proc Natl Acad Sci U S A. 2017:**114**(28):7456-7461.

https://doi.org/10.1073/pnas.1702613114

Mittler R. ROS Are Good. Trends Plant Sci. 2017:**22**(1):11-19.

https://doi.org/10.1016/j.tplants.2016.08.002

Mittler R, Zandalinas SI, Fichman Y, Van Breusegem F. Reactive oxygen species

signalling in plant stress responses. Nat Rev Mol Cell Biol. 2022:**23**(10):663-679.

https://doi.org/10.1038/s41580-022-00499-2

Nakabayashi R, Yonekura-Sakakibara K, Urano K, Suzuki M, Yamada Y,

Nishizawa T, Matsuda F, Kojima M, Sakakibara H, Shinozaki K, et al.

Enhancement of oxidative and drought tolerance in Arabidopsis by overaccumulation

of antioxidant flavonoids. Plant J. 2014:**77**(3):367-379.

https://doi.org/10.1111/tpj.12388

Neill SO, Gould KS, Kilmartin PA, Mitchell KA, Markham KR. Antioxidant

activities of red versus green leaves in *Elatostema rugosum*. Plant Cell Environ.

2002:**25**(4):539-547. https://doi.org/10.1046/j.1365-3040.2002.00837.x

Ni J, Wang S, Yu W, Liao Y, Pan C, Zhang M, Tao R, Wei J, Gao Y, Wang D, et al.

The ethylene-responsive transcription factor PpERF9 represses *PpRAP2.4* and

PpMYB114 via histone deacetylation to inhibit anthocyanin biosynthesis in pear. Plant

Cell. 2023:**35**(6):2271-2292. https://doi.org/10.1093/plcell/koad077

Olsson AS, Engström P, Söderman E. The homeobox genes *ATHB12* and *ATHB7*

encode potential regulators of growth in response to water deficit in *Arabidopsis*. Plant

Mol Biol. 2004:**55**(5):663-677. https://doi.org/10.1007/s11103-004-1581-4

Qu Y, Bai X, Zhu YJ, Qi R, Tian G, Wang Y, Li YH, Zhang KM. Reactive Oxygen

Species Acts as an Important Inducer in Low-temperature-induced Anthocyanin

Biosynthesis in *Begonia semperflorens*. J Am Soc Hortic Sci. 2018:**143**(6):486-493.

https://doi.org/10.21273/JASHS04488-18

Ren H, Rao J, Tang M, Li Y, Dang X, Lin D. PP2A interacts with KATANIN to

promote microtubule organization and conical cell morphogenesis. J Integr Plant Biol.

2022:**64**(8):1514-1530. https://doi.org/10.1111/jipb.13281

Ruediger R, Hentz M, Fait J, Mumby M, Walter G. Molecular model of the A subunit

of protein phosphatase 2A interaction with other subunits and tumor antigens. J Virol.

1994:**68**(1):123-129. https://doi.org/10.1128/JVI.68.1.123-129.1994

Schaart JG, Dubos C, De La Fuente IR, van Houwelingen AMML, de Vos RCH,

Jonker HH, Xu WJ, Routaboul JM, Lepiniec L, Bovy AG. Identification and

characterization of MYB‐bHLH‐WD40 regulatory complexes controlling

proanthocyanidin biosynthesis in strawberry (*Fragaria × ananassa*) fruits. New

Phytol. 2012:**197**(2):454-467. https://doi.org/10.1111/nph.12017

Shan X, Zhang Y, Peng W, Wang Z, Xie D. Molecular mechanism for

jasmonate-induction of anthocyanin accumulation in *Arabidopsis*. J Exp Bot.

2009:**60**(13):3849-3860. https://doi.org/10.1093/jxb/erp223

Shi H, Liu G, Wei Y, Chan Z. The zinc-finger transcription factor ZAT6 is essential for

hydrogen peroxide induction of anthocyanin synthesis in *Arabidopsis*. Plant Mol Biol.

2018:**97**(1-2):165-176. https://doi.org/10.1007/s11103-018-0730-0

Soderman E, Mattsson J, Engstrom P. The *Arabidopsis* homeobox gene *ATHB-7* is

induced by water deficit and by abscisic acid. Plant J. 1996:**10**(2):375-381.

https://doi.org/10.1046/j.1365-313x.1996.10020375.x

Söderman E, Hjellström M, Fahleson J, Engström P. The HD-Zip gene *ATHB6* in

Arabidopsis is expressed in developing leaves, roots and carpels and up-regulated by

water   deficit   conditions.   Plant   Mol   Biol.   1999:**40**(6):1073-1083.

https://doi.org/10.1023/A:1006267013170

Steyn WJ, Wand SJE, Holcroft DM, Jacobs G. Anthocyanins in vegetative tissues: a

proposed unified function in photoprotection. New Phytol. 2002:**155**(3):349-361.

https://doi.org/10.1046/j.1469-8137.2002.00482.x

Takos AM, Jaffe FW, Jacob SR, Bogs J, Robinson SP, Walker AR. Light-induced

expression of a *MYB* gene regulates anthocyanin biosynthesis in red apples. Plant

Physiol. 2006:**142**(3):1216-1232. https://doi.org/10.1104/pp.106.088104

Uhrig RG, Labandera AM, Moorhead GB. *Arabidopsis* PPP family of serine/threonine

protein phosphatases: many targets but few engines. Trends Plant Sci.

2013:**18**(9):505-513. https://doi.org/10.1016/j.tplants.2013.05.004

Wang L, Leister D, Guan L, Zheng Y, Schneider K, Lehmann M, Apel K, Kleine T.

The *Arabidopsis* SAFEGUARD1 suppresses singlet oxygen-induced stress responses

by protecting grana margins. Proc Natl Acad Sci U S A. 2020:**117**(12):6918-6927.

https://doi.org/10.1073/pnas.1918640117

Winkel-Shirley B. Flavonoid biosynthesis. A colorful model for genetics, biochemistry,

cell biology, and biotechnology. Plant Physiol. 2001:**126**(2):485-493.

https://doi.org/10.1104/pp.126.2.485

Xie XB, Li S, Zhang RF, Zhao J, Chen YC, Zhao Q, Yao YX, You CX, Zhang XS,

Hao YJ. The bHLH transcription factor MdbHLH3 promotes anthocyanin

accumulation and fruit colouration in response to low temperature in apples. Plant Cell

Environ. 2012:**35**(11):1884-1897. https://doi.org/10.1111/j.1365-3040.2012.02523.x

Xu H, Zou Q, Yang G, Jiang S, Fang H, Wang Y, Zhang J, Zhang Z, Wang N, Chen

X. MdMYB6 regulates anthocyanin formation in apple both through direct inhibition

of the biosynthesis pathway and through substrate removal. Hortic Res. 2020:**7**(1):72.

https://doi.org/10.1038/s41438-020-0294-4

Xu W, Dubos C, Lepiniec L. Transcriptional control of flavonoid biosynthesis by

MYB-bHLH-WDR complexes. Trends Plant Sci. 2015:**20**(3):176-185.

https://doi.org/10.1016/j.tplants.2014.12.001

Xu Z, Mahmood K, Rothstein SJ. ROS Induces Anthocyanin Production Via Late

Biosynthetic Genes and Anthocyanin Deficiency Confers the Hypersensitivity to

ROS-Generating Stresses in Arabidopsis. Plant Cell Physiol. 2017:**58**(8):1364-1377.

https://doi.org/10.1093/pcp/pcx073

Zhang D, Yu B, Bai J, Qian M, Shu Q, Su J, Teng Y. Effects of high temperatures on

UV-B/visible irradiation induced postharvest anthocyanin accumulation in 'Yunhongli

No. 1' (*Pyrus pyrifolia* Nakai) pears. Sci Hortic. 2012:**134**:53-59.

https://doi.org/10.1016/j.scienta.2011.10.025

Zhang L, Tao R, Wang S, Gao Y, Wang L, Yang S, Zhang X, Yu W, Wu X, Li K, et al.

PpZAT5 suppresses the expression of a B-box gene *PpBBX18* to inhibit anthocyanin

biosynthesis in the fruit peel of red pear. Front Plant Sci. 2022:**13**:1022034.

https://doi.org/10.3389/fpls.2022.1022034

Zhou HW, Nussbaumer C, Chao Y, DeLong A. Disparate roles for the regulatory A

subunit isoforms in Arabidopsis protein phosphatase 2A. Plant Cell.

2004:**16**(3):709-722. https://doi.org/10.1105/tpc.018994

1351

1352

Figure 1. High light stress-induced anthocyanin biosynthesis is ROS-dependent. A-K), Pear seedlings were treated with high light stress treatment (photon flux density: approximately 160 µmol·s^{-1}·m^{-2}) or high light stress supplemented with N, N'-dimethylthiourea (DMTU) treatment. DMTU was applied as ROS scavenger, and pear seedlings under appropriate light condition were used as Mock (photon flux density: approximately 50 µmol·s^{-1}·m^{-2}). Pear seedlings were sampled at 0 h, 6 h, 12 h, 24 h, 48 h, 72 h, 96 h. **A)**, Phenotype of pear seedlings (scale bar, 1 cm). **B)**, The anthocyanin content for pear seedlings. **C and D)**, NBT staining for superoxide and DAB staining for H$_2$O$_2$ in pear seedlings leaves, respectively (scale bar, 0.5 mm). **E)**, H$_2$O$_2$ quantification of pear seedlings. **F-K)**, The expression levels of *PuCHI*, *PuCHS*, *PuDFR*, *PuF3H*, *PuANS*, *PuMYB10* in pear seedlings under different treatments. *PuACTIN* was amplified as an internal control. Data are presented as means ± s.d. of 3 biological replicates. Asterisks indicate significant differences compared with Mock group (two-tailed Student's t-test, *P < 0.05, **P < 0.01), all exact P values are shown in Supplemental Data Set S5.

Figure 2. Identification of PuMYB123-like and its interaction with PubHLH3. A) The expression levels of *PuMYB123-like* in pear seedlings under high-light stress treatment or high-light stress supplemented with DMTU treatment, and pear seedlings under appropriate light condition were used as Mock. *PuACTIN* was amplified as an internal control. **B)** Phylogenetic analysis of PuMYB123-like. The phylogenetic tree was generated using the Maximum-likelihood method of MEGA (version 10.1.8). Bootstrap values from 1000 replicates for each branch are shown. The protein sequences from pear (Pu, *Pyrus pyrifolia*, highlighted with blue dots), apple (Md, *Malus × domestica*, highlighted with green dots), Arabidopsis (At, *Arabidopsis thaliana*, highlighted with rose red dots) and strawberry (Fa, *Fragaria × ananassa*, highlighted with purple dots). PuMYB123-like (EVM0034726.1) is bolded and enlarged. **C)** Subcellular localization of PuMYB123-like in *N. benthamiana* leaf cells (scale bar, 25 μm). **D)** Trans-acting activity of PuMYB123-like in yeast cells. **E)** Interaction between PuMYB123-like and PubHLH3 in a yeast two-hybrid assay, with pGADT7-T and pGBKT7-53 as positive controls. DDO, SD/-Trp/-Leu medium; QDO/X/A, SD/-Trp/-Leu/-Ade/-His medium with X-α-gal and Aureobasidin. **F)** The interaction between PuMYB123-like and PubHLH3 was detected by luciferase complementation imaging assays. The pseudocolour scale bar indicates the range of luminescence intensity. **G)** The interaction between PuMYB123-like and PubHLH3 analyzed using a pull-down assay. PuMYB123-like-His was incubated with GST or PubHLH3-GST. The interaction proteins were precipitated by GST-Trap agarose beads and analysed by immunoblotting with an anti-GST or anti-His antibody. **H)** A dual-luciferase assay demonstrated that PuMYB123-like promotes the promoter activity of anthocyanin-related genes (*PuCHS, PuANS, PuDFR, PuUFGT*). The promoter of anthocyanin-related genes was cloned into the pGreenII 0800-LUC (firefly luciferase) vector, and the full-length CDS of *PuMYB123-like* was cloned into the pGreenII 0029 62-SK vector. The empty vector of pGreenII 0029 62-SK was used as control. **I)** Dual-luciferase assay showed that interaction between PuMYB123-like and PubHLH3 significantly enhanced activation of PuMYB123-like on *PuCHS* and *PuANS* promoters. Data are presented as means ± s.d. of 3 biological replicates **A)** and 6 biological replicates **H) a**nd **I)**. Asterisks indicate significant differences compared with Mock **A)**, empty vector **H)** and PuMYB123-like **I)** (two-tailed Student's t-test, *P < 0.05, **P < 0.01), all exact P values are shown in Supplemental Data Set S5.

Figure 3. PuMYB123-like promotes anthocyanin biosynthesis. A) Phenotype of pear calli overexpressing and silencing *PuMYB123-like* (scale bar, 1 cm). Wild type pear calli were used as the negative control. Pear calli were incubated under continuous white light at 17°C for 3 d. **B)** Expression levels of anthocyanin-related genes and *PuMYB123-like* in *PuMYB123-like*-OE *and PuMYB123-like*-RNAi pear calli after light treatment. *PcACTIN* was amplified as an internal control. **C)** Anthocyanin contents in *PuMYB123-like*-OE and *PuMYB123-like*-RNAi pear calli after light treatment. **D)** Phenotype of *Arabidopsis* seedlings overexpressing *PuMYB123-like* (scale bar, 2 mm). Wild-type 'Columbia 0' was used as the negative control. *Arabidopsis* seedlings were grown on 1/2 MS medium for 5 days. **E)** Expression levels of *PuMYB123-like* in transgenic *Arabidopsis* lines. *AtPP2A* was amplified as an internal control. **F)** Anthocyanin contents in transgenic *Arabidopsis* lines. **G)** The hypocotyl length of transgenic *Arabidopsis* seedlings. **H)** Expression levels of anthocyanin-related genes in transgenic *Arabidopsis* lines. Data are presented as means ± s.d. of 3 biological replicates **B)**, **C)**, **E)**, **F)** and **H)** and 15 biological replicates **G)**. Asterisks indicate significant differences compared with wild type pear calli **B)** and **C)**, wild type *Arabidopsis* **E-H)**, (two-tailed Student's *t*-test, $*P < 0.05$, $**P < 0.01$), all exact *P* values are shown in Supplemental Data Set S5.

Figure 4. Identification of PuHB40 and its response to ROS. **A)** Distribution of potential binding sites of PuHB40 in *PuMYB123-like* promoter region. Vertical red lines indicate the position of site1 to site4. **B)** The expression levels of *PuHB40* in pear seedlings under high-light stress treatment or high-light stress supplemented with DMTU treatment, and pear seedlings under appropriate light condition were used as Mock. *PuACTIN* was amplified as an internal control. **C)** Changes in the PuHB40 abundance in pear seedlings under high-light stress treatment or high-light stress supplemented with DMTU treatment for 0 h, 6 h, 24 h, 48 h. The PuHB40 protein was detected by immunoblotting with an affinity-purified specific anti-PuHB40 polyclonal antibody. **D)** Phenotype of pear calli after 4 mM H_2O_2 treatment for 2 days (scale bar, 1 cm). Pear calli treated with 0 mM H_2O_2 were used as control. **E)** Anthocyanin contents in pear calli treated with 0 mM or 4 mM H_2O_2 for 2 days. **F)** Expression levels of *PcHB40*, *PcMYB123-like* and anthocyanin-related genes in pear calli treated with 0 mM or 4 mM H_2O_2 for 2 days. *PcACTIN* was amplified as an internal control. **G)** Expression levels of *PcHB40* in pear calli after 0 mM or 4 mM H_2O_2 treatment for 0 h, 3 h, 6 h, 12 h, 24 h, 72 h. *PcACTIN* was amplified as an internal control. **H)** Trans-acting activity of PuHB40 in yeast cells. **I)** Subcellular localization of PuHB40 in *N. benthamiana* leaf cells (scale bar, 25 μm). Data are presented as means ± s.d. of 3 biological replicates. Asterisks indicate significant differences compared with Mock **B)**, 4 mM H_2O_2 **E-F)** (two-tailed Student's *t*-test, *$P < 0.05$, **$P < 0.01$), all exact *P* values are shown in Supplemental Data Set S5.

Figure 5. PuHB40 positively regulates anthocyanin biosynthesis by promoting *PuMYB123-like* **transcription. A)** Effect of potential binding sites mutation on the activation of *PuMYB123-like* by PuHB40. Four potential binding sites (site1 to site4) were mutated into TTTTTTTTT, respectively. The promoters of *PuMYB123-like* with or without potential binding site mutation were cloned into the pGreenII 0800-LUC (firefly luciferase) vector, and the full-length CDS of *PuHB40* was cloned into the pGreenII 0029 62-SK vector. The empty vector of pGreenII 0029 62-SK was used as control. **B)** An EMSA confirmed that PuHB40 binds directly to the site4 in the *PuMYB123-like* promoter. The *ProPuMYB123-like* probe was the biotin-labeled fragment of *PuMYB123-like* promoter containing site4 (AAATTATTG), whereas the cold probe was an unlabeled probe (50-, 100-, and 200-fold molar excess). *ProPuMYB123-like*-mut probe was the same as the *ProPuMYB123-like* probe but with site4 (AAATTATTG) mutated to TTTTTTTTT. The His-tagged PuHB40 protein was purified. **C)** ChIP-qPCR assay results reflected the in vivo

binding of PuHB40 to the *PuMYB123-like* promoter. Cross-linked chromatin samples were extracted from *PuHB40*-MYC-overexpressing pear calli with 3 biological replicates and precipitated with an anti-MYC antibody. Eluted DNA was used to amplify *PuMYB123-like* promoter fragments and eight regions (P1 to P8) were examined, among which P6 containing site4. Pear calli overexpressing the MYC sequence were used as a negative control. Three biological replicates of pear calli were used in the ChIP assay. **D)** Transient overexpression of *PuHB40* with or without transient silencing of *PuMYB123-like* in pear seedlings. The full-length CDS of *PuHB40* and reverse complementary gene fragment (~200 to 300 bp) of *PuMYB123-like* were inserted into the pCAMBIA1301 vector for the subsequent expression under the control of the 35S promoter. The plasmids were transformed into *A. tumefaciens* GV3101 and infiltrated into pear seedlings by vacuum. The infiltrated pear seedlings were firstly incubated in darkness for 2 days and then exposed to continuous white light for 3 days. Pear seedlings infiltrated with an empty pCAMBIA1301 vector were used as a control. **E)** Anthocyanin content in pear seedlings. **F)** Expression levels of *PuHB40*, *PuMYB123-like* and anthocyanin-related genes in pear seedlings. Data are presented as means ± s.d. of 3 biological replicates **C), E)**, **F)** and 6 biological replicates **A)**. Asterisks indicate significant differences compared with empty vector **A)** and control **C), E)**, **F)** (two-tailed Student's *t-test*, $*P < 0.05$, $**P < 0.01$; ns, no significance, $P > 0.05$), all exact P values are shown in Supplemental Data Set S5.

Figure 6. PuHB40 promotes anthocyanin biosynthesis. A) Phenotype of pear calli overexpressing and silencing *PuHB40* (scale bar, 1 cm). Wild type pear calli were used as the negative control. Pear calli were incubated under continuous white light at 17°C for 3 d. **B) a**nd **D)** Expression levels of *PuHB40*, *PpMYB123-like* and anthocyanin-related genes in *PuHB40*-OE and *PuHB40*-RNAi pear calli after light treatment. *PcACTIN* was amplified as an internal control. **C)** Anthocyanin contents in *PuHB40*-OE and *PuHB40*-RNAi pear calli after light treatment. **E)** Phenotype of *Arabidopsis* seedlings overexpressing *PuHB40* (scale bar, 2 mm). Wild-type 'Columbia 0' was used as the negative control. *Arabidopsis* seedlings were grown on 1/2 MS medium for 5 days. **F)** Expression levels of *PuHB40* in transgenic *Arabidopsis* lines. *AtPP2A* was amplified as an internal control. **G)** Anthocyanin contents in transgenic *Arabidopsis* lines. **H)** The hypocotyl length of transgenic *Arabidopsis* seedlings. **I)** Expression levels of anthocyanin-related genes in transgenic *Arabidopsis* lines. Data are presented as means ± s.d. of 3 biological replicates **B-D)**, **F), G), I)** and 15 biological replicates **H)**. Asterisks indicate significant differences compared with wild type pear calli **B-D)** and wild type *Arabidopsis* **F-I)** (two-tailed Student's *t-test*, *$P <$ 0.05, **$P <$ 0.01), all exact P values are shown in Supplemental Data Set S5.

Figure 7. Identification of PuPP2AA2 and its interaction with PuHB40. A) Phylogenetic analysis of PuPP2AA2. The phylogenetic tree was generated using the Maximum-likelihood method of MEGA (version 10.1.8). Bootstrap values from 1000 replicates for each branch are shown. The protein sequences from pear (Pu, *Pyrus pyrifolia*, highlighted with blue dots) and Arabidopsis (*Arabidopsis thaliana*, highlighted with rose red dots). PuPP2AA2 (EVM0003533.1) is bolded and enlarged. **B)** Subcellular localization of PuPP2AA2 in *N. benthamiana* leaf cells (scale bar, 25 μm). **C)** The expression levels of *PuPP2AA2* in pear seedlings under high-light stress treatment or high-light stress supplemented with DMTU treatment. *PuACTIN* was amplified as an internal control. **D)** Changes in the PuPP2AA2 abundance in pear seedlings. Pear seedlings transiently overexpressed MYC-tagged *PuPP2AA2* and were exposed to high-light stress treatment or high-light stress supplemented with DMTU treatment for 0 h, 3 h, 6 h, 12 h. The PuPP2AA2 protein was detected by immunoblotting with anti-MYC monoclonal antibody. **E)** The interaction between PuPP2AA2 and PuHB40 using a pull-down assay. PuHB40-His was incubated with GST or PuPP2AA2-GST. The interaction proteins were precipitated by GST-Trap agarose beads and analysed by immunoblotting with an anti-GST or anti-His antibody. **F)** The interaction between PuPP2AA2 and PuHB40 was detected by luciferase complementation imaging assays. The pseudocolour scale bar indicates the range of luminescence intensity. **G)** Physical association between PuPP2AA2 and PuHB40 confirmed by a BiFC assay. Fluorescence indicates positive interactions (scale bar, 25 μm). Data are presented as means ± s.d. of 3 biological replicates. Asterisks indicate significant differences compared with high-light stress supplemented with DMTU treatment (two-tailed Student's *t-test*, *$P < 0.05$), all exact *P* values are shown in Supplemental Data Set S5.

Figure 8. PuHB40 was dephosphorylated by PP2A phosphatase. A) Phos-tag analysis of phosphorylation of PuHB40 after cantharidin treatment. *A. tumefaciens* GV3101 containing PuHB40-GFP plasmid infiltrated into *N. benthamiana* leaves by injection and 50 μM cantharidin was injected into the same position as the agrobacteria after 24 h. *N. benthamiana* leaves injected corresponding concentration of DMSO were used as negative control. Protein samples were extracted from *N. benthamiana* leaves after 24 h and were separated in a Phos-tag gel (upper panel), and PuHB40-GFP with different phosphorylated status was detected using the anti-GFP antibody. Total PuHB40-GFP proteins were analyzed by SDS-PAGE without Phos-tag (indicated as "Regular"), and were detected using anti-GFP antibody. P+ and P- indicate phosphorylated and non-phosphorylated form of PuHB40, respectively. The intensity of the bands was measured with the Image J software and was calculated as the ratio of the P+/P- form. **B)** Immunoblotting analysis showing that the PP2A complex directly dephosphorylates PuHB40 *in vitro*. MYC, PP2A complex, and PuHB40-GFP were immunopurified from pear seedlings transiently overexpressing MYC, PuPP2AA2-MYC, and PuHB40-GFP using MYC-Trap or GFP-Trap, respectively.

After incubation of PP2A complexes with PuHB40-GFP (MYC with PuHB40-GFP was set as control), protein samples in the reaction system were separated in Phos-tag gel (upper panel) and detected using the anti-PuHB40 antibody. The intensity of the bands was measured with the Image J software and calculated as the ratio of the P+/P-. Total proteins in the reaction system were analyzed by SDS-PAGE without Phos-tag (indicated as "Regular"). MYC, PuPP2AA2-MYC, and PuHB40-GFP were set as the loading controls and detected by the anti-MYC or anti-PuHB40 antibody, respectively. **C)** Subcellular localization of PuPP2AA2D in *N. benthamiana* leaf cells (scale bar, 25 μm). PuPP2AA2D is PuPP2AA2 with heat repeats 14 and 15 in the C-terminal deleted. **D)** The interaction between PuPP2AA2D and PuHB40 confirmed by a BiFC assay. Fluorescence indicates positive interactions (scale bar, 25 μm). **E)** The interaction between PuPP2AA2D and PuHB40 was detected by luciferase complementation imaging assays. The pseudocolour scale bar indicates the range of luminescence intensity. **F)** Immunoblotting analysis showing that PP2A dephosphorylates PuHB40 *in vivo*. Protein extracts from *N. benthamiana* leaves co-infiltrated PuHB40-GFP with MYC, PuPP2AA2-MYC, or PuPP2AA2D-MYC were separated in Phos-tag gel (upper panel) and detected using the anti-GFP antibody (MYC with PuHB40-GFP was set as control). The intensity of the bands was measured with the Image J software and calculated the ratio of P+/P−. Total PuHB40-GFP, MYC, PuPP2AA2-MYC and PuPP2AA2D-MYC proteins were analyzed by SDS-PAGE without Phos-tag (indicated as "Regular"). PuHB40-GFP, MYC, PuPP2AA2-MYC and PuPP2AA2D-MYC were set as the loading controls and detected by the anti-GFP or anti-MYC antibody, respectively.

G) Dual-luciferase assay showed that dephosphorylation of PuHB40 by PP2A significantly inhibited the activation of *PuMYB123-like* promoter by PuHB40. The *PuMYB123-like* promoter was cloned into the pGreenII 0800-LUC (firefly luciferase) vector, and the full-length CDS of *PuHB40*, PuPP2AA2 and PuPP2AA2D was cloned into the pGreenII 0029 62-SK vector. The PuHB40 was used as control. Data are presented as means ± s.d. of 6 biological replicates **G)**. Asterisks indicate significant differences compared PuHB40 **G)** (two-tailed Student's *t-test*, **$P <$ 0.01; ns, no significance, $P > 0.05$), all exact P values are shown in Supplemental Data Set S5.

Figure 9. Attenuation of PuHB40-induced anthocyanin biosynthesis by PuPP2AA2. **A)** Phenotype of pear calli overexpressing *PuPP2AA2* (scale bar, 1 cm). Wild type pear calli were used as the negative control. Pear calli were incubated under continuous white light at 17°C for 3 d. **B)** Anthocyanin contents in *PuPP2AA2*-OE after light treatment. **C)** Expression levels of *PuPP2AA2* and anthocyanin-related genes in *PuPP2AA2*-OE pear calli after light treatment. *PcACTIN* was amplified as an internal control. **D)** Transient overexpression of *PuHB40* with or without transient overexpressing of *PuPP2AA2* in pear seedlings. The full-length CDS of *PuHB40* and *PuPP2AA2* were inserted into the pCAMBIA1301 vector for the subsequent expression under the control of the 35S promoter. The plasmids were transformed into *A. tumefaciens* GV3101 and infiltrated into pear seedlings by vacuum. The infiltrated pear seedlings were firstly incubated in darkness for 2 days and then exposed to continuous white light for 3 days. Pear seedlings infiltrated with an empty pCAMBIA1301 vector were used as a control. **E)** Anthocyanin content in pear seedlings. **F)** Expression levels of *PuHB40*, *PuPP2AA2*, *PuMYB123-like* and anthocyanin-related genes in pear seedlings. **G)** Transient overexpression of *PuHB40* with or without transient overexpressing of *PuPP2AA2* in fruit (scale bar, 1 cm). The full-length CDS of *PuHB40* and *PuPP2AA2* were cloned to the pGreenII 0029 62-SK vector for the subsequent expression under the control of the 35S promoter. Pear fruits were infiltrated with *A. tumefaciens* GV3101 cells containing the recombinant plasmid using a needle-free syringe. Fruits infiltrated with an empty pGreenII 0029 62-SK vector were used as a control. The phenotypes were examined after a 1 day dark and 4 days light treatment. **H)** Total anthocyanin content in pear fruits. **I)** Expression levels of *PuHB40*, *PuPP2AA2* and anthocyanin-related genes in pear fruits. Data are presented as means ± s.d. of 3 biological replicates. Asterisks indicate significant differences compared with wild type pear calli **B)** and **C)**, empty vector **E), F), H), I)** (two-tailed Student's *t-test*, *$P < 0.05$, **$P < 0.01$; ns, no significance, $P > 0.05$), all exact *P* values are shown in Supplemental Data Set S5.

Figure 10. Proposed model for PuHB40-mediated ROS-dependent anthocyanin biosynthesis in pears. PuHB40 is the key transcription factor in ROS-induced anthocyanin biosynthesis. ROS promote PuHB40 expression and then positively regulate its transcriptional activity on PuMYB123-like/PubHLH3 complex to induce anthocyanin biosynthesis. The transcriptional activity of PuHB40 depends on its phosphorylation level which is regulated by PP2A. ROS maintain phosphorylation of PuHB40 at a high level, and enhance the transcriptional activity of PuHB40 on *PuMYB123-like* by reducing the transcriptional and translational levels of PuPP2AA2 under high-light stress, and thus promote anthocyanin biosynthesis. The lightning indicates the intensity of transcriptional regulation of *PuMYB123-like* by PuHB40.

Parsed Citations

Acero N, Gradillas A, Beltran M, Garcia A, Munoz Mingarro D. Comparison of phenolic compounds profile and antioxidant properties of different sweet cherry (Prunus avium L.) varieties. Food Chem. 2019:279:260-271. https://doi.org/10.1016/j.foodchem.2018.12.008
  Google Scholar: Author Only Title Only Author and Title

Altangerel N, Ariunbold GO, Gorman C, Alkahtani MH, Borrego EJ, Bohlmeyer D, Hemmer P, Kolomiets MV, Yuan JS, Scully MO. In vivo diagnostics of early abiotic plant stress response via Raman spectroscopy. Proc Natl Acad Sci U S A. 2017:114(13):3393-3396. https://doi.org/10.1073/pnas.1701328114
  Google Scholar: Author Only Title Only Author and Title

An JP, Wang XF, Zhang XW, Xu HF, Bi SQ, You CX, Hao YJ. An apple MYB transcription factor regulates cold tolerance and anthocyanin accumulation and undergoes MIEL1-mediated degradation. Plant Biotechnol J. 2020:18(2):337-353. https://doi.org/10.1111/pbi.13201
  Google Scholar: Author Only Title Only Author and Title

An XH, Tian Y, Chen KQ, Liu XJ, Liu DD, Xie XB, Cheng CG, Cong PH, Hao YJ. MdMYB9 and MdMYB11 are involved in the regulation of the JA-induced biosynthesis of anthocyanin and proanthocyanidin in apples. Plant Cell Physiol. 2015:56(4):650-662. https://doi.org/10.1093/pcp/pcu205
  Google Scholar: Author Only Title Only Author and Title

Ariel FD, Manavella PA, Dezar CA, Chan RL. The true story of the HD-Zip family. Trends Plant Sci. 2007:12(9):419-426. https://doi.org/10.1016/j.tplants.2007.08.003
  Google Scholar: Author Only Title Only Author and Title

Bai S, Tao R, Yin L, Ni J, Yang Q, Yan X, Yang F, Guo X, Li H, Teng Y. Two B-box proteins, PpBBX18 and PpBBX21, antagonistically regulate anthocyanin biosynthesis via competitive association with Pyrus pyrifolia ELONGATED HYPOCOTYL 5 in the peel of pear fruit. Plant J. 2019a:100(6):1208-1223. https://doi.org/10.1111/tpj.14510
  Google Scholar: Author Only Title Only Author and Title

Bai S, Tao R, Tang Y, Yin L, Ma Y, Ni J, Yan X, Yang Q, Wu Z, Zeng Y, et al. BBX16, a B-box protein, positively regulates light-induced anthocyanin accumulation by activating MYB10 in red pear. Plant Biotechnol J. 2019b:17(10):1985-1997. https://doi.org/10.1111/pbi.13114
  Google Scholar: Author Only Title Only Author and Title

Bi X, Zhang J, Chen C, Zhang D, Li P, Ma F. Anthocyanin contributes more to hydrogen peroxide scavenging than other phenolics in apple peel. Food Chem. 2014:152:205-209. https://doi.org/10.1016/j.foodchem.2013.11.088
  Google Scholar: Author Only Title Only Author and Title

Bian C, Guo X, Zhang Y, Wang L, Xu T, DeLong A, Dong J. Protein phosphatase 2A promotes stomatal development by stabilizing SPEECHLESS in Arabidopsis. Proc Natl Acad Sci U S A. 2020:117(23):13127-13137. https://doi.org/10.1073/pnas.1912075117
  Google Scholar: Author Only Title Only Author and Title

Byrne DP, Shrestha S, Galler M, Cao M, Daly LA, Campbell AE, Eyers CE, Veal EA, Kannan N, Eyers PA. Aurora A regulation by reversible cysteine oxidation reveals evolutionarily conserved redox control of Ser/Thr protein kinase activity. Sci Signal. 2020:13(613). https://doi.org/10.1126/scisignal.aax2713
  Google Scholar: Author Only Title Only Author and Title

Catala R, Medina J, Salinas J. Integration of low temperature and light signaling during cold acclimation response in Arabidopsis. Proc Natl Acad Sci U S A. 2011:108(39):16475-16480. https://doi.org/10.1073/pnas.1107161108
  Google Scholar: Author Only Title Only Author and Title

Chan KX, Mabbitt PD, Phua SY, Mueller JW, Nisar N, Gigolashvili T, Stroeher E, Grassl J, Arlt W, Estavillo GM, et al. Sensing and signaling of oxidative stress in chloroplasts by inactivation of the SAL1 phosphoadenosine phosphatase. Proc Natl Acad Sci U S A. 2016:113(31):E4567-4576. https://doi.org/10.1073/pnas.1604936113
  Google Scholar: Author Only Title Only Author and Title

Choi S, Kwon YR, Hossain MA, Hong S-W, Lee B-h, Lee H. A mutation in ELA1, an age-dependent negative regulator of PAP1/MYB75, causes UV- and cold stress-tolerance in Arabidopsis thaliana seedlings. Plant Sci. 2009:176(5):678-686. https://doi.org/10.1016/j.plantsci.2009.02.010
  Google Scholar: Author Only Title Only Author and Title

Croce R, van Amerongen H. Natural strategies for photosynthetic light harvesting. Nat Chem Biol. 2014:10(7):492-501. https://doi.org/10.1038/nchembio.1555
  Google Scholar: Author Only Title Only Author and Title

Das PK, Shin DH, Choi SB, Yoo SD, Choi G, Park YI. Cytokinins enhance sugar-induced anthocyanin biosynthesis in Arabidopsis. Mol Cells. 2012:34(1):93-101. https://doi.org/10.1007/s10059-012-0114-2
  Google Scholar: Author Only Title Only Author and Title

DeLong A. Switching the flip: protein phosphatase roles in signaling pathways. Curr Opin Plant Biol. 2006:9(5):470-477. https://doi.org/10.1016/j.pbi.2006.07.015
Google Scholar: Author Only Title Only Author and Title

Dubos C, Stracke R, Grotewold E, Weisshaar B, Martin C, Lepiniec L. MYB transcription factors in Arabidopsis. Trends Plant Sci. 2010:15(10):573-581. https://doi.org/10.1016/j.tplants.2010.06.005
Google Scholar: Author Only Title Only Author and Title

Durian G, Rahikainen M, Alegre S, Brosche M, Kangasjarvi S. Protein Phosphatase 2A in the Regulatory Network Underlying Biotic Stress Resistance in Plants. Front Plant Sci. 2016:7:812. https://doi.org/10.3389/fpls.2016.00812
Google Scholar: Author Only Title Only Author and Title

Espley RV, Hellens RP, Putterill J, Stevenson DE, Kutty-Amma S, Allan AC. Red colouration in apple fruit is due to the activity of the MYB transcription factor, MdMYB10. Plant J. 2007:49(3):414-427. https://doi.org/10.1111/j.1365-313X.2006.02964.x
Google Scholar: Author Only Title Only Author and Title

Farkas I, Dombradi V, Miskei M, Szabados L, Koncz C. Arabidopsis PPP family of serine/threonine phosphatases. Trends Plant Sci. 2007:12(4):169-176. https://doi.org/10.1016/j.tplants.2007.03.003
Google Scholar: Author Only Title Only Author and Title

Feild TS, Lee DW, Holbrook NM. Why Leaves Turn Red in Autumn. The Role of Anthocyanins in Senescing Leaves of Red-Osier Dogwood. Plant Physiol. 2001:127(2):566-574. https://doi.org/10.1104/pp.010063
Google Scholar: Author Only Title Only Author and Title

Feng S, Wang Y, Yang S, Xu Y, Chen X. Anthocyanin biosynthesis in pears is regulated by a R2R3-MYB transcription factor PyMYB10. Planta. 2010:232(1):245-255. https://doi.org/10.1007/s00425-010-1170-5
Google Scholar: Author Only Title Only Author and Title

Galvez-Valdivieso G, Fryer MJ, Lawson T, Slattery K, Truman W, Smirnoff N, Asami T, Davies WJ, Jones AM, Baker NR, et al. The high light response in Arabidopsis involves ABA signaling between vascular and bundle sheath cells. Plant Cell. 2009:21(7):2143-2162. https://doi.org/10.1105/tpc.108.061507
Google Scholar: Author Only Title Only Author and Title

Gould KS, McKelvie J, Markham KR. Do anthocyanins function as antioxidants in leaves? Imaging of H2O2 in red and green leaves after mechanical injury. Plant Cell Environ. 2002:25(10):1261-1269. https://doi.org/10.1046/j.1365-3040.2002.00905.x
Google Scholar: Author Only Title Only Author and Title

Groves MR, Hanlon N, Turowski P, Hemmings BA, Barford D. The structure of the protein phosphatase 2A PR65/A subunit reveals the conformation of its 15 tandemly repeated HEAT motifs. Cell. 1999:96(1):99-110. https://doi.org/10.1016/S0092-8674(00)80963-0
Google Scholar: Author Only Title Only Author and Title

Gupta R, Luan S. Redox control of protein tyrosine phosphatases and mitogen-activated protein kinases in plants. Plant Physiol. 2003:132(3):1149-1152. https://doi.org/10.1104/pp.103.020792
Google Scholar: Author Only Title Only Author and Title

Harris JC, Hrmova M, Lopato S, Langridge P. Modulation of plant growth by HD-Zip class I and II transcription factors in response to environmental stimuli. New Phytol. 2011:190(4):823-837. https://doi.org/10.1111/j.1469-8137.2011.03733.x
Google Scholar: Author Only Title Only Author and Title

Himmelbach A, Hoffmann T, Leube M, Hohener B, Grill E. Homeodomain protein ATHB6 is a target of the protein phosphatase ABI1 and regulates hormone responses in Arabidopsis. Embo J. 2002:21(12):3029-3038. https://doi.org/10.1093/emboj/cdf316
Google Scholar: Author Only Title Only Author and Title

Hipsch M, Lampl N, Zelinger E, Barda O, Waiger D, Rosenwasser S. Sensing stress responses in potato with whole-plant redox imaging. Plant Physiol. 2021:187(2):618-631. https://doi.org/10.1093/plphys/kiab159
Google Scholar: Author Only Title Only Author and Title

Hoch WA, Zeldin EL, McCown BH. Physiological significance of anthocyanins during autumnal leaf senescence. Tree Physiol. 2001:21(1):1-8. https://doi.org/10.1093/treephys/21.1.1
Google Scholar: Author Only Title Only Author and Title

Honkanen RE. Cantharidin, another natural toxin that inhibits the activity of serine/threonine protein phosphatases types 1 and 2A. FEBS Lett. 1993:330(3):283-286. https://doi.org/10.1016/0014-5793(93)80889-3
Google Scholar: Author Only Title Only Author and Title

Jaakola L. New insights into the regulation of anthocyanin biosynthesis in fruits. Trends Plant Sci. 2013:18(9):477-483. https://doi.org/10.1016/j.tplants.2013.06.003
Google Scholar: Author Only Title Only Author and Title

Jalmi SK, Sinha AK. ROS mediated MAPK signaling in abiotic and biotic stress- striking similarities and differences. Front Plant Sci. 2015:6:769. https://doi.org/10.3389/fpls.2015.00769

Google Scholar: Author Only Title Only Author and Title

Janssens V, Goris J. Protein phosphatase 2A: a highly regulated family of serine/threonine phosphatases implicated in cell growth and signalling. Biochem J. 2001:353(3):417-439. https://doi.org/10.1042/0264-6021:3530417

Google Scholar: Author Only Title Only Author and Title

Jia M, Li X, Wang W, Li T, Dai Z, Chen Y, Zhang K, Zhu H, Mao W, Feng Q, et al. SnRK2 subfamily I protein kinases regulate ethylene biosynthesis by phosphorylating HB transcription factors to induce ACO1 expression in apple. New Phytol. 2022:234(4):1262-1277. https://doi.org/10.1111/nph.18040

Google Scholar: Author Only Title Only Author and Title

Jiang T, He Y, Wu Z, Cui Y, Wang X, Huang H, Fan Y, Han M, Wang J, Wang S, et al. Enhancing stimulation of cyaniding, GhLDOX3 activates reactive oxygen species to regulate tolerance of alkalinity negatively in cotton. Ecotox Environ Safe. 2023:267:115655. https://doi.org/10.1016/j.ecoenv.2023.115655

Google Scholar: Author Only Title Only Author and Title

Kamibayashi C, Lickteig RL, Estes R, Walter G, Mumby MC. Expression of the A subunit of protein phosphatase 2A and characterization of its interactions with the catalytic and regulatory subunits. J Biol Chem. 1992:267(30):21864-21872. https://doi.org/10.1016/s0021-9258(19)36692-x

Google Scholar: Author Only Title Only Author and Title

Kerchev P, Waszczak C, Lewandowska A, Willems P, Shapiguzov A, Li Z, Alseekh S, Muhlenbock P, Hoeberichts FA, Huang J, et al. Lack of GLYCOLATE OXIDASE1, but Not GLYCOLATE OXIDASE2, Attenuates the Photorespiratory Phenotype of CATALASE2-Deficient Arabidopsis. Plant Physiol. 2016:171(3):1704-1719. https://doi.org/10.1104/pp.16.00359

Google Scholar: Author Only Title Only Author and Title

Koes R, Verweij W, Quattrocchio F. Flavonoids: a colorful model for the regulation and evolution of biochemical pathways. Trends Plant Sci. 2005:10(5):236-242. https://doi.org/10.1016/j.tplants.2005.03.002

Google Scholar: Author Only Title Only Author and Title

Kovtun Y, Chiu WL, Tena G, Sheen J. Functional analysis of oxidative stress-activated mitogen-activated protein kinase cascade in plants. Proc Natl Acad Sci U S A. 2000:97(6):2940-2945. https://doi.org/10.1073/pnas.97.6.2940

Google Scholar: Author Only Title Only Author and Title

Landi M, Tattini M, Gould KS. Multiple functional roles of anthocyanins in plant-environment interactions. Environ Exp Bot. 2015:119:4-17. https://doi.org/10.1016/j.envexpbot.2015.05.012

Google Scholar: Author Only Title Only Author and Title

Lee DW, Gould KS. Why leaves turn red: pigments called anthocyanins probably protect leaves from light damage by direct shielding and by scavenging free radicals. AM SCI. 2002:90(6):524-531. https://doi.org/10.1511/2002.6.524

Google Scholar: Author Only Title Only Author and Title

Lesk C, Rowhani P, Ramankutty N. Influence of extreme weather disasters on global crop production. Nature. 2016:529(7584):84-87. https://doi.org/10.1038/nature16467

Google Scholar: Author Only Title Only Author and Title

Li Y, Wang Y, Tan S, Li Z, Yuan Z, Glanc M, Domjan D, Wang K, Xuan W, Guo Y, et al. Root Growth Adaptation is Mediated by PYLs ABA Receptor-PP2A Protein Phosphatase Complex. Adv Sci. 2020:7(3):1901455. https://doi.org/10.1002/advs.201901455

Google Scholar: Author Only Title Only Author and Title

Li YY, Mao K, Zhao C, Zhao XY, Zhang HL, Shu HR, Hao YJ. MdCOP1 ubiquitin E3 ligases interact with MdMYB1 to regulate light-induced anthocyanin biosynthesis and red fruit coloration in apple. Plant Physiol. 2012:160(2):1011-1022. https://doi.org/10.1104/pp.112.199703

Google Scholar: Author Only Title Only Author and Title

Lillo C, Kataya AR, Heidari B, Creighton MT, Nemie-Feyissa D, Ginbot Z, Jonassen EM. Protein phosphatases PP2A, PP4 and PP6: mediators and regulators in development and responses to environmental cues. Plant Cell Environ. 2014:37(12):2631-2648. https://doi.org/10.1111/pce.12364

Google Scholar: Author Only Title Only Author and Title

Lin-Wang K, Micheletti D, Palmer J, Volz R, Lozano L, Espley R, Hellens RP, Chagne D, Rowan DD, Troggio M, et al. High temperature reduces apple fruit colour via modulation of the anthocyanin regulatory complex. Plant Cell Environ. 2011:34(7):1176-1190. https://doi.org/10.1111/j.1365-3040.2011.02316.x

Google Scholar: Author Only Title Only Author and Title

Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the $2-\triangle\triangle CT$ method. Methods. 2001:25(4):402-408. https://doi.org/10.1006/meth.2001.1262

Google Scholar: Author Only Title Only Author and Title

Mine A, Berens ML, Nobori T, Anver S, Fukumoto K, Winkelmuller TM, Takeda A, Becker D, Tsuda K. Pathogen exploitation of an abscisic acid- and jasmonate-inducible MAPK phosphatase and its interception by Arabidopsis immunity. Proc Natl Acad Sci U S A. 2017:114(28):7456-7461. https://doi.org/10.1073/pnas.1702613114
  Google Scholar: Author Only Title Only Author and Title

Mittler R. ROS Are Good. Trends Plant Sci. 2017:22(1):11-19. https://doi.org/10.1016/j.tplants.2016.08.002
  Google Scholar: Author Only Title Only Author and Title

Mittler R, Zandalinas SI, Fichman Y, Van Breusegem F. Reactive oxygen species signalling in plant stress responses. Nat Rev Mol Cell Biol. 2022:23(10):663-679. https://doi.org/10.1038/s41580-022-00499-2
  Google Scholar: Author Only Title Only Author and Title

Nakabayashi R, Yonekura-Sakakibara K, Urano K, Suzuki M, Yamada Y, Nishizawa T, Matsuda F, Kojima M, Sakakibara H, Shinozaki K, et al. Enhancement of oxidative and drought tolerance in Arabidopsis by overaccumulation of antioxidant flavonoids. Plant J. 2014:77(3):367-379. https://doi.org/10.1111/tpj.12388
  Google Scholar: Author Only Title Only Author and Title

Neill SO, Gould KS, Kilmartin PA, Mitchell KA, Markham KR. Antioxidant activities of red versus green leaves in Elatostema rugosum. Plant Cell Environ. 2002:25(4):539-547. https://doi.org/10.1046/j.1365-3040.2002.00837.x
  Google Scholar: Author Only Title Only Author and Title

Ni J, Wang S, Yu W, Liao Y, Pan C, Zhang M, Tao R, Wei J, Gao Y, Wang D, et al. The ethylene-responsive transcription factor PpERF9 represses PpRAP2.4 and PpMYB114 via histone deacetylation to inhibit anthocyanin biosynthesis in pear. Plant Cell. 2023:35(6):2271-2292. https://doi.org/10.1093/plcell/koad077
  Google Scholar: Author Only Title Only Author and Title

Olsson AS, Engström P, Söderman E. The homeobox genes ATHB12 and ATHB7 encode potential regulators of growth in response to water deficit in Arabidopsis. Plant Mol Biol. 2004:55(5):663-677. https://doi.org/10.1007/s11103-004-1581-4
  Google Scholar: Author Only Title Only Author and Title

Qu Y, Bai X, Zhu YJ, Qi R, Tian G, Wang Y, Li YH, Zhang KM. Reactive Oxygen Species Acts as an Important Inducer in Low-temperature-induced Anthocyanin Biosynthesis in Begonia semperflorens. J Am Soc Hortic Sci. 2018:143(6):486-493. https://doi.org/10.21273/JASHS04488-18
  Google Scholar: Author Only Title Only Author and Title

Ren H, Rao J, Tang M, Li Y, Dang X, Lin D. PP2A interacts with KATANIN to promote microtubule organization and conical cell morphogenesis. J Integr Plant Biol. 2022:64(8):1514-1530. https://doi.org/10.1111/jipb.13281
  Google Scholar: Author Only Title Only Author and Title

Ruediger R, Hentz M, Fait J, Mumby M, Walter G. Molecular model of the A subunit of protein phosphatase 2A interaction with other subunits and tumor antigens. J Virol. 1994:68(1):123-129. https://doi.org/10.1128/JVI.68.1.123-129.1994
  Google Scholar: Author Only Title Only Author and Title

Schaart JG, Dubos C, De La Fuente IR, van Houwelingen AMML, de Vos RCH, Jonker HH, Xu WJ, Routaboul JM, Lepiniec L, Bovy AG. Identification and characterization of MYB-bHLH-WD40 regulatory complexes controlling proanthocyanidin biosynthesis in strawberry (Fragaria × ananassa) fruits. New Phytol. 2012:197(2):454-467. https://doi.org/10.1111/nph.12017
  Google Scholar: Author Only Title Only Author and Title

Shan X, Zhang Y, Peng W, Wang Z, Xie D. Molecular mechanism for jasmonate-induction of anthocyanin accumulation in Arabidopsis. J Exp Bot. 2009:60(13):3849-3860. https://doi.org/10.1093/jxb/erp223
  Google Scholar: Author Only Title Only Author and Title

Shi H, Liu G, Wei Y, Chan Z. The zinc-finger transcription factor ZAT6 is essential for hydrogen peroxide induction of anthocyanin synthesis in Arabidopsis. Plant Mol Biol. 2018:97(1-2):165-176. https://doi.org/10.1007/s11103-018-0730-0
  Google Scholar: Author Only Title Only Author and Title

Soderman E, Mattsson J, Engstrom P. The Arabidopsis homeobox gene ATHB-7 is induced by water deficit and by abscisic acid. Plant J. 1996:10(2):375-381. https://doi.org/10.1046/j.1365-313x.1996.10020375.x
  Google Scholar: Author Only Title Only Author and Title

Söderman E, Hjellström M, Fahleson J, Engström P. The HD-Zip gene ATHB6 in Arabidopsis is expressed in developing leaves, roots and carpels and up-regulated by water deficit conditions. Plant Mol Biol. 1999:40(6):1073-1083. https://doi.org/10.1023/A:1006267013170
  Google Scholar: Author Only Title Only Author and Title

Steyn WJ, Wand SJE, Holcroft DM, Jacobs G. Anthocyanins in vegetative tissues: a proposed unified function in photoprotection. New Phytol. 2002:155(3):349-361. https://doi.org/10.1046/j.1469-8137.2002.00482.x
  Google Scholar: Author Only Title Only Author and Title

Takos AM, Jaffe FW, Jacob SR, Bogs J, Robinson SP, Walker AR. Light-induced expression of a MYB gene regulates anthocyanin

biosynthesis in red apples. Plant Physiol. 2006:142(3):1216-1232. https://doi.org/10.1104/pp.106.088104
Google Scholar: Author Only Title Only Author and Title

Uhrig RG, Labandera AM, Moorhead GB. Arabidopsis PPP family of serine/threonine protein phosphatases: many targets but few engines. Trends Plant Sci. 2013:18(9):505-513. https://doi.org/10.1016/j.tplants.2013.05.004
Google Scholar: Author Only Title Only Author and Title

Wang L, Leister D, Guan L, Zheng Y, Schneider K, Lehmann M, Apel K, Kleine T. The Arabidopsis SAFEGUARD1 suppresses singlet oxygen-induced stress responses by protecting grana margins. Proc Natl Acad Sci U S A. 2020:117(12):6918-6927. https://doi.org/10.1073/pnas.1918640117
Google Scholar: Author Only Title Only Author and Title

Winkel-Shirley B. Flavonoid biosynthesis. A colorful model for genetics, biochemistry, cell biology, and biotechnology. Plant Physiol. 2001:126(2):485-493. https://doi.org/10.1104/pp.126.2.485
Google Scholar: Author Only Title Only Author and Title

Xie XB, Li S, Zhang RF, Zhao J, Chen YC, Zhao Q, Yao YX, You CX, Zhang XS, Hao YJ. The bHLH transcription factor MdbHLH3 promotes anthocyanin accumulation and fruit colouration in response to low temperature in apples. Plant Cell Environ. 2012:35(11):1884-1897. https://doi.org/10.1111/j.1365-3040.2012.02523.x
Google Scholar: Author Only Title Only Author and Title

Xu H, Zou Q, Yang G, Jiang S, Fang H, Wang Y, Zhang J, Zhang Z, Wang N, Chen X. MdMYB6 regulates anthocyanin formation in apple both through direct inhibition of the biosynthesis pathway and through substrate removal. Hortic Res. 2020:7(1):72. https://doi.org/10.1038/s41438-020-0294-4
Google Scholar: Author Only Title Only Author and Title

Xu W, Dubos C, Lepiniec L. Transcriptional control of flavonoid biosynthesis by MYB-bHLH-WDR complexes. Trends Plant Sci. 2015:20(3):176-185. https://doi.org/10.1016/j.tplants.2014.12.001
Google Scholar: Author Only Title Only Author and Title

Xu Z, Mahmood K, Rothstein SJ. ROS Induces Anthocyanin Production Via Late Biosynthetic Genes and Anthocyanin Deficiency Confers the Hypersensitivity to ROS-Generating Stresses in Arabidopsis. Plant Cell Physiol. 2017:58(8):1364-1377. https://doi.org/10.1093/pcp/pcx073
Google Scholar: Author Only Title Only Author and Title

Zhang D, Yu B, Bai J, Qian M, Shu Q, Su J, Teng Y. Effects of high temperatures on UV-B/visible irradiation induced postharvest anthocyanin accumulation in 'Yunhongli No. 1' (Pyrus pyrifolia Nakai) pears. Sci Hortic. 2012:134:53-59. https://doi.org/10.1016/j.scienta.2011.10.025
Google Scholar: Author Only Title Only Author and Title

Zhang L, Tao R, Wang S, Gao Y, Wang L, Yang S, Zhang X, Yu W, Wu X, Li K, et al. PpZAT5 suppresses the expression of a B-box gene PpBBX18 to inhibit anthocyanin biosynthesis in the fruit peel of red pear. Front Plant Sci. 2022:13:1022034. https://doi.org/10.3389/fpls.2022.1022034
Google Scholar: Author Only Title Only Author and Title

Zhou HW, Nussbaumer C, Chao Y, DeLong A. Disparate roles for the regulatory A subunit isoforms in Arabidopsis protein phosphatase 2A. Plant Cell. 2004:16(3):709-722. https://doi.org/10.1105/tpc.018994
Google Scholar: Author Only Title Only Author and Title