Linked *OXTR* Variants Are Associated with Social Behavior Differences in Bonobos (*Pan paniscus*)

Sara A. Skiba*[1], Alek Hansen[2], Ryan McCall[2], Azeeza Byers[2,3], Sarah Waldron[2],

Amanda J. Epping[1], Jared P. Taglialatela[1,3], & Martin L. Hudson[2]

1. Ape Cognition and Conservation Initiative (Ape Initiative)
Des Moines, IA
2. Kennesaw State University
Department of Molecular and Cellular Biology
Kennesaw, GA
3. Kennesaw State University
Department of Ecology, Evolution, and Organismal Biology
Kennesaw, GA

*Corresponding Author
4200 Evergreen Ave
Des Moines, IA 50320
Email: sara@apeinitiative.org
ORCID ID: 0000-0002-0014-099X

Abstract

Single-nucleotide polymorphisms (SNPs) in forkhead box protein P2 (*FOXP2*) and oxytocin receptor (*OXTR*) genes have been associated with linguistic and social development in humans, as well as to symptom severity in autism spectrum disorder (ASD). Studying biobehavioral mechanisms in the species most closely related to humans can provide insights into the origins of human communication, and the impact of genetic variation on complex behavioral phenotypes. Here, we aimed to determine if bonobos (*Pan paniscus*) exhibit individual variation in *FOXP2* and *OXTR* loci that have been associated with human social development and behavior. Although the ASD-related variants were reported in 13-41% of the human population, we did not find variation at these loci in our sample of 13 bonobos. However, we did identify a novel variant in bonobo *FOXP2*, as well as four novel variants in bonobo *OXTR* that were 17-184 base pairs from the human ASD variants. We also found the same linked, homozygous allelic combination across the 4 novel *OXTR* SNPs (homozygous TGTC) in 6 of the 13 bonobos, indicating that this combination may be under positive selection. When comparing the combined *OXTR* genotypes, we found significant group differences in social behavior; bonobos with zero copies of the TGTC combination were less social than bonobos with one copy of the TGTC combination. Taken together, our findings suggest that these *OXTR* variants may influence individual-level social behavior in bonobos and support the notion that linked genetic variants are promising risk factors for social communication deficits in humans.

Keywords: *Autism, Bonobo, Genetics, Great Ape, Oxytocin, Social Behavior, Social Communication*

Introduction

Autism spectrum disorder (ASD) is characterized by social communication deficits and restricted, repetitive behaviors (RRBs) that impact daily functioning and can persist into adulthood. Determining the genetic factors that impact individual-level social communication is critical to our understanding of ASD and other neurodevelopmental disorders and may aid in identifying children at risk of developing social and linguistic impairments. The Simons Foundation created a database of genes associated with aspects of the ASD behavioral phenotype – SFARI Gene[1] – providing a systematic assessment of evidence for individual ASD-related genes[2]. This growing database highlights the complexity of human neurodevelopmental disorders, like autism, and how challenging it can be to identify biomarkers in humans.

A potential biological factor underlying individual differences in social communication is the forkhead box protein P2 (*FOXP2*). *FOXP2* is one of the first genes to be associated with human language disorders and fine orofacial motor control[3,4,5]. Most notably, researchers have determined that *FOXP2* is critical to the developmental processes underlying speech and language[3,4]. In addition, Haghighatfard and colleagues (2022) found that lower *FOXP2* expression was associated with executive dysfunction in children diagnosed with ASD[6]. Thus, it is possible that polymorphisms in *FOXP2* that affect gene regulation and protein expression may underlie individual-level differences in social communication[7].

Much like in the case of *FOXP2*, several studies have demonstrated the important role that the oxytocin receptor gene (*OXTR*) plays in social bond formation and social motivation[8,9,10,11]. In particular, researchers have documented relations

between OXTR variation and social affiliation[12], vocal symptoms[13], as well as social

communication impairments associated with ASD[14]. There is also considerable

evidence that *OXTR* SNPs are related to empathy, prosocial temperament, social

sensitivity, and stress reactivity in individuals diagnosed with ASD[15]. Collectively, these

studies suggest that variation in *FOXP2* and *OXTR* influence social communication

development and social functioning in humans.

To better understand the impact of genetic variants on complex behavioral

phenotypes, burgeoning evidence supports the study of these biobehavioral

mechanisms in nonhuman animals[16,17,18]. Indeed, variation in *FOXP2* and *OXTR* have

been associated with differences in social behavior and/or social communication in

rodents[19,20], zebrafish[21], great apes[22,23], and zebra finches[24]. Seminal work in rodents,

including knock-out experiments, account for much of what we know about ASD-

associated variants and other social communication related genes[25,26,27,28,29]. However,

recent evidence suggests that common animal models of neurodevelopmental disorders

(e.g., rodents and invertebrates) are limited in their comparability to human social

communication[30,31] and may be too phylogenetically distant from humans to advance

early identification and intervention techniques[32].

Bonobos, along with chimpanzees, are the closest living relatives to humans and

are regarded as having one of the most complex social communication systems in the

animal kingdom. While bonobos cannot be diagnosed with human neurodevelopmental

disorders, they do show significant individual-level variability in social engagement[33,34],

communicative production[35,36], and repetitive behaviors[34,37]. Several studies have

identified links between ASD-associated genes and social communication in

bonobos[9,38,39]. There is also evidence that oxytocin is related to socio-sexual behavior in

female bonobos[40]. Thus, bonobos are a promising candidate for investigating the impact

of genetic variants on human neurodevelopment and social communication.

Identifying biological factors that underlie individual-level social communication is

critical to our understanding of autism and other neurodevelopmental disorders. To this

end, we aimed to determine if bonobos - our closest living relatives - exhibit single

nucleotide variation in *FOXP2* and *OXTR* at known human loci that have been

implicated in autism or differences in social communication. Given that bonobos live in

large, dynamic social groups and that they produce complex communicative signals of

various types (vocalizations, facial expressions, manual gestures, and multi-source

signals), we hypothesized that bonobos would exhibit allelic variation in *FOXP2* and

OXTR.

Materials and Methods

<u>Genetic Analyses</u>

Biological samples were collected from 29 bonobos (7 whole blood samples and

22 buccal samples) living at the Ape Cognition and Conservation Initiative (n=7; IACUC

protocol #210305-01), the Columbus Zoo and Aquarium (n=6), and the Milwaukee

County Zoo (n=16). Whole blood samples were collected under anesthesia during the

bonobo's routine physical exam (n=6), except for one blood sample that was collected

from an awake bonobo via voluntary presentation (n=1). Buccal samples were collected

by swabbing the inner cheek or lower lip for 10-15 seconds with a QIAGEN OmniSwab

(n=22). All buccal samples were taken from awake bonobos that presented voluntarily

for sample collection.

Autism-associated genes included in this study were *FOXP2* and *OXTR*. Primer

pairs were designed using NCBI Primer-BLAST[41] and ApE[42] to flank each SNP by ~250

base pairs (giving approximately 500bp amplicons) and ordered from Thermo Fisher

Scientific. A total of 9 human SNP loci were included: *FOXP2* rs6980093, and *OXTR*

rs2270463, rs237877, rs237878, rs35062132, rs2254295, rs237894, rs237895, and

rs237900 (Table 1).

Table 1: Selected Human SNPs

Single Nucleotide Polymorphism	Position in Human Reference Genome	Allele Frequency in Humans	Variant Type
FOXP2 rs6980093	chr7:114522685	G (.41) / A (.59)	Intron
OXTR rs2270463	chr3:8733391	G (.77) / T (.23)	Intron
OXTR rs237877	chr3:8741201	C (.69) / T (.31)	Intron
OXTR rs237878	chr3:8741312	T (.78) / C (.22) / A (.00)	Intron
OXTR rs35062132	chr3:8753201	G (1.00) / A (.00) / C (.00)	Missense
OXTR rs2254295	chr3:8760606	T (.87) / C (.13)	Intron
OXTR rs237894	chr3:8764845	G (.76) / C (.24)	Intron
OXTR rs237895	chr3:8765737	T (.37) / C (.63)	Intron
OXTR rs237900	chr3:8767010	G (.64) / A (.36)	Intron

Table 1: The name, position, relative allele frequency, and variant type for the selected

human single nucleotide polymorphisms.

DNA was extracted from bonobo whole blood and buccal samples, amplified by

PCR, resolved by gel electrophoresis, gel purified (Zymo Gel DNA Recovery Kit) and

sent to Genewiz for Sanger sequencing (see Supplementary Material 1 for details). To

determine SNP presence, individual Sanger sequences were aligned to the bonobo

reference genome using ApE and the human reference genome [Dec. 2013

(GRCh38/hg38)] using the BLAT tool[43]. Heterozygotes were identified by visual

inspection of the Sanger chromatogram and confirmed by sequencing the reverse

strand.

Behavioral Data

To determine if any observed genetic variation was linked to social behavior, we

utilized previously collected observational data that were available for 12 of the 13

subjects[34,44]. In short, eight 10-minute focal observations (i.e., observing the behavior of

a single individual) from each subject were used to assess group differences in social

proximity – an established method for measuring social relationships in nonhuman

primates that encapsulates both social interest and engagement[45,46,47].

Statistical Analyses

For each observation, a social proximity score (ranging from 0-3), was calculated

using the following formula, where 11 is the total number of proximity data points per

focal follow:

$$\frac{(3*\text{N touching data points})+(2*\text{N socially close data points})+(1*\text{N solitary data points})+(0*\text{N isolated data points})}{(11-\text{N cannot be determined data points})}$$

A Kruskal-Wallis H test and a Wilcoxon rank test were utilized to assess genetic

differences in social behavior.

Results

Genetic Variation

Of the 29 biological samples, 13 were of sufficient quality for Sanger sequencing

(whole blood n=7, buccal swab n=6). See Supplementary Material 2 for the coefficients

of relatedness between each subject. Analyses revealed a novel SNP in bonobo

FOXP2, 75bp to the right of *FOXP2* rs6980093 (human chr7:114522685; A/G). Three

bonobos were heterozygous at this location (G/A), while the rest were homozygous

(A/A; bonobo 5' flanking sequence CACTCGTATCACATTATAAT A/G; Figure 1). Both

genotypes differ from the bonobo reference genome (G/G).

In addition, analyses revealed genetic variation in bonobo OXTR at four novel

loci (Figure 2A). Specifically, we identified a novel SNP 78bp to the left of rs237877

(chr3:8741201; C/T) and 184bp left of rs237878 (chr3:8741312; T/A/C); 12 of the 13

bonobos were homozygous (T/T) and 1 individual was heterozygous (T/G; 5' flanking

sequence TTGCAGCTATCACCTCATTT T/G). We also identified a novel SNP 19bp to

the right of human SNP rs35062132 (chr3:8753201; G/A/C). In our sample, 11 bonobos

were homozygous (G/G), and 2 bonobos were heterozygous (G/A; 5' flanking sequence

CGATGGCTCAGGACAAAGGA G/A). In addition, a novel SNP was observed 17bp to

the left of rs2254295 (chr3:8760606; T/C; Figure 2B-D) and adjacent to rs2254298

(chr3:8760542; G/A). Ten of the 13 bonobos in our sample were homozygous (T/T) and

3 bonobos were heterozygous (T/C; 5' flanking sequence

GGCACTGGATGAGGCTGCC T/C). The fourth novel SNP we observed in bonobo

OXTR was 19bp to the left of rs237900 (chr3:8767010; G/A). Analyses revealed 3

genotypes at this locus; 2 bonobos were homozygous (A/A), 2 bonobos were

heterozygous (C/A), and 9 bonobos were homozygous (C/C; 5' flanking sequence

GCCCAAGGACTGTGCTAAGG A/C). Collectively, we observed the same allelic

combination across the 4 novel *OXTR* SNPs (homozygous TGTC) in 6 of the 13

bonobos (Figure 2E). See Table 2 for a complete list of allele types and frequencies.

Table 2: Individual Sequencing Data

Bonobo ID	F14	F16	M14	F13	M10	F05	M13	F18	M15	M12	M18	F10	M19
Novel SNP FOXP2	A/A	A/A	A/A	A/A	A/A	G/A	A/A	A/A	G/A	G/A	G/A	G/A	A/A
Novel SNP OXTR #1	T/T	T/T	T/T	T/T	T/T	T/T	T/G	T/T	T/T	T/T	T/T	T/T	T/T
Novel SNP OXTR #2	G/G	G/A	G/G	G/G	G/G	G/G	G/A	G/G	G/G	G/G	G/G	G/G	G/G
Novel SNP OXTR #3	C/T	T/T	T/T	T/T	T/T	C/T	T/T	T/T	C/T	T/T	T/T	T/T	T/T
Novel SNP OXTR #4	C/C	A/A	C/C	C/C	C/C	C/C	A/A	C/A	C/C	C/A	C/C	C/C	C/C
FOXP2 rs6980093	A/A	A/A	A/A	A/A	A/A	A/A	A/A	A/A	A/A	A/A	A/A	A/A	A/A
OXTR rs2270463	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G
OXTR rs237877	T/T	T/T	T/T	T/T	T/T	T/T	T/T	T/T	T/T	T/T	T/T	T/T	T/T
OXTR rs237878	C/C	C/C	C/C	C/C	C/C	C/C	C/C	C/C	C/C	C/C	C/C	C/C	C/C
OXTR rs35062132	GG	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G
OXTR rs2254295	T/T	T/T	T/T	T/T	T/T	T/T	T/T	T/T	T/T	T/T	T/T	T/T	T/T
OXTR rs237894	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G
OXTR rs237895	C/C	C/C	C/C	C/C	C/C	C/C	C/C	C/C	C/C	C/C	C/C	C/C	C/C
OXTR rs237900	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G	G/G

<u>Table 2</u>: Individual sequencing data for the novel SNPs identified in bonobo FOXP2 and

OXTR and for the selected human loci.

<u>Behavioral Differences</u>

For *OXTR*, individual bonobos were grouped based on the number of *OXTR*

TGTC copies (Figure 2E) – zero (n=2), one (n=4), or two (n=6). Kruskal-Wallis results

indicated a significant *OXTR* group difference in social proximity score (H (2) = 7.2991,

p = 0.026; Figure 3; Table 3). Bonobos with zero copies of TGTC were less social (*Mdn*

= 1.45) than individuals with one copy of TGTC (*Mdn* = 2.50) and individuals with two

copies of TGTC (*Mdn* = 2.00). A post-hoc Wilcoxon rank test with Benjamini-Hochberg

adjustment revealed a significant difference in social proximity scores between bonobos

with zero TGTC copies and bonobos with one TGTC copy (p = 0.049), but not between

bonobos with zero copies and two copies (p = 0.339) or between one copy and two

copies (p = 0.510). For *FOXP2*, individuals were grouped based on whether they were

homozygous (A/A) or heterozygous (G/A) at the *FOXP2* SNP locus. Wilcoxon rank test

results indicated there was no significant *FOXP2* group difference in social proximity

score (W = 1164.5, p = 0.275).

<u>Table 3: Social Proximity Scores</u>

Bonobo ID	F14	F16	M14	F13	M10	F05	M13	F18	M12	M18	F10	M19
Proximity Score 1	2.73	1.70	1.18	0.64	1.82	3.00	3.00	1.64	0.00	0.55	3.00	2.36
Proximity Score 2	2.27	0.82	1.00	0.55	2.18	3.00	1.36	3.00	1.27	1.00	3.00	2.09
Proximity Score 3	3.00	1.45	1.09	0.82	0.64	2.73	2.09	3.00	0.00	0.00	3.00	2.00
Proximity Score 4	2.36	3.00	1.09	3.00	2.73	3.00	1.45	2.27	3.00	0.00	3.00	2.46
Proximity Score 5	1.45	2.18	1.91	2.82	1.64	3.00	0.27	3.00	0.45	0.45	3.00	2.18

Proximity Score 6	1.50	2.82	2.36	2.64	2.00	2.45	1.27	1.82	1.73	1.55	3.00	1.55
Proximity Score 7	2.90	1.00	2.00	0.82	0.00	3.00	0.36	2.91	2.18	2.18	2.45	2.55
Proximity Score 8	2.82	1.73	1.45	1.73	1.55	1.18	1.00	2.55	1.73	2.73	2.27	2.82

173 Table 3: Social proximity scores for each of the bonobo's focal observations.

Discussion

Determining the biological factors underlying individual-level social

communication is key to understanding autism and other neurodevelopmental disorders

and may aid in identifying children at risk of developing social communication

impairments. Studying biobehavioral mechanisms in captive, nonhuman animals,

permits researchers to elucidate the genetic contributions to complex behavioral

phenotypes, while minimizing confounding rearing and environmental factors. Thus, we

aimed to determine if one of the species most closely related to humans, the bonobo,

exhibits SNPs in *FOXP2* and *OXTR* at known human loci.

Despite whole genomic investigations in all four nonhuman species of great ape,

no SNPs have been identified in bonobo *FOXP2* to date. We are the first to report a

SNP in bonobo *FOXP2* (near rs6980093) and we identified two novel genotypes in our

sample. Although we did not find social behavior differences based on the bonobos'

FOXP genotypes, we are encouraged by the identification of variation in bonobo *FOXP2*

(75bp from the ASD-linked human SNP) and believe that further investigation into the

bonobo *FOXP2* gene and measures of bonobo vocal communication might yield

promising results.

The human *FOXP2* SNP rs6980093 is an intronic polymorphism, suggesting it

may be involved in regulating *FOXP2* expression[7,48]. Several studies have identified

relations between *FOXP2* rs6980093 variation and speech production[49], cortical

activation in language-related regions[50], as well as speech and language learning

abilities[7,51] The identification of variation in bonobo *FOXP2* is a foundational step in

understanding the impact of *FOXP2* on social communication in humans and our

closest living relatives. Bonobos have the largest vocal repertoire size of the nonhuman

great apes[52], and modify their communicative signals depending on social

context[53,54,55]. In addition, multiple reports in a single bonobo support the notion that

bonobos are able to understand spoken English words and sentences[56,57,58], as well as

degraded and computer-generated speech[59]. Thus, it is possible that *FOXP2* alleles

modulate *FOXP2* expression in bonobos and that differential *FOXP2* regulation impacts

individual variability in bonobo vocal communication.

Specific *OXTR* variants have been linked to social functioning[60] and symptom

severity[14,15] in autism. In our sample of bonobos, we identified a novel SNP between

human *OXTR* rs237877 and rs237878. Variants at these loci have been linked to level

of extraversion[8] and reward responsiveness[61] in typically developed adults, as well as

neurological responses to oxytocin in autistic adults[62]. In addition, we identified a novel

SNP in bonobo *OXTR* close to the human SNP rs35062132. Eleven of the 13 bonobos

in our sample exhibited the G/G genotype. In humans, the C/G genotype of *OXTR*

rs35062132 was associated with an increased risk of ASD and proposed to be a

biological basis for individual differences in social behavior[63]. Although they identified

three genotypes in their sample, Egawa and colleagues did not find a relation between

these genetic variants and ASD[64]. Hence, further investigations are needed to

understand the role rs35062132 variation plays in social communication development

and to determine if the rs35062132 G allele is a risk factor for autism[63,64].

We also found novel SNPs in bonobo *OXTR* between rs2254295 and rs2254298

and adjacent to rs237900. Links between these genetic variants and social functioning

are a well-established finding, particularly for rs2254298[65,66,67,68]. For example,

rs2254295 and rs2254298 variants were associated with nonverbal communication

scores in Japanese adult males diagnosed with ASD[11]. Yang and colleagues found

differences in serum oxytocin levels based on rs2254298 genotype[11]. However, the role

that specific rs2254298 variants play in autism remains unclear. Specifically, results

differ among human populations; the "A" allele was associated with autism in

Japanese[69] and Chinese[70] populations, whereas the "G" allele was considered a risk

factor in a Caucasian population of autistic children and adolescents[71]. Parker and

colleagues also identified links between rs2254298 variants and social impairments in

children with and without ASD[72]. In both groups, individuals with the "A" allele exhibited

greater social impairments than individuals without the "A" allele. Collectively, these

findings suggest that specific *OXTR* variants might be promising biomarkers for social

communication dysfunction in humans and highlight the need for alternative approaches

to assessing the impact *OXTR* variants have on complex behavioral phenotypes, like

those observed in ASD[73,74].

Most notably, we observed the same allelic combination across the four novel

OXTR SNPs (homozygous TGTC) in six of the 13 bonobos – demonstrating linkage

between these *OXTR* variants. Although we did not find variability at the selected

human loci, the high prevalence of the homozygous TGTC genotype suggests that

these related variants influence individual-level social communication in bonobos.

Linked *OXTR* variants have also been found in children, adolescents, and young adults

diagnosed with ASD[12]. In addition, Wu and colleagues identified linkage among two

OXTR variants in autistic people from the Chinese Han population[70]. Thus, we

encourage researchers interested in the biomarkers of human social communication

disorders to consider the relative influence of individual SNPs and their collective

contribution to complex behavioral phenotypes.

If the observed TGTC combination was selected for in bonobos, we would expect

to see behavioral differences based on these genotypes. To test this hypothesis, we

utilized previously collected observational data that were available for 12 of the bonobos

in our sample. By grouping bonobos based on the number of TGTC copies (zero, one,

or two), we were able to investigate genetic differences in social behavior. Interestingly,

bonobos with zero copies of the TGTC combination had lower social proximity scores

(i.e., they spent less time in close proximity to conspecific social partners) than bonobos

with one copy of the TGTC combination. This result supports previous conclusions that

OXTR plays a pivotal role in bonobo social behavior[9,23,38] and is consistent with findings

in humans that indicate that linked *OXTR* variants are associated with greater

impairments on the social responsiveness and repetitive behavior scales in autistic

children[75]. Our results are also similar to data collected from children, adolescents, and

young adults diagnosed with ASD; a haplotype comprised of four *OXTR* loci was

associated with greater impairments in social interaction and communication in autistic

individuals[12]. These collective findings highlight the importance of considering multiple

genetic variants in a given study and the benefits of multi-loci investigations in

nonhuman great apes. All told, our results suggest that the *OXTR* TGTC combination

was selected for in bonobos.

Given the documented relations between *OXTR* and social functioning,

researchers have proposed that oxytocin can help facilitate social information

processing in individuals with ASD. Indeed, evidence exists that oxytocin treatment can

improve social abilities in children diagnosed with ASD[76] and that oxytocin infusions can

reduce RRBs in autistic adults[62,77]. Researchers have also determined that oxytocin

treatment efficacy differs between people, with the greatest improvements to social

behavior occurring in individuals with the lowest pretreatment oxytocin blood

concentration levels[76]. Along with previous evidence of a low social, high RRB

phenotype in bonobos[34], our findings suggest that bonobos are an exemplary species

for evaluating the efficacy of oxytocin interventions in the treatment of social

communication dysfunction. Notwithstanding, further studies are needed to determine

the specific characteristics that impact oxytocin efficacy and to identify biomarkers that

predict which individuals will benefit the most from oxytocin treatments[62,76].

Seminal work in rodents accounts for much of what we know about *OXTR* and

other ASD-related genes. However, many of these studies require substantial

modification of the gene and/or the receptors or are limited in their translatability to the

complex phenotype of ASD[26,28,30]. In this study, we identified a naturally occurring

linkage among 4 novel *OXTR* variants and documented differences in bonobo social

behavior based on this combined *OXTR* genotype. We also demonstrated that it is

possible to detect genetic variability, variant linkage, and behavioral differences in even

small samples of nonhuman great apes. Thus, we encourage the incorporation of

bonobos in future investigations and emphasize the need for a publicly accessible

database to report SNPs identified in nonhuman primates.

Here, we are the first to report a SNP in bonobo *FOXP2* – a gene necessary for

typical linguistic development in humans. We also identified four novel SNPs in bonobo

OXTR and demonstrated linkage among these *OXTR* variants. Our results indicate that

individuals without the *OXTR* TGTC combination are less social than individuals with

one copy of the TGTC combination. Our collective findings suggest that these *OXTR*

variants influence individual-level social communication in bonobos and support the

notion that linked *OXTR* variants could be promising biological factors for identifying

humans at risk of developing social communication deficits. This study also highlights

the advantages of studying biobehavioral mechanisms in the species most closely

related to humans and indicates that bonobos are a suitable model for testing

hypotheses about the etiology of ASD and other human neurodevelopmental disorders.

Acknowledgements

This work was supported in part by NIH grants R15GM140472-01 (MLH),

T34GM140948-01A1 (MLH), and R25GM111565 (AB), Autism Speaks Postdoctoral

Fellowship 13439 (SAS), and generous donations from Ape Initiative. Approval was

obtained from each institution and all procedures followed the institution's animal care

and use protocols (Ape Initiative IACUC Protocol #210305-01). Thank you to Ape

Initiative, the Columbus Zoo and Aquarium, and the Milwaukee County Zoo for

providing biological samples. We would like to acknowledge that bonobos are an

endangered species, with human activity as the greatest threat to their survival.

Figures

<u>Figure 1</u>

<u>Figure 1</u>: Identification of a single-nucleotide polymorphism in bonobo *FOXP2* (A)

Diagram of the human *FOXP2* gene showing relative location of the SNP rs6980093.

Arrow shows direction of gene transcription. Short vertical bars (grey) show

transcriptional start sites. Black vertical bars show exons. (B) Alignment of the human

and bonobo references genomes, along with representative sequencing data from

subjects in this study. Numbers refer to the human reference genome location on

chromosome seven. (C, D) Representative Sanger sequencing chromatograms across

the SNP, showing (C) a homozygous sample (Female 14), and (D) a heterozygous

sample (Female 10).

<u>Figure 2</u>

<u>Figure 2</u>: Identification of single-nucleotide polymorphisms in bonobo *OXTR* (A)

Diagram of the human *OXTR* gene showing relative location of SNPs rs237877,

rs237878, rs35062132, rs2254298, rs2254295, and rs237900, as well as the novel

bonobo SNPs. Arrow shows direction of gene transcription. Untranslated regions are

shown in grey. Black oblongs show exons. (B) Alignment of the human and bonobo

references genomes, along with representative sequencing data from subjects in this

study. Numbers refer to the human reference genome location on chromosome three.

(C, D) Representative Sanger sequencing chromatograms across the novel SNP,

showing (C) a homozygous sample (Female 10), and (D) a heterozygous sample

(Female 14). (E) Pedigree of the 13 subjects, with corresponding *OXTR* genotypes at

the four novel SNPs in bonobo *OXTR*.

<u>Figure 3</u>

<u>Figure 3</u>: Observed group differences in social behavior based on the number of *OXTR*

TGTC copies. Average social proximity scores with corresponding standard error bars

(A) and raw social proximity scores (B) for bonobos with zero, one, and two copies of

the TGTC combination.

References

1. *Human Gene module. SFARI Gene. https://gene.sfari.org/database/human-gene/*
2. Abrahams BS, Arking DE, Campbell DB, Mefford HC, Morrow EM, Weiss LA, Menashe I, Wadkins T, Banerjee-Basu S, Packer A. SFARI Gene 2.0: a community-driven knowledgebase for the autism spectrum disorders (ASDs). Molecular Autism. 2013 Dec;4(1):1-3. https://doi.org/10.1186/2040-2392-4-36
3. Fisher SE, Scharff C. FOXP2 as a molecular window into speech and language. Trends in Genetics. 2009 Apr 1;25(4):166-77. https://doi.org/10.1016/j.tig.2009.03.002
4. Lai CS, Fisher SE, Hurst JA, Vargha-Khadem F, Monaco AP. A forkhead-domain gene is mutated in a severe speech and language disorder. Nature. 2001 Oct 4;413(6855):519-23. https://doi.org/10.1038/35097076
5. Shriberg LD, Ballard KJ, Tomblin JB, Duffy JR, Odell KH, Williams CA. Speech, prosody, and voice characteristics of a mother and daughter with a 7;13 translocation affecting *FOXP2*. J Speech Lang Hear Res [Internet]. 2006;49(3):500–25. http://dx.doi.org/10.1044/1092-4388(2006/038)
6. Haghighatfard A, Yaghoubi asl E, Bahadori RA, Aliabadian R, Farhadi M, Mohammadpour F, Tabrizi Z. FOXP2 down expression is associated with executive dysfunctions and electrophysiological abnormalities of brain in Autism spectrum disorder; a neuroimaging genetic study. Autism & Developmental Language Impairments. 2022 Sep;7:23969415221126391. https://doi.org/10.1177/23969415221126391
7. Mozzi A, Riva V, Forni D, Sironi M, Marino C, Molteni M, Riva S, Guerini FR, Clerici M, Cagliani R, Mascheretti S. A common genetic variant in FOXP2 is associated with language-based learning (dis) abilities: Evidence from two Italian independent samples. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics. 2017 Jul;174(5):578-86. https://doi.org/10.1002/ajmg.b.32546
8. Haram M, Tesli M, Dieset I, Steen NE, Røssberg JI, Djurovic S, Andreassen OA, Melle I. An attempt to identify single nucleotide polymorphisms contributing to possible relationships between personality traits and oxytocin-related genes. Neuropsychobiology. 2014 Jan 22;69(1):25-30. https://doi.org/10.1159/000356965
9. Theofanopoulou C, Andirkó A, Boeckx C, Jarvis ED. Oxytocin and vasotocin receptor variation and the evolution of human prosociality. Comprehensive Psychoneuroendocrinology. 2022 Aug 1;11:100139. https://doi.org/10.1016/j.cpnec.2022.100139
10. Wu N, Su Y. Variations in the oxytocin receptor gene and prosocial behavior: moderating effects of situational factors. Integrative Zoology. 2018 Nov;13(6):687-97. https://doi.org/10.1111/1749-4877.12336
11. Yang S, Dong X, Guo X, Han Y, Song H, Gao L, Dai W, Su Y, Zhang X. Serum oxytocin levels and an oxytocin receptor gene polymorphism (rs2254298) indicate social deficits in children and adolescents with autism spectrum disorders. Frontiers in Neuroscience. 2017 Apr 21;11:221. https://doi.org/10.3389/fnins.2017.00221

12. Wermter AK, Kamp-Becker I, Hesse P, Schulte-Körne G, Strauch K, Remschmidt H. Evidence for the involvement of genetic variation in the oxytocin receptor gene (OXTR) in the etiology of autistic disorders on high-functioning level. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics. 2010 Mar;153(2):629-39. https://doi.org/10.1002/ajmg.b.31032

13. Holmqvist Jämsen S, Johansson A, Westberg L, Santtila P, von der Pahlen B, Simberg S. Associations between vocal symptoms and genetic variants in the oxytocin receptor and arginine vasopressin 1A receptor gene. Journal of Speech, Language, and Hearing Research. 2017 Jul 12;60(7):1843-54. https://doi.org/10.1044/2016_JSLHR-S-16-0059

14. Oztan O, Jackson LP, Libove RA, Sumiyoshi RD, Phillips JM, Garner JP, Hardan AY, Parker KJ. Biomarker discovery for disease status and symptom severity in children with autism. Psychoneuroendocrinology. 2018 Mar 1;89:39-45. https://doi.org/10.1016/j.psyneuen.2017.12.022

15. Cataldo I, Azhari A, Esposito G. A review of oxytocin and arginine-vasopressin receptors and their modulation of autism spectrum disorder. Frontiers in Molecular Neuroscience. 2018 Feb 13;11:27. https://doi.org/10.3389/fnmol.2018.00027

16. De Abreu MS, Genario R, Giacomini AC, Demin KA, Lakstygal AM, Amstislavskaya TG, Fontana BD, Parker MO, Kalueff AV. Zebrafish as a model of neurodevelopmental disorders. Neuroscience. 2020 Oct 1;445:3-11. https://doi.org/10.1016/j.neuroscience.2019.08.034

17. Crawley JN. Translational animal models of autism and neurodevelopmental disorders. Dialogues in Clinical Neuroscience. 2022 Apr 1. https://doi.org/10.31887/DCNS.2012.14.3/jcrawley

18. Eaton SL, Wishart TM. Bridging the gap: large animal models in neurodegenerative research. Mammalian Genome. 2017 Aug;28:324-37. https://doi.org/10.1007/s00335-017-9687-6

19. Chabout J, Sarkar A, Patel SR, Radden T, Dunson DB, Fisher SE, Jarvis ED. A Foxp2 mutation implicated in human speech deficits alters sequencing of ultrasonic vocalizations in adult male mice. Frontiers in Behavioral Neuroscience. 2016 Oct 20;10:197. https://doi.org/10.3389/fnbeh.2016.00197

20. Medvedeva VP, Rieger MA, Vieth B, Mombereau C, Ziegenhain C, Ghosh T, Cressant A, Enard W, Granon S, Dougherty JD, Groszer M. Altered social behavior in mice carrying a cortical Foxp2 deletion. Human Molecular Genetics. 2019 Mar 1;28(5):701-17. https://doi.org/10.1093/hmg/ddy372

21. Gemmer A, Mirkes K, Anneser L, Eilers T, Kibat C, Mathuru A, Ryu S, Schuman E. Oxytocin receptors influence the development and maintenance of social behavior in zebrafish (*Danio rerio*). Scientific Reports. 2022 Mar 12;12(1):4322. https://doi.org/10.1038/s41598-022-07990-y

22. Staes N, Sherwood CC, Wright K, De Manuel M, Guevara EE, Marques-Bonet T, Krützen M, Massiah M, Hopkins WD, Ely JJ, Bradley BJ. FOXP2 variation in great ape populations offers insight into the evolution of communication skills. Scientific Reports. 2017 Dec 4;7(1):16866. https://doi.org/10.1038/s41598-017-16844-x

23. Staes N, Stevens JM, Helsen P, Hillyer M, Korody M, Eens M. Oxytocin and
vasopressin receptor gene variation as a proximate base for inter-and
intraspecific behavioral differences in bonobos and chimpanzees. PLoS One.
2014 Nov 18;9(11):e113364. https://doi.org/10.1371/journal.pone.0113364
24. Heston JB, White SA. Behavior-linked FoxP2 regulation enables zebra finch
vocal learning. Journal of Neuroscience. 2015 Feb 18;35(7):2885-94.
https://doi.org/10.1523/JNEUROSCI.3715-14.2015
25. Donaldson ZR, Young LJ. The relative contribution of proximal 5′ flanking
sequence and microsatellite variation on brain vasopressin 1a receptor (Avpr1a)
gene expression and behavior. PLoS Genetics. 2013 Aug 29;9(8):e1003729.
https://doi.org/10.1371/journal.pgen.1003729
26. Fischer J, Hammerschmidt K. Ultrasonic vocalizations in mouse models for
speech and socio-cognitive disorders: insights into the evolution of vocal
communication. Genes, Brain and Behavior. 2011 Feb;10(1):17-27.
https://doi.org/10.1111/j.1601-183X.2010.00610.x
27. Johnson ZV, Walum H, Xiao Y, Riefkohl PC, Young LJ. Oxytocin receptors
modulate a social salience neural network in male prairie voles. Hormones and
Behavior. 2017 Jan 1;87:16-24. https://doi.org/10.1016/j.yhbeh.2016.10.009
28. Pobbe RL, Pearson BL, Blanchard DC, Blanchard RJ. Oxytocin receptor and
Mecp2308/Y knockout mice exhibit altered expression of autism-related social
behaviors. Physiology & Behavior. 2012 Dec 5;107(5):641-8.
https://doi.org/10.1016/j.physbeh.2012.02.024
29. Shu W, Cho JY, Jiang Y, Zhang M, Weisz D, Elder GA, Schmeidler J, De
Gasperi R, Sosa MA, Rabidou D, Santucci AC. Altered ultrasonic vocalization in
mice with a disruption in the Foxp2 gene. Proceedings of the National Academy
of Sciences. 2005 Jul 5;102(27):9643-8.
https://doi.org/10.1073/pnas.0503739102
30. Bey AL, Jiang YH. Overview of mouse models of autism spectrum disorders.
Current Protocols in Pharmacology. 2014 Sep;66(1):5-66.
https://doi.org/10.1002/0471141755.ph0566s66
31. Silverman JL, Yang M, Lord C, Crawley JN. Behavioural phenotyping assays for
mouse models of autism. Nature Reviews Neuroscience. 2010 Jul;11(7):490-
502. https://doi.org/10.1038/nrn2851
32. Berendzen KM, Sharma R, Mandujano MA, Wei Y, Rogers FD, Simmons TC,
Seelke AM, Bond JM, Larios R, Goodwin NL, Sherman M. Oxytocin receptor is
not required for social attachment in prairie voles. Neuron. 2023 Mar
15;111(6):787-96. https://doi.org/10.1016/j.neuron.2022.12.011
33. Surbeck M, Hohmann G. Social preferences influence the short-term exchange
of social grooming among male bonobos. Animal Cognition. 2015 Mar;18:573-9.
https://doi.org/10.1007/s10071-014-0826-0
34. Skiba, SA. Behavioral, cognitive, and genetic factors underlying socio-
communicative development in bonobos. Dissertation, Georgia State University;
2021 Jul 23. https://doi.org/10.57709/23988841
35. Pika S, Liebal K, Tomasello M. Gestural communication in subadult bonobos
(Pan paniscus): repertoire and use. American Journal of Primatology: Official

Journal of the American Society of Primatologists. 2005 Jan;65(1):39-61. https://doi.org/10.1002/ajp.20096

36. Schamberg I, Cheney DL, Clay Z, Hohmann G, Seyfarth RM. Call combinations, vocal exchanges and interparty movement in wild bonobos. Animal Behaviour. 2016 Dec 1;122:109-16. https://doi.org/10.1016/j.anbehav.2016.10.003

37. Laméris DW, Staes N, Salas M, Matthyssen S, Verspeek J, Stevens JM. The influence of sex, rearing history, and personality on abnormal behaviour in zoo-housed bonobos (Pan paniscus). Applied Animal Behaviour Science. 2021 Jan 1;234:105178. https://doi.org/10.1016/j.applanim.2020.105178

38. Kovalaskas S, Rilling JK, Lindo J. Comparative analyses of the Pan lineage reveal selection on gene pathways associated with diet and sociality in bonobos. Genes, Brain and Behavior. 2021 Mar;20(3):e12715. https://doi.org/10.1111/gbb.12715

39. Staes N, Guevara EE, Helsen P, Eens M, Stevens JM. The Pan social brain: An evolutionary history of neurochemical receptor genes and their potential impact on sociocognitive differences. Journal of Human Evolution. 2021 Mar 1;152:102949. https://doi.org/10.1016/j.jhevol.2021.102949

40. Moscovice LR, Surbeck M, Fruth B, Hohmann G, Jaeggi AV, Deschner T. The cooperative sex: sexual interactions among female bonobos are linked to increases in oxytocin, proximity and coalitions. Hormones and Behavior. 2019 Nov 1;116:104581. https://doi.org/10.1016/j.yhbeh.2019.104581

41. Primer designing tool. Nih.gov. https://www.ncbi.nlm.nih.gov/tools/primer-blast/

42. ApE- A plasmid Editor. Utah.edu. https://jorgensen.biology.utah.edu/wayned/ape/

43. Kent WJ. BLAT—the BLAST-like alignment tool. Genome research. 2002 Apr1;12(4):656-64. http://www.genome.org/cgi/doi/10.1101/gr.229202.

44. Skiba SA. The Adaptive Value of Complex Socio-Communicative Behavior. Thesis. Kennesaw State University; 2017 Jul 14. https://digitalcommons.kennesaw.edu/integrbiol_etd/23/

45. Furuichi T. Social interactions and the life history of female Pan paniscus in Wamba, Zaire. International journal of Primatology. 1989 Jun;10:173-97. https://doi.org/10.1007/BF02735199

46. Gelardi V, Godard J, Paleressompoulle D, Claidiere N, Barrat A. Measuring social networks in primates: wearable sensors versus direct observations. Proceedings of the Royal Society A. 2020 Apr 29;476(2236):20190737. https://doi.org/10.1098/rspa.2019.0737

47. Taglialatela JP, Skiba SA, Evans RE, Bogart S, Schwob NG. Social behavior and social tolerance in chimpanzees and bonobos. Chimpanzee in Context: A Comparative Perspecitve on Chimpanzee Behavior, Congitin, Conservation, and Welfare. 2020:95-114. https://doi.org/10.7208/9780226728032-007

48. Cooper DN. Functional intronic polymorphisms: Buried treasure awaiting discovery within our genes. Human Genomics. 2010 Dec;4(5):1-5. https://doi.org/10.1186/1479-7364-4-5-284

49. Zhang S, Zhao J, Guo Z, Jones JA, Liu P, Liu H. The association between genetic variation in FOXP2 and sensorimotor control of speech production. Frontiers in Neuroscience. 2018 Sep 20;12:666. https://doi.org/10.3389/fnins.2018.00666

50. Pinel P, Fauchereau F, Moreno A, Barbot A, Lathrop M, Zelenika D, Le Bihan D,
Poline JB, Bourgeron T, Dehaene S. Genetic variants of FOXP2 and
KIAA0319/TTRAP/THEM2 locus are associated with altered brain activation in
distinct language-related regions. Journal of Neuroscience. 2012 Jan
18;32(3):817-25. https://doi.org/10.1523/JNEUROSCI.5996-10.2012
51. Chandrasekaran B, Yi HG, Blanco NJ, McGeary JE, Maddox WT. Enhanced
procedural learning of speech sound categories in a genetic variant of FOXP2.
Journal of Neuroscience. 2015 May 20;35(20):7808-12.
https://doi.org/10.1523/JNEUROSCI.4706-14.2015
52. McComb K, Semple S. Coevolution of vocal communication and sociality in
primates. Biology Letters. 2005 Dec 22;1(4):381-5.
https://doi.org/10.1098/rsbl.2005.0366
53. Clay Z, Archbold J, Zuberbühler K. Functional flexibility in wild bonobo vocal
behaviour. PeerJ. 2015 Aug 4;3:e1124. https://doi.org/10.7717/peerj.1124
54. Clay Z, Zuberbühler K. Communication during sex among female bonobos:
effects of dominance, solicitation and audience. Scientific Reports. 2012 Mar
1;2(1):291. https://doi.org/10.1038/srep00291
55. Hohmann G, Fruth B. Structure and use of distance calls in wild bonobos (Pan
paniscus). International Journal of Primatology. 1994 Oct;15:767-82.
https://doi.org/10.1007/BF02737430
56. Margiotoudi K, Bohn M, Schwob N, Taglialatela J, Pulvermüller F, Epping A,
Schweller K, Allritz M. Bo-NO-bouba-kiki: picture-word mapping but no
spontaneous sound symbolic speech-shape mapping in a language trained
bonobo. Proceedings of the Royal Society B. 2022 Feb 9;289(1968):20211717.
https://royalsocietypublishing.org/doi/full/10.1098/rspb.2021.1717
57. Brakke KE, Savage-Rumbaugh ES. The development of language skills in
bonobo and chimpanzee: I. Comprehension. Language & Communication. 1995
Apr. https://psycnet.apa.org/doi/10.1016/0271-5309(95)00001-7
58. Savage-Rumbaugh ES, Murphy J, Sevcik RA, Brakke KE, Williams SL,
Rumbaugh DM, Bates E. Language comprehension in ape and child.
Monographs of the Society for Research in Child Development. 1993 Jan 1:i-252.
https://doi.org/10.2307/1166068
59. Lahiff NJ, Slocombe KE, Taglialatela J, Dellwo V, Townsend SW. Degraded and
computer-generated speech processing in a bonobo. Animal Cognition. 2022
Dec;25(6):1393-8. https://doi.org/10.1007/s10071-022-01621-9
60. Baribeau DA, Dupuis A, Paton TA, Scherer SW, Schachar RJ, Arnold PD,
Szatmari P, Nicolson R, Georgiades S, Crosbie J, Brian J. Oxytocin receptor
polymorphisms are differentially associated with social abilities across
neurodevelopmental disorders. Scientific Reports. 2017 Sep 14;7(1):11618.
https://doi.org/10.1038/s41598-017-10821-0
61. Davis C, Zai CC, Adams N, Bonder R, Kennedy JL. Oxytocin and its association
with reward-based personality traits: A multilocus genetic profile (MLGP)
approach. Personality and Individual Differences. 2019 Feb 1;138:231-6.
https://doi.org/10.1016/j.paid.2018.09.002
62. Watanabe T, Otowa T, Abe O, Kuwabara H, Aoki Y, Natsubori T, Takao H,
Kakiuchi C, Kondo K, Ikeda M, Iwata N. Oxytocin receptor gene variations predict

neural and behavioral response to oxytocin in autism. Social Cognitive and Affective Neuroscience. 2017 Mar;12(3):496-506. https://doi.org/10.1093/scan/nsw150

63. Ma WJ, Hashii M, Munesue T, Hayashi K, Yagi K, Yamagishi M, Higashida H, Yokoyama S. Non-synonymous single-nucleotide variations of the human oxytocin receptor gene and autism spectrum disorders: a case–control study in a Japanese population and functional analysis. Molecular Autism. 2013 Dec;4(1):1-4. https://doi.org/10.1186/2040-2392-4-22

64. Egawa J, Watanabe Y, Shibuya M, Endo T, Sugimoto A, Igeta H, Nunokawa A, Inoue E, Someya T. Resequencing and association analysis of OXTR with autism spectrum disorder in a Japanese population. Psychiatry and Clinical Neurosciences. 2015 Mar;69(3):131-5. https://doi.org/10.1111/pcn.12205

65. Bozorgmehr A, Alizadeh F, Sadeghi B, Shahbazi A, Ofogh SN, Joghataei MT, Razian S, Heydari F, Ghadirivasfi M. Oxytocin moderates risky decision-making during the Iowa gambling task: A new insight based on the role of oxytocin receptor gene polymorphisms and interventional cognitive study. Neuroscience Letters. 2019 Aug 24;708:134328. https://doi.org/10.1016/j.neulet.2019.134328

66. Costa B, Pini S, Gabelloni P, Abelli M, Lari L, Cardini A, Muti M, Gesi C, Landi S, Galderisi S, Mucci A. Oxytocin receptor polymorphisms and adult attachment style in patients with depression. Psychoneuroendocrinology. 2009 Nov 1;34(10):1506-14. https://doi.org/10.1016/j.psyneuen.2009.05.006

67. Thompson RJ, Parker KJ, Hallmayer JF, Waugh CE, Gotlib IH. Oxytocin receptor gene polymorphism (rs2254298) interacts with familial risk for psychopathology to predict symptoms of depression and anxiety in adolescent girls. Psychoneuroendocrinology. 2011 Jan 1;36(1):144-7. https://doi.org/10.1016/j.psyneuen.2010.07.003

68. Wu N, Li Z, Su Y. The association between oxytocin receptor gene polymorphism (OXTR) and trait empathy. Journal of Affective Disorders. 2012 May 1;138(3):468-72. https://doi.org/10.1016/j.jad.2012.01.009

69. Liu X, Kawamura Y, Shimada T, Otowa T, Koishi S, Sugiyama T, Nishida H, Hashimoto O, Nakagami R, Tochigi M, Umekage T. Association of the oxytocin receptor (OXTR) gene polymorphisms with autism spectrum disorder (ASD) in the Japanese population. Journal of Human Genetics. 2010 Mar;55(3):137-41. https://doi.org/10.1038/jhg.2009.140

70. Wu S, Jia M, Ruan Y, Liu J, Guo Y, Shuang M, Gong X, Zhang Y, Yang X, Zhang D. Positive association of the oxytocin receptor gene (OXTR) with autism in the Chinese Han population. Biological Psychiatry. 2005 Jul 1;58(1):74-7. https://doi.org/10.1016/j.biopsych.2005.03.013

71. Jacob S, Brune CW, Carter CS, Leventhal BL, Lord C, Cook Jr EH. Association of the oxytocin receptor gene (OXTR) in Caucasian children and adolescents with autism. Neuroscience letters. 2007 Apr 24;417(1):6-9. https://doi.org/10.1016/j.neulet.2007.02.001

72. Parker KJ, Garner JP, Libove RA, Hyde SA, Hornbeak KB, Carson DS, Liao CP, Phillips JM, Hallmayer JF, Hardan AY. Plasma oxytocin concentrations and OXTR polymorphisms predict social impairments in children with and without

autism spectrum disorder. Proceedings of the National Academy of Sciences.
2014 Aug 19;111(33):12258-63. https://doi.org/10.1073/pnas.1402236111
73. LoParo D, Waldman ID. The oxytocin receptor gene (OXTR) is associated with
autism spectrum disorder: a meta-analysis. Molecular Psychiatry. 2015
May;20(5):640-6. https://doi.org/10.1038/mp.2014.77
74. Ylisaukko-oja T, Alarcón M, Cantor RM, Auranen M, Vanhala R, Kempas E, von
Wendt L, Järvelä I, Geschwind DH, Peltonen L. Search for autism loci by
combined analysis of Autism Genetic Resource Exchange and Finnish families.
Annals of Neurology. 2006 Jan;59(1):145-55. https://doi.org/10.1002/ana.20722
75. Harrison AJ, Gamsiz ED, Berkowitz IC, Nagpal S, Jerskey BA. Genetic variation
in the oxytocin receptor gene is associated with a social phenotype in autism
spectrum disorders. American Journal of Medical Genetics Part B:
Neuropsychiatric Genetics. 2015 Dec;168(8):720-9.
https://doi.org/10.1002/ana.20722
76. Parker KJ, Oztan O, Libove RA, Sumiyoshi RD, Jackson LP, Karhson DS,
Summers JE, Hinman KE, Motonaga KS, Phillips JM, Carson DS. Intranasal
oxytocin treatment for social deficits and biomarkers of response in children with
autism. Proceedings of the National Academy of Sciences. 2017 Jul
25;114(30):8119-24. https://doi.org/10.1073/pnas.1705521114
77. Hollander E, Novotny S, Hanratty M, Yaffe R, DeCaria CM, Aronowitz BR,
Mosovich S. Oxytocin infusion reduces repetitive behaviors in adults with autistic
and Asperger's disorders. Neuropsychopharmacology. 2003 Jan;28(1):193-8.
https://doi.org/10.1038/sj.npp.1300021

 Supplementary Material 1: Genetic Analyses

Bonobo DNA extraction was completed using the PureLink® Genomic DNA Mini Kit. For
whole blood samples, 3 ml of blood was collected from each bonobo and stored in a
freezer at -20 degrees Celsius. The blood lysate started as a combination of 200 µl of
frozen whole blood samples, 20 µl Proteinase K, and 20 µl RNase A in a 1.5 ml
microcentrifuge tube. The tube was then vortexed briefly and incubated at room
temperature for 2 minutes. Next, 200 µl PureLink® Genomic Lysis /Binding Buffer was
added to the lysate, and the lysate was vortexed until homogeneity. Once the lysate
was homogenous, it was incubated at 55 degrees Celsius for 10 minutes in a hot bead
bath to promote protein digestion. Then 200 µl 96-100% ethanol was added to the
lysate, and vortexed by 5 seconds to yield a homogenous solution. To ensure DNA
binding conditions were still met while removing salt and protein contaminants, 96-100%
ethanol was added to PureLink® Genomic Wash Buffer 1 and 2. Next, the lysate
prepared with the genomic lysis/binding buffer and ethanol (~ 640 µl) was transferred to
a PureLink® Spin Column in a collection tube. The column was then centrifuged at
10,000 x g for 1 minute at room temperature. The collection tube was discarded, and
the spin column was placed in a new sterile collection tube.

For buccal swab samples, the Qiagen Omni Swab tips were ejected and placed in their
own 2 ml microcentrifuge tube, and then 600 µl of phosphate buffered saline (PBS) was
added to the tube. Next, 20 µl of Proteinase K was added to a large 15 ml centrifuge
tube, as well as 600 µl of swab lysate, which were mixed well via pipetting. Afterwards,
600 µl of PureLink® Genomic Lysis/Binding Buffer was added to the lysate and mixed
by vortexing. The lysate was incubated at 55 degrees Celsius in a hot bead bath for 15
minutes. The tube was centrifuged to collect any leftover lysate that may be on the cap,
and then 200 µl of 96-100% ethanol was added to the lysate and vortexed for 5 seconds
to get a homogenous solution.

For DNA washing, 500 µl of ethanol prepared wash buffer 1 was added to the spin
column. Then the column was centrifuged at 10,000 x g at room temperature for 1
minute. The collection tube was discarded afterwards and replaced with a new sterile
tube. The DNA was washed a second time with ethanol prepared wash buffer 2 and
centrifuged at maximum speed, 15,000 x g, for 3 minutes at room temperature, and
then the collection tube was discarded. To elute the DNA, the spin column was placed
in a 1.5 ml microcentrifuge tube. Then 100 µl of PureLink® Genomic Elution Buffer was
added to the column and incubated at room temperature for 1 minute before
centrifugation at maximum speed. After centrifugation, the 1.5 ml microcentrifuge tube
contained purified genomic DNA. At the end, the DNA concentrations extracted from
blood were checked with a nanodrop.

Conventional polymerase chain reaction (PCR; PCR Master Mix (2X) from Thermo
Scientific) was used to amplify the target DNA sequence. A combination of 25 µl of PCR
Master Mix, 1 µM of forward primer, 1 µM of reverse primer, and 1 µg of template DNA
was then added to PCR tube kept on ice. Next, 22 µL of water was added to the mix
making it a total of 50 µL. The mix ratio was repeated for each bonobo DNA sample and

placed into the thermal cycler upon completion. The thermocycler conditions were set
as follows:
1. 1.) Initial denaturation - temperature: 94 degrees Celsius, time: 1 minute
2. 2.) Denaturation – temperature: 94 degrees Celsius, time: 20 seconds
3. 3.) Annealing – temperature: 55 degrees Celsius, time: 20 seconds
4. 4.) Extension – temperature: 72 degrees Celsius, time: 20 seconds
5. 5.) Final extension – 72 degrees Celsius, time: hold
6.

The initial denaturation and final extension phases ran for one cycle each while the
annealing and extension phases ran for a total of 35 cycles. Once the thermocycler
completed the protocol, the PCR products were loaded onto 1.3% mini agarose gels for
visualization. The visualized images served as a check to be sure that the PCR reaction
was successful and produced the target amplicon. The gel was made of 1.3 g of
agarose, 100 ml of TAE buffer, and 2 µl of ethidium bromide. The TAE buffer was a
combination of 20 ml TAE buffer, 980 ml of dH2O, and 2 µl of ethidium bromide. 250 ml
of TAE buffer and 2.5 µl of ethidium bromide was added to the electrophoresis cell. For
each well, the PCR product and purple loading dye was loaded with a ratio of 10:2 as
suggested by the Quick-Load Purple 100 bp DNA ladder guide. The cells ran at 90 volts
and 400 amps for 55 minutes. Afterwards the gels were placed in a Bio-Rad ChemiDoc
XRS+ System.
730
The third objective, gel purification, was conducted using a ZymocleanTM Gel DNA
Recovery kit. The gels were placed on a UV transilluminator, and then each band was
excised using a scalpel and transferred to a 1.5 ml microcentrifuge tubes. The
microcentrifuge tubes' mass was 1.00 gram each. The total mass of the gel was
calculated by taking the mass of the gel piece inside the microcentrifuge tube and
subtracting the microcentrifuge tubes' mass. The mass number was then multiplied by 3
to figure out how many volumes (µl) of agarose dissolving buffer (ADB) to add to the
microcentrifuge tube. Once ADB was added to the microcentrifuge tubes, they were
incubated in a 55 degrees Celsius hot bead bath until the gel piece was completely
dissolved and the solution was homogenous. Afterwards, the melted agarose solution
was transferred to a Zymo-SpinTM Column in a collection tube and centrifuged for 60
seconds at 15,000 x g. The flow through was discarded from the collection tube so that
it could be used again. For DNA washing, 24 ml of 96-100% ethanol was added to the 6
ml DNA wash buffer and 96 ml of 96-100% ethanol was added to the 24 ml DNA wash
buffer. Next, 200 µl of DNA wash buffer was added to the column, and then the column
was spun again for 30 seconds 15,000 x g. The flow through was discarded and the
washing step was repeated. The final steps included placing the spin column into a 1.5
ml microcentrifuge tube, adding 10 µl of DNA elution Buffer directly to the center of the
spin column, and centrifuging for 60 seconds. The final mass of the purified DNA
products was determined using both gel visualization (using 1 µl of DNA mixed with 2 µl
dye and 3 µl water for clarity), and a nanodrop machine. The remaining extracted DNA
from the gels were sent to GENEWHIZ for Sanger sequencing.
753
754
755

Supplementary Material 2: Kinship Coefficients

Bonobo ID	F14	F16	M14	F13	M10	F05	M13	F18	M15	M12	M18	F10	M19
F14	1	0	0	0.125	0	0	0.125	0.031	0	0.063	0.094	0	0.25
F16	0	1	0	0	0.031	0	0	0.016	0	0	0	0	0
M14	0	0	1	0	0	0	0	0	0.125	0	0	0	0.25
F13	0.125	0	0	1	0	0	0.25	0.031	0	0.188	0.344	0	0.063
M10	0	0.031	0	0	1	0	0	0.063	0	0.063	0.031	0	0
F05	0	0	0	0	0	1	0	0	0.25	0	0	0	0
M13	0.125	0	0	0.25	0	0	1	0.031	0	0.188	0.219	0	0.063
F18	0.031	0.016	0	0.031	0.063	0	0.031	1	0	0.031	0.031	0	0.016
M15	0	0	0.125	0	0	0.25	0	0	1	0	0	0	0.063
M12	0.063	0	0	0.188	0.063	0	0.188	0.031	0	1	0.344	0	0.031
M18	0.094	0	0	0.344	0.031	0	0.219	0.031	0	0.344	1	0	0.047
F10	0	0	0	0	0	0	0	0	0	0	0	1	0
M19	0.25	0	0.25	0.063	0	0	0.063	0.016	0.063	0.031	0.047	0	1

Supplementary Material 2: Cell value is the kinship coefficient between the corresponding individuals. Each individual has a kinship coefficient of 1 with themselves.