Acclimation Capacity to Global Warming of Amphibians and Freshwater Fishes: Drivers, Patterns, and Data Limitations

Running title: *Acclimation: Amphibians and Freshwater Fishes*

Katharina Ruthsatz[1,2], Flemming Dahlke[3], Katharina Alter[4], Sylke Wohlrab[5,6], Paula C. Eterovick[1], Mariana L. Lyra[7,8], Sven Gippner[1], Steven J. Cooke[9], Myron A. Peck[4,10]

[1]*Zoological Institute, Technische Universität Braunschweig, Mendelssohnstraße 4, 38106 Braunschweig, Germany*

[2] *Institute of Animal Cell and Systems Biology, Universität Hamburg, Martin-Luther-King-Platz 3, 20146 Hamburg, Germany*

[3]*Ecology of Living Marine Resources, Universität Hamburg, Große Elbstraße 133, 22767 Hamburg, Germany*

[4]*Department of Coastal Systems, Royal Netherlands Institute for Sea Research, PO Box 59 1790 AB, Den Burg (Texel), the Netherlands*

[5]*Alfred Wegner Institute Helmholtz Center for Polar and Marine Research, 27570 Bremerhaven, Germany*

[6]*Helmholtz Institute for Functional Marine Biodiversity at the University of Oldenburg (HIFMB), 23129 Oldenburg, Germany*

[7]*New York University Abu Dhabi, Saadiyat Island, Abu Dhabi, P.O. Box 129188, United Arab Emirates*

[8]*Center for Research on Biodiversity Dynamics and Climate Change, State University of São Paulo-UNESP, Rio Claro, SP, Brazil.*

[9]*Fish Ecology and Conservation Physiology Laboratory, Department of Biology and Institute of Environmental and Interdisciplinary Science, Carleton University, Ottawa, ON K1S 5B6, Canada*

[10]*Marine Animal Ecology Group, Department of Animal Sciences, Wageningen University, PO*

Box 338, 6700 AH, Wageningen, the Netherlands

Key words: *acclimation response ratio, CT_{max}, thermal tolerance plasticity, developmental*

phenotypic plasticity, thermal bottleneck, metamorphosis, Bogert Effect, climate variability

hypothesis

Corresponding author: **Katharina Ruthsatz**; ORCID: 0000-0002-3273-2826. Current address:

Zoological Institute, Technische Universität Braunschweig, Mendelssohnstraße 4, 38106

Braunschweig, Germany. Phone: 0049 531 3912393. Email: katharinaruthsatz@gmail.com.

Word count: 7511

Abstract

Amphibians and fishes play a central role in shaping the structure and function of freshwater

environments. These organisms have a limited capacity to disperse across different habitats and

the thermal buffer offered by freshwater systems is small. Understanding determinants and

patterns of their physiological sensitivity across life history is, therefore, imperative to

predicting the impacts of climate change in freshwater systems. Based on a systematic literature

review including 345 experiments with 998 estimates on 96 amphibian (Anura/Caudata) and

93 freshwater fish species (Teleostei), we conducted a quantitative synthesis to explore

phylogenetic, ontogenetic, and biogeographic (thermal adaptation) patterns in upper thermal

tolerance (CT_{max}) and thermal acclimation capacity (Acclimation Response Ratio, ARR) as well

as the influence of the methodology used to assess these thermal traits using a conditional

inference tree analysis. We found globally consistent patterns in CT_{max} and ARR, with

phylogeny (taxa/order), experimental methodology, climatic origin, and life stage as significant

determinants of thermal traits. The analysis demonstrated that CT_{max} does not primarily depend

on the climatic origin but on experimental acclimation temperature and duration, and life stage.

Higher acclimation temperatures and longer acclimation times led to higher CT_{max} values,

whereby Anuran larvae revealed a higher CT_{max} than older life stages. The ARR of freshwater

fishes was more than twice that of amphibians. Differences in ARR between life stages were

not significant. In addition to phylogenetic differences, we found that ARR also depended on

acclimation duration, ramping rate, and adaptation to local temperature variability. However,

the amount of data on early life stages is too small, methodologically inconsistent, and

phylogenetically unbalanced to identify potential life cycle bottlenecks in thermal traits. We

therefore propose methods to improve the robustness and comparability of CT_{max}/ARR data

across species and life stages, which is crucial for the conservation of freshwater biodiversity

under climate change.

1. Introduction

Global climate change is not only causing an increase in mean air and water temperatures, but also an increased magnitude and frequency of extreme climatic events (Lee et al. 2023). As a result, ectotherms are more likely to experience temperatures beyond their critical thermal maximum (CT_{max}) in both terrestrial and aquatic habitats (Duarte et al. 2012; Sunday et al. 2014). This is particularly true for populations already living close to their upper thermal limit. Consequently, the ability to mitigate thermal stress through either migration, evolutionary genetic adaptation or acclimation is crucial for the persistence of species in a changing climate (Franks & Hoffmann 2002; Huey et al. 2012; Seebacher et al. 2015). Given the limited dispersal ability of many species (e.g., freshwater species; Woodward et al. 2010) and rapid pace of global warming (Hoffmann & Sgró 2011), physiological acclimation is arguably the most important mechanism for coping with climate change (Gunderson & Leal 2015). Understanding differences in acclimation capacity of species and identifying global patterns can therefore help to identify climate change risks to biodiversity and develop effective conservation measures (Somero 2010).

As an adaptive response to larger seasonal differences in temperature, thermal tolerance and acclimation capacity of ectothermic species or populations tend to increase with increasing latitude from tropical through temperate climate zones (e.g., Somero 2005; Deutsch et al. 2008; Sunday et al. 2011; Peck et al. 2014; Rohr et al. 2018; Cicchino et al. 2023; but see: Sørensen et al. 2016; Gunderson & Stillman 2015) and from higher to lower elevations (Enriquez-Urzelai et al. 2020; but not: Sunday et al. 2019; Gutiérrez-Pesquera et al. 2022). This biogeographical pattern is consistent with the *climate variability hypothesis* (Janzen 1967; Ghalambor et al. 2006), suggesting that climatic differences across altitudinal and latitudinal gradients lead to corresponding adaptations in thermal physiology (but see: Gutiérrez-Pesquera et al. 2022). Low-latitude species adapted to relatively stable temperature conditions may have a lower acclimation capacity and, therefore, may be more vulnerable to climate change (Tewksbury et al. 2008; Sunday et al. 2014; but see: Bovo et al. 2023). However, there is still little empirical evidence supporting the climate variability hypothesis, possibly due to the limited geographical and phylogenetic coverage of observations, and because of the inconsistency of the methods applied to measure the acclimation capacity of different species and life stages (Gutiérrez-Pesquera et al. 2016; Shah et al. 2017). Moreover, the use of different methods or protocols might impact the estimation of thermal traits (Terblanche et al. 2007; Chown et al. 2009; Rohr et al. 2018; Pottier et al. 2022a; but not: Sunday et al. 2019). For example, acclimation duration

(i.e., how long organisms were held at an acclimation temperature before being exposed to the
test temperature; Rohr et al. 2018; Ruthsatz et al. 2022a) and ramping protocol (i.e., heating or
cooling rate in thermal tolerance trials; Illing et al. 2020; Penman et al. 2023) have been
suggested to influence measurements of acclimation capacity, as the underlying physiological
processes occur over certain time periods.

In animals with complex life histories, thermal tolerance and acclimation capacity are thought
to change during ontogeny according to physiological and morphological reorganizations and
concomitant aerobic capacities in relation to oxygen demand (Pörtner 2002; Pörtner & Peck
2010; Ruthsatz et al. 2020a,b, 2022a) as well as energetic costs associated with developmental
processes (Ruthsatz et al. 2019). Furthermore, life stages might differ in their ability for
behavioral thermoregulation (Navas et al. 2008; Little & Seebacher 2017) resulting in stage-
specific adaptations in thermal traits (Huey et al. 1999). Therefore, determining taxon-specific
acclimation capacity at different ontogenetic stages should be taken into account when studying
climate adaptation of ectothermic species, as it will help identify life cycle bottlenecks and
provide robust data on the vulnerability of populations or species to global warming
(Bodensteiner et al. 2021; Pottier et al. 2022a; Dahlke et al. 2022). To date, most studies have
not distinguished among life stages or have only focused on adults when addressing the *climate*
variability hypothesis (e.g., Gunderson & Stillman 2015; Sunday et al. 2011, 2014, 2019; Rohr
et al. 2018) and the vulnerability of species to global change (e.g., Calosi et al. 2008; Comte &
Olden 2017a; Morley et al. 2019; Molina et al. 2023). Moreover, most studies have pooled
several pre- and post-metamorphic life stages (Pottier et al. 2022a; Weaving et al. 2022) thereby
risking to overlook a critical thermal bottleneck in the life cycle (Dahlke et al. 2020). The extent
to which thermal tolerance and acclimation capacity change during ontogeny is, therefore, not
clear for many taxa.

Amphibians and freshwater fishes tend to live in relatively shallow waters (e.g., wetlands,
ponds, rivers, lakes) and may, therefore, experience strong seasonal and daily temperature
fluctuations (Capon et al. 2021) and climate extremes such as heat waves (IPCC 2021). In
addition, both taxa have a limited ability to disperse over larger distances and habitats to avoid
unfavorable climatic conditions (Albert et al. 2011; Yu et al. 2013; Campos et al. 2021).
Consequently, as a result of local adaptation (Meek et al. 2023), there should be a close
correspondence between the capacity for thermal acclimation and the climatic conditions that
amphibians and freshwater fishes experience during their life cycle. However, freshwater
systems offer a wide range of thermal microhabitats that enable behavioral thermoregulation

(Campos et al. 2021), especially for (post-metamorphic) amphibians that can switch between water and land. The potential of behavioral thermoregulation could reduce the need for physiological adaptations (also known as the *Bogert Effect*, (Bogert 1949)) and thus counteract the emergence of geographical patterns in thermal acclimation capacity and/or thermal tolerance. Given the central role that amphibians and freshwater fishes play in shaping the structure and function of these ecosystems (Closs et al. 2016; Hocking & Babbitt 2014), understanding determinants and patterns of their physiological sensitivity is imperative to predicting the impacts of climate change on freshwater systems.

Here, we aimed to define the determinants and patterns of acclimation capacity in upper thermal tolerance in amphibians and freshwater fishes addressing, for the first time, ontogeny dependent variation. To do so, we compiled literature on upper thermal tolerance and collected empirical data for CT_{max} in four amphibian (i.e., larvae, metamorphs, juveniles, and adults) and three freshwater fish (i.e., larvae, juveniles, and adults) life stages acclimated to at least two different temperatures. Next, we calculated the population-specific acclimation capacity, i.e., mean acclimation response ratio (ARR) of upper thermal limits (i.e., the slope of the linear function describing the change in thermal tolerance with a given change in acclimation temperature; e.g., Hutchison 1961; Claussen 1977), and conducted a quantitative synthesis on the acclimation capacity of amphibians and freshwater fishes to test for differences among taxonomic groups, among life stages, and across thermal characteristics of populations, i.e., biogeographic differences based on local thermal adaptation. Further, we investigated how the methodological context (i.e., acclimation duration and temperature, ramping rate) affects estimates of thermal traits. Finally, we summarize methodological concerns, highlight key knowledge gaps, and provide research recommendations for generating reliable and comparable data on the acclimation capacity of ectothermic species and their life stages. This will improve our ability to predict future climate vulnerability of species and populations.

2. Materials and Methods

2.1 Systematic literature review

We conducted a systematic literature review using ISI Web of Science (ISI WOS, 2021) on 2022/06/30 and did not apply a timespan limit. The following Boolean search string was used to capture articles with experimental studies manipulating acclimation temperatures of amphibians and freshwater fishes at different life stages, and subsequently measured their CT_{max}: ("amphibian*" OR "newt*" OR "frog*" OR "toad*" OR "salamander*" OR "freshwater" AND "fish*") AND ("early" OR "young" OR "life stage*" OR "ontogen*" OR

"development*" OR "hatchling*" OR "alevin*" OR "larv*" OR "tadpole*" OR "metamorph*" OR "postmetamorph*" OR "post-metamorph*" OR "postlarva*" OR "post-larva*" OR "fry*" OR "parr*" OR "smolt*" OR "subadult*" OR "sub-adult" OR "juvenile" OR "fingerling*" OR "adult*") AND ("thermal" OR "temperature" OR "acclimat*" OR "heat" OR "warm*") AND ("tolerance*" OR "thermal tolerance*" OR "temperature tolerance*" OR "warming tolerance*" OR "heat tolerance*" OR "thermal stress tolerance*" OR "heat stress tolerance*" OR "temperature stress tolerance*" OR "limit*" OR "temperature stress*" OR "thermal limit*" OR "critical temperature*" OR "CT max" OR "critical thermal m*" OR "thermal performance breadth*" OR "thermal breadth*" OR "performance breadth*" OR "thermal range*" OR "thermal window*" OR "thermal tolerance window*" OR "tolerance window*" OR "sensitivity*" OR "thermal sensitivity").

Our search resulted in 11,057 documents (Fig. S1). After removing book chapters, conference contributions, reviews, meeting abstracts, editorial material, preprints, and proceedings articles, 10,740 published peer-reviewed articles remained in our initial database. Additionally, we manually added articles included in the meta-analyses of Claussen (1977), Gunderson and Stillman (2015), Comte & Olden (2017b), Morley et al. (2019) and Dahlke et al. (2020) that met our inclusion criteria (specified below) but were not obtained through the ISI Web of Science search. After an initial subjective evaluation of titles, 3,991 articles potentially containing results matching the objective of the present study were kept and further assessed for eligibility using the abstract. Thirty-four articles were not accessible. We contacted the authors of the original articles to request missing information and heard back from two authors. Finally, a total of 93 articles (34 on amphibians; 59 on freshwater fishes) met our inclusion criteria. Search methods are summarized in a PRISMA flowchart (Fig. S1), and a list of included articles and experimental studies therewithin is available in the figshare data repository under https://figshare.com/s/57775032a0b79c416ef2 (DOI: 10.6084/m9.figshare.24872133; published after acceptance).

2.2 Inclusion criteria

Experimental studies were selected based on the following seven inclusion criteria:

(1) Experimental studies were conducted on amphibians (anurans, caudates, or gymnophiones) or freshwater fishes (teleosts).

(2) Animals were acclimated to at least two constant acclimation temperatures under laboratory conditions prior to the CT_{max} measurements. Fluctuating temperature treatments were not

considered. Therefore, field experiments were not considered since the acclimation temperature under field conditions is likely to fluctuate.

(3) Articles provided comprehensive information on methodology (acclimation temperatures and duration, ramping rate), phylogeny (species names), sampling location (GPS location), and life stage. If no GPS coordinates were provided but a concrete sampling location was stated (e.g., Central Park, New York City, NY, USA), we searched for the coordinates of the respective location on Google Maps.

(4) Animals were collected from their natural habitat. Data were excluded if measurements were taken from specimens bred artificially to reduce confounding issues associated with artificial selective history (Bennett et al. 2018). Experimental studies were also included if adult animals were collected to immediately reproduce in the lab to obtain larvae.

(5) The critical thermal maximum (CT_{max}) was used as a standard measure of heat tolerance (Lutterschmidt & Hutchison 1997a). Experimental studies using other measures such as voluntary thermal maximum, time to death, heat knockdown, or lethal temperatures as well as extrapolations from thermal performance curves were not considered. In aquatic and terrestrial ectotherms such as amphibians and fish, CT_{max} is generally measured as loss of equilibrium (LOE) or loss of rightening response (LRR) following a steady increase in water or air temperature [dynamic method according to Fry (1947)]. In comparison to other endpoints such as the onset of spasms (Lutterschmidt & Hutchison 1997b), measuring CT_{max} as LOE or LRR is a non-lethal, robust method at various body sizes that is repeatable within individuals (Morgan et al. 2018).

(6) At CT_{max} measurements, the animals could be classified into one of four different categories representing the consecutive life stages of amphibians and freshwater fish: (a) larva (pre-metamorphic; amphibians: < Gosner stage 42), (b) metamorph (only for amphibians: Gosner stage 42-46), (c) juvenile (post-metamorphic), and (d) adult (after reaching sexual maturity). Embryos were not included in the present study because the assessment of acute heat tolerance in non-mobile life stages requires different endpoints that may not be directly comparable to the LOE/LRR-based CT_{max} of other life stages (Cowan et al. 2023; Lechner et al. 2023).

(7) Food was provided *ad libitum* during the acclimation time since food deprivation might decrease thermal tolerance and/or acclimation capacity (Lee et al. 2016).

2.3 Data extraction

When all inclusion criteria were met, data were collated in a spreadsheet. We extracted mean CT_{max} for all acclimation temperatures resulting in 998 single data points (513 for amphibians; 485 for freshwater fish). Some of the selected articles performed different experimental studies on e.g., different populations of one species, different species, or different life stages. Therefore, all available datasets were included, resulting in 345 experimental studies from 93 articles with 345 paired effect sizes CT_{max} and acclimation capacity. Data presented in the text or tables were directly extracted from the articles. When only raw data were available, mean values were calculated. For articles that presented results in figures instead of tables, Engauge Digitizer 12.1 was used (Mitchell et al. 2021) to extract data from the graphs. In addition to CT_{max} data, information on the methodology (i.e., acclimation temperatures and duration, ramping protocol), as well as on variables representing sampling location as detailed as possible (i.e., GPS coordinates), phylogeny (i.e., scientific classification according to the Linnean classification), and life stage at CT_{max} assessment was extracted. The data extractions were performed by KR, KA and PCE, followed by an accuracy check of the data (KA: freshwater fish sub dataset; KR: amphibian sub datatset; PCE: both sub datasets).

2.4 Bioclimatic variables

For each sampling location, 19 bioclimatic metrics, related to temperature and precipitation, and elevation were extracted using the WorldClim 2 database (http://www.worldclim.org/; Fick & Hijmans 2017) for the average of the years 1970–2000. The data were extracted at a spatial resolution of 30 seconds (~1 km^2), using package 'geodata' (Hijmans 2021) in R (version 4.2.1; R Core Team, 2020). Bioclimatic variables are coded as follows (name of the variable and the unit used as input is included in parenthesis): Annual Mean Temperature (Bio1; °C), Mean Diurnal Range (Bio 2; °C), Isothermality (Bio 3; percentage), actual Temperature Seasonality (Bio 4; standard deviation in °C), Maximum Temperature of Warmest Month (Bio 5; °C), Minimum Temperature of Coldest Month (Bio 6; °C), Annual Temperature Range (Bio 7; °C), Mean Temperature of Wettest Quarter (Bio 8; °C), Mean Temperature of Driest Quarter (Bio 9; °C), Mean Temperature of Warmest Quarter (Bio 10; °C), Mean Temperature of Coldest Quarter (Bio 11; °C), Annual Precipitation (Bio 12; mm), Precipitation of Wettest Month (Bio 13; mm), Precipitation of Driest Month (Bio 14; mm), Precipitation Seasonality (Bio 15; coefficient of variation expressed in percent), Precipitation of Wettest Quarter (Bio 16; mm), Precipitation of Driest Quarter (Bio 17; mm), Precipitation of Warmest Quarter (Bio 18; mm), and Precipitation of Coldest Quarter (Bio 19; mm) (http://www.worldclim.org/data/bioclim.html). These macroclimatic data were used as

approximations to identify patterns of local adaptation in amphibians and freshwater fish, as microclimatic data (e.g. site-specific temperatures) were not available in original articles or in the WorldClim database. Following previous studies (Morley et al. 2019; Carilo Filho et al. 2022), mean near-surface air temperature was assumed to reflect the temperature profile of freshwater systems and used to analyze thermal adaptation in both taxa. We are aware that these macroclimatic data as provided by the WorldClim database (representing surface air temperature) present some limitations in reflecting microclimatic data (see Conclusion section for methods to improve future studies). In contrast to marine habitats, the temperature of small or shallow bodies of water might fluctuate with the surface air temperature and animals at the surface of freshwater systems might be further exposed to high temperatures. Therefore, we assume that the average near-surface air temperature is a suitable estimate of the temperature of freshwater systems, thereby reflecting the thermal local adaptation of investigated amphibian and freshwater populations. This is in accordance with previous studies testing the effect of thermal adaptation on the thermal physiology of various taxa (e.g., Gutiérrez-Pesquera et al. 2016; Morley et al. 2019; Carilo Filho et al. 2022; Sinai et al. 2022). Sampling locations were assigned to latitudinal groups based on the absolute latitude (°N/S) and were categorized as either tropical (0–25°), sub-tropical (>25–40°), temperate (>40–53.55°) or polar (>53.55°; Morley et al. 2019).

2.5 Effect size calculation: acclimation response ratio (ARR)

A well-established method to measure acclimation capacity in thermal tolerance in ectothermic animals is the calculation of the acclimation response ratio (ARR), i.e., the slope of the linear function describing the change in thermal tolerance with a given change in acclimation temperature (e.g., Hutchison 1961; Claussen 1977; Gunderson & Stillman 2015; Morley et al. 2019). We separately calculated the ARR for CT_{max} within each study using the equation according to Claussen (1977):

$$ARR = \frac{CTmax_{[T2]} - CTmax_{[T1]}}{T2 - T1},$$

where T represents the acclimation temperature (°C; with T2 = highest acclimation temperature and T1 = lowest acclimation temperature) and CT_{max} the heat tolerance estimates (°C). When data on more than two acclimation temperatures were presented, we calculated the ARR for each stepwise comparison (e.g., 18-20°C, 20-22°C, 22-24°C; Pottier et al. 2022a) and used the mean ARR of all comparisons in the statistical analysis. Higher absolute values of ARR correspond to higher plasticity in thermal tolerance limits (i.e. greater acclimation capacity;

Claussen 1977; Kingsolver & Huey 1998; Gunderson & Stillman 2015; van Heerwarden et al. 2016). An acclimation response ratio of 1.00 indicates a 100% acclimation in thermal tolerance to a temperature increase of 1°C (Morley et al. 2019).

2.6 Statistical analyses

For all statistical tests R 4.0.2 (R Core Team, 2020) was used. All plots were constructed using R packages 'ggplot2' (Wickham & Wickham 2009), 'ggtree' (Yu et al. 2017) and Adobe Illustrator CS6.

Conditional inference tree (CIT) analysis (R package 'partykit', Hothorn & Zeileis 2015) was used to assess the influence of geographic origin (bioclimatic variables, elevation), phylogeny (taxon and order level), experimental methodology (ramping rate, mean acclimation duration, mean acclimation temperature), and ontogeny on CT_{max} and ARR of amphibians and freshwater fishes. As an advantage over traditional (parametric) methods, CIT is a non-parametric method that handles complex non-linear relationships without making specific assumptions about data distribution or being sensitive to outliers. CIT involves recursive partitioning to split data into subsets based on the relevance of predictor variables. At each node of the tree, a permutation-based test (Monte Carlo method with Bonferroni correction) determines whether the split is statistically significant ($\alpha < 0.05$). The initial split in the tree indicates which predictor variable has the strongest correlation with the response variable (CT_{max} or ARR). The resulting tree provides a hierarchical structure and classification of significant predictor variables. A post-pruning strategy based on the Akaike Information Criterion (AIC, Akaike 1974) was used to avoid overfitting, i.e., removal of nodes that do not improve the overall fit of the model (Hothorn & Zeileis 2015). The raw data used for the analysis are provided in the electronic supplementary material (Table S1).

Phylogenetic trees for visualizations were created using the R package 'fishtree' (Chang et al. 2019) for freshwater fishes (Teleostei). The amphibian ultrametric tree was obtained from the timetree.org website in June 2022 (Kumar et al. 2022). To ensure the validity, we developed a workflow to filter out taxonomically invalid taxa. Firstly, a time tree was generated using the "Build a Timetree" function on timetree.org. The taxa names were then extracted from the generated time trees using the R package 'ape' (Paradis & Schliep 2019), excluding non-binomial names. The extracted list was cross-checked with the GBIF Backbone Taxonomy (GBIF Secretariat, 2021) using the species matching tool (https://www.gbif.org/tools/species-lookup, accessed June 2022). Matches at the species rank with an accuracy of 100% were extracted and matches with lower accuracy were manually verified. The resulting species list

was uploaded back onto timetree.org using the "Load a List of Species" function. The newly

generated time tree was then downloaded and used for visualizations.

The relationship between absolute latitude (°N/S), elevation (m), and the bioclimatic variables

that revealed a significant effect on CT_{max} (Bio14, Bio7) and ARR (Bio3) in the CIT analysis

was assessed using linear regressions. Differences in respective bioclimatic variables between

latitudinal groups (tropical, sub-tropical, temperate, polar) were compared using Kruskal-

Wallis tests followed by pairwise Mann-Whitney U- tests with false discovery rate (FDR)-

correction.

We were unable to test for a publication bias (i.e., whether statistically insignificant or negative

results are less likely to be published) in our data set given the use of a non-standardized effect

size metric (i.e., ARR). Therefore, it remains possible that this bias may also occur.

3. Results

The analysis of paired CT_{max} and ARR values included 201 experimental studies on 96

amphibian species (2 orders) (Fig. 1a, b) and 144 experimental studies on 93 teleost species (14

orders) (Fig. 1c, d), corresponding to a phylogenetic coverage of approximately 1% within each

taxon. No data were available for Gymnophiona. Most experimental studies determined CT_{max}

and ARR in adult animals. For freshwater fishes, only one study determining CT_{max} in larvae

was obtained, while 20% of the data covered juveniles (Fig. 1c). For amphibian species, 12%

of the data was available for one or several of the early-life stages, i.e. larvae, metamorphs,

and/or juveniles (Fig. 1a).The geographic origin of amphibian and fish species displayed a bias

towards temperate and subtropical regions, with 60% of all sampling sites located in North

America, 10% in Australia and New Zealand, 8% in South America, 6% in Europe, and 4% in

China (Fig. 1b, d). The CT_{max} protocols were variable in both taxa, with acclimation times

ranging from 0.5 to 150 d (mean = 11.2 d), acclimation temperatures ranging from 8.0 to 31.75

°C (mean = 19.45 °C), and ramping rates ranging from 0.02 to 1.0 °C min^{-1} (mean = 0.75 °C

min^{-1}).

Across all species studied, CT_{max} ranged from 20.75 °C to 46.00 °C (Table S1; Fig. S2a).

Acclimation capacity in upper thermal tolerance was positive in 97.1% of the experimental

studies with an average of 0.22 °C with every 1 °C increase in acclimation temperature across

all experimental studies and life stages, whereas the acclimation response for CT_{max} was

negative (i.e., negative ARR) in 2.0% of the experimental studies. Across all species studied,

ARR ranged from -0.08 to 1.68 (Table S1; Fig. S2b).

Figure 1. Phylogenetic, ontogenetic, and geographic background of critical thermal maximum (CT$_{max}$) and acclimation response ratio (ARR) measurements in (a, c) amphibians (Anura and Caudata) and (b, d) freshwater fishes (Teleostei). Phylogenetic trees contain (a) 96 amphibian and (b) 93 teleost species assigned to 2 and 14 orders, respectively. Grey shaded squares indicate which life stages of a species were studied. Maps show the geographical origin and local mean temperature of (c) amphibian and (d) fish species. Phylogenetic coverage was approximately 1% within each taxon. Silhouettes were taken from PhyloPic (www.phylopic.org).

Figure 2. Conditional inference tree (CIT) for critical thermal maximum (CT_{max}) of freshwater fish (Teleostei) and amphibians (Anura and Caudata). The sequence of internal nodes (splits) corresponds to a hierarchical structure of significant predictor variables. At each split, lines indicate the classification into groups with higher (black) and lower (red) CT_{max} values. Boxplots show the distribution of CT_{max} values (colored symbols) for each terminal node (n = number of experimental studies). Acclimation T = mean acclimation temperature used in an experiment before CT_{max} measurements (°C). Habitat T = mean annual temperature Bio 1 (°C). Acclimation duration = mean exposure time to acclimation temperatures before CT_{max} measurements (d).

For CT_{max} data of freshwater teleosts and amphibians (345 observations = number of experiments), the CIT model (73% explained variance) resulted in a classification tree with 8 internal splits and 9 terminal nodes (Fig. 2). Phylogeny (order/taxon), acclimation temperature and acclimation duration were the main discriminators of CT_{max}, followed by geographic origin (annual habitat temperature, bio1) and life stage. The first split separated a group of teleost studies (mainly salmonid species) with lower CT_{max} values compared to the mean of all other studies (p < 0.001). In this group of cold-water fish, CT_{max} increased significantly with acclimation duration (nodes 1-3), with classification thresholds of split 2 and 3 defining differences between short (< 6 days), intermediate (6-21 days) and long acclimation times (>

21 days). For the majority of teleost and amphibian data (308 observations), CT_{max} was significantly dependent on acclimation temperature (split 4, p < 0.001), mean annual habitat temperature (Bio 1; split 5, p = 0.001) and life stage (split 6, p < 0.001). Overall, there was a positive correlation between acclimation/mean annual habitat temperature and CT_{max} (node 4-9). Split 4 separated teleosts and amphibians acclimated below or above 28°C. In the colder group (split 5), CT_{max} was higher for teleosts and amphibians that originated from warmer habitats, with a classification threshold of 16.2°C (node 4 & 5). For species acclimated above 28°C, CT_{max} differed between ontogenetic life stages (split 6), with larvae having higher tolerance limits than juveniles, metamorphs and adults. However, this classification was phylogenetically unbalanced, as all larvae were anuran species (node 8 & 9). Splits 7 and 8 show that after accounting for ontogenetic differences, CT_{max} was higher at warmer acclimation temperature (>31 °C, node 6 & 7) and higher mean annual habitat temperatures (> 16.8 °C, node 8 & 9).

Figure 3. Conditional inference tree (CIT) for the acclimation response ratio (ARR) of freshwater fish (Teleostei) and amphibians (Anura and Caudata). The sequence of internal nodes (splits) corresponds to a hierarchical structure of significant predictor variables. At each internal node, lines indicate the classification into groups with higher (black) and lower (red) ARR values. Boxplots show the distribution of ARR values (colored symbols) for each terminal

node (n = number of experimental studies). In node 1, high outlier values (n = 2) were not shown for scaling reasons. Acclimation T = mean acclimation temperature used in an experiment before CT_{max} measurements (°C). Ramping rate = heating rate for CT_{max} measurements in °C per minute. Isothermality = local temperature variability Bio 3 (%). Acclimation duration = mean exposure time to acclimation temperatures before CT_{max} measurements (d).

For ARR data (345 observations), the CIT model (52% explained variance) produced a classification tree with 7 internal splits and 8 terminal nodes (Fig. 3). The main discriminators of ARR were phylogeny (order/taxon) and acclimation time, followed by CT_{max} methodology (ramping rate) and a minor influence of local temperature variability (isothermality, Bio 3). The first split of the classification tree separated a large number of teleost studies (102 observations, 68 species, node 1 & 2) with a higher ARR compared to the average of the other fish and amphibian studies in the dataset (p < 0.001). The relatively high thermal plasticity of the fish species in node 1 & 2 was not related to any of the variables examined. Within this group, however, ARR decreased at higher acclimation temperatures, with a classification threshold of 21°C (split 2, p = 0.003). Split 3 was based on CT_{max} methodology, classifying fish and amphibian studies into groups with an acclimation duration longer or shorter than 14 days (node 3-8). Longer acclimation generally resulted in higher ARR values (node 3-5, p < 0.001), but the acclimation effect differed according to ramping rate (split 4), phylogeny (split 5 & 6) and climate variability (split 7). When the acclimation duration exceeded 14 days, ARR studies were further classified according to ramping rate (split 4, p <0.001) and taxon (split 5, p = 0.031). These classifications imply higher ARR values of anurans at slow heating rates (< 0.1°C min^{-1}), and higher ARR values of teleosts compared to amphibians at ramping rates > 0.1°C min^{-1} (Fig. 3). If the acclimation period was less than 14 days, a subsequent split occurred according to phylogenetic order (split 6, p < 0.001) and isothermality (split 7), indicating a positive relationship between ARR and local temperature variability (node 7 & 8, p = 0.011). Ontogenetic differences in ARR were not detected.

Isothermality (Bio 3; R^2 = 0.708, p <0.001) and annual mean temperature (Bio 1; R^2 = 0.696, p < 0.001) decreased from the tropics to the polar latitudes (Fig. S4). Both bioclimatic variables did not correlate with elevation (Bio 1: R^2 = 0.137, p <0.001; Bio 3: R^2 = 0.012, p = 0.046).

4. Discussion

4.1 Context-Dependent Drivers and Broad-Scale Patterns of Physiological Limits

The most striking result from our analyses is that the acclimation capacity of freshwater fish was more than twice that of amphibians, indicating a strong phylogenetic signal. Our findings align with those reported in previous syntheses (Gunderson & Stillman 2015; Rohr et al. 2018; Morley et al. 2019; Pottier et al. 2022a), demonstrating a higher thermal plasticity in organisms inhabiting aquatic habitats compared to their terrestrial counterparts in general, and with fish (marine and freshwater) exhibiting greater thermal plasticity compared to amphibians (Gunderson & Stillman 2015). Compared to most (post-metamorphic) amphibians, freshwater fish are restricted to their aquatic habitat throughout their life cycle (Comte & Grenouillet 2013). As aquatic habitats tend to have less spatial variability in operative thermal conditions than terrestrial habitats (Gunderson & Stillman 2015; Sunday et al. 2014), behavioral thermoregulation is constrained, and freshwater fish are more likely than amphibians to use thermal plasticity to buffer against changing thermal conditions. Consequently, the capacity for thermal plasticity appears to be phylogenetically conserved between both taxa (Angilletta et al. 2002; Bodensteiner et al. 2021), depending on the ability for behavioral thermoregulation as explained by the *Bogert effect* (Bogert 1949; Cowles & Bogert 1944; Muñoz 2022), rather than on the level of thermal variation to which a population is exposed to (Huey et al. 1999) as suggested by the *climate variability hypothesis* (Janzen 1967). Moreover, unlike freshwater fishes, most amphibians undergo a habitat shift from a larval aquatic to a post-metamorphic terrestrial habitat (Shi 2000). As an adaptive response to often shallow or temporary larval habitats (Newman 1992), amphibian larvae display a high degree of plasticity in growth and development (Kulkarni et al. 2017; Ruthsatz et al. 2018a,b; Burraco et al. 2021; Sinai et al. 2022), providing a means for increasing fitness (Schlichting & Pigliucci 1998). Therefore, plasticity in timing of metamorphosis appears to be more important than that in thermal tolerance to reduce mortality risk (Rudolf & Rödel 2007) due to desiccation or temperature extremes (Burraco et al. 2022; Albecker et al. 2023). This might, at least in part, explain the lack of an ontogenetic pattern in acclimation capacity in amphibians.

Unlike acclimation capacity, geographic origin was one of the primary determinants of heat tolerance in both taxa, with a higher heat tolerance in populations from regions with a higher annual mean temperature compared to populations from regions with a lower annual mean temperature. These findings support, at least in part, the *climate variability hypothesis* (Janzen 1967; Bozinovic et al. 2011) since annual mean temperature decreases with latitude and is

highest in tropical low-elevation regions. Both taxa, therefore, exhibit physiological adaptations to latitude-dependent thermal regimes to which they are exposed. Our results agree with the findings of a large body of research that has confirmed the link between physiological limits and large-scale geography based on a species' local adaptation to temperature and other associated climatic variables (e.g., Gutiérrez-Pesquera et al. 2016; Sunday et al. 2011, 2019; Pintanel et al. 2022; but not: Addo-Bediako et al. 2000; Sørensen et al. 2016). Yet, in addition to the broader literature, our synthesis indicated that CT_{max} was not only *generally* higher in populations from warmer regions as often found in low latitudes but also, importantly, varied with ontogeny. These findings emphasize the importance of assessing life stage-specific sensitivity to thermal stress as well as spatial climatic differences in conservation science. Therefore, focusing on large-scale geographical patterns for predicting how biodiversity will respond to future environmental change might bear the risk of overlooking context-dependent variation in thermal traits and thus, intraspecific differences in vulnerability to changing thermal conditions (Duarte et al 2012, Gutierrez-Pesquera et al 2016). For example, Bovo et al. (2023) demonstrated that responses of tropical amphibians to climate variation were heterogenous as a consequence of intraspecific variation in physiological traits and spatial variation in climate with elevation. Furthermore, Sunday et al. (2011) and Pinsky et al. (2019) reported that the physiological sensitivity of ectotherms across all latitudes depended on the realm, with terrestrial ectotherms being less sensitive to warming than aquatic ectotherms due to their higher capacity for behavioral thermoregulation.

4.2 Life Stage-Specific Thermal Sensitivity as a Key Factor in Species Vulnerability to Climate Change

In species with complex life-histories such as amphibians and teleost fish, life stages differ in size, morphology, physiology, and behavior (Wilbur 1980). Therefore, selection might promote stage-specific adaptations in thermal physiology (Enriquez-Urzelai et al. 2019; Ruthsatz et al. 2022a -or b). Ignoring those life stage-specific differences in thermal physiology may drastically underestimate climate vulnerability of species with consequences for successful conservation actions. Here, we found that CT_{max} but not acclimation capacity differed between life stages in amphibians, with a higher CT_{max} in larval stages. However, this pattern was only evident when high acclimation temperatures were used in experimental set-ups and was most pronounced in larvae from habitats with a high annual mean temperature as those found in the tropics. Limnic larvae may have a reduced capacity for behavioral thermoregulation due to their limited body size impairing the movement between different microclimates (Kingsolver et al.

2011; Sinclair et al. 2016; Enriquez-Urzelai et al. 2019), making them more dependent on passive responses to temperature fluctuations. To cope with changes in temperatures, a high heat tolerance is therefore advantageous in early life stages (Ruthsatz et al. 2022a). This might be particularly true for animals that are already living close to their heat tolerance since the ontogenetic difference was only evident at high acclimation temperatures and strongest in anurans from habitats with a high annual mean temperature. In contrast, post-metamorphic stages might rather be able to select favorable microclimates by behavioral thermoregulation (Navas et al. 2007; Haesemeyer 2020). This is particularly true for amphibians, as their post-metamorphic terrestrial habitats offer much spatial variability in operative thermal conditions (Gunderson & Stillman 2015), while juvenile and adult (freshwater) fish are able to behaviorally thermoregulate by performing vertical and horizontal movements (Amat-Trigo et al. 2023; Breau et al. 2007; but not: Clark et al. 2022). Our findings are in line with the pattern found for aquatic larvae by Cupp (1980), Sherman (1980), Enriquez-Urzelai et al. (2019), and Ruthsatz et al. (2022a), who demonstrated a higher CT_{max} in amphibian larvae than in post-metamorphic stages. In contrast, Dahlke et al. (2020) found no difference in heat tolerance between larval and adult stages in marine and freshwater fish. Notably, our synthesis yielded only one estimate for larval CT_{max} and acclimation capacity in freshwater fish and, thus, we lack the data for any conclusion. Given that small body sizes of larvae restrict their capacity for behavioral thermoregulation, we would would expect freshwater fish to exhibit the same life stage-specific differences in thermal sensitivity observed in amphibians. In a recent study on the European common frog (*Rana temporaria*), young larvae may define the climate sensitivity of populations since that life stage exhibited the lowest acclimation capacity (Ruthsatz et al. 2022a). Furthermore, it is worth noting that our synthesis did neither encompass embryos nor gametes producing/spawning stages, which have recently been reported to have the lowest heat tolerance in fish (Dahlke et al. 2020, 2022; Pottier et al. 2022b) and the lowest acclimation capacity across ectotherms (Pottier et al. 2022a). To better identify potential life history bottlenecks in thermal sensitivity in amphibian, fish and other taxa inhabiting freshwater, future studies should adopt a more comprehensive approach by considering a wider range of life stages within species.

4.3 Understanding Context-Dependent Physiological Adaptation in Ectotherms

Global syntheses on physiological studies can help us determine the winners and losers of climate change through assessment of broad-scale patterns of species' thermal limits and acclimation capacity for modifying their thermal tolerance (Somero 2010). This knowledge, in

turn, enables us to develop suitable conservation strategies to mitigate the negative effects of climate change. However, our key findings emphasize that assessing species' vulnerability to changing thermal conditions based on large-scale geographic and/or phylogenetic patterns in thermal traits might cover up context-dependent physiological adaptations. In other words, tropical ectothermic species are considered particularly vulnerable to global warming as they live close to their physiological limits and have poor acclimation ability (Tewksbury et al. 2008; Huey et al. 2009; Sunday et al. 2014), but such generalizations might for instance ignore intraspecific variation in physiological limits across variation in climate with elevation (Bovo et al. 2023). Physiological adaptations are driven by the interplay between microclimate temperature heterogeneity and the behavioral thermoregulatory abilities of ectotherms (Huey et al. 2012; Pincebourde et al. 2016; present study) depending on their habitat characteristics (Pinsky et al. 2019; Kulkarni et al. 2017), ontogeny (Enriquez-Urzelai et al. 2019; Ruthsatz et al. 2022a), life history traits such as body size (Rubalcaba et al. 2020; Peralta-Maraver & Rezende 2021) or activity patterns (Navas et al. 2007; Ruthsatz et al. 2022b), and/or energy balance (Pörtner et al. 2005; Muñoz et al. 2022). Physiological traits and limits are consequently rather evolutionary driven dynamic concepts than fixed values for a species (Bovo et al. 2018, 2023; Navas et al. 2022). In order to improve predictions of climate change impacts on biodiversity, it is imperative to deepen our understanding of context-dependent physiological adaptations (Meek et al. 2023), thereby advancing the development of suitable conservation measures/strategies that incorporate evolutionarily enlightened perspectives (Ashley et al. 2003; Cook & Sgró 2018) beyond the species level (Fig. 4).

4.4 Conclusion

There is a growing body of physiological studies assessing thermal limits and acclimation capacity of species, investigating physiological systems setting these limits to better predict shifts in the productivity and species distribution patterns in a warming world. Our synthesis points to representation biases in taxonomy, species' biogeographic distribution, life stage, and biases resulting from non-standardized study design. Here, we found the influence of life stage, phylogeny, and thermal adaptation to depend on acclimation duration and acclimation temperature of the animals and the ramping rate used, underscoring the importance of a thoughtful selection of the methodological approach. Notably, due to the biases present in existing data sets, most results from subsequent analyses (including those presented here) cannot be generalized to all geographic regions, life stages, and (taxonomic) groups, which limits the ability to predict organismal responses to global change (White et al. 2021).

Therefore, we conclude our synthesis by addressing those data inadequacies and proposing methods to enhance data collection presented in five themes (Fig. 4):

- **Biogeographic and taxonomic coverage**

We found geographical trends in physiological limits associated with bioclimatic conditions at different latitudes. Thus, additional research is needed in poorly represented (mostly tropical) regions that generally correspond to low- and middle-income countries (King 2004). Addressing this issue can yield additional, important insights, as such geographic bias in existing data sets has been shown to limit our ability to predict organismal responses in these less-studied regions (White et al. 2021). Our synthesis highlights that most experiments have been conducted in North America, Europe, and Australia and information gaps exist for most parts of Neotropics, Africa and Sino-Oriental regions. Those regions harbor the highest diversity of species for both anurans and freshwater fishes (Jenkins et al. 2013, Van der Sleen & Albert, 2022). Such regional differences in research effort are common in conservation science (Schiesari et al. 2007; Winter et al. 2016; White et al. 2021; McLaughlin et al. 2022; Sinai et al. 2022) despite the fact that under-studied regions contain the vast majority of global biodiversity hotspots (Mittermeier et al. 2011). Furthermore, most of these experiments used species that are common, widely distributed, and/or easily obtained by researchers. Studies on other species (particularly those already in decline) are needed to avoid taxonomic bias and reach stronger conclusions on whether specific taxa might be more sensitive to global warming (da Silva et al. 2020). Additionally, there may be an elevation bias in sampling, as most available data represent lowland rather than mountain species. As, for example, CT_{max} of tropical and subtropical amphibian species has been shown to vary within the same latitude (Duarte et al 2012, Gutierrez-Pesquera et al 2016), we urge future studies to focus on elevation gradients as well as cross-latitudinal sampling, especially in tropical high-mountain populations (Bovo et al., 2023).

- **Relevance of life stage and sex**

We urge future studies to measure a wider range of life stages and to measure both sexes within species to better identify thermal bottlenecks and assess climate change risks (Klockmann et al. 2017; Dahlke et al. 2020). All of the summaries to date likely overestimate physiological limits since most studies have been performed on adults, and thermal tolerance increases (Klockmann et al. 2017; Rubalcaba & Olalla-Tárraga 2020; but not: di Santo & Lobel 2017) and acclimation capacity decreases (Pottier et al. 2021) with body size. Furthermore, after sexual maturity,

females and males differ in a wide range of morphological, physiological, and behavioral
aspects as well as in their energetic investment in gamete production (Hayward & Gillooly
2011). A recent meta-analysis across ectothermic taxa revealed that the acclimation capacity
differed between adult males and females in wild-caught animals (Pottier et al. 2021).
Furthermore, Dahlke et al. (2020) found narrower thermal tolerance ranges in spawning females
and van Heerwaarden and Sgrò (2021) demonstrated that a low heat tolerance of male fertility
is a critical bottleneck in insects.

- **Ecological relevance**

Future laboratory experiments should adopt a more comprehensive and multifaceted approach
for higher ecological relevance of thermal trait estimates (Desforges et al. 2023). Under natural
conditions, organisms must often cope with multiple simultaneously occurring environmental
stressors (Rohr & Palmer 2013; Gunderson et al. 2016) such as declining dissolved oxygen
levels in freshwater habitats due to climate-induced temperature increases (Pörtner & Peck
2010). As thermal limits are shaped by oxygen availability (Pörtner 2001, 2010), organisms
might exhibit lower thermal limits under natural conditions. Moreover, exposure to pollutants
might reduce thermal tolerance (Little & Seebacher 2015) or acclimation capacity (Ruthsatz et
al. 2018b) due to increased metabolic demands of detoxification processes or disruption of
endocrine pathways involved in physiological acclimation. Physiological responses to
increased water temperature as performed in the experimental studies summarized here (i.e., a
single stressor) may not align with observed responses of individuals in the natural
environments with multiple stressors (Katzenberger et al. 2014; Potts et al. 2021).

- **Methodological approach**

Customizing protocols to account for organismal and context-dependent variations in
physiological limits (e.g., body size, life stage, sex, thermal history) will allow researchers to
obtain more ecologically relevant estimates to inform conservation efforts. The application of
an acute thermal ramping rate and a standardized endpoint such as the loss of equilibrium are
used to measure critical thermal limits (Becker & Genoway 1979). The estimates are sensitive
to differences in the methods. For example, faster ramping (heating) rates tend to yield higher
thermal tolerance estimates compared to slower ramping rates (Moyano et al. 2017; Kovacevic
et al. 2019; Penham et al. 2023). Furthermore, measuring several endpoints (loss of
equilibrium/rightening response, onset of spasms/heat rigor) if possible (Wu & Kam 2005;
Cowan et al. 2023) might facilitate comparisons between different taxonomic groups and/or

experiments (Lutterschmidt & Hutchison 1997). Using wild-collected animals is important as
those reared in the laboratory may have physiology traits that differ from wild conspecifics
(Pottier et al. 2021; Morgan et al. 2022). Methodological recommendations have been recently
published and comparable methods are needed to compare thermal limits of different life stages
(Cowan et al. 2023; Desforges et al. 2023).

- **Through reporting of research details**

Thorough and transparent reporting of experimental details in empirical studies such as
sampling location and animal origin, among others, is required to enhance the comparability of
studies on thermal traits. Furthermore, most studies working on adults did not report the sex of
animals despite the potential for sex (or reproductive state) to be important factors in thermal
sensitivity. Additionally, reporting microhabitat temperatures (i.e., in-site temperatures) will
improve our understanding of thermal adaptation at the population-level. The ability to make
broad-scale comparisons of thermal tolerance across taxa, life stages, and regions will be
enhanced when studies report as much methodological detail as possible. Consequently, these
future studies will contribute more robust estimates of climate vulnerability needed to guide
climate change interventions.

By considering our recommendations, future studies will be more comparable, facilitating the
utilization of respective findings in large-scale studies and models that assess species
vulnerability and thus, population dynamics under global warming.

Figure 4. Synthesis of biological and methodological determinants as well as key directions for future research on acclimation capacity in amphibians and freshwater fish to advance the application of thermal traits in assessing species' and populations' vulnerability to climate change. See text for further details.

5. Data availability

Data associated with this study are available in the figshare data repository under https://figshare.com/s/57775032a0b79c416ef2 (DOI: 10.6084/m9.figshare.24872133; published after acceptance).

6. References

Akaike, H. (1974). A new look at the statistical model identification. IEEE transactions on automatic control, 19(6), 716-723.

Albecker, M. A., Strobel, S. M., & Womack, M. C. (2023). Developmental Plasticity in Anurans: Meta-analysis Reveals Effects of Larval Environments on Size at Metamorphosis And Timing of Metamorphosis. Integrative and Comparative Biology, icad059.

Albert, J. S., Carvalho, T. P., Petry, P., Holder, M. A., Maxime, E. L., Espino, J., ... & Reis, R. E. (2011). Aquatic biodiversity in the Amazon: habitat specialization and geographic isolation promote species richness. Animals, 1(2), 205-241.

Amat-Trigo, F., Andreou, D., Gillingham, P. K., & Britton, J. R. (2023). Behavioural thermoregulation in cold-water freshwater fish: Innate resilience to climate warming?. Fish and Fisheries, 24(1), 187-195.

Angilletta Jr, M. J., Niewiarowski, P. H., & Navas, C. A. (2002). The evolution of thermal physiology in ectotherms. Journal of thermal Biology, 27(4), 249-268.

Angilletta, M. J. (2009). Thermal adaptation: a theoretical and empirical synthesis.

Ashley, M. V., Willson, M. F., Pergams, O. R., O'Dowd, D. J., Gende, S. M., & Brown, J. S. (2003). Evolutionarily enlightened management. Biological Conservation, 111(2), 115-123.

Becker, C. D., & Genoway, R. G. (1979). Evaluation of the critical thermal maximum for determining thermal tolerance of freshwater fish. Environmental Biology of Fishes, 4, 245-256.

Bodensteiner, B. L., Agudelo-Cantero, G. A., Arietta, A. A., Gunderson, A. R., Muñoz, M. M., Refsnider, J. M., & Gangloff, E. J. (2021). Thermal adaptation revisited: How conserved are thermal traits of reptiles and amphibians?. Journal of Experimental Zoology Part A: Ecological and Integrative Physiology, 335(1), 173-194.

Bogert, C. M. (1949). Thermoregulation in reptiles, a factor in evolution. Evolution, 3(3), 195-211.

Bovo, R. P., Navas, C. A., Tejedo, M., Valença, S. E., & Gouveia, S. F. (2018). Ecophysiology of amphibians: Information for best mechanistic models. Diversity, 10(4), 118.

Bovo, R. P., Simon, M. N., Provete, D. B., Lyra, M., Navas, C. A., & Andrade, D. V. (2023). Beyond Janzen's Hypothesis: How Amphibians That Climb Tropical Mountains Respond to Climate Variation. Integrative Organismal Biology, 5(1), obad009.

Bozinovic, F., Calosi, P., & Spicer, J. I. (2011). Physiological correlates of geographic range in animals. Annual Review of Ecology, Evolution, and Systematics, 42, 155-179.

Breau, C., Cunjak, R. A., & Bremset, G. (2007). Age-specific aggregation of wild juvenile Atlantic salmon Salmo salar at cool water sources during high temperature events. Journal of Fish Biology, 71(4), 1179-1191.

Burraco, P., Laurila, A., & Orizaola, G. (2021). Limits to compensatory responses to altered phenology in amphibian larvae. Oikos, 130(2), 231-239.

Burraco, P., Rendón, M. A., Díaz-Paniagua, C., & Gomez-Mestre, I. (2022). Maintenance of phenotypic plasticity is linked to oxidative stress in spadefoot toad larvae. Oikos, 2022(5), e09078.

Calosi, P., Bilton, D. T., & Spicer, J. I. (2008). Thermal tolerance, acclimatory capacity and vulnerability to global climate change. Biology Letters, 4(1), 99-102.

Campos, D. F., Amanajás, R. D., Almeida-Val, V. M., & Val, A. L. (2021). Climate vulnerability of South American freshwater fish: Thermal tolerance and acclimation. Journal of Experimental Zoology Part A: Ecological and Integrative Physiology, 335(9-10), 723-734.

Capon, S. J., Stewart-Koster, B., & Bunn, S. E. (2021). Future of freshwater ecosystems in a 1.5 C warmer world. Frontiers in Environmental Science, 9, 596.

Carilo Filho, L. M., Gomes, L., Katzenberger, M., Sole, M., & Orrico, V. G. (2022). There and back again: A meta-analytical approach on the influence of acclimation and altitude in the upper thermal tolerance of amphibians and reptiles. Frontiers in Ecology and Evolution, 10, 1017255.

Carpenter, S. R., Fisher, S. G., Grimm, N. B., & Kitchell, J. F. (1992). Global change and freshwater ecosystems. Annual Review of Ecology and Systematics, 23(1), 119-139.

Chang, J., Rabosky, D. L., Smith, S. A., & Alfaro, M. E. (2019). An R package and online resource for macroevolutionary studies using the ray-finned fish tree of life. Methods in Ecology and Evolution, 10(7), 1118-1124.

Chown, S. L., Jumbam, K. R., Sørensen, J. G., & Terblanche, J. S. (2009). Phenotypic variance, plasticity and heritability estimates of critical thermal limits depend on methodological context. Functional Ecology, 23(1), 133-140.

Cicchino , A. S., Shah, A. A., Forester, B. R., Dunham, J. B., Poff, N. L., Ghalambor, C. K., & Funk, W. C. (2023). Acclimation capacity of critical thermal maximum varies among populations: Consequences for estimates of vulnerability. Ecosphere, 14(11), e4691.

Clark, T. D., Scheuffele, H., Pratchett, M. S., & Skeeles, M. R. (2022). Behavioural temperature regulation is a low priority in a coral reef fish (*Plectropomus leopardus*): insights from a novel behavioural thermoregulation system. Journal of Experimental Biology, 225(18), jeb244212.

Claussen, D. L. (1977). Thermal acclimation in ambystomatid salamanders. Comparative Biochemistry and Physiology--Part A: Physiology, 58(4), 333-340.Closs et al. 2016

Comte, L., & Grenouillet, G. (2013). Do stream fish track climate change? Assessing distribution shifts in recent decades. Ecography, 36(11), 1236-1246.

Comte, L., & Olden, J. D. (2017a). Climatic vulnerability of the world's freshwater and marine fishes. Nature Climate Change, 7(10), 718-722.

Comte, L., & Olden, J. D. (2017b). Evolutionary and environmental determinants of freshwater fish thermal tolerance and plasticity. Global Change Biology, 23(2), 728-736.

Cook, C. N., & Sgrò, C. M. (2018). Understanding managers' and scientists' perspectives on opportunities to achieve more evolutionarily enlightened management in conservation. Evolutionary Applications, 11(8), 1371-1388.

Cowan, Z. L., Andreassen, A. H., De Bonville, J., Green, L., Binning, S. A., Silva-Garay, L., ... & Sundin, J. (2023). A novel method for measuring acute thermal tolerance in fish embryos. Conservation Physiology, 11(1), coad061.

Cowles, R. B., & Bogert, C. M. (1944). A preliminary study of the thermal requirements of desert reptiles. Bulletin of the AMNH; v. 83, article 5.

da Silva, A. F., Malhado, A. C., Correia, R. A., Ladle, R. J., Vital, M. V., & Mott, T. (2020). Taxonomic bias in amphibian research: Are researchers responding to conservation need?. Journal for Nature Conservation, 56, 125829.

Dahlke, F. T., Wohlrab, S., Butzin, M., & Pörtner, H. O. (2020). Thermal bottlenecks in the life cycle define climate vulnerability of fish. Science, 369(6499), 65-70.

Dahlke, F., Butzin, M., Wohlrab, S., & Pörtner, H. O. (2022). Reply to: methodological

inconsistencies define thermal bottlenecks in fish life cycle. Evolutionary Ecology, 36(2), 293-

298.Deutsch et al. 2008

Desforges, J. E., Birnie‑Gauvin, K., Jutfelt, F., Gilmour, K. M., Eliason, E. J., Dressler, T. L.,

... & Cooke, S. J. (2023). The ecological relevance of critical thermal maxima methodology for

fishes. Journal of Fish Biology, 102(5), 1000-1016.

Di Santo, V., & Lobel, P. S. (2017). Body size and thermal tolerance in tropical gobies. Journal

of Experimental Marine Biology and Ecology, 487, 11-17.

Duarte, H., Tejedo, M., Katzenberger, M., Marangoni, F., Baldo, D., Beltrán, J. F., ... &

Gonzalez-Voyer, A. (2012). Can amphibians take the heat? Vulnerability to climate warming

in subtropical and temperate larval amphibian communities. Global change biology, 18(2), 412-

421.

Enriquez-Urzelai, U., Sacco, M., Palacio, A. S., Pintanel, P., Tejedo, M., & Nicieza, A. G.

(2019). Ontogenetic reduction in thermal tolerance is not alleviated by earlier developmental

acclimation in *Rana temporaria*. Oecologia, 189, 385-394.

Enriquez-Urzelai, U., Tingley, R., Kearney, M. R., Sacco, M., Palacio, A. S., Tejedo, M., &

Nicieza, A. G. (2020). The roles of acclimation and behaviour in buffering climate change

impacts along elevational gradients. Journal of Animal Ecology, 89(7), 1722-1734.

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: new 1-km spatial resolution climate surfaces

for global land areas. International Journal of Climatology, 37(12), 4302-4315.

Franks, S. J., & Hoffmann, A. A. (2012). Genetics of climate change adaptation. Annual

Review of Genetics, 46, 185-208.

Fry, F. (1947). Effects of the environment on animal activity. Publ. Out. Fish. Res. Lab., 55(68),

1-62.

Ghalambor, C. K., Huey, R. B., Martin, P. R., Tewksbury, J. J., & Wang, G. (2006). Are

mountain passes higher in the tropics? Janzen's hypothesis revisited. Integrative and

comparative biology, 46(1), 5-17.

Griffith, A. W., & Gobler, C. J. (2020). Harmful algal blooms: A climate change co-stressor in

marine and freshwater ecosystems. Harmful Algae, 91, 101590.

Gunderson, A. R., & Leal, M. (2016). A conceptual framework for understanding thermal
constraints on ectotherm activity with implications for predicting responses to global change.
Ecology Letters, 19(2), 111-120.

Gunderson, A. R., & Stillman, J. H. (2015). Plasticity in thermal tolerance has limited potential
to buffer ectotherms from global warming. Proceedings of the Royal Society B: Biological
Sciences, 282(1808), 20150401.

Gunderson, A. R., Armstrong, E. J., & Stillman, J. H. (2016). Multiple stressors in a changing
world: the need for an improved perspective on physiological responses to the dynamic marine
environment. Annual Review of Marine Science, 8, 357-378.

Gutiérrez-Pesquera, L. M., Tejedo, M., Olalla-Tárraga, M. Á., Duarte, H., Nicieza, A., & Solé,
M. (2016). Testing the climate variability hypothesis in thermal tolerance limits of tropical and
temperate tadpoles. Journal of Biogeography, 43(6), 1166-1178.

Gutiérrez-Pesquera, L. M., Tejedo, M., Camacho, A., Enriquez-Urzelai, U., Katzenberger, M.,
Choda, M., ... & Nicieza, A. G. (2022). Phenology and plasticity can prevent adaptive clines in
thermal tolerance across temperate mountains: The importance of the elevation‑time axis.
Ecology and Evolution, 12(10), e9349.

Haddaway, N. R., Page, M. J., Pritchard, C. C., & McGuinness, L. A. (2022). PRISMA2020:
An R package and Shiny app for producing PRISMA 2020-compliant flow diagrams, with
interactivity for optimised digital transparency and Open Synthesis. Campbell Systematic
Reviews, 18(2), e1230.

Haesemeyer, M. (2020). Thermoregulation in fish. Molecular and Cellular Endocrinology, 518,
110986.

Hayward, A., & Gillooly, J. F. (2011). The cost of sex: quantifying energetic investment in
gamete production by males and females. PLoS One, 6(1), e16557.

Hocking, D. J., & Babbitt, K. J. (2014). Amphibian contributions to ecosystem services.
Herpetological Conservation and Biology, 9(1), 1−17.

Hoffmann, A. A., & Sgró, C. M. (2011). Climate change and evolutionary adaptation. Nature,
470: 479–485.

Hothorn, T., & Zeileis, A. (2015). partykit: A modular toolkit for recursive partytioning in R.
The Journal of Machine Learning Research, 16(1), 3905-3909.

Huey, R. B., Berrigan, D., Gilchrist, G. W., & Herron, J. C. (1999). Testing the adaptive significance of acclimation: a strong inference approach. American Zoologist, 39(2), 323-336.

Huey, R. B., Deutsch, C. A., Tewksbury, J. J., Vitt, L. J., Hertz, P. E., Álvarez Pérez, H. J., & Garland Jr, T. (2009). Why tropical forest lizards are vulnerable to climate warming. Proceedings of the Royal Society B: Biological Sciences, 276(1664), 1939-1948.

Huey, R. B., Kearney, M. R., Krockenberger, A., Holtum, J. A., Jess, M., & Williams, S. E. (2012). Predicting organismal vulnerability to climate warming: roles of behaviour, physiology and adaptation. Philosophical Transactions of the Royal Society B: Biological Sciences, 367(1596), 1665-1679.

Hutchison, V. H. (1961). Critical thermal maxima in salamanders. Physiological zoology, 34(2), 92-125.

Illing, B., Downie, A. T., Beghin, M., & Rummer, J. L. (2020). Critical thermal maxima of early life stages of three tropical fishes: Effects of rearing temperature and experimental heating rate. Journal of Thermal Biology, 90, 102582.

IPCC (2021). Summary for Policymakers. In: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge: Cambridge University Press.

ISI WOS, Thomson Scientific, No Date. Web of Science. http://scientific.thomsonreuters.com/products/wos/

Janzen, D. H. (1967). Why mountain passes are higher in the tropics. The American Naturalist, 101(919), 233-249.

Jenkins, C. N., Pimm, S. L., & Joppa, L. N. (2013). Global Patterns of Terrestrial Vertebrate Diversity and Conservation. PNAS 110(28): E2602-E2610. doi: 10.1073/pnas.1302251110

Katzenberger, M., Hammond, J., Duarte, H., Tejedo, M., Calabuig, C., & Relyea, R. A. (2014). Swimming with predators and pesticides: how environmental stressors affect the thermal physiology of tadpoles. PloS one, 9(5), e98265.

Killen, S. S., Adriaenssens, B., Marras, S., Claireaux, G., & Cooke, S. J. (2016). Context dependency of trait repeatability and its relevance for management and conservation of fish populations. Conservation Physiology, 4(1), cow007.

Kingsolver, J. G., & Huey, R. B. (1998). Evolutionary analyses of morphological and
physiological plasticity in thermally variable environments. American Zoologist, 38(3), 545-
560.

Kingsolver, J. G., Arthur Woods, H., Buckley, L. B., Potter, K. A., MacLean, H. J., & Higgins,
J. K. (2011). Complex life cycles and the responses of insects to climate change.

Klockmann, M., Günter, F., & Fischer, K. (2017). Heat resistance throughout ontogeny: body
size constrains thermal tolerance. Global Change Biology, 23(2), 686-696.

Kovacevic, A., Latombe, G., & Chown, S. L. (2019). Rate dynamics of ectotherm responses to
thermal stress. Proceedings of the Royal Society B, 286(1902), 20190174.

Kumar, S., Suleski, M., Craig, J. M., Kasprowicz, A. E., Sanderford, M., Li, M., ... & Hedges,
S. B. (2022). TimeTree 5: an expanded resource for species divergence times. Molecular
Biology and Evolution, 39(8), msac174.

Lechner, E. R., Stewart, E. M., Wilson, C. C., & Raby, G. D (2023). CTmax in brook trout
(Salvelinus fontinalis) embryos shows an acclimation response to developmental temperatures
but is more variable than in later life stages. Journal of Fish Biology, 1–5.

Lee, S., Hung, S. S., Fangue, N. A., Haller, L., Verhille, C. E., Zhao, J., & Todgham, A. E.
(2016). Effects of feed restriction on the upper temperature tolerance and heat shock response
in juvenile green and white sturgeon. Comparative Biochemistry and Physiology Part A:
Molecular & Integrative Physiology, 198, 87-95.

Lee, H., Calvin, K., Dasgupta, D., Krinner, G., Mukherji, A., Thorne, P., ... & Zommers, Z.
(2023). Climate change 2023: synthesis report. Contribution of working groups I, II and III to
the sixth assessment report of the intergovernmental panel on climate change.

Little, A. G., & Seebacher, F. (2015). Temperature determines toxicity: bisphenol A reduces
thermal tolerance in fish. *Environmental Pollution*, *197*, 84-89.

Little, A. G., & Seebacher, F. (2017). Acclimation, acclimatization, and seasonal variation in
amphibians and reptiles. In Amphibian and reptile adaptations to the environment. CRC Press,
41-62.

Lutterschmidt, W. I., & Hutchison, V. H. (1997a). The critical thermal maximum: history and
critique. Canadian Journal of Zoology, 75(10), 1561-1574.

852 Lutterschmidt, W. I., & Hutchison, V. H. (1997b). The critical thermal maximum: data to
853 support the onset of spasms as the definitive end point. Canadian Journal of Zoology, 75(10),
854 1553-1560.

855 McLaughlin, B. C., Skikne, S. A., Beller, E., Blakey, R. V., Olliff-Yang, R. L., Morueta-Holme,
856 N., ... & Zavaleta, E. S. (2022). Conservation strategies for the climate crisis: An update on
857 three decades of biodiversity management recommendations from science. Biological
858 Conservation, 268, 109497.

859 McMenamin, S. K., & Parichy, D. M. (2013). Metamorphosis in teleosts. Current topics in
860 developmental biology, 103, 127-165.

861 Meek, M. H., Beever, E. A., Barbosa, S., Fitzpatrick, S. W., Fletcher, N. K., Mittan-Moreau,
862 C. S., ... & Hellmann, J. J. (2023). Understanding local adaptation to prepare populations for
863 climate change. BioScience, 73(1), 36-47.

864 Mitchell, M., et al, "Engauge Digitizer Software." Webpage:
865 http://markummitchell.github.io/engauge-digitizer, Last Accessed: October 27, 2023

866 Mittermeier, R. A., Turner, W. R., Larsen, F. W., Brooks, T. M., & Gascon, C. (2011). Global
867 biodiversity conservation: the critical role of hotspots. In Biodiversity hotspots: distribution and
868 protection of conservation priority areas. Springer Berlin Heidelberg, 3-22.

869 Molina, A. N., Pulgar, J. M., Rezende, E. L., & Carter, M. J. (2023). Heat tolerance of marine
870 ectotherms in a warming Antarctica. Global Change Biology, 29(1), 179-188.

871 Morgan, R., Andreassen, A. H., Åsheim, E. R., Finnøen, M. H., Dresler, G., Brembu, T., ... &
872 Jutfelt, F. (2022). Reduced physiological plasticity in a fish adapted to stable temperatures.
873 Proceedings of the National Academy of Sciences, 119(22), e2201919119.

874 Morgan, R., Finnøen, M. H., & Jutfelt, F. (2018). CTmax is repeatable and doesn't reduce
875 growth in zebrafish. Scientific Reports, 8(1), 7099.

876 Morley, S. A., Peck, L. S., Sunday, J. M., Heiser, S., & Bates, A. E. (2019). Physiological
877 acclimation and persistence of ectothermic species under extreme heat events. Global Ecology
878 and Biogeography, 28(7), 1018-1037.

879 Moyano, M., Candebat, C., Ruhbaum, Y., Alvarez-Fernandez, S., Claireaux, G., Zambonino-
880 Infante, J. L., & Peck, M. A. (2017). Effects of warming rate, acclimation temperature and

ontogeny on the critical thermal maximum of temperate marine fish larvae. PLoS One, 12(7),
e0179928.

Muñoz, M. M. (2022). The Bogert effect, a factor in evolution. Evolution, 76(s1), 49-66.

Navas, C. A., & Otani, L. (2007). Physiology, environmental change, and anuran conservation.
Phyllomedusa: Journal of Herpetology, 6(2), 83-103.

Navas, C. A., Agudelo-Cantero, G. A., & Loeschcke, V. (2022). Thermal boldness: Volunteer
exploration of extreme temperatures in fruit flies. Journal of Insect Physiology, 136, 104330.

Navas, C. A., Gomes, F. R., & Carvalho, J. E. (2008). Thermal relationships and exercise
physiology in anuran amphibians: integration and evolutionary implications. Comparative
Biochemistry and Physiology Part A: Molecular & Integrative Physiology, 151(3), 344-362.

Newman, R. A. (1992). Adaptive plasticity in amphibian metamorphosis. BioScience, 42(9),
671-678.

Oyamaguchi, H. M., Vo, P., Grewal, K., Do, R., Erwin, E., Jeong, N., ... & Gridi-Papp, M.
(2018). Thermal sensitivity of a Neotropical amphibian (*Engystomops pustulosus*) and its
vulnerability to climate change. Biotropica, 50(2), 326-337.

Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. Bioinformatics, 35(3), 526-528.

Peck, L. S., Morley, S. A., Richard, J., & Clark, M. S. (2014). Acclimation and thermal
tolerance in Antarctic marine ectotherms. Journal of Experimental Biology, 217(1), 16-22.

Penman, R. J., Bugg, W., Rost-Komiya, B., Earhart, M. L., & Brauner, C. J. (2023). Slow heating rates
increase thermal tolerance and alter mRNA HSP expression in juvenile white sturgeon (*Acipenser
transmontanus*). Journal of Thermal Biology, 103599.

Peralta-Maraver, I., & Rezende, E. L. (2021). Heat tolerance in ectotherms scales predictably
with body size. Nature Climate Change, 11(1), 58-63.

Pincebourde, S., Murdock, C. C., Vickers, M., & Sears, M. W. (2016). Fine-scale microclimatic
variation can shape the responses of organisms to global change in both natural and urban
environments. Integrative and Comparative Biology, 56(1), 45-61.

Pinsky, M. L., Eikeset, A. M., McCauley, D. J., Payne, J. L., & Sunday, J. M. (2019). Greater
vulnerability to warming of marine versus terrestrial ectotherms. Nature, 569(7754), 108-111.

Pintanel, P., Tejedo, M., Merino-Viteri, A., Almeida-Reinoso, F., Salinas-Ivanenko, S., López-Rosero, A. C., ... & Gutiérrez-Pesquera, L. M. (2022). Elevational and local climate variability predicts thermal breadth of mountain tropical tadpoles. Ecography, 2022(5), e05906.

Pörtner, H. (2001). Climate change and temperature-dependent biogeography: oxygen limitation of thermal tolerance in animals. Naturwissenschaften, *88*, 137-146.

Pörtner, H. O. (2002). Climate variations and the physiological basis of temperature dependent biogeography: systemic to molecular hierarchy of thermal tolerance in animals. Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology, 132(4), 739-761.

Pörtner, H. O. (2010). Oxygen-and capacity-limitation of thermal tolerance: a matrix for integrating climate-related stressor effects in marine ecosystems. Journal of Experimental Biology, 213(6), 881-893.

Pörtner, H. O., & Peck, M. A. (2010). Climate change effects on fishes and fisheries: towards a cause-and-effect understanding. Journal of Fish Biology, 77(8), 1745-1779.

Pottier, P., Burke, S., Drobniak, S. M., & Nakagawa, S. (2022b). Methodological inconsistencies define thermal bottlenecks in fish life cycle: a comment on Dahlke et al. 2020. Evolutionary Ecology, 36(2), 287-292.

Pottier, P., Burke, S., Drobniak, S. M., Lagisz, M., & Nakagawa, S. (2021). Sexual (in) equality? A meta-analysis of sex differences in thermal acclimation capacity across ectotherms. Functional Ecology, 35(12), 2663-2678.

Pottier, P., Burke, S., Zhang, R. Y., Noble, D. W., Schwanz, L. E., Drobniak, S. M., & Nakagawa, S. (2022a). Developmental plasticity in thermal tolerance: Ontogenetic variation, persistence, and future directions. Ecology Letters, 25(10), 2245-2268.

Potts, L. B., Mandrak, N. E., & Chapman, L. J. (2021). Coping with climate change: phenotypic plasticity in an imperilled freshwater fish in response to elevated water temperature. Aquatic Conservation: Marine and Freshwater Ecosystems, 31(10), 2726-2736.

R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Rohr, J. R., Civitello, D. J., Cohen, J. M., Roznik, E. A., Sinervo, B., & Dell, A. I. (2018). The complex drivers of thermal acclimation and breadth in ectotherms. Ecology Letters, 21(9), 1425-1439.

Rubalcaba, J. G., & Olalla-Tárraga, M. Á. (2020). The biogeography of thermal risk for
terrestrial ectotherms: Scaling of thermal tolerance with body size and latitude. Journal of
Animal Ecology, 89(5), 1277-1285.

Rudolf, V. H., & Rödel, M. O. (2007). Phenotypic plasticity and optimal timing of
metamorphosis under uncertain time constraints. Evolutionary Ecology, 21, 121-142.

Ruthsatz, K., Dausmann, K. H., Paesler, K., Babos, P., Sabatino, N. M., Peck, M. A., & Glos,
J. (2020a). Shifts in sensitivity of amphibian metamorphosis to endocrine disruption: the
common frog (*Rana temporaria*) as a case study. Conservation Physiology, 8(1), coaa100.

Ruthsatz, K., Dausmann, K. H., Peck, M. A., & Glos, J. (2022a). Thermal tolerance and
acclimation capacity in the European common frog (*Rana temporaria*) change throughout
ontogeny. Journal of Experimental Zoology Part A: Ecological and Integrative Physiology,
337(5), 477-490.

Ruthsatz, K., Dausmann, K. H., Peck, M. A., Drees, C., Sabatino, N. M., Becker, L. I., ... &
Glos, J. (2018b). Thyroid hormone levels and temperature during development alter thermal
tolerance and energetics of *Xenopus laevis* larvae. Conservation Physiology, 6(1), coy059.

Ruthsatz, K., Dausmann, K. H., Reinhardt, S., Robinson, T., Sabatino, N. M., Peck, M. A., &
Glos, J. (2019). Endocrine disruption alters developmental energy allocation and performance
in *Rana temporaria*. Integrative and Comparative Biology, 59(1), 70-88.

Ruthsatz, K., Dausmann, K. H., Reinhardt, S., Robinson, T., Sabatino, N. M., Peck, M. A., &
Glos, J. (2020b). Post-metamorphic carry-over effects of altered thyroid hormone level and
developmental temperature: physiological plasticity and body condition at two life stages in
Rana temporaria. Journal of Comparative Physiology B, 190, 297-315.

Ruthsatz, K., Peck, M. A., Dausmann, K. H., Sabatino, N. M., & Glos, J. (2018a). Patterns of
temperature induced developmental plasticity in anuran larvae. Journal of Thermal Biology, 74,
123-132.

Ruthsatz, K., Rakotoarison, A., Razafimampiandra, J. C., Randriamahefa, V. S.,
Rabemananjara, F. C., Rakotondraparany, F., ... & Vences, M. (2022b). Field body
temperatures in Malagasy rainforest frogs. Herpetology Notes, 15, 565-578.

Schiesari, L., Grillitsch, B., & Grillitsch, H. (2007). Biogeographic biases in research and their consequences for linking amphibian declines to pollution. Conservation Biology, 21(2), 465-471.

Schlichting, C. D., & Pigliucci, M. (1998). Phenotypic evolution: a reaction norm perspective. Sinauer associates incorporated. In: Phenotypic evolution: a reaction norm perspective, Sinauer Associates Incorporated, Sunderland, USA.

Seebacher, F., White, C. R., & Franklin, C. E. (2015). Physiological plasticity increases resilience of ectothermic animals to climate change. Nature Climate Change, 5(1), 61-66.

Shah, A. A., Gill, B. A., Encalada, A. C., Flecker, A. S., Funk, W. C., Guayasamin, J. M., ... & Ghalambor, C. K. (2017). Climate variability predicts thermal limits of aquatic insects across elevation and latitude. Functional Ecology, 31(11), 2118-2127.

Sherman, E. (1980). Ontogenetic change in thermal tolerance of the toad *Bufo woodhousii* fowleri. Comparative Biochemistry and Physiology Part A: Physiology, 65(2), 227-230.

Shi, Y. B. (2000). Amphibian metamorphosis: from morphology to molecular biology (No. 19638). Wiley-Liss.

Sinai, N., Glos, J., Mohan, A. V., Lyra, M. L., Riepe, M., Thöle, E., ... & Ruthsatz, K. (2022). Developmental plasticity in amphibian larvae across the world: Investigating the roles of temperature and latitude. Journal of Thermal Biology, 106, 103233.

Sinclair, B. J., Marshall, K. E., Sewell, M. A., Levesque, D. L., Willett, C. S., Slotsbo, S., ... & Huey, R. B. (2016). Can we predict ectotherm responses to climate change using thermal performance curves and body temperatures?. Ecology letters, 19(11), 1372-1385.

Somero, G. N. (2005). Linking biogeography to physiology: evolutionary and acclimatory adjustments of thermal limits. Frontiers in Zoology, 2(1), 1-9.

Somero, G. N. (2010). The physiology of climate change: how potentials for acclimatization and genetic adaptation will determine 'winners' and 'losers'. Journal of Experimental Biology, 213(6), 912-920.

Sørensen, J. G., Kristensen, T. N., & Overgaard, J. (2016). Evolutionary and ecological patterns of thermal acclimation capacity in *Drosophila*: is it important for keeping up with climate change?. Current Opinion in Insect Science, 17, 98-104.

Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2011). Global analysis of thermal tolerance and latitude in ectotherms. Proceedings of the Royal Society B: Biological Sciences, 278(1713), 1823-1830.

Sunday, J. M., Bates, A. E., Kearney, M. R., Colwell, R. K., Dulvy, N. K., Longino, J. T., & Huey, R. B. (2014). Thermal-safety margins and the necessity of thermoregulatory behavior across latitude and elevation. Proceedings of the National Academy of Sciences, 111(15), 5610-5615.

Sunday, J., Bennett, J. M., Calosi, P., Clusella-Trullas, S., Gravel, S., Hargreaves, A. L., ... & Morales-Castilla, I. (2019). Thermal tolerance patterns across latitude and elevation. Philosophical Transactions of the Royal Society B, 374(1778), 20190036.

Terblanche, J. S., Deere, J. A., Clusella-Trullas, S., Janion, C., & Chown, S. L. (2007). Critical thermal limits depend on methodological context. Proceedings of the Royal Society B: Biological Sciences, 274(1628), 2935-2943.

Tewksbury, J. J., Huey, R. B., & Deutsch, C. A. (2008). Putting the heat on tropical animals. Science, 320(5881), 1296-1297.

van der Sleen, P., & Albert, J.S. (2022). Patterns in freshwater fish diversity. In Encyclopedia of Inland Waters (Second edition), Elsevier, Oxford, Mehner, T., & Tockner K. (Eds.), pp. 243-255

van Heerwaarden, B., Kellermann, V., & Sgrò, C. M. (2016). Limited scope for plasticity to increase upper thermal limits. Functional Ecology, 30(12), 1947-1956.

van Heerwaarden, B., & Sgrò, C. M. (2021). Male fertility thermal limits predict vulnerability to climate warming. Nature Communications, 12(1), 2214.

Weaving, H., Terblanche, J. S., Pottier, P., & English, S. (2022). Meta-analysis reveals weak but pervasive plasticity in insect thermal limits. Nature Communications, 13(1), 5292.

Wickham, H., & Wickham, H. (2009). Getting started with qplot. ggplot2: elegant graphics for data analysis, 9-26.

Wilbur, H. M. (1980). Complex life cycles. Annual review of Ecology and Systematics, 11(1), 67-93.

Winter, M., Fiedler, W., Hochachka, W. M., Koehncke, A., Meiri, S., & De la Riva, I. (2016).
Patterns and biases in climate change research on amphibians and reptiles: a systematic review.
Royal Society Open Science, 3(9), 160158.

Woods, H. A., Dillon, M. E., & Pincebourde, S. (2015). The roles of microclimatic diversity
and of behavior in mediating the responses of ectotherms to climate change. Journal of Thermal
Biology, 54, 86-97.

Woodward, G., Perkins, D. M., & Brown, L. E. (2010). Climate change and freshwater
ecosystems: impacts across multiple levels of organization. Philosophical Transactions of the
Royal Society B: Biological Sciences, 365(1549), 2093-2106.

Yu, G., Smith, D. K., Zhu, H., Guan, Y., & Lam, T. T. Y. (2017). ggtree: an R package for
visualization and annotation of phylogenetic trees with their covariates and other associated
data. Methods in Ecology and Evolution, 8(1), 28-36.

7. Yu, S., Wages, M. R., Cobb, G. P., & Maul, J. D. (2013). Effects of chlorothalonil on
development and growth of amphibian embryos and larvae. Environmental pollution,
181, 329-334.
Acknowledgements

We would like to thank the German Research Foundation (DFG) for providing financial support
to KR and PCE for this study and the Universität Hamburg for providing financial support to
KR by the PostDoc1st award. We thank M. Multsch for assisting with the data visualization
and U. Enriquez-Urzelai and M. H. Bernal for providing data on request. We thank Trina
Rytwinski for her guidance on the statistical approach. We are grateful for the valuable
comments provided by two anonymous reviewers. We are particularly grateful to all authors of
the articles included in our systematic literature review, whose effort and work made the present
study possible.

8. Author contributions

Conceptualization: KR and MAP. Methodology: KR, KA, FD, SW, SG, MAP. Data Extraction
and Quality Check: KR, KA, PCE, MLL, FD and SW. Formal Analysis: FD and SW.
Investigation: KR, FD, and SW. Data Curation: KR. Visualization: KR, FD, and SW. Writing
– Original Draft: KR, FD, and MAP. Writing – Review and Editing: all authors. Funding
Acquisition: KR. Project Administration: KR. Supervision: KR and MAP. All authors gave
their final approval for submission.

9. Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or
financial relationships that could be construed as a potential conflict of interest.

10. Statement of Ethics

The authors have no ethical conflicts to disclose.

11. Funding

KR was supported by a PhD award (PostDoc1st) from the University of Hamburg. The German
Research Foundation (DFG) project (459850971; A new perspective on amphibians and global
change: Detecting sublethal effects of environmental stress as agents of silent population
declines) supported KR and PCE. MLL was supported by the São Paulo Research Foundation
in Brazil (FAPESP #2021/10639-5).