Piezo initiates transient production of collagen IV to repair damaged basement
membranes

Aubrie M. Stricker[1], M. Shane Hutson[2], Andrea Page-McCaw[1,3]

[1]Department of Cell and Developmental Biology, Center for Matrix Biology, Program in
Developmental Biology, Vanderbilt University, Nashville, TN, USA

[2]Department of Physics and Astronomy, Department of Biological Sciences, Vanderbilt
University, Nashville, TN, USA

[3]Lead contact: andrea.page-mccaw@vanderbilt.edu

Summary

Basement membranes are sheets of extracellular matrix separating tissue layers and providing mechanical support. Their mechanical properties are determined largely by their most abundant protein, Collagen IV (Col4). Although basement membranes are repaired after damage, little is known about how. To wit, since basement membrane is extracellular it is unknown how damage is detected, and since Col4 is long-lived it is unknown how it is regulated to avoid fibrosis. Using the basement membrane of the adult *Drosophila* midgut as a model, we show that repair is distinct from maintenance. In healthy conditions, midgut Col4 originates from the fat body, but after damage, a subpopulation of enteroblasts we term "matrix menders" transiently express Col4, and Col4 from these cells is required for repair. Activation of the mechanosensitive channel Piezo is required for matrix menders to upregulate Col4, and the signal to initiate repair is a reduction in basement membrane stiffness. Our data suggests that mechanical sensitivity may be a general property of Col4-producing cells.

Keywords: Basement membrane, *Drosophila*, Collagen IV, midgut, Piezo, matrix, tensional stiffness, enteroblasts, homeostasis, damage, repair

Introduction

Extracellular matrix (ECM) is an essential conserved component of all animal tissues. In addition to providing structural support, ECM influences cell adhesion, shape, signaling, and behaviors. When tissues are damaged, ECM needs to be repaired, and many human diseases have pathological damage to ECM[1]. Despite the critical importance of ECM repair, little is known about these cellular and molecular processes. At the most fundamental level, it is unknown how cells detect ECM damage. Moreover, ECM proteins, including collagens, are among the most stable proteins in the body, with the longest-lived exhibiting half-lives on the order of an animal's lifetime[2]. ECM longevity presents a special challenge, as repairing with too much ECM causes fibrosis. In this study, we identify a mechanism cells use to sense damaged matrix, and they respond by transiently producing matrix until repair is completed.

ECM can be subdivided into stromal matrix and basement membrane (basal lamina). Basement membranes, the focus of this study, are the most ancient and conserved ECM, arising along with multicellularity itself[3]. Basement membranes are thin sheets of matrix that underlie epithelia and wrap around muscles and organs. Their most abundant protein is collagen IV (Col4), a heterotrimer that self-assembles *in vitro* into 2D networks and provides structural support[4]. *In vivo*, networks of Col4 are apposed to networks of laminin, and like Col4, laminin self-assembles *in vitro* into a 2D sheet[5]. Self-assembly raises questions about the nature of basement membrane repair. Is there a continuing self-assembly process for maintenance that also functions passively to repair damage? Or is repair an active process, with cells sensing damage and directing repair?

Drosophila offers an excellent model to address questions of basement

membrane homeostasis and repair. Flies have the same basement membrane structure

and composition as mammals, with all the main components but fewer copies of each

one: one collagen heterotrimer, two laminin heterotrimers, one nidogen gene, and one

perlecan gene[6]. The endogenous Col4 a2 (Vkg) has been "trapped" with a GFP exon,

generating a functional Vkg-GFP protein, a tremendously powerful reagent for imaging

and genetics. To analyze basement membrane repair, we developed the *Drosophila*

posterior midgut as a model tissue. The fly gut is a monolayer epithelium derived from

stem cells that generate two lineages, enterocytes and enteroendocrine cells, with

intermediates in between[7–9]. The gut epithelium is surrounded by peristalsis muscles,

and underlying the epithelium and surrounding the muscles are basement membranes

(Fig. 1A). When adult flies ingest dextran sodium sulfate (DSS), it gets visibly lodged in

the gut basement membrane, damages its structure, and reduces its stiffness[10]. After

DSS removal, repair occurs within two days, providing a window for analyzing genetic

requirements. How basement membrane repair is regulated in the fly gut, or in any

tissue, is unclear.

In this study, we find that genes encoding basement membrane components are

required for both repair and maintenance of the adult basement membrane. However,

after damage, the source of Col4 for the midgut basement membrane changes: without

damage, Col4 originates from the fat body, but after damage, local epithelial cells

transiently upregulate Col4 and its crosslinking enzyme Peroxidasin. These "matrix

mender" cells are specialized enteroblasts, and knocking down Col4 in enteroblasts

inhibits gut basement membrane repair. Damage-induced expression of Col4 in matrix

mender cells requires the mechanosensitive channel *Piezo*, suggesting they sense and

respond to tissue mechanics. Indeed, reducing basement membrane stiffness by

reducing collagen crosslinking is sufficient to induce Col4 expression in matrix mender

cells in a Piezo-dependent manner. Interestingly, most or all the Col4 in the gut

originates from Piezo-expressing cells, suggesting that Piezo-based mechano-

regulation may be a general strategy to regulate Col4 production.

Results

Basement membrane repair and maintenance require many of the same

components.

DSS directly damages the posterior midgut basement membrane of *Drosophila*[10].

In brief, after ingestion, DSS is transported through enterocytes to accumulate in the

underlying basement membrane (Fig. 1A). This mechanism is different from mice,

where the epithelium excludes DSS while it is intact[10,11]. In flies, DSS-containing

basement membranes swell, becoming thicker and less dense as assayed by TEM and

super resolution microscopy, and losing their tensile stiffness as assayed in a

mechanical tensile strain assay[10]. The mechanical damage to the gut basement

membranes alters the gut muscle shape. When wrapped by weakened basement

membranes, the gut muscles hyper-contract (Fig. 1A, right side). Identical changes to

gut muscle shape are observed in animals lacking the Col4 covalent-crosslinking

activity of Peroxidasin (Pxn)[10,12,13], and basement membranes from such animals also

have reduced tensile stiffness in the mechanical stretching assay[10,13,14]. Thus,

basement membrane stiffness can be inferred from gut muscle shape, visible either as

outlined by basement membrane (Fig. 1A) or by as stained by actin (Fig. 3F-I). After

removing DSS, basement membrane is repaired in two days, restoring both the

thickness/structure of the basement membrane itself and the aspect ratio of the

muscles[10]. Repair requires basement membrane protein synthesis, including *vkg* and

LanB1, as well as *Peroxidasin* (*Pxn*), suggesting damaged basement membrane is

replaced with new matrix[10].

We asked whether other conserved basement membrane proteins were required

for repair by targeting Perlecan (*trol*), Nidogen (*Ndg*), and the Col4 extracellular

chaperone, *SPARC*. These genes are required for basement membrane assembly

during development, so we knocked them down with dsRNA in 6-day old adults, well

after adult basement membranes were formed. Knockdown was initiated ubiquitously

with *TubP-Gal4* for two days, then flies were fed DSS for two days, then repair was

allowed for two days before the guts were fixed and analyzed. Assessing basement

membrane function by analyzing muscle shape (aspect ratio), we found *Perlecan/trol*,

Ndg, and *SPARC* were all required for repair (Fig. 1B-G, Fig. S1A-C).

To probe the relationship between repair and maintenance, we asked if these

genes were required for basement membrane maintenance, without external damage,

by knocking them down for longer. In the repair assay, the undamaged controls showed

that *trol*, *Ndg* and *SPARC* were not required for basement membrane maintenance over

6 days of knockdown. However, when we knocked down *SPARC*, *trol*, and *Ndg* for 10

days, without external damage, the gut muscles were dysmorphic, indicating that all

three genes are required for maintaining basement membrane over 10 days in adults

(Fig. 1H-M, Fig. S1D-F). In contrast, knocking down non-matrix extracellular proteins

Gbp1 or *Gbp2*[15] had no effect on basement membrane maintenance or repair (Fig.

S1G-R). Thus, basement membrane is replaced faster during repair than during normal

homeostatic maintenance.

Col4 and Peroxidasin are upregulated in some gut cells following damage.

Although basement membranes must be maintained all throughout the body,

damage-triggered repair should be a localized process. The Col4 in many *Drosophila*

tissues originates from a distant organ, the fat body[16,17], and we found that was true

also for the undamaged gut by targeting the GFP moiety in *vkg-GFP/+* heterozygotes.

vkg-GFP/+ gut basement membranes fluorescence (Fig. 2A), but when dsRNA against

GFP was expressed in all cells with *TubP-Gal4*, there was virtually no gut fluorescence

(Fig. 2B). When GFP was knocked down specifically in the fat body with *c564-Gal4*, gut

fluorescence was also extremely low (Fig. 2C,D), suggesting that most of the gut Col4

originates from the fat body, a distant rather than local source. Importantly, the

untagged *vkg* allele was not affected by this knockdown strategy, and *vkg*

heterozygotes are viable and healthy[17], so the gut basement membrane was still

functional.

To identify a potential local source of Col4 after DSS damage, we used *Cg-Gal4*,

where Gal4 is under control of an enhancer that lies between the two head-to-head

promoters of Col4a1 and Col4a2, which together form the single fly Col4 heterotrimer.

In controls, only a few cells faintly expressed *Cg>GFP* in the gut*,* but after damage the

number of *Cg>GFP* cells in the gut increased dramatically, from 1.8% to 5.7% of gut

epithelial cells (Fig. 2E-G). Further, on a per cell basis the level of expression was

higher after damage (Fig. 2H). Interestingly, reporters for *Pxn* and *Laminin* (*Pxn>GFP*

and *LanB1>GFP*) were active in cells of the gut even without basement membrane

damage (Fig. S2), although the number of *Pxn>GFP* cells doubled after damage (Fig.

2I,J). We dubbed these *Cg>GFP* cells "matrix menders". Interestingly, after the gut

basement membrane was repaired, matrix menders were not observed (Fig. 2G). Thus,

in response to basement membrane damage these cells synthesize new Col4, but

synthesis is halted once repair is complete.

Lineage tracing indicates matrix menders are a subset of enteroblasts.

To determine the cellular identity of matrix menders, intestinal stem cells were

stained with anti-Delta: ~70% of *Cg>GFP* cells were adjacent to stem cells (86/122, see

Fig. 3A). This suggested that matrix menders are the daughters of stem cells, either

enteroblasts or pre-enteroendocrine (pre-EE) cells. For lineage-tracing, we turned to the

Gal4 technique for real-time and clonal expression (G-TRACE)[18]. In this system, "real-

time" Col4 expression was reported by *Cg-Gal4* driving *UAS-nlsRFP*, whereas nuclei

that previously expressed Col4 ("lineage expression") were permanently labeled with

nlsGFP (Fig. 3B). Anti-Prospero, which labels pre-EE and enteroendocrine cells, did not

colocalize with either Col4-expressing (RFP+) cells or their lineage (GFP+) (Fig. 3C-

C'''). However, the enterocyte-specific antibody Pdm1 stained about a quarter of GFP+

cells (Fig. 3D-D''', E-E''') after repair, while almost never labeling RFP+ cells. These

results indicate that the matrix menders are enteroblasts, but they stop expressing Col4

before differentiating into enterocytes. Enteroblasts are small protrusive cells that

migrate away from the stem cells (Fig. 3F,G, and [19]) and are in contact with the

basement membrane (Fig. S3A,B), as are the matrix menders (Fig. 3A,H,I). Not all

enteroblasts are matrix menders, however: DSS treatment gives rise to ~2.5 times as

many enteroblasts labeled with *Su(H)>GFP* than matrix mender cells labeled with

Cg>GFP (Fig. S3C).

Matrix menders are dispensable after repair.

G-TRACE gave insight into the lifespan of matrix mender cells after they stopped

expressing collagen. The number of Col4-expressing RFP+ nuclei increased in

response to DSS and then declined after repair (Fig. 3J), similar to Cg>GFP cells (Fig.

2G), although because nuclear RFP lasts longer than cytoplasmic GFP, more RFP+

cells are observed for longer. Contrary to expectations, however, RFP+ cells did not

transition into GFP-only lineage-labeled cells: although RFP+ cells increase in number 2

days after DSS removal, there is no corresponding delayed pulse of GFP+ lineage

label, indicating that most red cells are removed (Fig. 3K). Thus, it appears matrix

mender cells are short-lived after they have completed repair.

Collagen IV from enteroblasts is required for repair.

We asked whether Col4 synthesized by the matrix menders is required to repair

the gut basement membrane by knocking down collagen specifically in enteroblasts and

assaying repair. When *Col4a1* was knocked down in enteroblasts with *Su(H)-Gal4*, only

29% (n = 7) of the animals repaired their gut basement membranes, indicating that the

Col4 from the matrix menders is required for gut basement membrane repair (Fig. 3L).

Piezo is required for basement membrane repair and for Col4 upregulation

How does the gut sense damage to turn on Col4? DSS makes basement

membranes less stiff[10], and we envision most types of matrix damage would result in

reduced stiffness; thus, we reasoned a mechanosensitive mechanism might detect

basement membrane damage. Piezo is a mechanosensitive ion channel active in the

gut[7,20,21], so we tested its role in basement membrane maintenance and repair. *Piezo*^{KO}

null flies are homozygous viable, and the basement membranes in aged *Piezo*^{KO} adult

animals appeared healthy (Fig. 4D-F), indicating no role for Piezo in maintenance.

However, after feeding them DSS for two days, *Piezo*^{KO} flies were unable to repair their

basement membranes (Fig. 4A-C). *Piezo* is the first gene identified required specifically

for repair and not maintenance.

These results indicate that basement membrane repair is triggered by the

presence of damage and detected by a Piezo-based mechanosensitive mechanism. To

determine if Piezo was required for local Col4 production in matrix mender cells, we

quantified the number of *Cg>GFP* cells in *Piezo*^{KO} versus control midguts after DSS

damage. In controls, 5.8% of epithelial midgut cells expressed GFP, but in *Piezo*^{KO} only

1.9% expressed GFP (Fig. 4G-I). In addition to having fewer matrix menders, there was

a 2.4-fold decrease in *Cg>GFP* expression levels in the *Piezo*^{KO} guts (Fig. 4J). Thus,

Piezo is required for Col4 upregulation in response to damage.

Piezo detects decreased basement membrane stiffness to activate matrix

menders.

What aspect of basement membrane damage does Piezo respond to? To test

the stiffness hypothesis, we altered basement membrane stiffness another way, by

inhibiting Pxn crosslinking, known to contribute substantially to basement membrane

tensile stiffness[13,14]. Peroxidasin is irreversibly inhibited by Phloroglucinol (PHG), and

we previously showed that 100 µm PHG decreases the tensile stiffness (viscoelastic

modulus) of adult fly basement membranes by half[13]. PHG-induced decrease in

stiffness was evident in the muscle aspect ratio, and even more pronounced with 5 mM

PHG (Fig. 4M). Excitingly, 100 µM PHG activated *Cg>GFP* in 3.5% of gut epithelial

cells, and 5 mM PHG activated *Cg>GFP* in 5.3%, levels similar to DSS (Fig. 4K-N).

Further, stiffness-induced activation of Col4 in matrix menders required *Piezo*, as

Piezo^KO flies had no increase in *Cg>Gal4* cells with PHG (Fig. 4N-P). Without *Piezo*, the

levels of reporter expression actually fell in response to PHG (Fig S4A,B). These results

indicate that Piezo responds to the reduction in basement membrane stiffness to

activate matrix synthesis.

Piezo must be activated in Col4-expressing cells for repair to occur.

To determine specifically if Piezo activation was required for repair, we used the

inhibitor *Grammostola spatulata* mechanotoxin #4 (GsMTx4) from tarantula toxin, which

inserts in the plasma membrane and reduces local membrane tension, inhibiting

tension-based activation of Piezo and other mechanosensitive channels[22–24]. Although

wild-type GsMTx4 acts non-cell autonomously, the mutant GsMTx4-AP[25] is cell

autonomous. We compared the ubiquitous expression of GsMTx4-AP to homozygous

Piezo^KO mutants for their ability to repair after damage. Only 33% of the GsMTx4-AP

expressing guts were able to repair (n=12), similar to 22% repair in homozygous

Piezo^KO mutants, indicating that Piezo must be activated for basement membrane repair

(Fig. 4Q). Further, repair was completely blocked when GsMTx4-AP was expressed

specifically in Cg-expressing cells, yet repair was not impaired at all when GsMTx4-AP

was expressed in tracheal cells or neurons, cell types that also express Piezo ([26] and

Fig. S4C) and reside in the gut (Fig. 4Q). Thus, basement membrane repair requires

Piezo activation specifically in cells that express Col4.

Piezo positive cells are the source of Col4.

As an orthogonal approach to analyzing the relationship between Piezo and

Col4, we asked whether the collagen from Piezo-expressing cells was incorporated into

the basement membrane of the midgut. We knocked down GFP in a heterozygous *Vkg-*

GFP/+ animal, throughout development into adulthood. As expected, ubiquitously

knocking down GFP resulted in basement membranes with almost no GFP, both in

undamaged and damaged conditions. Interestingly, knocking down GFP with *Piezo-*

Gal4 gave similarly low levels of GFP in gut basement membranes (Fig. 4R-T) –

suggesting that all cells that contribute Col4 to the gut also express Piezo. Similar

results were found after DSS damage, indicating that both in the absence and presence

of damage, midgut Col4 comes from Piezo-expressing cells (Fig. 4T).

Discussion

Using the basement membrane around the *Drosophila* posterior midgut as a

model, we find that basement membrane damage is repaired by a distinct set of local

cells, matrix mender cells, that respond to basement membrane stiffness to initiate Col4

expression. Upregulation of Col4 in matrix menders requires activation of the

mechanosensitive channel Piezo. Because Piezo is known to be activated by a

reduction in membrane tension[27], our results imply that a reduction in basement

membrane stiffness causes stretching of the plasma membrane. The matrix mender

cells have a large area of contact with the basement membrane they monitor (Fig. 3H,I),

and we envision that when the basement membrane tensional stiffness is reduced – i.e.

when the basement membrane stretches more – the plasma membrane of the

neighboring cells is also stretched. Contraction of the gut peristalsis muscles would

exacerbate membrane stretching in the matrix menders.

Lineage tracing indicates that matrix menders are short-lived enteroblasts, with

some differentiating into enterocytes and others removed from the population,

presumably by dying. Knocking down Col4 specifically in enteroblasts prevented

basement membrane repair, confirming that matrix menders are enteroblasts. These

cells synthesize collagen for less than two days after DSS removal, consistent with the

48 h time frame for repair[10], and lineage tracing indicates they stop producing collagen

before they differentiate or die, suggesting that matrix menders are specialized for their

function of repairing the matrix. The short duration of collagen expression is important

because collagen itself is a long-lived molecule, and too much of it would alter tissue

mechanical properties causing fibrosis. Thus, matrix mender cells appear to be

exquisitely balanced to produce enough Col4 for repair, but not too much.

We speculate that the damage status of many basement membranes may be

monitored by Piezo. When Vkg-GFP was knocked down specifically in *Piezo*-expressing

cells, the gut basement membranes incorporated little Vkg-GFP, even the undamaged

ones that get Col4 from the fat body. These data indicate that fat body cells also

express Piezo. Like matrix mender cells in the gut, fat body cells are in contact with a

local basement membrane[28,29]. These results raise the possibility of a general model

that collagen-producing cells use a Piezo-dependent mechanism for

mechanosensitivity. In support of this concept, we previously observed that when Col4

crosslinking is inhibited in larvae, the whole-body levels of Col4 are increased (McCall

et al, Fig S7), a result that supports the idea that Col4 levels respond to of basement

membrane stiffness.

Acknowledgements

We thank K. Elkie Peebles, Kimberly LaFever, Jordyn Sanner Barr, and Leah Caplan

for technical help and Barry Denholm, BDSC, VDRC, and Kyoto *DSC* for fly stocks.

Anti-Pdm1 was a gift from Yang Xiaohang. Work was supported by NIGMS

(R01GM137595 to A.P-M., and F31GM148021 to A.S.)

Author Contributions

AS: Conceptualization, methodology, validation, formal analysis, investigation,

resources, writing and editing, visualization. MSH: Methodology, software. APM:

Conceptualization, methodology, writing and editing, visualization, supervision, project

administration.

Competing interests

The authors declare no competing interests.

Figure Legends

Figure 1. Adult basement membrane repair and maintenance require new matrix

components.

(A) Left: *Drosophila* midgut. The gut tube is enclosed by peristalsis muscles

(magenta), each wrapped in basement membrane (green). Inside the muscle layer, the

gut monolayer epithelium (blue nuclei/DAPI stain) comprises absorptive enterocytes

(light blue), stem cells (brown), enteroblasts (dark blue), and pre-EE and EE cells

(orange), all siting on basement membrane continuous with muscle basement

membrane (green).

Right: DSS damage. DSS (orange dots) is ingested, transported through the

enterocytes, and deposited in the underlying basement membrane where it causes

damage, making basement membrane less dense and less stiff, resulting in visibly

dysmorphic muscles. Basement membranes and muscle shapes are fully repaired 2 d

after DSS is removed. For details see Howard et al., (2019). Micrograph bars, 10 μm.

(B-G) *trol* and *Ndg* are required for basement membrane repair in adult guts. Damage is

quantified as muscle shape. Mean +/- SEM, significance evaluated by ANOVA. Bar, 10

μm.

(H-M) *trol* and *Ndg* are required for basement membrane maintenance without external

damage in adult guts. Significance evaluated by *t*-test.

Figure 2. "Matrix mender" gut cells upregulate Col4 after basement membrane

damage.

(A-D) In undamaged conditions, midgut Col4 comes from the fat body. Surface view of

gut basement membrane in *Vkg-GFP/+* controls (A). Ubiquitous knockdown of GFP

eliminated Vkg-GFP fluorescence in the gut (B). Similar fluorescence loss was observed

with fat-body specific GFP knockdown, using *c564-Gal4* (C,D). Bar, 10 μm.

(E-H) Col4 transcriptional reporter, *Cg-Gal4,* was activated in unknown epithelial cells

(matrix menders) of the posterior midgut in response to DSS-induced basement

membrane damage (E-F). *Cg>GFP* expression stops after repair, 2 d after DSS removal

(G). DSS increases both the number of Cg-expressing cells (G), and the intensity of

expression (H). Bar, 20 μm. Significance evaluated by ANOVA in G and *t*-test in H.

(I,J) Peroxidasin transcriptional reporter, *pxn-Gal4*, was activated in some epithelial

cells of the posterior midgut in response to DSS-induced basement membrane damage,

appearing similar to *Cg-Gal4* expressing cells. Bar, 20 μm.

Figure 3. Matrix menders are enteroblasts that provide Col4 necessary for repair

and die soon after repair is completed.

(A) *Cg>GFP* matrix mender cells localized near but not with Delta-labeled intestinal

stem cells. Bar, 20 μm.

(B) Overview of *Gal4* technique for real-time and clonal expression (G-TRACE).

Adapted from Evans et al., 2009.

(C-E) *Cg>G-TRACE* real-time expressing cells (red) and lineage expressing cells

(green). *Cg>G-TRACE* cells did not colocalize with the EE/pre-EE labeled with anti-

Prospero (C). Some *Cg>G-TRACE* cells colocalized with enterocytes labeled with anti-

Pdm1 after 2d recovery from DSS (D - No recovery, E - 2 d recovery). Bar, 10 μm.

(F-I) Enteroblasts (F-G) and matrix mender cells (H-I) have extensive contact with

basement membrane, both with and without DSS. Actin staining labels muscles

wrapped in basement membrane. Bar, 10 µm.

(J,K) After DSS damage and repair, most *Cg>G-TRACE* real-time expressing cells (red,

J) do not transition into GFP-only lineage expressing cells (green, K), indicating that Cg-

expressing cells are lost from the epithelium after repair. All real-time expressing cells

(red) whether or not they were also lineage labeled (green) are shown in J; only green-

labeled cells, not red, are shown in K. Together, panels J and K represent each cell only

once.

(L) Col4 from enteroblasts is required for repair. Percentage of guts with healthy

basement membrane morphology when *Col4a1* was knocked down in enteroblasts

using *Su(H)-Gal4*. Flies were fed DSS or not, followed by 2 d of recovery. Morphology

was assessed on a per-gut basis by the muscle shape and scored blinded to sample

identity.

Figure 4. Piezo is required for basement membrane repair, by activating Col4

synthesis in the matrix mender cells.

(A-F) *Piezo* is required for basement membrane repair in adult guts, assayed 2 d after

removing DSS, but not required for basement membrane maintenance, assayed in

aged 10 d *Piezo^{KO}* adults with no external damage. Basement membrane and muscle

shape visualized by Vkg-GFP. Mean +/-SEM. Significance evaluated by ANOVA (C) or

t-test (F). Bar, 10 µm.

(G-J) *Piezo* is required for matrix mender cells to express Col4 (*Cg>GFP*) in response

to DSS damage. The number of Col4-expressing cells (*Cg>GFP)* and the level of

expression were both significantly reduced in *Piezo^{KO}* flies. Significance evaluated by

ANOVA (I) and *t-test* (J). Bar, 20 μm.

(K-P) The crosslinking inhibitor PHG reduced basement membrane stiffness, evidenced

by dysmorphic muscles (M) and tensile stiffness measurements (Peebles et al, 2023).

PHG-induced changes to membrane stiffness activated matrix mender cells (K,L,N), in

a Piezo-dependent manner (N-P) Significance evaluated by ANOVA. Bar, 20 μm.

(Q) *Piezo* is required in Col4-expressing cells for basement membrane repair. Piezo

activity was inhibited in indicated cell populations with GsMTx4. Flies were fed DSS or

not for 2 d, followed by 2 d of recovery. Repair was assessed on a per-gut basis by the

muscle shape and scored blinded to sample identity.

(R-T) Even without damage, Col4 in the gut basement membrane is synthesized in cells

that also express *Piezo*. Surface view of *Vkg-GFP/+* heterozygous control guts (R,S);

Vkg-GFP levels are almost undetectable when GFP is knocked down with *Piezo-Gal4*

(S,T). Bar, 20 μm. Significance evaluated by ANOVA.

Materials and Methods

Fly husbandry

Flies were maintained on cornmeal-molasses food at 25° C unless otherwise

specified. For temperature-sensitive RNAi studies under *Gal80^{ts}* control, flies were

maintained at 18° C until they eclosed. Female flies were then aged at 18° C until they

were 6-day adults (mated) before moving them to 29° C to activate the Gal4. Fly stocks

used were:

Genotype	Source
VkgGFP^{791}; tubGal4, tubGal80/S-T	this study
UAS-Nidogen^{RNAi}	VDRC 109625
UAS-Perlecan^{RNAi}	VDRC 105502
UAS-SPARC^{RNAi}	BSC 40885
UAS-Gbp1^{RNAi}	VDRC KK168755
UAS-Gbp2^{RNAi}	VDRC GD16696
UAS-GFP	BSC 1522
UAS-mCD8 GFP	Kyoto 108068
Cg-Gal4	BSC 7011
Pxn-Gal4/TM3	BSC 22805
LanB1-Gal4/CyO	BSC 63941
UAS-Stinger	Cory Evans
VkgGFP^{791}	Carnegie Protein trap library, CC00971
Piezo^{KO}	BSC 58770
Piezo^{KO}, Cg-Gal4/CyO	this study
Piezo^{KO}; UAS-GFP/TM3	this study
Piezo^{KO}, VkgGFP^{791}	this study
w^{1118}	Todd Laverty, Janelia Farm
UAS-GsMTx4 (active peptide)	Denholm Lab
Elav-Gal4	BSC 8765
Btl-Gal4	Greg Beitel
VkgGFP, UAS-GFP^{RNAi}/CyO, mCh	BSC 9330
Tubulin-Gal4	Kyoto 108069
Piezo-Gal4	BSC 59266
UAS-Piezo^{RNAi}	VDRC 2796

UAS-Cg25c^{RNAi}	VDRC 28369
Su(H)Gal4, UAS-mCD8GFP	BSC 87733

399 DSS feeding

As described in Amcheslavsky et al. (2009) and Howard et al. (2019), a 2.5 cm x

3 cm piece of chromatography paper (Whatman 3030-861, Grade 3 MM CHR) was

placed in an empty vial. 500 µl of a 5% sucrose solution with or without 3% 36-50 kDa

DSS (dextran sulfate sodium salt colitis grade, MP Biomedicals, CAS number 9011-18-

1, 36,000 – 50,000 MW) was added to the paper in the vial. Flies utilizing the Gal80^{ts}

system were moved to 29° C for 2 days to ensure the Gal4 was active prior to starting

DSS feeding treatments. Flies not utilizing the Gal80^{ts} system were aged 3-5 days at

25° C prior to starting DSS feeding treatments. Female flies were anesthetized and

carefully placed in the vial to ensure they did not contact the liquid. Flies were added to

new vials with fresh media each day for 2 days. For recovery experiments, flies were

transferred to a vial containing standard cornmeal-molasses food for 2 days. For

imaging in 3 dimensions (Z-stacks), flies on standard cornmeal-molasses food were fed

5% sucrose for 4 hours before dissection to reduce autofluorescence from the food.

413 Phloroglucinol feeding

414   Non-cornmeal molasses food was freshly made with or without 100 µM or 5 mM

415 Phloroglucinol (PHG) (Sigma-Aldrich, CAS number 108-73-6) and solidified in a vial

416 overnight. 20 aged female flies and 5 male flies were added to the vials with or without

417 PHG. Flies were kept at 25° C for 5 days. Prior to dissection, flies were fed 5% sucrose

418 for 4 hours before dissection to reduce autofluorescence from the food.

419 CNA35 staining

Immediately after gut dissections, CNA35 green monomer (1:100) was added to

guts for 2 hours, followed by a 1 hour fixation with 8% PFA.

Gut dissections and preparations

Adult females were placed in cold Grace's media and pinched between the

abdomen and thorax using sharp #5 dissecting forceps (Dunmont). The abdomen was

separated from the thorax, and the integument surrounding the abdomen was peeled

away, exposing the gut and Malpighian tubules, being careful not to pinch or stretch the

gut. The Grace's media was removed using a Pasteur pipette, followed by one quick

wash with 1x PBS (137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4, and 1.8 mM KH_2PO_4).

The guts were fixed with 4% paraformaldehyde (Ted Pella, Paraformaldehyde 16%,

Product #18505) in PBS for 20 minutes at room temperature, or 8% paraformaldehyde

in PBS for 60 minutes at room temperature, followed by 3 quick washes in PBS, and 4 x

15 min washes with PBS or PBT.

For immunofluorescence experiments, blocking buffer was added to fixed guts

for 2 hours at room temperature or overnight at 4° C then incubated in primary antibody

and 0.2 µM SiRActin (Spirochrome Cat# SC001) in PBS or PBT overnight at 4° C,

followed by 4 x 30 min washes with 1x PBS. Secondary antibodies were added for 2

hours at room temperature, followed by 4 x 30 min washes. Guts were then mounted in

DAPI-containing mounting media (Vectashield, Vector Laboratories, H1200). Antibodies

used were:

• Rabbit anti-GFP (1:200, Torrey Pines, TP401)

• Mouse IgG1 anti-Delta (1:50, DSHB SA4)

• Mouse IgG1 anti-Prospero (1:50, DSHB MR1A)

• Rabbit anti-Pdm1 (1:1000, gift from Yang Xiaohang)

• Cy3 goat anti-mouse IgG1 (1:200, Jackson ImmunoResearch Laboratory Inc.)

• Cy3 donkey anti-rabbit (1:200, Jackson ImmunoResearch Laboratory Inc.)

• Cy5 goat anti-mouse 1gG1 (1:200, Jackson ImmunoResearch Laboratory Inc.)

• Cy5 donkey anti-rabbit (1:200, Jackson ImmunoResearch Laboratory Inc.)

Light Microscopy

Single optical sections of the guts were imaged using a Zeiss Apotome mounted

to an Axio Image M2 with a 63x/1.4 oil Plan-Apochromat objective. Z-stacks were

acquired using a 40x/1.3 oil Plan-Apochromat objective at a 0.75 µm step size taken

through the top half of the gut tube (lumen through the top of the circumferential muscle

layer). Images were taken using an AxioCam MRm camera (Zeiss), X-Cite 120Q light

source (Excelitas Technologies), and AxioVision 4.8 software. ImageJ (version 1.53v,

National Institutes of Health) 16-bit, ZVI files were used for analysis and figures for

images.

The fluorescence intensity of the basement membrane was determined by taking

single optical slices *en face* through the middle of the circumferential muscles to ensure

the same plane of the basement membrane was imaged, and the exposures were kept

the same. The fluorescence intensity of the basement membrane was measured in

ImageJ using the "box" tool to measure the average intensity of 3 representative regions

of the gut basement membrane. Similarly, the background intensity was determined by

subtracting the average of 3 boxes outside of the gut region. An AVOVA test was

performed (GraphPad Prism 9.5).

466   Fluorescence intensity measurements of gut cells to be compared were acquired

467 at the same exposure from at least 3 experiments done on different days. Maximum

468 intensity projections were created using ImageJ and the line tool was used to draw

469 through the center of the 3 brightest nuclei in the GFP channel and the fluorescence

470 intensity was measured and averaged. Background fluorescence was subtracted by

471 taking the average intensity of 3 lines within the gut region that did not overlap with GFP

472 nuclei. A *t-test* was performed.

474 **Measurement of muscle shape**

475   Individual muscle cells were identified and segmented using the software

476 Seedwater Segmenter (Mashburn et al. 2012). Custom code in Mathematica (Wolfram

477 Inc., Champaign, IL) was then used to analyze the segmented images and determine

478 the aspect ratio of each muscle cell.

480 **Quantifying number of cells**

481   GFP+ cells were manually quantified by creating a maximum intensity projection

482 and confirmed by the presence of a nucleus. The total number of nuclei in the posterior

483 midgut was quantified by creating a maximum intensity projection and setting a

484 threshold to select the DAPI-stained nuclei. Images were "despeckled," and a

485 watershed was applied to the image. Using the ImageJ "analyze particle" feature, the

486 number of particles larger than 10 μm^2 was quantified. The areas where nuclei were

487 fused together or unable to be accurately quantified by ImageJ were manually

488 quantified.

References

1. Iozzo, R.V., and Gubbiotti, M.A. (2018). Extracellular matrix: The driving force of mammalian diseases. Matrix Biology *71–72*, 1–9. 10.1016/j.matbio.2018.03.023.

2. Liu, P., Xie, X., and Jin, J. (2020). Isotopic Nitrogen-15 Labeling of Mice Identified Long-lived Proteins of the Renal Basement Membranes. Scientific Reports *10*, 1–7. 10.1038/s41598-020-62348-6.

3. Fidler, A.L., Vanacore, R.M., Chetyrkin, S.V., Pedchenko, V.K., Bhave, G., Yin, V.P., Stothers, C.L., Rose, K.L., McDonald, W.H., Clark, T.A., et al. (2014). A unique covalent bond in basement membrane is a primordial innovation for tissue evolution. Proceedings of the National Academy of Sciences of the United States of America *111*, 331–336. 10.1073/pnas.1318499111.

4. Yurchenco, P.D., and Furthmayr, H. (1984). Self-Assembly of Basement Membrane Collagen. Biochemistry *23*, 1839–1850. 10.1021/bi00303a040.

5. Yurchenco, P.D., Tsilibary, E.C., Charonis, A.S., and Furthmayr, H. (1986). Models for the self-assembly of basement membrane. Journal of Histochemistry and Cytochemistry *34*, 93–102. 10.1177/34.1.3510247.

6. Ramos-Lewis, W., and Page-McCaw, A. (2019). Basement membrane mechanics shape development: Lessons from the fly. Matrix Biology *75–76*, 72–81. 10.1016/j.matbio.2018.04.004.

7. He, L., Si, G., Huang, J., Samuel, A.D.T., and Perrimon, N. (2018). Mechanical regulation of stem cell differentiation through stretch-activated Piezo channel. Nature *555*, 103–106. 10.1038/nature25744.

8. Martin, J.L., Sanders, E.N., Moreno-Roman, P., Koyama, L.A.J., Balachandra, S., Du, X., and O'Brien, L.E. (2018). Long-term live imaging of the Drosophila adult midgut reveals real-time dynamics of division, differentiation and loss. eLife *7*. 10.7554/ELIFE.36248.

9. Micchelli, C.A., and Perrimon, N. (2006). Evidence that stem cells reside in the adult Drosophila midgut epithelium. Nature *439*, 475–479. 10.1038/nature04371.

10. Howard, A.M., Lafever, K.S., Fenix, A.M., Scurrah, C.R., Lau, K.S., Burnette, D.T., Bhave, G., Ferrell, N., and Page-McCaw, A. (2019). DSS-induced damage to basement membranes is repaired by matrix replacement and crosslinking. Journal of Cell Science *132*. 10.1242/jcs.226860.

11. Shimshoni, E., Adir, I., Afik, R., Solomonov, I., Shenoy, A., Adler, M., Puricelli, L., Sabino, F., Savickas, S., Mouhadeb, O., et al. (2021). Distinct extracellular–matrix remodeling events precede symptoms of inflammation. Matrix Biology *96*, 47–68. 10.1016/j.matbio.2020.11.001.

12. McCall, A.S., Cummings, C.F., Bhave, G., Vanacore, R., Page-Mccaw, A., and Hudson, B.G. (2014). Bromine is an essential trace element for assembly of collagen IV scaffolds in tissue development and architecture. Cell *157*, 1380–1392. 10.1016/j.cell.2014.05.009.

13. Peebles, K.E., LaFever, K.S., Page-McCaw, P.S., Colon, S., Wang, D., Stricker, A.M., Ferrell, N., Bhave, G., and Page-McCaw, A. (2024). Peroxidasin is required for full viability in development and for maintenance of tissue mechanics in adults. Matrix Biology *125*, 1–11. 10.1016/j.matbio.2023.11.005.

14. Bhave, G., Colon, S., and Ferrell, N. (2017). The sulfilimine cross-link of collagen IV contributes to kidney tubular basement membrane stiffness. American Journal of Physiology - Renal Physiology *313*, F596–F602. 10.1152/ajprenal.00096.2017.

15. O'Connor, J.T., Stevens, A.C., Shannon, E.K., Akbar, F.B., LaFever, K.S., Narayanan, N.P., Gailey, C.D., Hutson, M.S., and Page-McCaw, A. (2021). Proteolytic activation of Growth-blocking peptides triggers calcium responses through the GPCR Mthl10 during epithelial wound detection. Developmental Cell *56*, 2160-2175.e5. 10.1016/j.devcel.2021.06.020.

16. Pastor-Pareja, J.C., and Xu, T. (2011). Shaping Cells and Organs in Drosophila by Opposing Roles of Fat Body-Secreted Collagen IV and Perlecan. Developmental Cell *21*, 245–256. 10.1016/j.devcel.2011.06.026.

17. Ramos-Lewis, W., LaFever, K.S., and Page-McCaw, A. (2018). A scar-like lesion is apparent in basement membrane after wound repair in vivo. Matrix Biology *74*, 101–120. 10.1016/j.matbio.2018.07.004.

18. Evans, C.J., Olson, J.M., Ngo, K.T., Kim, E., Lee, N.E., Kuoy, E., Patananan, A.N., Sitz, D., Tran, P.T., Do, M.T., et al. (2009). G-TRACE: Rapid Gal4-based cell lineage analysis in Drosophila. Nature Methods *6*, 603–605. 10.1038/nmeth.1356.

19. Antonello, Z.A. (2015). Robust intestinal homeostasis relies on cellular plasticity in enteroblasts mediated by miR-8–Escargot switch. The EMBO Journal *34*, 2025–2041. 10.15252/embj.201591517.

20. Min, S., Oh, Y., Verma, P., Whitehead, S.C., Yapici, N., Van Vactor, D., Suh, G.S.B., and Liberles, S.D. (2021). Control of feeding by piezo-mediated gut mechanosensation in Drosophila. eLife *10*, 1–30. 10.7554/eLife.63049.

21. Lee, J., Cabrera, A.J.H., Nguyen, C.M.T., and Kwon, Y.V. (2020). Dissemination of Ras V12-transformed cells requires the mechanosensitive channel Piezo. Nature Communications *11*, 1–13. 10.1038/s41467-020-17341-y.

22. Suchyna, T.M., Johnson, J.H., Hamer, K., Leykam, J.F., Gage, D.A., Clemo, H.F., Baumgarten, C.M., and Sachs, F. (2000). Identification of a peptide toxin from Grammostola spatulata spider venom that blocks cation-selective stretch-activated channels. Journal of General Physiology *115*, 583–598. 10.1085/jgp.115.5.583.

23. Suchyna, T.M., Tape, S.E., Koeppe, R.E., Andersen, O.S., Sachs, F., and Gottlieb,
P.A. (2004). Bilayer-dependent inhibition of mechanosensitive channels by
neuroactive peptide enantiomers. Nature *430*, 235–240. 10.1038/nature02743.

24. Bae, C., Sachs, F., and Gottlieb, P.A. (2011). The Mechanosensitive Ion Channel
Piezo1 is Inhibited by the Peptide GsMTx4. Biochemistry *50*, 6295–6300.
10.1021/bi200770q.

25. Beqja, D., Haidar, S., Nikolaev, M., Shen, Y., and Denholm, B. (2020). Transgenic
Tarantula Toxin: A novel tool to study mechanosensitive ion channels in Drosophila.
Journal of Insect Physiology *127*, 104116. 10.1016/j.jinsphys.2020.104116.

26. Song, Y., Li, D., Farrelly, O., Miles, L., Li, F., Kim, S.E., Lo, T.Y., Wang, F., Li, T.,
Thompson-Peer, K.L., et al. (2019). The Mechanosensitive Ion Channel Piezo
Inhibits Axon Regeneration. Neuron *102*, 373-389.e6.
10.1016/j.neuron.2019.01.050.

27. Coste, B., Xiao, B., Santos, J.S., Syeda, R., Grandl, J., Spencer, K.S., Kim, S.E.,
Schmidt, M., Mathur, J., Dubin, A.E., et al. (2012). Piezo proteins are pore-forming
subunits of mechanically activated channels. Nature *483*, 176–181.
10.1038/nature10812.

28. Dai, J., Ma, M., Feng, Z., and Pastor-Pareja, J.C. (2017). Inter-adipocyte Adhesion
and Signaling by Collagen IV Intercellular Concentrations in Drosophila. Current
Biology *27*, 2729-2740.e4. 10.1016/j.cub.2017.08.002.

29. Shahab, J., Baratta, C., Scuric, B., Godt, D., Venken, K.J.T., and Ringuette, M.J.
(2015). Loss of SPARC dysregulates basal lamina assembly to disrupt larval fat
body homeostasis in Drosophila melanogaster. Developmental Dynamics *244*, 540–
552. 10.1002/dvdy.24243.

Graphical abstract

Healthy basement membrane

Repair

3. Basement membrane is repaired, collagen IV turns off, matrix mender cells differentiate or die

1. Damage decreases basement membrane stiffness

Damage

2. Piezo detects basement membrane stretching, activates Collagen IV synthesis in local matrix mender cells

Detection

Figure 1

A

Posterior midgut structure

Basement membrane

Muscle Actin

Epithelial nuclei

Lumen

Body cavity

Healthy basement membrane (green) and muscles (magenta)

Ingested DSS damages gut basement membranes

Lumen   DSS

Basement membrane

Basement membrane

Repair

Damaged basement membrane

Repair: 2 d after DSS damage

trol^RNAi

B *control*

C *trol^RNAi*

D

Muscle Shape (h/w)

Control (4 guts)   No DSS   *trol^RNAi* (4 guts)   Control (3 guts)   DSS feeding   *trol^RNAi* (6 guts)

***

Ndg^RNAi

E *control*

F *Ndg^RNAi*

G

Muscle Shape (h/w)

Control (9 guts)   No DSS   *Ndg^RNAi* (9 guts)   Control (8 guts)   DSS feeding   *Ndg^RNAi* (10 guts)

****

Maintenance: 10 d knockdown

H *control*

I *trol^RNAi*

J

Muscle Shape (h/w)

Control (4 guts)   *trol^RNAi* (5 guts)   10 d KD

**

K *control*

L *Ndg^RNAi*

M

Muscle Shape (h/w)

Control (10 guts)   *Ndg^RNAi* (12 guts)   10 d KD

****

Figure 2

Figure 3

Figure 4

Repair requires Piezo

A control — Vkg-GFP
B Piezo^{KO}
C Muscle Shape (h/w) — **** — Control (7 guts), Piezo^{KO} (10 guts), No DSS; Control (5 guts), Piezo^{KO} (9 guts), DSS feeding

Maintenance does not require Piezo

D control
E Piezo^{KO}
F Muscle Shape (h/w) — ns — Control (11 guts), Piezo^{KO} (10 guts), 10 d aged

Matrix menders require Piezo

G DSS — Cg>GFP
H DSS — Piezo^{KO},Cg>GFP
I # gut cells expressing Col4 — # Cg>GFP cells/100 — **** — ** — No DSS, DSS (Cg>GFP); No DSS, DSS (Piezo^{KO}, Cg>GFP)
J Col4 reporter intensity — GFP intensity (A.U.) — * — Cg>GFP DSS, Piezo^{KO}, Cg>GFP DSS

Decreased matrix stiffness activates matrix menders via Piezo

K No PHG — Cg>GFP
L 5 mM PHG — Cg>GFP
M Muscle Shape (h/w) — **** — *** — * — No PHG, 100 µM PHG, 5 mM PHG
N # gut cells expressing Col4 — Cg>GFP cells/100 — ns — ** — ** — ns — No PHG, 100 µM PHG, 5 mM PHG (Cg>GFP); No PHG, 5 mM PHG (Piezo^{KO}, Cg>GFP)
O No PHG — Piezo^{KO}, Cg>GFP
P 5 mM PHG — Piezo^{KO}, Cg>GFP
Q Piezo needs to be activated in Col4-expressing cells — % guts undamaged/repaired (scored blinded) — DSS: − + — n=9, n=10 ∅>GsMTx4 (no expression); n=7, n=12 Tub>GsMTx4 (all cells); n=8, n=7 (0) Cg>GsMTx4 (Collagen IV-expressing cells); n=8, n=8 Elav>GsMTx4 (neurons); n=6, n=7 Btl>GsMTx4 (trachea); n=10, n=10 Vkg-GFP; n=9, n=9 Vkg-GFP, Piezo^{KO}

Col4 is from Piezo+ cells

R No KD — Gut, Malpighian tubule
S Piezo>GFP^{RNAi} — Gut, Malpighian tubule — Vkg-GFP DAPI
T No DSS — ** ; DSS — **** — Fluorescence Intensity (A.U.) — No KD, TubP>GFP^{RNAi}, Piezo>GFP^{RNAi} (Vkg-GFP KD); No KD, TubP>GFP^{RNAi}, Piezo>GFP^{RNAi} (Vkg-GFP KD)