The everting *Drosophila* wing disc is a shape-programmed material

Jana F. Fuhrmann[1,2], Abhijeet Krishna[1,2,3], Joris Paijmans[4], Charlie Duclut[4,5], Suzanne Eaton[1,2,3,6], Marko Popovic[2,3,4], Frank Julicher[2,3,4], Carl D. Modes[1,2,3*] and Natalie A. Dye[1,2,7*]

[1*]Max-Planck Institute for Molecular Cell Biology and Genetics, MPI-CBG, Pfotenhauerstrasse 108, Dresden, 01307, Germany.
[2]Excellence Cluster, Physics of Life, Technische Universität Dresden, Arnoldstrasse 18, Dresden, 01307, Germany.
[3]Center for Systems Biology, Pfotenhauerstrasse 108, Dresden, 01307, Germany.
[4]Max-Planck Institute for Physics of Complex Systems, MPI-PKS, Nöthnitzer Str. 38, Dresden, 01187, Germany.
[5]Laboratoire Physico-Chimie Curie, CNRS UMR 168, Institut Curie, Université PSL, Sorbonne Université, Paris, 75005, France.
[6]Biotechnologisches Zentrum, Technische Universität Dresden, Tatzberg 47-49, Dresden, 01307, Germany.
[7]Mildred Scheel Nachwuchszentrum P2, Medical Faculty, Technische Universität Dresden, Dresden, 01307, Germany.

*Corresponding author(s). E-mail(s): modes@mpi-cbg.de; natalie_anne.dye@tu-dresden.de;

Keywords: morphogenesis, Drosophila, wing disc, shape programming, cell rearrangements, MyoVI

Abstract

How complex 3D tissue shape emerges during animal development remains an important open question in biology and biophysics. Although it is well-established that tissue-scale patterns of in-plane cell behaviors can reshape a tissue in 2D, whether and how such patterns could also deform a tissue in 3D remains unclear. Meanwhile, recent advances in materials science, physics, and engineering have enhanced our ability to craft and understand inanimate "shape-programmable" materials capable of undergoing blueprinted 3D shape transformations arising from in-plane gradients of spontaneous strains. Here, we use this conceptual framework to understand eversion of the *Drosophila* wing disc, a complex 3D tissue reshaping event in wing development. Using quantification of spatial patterns of cell dynamics on an evolving 3D tissue surface, combined with physical modeling and genetic perturbation, we demonstrate that the wing can deform itself in 3D through large-scale patterning of in-plane cellular behaviors, primarily rearrangements. This work shows that concepts from shape-programming can be employed to understand animal tissue morphogenesis and suggests that there exist intricate patterns in nature that could present novel designs for shape-programmable materials.

Introduction

Epithelial tissues are sheets of tightly connected cells with apical-basal polarity that form the basic architecture of many animal organs. Deformations of animal epithelia in 3D can be mediated by external forces, either from neighboring tissue that induces buckling instabilities (e.g., [1–3]) or extracellular matrix that confines (e.g., [4]) or expands (e.g., [5]). Alternatively, local differences in mechanics at the apical and basal sides of the deforming epithelia itself can drive out-of-plane tissue shape changes (e.g., ventral furrow invagination in the *Drosophila* embryo (reviewed in [6]) and fold formation in *Drosophila* imaginal discs [7]).

Here, we describe a mechanism for generating complex 3D tissue shape involving tissue-scale patterning of in-plane deformations, analogous to the shape transformations of certain inanimate shape-programmable materials. These shape programmable materials experience *spontaneous strains*, where the local preferred lengths change in response to stimuli in a desired way. Globally patterned spontaneous strains can create a geometric incompatibility with the original shape, triggering a specific, desired 3D deformation ([8–10]). These ideas have already proved insightful to the understanding of differential growth mediated plant morphogenesis [11, 12]. Animal epithelia are more dynamic, however, able to change cell shape and size, as well as rearrange tissue topology. As these behaviours cause in-plane changes in local tissue dimensions, the ingredients for shape-programmability are, in principle, present.

To test these concepts in animal morphogenesis, we quantify tissue shape changes and cell behaviors in the *Drosophila* wing disc during a 3D morphogenetic process called eversion (Fig. 1a). Through eversion, the wing disc proper, an epithelial mono-layer, undergoes a shape deformation in which the future dorsal and ventral surfaces of the wing blade appose to form a bi-layer and escape the overlying squamous epithelium called the peripodial membrane. This process is triggered by a peak in circulating levels of the hormone 20-hydroxyecdysone (20E), analogous to an activator in shape-programming. This complex tissue shape change is independent of forces external to the wing disc, as demonstrated by its ability to occur in explant culture [13]. The shape changes of the disc proper also cannot be fully explained by removal of the peripodial membrane or extracellular matrix and appear to be self-sufficient, involving active cellular processes [14–18]. It has long been postulated that the eversion of wing (and leg) discs involves cell rearrangements and/or cell shape changes that resolve previously established morphological patterns [19–22]. Here, we test these hypotheses by systematic quantification and genetic perturbation of cell behaviors during eversion and demonstrate how cell behaviors contribute to tissue shaping using a physical model analogous to shape programming.

Although the mechanism presented here explains a specific morphogenetic process during *Drosophila* wing development, it has the potential to be a general mechanism of epithelial shaping. In addition, developing such a data-driven pipeline, exploiting both gradients in isotropic deformation as well as the anisotropic component of spontaneous strain, advances our ability to understand shape-programmable systems in general.

1 The wing pouch undergoes anisotropic curvature changes during eversion

We first sought to characterize the tissue shape changes happening during wing disc eversion. To this end, we explanted wing discs at fixed time intervals, from late larval stage (wL3) to 6 hours After Puparium Formation (hAPF). We imaged the wing discs using multi-angle light sheet microscopy and then reconstructed and analyzed the 3D image stack [23]. In this way, we capture the complex 3D shape changes happening throughout the wing disc during eversion (Fig. 1b, Supplementary Video 1). Focusing on the prospective wing blade region (so-called wing disc pouch), the most dramatic tissue shape changes can be seen in a central cross-section along the axis perpendicular to the dorsal ventral boundary (DVB), referred to as "across-DVB" (Fig. 1c,d, Extended Data Fig. S1b,c). We observe that the tissue becomes flatter as the dorsal and ventral surfaces appose, with curvature increasing at the DVB. In the perpendicular plane, taken through DVB (referred to as "along-DVB"), the tissue does not change as significantly, preserving curvature in this direction (Extended Data Fig. S1c-e, S5b,d). These data provide a quantitative view

of the tissue shape changes taking place during eversion, demonstrating the development of strong shape anisotropy that is robust from wing to wing.

Fig. 1 **a**, Schematic cross-sections along the long axis of the wing disc before and after eversion. Before eversion, the wing disc resembles an epithelial sac with apical facing inwards. The tissue consists of the Disc Proper (DP), which is a folded, thick, pseudo-stratified monolayer, and the Peripodial Membrane (PM), a thin squamous monolayer. After eversion, the PM is removed and the former pouch region of the DP forms the wing bilayer with apical facing out and dorsal and ventral on opposing sides. **b**, Example of a 3D segmentation of the DP in a head-on and side view before eversion (left, wL3) and after bilayer formation (right, 2hAPF). Pouch: blue; across-DVB: solid line; along-DVB: dashed line. **c,d**, Representative images for stages of eversion. Wing discs are labelled with Ecadherin-GFP. The pouch region is highlighted, colored by time. **c**, Projection view showing the dorsal side for early pupal stages and dorsal (down), ventral (up), and DVB for wL3. The position of the DVB is indicated with a dashed line. **d**, Across-DVB cross-section. The position of the DVB is indicated in white. Asterisk shows the rupture point of the PM, which gets removed around 4hAPF. Minimum 5 wing discs were analyzed for each time point; hAPF = hours after puparium formation; wL3 = wandering larval stage, 3rd instar; scale bars = 100 μm.

2 Programmable spring network as a model for epithelial morphogenesis

We developed a coarse-grained model of tissue shape changes, leveraging an analogy between tissue remodelling by internal processes and spontaneous strain-driven shape programming of nematic elastomers [24–26]. We use a double layer of interconnected programmable springs representing the apical

surface geometry and the material properties of an epithelial sheet, including a
bending rigidity introduced by the thickness of the double layer (Methods 7.8,
7.13). As an initial configuration, we use a stress-free spherical cap and then
assign new rest lengths to the springs. In a continuum limit, this corresponds
to introducing a spontaneous strain field $\underline{\lambda}(\underline{X})$, which depends on the spatial
coordinates $\underline{X}$ (Methods 7.9). To simplify notation, we write $\underline{\lambda}$ for $\underline{\lambda}(\underline{X})$ here-
after. To generate a final output shape, we quasi-statically relax the spring
network (Methods 7.8, 7.9). As with conventional elastic strain tensors, $\underline{\underline{\lambda}}$ can
be decomposed into isotropic (λ) and anisotropic ($\tilde{\lambda}$) modes (Fig. 2a).

We first wanted to understand how simple choices of spontaneous strain
patterns induce a shape change in our model. A simple gradient of λ, for
example, causes the spherical cap to balloon in the center or generate wrinkles
at the periphery (Fig. 2b.i. and ii). Changing the directions and gradients of
$\tilde{\lambda}$ leads to elongation of the cap, increase in the curvature at the tip or even
flattening of the curvature in the center, eventually leading to a saddle shape
(Fig. 2b.iii - viii.).

We propose that cell behaviors can give rise to a spontaneous strain field,
thereby shape-programming the wing disc pouch and driving 3D shape changes
during eversion. The spontaneous strains from these cell behaviors can be
represented in the programmable spring model by changes in the spring rest
lengths. We calculate a coarse-grained spontaneous strain from the underly-
ing cell behaviors, allowing us to leave the topology of the spring network
unchanged [27].

We build our spontaneous strains from the strains that we infer from
observed cell behaviors during eversion, which we refer to as observed strains
$\underline{\lambda}^*$. For the isotropic component, we focus on cell area changes (λ_A^*), as
cell division and cell death are minimal in the everting wing disc (Fig. 2ci)
[13, 28, 29]. The anisotropic components of spontaneous strain capture contri-
butions stemming from both changes in cell elongation ($\tilde{\lambda}_Q^*$) as well as from
cell rearrangements ($\tilde{\lambda}_R^*$) (Fig. 2cii).

Our model can therefore relate cell behaviors to spontaneous strains in
order to understand resulting tissue deformations. We now investigate these
quantities in the everting wing disc.

Fig. 2 **a**, A thick spherical cap as a model for an epithelial tissue. We define a radial coordinate r and basis vectors $\underline{e}_r, \underline{e}_\phi$, and $\underline{e}_h$. The thickness of the spring network h is constant everywhere and introduces a bending rigidity. The model tissue is an elastic medium implemented as a spring network with an initially stress-free state. We change the rest lengths of the springs by imposing a spontaneous strain field $\underline{\underline{\lambda}}$ and allow subsequent relaxation to take a new 3D output shape. Top and bottom springs at any position in the lattice have their rest lengths updated by the same amount. The spontaneous strain field $\underline{\underline{\lambda}}$ consists of an isotropic component λ and an anisotropic component $\tilde{\lambda}$. These components cause changes in area (A_i to A_f) or area-preserving changes in shape (L_i to L_f), respectively. **b**, Model realizations with simple patterns of spontaneous strains. For each realization, the input pattern of spontaneous strains is displayed above, with the magnitude of strain encoded by color. For anisotropic strain ($\tilde{\lambda}$), the bars indicate the orientation. Below is the output shape. In **b.i-ii**, we vary the isotropic contribution λ and keep $\tilde{\lambda} = 1$, while in **b.iii-viii**, we vary $\tilde{\lambda}$ and keep $\lambda = 1$. We probe the model output from input linear radial gradients in λ or $\tilde{\lambda}$, giving rise to cones with varying degrees of sharpness at the tip (**i ,v ,vi ,vii**) or saddle shapes (**ii, viii**). Using a spatially homogeneous pattern of $\tilde{\lambda}$, we observe an elongated spherical cap when patterned along a fixed direction (**iii**) and a blunt cone when patterned radially (**iv**). **c**, Schematics showing the calculation of apparent spontaneous strains from observed cellular behaviors. **c.i** For a patch of cells going from area of A_i to A_f, we extract λ. **c.ii** A patch of cells undergoing anisotropic deformation due to cell elongation changes or neighbor exchanges causes the length scale in one direction to change from L_i to L_f. From this change, $\tilde{\lambda}$ is extracted, while $\lambda = 1$, as there is no isotropic contribution.

3 Spatial patterns of cell dynamics in the everting wing pouch

From larval stages, we know that cell behaviors in the pouch are organized radially in the region outside of the dorsal ventral boundary (outDVB) and

parallel to the boundary in the region closest to the dorsal ventral boundary (DVB) [30–33]. During eversion, we observe that cell shapes and sizes are patterned similarly. In early stages, cell area follows a radial gradient that disappears by the end of eversion (4hAPF) (Fig. 3a, Methods 7.6). Cell elongation exhibits a global nematic order through 4hAPF before disordering at 6hAPF (Fig. 3b, Methods 7.6).

To compare spatial patterns of cell behaviors over eversion time and across experiments, we define a coordinate system on the evolving 3D geometry. To this end, we use the cellular network topology to define the distance measure on the tissue surface. The topological distance between two cells is defined as the number of cells on the shortest path through the network from one cell to the other (See Extended Data Fig. S2a). We then use topological distance to define a coordinate system in the outDVB and DVB regions (Fig. 3c, Extended Data Fig.S2 and S3a,b, and Methods 7.7). The outDVB region consists of the dorsal and ventral halves, and we identify a single cell that defines the origin in each half (O_D and O_V). In the DVB, we define the origin (O_{DV}) as a line of cells transversing the DVB. The topological distance k to the origin defines a radial topological coordinate in each region, see Fig. 3c,d.

During eversion, the tissue unfolds and more of the pouch comes into view. In order to compare cell behaviors at different time points, we need to identify a region of tissue that remains in the field of view throughout eversion. To this end, we count the number of cells N_{ROI} within the largest visible topological ring at wL3. The corresponding region of interest at later time points is then defined to be centered at the origin and containing the same number of cells. Since there are very few divisions and extrusions during eversion [13, 28], and because cells cannot flow across the DVB [34, 35], we expect that our regions of interest contain largely the same set of cells, and we refer to them as topologically tracked regions (Fig. 3d,g, Extended Data Fig. S5a,b).

Next, we quantify patterns of cell area A and cell elongation as a function of topological coordinate k throughout eversion (See Methods 7.6). We find that our topological coordinate system recapitulates previously reported gradients in cell area and radial cell elongation at earlier larval stages (Extended Data Fig. S3c,d). In outDVB at wL3, we observe a cell area gradient that relaxes gradually until 4hAPF. At the same time, cell elongation develops a gradient, with cells in the periphery elongating tangentially (Fig. 3e,f). Between 4h and 6hAPF, cells dramatically expand their area and tangential cell elongation completely relaxes. We do not observe gradients in cell area or cell elongation in the DVB. Instead, cell area expands globally in the DVB during eversion, while cell elongation along the DVB first increases up to 2hAPF and then decreases at 4hAPF (Fig. 3e,f).

Using topological distance allows us to extract spatial patterns of oriented cell rearrangements from snapshots of eversion. Radially oriented rearrangements lead to a decrease in the number of cells per k, whereas tangentially oriented rearrangements lead to an increase (see Fig. 3g). As a consequence, $k(N_{\mathrm{ROI}})$ changes based on the orientation and magnitude of rearrangements.

We find that $k(N_{\text{ROI}})$ increases with time (Fig. 3g), consistent with radially
oriented cell rearrangements in outDVB and rearrangements oriented along
the boundary in the DVB.
Together, these measured cell behaviors are a superposition of different
radial patterns with the additional complexity of the DVB. Next, using our
programmable spring model (Fig. 2), we asked how in-plane strains caused by
these cell behaviors could drive 3D shape changes during eversion.

Fig. 3 **a,b,d**, Cell measurements highlighted on the surface of representative examples of everting wing discs over time. At wL3, the full pouch is visible, whereas only the dorsal side is shown here for early pupal stages. **a**, Cells are colored by apical cell area. **b**, Bars highlight the orientation of locally averaged cell elongation Q (projected onto 2D), and color indicates the elongation magnitude $|Q|$ averaged over patches of size $350\mu m^2$. **c**, Segmentation of a wL3 pouch; the origins for the topological coordinates k (O_D and O_V in the outDVB region and a center line O_{DV} for the DVB) are highlighted in dark blue. The arrows indicate the direction of spatial coordinates that result from these origins, transversing along the DVB and radially for the outDVB. The inset shows the center region of the same wing pouch, where each cell up to $k = 5$ is colored by k. The origin cells are at $k = 0$ (dark blue). Note that the origin is a single cell for each side of the outDVB and a line of cells for the DVB. Due to the 3D nature of the wing pouch, the topological coordinate system is defined in one view for larval stages and in 4 separate imaging angles for early pupal stages (see schematic, right). **d**, The maximum k depends on the size of the segmented region (upper row). For the topologically tracked region, the maximum k may change due to rearrangements and is denoted $k(N_{\mathrm{ROI}})$ (lower row). **e-f**, Cell area (e) or cell elongation (f) spatially averaged over k (minimum five wing discs per stage). Dorsal and ventral are averaged together into 'outDVB'. Geometric representations (top) show outDVB as half-circles and the DVB as a central rectangular box. **e**, Geometric representations highlight cell area gradients ($A/\langle A \rangle$) within each stage. Lower panels show the cell area (A) as a function of k for all time points. **f**, The component of cell elongation (Q_{rr} for cells in the outDVB and $Q_{\rho\rho}$ for cells in the DVB) is calculated relative to the origin for each cell of the respective region. This makes Q_{rr} the radial component of cell elongation, whereas $Q_{\rho\rho}$ is effectively the cell elongation along the DVB (cartoon insets on the lower panel, see also Methods 7.7). Q_{rr} and $Q_{\rho\rho}$ are calculated as a function of k. In the upper panel, magnitudes are represented by color. **g**, Schematic (top) showing how we estimate cell rearrangements using topology. Each circle represents a cell in the outDVB region of the wing disc, colored by topological distance at the initial time point. If the number of cells per k decreases, the deformation by rearrangements is radial. Plots (bottom) show the number of cells in the wing disc pouch N contained within k. The horizontal line shows N_{ROI} for the wL3 stage; the vertical lines show corresponding ($k(N_{\mathrm{ROI}})$) for each stage. In **e-g**, solid lines indicate the mean, and ribbons show 95% confidence of the mean.

4 Cell rearrangements drive wing pouch eversion

To compare the output of the model to the 3D shape changes happening during eversion, we quantified the curvature and size dynamics of the apical surface of the wing pouch. We limit the analysis to the topologically tracked region and quantify the change in curvature from the wL3 stage along lines in the along-DVB and across-DVB directions (Fig. 1b-d, Extended Data Fig.S5, Methods 7.12). We focus on the stages between wL3 and 4hAPF, during which cell shape patterns have radial symmetry (Fig. 3b,e,f). We observe an overall curvature increase that is more pronounced in the across-DVB direction, peaking at the DVB, while flattening at the dorsal and ventral sides. Furthermore, the overall tissue area increases (Fig. 4a, Extended Data Fig. S5a).

Next, we measure the strain field ($\underline{\underline{\lambda}}^*$, Fig. 4b) resulting from cell behaviors as a function of the distance from the origin, r or ρ (Fig. 4b, Extended Data Fig. S6, Extended Data Fig. S7, Methods 7.11). We quantify the isotropic component resulting from cell area changes (λ_A^*). In the outDVB, we observe an area expansion ($\lambda_A^* > 1$) up to 2hAPF with a radially decreasing profile (Fig. 4b, Extended Data Fig. S7). In the DVB, we observe the buildup of a shallower gradient that is transiently paused from 0hAPF to 2hAPF. The contribution to the anisotropic component of $\underline{\underline{\lambda}}^*$ from changes in cell elongations $\tilde{\lambda}_Q^*$ is small compared to the contribution by cell rearrangements $\tilde{\lambda}_R^*$. While $\tilde{\lambda}_Q^*$ is tangential, following a shallow gradient, $\tilde{\lambda}_R^*$ is radial and increasing with the distance from origin in the outDVB and decreasing in the DVB (Fig. 4b, Extended Data Fig. S7, Methods 7.10).

We next use the programmable spring model to test how the observed in-plane cellular behaviors can cause tissue shape changes. We define the DVB and outDVB regions in the model, matching their relative sizes in the wing pouch (Fig. 4c, Extended Data Fig. S8). For each individual cell behavior and measured time point (wL3, 0hAPF, 2hAPF, and 4hAPF), we use the in-plane strain $\underline{\underline{\lambda}}^*$ that we infer from each observed class of cell behaviors as examples of spontaneous strain $\underline{\underline{\lambda}}$. We use these to program the spring lengths in the model. For each insertion of spontaneous strain (model time points: initial, t1, t2, final, corresponding to the experimental time points), we relax the spring network quasi-statically to a force balanced state (Methods 7.8, 7.9). As the effective bending modulus of the wing disc is experimentally inaccessible, we fit the thickness of the model in an example scenario where all observed cell behaviours are input as spontaneous strains and use the same thickness thereafter (Extended Data Fig. S9a, Methods 7.13).

We first consider cell rearrangements as a possible source of spontaneous strain. When we only input $\underline{\underline{\lambda}}_R = \{1, \tilde{\lambda}_R^*\}$ as spontaneous strain in the model, it alone creates a strong curvature increase, resembling many features of the data but without increasing tissue size (Fig. 4a, d). Note that $\underline{\underline{\lambda}}_R$ also breaks the symmetry between the two directions, across-DVB and along-DVB, at the final stage.

After relaxing the spring network to a force balanced state, stresses due to residual strains remain. The stresses corresponding to these residual strains can drive passive responses in cell behaviors. The residual strains appear as a mismatch of spontaneous strains (input to the model, $\underline{\lambda}$) and strains resulting from changes in spring length during relaxation of the network, $\underline{\underline{F}}$ (Fig. 4e.i, Methods 7.9).

When we calculate the residual strains generated by spontaneous strain from rearrangements, we find that the anisotropic component of the residual strain ($\tilde{\lambda}_R^{\mathrm{res}}$) is tangentially oriented (Fig. 4e.ii, Extended Data Fig. S10). This tangentially oriented strain is similar to the pattern of cell elongation changes ($\tilde{\lambda}_Q^*$)(compare Fig. 4b and 4e.ii), suggesting that these cell elongation changes are a passive response to spontaneous strain by rearrangements. To test this idea, we next consider cell elongation as possible source of spontaneous strain. When we only input $\underline{\underline{\lambda}}_Q = \{1, \tilde{\lambda}_Q^*\}$ as spontaneous strain in the model, we observe that the spring network shape flattens at the center rather than curve, and cell elongations themselves do not lead to any further residual strains (Fig. 4f,g, Extended Data Fig. S10). This result is consistent with cell elongation changes being a passive response to cell rearrangements and not driving tissue shape change during eversion.

Cell rearrangements as spontaneous strains also lead to residual isotropic compression (λ_R^{res}, Fig. 4e.ii, Extended Data Fig. S10). This residual could be compensated by spontaneous area change, which is also required by the observation that overall tissue size increases during eversion.

When we only input the isotropic strain, $\underline{\underline{\lambda}}_A = \{\lambda_A^*, 1\}$, from observed cell area changes as spontaneous strain in the model, overall size increases with minimal curvature change (Fig. 4h). This result indicates that although cell area changes are an active behavior and lead to overall size increase, they do not significantly contribute to changes in tissue shape. However, there does appear to be a transient effect of cell area changes on tissue curvature at time point t2. This transient effect in the scenario of spontaneous area strain only arises from the experimentally observed pause in cell area expansion in the DVB at 2hAPF as compared to the outDVB (Extended Data Fig. S7). It later disappears in the model when the cell area in the DVB expands to match the outDVB at 4hAPF (Fig. 4e). Measuring $\underline{\underline{\lambda}}_A^{\mathrm{res}}$, we find that cell area changes themselves create a small residual in the DVB (Fig. 4i, Extended Data Fig. S10). The anisotropic part of this residual could also contribute to the observed passive cell elongations.

Using these examples, we next infer the spontaneous strain patterns that drive tissue shape changes and govern cellular behaviors. We have found that both cell rearrangements and cell area changes are active and contribute to spontaneous strain. We therefore conclude that cell elongation is a passive elastic response and does not contribute to spontaneous strain. The total spontaneous strain, therefore, is composed of the anisotropic part of the observed strain due to rearrangements ($\tilde{\lambda}_R^*$, Fig. 4b) and the isotropic part of the

observed cell area changes (λ_A^*) compensated by the isotropic part of the residual strain due to cell rearrangements (λ_R^{res}, Fig. 4e.ii). Fig. 4j shows the shape changes generated by this total spontaneous strain, which accounts for the curvature and size changes observed from wL3 to 4hAPF in the everting wing disc (see also Extended Data Fig. S9b). The good qualitative agreement between model output and wing pouch curvature changes indicates that we have identified the most relevant active cellular events responsible for the pouch shape change during eversion. Furthermore, the patterns of residual strains generated by the model provide a prediction of the mechanical stresses present in the wing disc pouch after eversion. In particular, after eversion (at 4hAPF), cells experience elongation due to shear stress as well as area constriction due to compressive stresses (Fig. 4k, Extended Data Fig. S7).

Fig. 4 **a**, Overlay of a wL3 (white) and a 4hAPF (cyan) wing pouch (left) and plots of the average change in tissue curvature in the topologically tracked region in across-DVB (middle) and along-DVB (right) directions. **b**, Observed strain from cellular behaviors between time points wL3 to 4hAPF as a function of normalized distance from origin r and ρ. Observed strains arise from (left to right): rearrangements ($\tilde{\lambda}_R^*$), cell elongation changes ($\tilde{\lambda}_Q^*$), and cell area changes (λ_A^*). Half circles indicate the outDVB region; the rectangular box indicates the DVB. The color represents the magnitude of different strains; the bars visualize the direction of observed strain for $\tilde{\lambda}_R^*$ and $\tilde{\lambda}_Q^*$. **c**, The model coordinates are designed to match the geometry of spatial patterns in the wing disc pouch (See also Fig. 3c). **d,f,h** Observed in-plane strain from rearrangements (**d**, $\tilde{\lambda}_R^*$), cell elongation changes (**f**, $\tilde{\lambda}_Q^*$), and cell area changes (**h**, λ_A^*) are inserted in the model as spontaneous strains by a change in rest length of the springs (δ^o/δ^*). To compare the initial and final stages (corresponding to wL3 to 4hAPF), the model cross-section shows the shape in the across-DVB direction. The change in curvature of the model outcomes are plotted for all time points (right) in the across-DVB and along-DVB directions. The initial shape is a spherical cap with a radius resembling the wL3 stage. t1,t2, and final stages are the model results from the change in strains by 0,2, and 4hAPF. $\underline{\lambda}$ contains observed strains from the individual measured cell behaviors, while the other components (λ or $\tilde{\lambda}$) are set to 1. **e.i,g.i,i.i**, Input spontaneous strain for $\tilde{\lambda}_R^*$ (**e**), $\tilde{\lambda}_Q^*$ (**g**), and λ_A^* (**i**) at the final eversion time point (λ, $\tilde{\lambda}$) and the resulting strain that is achieved after relaxation of the model, which can be isotropic (F) and anisotropic ($\tilde{F}$). **e.ii,g.ii,i.ii**, Residual strain that remains at the final time point. The colors shows the magnitude of strain using the same range as indicated in Fig. 4b. Plots are split vertically to show the isotropic component (λ) on the left and the anisotropic component ($\tilde{\lambda}$) on the right. **j, k**, Model output and residual strains for the inferred spontaneous strains, following the same procedure as in **d-i**.

5 MyoVI knockdown alters active cell behaviors resulting in a tissue shape phenotype

So far, our analysis has revealed that active in-plane cell behaviors can drive tissue shape changes in the wing blade during eversion. We now use a genetic perturbation to confirm the role of active cell behaviors. Previous work in the wing disc pouch of earlier larval stages showed that cell rearrangements drive cell shape patterns [33]. This work suggested that patterns of active cell rearrangements self-organize via mechanosensitive feedback mediated by MyoVI. We therefore next investigate whether MyoVI knockdown in the wing pouch (Extended Data Fig. S11) alters cell rearrangements during eversion and leads to a tissue shape phenotype.

We observe that the MyoVI[RNAi] wing disc fails to form a flat bi-layer after eversion, even though its initial shape is similar to the *wild type* (wt) (Fig. 5a.i). This phenotype is best captured in the behavior of curvature in the across-DVB direction (Fig. 5a.ii). Here, the curvature decreases in the center, in contrast to wt, where it increases. In the along-DVB direction, the curvature remains unchanged over time in the MyoVI[RNAi] knockdown (Fig. 5a,b, Extended Data Fig. S11c-d). Other features of eversion, such as the opening of the folds and the removal of the peripodial membrane are unaffected by the MyoVI[RNAi] knockdown (Fig. 5a, see 4hAPF), indicating that the cause for the altered shape is pouch-intrinsic. This result further supports the idea that tissue shape changes in the wing pouch during eversion are independent of other morphogenetic events.

Next, we quantified cell behaviors in MyoVI[RNAi]. While initially the gradients in cell areas and elongation are similar to wt (Extended Data Fig. S11e,f), the inferred strains from individual types of cell behaviors $\underline{\lambda}^*$ differ (Fig. 5c, Extended Data Fig. S12). From work in earlier larval stages, we expect oriented rearrangements to be reduced [33]. Indeed, we find that MyoVI[RNAi] reduces the amount of radial cell rearrangements in the outDVB during eversion. However, in the DVB, rearrangements are of opposite orientation as compared to wt eversion. Notably, we also see a complete lack of cell area expansion in the DVB. The pattern of cell elongations in the outDVB is similar to wt, but in the DVB it is of perpendicular orientation.

Using the programmable spring model, we test how the reduction of spontaneous strain due to cell rearrangements affects tissue shape changes. When we input $\underline{\underline{\lambda}}_R = \{1, \tilde{\lambda}^*_R\}$ from cell rearrangements measured in MyoVI[RNAi] as spontaneous strain in the model, we see only a slight increase in curvature in the final time point in both along- and across-DVB directions (Fig. 5d). Thus, we conclude that the reduction of cell rearrangements in MyoVI[RNAi] as compared to wt contributes to the abnormal tissue shape changes happening during eversion in MyoVI[RNAi]. We find that the anisotropic component of the residual strain ($\tilde{\lambda}^{\text{res}}_R$) is small and tangentially oriented in the outDVB and radially in the DVB, similar to the cell elongation pattern (Fig. 5c,e, Extended Data Fig. S13). If we input measured cell elongation changes as spontaneous

strain in the model ($\underline{\underline{\lambda}}_Q = \{1, \lambda_Q^*\}$), we do not recapitulate the observed tis-   351
sue shape changes (Extended Data Fig. S14). This result suggests that the   352
cell elongation changes in MyoVI$^{\text{RNAi}}$ are a passive response to spontaneous   353
strain by cell rearrangements, as in wt.   354

     While the change in spontaneous strain due to rearrangements captures a   355
significant portion of the difference between the wt and MyoVI$^{\text{RNAi}}$, it fails to   356
recapitulate the finer progression of shape from wL3 to 4hAPF in MyoVI$^{\text{RNAi}}$.   357
In particular, the curvature at the final time point of the model calculation   358
is not flattened in the center of the across-DVB direction, and the curvature   359
increases slightly in both directions (Fig. 5b,d).   360

     Thus, we proceed to input the observed cell area changes as spontaneous   361
strains in the model ($\underline{\underline{\lambda}}_A = \{\lambda_A^*, 1\}$). We find that they produce shape changes   362
over time similar to those observed during MyoVI$^{\text{RNAi}}$ eversion, recapitulating   363
both the decrease in curvature in the center of the across-DVB direction and   364
the lack of curvature change in the along-DVB direction (Fig. 5f). We conclude,   365
therefore, that the subtle flattening in the pouch center in MyoVI$^{\text{RNAi}}$ during   366
eversion can be explained by the combination of cell area expansion in the   367
outDVB with no area expansion in the DVB. This result highlights that, while   368
cell area changes do not lead to a curvature change in wt, the difference in   369
area expansion between the tissue regions results in the MyoVI$^{\text{RNAi}}$ shape.   370
In addition, although we did not observe cell area expansion in the DVB,   371
the area expansion in the outDVB creates residual strains in both regions   372
(Fig. 5g, Extended Data Fig. S13). These residual strains have an anisotropic   373
component that, together with the residual strains from cell rearrangements,   374
account for the measured cell elongation patterns in MyoVI$^{\text{RNAi}}$ (compare Fig.   375
5c,e.ii,g.ii)).   376

Fig. 5 a, MyoVI^{RNAi} phenotype during eversion (scale bars = 100 μm). Representative across-DVB cross-sections (**a.i**) and comparison of apical shape between MyoVI^{RNAi} and control (**a.ii**). **b**, Overlay of a wL3 (white) and a 4hAPF (cyan) MyoVI^{RNAi} wing pouch (left) and plots of the average change in tissue curvature in the topologically tracked region for across-DVB (middle) and along-DVB (right) directions. **c**, Observed strain from cellular behaviors in MyoVI^{RNAi} wing discs between time points wL3 to 4hAPF. Plots are split vertically with the observed strains for *wild type* (WT) for comparison on the left and MyoVI^{RNAi} on the right. Measured strains come from $\tilde{\lambda}_R^*$, $\tilde{\lambda}_Q^*$, and λ_A^*. Quarter circles indicate the outDVB region, and the rectangular box indicates the DVB. The color represents the magnitude of different strains; the bars indicate the direction of observed strain for $\tilde{\lambda}*_R$ and $\tilde{\lambda}*_Q$. **d,f**, Observed in-plane behaviors are inserted in the model as spontaneous strains by a change in rest lengths of the springs (δ^o/δ^*). The initial stage is a spherical cap with the radius taken to resemble the shape at the *wild type* wL3 stage. t1,t2, and final stages are the model results after a change in spring rest length according to observed strains from 0,2, and 4hAPF for MyoVI^{RNAi}. $\underline{\underline{\lambda}}$ contains observed strains from rearrangements (**d**) or area changes (**f**), while the other components (λ_R or $\tilde{\lambda}_A$) are set to 1. **e.i,g.i**, Input spontaneous strain ((λ and $\tilde{\lambda}$) at the final eversion time point and comparison with the resulting strain that is achieved after relaxation of the model (F and $\tilde{F}$). **e.ii,g.ii**, Residual strain that remains from the difference between input and resulting spontaneous strain at the final time point, plotted in the same way as Fig. 4e,g,i,k.ii.

6 Discussion

In this work, we show that 3D epithelial tissue morphogenesis in the *Drosophila* wing disc is based on in-plane spontaneous strains generated by active cellular behaviors. We develop a metric-free, topological method to quantify patterns of cell dynamics on arbitrarily shaped tissue surfaces, as well as a theoretical

approach to tissue morphogenesis inspired by shape-programmable materials. 382
These advancements together reveal the mechanics of tissue shape changes 383
during wing disc eversion, showing that active rearrangements and active area 384
expansion govern the 3D tissue shape and size changes. 385

We hypothesize that the organization of active behaviors during wing ever- 386
sion arises from patterning during larval growth. First, the pre-patterned radial 387
cell area gradient resolves during eversion, giving rise to a gradient of spon- 388
taneous strain in the outDVB. Second, the orientation of cell rearrangements 389
follows that of earlier stages, indicating that the mechanosensitive feedback 390
that was revealed in previous work is still active during eversion. Overall, this 391
suggests a developmental mechanism through which mechanical cues at early 392
stages organize cell behavior patterns that later resolve, resulting in a shape 393
change. Such behavior would resemble biochemical pre-patterning, in which 394
cell fates are often defined long before differentiation. 395

Active, patterned rearrangements can robustly give rise to a specific target 396
shape if the tissue is solid on the time scale of morphogenesis. Our work there- 397
fore reveals that the everting wing disc behaves as an elastic solid undergoing 398
plastic deformation and demonstrates that the mere presence of rearrange- 399
ments should not be taken as a sign of a fluid tissue with a vanishing elastic 400
modulus. Many animal tissues with dynamic rearrangements could thus be in 401
the solid regime and therefore be pre-patterned towards a target shape. Our 402
work, inspired by shape programmability of complex materials, reveals prin- 403
ciples of shape generation that could be quite general. We therefore propose 404
that many other morphogenetic events could and should be considered – and 405
better understood – through the lens of shape-programmability. 406

7 Methods   407

7.1 Experimental model   408

All experiments were performed with publicly available *Drosophila* 409
melanogaster lines. Flies were maintained at 25°C under 12hr light/dark cycle 410
and fed with standard food containing cornmeal, yeast extract, soy flour, malt, 411
agar, methyl 4-hydroxybenzoate, sugar beet syrup, and propionic acid. Adult 412
flies were transferred to fresh food 2-3 times per week. Only males were stud- 413
ied for consistency and due to their smaller size. As *wild type*, we used the F1 414
offspring of a cross between *w-;ecad::GFP* and *w;nub-Gal4,ecad::GFP;;*. 415

7.2 *Drosophila melanogaster* lines   416

genotype	construct	origin
w-; ecad::GFP;;	Ecadherin::GFP	Bloomington #60584
w; nub-Gal4, ecad::GFP;;	nub-Gal4	Bloomington #86108
w-; ecad::GFP, myoVI^{RNAi};;	MyoVI^{RNAi}	VDRC #37534

7.3 Image acquisition and processing

Sample preparation: Wing discs of larval stages were dissected in culture medium as previously described [36], without surface sterilization or antibiotics. Prepupal stages from 0 to 6 hAPF required a slightly different dissection strategy. Prepupae were marked by the time of white Pupa formation and collected with a wet brush after the required time interval. Next, pupae were placed on a wet tissue, cleaned with a wet brush to remove residual food, and transferred into glass staining blocks (see Ref. [36]) filled with dissection medium. To dissect the wing disc, a small cut was performed with fine surgical scissors (2.5 mm, FST 15000-08, Fine Science Tools GmbH) at the posterior end, which creates a small hole to release pressure. This allowed for the next cut to be performed at half the anterior-posterior length, separating the anterior and posterior halves. Next, the anterior part of the puparium was first opened at the anterior end by administering a cut just posterior to the spiracles and then a second cut was performed on the ventral side along the PD-axis. The pupal case was then held open with one forceps, and a second forceps in the other hand was used to remove the wing disc . To dissect wing discs from 0hAPF, the pupa is still soft enough to be turned inside-out after the cut that separates anterior and posterior halves, similar to larval stages.

Imaging: Imaging was performed with a Zeiss Lightsheet 1 system. Wing discs were mounted in capillaries (Zeiss, Capillary size 1, inner diameter ca. 0.68 mm) with 1 % low melting-point agarose (LMA, Serva, CAS 9012-36-6). LMA was prepared by mixing 1:1 of Grace's insect medium and a 2% LMA stock solution in water. Wing discs were transferred into mounting medium in U-glass dishes, aspirated into the capillary at room temperature, and imaged immediately after the LMA solidified. The imaging chamber was filled with Grace's insect medium (measured refractive index = 1.3424). For pupal stages, four imaging angles (dorsal, ventral, and 2 lateral) with 90° rotation were acquired; for larval stages, three imaging angles (dorsal and ventral in one, and 2 lateral) were acquired. Data from dual illumination was fused on the microscope using a mean fusion.

Multiview reconstruction: Multiview reconstruction was based on the BigStitcher plugin in Fiji [23, 37]. Images were acquired without fluorescent beads, and multiview reconstruction was done using a semi-automated approach. Individual views are manually pre-aligned. Thereafter, precise multiview alignment was computed based on bright spots in the data with an affine transformation model using the iterative Closest Point (ICP) algorithm. Next, images were oriented to show the apical side in XY and lateral in ZY. Lastly, images were deconvolved using point spread functions extracted from the bright spots and saved as tif files with a manually specified bounding box.

Surface extraction of 3D images for visualization: Surfaces shown in Fig. 1 and Supplemental movies were extracted from 2hAPF and wL3 images. To do so, we first trained a pixel classifier on the strong apical signal of Ecadherin-GFP of a different image of the same stage with napari-accelerated-pixel-and-object-classification [38, 39]. Feature sizes of 1-5 pixels were used to

predict the foreground on the target image. Next, we used the pyclesperanto 462
library [40] to select the largest labels and close gaps in the segmentation with 463
the closing sphere algorithm. For additional gap-filling in the 2hAPF time 464
point, we used vedo [41] to generate a pointcloud and extract the pointcloud 465
density. When necessary, we applied some manual pruning of the segmenta- 466
tion in napari. We repeated this processing on the weak Ecadherin-GFP signal 467
from the lateral membrane and subtracted the apical segmentation from the 468
output. As a result, we achieve a full tissue segmentation that stops just below 469
the apical junction layer. We then extracted the surface by the napari-process- 470
points-and-surfaces [42] library and applied smoothing and filling holes. The 471
visualizations were generated using Paraview [43]. Regions and directions of 472
the cross-sections were annotated in Illustrator. Supplemental movies were 473
created using paraview and Fiji [37]. 474

7.4 Segmentation of the apical junction network   475

To analyze cell shapes, we use four angles separated by 90° for the segmenta- 476
tion of early pupal stages, and a single angle for larval wing discs (Extended 477
Data Fig. S4). Z-stacks from each imaging angle were denoised if necessary, 478
by using the N2V algorithm [44], and the signal to background ratio was 479
further improved by background subtraction tools in Fiji [37]. We made 2D 480
projections of the Ecadherin-GFP signal in the Disc Proper layer as previously 481
described [45]. Importantly, this algorithm also outputs a height-map image, 482
which encodes the 3D information in the intensity of each pixel. The cells in 483
the wing pouch were segmented using Tissue Analyzer and manually corrected 484
[46]. We chose a bond length cutoff of 2 pixels ($\sim 0.46\mu m$). The ventral side 485
for 0 hAPF was excluded from the analysis, as at this stage, the ventral region 486
is never fully in view from any imaging angle. The number of wing discs per 487
time point and the images for each region are indicated in Table 1. Images were 488
rotated to orient distal down. Height-map images were rotated accordingly 489
using imagemagickTM software (ImageMagick Development Team, 2021). We 490
use Fiji macros included with TissueMiner [47] to manually specify regions of 491
interest (ROIs). The DV boundary is identified based on Ecadherin-GFP sig- 492
nal intensity [48] and the dorsal vs. ventral pouch by their positions relative 493
to global tissue morphology. For larval stages, the DV boundary, dorsal, and 494
ventral regions are identified in one image. For images showing lateral views 495
of pupal stages, the DV boundary is identified, whereas for images showing 496
the outDVB region, the dorsal or ventral region and the cells next to the DV 497
boundary are labelled. The cells next to the DV boundary are required as a 498
landmark for topological analysis but are otherwise not analysed separately. 499
We then run the TissueMiner workflow to create a relational database. 500

Table 1 wing disc data

Genotype	Stage	Number of wing discs	ROI	Replicates
E-cad::GFP, nub-Gal4	96hAEL	7	DVB	7
			dorsal	7
			ventral	7
	120hAEL	5	DVB	5
			dorsal	5
			ventral	5
	wL3	5	DVB	5
			dorsal	5
			ventral	5
	0hAPF	7	DVB	7
			dorsal	7
	2hAPF	5	DVB	5
			dorsal	5
			ventral	5
	4hAPF	7	DVB	7
			dorsal	7
			ventral	7
	6hAPF	6	DVB	6
			dorsal	4
			ventral	4
E-cad::GFP, nub-Gal4, MyoVIRNAi	wL3	5	DVB	5
			dorsal	5
			ventral	5
	0hAPF	6	DVB	6
			dorsal	6
	2hAPF	6	DVB	6
			dorsal	5
			ventral	6
	4hAPF	5	DVB	5
			dorsal	5
			ventral	5
	6hAPF	7	DVB	7
			dorsal	6
			ventral	6

7.5 3D cellular network

We represent the configuration of the cellular network by positions of the cell vertices, where three or more cell bonds meet, and their topological relations as in TissueMiner [47]. We extend TissueMiner to the third dimension using the information extracted from height-maps, as described in Methods 7.4.

7.6 Measurement of cell area and cell elongation tensor

Each cell α in the 3D network contains N^α vertices $\underline{v}_i^\alpha$, defining the network geometry. For every cell, we define a centroid $\underline{R}^\alpha$, an area A^α, and a unit normal vector $\underline{\hat{N}}^\alpha$ as

$$\underline{R}^\alpha = \frac{1}{N^\alpha} \sum_{i=1}^{N^\alpha} \underline{v}_i, \quad A^\alpha = \frac{1}{2} \sum_{i=1}^{N^\alpha} \|\underline{n}_i^\alpha\|, \quad \underline{\hat{N}}^\alpha = \frac{1}{\|\sum_{i=1}^{N^\alpha} \underline{n}_i^\alpha\|} \sum_{i=1}^{N^\alpha} \underline{n}_i^\alpha \quad (1)$$

where $\underline{n}_i^\alpha = (\underline{v}_{i+1}^\alpha - \underline{v}_i^\alpha) \times (\underline{R}^\alpha - \underline{v}_i^\alpha)$ is the normal vector on the triangle formed 510
by one edge of the cell and the vector pointing from the cell vertex to the cell 511
centroid. It has a norm equal to twice the area of the triangle. 512

We then create a subcelluar triangulation by connecting the two consec- 513
utive vertices in every cell with its centroid $\{\underline{v}_i, \underline{v}_{i+1}, \underline{R}^\alpha\}$. This creates a 514
complete triangulation that depends both on the vertex positions and the 515
centroids of the cellular network. 516

Each triangle is defined by its three vertices $\{\underline{R}_0, \underline{R}_1, \underline{R}_2\}$, which define two 517
triangle vectors $\underline{E}_1$, $\underline{E}_2$ and its unit normal vector $\hat{\underline{N}}$ 518

$$E_1 = \underline{R}_1 - \underline{R}_0, \quad E_2 = \underline{R}_2 - \underline{R}_0, \quad \hat{\underline{N}} = \frac{\underline{E}_1 \times \underline{E}_2}{\|\underline{E}_1 \times \underline{E}_2\|}. \tag{2}$$

These vectors also define the local basis on the triangle. Using the triangle 519
vectors, we can define the area of the triangle and the rotation angles θ_x and 520
θ_y that rotate a vector parallel to the z-axis of the lab reference frame to the 521
vector normal to the plane of the triangle 522

$$A = \frac{1}{2}\|\underline{E}_1 \times \underline{E}_2\|, \quad \theta_x = -\arctan(N_y, N_z), \quad \theta_y = \arctan(N_x, 1 - N_x^2). \tag{3}$$

Here, $\arctan(x, y)$ is the element wised arc tangent of x/y, and N_i is a 523
component of the unit vector normal to the triangle plane. 524

For each triangle, we define the triangle shape tensor $\underline{\underline{S}}^{3d}$ as a tensor that 525
maps a reference equilateral triangle with area A_0 lying in the xy-plane, defined 526
by the vectors vectors $\underline{C}_i$ to the current triangle 527

$$\underline{E}_i = \underline{\underline{S}}^{3d}\underline{C}_i. \tag{4}$$

The vectors of the reference equilateral triangle are 528

$$C_1 = \begin{pmatrix} l \\ 0 \\ 0 \end{pmatrix}, \quad \underline{C}_2 = \begin{pmatrix} l/2 \\ \sqrt{3}/2l \\ 0 \end{pmatrix}, \quad \underline{C}_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \tag{5}$$

where the side length $l = \sqrt{4A_0/\sqrt{3}}$ with $A_0 = 1$. 529

The triangle shape tensor $\underline{\underline{S}}^{3d}$ can be written in terms of a planar state 530
tensor $\underline{\underline{S}}^{planar}$ in the reference frame of the triangle as 531

$$\underline{\underline{S}}^{3d} = \underline{\underline{R}}_x(\theta_x)\underline{\underline{R}}_y(\theta_y)\underline{\underline{S}}^{planar}, \tag{6}$$

where $\underline{\underline{R}}_x(\theta_x)$ and $\underline{\underline{R}}_y(\theta_y)$ are rotations around the x and y axis, respectively. 532
The angles θ_x and θ_y are defined in Eq. 3. The planar triangle state tensor, 533
represented by a 3x3 matrix with the z components set to 0, can be decomposed 534

as

$$\underline{\underline{S}}^{\text{planar}} = \sqrt{\frac{A}{A_0}} \underline{\underline{R}}_{\text{z}}(\phi) \exp(\|\widetilde{\underline{\underline{Q}}}\|\underline{\underline{\gamma}}) \underline{\underline{R}}_{\text{z}}(-\phi) \underline{\underline{R}}_{\text{z}}(\theta_z) \tag{7}$$

as in TissueMiner. Here, $\underline{\underline{\gamma}}$ is a diagonal matrix with diagonal elements $\{1, -1, 0\}$, and $\underline{\underline{R}}_{\text{z}}$ is the rotation matrix around the z-axis. A is the area of the triangle, $\|\widetilde{\underline{\underline{Q}}}\|$ the magnitude of the elongation tensor, ϕ the direction of elongation in the xy-plane, and θ_z is the rotation angle around the z-axis relative to the reference unilateral triangle. The 3D elongation tensor $\widetilde{\underline{\underline{Q}}}$ in the lab reference frame and the elongation tensor in the xy-plane of the triangle $\widetilde{\underline{\underline{Q}}}^{\text{planar}}$ are related by

$$\widetilde{\underline{\underline{Q}}} = \underline{\underline{R}}_{\text{x}}(\theta_x) \underline{\underline{R}}_{\text{y}}(\theta_y) \widetilde{\underline{\underline{Q}}}^{\text{planar}} \underline{\underline{R}}_{\text{x}}(-\theta_y) \underline{\underline{R}}_{\text{y}}(-\theta_x) \tag{8}$$

$$\widetilde{\underline{\underline{Q}}}^{\text{planar}} = \underline{\underline{R}}_{\text{z}}(\phi) \exp(\|\widetilde{\underline{\underline{Q}}}\|\underline{\underline{\gamma}}) \underline{\underline{R}}_{\text{x}}(-\phi). \tag{9}$$

The magnitude of elongation is calculated as [49]

$$\|\widetilde{\underline{\underline{Q}}}\| = \operatorname{arcsinh}\left(\frac{\|\underline{\underline{S}}^{ts}\|}{\sqrt{\|\underline{\underline{S}}^{ta}\|^2 - \|\underline{\underline{S}}^{ts}\|^2}}\right). \tag{10}$$

where $\|\underline{\underline{S}}^{ta}\|$ and $\|\underline{\underline{S}}^{ts}\|$ are the norms of the trace-antisymmetric and traceless-symmetric part of the planar triangle state tensor $\underline{\underline{S}}^{\text{planar}}$, respectively. The angle of the elongation tensor is given by

$$\phi = \frac{1}{2} \arctan 2\left(B_{xy}, B_{xx}\right), \tag{11}$$

where B_{ij} are the components of the nematic part of the triangle state tensor $\underline{\underline{S}}$ and $\arctan 2(x1, x2)$ the inverse tangent of $x1/x2$, where the sign of $x1$ and $x2$ is taken into account. In this way, one can select the branch the multivalued inverse tangent function that corresponds to the angle defined by the point $(x1, x2)$ in a plane.

We now define the cell elongation tensor as the area-weighted average of the corresponding triangle elongations

$$\underline{\underline{Q}}^\alpha = \frac{1}{A^\alpha} \sum_{t \in \text{cell}} a^t \underline{\underline{Q}}^t, \tag{12}$$

where A^α is the area of the cell, a^t the area of a triangle that overlaps with the cell, and $\underline{\underline{Q}}^t$ is the elongation tensor of that triangle.

To calculate the radial component of the cell elongation tensor relative to the origin in cell α, we first define the radial direction. To this end, we use a 3D vector $\underline{r}$ connecting the origin to the cell centroid and we project its direction $\hat{\underline{r}} = \underline{r}/\|\underline{r}\|$ into the tangent plane of the cell, which defines the in-plane radial

direction $\hat{\underline{r}}_{\text{tangent}}$. The tangent plane of the cell is defined by its normal vector   561
$\underline{\hat{N}}$ defined in Eq. 1. We calculate the radial components of the cell elongation   562
tensor as   563

$$Q_{rr} \equiv \hat{\underline{r}}_{\text{tangent}} \cdot \underline{\underline{Q}} \cdot \hat{\underline{r}}_{\text{tangent}} \tag{13}$$

relative to the origin.   564

In the DVB, multiple cells form the origin. To calculate $Q_{\rho\rho}$, the vector ρ 565
connects the cell centroid to the averaged position of the topologically nearest   566
cells of $k = 0$. We project its direction $\hat{\rho} = \rho/\|\rho\|$ into the tangent plane of   567
the cell α, which defines the in-plane direction $\hat{\underline{\rho}}_{\text{tangent}}$ from DVB origin. We   568
calculate the components of the cell elongation tensor as   569

$$Q_{\rho\rho} \equiv \hat{\underline{\rho}}_{\text{tangent}} \cdot \underline{\underline{Q}} \cdot \hat{\underline{\rho}}_{\text{tangent}}. \tag{14}$$

7.7 Topological distance coordinate system   570

To calculate topological distances between any two cells, we determine the   571
topological network using the python-igraph library [50].   572

In each of the tissue regions, we define separate origins: 573

- outDVB region: To define the origin of the outDVB regions, we first deter-   574
  mine the pouch margin cells as cells that live on the outermost row of the   575
  segmentation mask and do not overlap with the DVB ROI. Then, for each   576
  cell in the region, we calculate the shortest topological distance to the margin   577
  cells. This identifies the set of maximally distant cells that have the maximal   578
  shortest topological distance to the margin. The origin is then defined as   579
  the cell that is neighboring the DVB and is at the shortest metric distance   580
  to the averaged position of maximally distant cells. At larval stages, both   581
  dorsal and ventral sides of the outDVB region are visible, and an origin cell   582
  is defined on both sides.   583
- DVB region: We define the origin to consist of a line of cells transversing   584
  the DVB. At larval stages, the origin cells are defined as those cells within   585
  $\sqrt{A_{\text{cell}}/\pi} * 1.2$ distance to a straight line connecting the dorsal and ventral   586
  center cells. For pupal stages, the origin cells for the DVB are defined as the   587
  first row of cells next to the margin of the segmentation mask on the distal   588
  side.   589

The so-identified origin cells serve as the origin for the topological distance (k) 590
for each cell in the tissue. In this way, k follows the radial direction along the   591
surface for the outDVB and the path along the the DVB for the DVB.   592
3D visualization of cell properties: We visualize cellular properties and   593
cell elongation tensors on the 3D segmentation mask using paraview [43].   594

To plot a rank 2 tensor, like the cell elongation tensor, we take the largest   595
eigenvalue of $\underline{\underline{Q}}^{\alpha}$ as the norm of elongation and the corresponding eigenvector   596
as the direction of elongation that we can plot to the surface. Note that for   597

cells / patches that are reasonably flat, the eigenvector with the eigenvalue
closest to zero is (almost) parallel to the normal vector on the patch.
Spatial analysis of cell properties: We acquired data for 5 to 7 wing discs
of each developmental stage. Images that were not of segment-able quality were
excluded from the analysis. We average cell properties by k between dorsal
and ventral for the outDVB and between images from both sides of the DVB.
We use a cell area-weighted average for elongation.

605 7.8 Mechanics of the programmable spring lattice

We use a programmable spring lattice in the shape of a spherical cap to model
the wing disc pouch, which is an epithelial monolayer.
Approximating the wing disc pouch as a spherical cap: We calculate
the average radius of curvature of the apical side of the wing disc pouch at
wL3 stage in the topologically tracked region as $R = 77.66\mu m$. The angular
size of the spherical cap, denoted by θ_M, is given by

$$\theta_M = \frac{1}{2}\left(\frac{w_{DV} + 2w_{ODV}}{R}\right), \tag{15}$$

where w_{DV} is the width of the DVB and w_{ODV} is the average in-surface dis-
tance from the DVB to the periphery of the outDVB region (Extended Data
Fig. S8a). We calculate $w_{DV} = 15\mu m$ and $w_{ODV} = 59.77\mu m$. Using these cal-
culated dimensions, we determine $\theta_M = 49.63°$.
Generating the lattice : We first generate a triangular lattice in the shape
of a hollow sphere, keeping the radius of curvature R calculated above. This
lattice was obtained using the function *meshzoo.icosa_sphere* available in the
Python package Meshzoo (www.github.com/meshpro/meshzoo). In this func-
tion, we set the argument $refine\_factor = 30$, which leads to edges of length
$3.11 \pm 0.18\mu m$. This edge length was found to be small enough to prevent
computational errors in the simulations of this study. We then cropped the
spherical lattice to obtain a spherical cap of angular size θ_M (calculated above,
Extended Data Fig. S8b). Next, we place a second layer at the bottom of this
lattice at a separation of h. This new layer is identical to the original lattice
in terms of the topology of the lattice network but is rescaled to have a radius
of curvature of $R - h$. We connect the two layers with programmable springs
using the topology shown in the inset of Extended Data Fig. S8c. The lattice
obtained this way represents an elastic surface of thickness h, which can be
changed to tune the bending rigidity of the model. Vertices typically have 13
neighbors (6 on their own layer and 7 on the other layer). However, six to eight
vertices out of about 3220 vertices in the whole network form point defects.
These vertices have 11 neighbors.
In order to remove any possible effects coming from the lattice structure
(angle of edges or degree of connectivity), we perform simulations for each
condition by taking spherical caps from 50 different regions of the sphere and
averaging the result. We see only very small variability in the final shape,

quantified by the standard deviation of the curvature change profiles in our
model results. Thus, we conclude that the lattice structure does not affect our
results.

Elastic energy of model: The edges of the lattice act as overdamped elastic
springs with rest lengths equal to their initial lengths. Hence the model is
stress-free at $T = 0$

$$\delta_I^a = \|\Delta \underline{X}^a\| \tag{16}$$

where a denotes a single spring; $\Delta \underline{X}^a$ denotes the spring vector given by $\underline{X}^\beta - \underline{X}^\alpha$, where α and β are the vertices at the two ends of spring a and X^α denotes
the position vector of vertex α. During a consequent time step T, the rest
length of spring a (δ_T^a) can differ from its current length δ. The elastic energy
of this state for the whole lattice is given by

$$W = \frac{1}{2} \sum_a k (\delta^a - \delta_T^a)^2, \tag{17}$$

where the sum is over all springs of the network and k represents the spring
constant. At each computational time step T, the model tries to find a preferred
configuration by minimizing W, hence T acts as a "quasi-static time step". To
minimize the energy of the model at a given T, we use overdamped dynamics
with smaller time steps τ, which restart for each new quasi-static time step T.

$$\begin{aligned}
\frac{d\underline{\mathbf{x}}^\alpha}{d\tau} &= -\frac{1}{\gamma} \frac{\partial W}{\partial \underline{\mathbf{x}}^\alpha} \\
&= -\frac{k}{\gamma} \sum_a (\delta^a - \delta_T^a) \, \underline{\hat{\delta}}^a.
\end{aligned} \tag{18}$$

Here, γ represents the friction coefficient. $\mathbf{x}^\alpha$ corresponds to the current posi-
tion of the vertex α. δ^a is the length of the springs connected to vertex α.
$\underline{\hat{\delta}}^a = (\mathbf{x}^\alpha - \mathbf{x}^\beta)/\delta^a = (\Delta \mathbf{x}^a)/\delta^a$ represents the unit vector along the spring a
that connects vertices α and β.

We relax the model at each quasi-static time step T to achieve force balance
by updating the positions of the particles using

$$\underline{\mathbf{x}}^\alpha(\tau + d\tau) = \underline{\mathbf{x}}^\alpha(\tau) - d\tau \frac{k}{\gamma} \sum_a (\delta^a - \delta_T^a) \, \underline{\hat{\delta}}^a, \tag{19}$$

where $d\tau \frac{k}{\gamma}$ was set to 0.01 (ensuring no numerical artifacts).

The particles were moved until the average movement of the particles
$\langle \|\mathbf{x}^\alpha(\tau + d\tau) - \mathbf{x}^\alpha(\tau)\| \rangle / R$ reduced to 10^{-9}, where R is the radius of curvature
of the outer surface of the spherical cap in the initial stress-free state.

7.9 Spontaneous strain tensor

Tissue shape change during development is modelled in this work as the
appearance of spontaneous strains, a change in the ground state of local length

667 scales. This notion can be captured with a spontaneous strain tensor field,
668 $\underline{\underline{\lambda}}(\underline{X})$, a rank 2 tensor. Each component corresponds to the multiplicative fac-
669 tor by which the rest lengths of the material changes in a particular direction.
670 In some general coordinate system, we can write $\underline{\underline{\lambda}}$ as

$$\underline{\underline{\lambda}} = \begin{pmatrix} \lambda_{11} & \lambda_{12} & \lambda_{13} \\ \lambda_{21} & \lambda_{22} & \lambda_{23} \\ \lambda_{31} & \lambda_{32} & \lambda_{33} \end{pmatrix}. \tag{20}$$

671 We choose the coordinate system so that it aligns with our desired deformation
672 pattern. In this case, $\underline{\underline{\lambda}}(\underline{X})$ is in a diagonal representation :

$$\underline{\underline{\lambda}} = \lambda_{11}(\underline{e}_1 \otimes \underline{e}_1) + \lambda_{22}(\underline{e}_2 \otimes \underline{e}_2) + \lambda_{33}(\underline{e}_3 \otimes \underline{e}_3), \tag{21}$$

where the basis vectors are chosen such that $\underline{e}_1, \underline{e}_2$ are surface tangents while
$\underline{e}_3$ is surface normal. In general, we keep $\lambda_{33} = 1$, since we do not input any
spontaneous strains along the thickness of the model.
The surface components of $\underline{\underline{\lambda}}$ can be further broken down into isotropic
and anisotropic components. Isotropic deformation changes the local area of
the surface by changing the local lengths equally in all directions. Anisotropic
deformation increases the local length in one direction while decreasing the
local length in the other direction so as to preserve the local area. Thus,
we decompose the deformation as a product of isotropic and anisotropic
contributions.

$$\begin{aligned} \lambda_{11} &= \lambda\tilde{\lambda} \\ \lambda_{22} &= \lambda\tilde{\lambda}^{-1}. \end{aligned} \tag{22}$$

683 Then, the spontaneous deformation tensor can be written as

$$\underline{\underline{\lambda}} = \lambda\tilde{\lambda}(\underline{e}_1 \otimes \underline{e}_1) + \lambda\tilde{\lambda}^{-1}(\underline{e}_2 \otimes \underline{e}_2) + (\underline{e}_3 \otimes \underline{e}_3). \tag{23}$$

684 Finally, as $\underline{\underline{\lambda}}$ is a field, each of the components in the above equation generally
685 depend on the location on the surface, $\underline{X}$.
686 **Discretizing $\underline{\underline{\lambda}}$**: As our spring lattice is discrete in nature, we use the following
687 strategy to discretize $\underline{\underline{\lambda}}$. For a single spring, $\underline{\underline{\lambda}}$ is an average of the value of $\underline{\underline{\lambda}}$
688 on the two ends of the springs.

$$\underline{\underline{\lambda}}^a = \frac{1}{2}\left(\underline{\underline{\lambda}}(\underline{X}^\alpha) + \underline{\underline{\lambda}}(\underline{X}^\beta)\right), \tag{24}$$

689 where α and β are the two vertices of the spring a.
690 **Assigning new rest lengths to springs**: The initial length of spring a
691 connecting vertices α and β is given by

$$\delta_I^a = \|\underline{X}^\alpha - \underline{X}^\beta\| = \|\Delta\underline{X}^a\|. \tag{25}$$

To assign new rest lengths, we use   692

$$\delta_F^a = \|\underline{\underline{\lambda}}^a . \Delta\underline{\mathbf{X}}^a\|. \tag{26}$$

Note that we assign a new rest length to any spring a based on the positions   693
of its vertices ($\underline{X}^\alpha$ and $\underline{X}^\beta$), independent of the layer in which these vertices   694
lie (top and bottom).   695
Implementing shape change over time: We increase the spontaneous   696
strain slowly to model the slow build up of stresses due to cell behaviours.   697
Hence, we first calculate the target rest length of springs (δ_F^a). At each time   698
step, we assign a rest length δ_T^a and minimize the energy of the model. We   699
increase δ_T^a in a simple linear manner from δ_I^a to δ_F^a 700

$$\delta_T^a = \delta_I^a + (\delta_F^a - \delta_I^a)\frac{T}{T_F}, \tag{27}$$

where T_F is the number of quasi-static time steps in which the whole simulation   701
takes place. Note that within each time step, the lattice is brought to a force   702
balance state. The simulations were performed for different choices of T_F (1, 2,   703
5), but we found that the differences in output shapes were undetectable. Still,   704
$T_F = 5$ was chosen to simulate the slow appearance of spontaneous strains.   705
Measuring resulting strains in model: In our spring model, displacements   706
are defined by positions of vertices and we define the deformation gradient   707
tensor $\underline{\underline{F}}^\alpha$ at each vertex α of the network.   708
   For each spring a emerging from the vertex α, the deformation gradient   709
tensor should satisfy   710

$$\underline{x}^a = \underline{\underline{F}}^\alpha \underline{X}^a. \tag{28}$$

However, $\underline{\underline{F}}^\alpha$ contains 9 degrees of freedom, while there are 13 springs for each   711
vertex and therefore 13 independent equations to be satisfied. Note that six   712
to eight vertices out of about 3220 vertices in the whole network form point   713
defects and thus have 11 springs. Therefore, we define $\underline{\underline{F}}^\alpha$ as the tensor that   714
best satisfies conditions in Eqs. 28 by minimizing the sum of residuals squared   715

$$S = \sum_a \|\underline{\underline{F}}^\alpha \underline{X}^a - \underline{x}^a\|^2. \tag{29}$$

This is an ordinary least squares (OLS) problem split into three independent   716
basis vectors. We solved this OLS using the Numpy method *numpy.linalg.lstsq*   717
in cartesian coordinates [51]. We then express $\underline{F}$ in the coordinate system cor-   718
responding to vertex α in the model explained above. From this, we calculate   719
the isotropic (F) and anisotropic ($\tilde{F}$) components using   720

$$F = \sqrt{F_{rr}F_{\phi\phi}} \tag{30}$$

$$\tilde{F} = \sqrt{F_{rr}/F_{\phi\phi}}. \tag{31}$$

721 Finally, we compute $\underline{\underline{\lambda}}^{\text{res}}$ as

$$\underline{\underline{\lambda}}^{\text{res}} = \underline{\underline{F}}\,\underline{\underline{\lambda}}^{-1}. \tag{32}$$

722 The isotropic (λ^{res}) and anisotropic components ($\tilde{\lambda}^{\text{res}}$) of $\underline{\underline{\lambda}}^{\text{res}}$ are calculated
723 in the same way as for $\underline{\underline{F}}$.

7.10 Nematic director pattern on spherical surface

In the initial state of the model, we specify a coordinate system on the spherical
surface in different regions (outDVB and DVB). These coordinate systems are
chosen such that the observed nematic patterns of spontaneous strains ($\tilde{\lambda}$)
align with the major axes of the chosen coordinate systems.
We first define the origins in our model similar to the origins defined in
the data. To do so, we first measure $\theta_D V$ (Extended Data Fig. 12 a). The
coordinates of O_D and O_V are then given by ($\pm R\sin(\theta_{DV}/2), 0, R\cos(\theta_{DV}/2)$)
in the cartesian coordinate system. The center for the DVB region is given by
the line O_{DV} which joins O_D and O_V.
In the outDVB region, we have a coordinate system in which the basis vec-
tors are given by $\underline{e}_r, \underline{e}_\phi, \underline{e}_h$ (Extended Data Fig. S8). $\underline{e}_h$ is simply the normal
vector on the spherical surface. To calculate $\underline{e}_r$ at a point, we draw a vector
from the origin in this region (O_D or O_V) to the point. We then take a pro-
jection of this vector onto the tangent plane of the surface and normalize it to
give us a unit vector. In this way, we calculate $\underline{e}_r$ as a surface tangent vector
emanating radially outwards from the origins of the outDVB regions. $\underline{e}_\phi$ is then
the direction perpendicular to $\underline{e}_r$ and $\underline{e}_h$. For each point in the outDVB region,
we calculate the geodesic distance between the point and the center point of
its region. We then normalize this distance by the maximum geodesic distance
from the center calculated in this region. This gives us a scalar coordinate r
which varies from 0 to 1.
In the DVB region, the basis vectors are given by ($\underline{e}_\rho, \underline{e}_w, \underline{e}_h$) (Extended
Data Fig. S8). $\underline{e}_h$ is simply the normal vector on the spherical surface. To
calculate $\underline{e}_\rho$ at a point, we draw a vector from the nearest point on O_{DV} to
the point. We then take a projection of this vector onto the tangent plane of
the surface and normalize it to give us a unit vector. In this way, we calculate
$\underline{e}_\rho$ as a surface tangent vector emanating outwards from the center line of the
DVB region as well as parallel to the DVB. $\underline{e}_w$ is perpendicular to $\underline{e}_\rho$ and
$\underline{e}_h$. For each point in the outDVB region, we calculate the shortest distance
between the point and the center line of the DVB region. We then normalize
this distance by the maximum distance from the center line in the DVB region.
This gives us a scalar coordinate ρ.
For the simple examples presented in Fig 2c (except Fig 2c.iii), θ_{DV} was
set to be 0 to have a simple radial coordinate system. For Fig 2c.iii, $\theta_{DV} > \theta_M$.

7.11 Extracting the strain pattern from segmented images

To quantify the strain due to different cell behaviors along the basis vectors of the chosen coordinate system, we compare cells within topologically tracked bins between two different developmental stages.

Tracking location between developmental stages: We leverage the topological distance coordinate system to track locations between discs. Each topological ring k is given a value N which denotes the cumulative number of cells from the topological origin defined in each region (O_D, O_V, and O_{DV}). We use N to track the location in our static images of different discs at different developmental stages.

Observed strain due to cell area change: Cell area scales with square of the distance between cell vertices. Thus, the factor by which the local lengths change in all directions is given by

$$\lambda_A^*(N) = \sqrt{\frac{A(N, t + \Delta t)}{A(N, t)}}. \tag{33}$$

Here, t corresponds to an initial developmental stage, and $t + \Delta t$ corresponds to a later developmental stage. A refers to the average cell area evaluated at N.

Observed strain due to cell elongation change: Each cell is given a cell elongation tensor $\underline{\underline{Q}}$ that is the average of further subdivisions of the cell polygon into triangles (Methods 7.6). Each triangle can be circumscribed by an ellipse, the centroid of which coincides with the centroid of the triangle. According to [52], the length of the long axis of the ellipse is given by $l = r_o \exp(\|\underline{\underline{Q}}\|)$, where r_o is the radius of a reference equilateral triangle. The length of the short axis of the ellipse is given by $s = r_o \exp(-\|\tilde{\underline{\underline{Q}}}\|)$. The axes of the ellipse match with the radial and tangential directions if the off-diagonal components $Q_{r\phi}$ or $Q_{\rho\phi}$ are approximately 0. This was the case for our data as well. The length scale associated with the radial direction is l if Q_{rr} or $Q_{\rho\rho}$ is positive and s if Q_{rr} or $Q_{\rho\rho}$ is negative. Thus, we get a measure of the length scales along the radial direction, which we denote by L and is given by

$$L = \exp\left(\sigma\|\tilde{\underline{\underline{Q}}}\|\right), \tag{34}$$

where σ is the sign of Q_{rr} or $Q_{\rho\rho}$.

We then average L within each ring and compute a ratio of the length scales along the radial direction between two developmental stages by computing

$$\tilde{\lambda}_Q^*(N) = \frac{L(N, t + \Delta t)}{L(N, t)}. \tag{35}$$

Observed strain due to cell rearrangements: Rearrangements lead to

anisotropic deformation of the tissue. In our topological coordinate system,
radially oriented rearrangements lead to an increase in the number of rings
needed to accommodate some fixed number of cells (Extended Data Fig. S6).
Similarly, tangential rearrangements would lead to a decrease in the number of
topological rings. Thus, by measuring the change in the number of rings needed
to accommodate some fixed number of cells, we can estimate the deformation
due to the net effect of radial and tangential rearrangements.
In a tissue region at developmental stage t, let us consider a single ring with
index k and cumulative number of cells N. Ring k contains ΔN cells given by
$N(k,t) - N(k-1,t)$. By construction, the number of rings needed to contain
ΔN cells at location N is given by $n(N,t) = 1$. For a later developmental stage,
$t + \Delta t$, we estimate $n(N, t + \Delta t)$ which is the number of rings that contain
ΔN cells at the location N. This is done by taking the difference between k
values evaluated at $t + \Delta t$ and at locations $N(k-1,t)$ and $N(k,t)$ (see also
Extended Data Fig. S6)

$$n(N, t + \Delta t) = k(N(k,t), t + \Delta t) - k(N(k-1,t), t + \Delta t) \qquad (36)$$

807 As $n(N,t)$ and $n(N, t + \Delta t)$ are measures of the number of topological rings,
808 they represent the radial topological length scales that change due to cell
809 rearrangements. Thus, the strain due to cell rearrangements is quantified by

$$\lambda_R^*(N) = \frac{n(N, t + \Delta t)}{n(N, t)}. \qquad (37)$$

810 $\lambda_R^*(N) > 1$ represents radial extension of the tissue due to radially oriented
811 rearrangements, while $\lambda_R^*(N) < 1$ represents tangential extension.
812 **Observed strain due to combination of cell elongation change and**
813 **cell rearrangements**: The combined strain due to cell elongation change and
814 cell rearrangements is given by

$$\tilde{\lambda}^*(N) = \tilde{\lambda}_Q^*(N)\tilde{\lambda}_R^*(N). \qquad (38)$$

815
Mapping locations between wing disc images and model: In the model,
we have a dimensionless scalar coordinate in the outDVB and DVB regions
varying from 0 to 1. In the data as well, we prescribe a scalar coordinate to
each topological bin. To do so, we calculate the path length in μm of the
shortest path along cell centers from each cell to the origin and average this
path length for each topological bin. For each bin, we normalize this path
length by the average path length of the outermost topological bin in the
corresponding region (DVB or outDVB). We call this normalized path length
r for the outDVB region and ρ for the DVB region. Due to our normalization,
r and ρ run from 0 to 1, similar to the model.
Thus, we are able to map any topological ring (identified by k and N) to
a scalar coordinate r in outDVB or ρ in DVB. In Methods 7.10, we explain

the mapping between r and ρ to the cartesian coordinates of the vertices in the model given by $\underline{X}^\alpha$, where α is a vertex. Using this mapping, any strain component, for example $\tilde{\lambda}_R^*(\underline{X}^\alpha)$, on the model vertex α can be evaluated from a corresponding $\tilde{\lambda}_R^*(N)$.

7.12 Quantifying curvature of cross-sections

Tissue shape analysis is performed on multi-angle fused SPIM images. We used Fiji re-slicing tools to generate two orthogonal cross-sections along the apical-basal direction. Across-DVB is a cross-section along the center of the long axis of the wing disc. To find the center, we used the position of the sensory organ precursors and general morphology. The along-DVB cross-section follows the DVB and was identified by Ecadherin-GFP signal intensity. The apical pouch shape was outlined manually along both directions over the pouch region up to the HP-fold using custom Fiji macros. Subsequent pouch shape analysis was performed in Python. The tissue shape information was extracted form Fiji into Python using the Python 'read-roi' package.

The extracted apical shapes were aligned and rotated for each wing disc as follows. First, starting from the left-most point in the curve, we measure the arc length of the curve in the clockwise direction. The arc length of the ith point on the curve is given by

$$s(i) = \sum_{i=2}^{n} \|\underline{x}_i - \underline{x}_{i-1}\|, \tag{39}$$

where n is the number of points in the discrete curve and $\underline{x}_i = (x_i, y_i)$ is the position vector of the ith point. We keep $s(i = 1) = 0$.

Next, we define the center of the curve at the middle and offset the arc lengths to have $s = 0$ at the center. This leads to negative arc lengths on the left side of the center and positive arc lengths on the right side of the center (Extended Data Fig. S1).

In order to compute a mean curve from different wing discs of the same developmental stage, we translate and rotate the curves (Extended Data Fig. S1b). We translate each curve by setting their midpoints as the origin $(0, 0)$. To rotate the curve, we compute the center of mass of the curve. Then, we define the new y axis as the line that joins the center of mass to the origin. Finally, for each curve, we smoothen and interpolate between the discrete points using spline interpolation. We use the *scipy.interpolate.UnivariateSpline* function of scipy [53]. To smoothen the spline, we define five knot points, one being the mid-point of the curve, and two others being at three-fourth and half of length from mid-point from either sides. Next, we compute the curvature of each curve using the following expression

$$\kappa = \frac{x'y'' - y'x''}{(x'^2 + y'^2)^{\frac{3}{2}}}, \tag{40}$$

where $'$ refers to the derivative with respect to the parameter of the curve,
which is arc length in our case.

Finally, to compute an average curve, we get the average position vectors at
arc lengths starting from a minimum arc length until a maximum arc length in
intervals of $5\mu m$. We do similar averaging for curvature values to get average
curvature profiles.

To calculate the change in curvature, we normalize each curve from 0 to
1 and use a linear interpolation with 40 positions to subtract the initial from
subsequent curvatures. We then re-introduce the average arc length for each
each developmental stage for each of the normalized positions.

Model output: To quantify the curvature of the model output, we first isolate
the top layer of the lattice. Then, we take the along-DVB cross-section (XZ
plane) and the across-DVB cross-section (YZ plane). To take the cross-section,
we record the points of intersection of the in-surface springs with the respective
plane of the cross-section. From this, we get a discrete set of points that are
ordered along their horizontal position to get a counter-clockwise curve. This
curve data is now similar to the data we obtain from segmented images. Hence,
we apply the exact same procedure describe above to quantify the curvature
of the model output.

883 7.13 Tuning thickness

We tune the thickness of the model in order to change the bending modulus. We
first perform simulations by inputting all cell behaviours (cell area changes, cell
elongation changes and cell rearrangements) combined as spontaneous strains.
We perform this simulation for different thicknesses $h/R = 0.05, 0.1, 0.15$,
where h is the thickness of the model and R is the radius of curvature of the
top surface of the initial state of the model (Extended Data Fig. S9a). We find
that $h/R = 0.1$ gives us the best matching of the curvature change profiles
with the wing disc pouch. Fixing $h/R = 0.1$, we perform further analysis to
infer the spontaneous strains in the wing disc pouch. Inputting these inferred
spontaneous strains, we again performed simulations for different thicknesses.
We found $h/R = 0.1$ still matchs the wing disc pouch curvature change values
best (Extended Data Fig. S9b).

896 Acknowledgements

We thank Christian Dahmann, Anne Classen, and past/present members of the
Eaton, Dye, Modes, Julicher, and Popovic teams for discussions on the project
prior to publication. Miki Ebisuya gave thoughtful feedback on the manuscript.
We also thank the Light Microscopy Facility, the Computer Department, and
the Fly Keepers of the MPI-CBG, as well as the Bioimage Analysis group
at PoL for their support and expertise. This work was funded by Germany's
Excellence Strategy – EXC-2068 – 390729961– Cluster of Excellence Physics
of Life of TU Dresden, as well as grants awarded to SE from the Deutsche

Forschungsgemeinschaft (SPP1782, EA4/10-2) and core funding of the Max-Planck Society. NAD additionally acknowledges funding from the Deutsche Krebshilfe (MSNZ P2 Dresden). AK was funded through the Elbe PhD program. CD acknowledges the support of a postdoctoral fellowship from the LabEx "Who Am I?" (ANR-11-LABX-0071) and the Université Paris Cité IdEx (ANR-18-IDEX-0001) funded by the French Government through its "Investments for the Future". We dedicate this work to our coauthor Prof. Dr. Suzanne Eaton, who tragically passed away near the beginning of the project.

Author Contributions

JFF: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Data Curation, Writing - Original Draft, Visualization
AK: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Data Curation, Writing - Original Draft, Visualization
JP: Methodology, Software, Validation, Data Curation
CD: Methodology, Validation
SE: Conceptualization, Resources, Funding acquisition, Supervision
MP: Methodology, Validation, Writing - Original Draft
FJ: Conceptualization, Writing - Original Draft, Supervision
CDM: Conceptualization, Methodology, Resources, Writing - Original Draft, Project administration, Supervision, Funding acquisition
NAD: Conceptualization, Methodology, Resources, Writing - Original Draft, Project administration, Supervision, Funding acquisition

References

[1] Miriam Osterfield, Xinxin Du, Trudi Schüpbach, Eric Wieschaus, and Stanislav Y. Shvartsman. Three-dimensional epithelial morphogenesis in the developing Drosophila egg. *Dev Cell*, 24(4):400–410, February 2013.

[2] Noah P Mitchell, Dillon J Cislo, Suraj Shankar, Yuzheng Lin, Boris I Shraiman, and Sebastian J Streichan. Visceral organ morphogenesis via calcium-patterned muscle constrictions. *eLife*, 11, May 2022.

[3] Amy E. Shyer, Tuomas Tallinen, Nandan L. Nerurkar, Zhiyan Wei, Eun Seok Gil, David L. Kaplan, Clifford J. Tabin, and L. Mahadevan. Villification: How the gut gets its villi. *Science*, 342(6155):212–218, October 2013.

[4] David Bilder and Saori L. Haigo. Expanding the morphogenetic repertoire: Perspectives from the drosophila egg. *Developmental Cell*, 22(1):12–23, January 2012.

[5] Akankshi Munjal, Edouard Hannezo, Tony Y.-C. Tsai, Timothy J. Mitchison, and Sean G. Megason. Extracellular hyaluronate pressure shaped by

cellular tethers drives tissue morphogenesis. *Cell*, 184(26):6313–6325.e18, December 2021.

[6] Adam C Martin. The physical mechanisms of *Drosophila* gastrulation: Mesoderm and endoderm invagination. *Genetics*, 214(3):543–560, March 2020.

[7] Liyuan Sui, Silvanus Alt, Martin Weigert, Natalie Dye, Suzanne Eaton, Florian Jug, Eugene W. Myers, Frank Jülicher, Guillaume Salbreux, and Christian Dahmann. Differential lateral and basal tension drive folding of drosophila wing discs through two distinct mechanisms. *Nature Communications*, 9(1), November 2018.

[8] Carl Modes and Mark Warner. Shape-programmable materials. *Physics Today*, 69(1):32–38, January 2016. Publisher: American Institute of Physics.

[9] E. Efrati, E. Sharon, and R. Kupferman. Elastic theory of unconstrained non-euclidean plates. *Journal of the Mechanics and Physics of Solids*, 57(4):762–775, 2009.

[10] Taylor H. Ware, Michael E. McConney, Jeong Jae Wie, Vincent P. Tondiglia, and Timothy J. White. Voxelated liquid crystal elastomers. *Science*, 347(6225):982–984, 2015.

[11] Julien Dervaux and Martine Ben Amar. Morphogenesis of growing soft tissues. *Phys. Rev. Lett.*, 101:068101, Aug 2008.

[12] Utpal Nath, Brian C. W. Crawford, Rosemary Carpenter, and Enrico Coen. Genetic control of surface curvature. *Science*, 299(5611):1404–1407, 2003.

[13] James W. Fristrom, W. Robert Logan, and Collin Murphy. The synthetic and minimal culture requirements for evagination of imaginal discs of Drosophila melanogaster in vitro. *Developmental Biology*, 33(2):441–456, August 1973.

[14] L.I. Fessler, M.L. Condic, R.E. Nelson, J.H. Fessler, and J.W. Fristrom. Site-specific cleavage of basement membrane collagen IV during Drosophila metamorphosis. *Development*, 117(3):1061–1069, March 1993.

[15] Maria-del-Carmen Diaz-de-la Loza, Robert P. Ray, Poulami S. Ganguly, Silvanus Alt, John R. Davis, Andreas Hoppe, Nic Tapon, Guillaume Salbreux, and Barry J. Thompson. Apical and Basal Matrix Remodeling Control Epithelial Morphogenesis. *Developmental Cell*, 46(1):23–39.e5, July 2018.

[16] Eva Fekete, Dianne Fristrom, Istvan Kiss, and James W. Fristrom. The mechanism of evagination of imaginal discs of Drosophila melanogaster : II. Studies on trypsin-accelerated evagination. *Wilhelm Roux's Archives of Developmental Biology*, 178(2):123–138, June 1975.

[17] Silvia Aldaz, Luis M. Escudero, and Matthew Freeman. Dual role of myosin II during Drosophila imaginal disc metamorphosis. *Nature Communications*, 4:1761, 2013.

[18] D. Fristrom and C. Chihara. The mechanism of evagination of imaginal discs of Drosophila melanogaster. V. Evagination of disc fragments. *Developmental Biology*, 66(2):564–570, October 1978.

[19] Dianne Fristrom. The mechanism of evagination of imaginal discs of Drosophila melanogaster,: III. Evidence for cell rearrangement. *Developmental Biology*, 54(2):163–171, December 1976.

[20] M L Condic, D Fristrom, and J W Fristrom. Apical cell shape changes during Drosophila imaginal leg disc elongation: A novel morphogenetic mechanism. *Development*, 111:23–33, 1991.

[21] D. K. Fristrom and J. W. Fristrom. The metamorphic development of the adult epidermis. In *The Development of Drosophila Melanogaster*, volume 2, pages 843–897. Cold Spring Harbor Laboratory Press, 1993.

[22] Job Taylor and Paul N. Adler. Cell Rearrangement and Cell Division during the Tissue Level Morphogenesis of Evaginating Drosophila Imaginal Discs. *Developmental biology*, 313(2):739–751, January 2008.

[23] David Hoerl, Fabio Rojas Rusak, Friedrich Preusser, Paul Tillberg, Nadine Randel, Raghav K. Chhetri, Albert Cardona, Philipp J. Keller, Hartmann Harz, Heinrich Leonhardt, Mathias Treier, and Stephan Preibisch. BigStitcher: Reconstructing high-resolution image datasets of cleared and expanded samples. *Nat Methods*, 16(9):870–874, September 2019.

[24] C. D. Modes and M. Warner. Blueprinting nematic glass: Systematically constructing and combining active points of curvature for emergent morphology. *Physical Review E*, 84(2), Aug 2011.

[25] Hillel Aharoni, Eran Sharon, and Raz Kupferman. Geometry of thin nematic elastomer sheets. *Phys. Rev. Lett.*, 113:257801, Dec 2014.

[26] C. D. Modes, K. Bhattacharya, and M. Warner. Gaussian curvature from flat elastica sheets. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 467(2128):1121–1140, 2011.

[27] Carlos M. Duque and Carl D. Modes. Shape programming in entropic tissues, 2023.

[28] Margrit Schubiger and John Palka. Changing spatial patterns of DNA replication in the developing wing of Drosophila. *Developmental Biology*, 123(1):145–153, September 1987.

[29] Dianne Fristrom and James W. Fristrom. The mechanism of evagination of imaginal discs of Drosophila melanogaster: I. General considerations. *Developmental Biology*, 43(1):1–23, March 1975.

[30] Tinri Aegerter-Wilmsen, Alister C. Smith, Alix J. Christen, Christof M. Aegerter, Ernst Hafen, and Konrad Basler. Exploring the effects of mechanical feedback on epithelial topology. *Development (Cambridge, England)*, 137(3):499–506, February 2010.

[31] Loïc LeGoff, Hervé Rouault, and Thomas Lecuit. A global pattern of mechanical stress polarizes cell divisions and cell shape in the growing Drosophila wing disc. *Development*, 140(19):4051–4059, October 2013.

[32] Yanlan Mao, Alexander L Tournier, Andreas Hoppe, Lennart Kester, Barry J Thompson, and Nicolas Tapon. Differential proliferation rates generate patterns of mechanical tension that orient tissue growth. *The EMBO Journal*, 32(21):2790–2803, October 2013.

[33] Natalie A Dye, Marko Popović, K Venkatesan Iyer, Jana F Fuhrmann, Romina Piscitello-Gómez, Suzanne Eaton, and Frank Jülicher. Self-organized patterning of cell morphology via mechanosensitive feedback. *eLife*, 10:e57964, March 2021.

[34] Peter J. Bryant. Cell lineage relationships in the imaginal wing disc of Drosophila melanogaster. *Developmental Biology*, 22(3):389–411, October 1970.

[35] A. Garcia-Bellido, P. Ripoll, and G. Morata. Developmental Compartmentalisation of the Wing Disk of Drosophila. *Nature New Biology*, 245(147):251–253, October 1973.

[36] Natalie A. Dye. Cultivation and Live Imaging of Drosophila Imaginal Discs. *Methods in Molecular Biology (Clifton, N.J.)*, 2540:317–334, 2022.

[37] Johannes Schindelin, Ignacio Arganda-Carreras, Erwin Frise, Verena Kaynig, Mark Longair, Tobias Pietzsch, Stephan Preibisch, Curtis Rueden, Stephan Saalfeld, Benjamin Schmid, Jean-Yves Tinevez, Daniel James White, Volker Hartenstein, Kevin Eliceiri, Pavel Tomancak, and Albert Cardona. Fiji: An open-source platform for biological-image analysis. *Nature Methods*, 9(7):676–682, June 2012.

[38] Jannis Ahlers, Daniel Althviz Moré, Oren Amsalem, Ashley Anderson, Grzegorz Bokota, Peter Boone, Jordão Bragantini, Genevieve Buckley, Alister Burt, Matthias Bussonnier, Ahmet Can Solak, Clément Caporal, Draga Doncila Pop, Kira Evans, Jeremy Freeman, Lorenzo Gaifas, Christoph Gohlke, Kabilar Gunalan, Hagai Har-Gil, Mark Harfouche, Kyle I. S. Harrington, Volker Hilsenstein, Katherine Hutchings, Talley Lambert, Jessy Lauer, Gregor Lichtner, Ziyang Liu, Lucy Liu, Alan Lowe, Luca Marconato, Sean Martin, Abigail McGovern, Lukasz Migas, Nadalyn Miller, Hector Muñoz, Jan-Hendrik Müller, Christopher Nauroth-Kreß, Juan Nunez-Iglesias, Constantin Pape, Kim Pevey, Gonzalo Peña-Castellanos, Andrea Pierré, Jaime Rodríguez-Guerra, David Ross, Loic Royer, Craig T. Russell, Gabriel Selzer, Paul Smith, Peter Sobolewski, Konstantin Sofiiuk, Nicholas Sofroniew, David Stansby, Andrew Sweet, Wouter-Michiel Vierdag, Pam Wadhwa, Melissa Weber Mendonça, Jonas Windhager, Philip Winston, and Kevin Yamauchi. Napari: A multi-dimensional image viewer for Python, July 2023.

[39] Robert Haase, Dohyeon Lee, Draga Doncila Pop, and Laura Žigutytė. Haesleinhuepf/napari-accelerated-pixel-and-object- classification: 0.14.0, October 2023.

[40] Robert Haase, Pradeep Rajasekhar, Talley Lambert, grahamross123, Juan Nunez-Iglesias, Lachie, Clement Caporal, Christophe Avenel, ENicolay, and Elissavet Sandaltzopoulou. clEsperanto/pyclesperanto_prototype: 0.24.1, April 2023.

[41] Marco Musy, Guillaume Jacquenot, Giovanni Dalmasso, Jaewook Lee, Ruben de Bruin, Johannes Soltwedel, Mikael Tulldahl, Ahinoam Pollack, Brandon Hacha, Federico Claudi, Codacy Badger, Anton Sol, RobinEnjalbert, Zhi-Qiang Zhou, Andrey Yershov, Bane Sullivan, Brian Lerner, Daniel Hrisca, Diego Volpatto, Evan, Franco Matzkin, mkerrinrapid, Nico Schlömer, RichardScottOZ, Xushan Lu, and Oliver Schneider. marcomusy/vedo: 2023.4.6, June 2023.

[42] Robert Haase, Johannes Soltwedel, and Kevin Yamauchi. Haesleinhuepf/napari-process-points-and-surfaces: 0.5.0, April 2023.

[43] Charles Law James Ahrens, Berk Geveci. Paraview: An end-user tool for large-data visualization. In Charles D. Hansen and Chris R. Johnson, editors, *Visualization Handbook*, pages 717–731. Butterworth-Heinemann, January 2005.

[44] Alexander Krull, Tim-Oliver Buchholz, and Florian Jug. Noise2Void - Learning Denoising from Single Noisy Images, April 2019.

[45] Natalie A. Dye, Marko Popović, Stephanie Spannl, Raphaël Etournay, Dagmar Kainmüller, Suhrid Ghosh, Eugene W. Myers, Frank Jülicher,

and Suzanne Eaton. Cell dynamics underlying oriented growth of the *Drosophila* wing imaginal disc. *Development*, page dev.155069, January 2017.

[46] Benoit Aigouy, Daiki Umetsu, and Suzanne Eaton. Segmentation and Quantitative Analysis of Epithelial Tissues. *Methods in Molecular Biology (Clifton, N.J.)*, 1478:227–239, 2016.

[47] Raphaël Etournay, Matthias Merkel, Marko Popović, Holger Brandl, Natalie A Dye, Benoît Aigouy, Guillaume Salbreux, Suzanne Eaton, and Frank Jülicher. TissueMiner: A multiscale analysis toolkit to quantify how cellular processes create tissue dynamics. *eLife*, 5:e14334, May 2016.

[48] Manish Jaiswal, Namita Agrawal, and Pradip Sinha. Fat and Wingless signaling oppositely regulate epithelial cell-cell adhesion and distal wing development in Drosophila. *Development*, 133(5):925–935, March 2006.

[49] Matthias Merkel. From cells to tissues: Remodeling and polarity reorientation in epithelial tissues, 2014.

[50] Zanini, F., Nepusz, T., Horvát, S., and Traag, V. Igraph/python-igraph, January 2015.

[51] Charles R. Harris, K. Jarrod Millman, Stéfan J. van der Walt, Ralf Gommers, Pauli Virtanen, David Cournapeau, Eric Wieser, Julian Taylor, Sebastian Berg, Nathaniel J. Smith, Robert Kern, Matti Picus, Stephan Hoyer, Marten H. van Kerkwijk, Matthew Brett, Allan Haldane, Jaime Fernández del Río, Mark Wiebe, Pearu Peterson, Pierre Gérard-Marchant, Kevin Sheppard, Tyler Reddy, Warren Weckesser, Hameer Abbasi, Christoph Gohlke, and Travis E. Oliphant. Array programming with NumPy. *Nature*, 585(7825):357–362, September 2020.

[52] Matthias Merkel, Raphaël Etournay, Marko Popović, Guillaume Salbreux, Suzanne Eaton, and Frank Jülicher. Triangles bridge the scales: Quantifying cellular contributions to tissue deformation. *Physical Review E*, 95(3):032401, 2017.

[53] Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, Pearu Peterson, Warren Weckesser, Jonathan Bright, Stéfan J. van der Walt, Matthew Brett, Joshua Wilson, K. Jarrod Millman, Nikolay Mayorov, Andrew R. J. Nelson, Eric Jones, Robert Kern, Eric Larson, C J Carey, İlhan Polat, Yu Feng, Eric W. Moore, Jake VanderPlas, Denis Laxalde, Josef Perktold, Robert Cimrman, Ian Henriksen, E. A. Quintero, Charles R. Harris, Anne M. Archibald, Antônio H. Ribeiro, Fabian Pedregosa, Paul van Mulbregt, and SciPy 1.0 Contributors. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nature Methods*, 17:261–272, 2020.