Senescent cells inhibit muscle differentiation via the lipid- SASP 15d-PGJ$_2$ mediated modification and control of HRas

Swarang Sachin Pundlik[1,2], Snehasudha Subhadarshini Sahoo[1,3], Alok Barik[1], Ashwin Venkateshvaran[1], Mahapatra Anshuman Jaysingh[4,5], Raviswamy G H Math[6], Arvind Ramanathan[1,*]

1. Metabolic Regulation of Cell Fate (RCF), Institute for Stem Cell Science and Regenerative Medicine (InStem), Bangalore Life Science Cluster, GKVK-Post, Bellary Road, Bengaluru, Karnataka-560065, India.
2. Manipal Academy of Higher Education (MAHE), Tiger Circle Road, Madhav Nagar, Manipal, Karnataka-576104, India.
3. University of North Carolina at Chapel Hill, 216 Lenoir Dr, Chapel Hill, North Carolina-27599 U. S. A.
4. Department of Biological Sciences, Indian Institute of Science Education and Research Kolkata (IISER-K), Campus Road, Mohanpur, West Bengal-741246, India.
5. Division of Biology and Biomedical Sciences, Washington University in St Louis, 1 Brookings Dr, St. Louis, Missouri-63130, U. S. A.
6. National Centre for Biological Sciences (NCBS), Rajiv Gandhi Nagar, Kodigehalli, Bengaluru, Karnataka-560065, India.

* All correspondence should be written to Dr. Arvind Ramanathan at arvind@instem.res.in

Abstract

Senescent cells which are characterized by multiple features such as increased
expression of SA β-gal and cell cycle inhibitors such as p21 or p16, accumulate with
tissue damage and dysregulate tissue homeostasis. In the context of skeletal
muscle, it is known that agents used for chemotherapy such as doxorubicin cause
buildup of senescent cells, leading to the inhibition of tissue regeneration. Senescent
cells influence neighboring cells via numerous secreted factors which form the
senescence-associated secreted phenotype (SASP). Lipids are emerging as a key
component of SASP that can control tissue homeostasis. Arachidonic acid-derived
lipids have been shown to accumulate within senescent cells, specifically 15d-PGJ$_2$,
which is an electrophilic lipid produced by the non-enzymatic dehydration of the
prostaglandin PGD$_2$. In this study, we show that 15d-PGJ$_2$ is also released by
Doxorubicin-induced senescent cells as a SASP factor. Treatment of skeletal muscle
myoblasts with the conditioned medium from these senescent cells inhibits myoblast
fusion during differentiation. Inhibition of L-PTGDS, the enzyme that synthesizes
PGD$_2$, diminishes the release of 15d-PGJ$_2$ by senescent cells and restores muscle
differentiation. We further show that this lipid post-translationally modifies Cys184 of
HRas in skeletal muscle cells, causing a reduction in the localization of HRas to the
Golgi, increased HRas binding to RAF RBD, and activation of cellular MAPK-Erk
signaling (but not the Akt signaling). Mutating C184 of HRas prevents the ability of
15d- PGJ$_2$ to inhibit the differentiation of muscle cells and control the activity of
HRas. This work shows that 15d-PGJ$_2$ released from senescent cells could be
targeted to restore muscle homeostasis after chemotherapy.

Introduction

Senescent cells are important drivers of aging and damage-associated loss of tissue homeostasis(Childs et al., 2015). Anti-cancer chemotherapy presents an important context where treatment with chemotherapeutics such as Doxorubicin (Doxo) causes widespread cellular senescence which inhibits tissue homeostasis and regeneration, including in skeletal muscles(Francis et al., 2022). It has been shown that Doxo causes systemic inflammation and leads to the emergence of senescent cells across tissues(Di Leonardo et al., 1994; Hu and Zhang, 2019; Robles and Adami, 1998). Senescent cells negatively affect tissue homeostasis and regeneration by releasing factors including proteins like growth factors, matrix metalloproteases, cytokines, and chemokines, and small molecules like fatty acid derivatives(Campisi, 2005; Coppé et al., 2010; Dilley et al., 2003; Krtolica et al., 2001; Parrinello et al., 2005; Shelton et al., 1999; Yang et al., 2006). The release of these factors from senescent cells is called the Senescence Associated Secretory Phenotype (SASP). It is expected that these SASP factors and their mechanisms of action will vary depending on cellular and tissue contexts. Identifying SASP factors and their underlying mechanistic targets will be critical for building an understanding of how senescent cells control tissue homeostasis(Coppé et al., 2010; Davalos et al., 2010). Lipids are a less explored family of SASP factors, and it is important to understand how they affect tissue regeneration(Hamsanathan and Gurkar, 2022). We have previously shown that senescent cells have increased intracellular levels of prostaglandin 15d-PGJ$_2$(Wiley et al., 2021), a non-enzymatic dehydration product of prostaglandin PGD$_2$(Shibata et al., 2002). In the context of skeletal muscle, PGD$_2$ and 15d-PGJ$_2$ have been shown to negatively regulate muscle differentiation via mechanisms that do not depend on a cognate receptor(Hunter et al., 2001; Veliça et al., 2010). Here we study the role of 15d-PGJ$_2$ as a member of SASP and identify mechanisms by which it might negatively affect muscle regeneration. 15d-PGJ$_2$ has been previously shown to covalently modify multiple proteins like MAPK1, MCM4, EIF4A-I, PKM1, GFAP etc. endothelial and neuronal cells(Marcone and Fitzgerald, 2013; Yamamoto et al., 2011). 15d-PGJ$_2$ was shown to be covalently modifying HRas in NIH3T3, Cos1, and IMR90 cell lines(Luis Oliva et al., 2003; Wiley et al., 2021). We further studied HRas as an important target that might mediate the effects of 15d-PGJ$_2$ on muscle differentiation via covalent modification. We investigated HRas as a possible

effector of 15d-PGJ$_2$ because (i) HRas belongs to the Ras superfamily of small molecule GTPases and is known regulator of key cellular processes(BOHR Orthopredic Hospital et al., 1964; Davis et al., 1983; Kirsten and Mayer, 1967; Vetter and Wittinghofer, 2001). (ii) constitutively active HRas mutant (HRas V12) has been shown to inhibit the differentiation of myoblasts by inhibiting MyoD and Myogenin expression(Konieczny et al., 1989; Lassar et al., 1989; Olson,' et al., 1987; Van Der Burgt et al., 2007). (iii) Downstream signaling of HRas is important for muscle homeostasis as skeletal and cardiac myopathies are observed in individuals carrying constitutively active mutants of HRas(Engler et al., 2021; Konieczny et al., 1989; Lee et al., 2010; Olson,' et al., 1987; Scholz et al., 2009; Van Der Burgt et al., 2007). HRas is highly regulates by lipid modifications, it undergoes reversible palmitoylation and de-palmitoylation at C-terminal cysteines, which regulate the intracellular distribution and activity of HRas(Gutierrez et al., 1989; Lu and Hofmann, 1995; Rocks et al., 2005). In this study, we show that 15d-PGJ$_2$ is synthesized and released by senescent myoblasts upon treatment with Doxo. 15d-PGJ$_2$, taken up by the myoblasts, covalently modifies HRas at cysteine 184 and activates it. We also show that previously reported inhibition of differentiation of myoblasts by 15d-PGJ$_2$ depends on HRas C-terminal cysteines, notably cysteine 184. This study provides a mechanism by which prostaglandins secreted as SASP inhibit the differentiation of myoblasts, affecting muscle homeostasis in patients undergoing chemotherapy.

Results

Doxorubicin (Doxo) treatment induces senescence in mouse skeletal muscles and C2C12 mouse myoblasts.

Doxorubicin-mediated DNA damage has been shown to induce senescence in cells(Di Leonardo et al., 1994; Hu and Zhang, 2019; Robles and Adami, 1998). Therefore, we injected B6J mice with Doxorubicin (Doxo) (5mg/kg) every 3 days for 9 days and observed induction of DNA damage-mediated senescence in hindlimb skeletal muscles (Fig. S1A). We observed an increase in the expression of p21 and increased nuclear levels of the DNA damage marker γH2A.X in mouse Gastrocnemius muscles (Fig. 1A and B). We also observed a significant increase in the mRNA levels of known senescence markers (p16 and p21), SASP factors (CXCL1, CXCL2, TNF1α, IL6, TGFβ1) in skeletal muscles of mice treated with Doxo

compared to that of mice treated with saline (Fig. 1C). These observations suggest that there is induction of senescence in skeletal muscles of mice upon treatment with Doxo.

We then measured the induction of senescence *in vitro* in C2C12 mouse myoblasts after treatment with Doxo (Fig. S1B). We treated C2C12 myoblasts with Doxo (150 nM) and observed a flattened morphology and increased cell size (Fig. S1C). We also observed a significant increase in the size of nuclei in C2C12 cells after treatment with Doxo (Fig. 1D) and increased senescence-associated β galactosidase (SA-β gal) activity in cells treated with Doxo (Fig. S1D). These observations suggest the induction of senescence in C2C12 cells after treatment with Doxo.

Doxo-mediated senescence induces synthesis and release of 15d-PGJ$_2$ in C2C12 myoblasts and mouse skeletal muscle.

Synthesis of prostaglandins by senescent cells has previously been reported(Wiley et al., 2021; Wiley and Campisi, 2021). Specifically, levels of PGD$_2$ and its metabolite 15d-PGJ$_2$ have been shown to be significantly increased. Therefore, we measured the levels of mRNA of enzymes involved in the synthesis of PGD$_2$/15d-PGJ$_2$ (PTGS1, PTGS2, and PTGDS), in the gastrocnemius muscle of mice after treatment with Doxo. We observed a significant increase in the mRNA levels of PTGS1, PTGS2, and PTGDS enzymes in the skeletal muscle of mice treated with Doxo (Fig. 1C). We also observed a time-dependent increase in the mRNA levels of PTGS1, PTGS2, PTGDS, and PTGES enzymes in C2C12 cells treated with Doxo compared to Day 0 (Fig. 1E). Expression of enzyme PTGES was elevated on Day 4, whereas the expression of Prostaglandin D synthase (PTGDS) increased only after Day 8, reaching maximum expression on Day 12. These observations suggest an increase in the synthesis of prostaglandins in senescent cells.

15d-PGJ$_2$ is a non-enzymatic dehydration product of PGD$_2$(Shibata et al., 2002). We observed an increase in the mRNA levels synthetic enzymes of 15d-PGJ$_2$ in senescent C2C12 cells. Therefore, we measured the levels of extracellular 15d-PGJ$_2$. We collected the conditioned medium from senescent and quiescent C2C12 cells and observed an increase in the concentration of 15d-PGJ$_2$ in the medium of senescent cells (Fig. 1F). This shows an increased release of 15d-PGJ$_2$ by senescent myoblasts.

Prostaglandin PGD_2 and its metabolites in the conditioned medium of senescent cells inhibit the differentiation of C2C12 myoblasts.

15d-PGJ_2 (the final non-enzymatic dehydration product of PGD_2) has been shown to inhibit the differentiation of myoblasts(Hunter et al., 2001). We observed the release of 15d-PGJ_2 by senescent cells, showing that 15d-PGJ_2 is a SASP factor (Fig. 1F). Conditioned medium of senescent cells inhibits the differentiation of myoblasts in myotonic dystrophy type 1(Conte et al., 2023). Therefore, we tested whether 15d-PGJ_2, the terminal dehydration product of PGD_2, is required for the inhibitory effect of SASP on the differentiation of myoblasts. We treated C2C12 myoblasts with the conditioned medium of senescent cells or senescent cells treated with 30 µM of AT-56 (a well-characterized inhibitor of prostaglandin D synthase (PTGDS))(Hu et al., 2021; S. Hu et al., 2023; Shunfeng Hu et al., 2023; Irikura et al., 2009). We observed a decrease in the fusion index of the C2C12 myoblasts treated with the conditioned medium of senescent cells (Fig. 2A), suggesting that SASP factors decrease the differentiation of myoblasts. This decrease in inhibition was rescued in myoblasts treated with the conditioned medium of senescent cells treated with AT-56 (Fig. 2A). This suggests that prostaglandins PGD_2/15d-PGJ_2 released by senescent cells as SASP factors can inhibit the differentiation of myoblasts.

15d-PGJ_2 inhibits the proliferation and differentiation of mouse and human myoblasts.

15d-PGJ_2 has been shown to affect the proliferation of cancer cell lines, both positively and negatively (Chen et al., 2003; Choi et al., 2020; Slanovc et al., 2024; Yen et al., 2014). We measured the effect of 15d-PGJ_2 on the proliferation of C2C12 myoblasts. We treated C2C12 myoblasts with 15d-PGJ_2 (10 µM) or DMSO for 72 hours. We observed a decrease in the proliferation of C2C12 cells after treatment with 15d-PGJ_2 (Fig. 2B). The doubling time of C2C12 cells was also increased upon treatment with 15d-PGJ_2 (57.24 hours) compared to DMSO (13.76 hours). This suggests that 15d-PGJ_2 decreases the proliferation of C2C12 myoblasts.

We measured the differentiation of C2C12 mouse and primary human myoblasts after treatment with 15d-PGJ_2. C2C12 myoblasts showed cell death after 24 hours and 48 hours upon treatment with 15d-PGJ_2 (10 µM and 5 µM) respectively in the C2C12 differentiation medium (Fig. S2A). Therefore, we treated differentiating

myoblasts with 15d-PGJ$_2$ (1 µM, 2 µM, and 4 µM) for 5 days. We observed a dose-

dependent decrease in the mRNA levels of MyoD, MyoG, and MHC in differentiating

C2C12 cells after treatment with 15dPGJ$_2$ (Fig. 2C). We also observed a decrease in

the fusion of myoblasts into myotubes in C2C12 cells treated with 15d-PGJ$_2$ (4 µM)

compared to DMSO (Fig. 2D). We also observed a dose-dependent decrease in the

protein levels of MHC in differentiating primary human myoblasts upon treatment

with 15d-PGJ$_2$ (Fig. 2E). These observations suggest that 15d-PGJ$_2$ inhibits the

differentiation of both mouse and human myoblasts.

Biotinylated 15d-PGJ$_2$ covalently modifies HRas at Cysteine 184.

15d-PGJ$_2$ has been shown to covalently modify several proteins including p53 and

NF-κB, which are involved in several key biological processes(Marcone and

Fitzgerald, 2013). HRas was identified to be covalently modified by 15d-PGJ$_2$ at

cysteine 184 in NIH3T3 and Cos1 cells(Luis Oliva et al., 2003). Therefore, we tested

whether 15d-PGJ$_2$ could covalently modify HRas in C2C12 cells. We treated C2C12

cells expressing the EGFP-tagged wild-type HRas with biotinylated 15d-PGJ$_2$ (5

µM). We then immunoprecipitated biotinylated 15d-PGJ$_2$ using streptavidin. We

observed a significant increase in the pulldown of HRas upon treatment with 15d-

PGJ$_2$-biotin compared to DMSO (Fig. 3A), suggesting an interaction between 15d-

PGJ$_2$ and HRas. To measure the role of individual C-terminal cysteines in the binding

of HRas with 15d-PGJ$_2$, we treated C2C12 cells expressing the EGFP-tagged

C181S and C184S mutants of HRas with biotinylated 15d-PGJ$_2$ (5 µM), and

immunoprecipitated using streptavidin. We observed that the intensity of EGFP-

tagged HRas was significantly decreased in cells expressing the C184S mutant but

not in those expressing the C181S mutant (Fig. 3A). This suggests that 15d-PGJ$_2$

covalently modifies HRas at cysteine 184 in C2C12 cells.

15d-PGJ$_2$ increases the FRET between EGFP-HRas and mCherry-RAF RBD in

wild-type and C181S mutant but not in the C184S mutant of HRas.

We next tested the effect of covalent modification of HRas by 15d-PGJ$_2$ on HRas

GTPase activity using FRET. mCherry RAF RBD is a well characterized sensor of

activity of HRas. RAF RBD binds to the activated HRas upon activation of HRas,

allowing FRET between EGFP and mCherry(Rocks et al., 2005). We co-expressed

EGFP-tagged HRas (EGFP-HRas) with mCherry-tagged Ras Binding Domain of

RAF kinase (mCherry-RAF RBD) in C2C12 myoblasts (Fig. 3B). We measured the

efficiency of FRET between EGFP and mCherry using an ImageJ plugin, FRET

analyzer(Hachet-Haas et al., 2006). We compared the mean acceptor normalized

FRET index in C2C12 myoblasts expressing EGFP-HRas WT before and after

treatment of 15d-PGJ$_2$ (10 µM) for 1 hour. We observed an increase in the mean

acceptor normalized FRET index upon treatment with 15d-PGJ$_2$ (Fig. 3C). This

suggests that 15d-PGJ$_2$ activates HRas. To measure the role of individual C-terminal

cysteines in 15d-PGJ$_2$ mediated activation of HRas, we co-expressed EGFP-HRas

C181S or C184S with mCherry-RAF RBD in C2C12 myoblasts. We measured the

mean acceptor normalized FRET index before and after 1 hour of treatment with

15d-PGJ$_2$ (10 µM). We observed an increase in the mean acceptor normalized FRET

index in cells expressing EGFP-HRas C181S upon treatment with 15d-PGJ$_2$ but not

in cells expressing EGFP-HRas C184S (Fig. 3D). These observations suggest that

activation of HRas by 15d-PGJ$_2$ occurs in a cysteine 184 dependent manner.

15d-PGJ$_2$ increases phosphorylation of Erk (Thr202/Tyr204) but not Akt (S473)

in C2C12 myoblasts.

HRas regulates two major downstream signaling pathways, the MAP kinase (MAPK)

pathway and the PI3 kinase (PI3K) pathway(Pylayeva-Gupta et al., 2011). We tested

the effects of treatment with 15d-PGJ$_2$ on these two downstream signaling pathways

by measuring the phosphorylation of Erk (42 kDa and 44 kDa) and Akt proteins in

C2C12 cells. We treated C2C12 cells with 15d-PGJ$_2$ (5 µM and 10 µM) or DMSO for

1 hr (after 24 hrs. of serum starvation) and observed a dose-dependent increase in

the phosphorylation of Erk (T202/Y204) (42 kDa) but not of Erk (44 kDa) (Fig. 3E).

We did not observe an increase in the phosphorylation of Akt (S473) in C2C12 cells

after treatment with 15d-PGJ$_2$ (Fig. S3C). These observations suggest that 15d-PGJ$_2$

activates the MAPK signaling pathway, but not the PI3K signaling pathway.

15d-PGJ$_2$ contains a reactive electrophilic center in its cyclopentenone ring, that can

react with cysteine residues of proteins(Luis Oliva et al., 2003). We tested its role in

activating the MAPK signaling pathway. We measured the phosphorylation of Erk

(42kDa and 44 kDa) in C2C12 cells after treatment with cells with 9,10-dihydro-15d-

PGJ$_2$ (10 µM), a 15d-PGJ$_2$ analog which is devoid of the electrophilic center, for 1 hr

(after 24 hr. of serum starvation). We observed that the phosphorylation of Erk (42

kDa and 44 kDa) in C2C12 cells treated with 9,10-dihydro-15d-PGJ$_2$ was
significantly reduced as compared to the treatment with 15d-PGJ$_2$ (Fig. 3F). This
shows that 15d-PGJ$_2$ activates the HRas-MAPK signaling pathway via the
electrophilic center in its cyclopentenone ring.

15d-PGJ$_2$ increases the localization of EGFP-tagged HRas at the plasma
membrane compared to the Golgi in a C-terminal cysteine-dependent manner.

15d-PGJ$_2$ covalently modifies cysteine 184 and activates HRas signaling (Fig. 3).
Reversible palmitoylation of cysteine 181 and cysteine 184 in the C-terminal tail of
HRas regulate intracellular distribution and signaling of HRas. Inhibition of
palmitoylation of the C-terminal cysteine 181, either by a palmitoylation inhibitor 2-
Bromopalmitate or by mutation to serine, causes accumulation of HRas at the Golgi
compared to the plasma membrane and alters activity(Rocks et al., 2005).
Therefore, we tested whether the modification of 15d-PGJ$_2$ alters the intracellular
distribution of HRas. We co-expressed the EGFP-tagged wild type and the C181S
and the C184S mutants of HRas with a previously reported marker of Golgi(Shaner
et al., 2008) in C2C12 cells and stained the cells with plasma membrane marker
WGA-633 (Fig. 4A). We compared R$_{mean}$, the ratio of mean EGFP-HRas intensity at
the Golgi to the mean HRas intensity at the plasma membrane, to measure the
distribution of HRas between the plasma membrane and the Golgi. We measured
the intracellular distribution of HRas between the Golgi and the plasma membrane in
C2C12 cells after treatment with 15d-PGJ$_2$ (10 μM) for 24 hours and observed a
decrease in the R$_{mean}$ of C2C12 cells expressing the wild-type HRas after treatment
with 15d-PGJ$_2$ (Fig. 4B). However, we did not observe a change in the R$_{mean}$ of
C2C12 cells expressing HRas C181S or HRas C184S after treatment with 15d-PGJ$_2$
(Fig. 4C). These observations suggest that 15d-PGJ$_2$ increases the localization of
HRas at the plasma membrane as compared to that in the Golgi in an HRas C-
terminal cysteine-dependent manner.

15d-PGJ$_2$ mediated inhibition of differentiation of C2C12 cells is rescued by
C181S and C184S mutants of HRas.

247 HRas inhibits the differentiation of C2C12 myoblasts(Engler et al., 2021; Konieczny
248 et al., 1989; Lassar et al., 1989; Lee et al., 2010; Olson,' et al., 1987; Scholz et al.,
249 2009; Van Der Burgt et al., 2007).15d-PGJ$_2$ covalently modifies cysteine 184 and

activates HRas (Fig. 3). Therefore, we tested whether the inhibition of myoblast differentiation by 15d-PGJ$_2$ depends on the activation of HRas signaling by modification of the C-terminal cysteine 184. We expressed the wild-type and the cysteine mutants of HRas (EGFP-HRas WT/C181S/C184S) in C2C12 myoblasts and treated the cells with 15d-PGJ$_2$ (4 µM) or DMSO during differentiation. We observed a decrease in the levels of mRNA of MHC in C2C12 cells expressing HRas WT and HRas C181S after 5 days of treatment with 15d-PGJ$_2$. We did not observe this in expressing HRas C184S (Fig. 4D). We also observed a significant decrease in the protein levels of MHC in differentiating C2C12 cells expressing HRas WT and HRas C181S after treatment with 15d-PGJ$_2$ (Fig. 4E). This decrease was partially rescued in cells expressing HRas C184S (Fig. 4E). These observations suggest that the inhibition of myoblast differentiation by 15d-PGJ$_2$ depends on modification of HRas C-terminal cysteine 184.

Discussion

Senescence is characterized by an irreversible arrest in cell proliferation(Hayflick, 1965). Cells undergo senescence because of a myriad of stresses, including DNA damage, mitochondrial damage, and oncogene overexpression(Bihani et al., 2007, 2004; Casar et al., 2018; Chen and Ames, 1994; Chen et al., 1998; Coppé et al., 2008; D'Adda Di Fagagna, 2008; D'Adda Di Fagagna et al., 2003; Di Leonardo et al., 1994; Franza et al., 1986; Land et al., 1983; Robles and Adami, 1998; Serrano et al., 1997; Wiley et al., 2016; Woods et al., 1997). Senescent cells exhibit a multi-faceted physiological response, where they exhibit a flattened morphology, increase in cell size(Chen and Ames, 1994; Serrano et al., 1997), upregulation of tumor suppressor proteins(Calabrese et al., 2009; Lowe et al., 2004; Stein et al., 1990; Zindy et al., 2003), expression of neutral pH active β-galactosidase(Dimri et al., 1995; Lee et al., 2006), and altered metabolic state(Bittles and Harper, 1984; Jones et al., 2005; Wiley and Campisi, 2021, 2016; Zwerschke et al., 2003). Arachidonic acid metabolism is upregulated in senescent cells, which leads to increased synthesis of eicosanoid prostaglandins, which regulate the physiology of senescent cells(Wiley et al., 2021; Wiley and Campisi, 2021). Senescent cells exhibit a secretory phenotype (SASP) consisting of a variety of bioactive molecules including cytokines and chemokines, growth factors, matrix metalloproteases etc(Coppé et al., 2008). Senescent cells influence the surrounding cells via the SASP factors, which regulate

proliferation, migration, and other cell biological processes in the neighboring
cells(Campisi, 2005). SASP-mediated perturbations in the microenvironment are
implicated in several senescence-associate pathologies(Wiley and Campisi, 2021).
Senescent fibroblasts increase the proliferation of premalignant and malignant
epithelial cells(Krtolica et al., 2001). Conditioned medium of senescent fibroblasts
promoted tumorigenesis in mouse keratinocytes(Dilley et al., 2003). Senescent
fibroblasts transform pre-malignant breast cancer cells into invasive, tumor-forming
cells(Parrinello et al., 2005). Senescence in muscle stem cells induces sarcopenia
via activation of p38 MAP kinase pathway and transient inhibition of the p38 MAP
kinases rejuvenates aged muscle stem cells to ameliorate sarcopenia(Cosgrove et
al., 2014). Senescent cells inhibit the differentiation of myoblasts by secretion of IL6
by senescent muscle stem cells in myotonic dystrophy(Conte et al., 2023).

In this study, we show that senescent myoblasts synthesize and release eicosanoid
prostaglandin 15-deoxy-$\Delta^{12,14}$-prostaglandin J_2 (15d-PGJ$_2$) (Fig. 1E and F), the
terminal non-enzymatic dehydration product of prostaglandin PGD$_2$(Shibata et al.,
2002). We used Doxorubicin (Doxo) to induce senescence in C2C12 myoblasts and
showed that the conditioned medium of senescent C2C12 cells inhibits differentiation
of C2C12 myoblasts (Fig. 2A). Inhibition of synthesis of PGD$_2$ by treatment of
senescent cells with AT-56, a well-characterized inhibitor of prostaglandin D
synthase(Hu et al., 2021; S. Hu et al., 2023; Shunfeng Hu et al., 2023; Irikura et al.,
2009), rescued this inhibitory effect of the conditioned medium (Fig. 2A). A study has
shown that prostaglandin PGD$_2$ inhibits differentiation of C2C12 myoblasts(Veliça et
al., 2010), but the authors noted that knockout of DP1 and DP2 (the known receptors
of prostaglandins PGD$_2$) does not abrogate inhibition of differentiation of myoblasts
by PGD$_2$. This observation suggested that PGD$_2$ might inhibit the differentiation of
myoblasts by a receptor-independent mechanism, possibly by its spontaneous non-
enzymatic dehydration to 15d-PGJ$_2$. 15d-PGJ$_2$ has been suggested to be an
endogenous ligand of PPARγ(Li et al., 2019). However, the inhibition of PPARγ did
not abrogate the inhibition of differentiation of C2C12 myoblasts by 15d-PGJ$_2$,
suggesting existence other possible mechanisms(Hunter et al., 2001). 15d-PGJ$_2$ has
varied effects on cell physiology in a context dependent manner. On one hand, 15d-
PGJ$_2$ promotes tumorigenesis by inducing epithelial to mesenchymal transition in
breast cancer cell line MCF7(Choi et al., 2020), 15d-PGJ$_2$ inhibits proliferation of

A549, H1299, and H23 lung adenocarcinoma cells via induction of ROS and activation of apoptosis(Slanovc et al., 2024). Here, we show that 15d-PGJ$_2$ inhibits the proliferation and the differentiation of C2C12 myoblasts (Fig. 2B, C and D).

15d-PGJ$_2$ contains an electrophilic cyclopentenone ring in its structure, allowing 15d-PGJ$_2$ to covalently modify and form a Michael adducts with cysteine residues of proteins(Shibata et al., 2002). A previous proteomic study in endothelial cells showed biotinylated 15d-PGJ$_2$ covalently modified over 300 proteins, which regulate several physiological processes including cell cycle (MAPK1, MCM4), cell metabolism (Fatty acid synthase, Isocitrate dehydrogenase), apoptosis (PDCD6I), translation regulation (Elongation factor 1 and 2, EIF4A-I), intracellular transport (Importin subunit β1, Exportin 2, Kinesin 1 heavy chain)(Marcone and Fitzgerald, 2013). Another proteomic study in neuronal cells suggested that 15d-PGJ$_2$ modifies several proteins including chaperone HSP8A, glycolytic proteins Enolase 1 and 2, GAPDH, PKM1, cytoskeleton proteins Tubulin β2b, β actin, GFAP etc(Yamamoto et al., 2011). This study also showed modification of a peptide fragments homologous to IκB kinase β, Thioredoxin, and a small molecule GTPase HRas. 15d-PGJ$_2$ modifies HRas in NIH3T3 and Cos1 cells(Luis Oliva et al., 2003) and IMR90 cells(Wiley et al., 2021). Modification by 15d-PGJ$_2$ led to activation of HRas, judged by an increase in GTP bound HRas. It is clear that 15d-PGJ$_2$ is capable of modifying numerous proteins in different contexts. In spite of these observations, the functional relevance of these modifications in numerous contexts remains to be mapped. Here we focused on the role of 15d-PGJ$_2$ in the context of senescence and skeletal muscle differentiation. In this study, we showed that 15d-PGJ$_2$ covalently modifies HRas at cysteine 184 but not cysteine 181 in C2C12 myoblasts (Fig. 3A). We showed by FRET microscopy that modification of HRas by 15d-PGJ$_2$ in HRas WT and HRas C181S activates HRas in C2C12 cells, but 15d-PGJ$_2$ is unable to activate HRas C184S in this context (Fig. 3C and D). This observation shows a direct link between modification of HRas by 15d-PGJ$_2$ and activation of HRas GTPase.

HRas activates two major downstream signaling pathways, the HRas-MAPK and the HRas-PI3K pathway(Pylayeva-Gupta et al., 2011). We showed that covalent modification of HRas by 15d-PGJ$_2$ via the electrophilic cyclopentenone ring activates HRas (Fig. 3C and D) and activates the HRas-MAPK pathway, demonstrated by an increase in the phosphorylation of Erk after treatment with 15d-PGJ$_2$ (Fig. 3E and

F). However, we did not observe activation of the HRas-PI3K pathway, as we did not

see an increase in the phosphorylation of Akt after treatment with 15d-PGJ$_2$ (Fig.

S3C). MAPK and PI3K pathways are known regulators of muscle

differentiation(Bennett and Tonks, 1997; Rommel et al., 1999), where inhibition of the

RAF-MEK-Erk pathway or activation of PI3K pathway promotes differentiation of

myoblasts. Preferential activation of HRas-MAPK over HRas-PI3K pathway after

treatment with 15d-PGJ$_2$ can be a possible mechanism by which 15d-PGJ$_2$ can

inhibit differentiation of myoblasts. HRas is known to regulate differentiation of

myoblasts in different contexts. Constitutively active HRas signaling by oncogenic

HRas mutant (HRas V12) leads to inhibition of differentiation of myoblasts(Konieczny

et al., 1989; Lassar et al., 1989; Olson,' et al., 1987; Van Der Burgt et al., 2007).

Here we showed that inhibition of differentiation of myoblasts after 15d-PGJ$_2$ is

partially rescued in cells expressing C184S mutants of HRas but not the wild type or

C181S mutant (Fig. 4D and E and S4E). HRas C184S did not get modified by 15d-

PGJ$_2$ (Fig. 3A). These observations suggest that the inhibition of differentiation of

myoblasts by 15d-PGJ$_2$ is partially dependent on modification of HRas by 15d-PGJ$_2$.

Cysteine 181 and 184 in the C-terminal of HRas regulate the intracellular distribution

of HRas between the plasma membrane and the Golgi by reversible palmitoylation

and depalmitoylation(Rocks et al., 2005). Inhibition of palmitoylation of C-terminal

cysteine 181, wither by treatment with protein palmitoylation inhibitor 2-

bromopalmitate or mutation of cysteine to serine, leads to accumulation of HRas at

the Golgi. Intracellular localization of HRas maintains two distinct pools of HRas

activity, where the plasma membrane pool shows a faster activation followed by

short kinetics and the Golgi pool shows a slower activation but a sustained

activation(Agudo-Ibáñez et al., 2015; Busquets-Hernández and Triola, 2021;

Lorentzen et al., 2010; Rocks et al., 2005). We showed that modification of HRas by

15d-PGJ$_2$ alters the intracellular distribution of HRas. We showed that modification of

HRas by 15d-PGJ$_2$ leads to an increase in the localization of the wild type HRas at

the plasma membrane compared to the Golgi (Fig. 4B). We did not observe any

changes in the intracellular distribution of HRas C181S or HRas C184S after

treatment with 15d-PGJ$_2$ (Fig. 4C). HRas C184S is not modified by 15d-PGJ$_2$, but

HRas C181S is modified by 15d-PGJ$_2$ (Fig. 3A). This suggests that intracellular

redistribution of HRas due to modification by 15d-PGJ$_2$ at cysteine 184 requires

palmitoylation of cysteine 181.

Previous reports suggest that downstream signaling of HRas depends on the

intracellular localization of HRas(Rocks et al., 2005; Santra et al., 2019). For

example, targeted localization of HRas at the ER membrane induced expression of

cell-migration genes. Localization of HRas at the plasma membrane showed strong

correlation with expression of cell cycle genes, particularly the MAPK signaling

pathway. Localization of HRas at the plasma membrane also showed negative

correlation with genes associated with the PI3K-Akt pathway. Here we showed that

the intracellular distribution of HRas regulates differentiation of myoblasts. in order to

show this, we used the constitutively active mutant of HRas (HRas V12) which has

been shown to inhibit differentiation of myoblasts(Engler et al., 2021; Konieczny et

al., 1989; Lassar et al., 1989; Olson,' et al., 1987; Scholz et al., 2009; Van Der Burgt

et al., 2007). We expressed cysteine mutants of HRas V12 in C2C12 myoblasts and

found that HRas V12 C181S localized predominantly at the Golgi whereas HRas

V12 and HRas V12 C184S localized at both the plasma membrane and the Golgi

(Fig. S4A). When differentiated, we observed that C2C12 cells expressing HRas V12

C181S differentiated but HRas V12 or HRas V12 C184S did not differentiate (Fig.

S4B, C, and D). These observations suggest alteration of intracellular distribution of

HRas affects HRas mediated inhibition of differentiation of myoblasts.

Doxorubicin (Doxo) is a widely used chemotherapy agent for treatment of

cancers(Johnson-Arbor and Dubey, 2022). Treatment with Doxo induces

senescence. Doxo mediated DNA damage leads to p53, p16, and p21 dependent

senescence in human fibroblasts(Di Leonardo et al., 1994; Robles and Adami,

1998). On the other hand, treatment with doxorubicin leads to decrease in muscle

mass and cross-sectional area, leading to chemotherapy induced cachexia(Hiensch

et al., 2020). Several mechanisms have been proposed behind chemotherapy

induced cachexia, including generation of reactive oxygen species(Gilliam and St.

Clair, 2011), activation of proteases like calpain and caspases(Gilliam et al., 2012;

Smuder et al., 2011), and impaired insulin signaling(de Lima Junior et al., 2016). This

study provides a possible mechanism behind chemotherapy induced loss of muscle

mass and functioning. Induction of senescence in myoblasts by treatment with Doxo

could lead to increased synthesis and release of 15d-PGJ$_2$ by senescent cells which

could be taken up by myoblasts in the microenvironment. The lipid could covalently modify and activates HRas at cysteine 184 to inhibit differentiation of myoblasts. Therefore, targeting the synthesis and release of 15d-PGJ$_2$ by senescent cells could serve as an important target to promote skeletal muscle homeostasis in cancer patients.

Figures

Figure 1. Prostaglandin synthesis and release by Doxo-induced senescent cells inhibits myoblast differentiation.

A. Expression and localization of tumor suppressor protein p21, measured by immunofluorescence, in hindlimb skeletal muscles of mice after 11 days of treatment with Doxo (5 mg/kg) or Saline.

B. Representative confocal micrograph of expression of γH2A.X in the gastrocnemius muscle of mice treated with Doxo (5 mg/kg) or Saline.

C. Expression of mRNAs of senescence markers (p16 and p21), SASP factors (CXCL1, CXCL2, TNF1α, IL6, TGFβ1), and enzymes involved in the biosynthesis of prostaglandin PGD_2/15d-PGJ_2 (PTGS1, PTGS2, PTGDS), measured by qPCR, in hindlimb skeletal muscles of mice after 11 days of treatment with Doxo (5 mg/kg) or Saline.

D. A representative confocal micrograph and a scatter plot of the nuclear area of C2C12 myoblasts, measured by immunofluorescence, after 16 days of treatment with Doxo (150 nM) or DMSO.

E. Expression of mRNAs of prostaglandin biosynthetic enzymes, measured by qPCR, in C2C12 myoblasts after treatment with Doxo (150 nM) or DMSO.

F. Concentration of 15d-PGJ_2 released from quiescent or senescent C2C12 cells.

(Statistical significance was tested by the two-tailed student's t-test ns=$p>0.05$, *=$p<0.05$, **=$p<0.01$, ***=$p<0.001$, ****=$p<0.0001$)

Figure 2. 15d-PGJ$_2$ inhibits differentiation of myoblasts.

A. Expression of MHC protein and the fusion of myoblasts in myotubes, measured by immunofluorescence, after treatment with conditioned medium of senescent cells treated with PTGDS inhibitor AT-56 (30 μM) or DMSO

B. Normalized number of C2C12 myoblasts treated with 15d-PGJ$_2$ (10 μM) or DMSO

C. Expression of mRNAs of markers of differentiation (MyoD, MyoG, and MHC), measured by qPCR, in C2C12 myoblasts treated with 15d-PGJ$_2$ (1 μM, 2 μM, or 4 μM) or DMSO.

D. Expression of MHC protein and the fusion of myoblasts in syncytial myotubes, measured by immunofluorescence, after treatment with 15d-PGJ$_2$ (4 μM) or DMSO.

E. Expression of MHC protein, measured by immunoblotting, in primary human skeletal myoblasts after treatment with 15d-PGJ$_2$ (1 μM, 2 μM, or 4 μM) or DMSO for 5 days.

(Statistical significance was tested by the two-tailed student's t-test ns=$p > 0.05$, *=$p < 0.05$, **=$p < 0.01$, ***=$p < 0.001$, ****=$p < 0.0001$)

Figure 3. 15d-PGJ$_2$ covalently modifies HRas at Cysteine 184 and activates the HRas-MAPK pathway via the electrophilic cyclopentenone ring.

A. Streptavidin-immunoprecipitation of EGFP-HRas, measured by immunoblotting, in C2C12 cells after 3 hours of treatment with 15d-PGJ$_2$-Biotin (5 µM).

B. Representative confocal micrograph of Fluorescence Resonance Energy Transfer (FRET) between EGFP-tagged HRas (EGFP-HRas) and mCherry-tagged Ras binding domain (RBD) of RAF kinase (mCherry-RAF RBD).

C. Activation of the EGFP-tagged wild type HRas (HRas WT), measured by FRET, before and after 1 hour of treatment with 15d-PGJ$_2$ (10 µM) after starvation for 24 hours.

D. Activation of the EGFP-tagged C-terminal cysteine mutants of HRas (HRas C181S and HRas C184S), measured by FRET, before and after 1 hour of treatment with 15d-PGJ$_2$ (10 µM) after starvation for 24 hours.

E. Phosphorylation of Erk (42 kDa and 44 kDa), measured by immunoblotting, in C2C12 cells after 1 hour of treatment with 15d-PGJ$_2$ (5 µM, 10 µM) or DMSO after starvation for 24 hours.

F. Phosphorylation of Erk (42 kDa and 44 kDa), measured by immunoblotting, in C2C12 cells after 1 hour of treatment with 15d-PGJ$_2$ (10 µM)/ 9,10-dihydro-15d-PGJ$_2$ (10 µM) or DMSO after starvation for 24 hours.

(Statistical significance was tested by the two-tailed student's t-test ns=$p>0.05$, *=$p<0.05$, **=$p<0.01$, ***=$p<0.001$, ****=$p<0.0001$)

$$R_{mean} = \frac{mean\ HRas\ at\ Golgi}{mean\ HRas\ at\ Plasma\ Membrane}$$

Figure 4. 15d-PGJ$_2$ controls the intracellular distribution of HRas and differentiation of C2C12 cells in an HRas C-terminal cysteine-dependent manner.

A. Representative confocal micrograph of C2C12 myoblasts showing localization of EGFP-tagged HRas between the plasma membrane (stained with Alexa Fluor 633 conjugated Wheat Germ Agglutinin) and the Golgi (labelled with TagRFP-tagged Golgi resident GalT protein). A statistic R$_{mean}$ was defined as

the ratio of mean HRas intensity at the Golgi to the mean HRas intensity at the plasma membrane.

B. Distribution of the wild-type HRas between the Golgi and the plasma membrane, measured by R_{mean}, in C2C12 myoblasts treated with 15d-PGJ$_2$ (10 µM) or DMSO for 24 hours.

C. Distribution of the C-terminal cysteine mutants of HRas between the Golgi and the plasma membrane, measured by R_{mean}, in C2C12 myoblasts treated with 15d-PGJ$_2$ (10 µM) or DMSO for 24 hours.

D. Expression of mRNAs of known markers of differentiation (MyoD, MyoG, and MHC), measured by qPCR, in differentiating C2C12 myoblasts expressing the EGFP-tagged wild-type and the C-terminal cysteine mutants of HRas after treatment with 15d-PGJ$_2$ (4 µM) or DMSO for 5 days.

E. Expression of MHC protein, measured by immunoblotting, in differentiating C2C12 myoblasts expressing the EGFP-tagged wild-type and the C-terminal cysteine mutants of HRas after treatment with 15d-PGJ$_2$ (4 µM) or DMSO for 5 days.

(Statistical significance was tested by the two-tailed student's t-test ns=$p>0.05$, *=$p<0.05$, **=$p<0.01$, ***=$p<0.001$, ****=$p<0.0001$)

Acknowledgements

We thank Prof. Satyajit Mayor (NCBS), Prof. Phillipe Bastiens, and Prof. Apurva Sarin (InStem) for providing the wid-type HRas construct, the mCherry-RAF RBD construct and the vector backbones respectively. We thank Dr. Neetu Saini (InStem) for her help with setting up the cell culture facility. We thank Mr. Heera Lal for his help with the animal work. We thank Dr. Kamlesh Kumar Yadav and Ms. Sudeshna Saha for their help during the project. We thank the Central Imaging and Flow Cytometry Facility (CIFF) (NCBS-InStem) for their support with microscopy. We thank the Animal Care and Resource Centre (ACRC) (NCBS-InStem) for their support with mouse experiments. We thank the Mass Spectrometry facility (NCBS-InStem) for their support with the mass spectrometry work.

Funding

This work was supported by SERB SUPRA grant to Dr. Arvind Ramanathan. SSP and AB are supported by GS program (InStem), AV is supported by DBT-JRF grant.

Author Contribution

SSP: Project conceptualization, Cell culture treatments and assays, Biochemistry, Microscopy, Image Processing, and analysis, Writing: original draft, review, and edits.

SSS: Cell culture treatments and assays, Biochemistry, Writing: review and edits.

AB: Animal work, Cell culture treatments and assays, Biochemistry, Mass Spectrometry, Writing: review and edits.

AV: Cell culture treatments and assays, Image processing and analysis, writing: review and edits.

MAJ: Image processing and analysis.

RGHM: Mass Spectrometry.

AR: Project Conceptualization, Writing: original draft, review, and edits. Corresponding Author.

Competing Interests

The authors declare no competing interests.

References

Agudo-Ibáñez L, Herrero A, Barbacid M, Crespo P. 2015. H-Ras Distribution and Signaling in Plasma Membrane Microdomains Are Regulated by Acylation and Deacylation Events. *Mol Cell Biol* **35**:1898–1914. doi:10.1128/mcb.01398-14

Bennett AM, Tonks NK. 1997. Regulation of distinct stages of skeletal muscle differentiation by mitogen-activated protein kinases. *Science (80-)* **278**:1288–1291. doi:10.1126/SCIENCE.278.5341.1288

Bihani T, Chicas A, Lo CPK, Lin AW. 2007. Dissecting the senescence-like program in tumor cells activated by ras signaling. *J Biol Chem* **282**:2666–2675. doi:10.1074/jbc.M608127200

Bihani T, Mason DX, Jackson TJ, Chen SC, Boettner B, Lin AW. 2004. Differential Oncogenic Ras Signaling and Senescence in Tumor Cells. *Cell Cycle* **3**:1201–1207.

Bittles AH, Harper N. 1984. Increased glycolysis in ageing cultured human diploid fibroblasts. *Biosci Rep* **4**:751–756. doi:10.1007/BF01128816

BOHR Orthopredic Hospital H, Knudsen H, Kat J. 1964. An Unidentified Virus which causes the Rapid Production of Tumours in Mice. *Nat 1964 2044963* **204**:1104–1105. doi:10.1038/2041104b0

Busquets-Hernández C, Triola G. 2021. Palmitoylation as a Key Regulator of Ras Localization and Function. *Front Mol Biosci* **8**:1–8. doi:10.3389/fmolb.2021.659861

Calabrese V, Mallette FA, Deschênes-Simard X, Ramanathan S, Gagnon J, Moores A, Ilangumaran S, Ferbeyre G. 2009. SOCS1 Links Cytokine Signaling to p53 and Senescence. *Mol Cell* **36**:754–767. doi:10.1016/j.molcel.2009.09.044

Campisi J. 2005. Senescent cells, tumor suppression, and organismal aging: Good citizens, bad neighbors. *Cell* **120**:513–522. doi:10.1016/j.cell.2005.02.003

Casar B, Badrock AP, Jiménez I, Arozarena I, Colón-bolea P, Lorenzo-martín LF, Barinaga-rementería I, Barriuso J, Cappitelli V, Donoghue DJ, Bustelo XR, Hurlstone A, Crespo P. 2018. RAS at the Golgi antagonizes malignant transformation through PTPR κ -mediated inhibition of ERK activation. *Nat Commun* 1–17. doi:10.1038/s41467-018-05941-8

Chen Q, Ames BN. 1994. Senescence-like growth arrest induced by hydrogen peroxide in human diploid fibroblast F65 cells. *Proc Natl Acad Sci U S A* **91**:4130–4134. doi:10.1073/pnas.91.10.4130

Chen QM, Bartholomew JC, Campisi J, Acosta M, Reagan JD, Ames BN. 1998. Molecular analysis of H2O2-induced senescent-like growth arrest in normal human fibroblasts: p53 and Rb control G1 arrest but not cell replication. *Biochem J* **332**:43–50. doi:10.1042/bj3320043

Chen YX, Zhong XY, Qin YF, Bing W, He LZ. 2003. 15d-PGJ2 inhibits cell growth and induces apoptosis of MCG-803 human gastric cancer cell line. *World J Gastroenterol* **9**:2149. doi:10.3748/WJG.V9.I10.2149

Childs BG, Durik M, Baker DJ, van Deursen JM. 2015. Cellular senescence in aging and age-related disease: from mechanisms to therapy disease: from

mechanisms to therapy. doi:10.1038/nm.4000

Choi J, Suh J-Y, Kim D-H, Na H-K, Surh Y-J. 2020. 15-Deoxy-Δ12,14-prostaglandin
J2 Induces Epithelial-tomesenchymal Transition in Human Breast Cancer Cells
and Promotes Fibroblast Activation. *J Cancer Prev* **25**:152–163.
doi:10.15430/JCP.2020.25.3.152

Conte TC, Duran-Bishop G, Orfi Z, Mokhtari I, Deprez A, Côté I, Molina T, Kim TY,
Tellier L, Roussel MP, Maggiorani D, Benabdallah B, Leclerc S, Feulner L,
Pellerito O, Mathieu J, Andelfinger G, Gagnon C, Beauséjour C, McGraw S,
Duchesne E, Dumont NA. 2023. Clearance of defective muscle stem cells by
senolytics restores myogenesis in myotonic dystrophy type 1. *Nat Commun*
2023 141 **14**:1–17. doi:10.1038/s41467-023-39663-3

Coppé JP, Desprez PY, Krtolica A, Campisi J. 2010. The senescence-associated
secretory phenotype: The dark side of tumor suppression. *Annu Rev Pathol*
Mech Dis **5**:99–118. doi:10.1146/annurev-pathol-121808-102144

Coppé JP, Patil CK, Rodier F, Sun Y, Muñoz DP, Goldstein J, Nelson PS, Desprez
PY, Campisi J. 2008. Senescence-associated secretory phenotypes reveal cell-
nonautonomous functions of oncogenic RAS and the p53 tumor suppressor.
PLoS Biol **6**. doi:10.1371/journal.pbio.0060301

Cosgrove BD, Gilbert PM, Porpiglia E, Mourkioti F, Lee SP, Corbel SY, Llewellyn
ME, Delp SL, Blau HM. 2014. Rejuvenation of the muscle stem cell population
restores strength to injured aged muscles. *Nat Med 2014 203* **20**:255–264.
doi:10.1038/nm.3464

D'Adda Di Fagagna F. 2008. Living on a break: Cellular senescence as a DNA-
damage response. *Nat Rev Cancer* **8**:512–522. doi:10.1038/nrc2440

D'Adda Di Fagagna F, Reaper PM, Clay-Farrace L, Fiegler H, Carr P, Von Zglinicki
T, Saretzki G, Carter NP, Jackson SP. 2003. A DNA damage checkpoint
response in telomere-initiated senescence. *Nature* **426**:194–198.
doi:10.1038/nature02118

Davalos AR, Coppe JP, Campisi J, Desprez PY. 2010. Senescent cells as a source
of inflammatory factors for tumor progression. *Cancer Metastasis Rev* **29**:273–

283. doi:10.1007/s10555-010-9220-9

Davis M, Malcolm S, Hall A, Marshall CJ. 1983. Localisation of the human N-ras oncogene to chromosome 1cen - p21 by in situ hybridisation. *EMBO J* **2**:2281–2283. doi:10.1002/J.1460-2075.1983.TB01735.X

de Lima Junior EA, Yamashita AS, Pimentel GD, De Sousa LGO, Santos RVT, Gonçalves CL, Streck EL, de Lira FS, Rosa Neto JC. 2016. Doxorubicin caused severe hyperglycaemia and insulin resistance, mediated by inhibition in AMPk signalling in skeletal muscle. *J Cachexia Sarcopenia Muscle* **7**:615–625. doi:10.1002/JCSM.12104

Di Leonardo A, Linke SP, Clarkin K, Wahl GM. 1994. DNA damage triggers a prolonged p53-dependent G1 arrest and long-term induction of Cip1 in normal human fibroblasts. *Genes Dev* **8**:2540–2551. doi:10.1101/gad.8.21.2540

Dilley TK, Bowden GT, Chen QM. 2003. Novel mechanisms of sublethal oxidant toxicity: Induction of premature senescence in human fibroblasts confers tumor promoter activity. *Exp Cell Res* **290**:38–48. doi:10.1016/S0014-4827(03)00308-2

Dimri GP, Leet X, Basile G, Acosta M, Scorrt G, Roskelley C, Medrano EE, Linskensi M, Rubeljii I, Pereira-Smithii O, Peacocket M, Campisi J, Pardee B. 1995. A biomarker that identifies senescent human cells in culture and in aging skin in vivo (replicative senescence/tumor suppression/18-galactosidase) Communicated by Arthur. *Cell Bioiogy* **92**:9363–9367.

Engler M, Fidan M, Nandi S, Cirstea IC. 2021. Senescence in RASopathies, a possible novel contributor to a complex pathophenoype. *Mech Ageing Dev*. doi:10.1016/j.mad.2020.111411

Francis TG, Jaka O, Ellison-Hughes GM, Lazarus NR, Harridge SDR. 2022. Human primary skeletal muscle-derived myoblasts and fibroblasts reveal different senescent phenotypes. *JCSM Rapid Commun* **5**:226–238. doi:10.1002/RCO2.67

Franza BR, Maruyama K, Garrels JI, Ruley HE. 1986. In vitro establishment is not a sufficient prerequisite for transformation by activated ras oncogenes. *Cell*

44:409–418. doi:10.1016/0092-8674(86)90462-9

Gilliam LAA, Moylan JS, Patterson EW, Smith JD, Wilson AS, Rabbani Z, Reid MB. 2012. Doxorubicin acts via mitochondrial ROS to stimulate catabolism in C2C12 myotubes. *Am J Physiol - Cell Physiol* **302**:195–202. doi:10.1152/AJPCELL.00217.2011/ASSET/IMAGES/LARGE/ZH00121167990008.JPEG

Gilliam LAA, St. Clair DK. 2011. Chemotherapy-Induced Weakness and Fatigue in Skeletal Muscle: The Role of Oxidative Stress. *https://home.liebertpub.com/ars* **15**:2543–2563. doi:10.1089/ARS.2011.3965

Gutierrez L, Magee AI, Marshall CJ, Hancock JF. 1989. Post-translational processing of p21ras is two-step and involves carboxyl-methylation and carboxy-terminal proteolysis. *EMBO J* **8**:1093–1098. doi:10.1002/J.1460-2075.1989.TB03478.X

Hachet-Haas M, Converset N, Marchal O, Matthes H, Gioria S, Galzi J-L, Lecat S. 2006. FRET and colocalization analyzer—A method to validate measurements of sensitized emission FRET acquired by confocal microscopy and available as an ImageJ Plug-in. *Microsc Res Tech* **69**:941–956. doi:10.1002/jemt.20376

Hamsanathan S, Gurkar AU. 2022. Lipids as Regulators of Cellular Senescence. *Front Physiol* **13**:796850. doi:10.3389/FPHYS.2022.796850/BIBTEX

Hayflick L. 1965. The limited in vitro lifetime of human diploid cell strains. *Exp Cell Res* **37**:614–636. doi:10.1016/0014-4827(65)90211-9

Hiensch AE, Bolam KA, Mijwel S, Jeneson JAL, Huitema ADR, Kranenburg O, van der Wall E, Rundqvist H, Wengstrom Y, May AM. 2020. Doxorubicin-induced skeletal muscle atrophy: Elucidating the underlying molecular pathways. *Acta Physiol* **229**:e13400. doi:10.1111/APHA.13400

Hu S., Liu B, Shang J, Lu T, Cai Y, Ding M, Zhou X, Wang X. 2023. PTGDS PROMOTES TUMORIGENESIS OF PERIPHERAL T CELL LYMPHOMA THROUGH REGULATING IRON METABOLISM. *Hematol Oncol* **41**:625–625. doi:10.1002/HON.3165_475

Hu Shunfeng, Liu B, Shang J, Lu T, Ding M, Zhou X, Wang X. 2023. P1210:

TARGETING PTGDS AS A NOVEL THERAPEUTIC APPROACH IN
PERIPHERAL T CELL LYMPHOMA THROUGH REGULATING IRON
METABOLISM. *HemaSphere* **7**:e6034196.
doi:10.1097/01.HS9.0000971736.60341.96

Hu S, Ren S, Cai Y, Liu J, Han Y, Zhao Y, Yang J, Zhou X, Wang X. 2021.
Glycoprotein PTGDS promotes tumorigenesis of diffuse large B-cell lymphoma
by MYH9-mediated regulation of Wnt–β-catenin–STAT3 signaling. *Cell Death*
Differ 2021 293 **29**:642–656. doi:10.1038/s41418-021-00880-2

Hu X, Zhang H. 2019. Doxorubicin-Induced Cancer Cell Senescence Shows a Time
Delay Effect and Is Inhibited by Epithelial-Mesenchymal Transition (EMT). *Med*
Sci Monit **25**:3617–3623. doi:10.12659/MSM.914295

Hunter JG, Van Delft MF, Rachubinski RA, Capone JP. 2001. Peroxisome
Proliferator-activated Receptor γ Ligands Differentially Modulate Muscle Cell
Differentiation and MyoD Gene Expression via Peroxisome Proliferator-activated
Receptor γ-dependent and -independent Pathways. *J Biol Chem* **276**:38297–
38306. doi:10.1074/JBC.M103594200

Irikura D, Aritake K, Nagata N, Maruyama T, Shimamoto S, Urade Y. 2009.
Biochemical, Functional, and Pharmacological Characterization of AT-56, an
Orally Active and Selective Inhibitor of Lipocalin-type Prostaglandin D Synthase.
J Biol Chem **284**:7623–7630. doi:10.1074/JBC.M808593200

Johnson-Arbor K, Dubey R. 2022. Doxorubicin. *xPharm Compr Pharmacol Ref* 1–5.
doi:10.1016/B978-008055232-3.61650-2

Jones KR, Elmore LW, Jackson-Cook C, Demasters G, Povirk LF, Holt SE, Gewirtz
DA. 2005. p53-dependent accelerated senescence induced by ionizing radiation
in breast tumour cells. *Int J Radiat Biol* **81**:445–458.
doi:10.1080/09553000500168549

Kirsten WH, Mayer LA. 1967. Morphologic Responses to a Murine Erythroblastosis
Virus. *JNCI J Natl Cancer Inst* **39**:311–335. doi:10.1093/JNCI/39.2.311

Konieczny SF, Drobes BL, Menke SL, Taparowsky EJ. 1989. Inhibition of myogenic
differentiation by the H-ras oncogene is associated with the down regulation of

the MyoD1 gene. *Oncogene* **4**:473–481.

Krtolica A, Parrinello S, Lockett S, Desprez PY, Campisi J. 2001. Senescent fibroblasts promote epithelial cell growth and tumorigenesis: A link between cancer and aging. *Proc Natl Acad Sci U S A* **98**:12072–12077. doi:10.1073/pnas.211053698

Land H, Parada LF, Weinberg RA. 1983. Tumorigenic conversion of primary embryo fibroblasts requires at least two cooperating oncogenes. *Nature* **304**:596–602. doi:10.1038/304596a0

Lassar AB, Thayer MJ, Overell RW, Weintraub H. 1989. Transformation by activated ras or fos prevents myogenesis by inhibiting expression of MyoD1. *Cell* **58**:659–667. doi:10.1016/0092-8674(89)90101-3

Lee BY, Han JA, Im JS, Morrone A, Johung K, Goodwin EC, Kleijer WJ, DiMaio D, Hwang ES. 2006. Senescence-associated β-galactosidase is lysosomal β-galactosidase. *Aging Cell* **5**:187–195. doi:10.1111/j.1474-9726.2006.00199.x

Lee J, Choi KJ, Lim MJ, Hong F, Choi TG, Tak E, Lee S, Kim YJ, Chang SG, Cho JM, Ha J, Kim SS. 2010. Proto-oncogenic H-Ras, K-Ras, and N-Ras are involved in muscle differentiation via phosphatidylinositol 3-kinase. *Cell Res* **20**:919–934. doi:10.1038/cr.2010.92

Li J, Guo C, Wu J. 2019. 15-Deoxy-Δ-12,14-Prostaglandin J2 (15d-PGJ2), an Endogenous Ligand of PPAR-γ: Function and Mechanism. *PPAR Res* **2019**. doi:10.1155/2019/7242030

Lorentzen A, Kinkhabwala A, Rocks O, Vartak N, Bastiaens PIH. 2010. Regulation of Ras localization by acylation enables a mode of intracellular signal propagation. *Sci Signal* **3**. doi:10.1126/SCISIGNAL.20001370/SUPPL_FILE/3_RA68_SM.PDF

Lowe SW, Cepero E, Evan G. 2004. Intrinsic tumour suppression. *Nature* **432**:307–315. doi:10.1038/nature03098

Lu JY, Hofmann SL. 1995. Depalmitoylation of CAAX motif proteins. Protein structural determinants of palmitate turnover rate. *J Biol Chem* **270**:7251–7256. doi:10.1074/JBC.270.13.7251/ATTACHMENT/56A70B86-B9FE-4517-9BAB-

74A5749981A0/TBL1

Luis Oliva J, Pé Rez-Sala D, Castrillo A, Martínez N, Cañ Ada ‡ FJ, Boscá L, Rojas
JM. 2003. The cyclopentenone 15-deoxy-12,14-prostaglandin J 2 binds to and
activates H-Ras.

Marcone S, Fitzgerald DJ. 2013. Proteomic identification of the candidate target
proteins of 15-deoxy-delta12,14-prostaglandin J2. *Proteomics* **13**:2135–2139.
doi:10.1002/pmic.201200289

Morgenstern J, Fleming T, Kadiyska I, Brings S, Groener JB, Nawroth P, Hecker M,
Brune M. 2018. Sensitive mass spectrometric assay for determination of 15-
deoxy-Δ12,14-prostaglandin J2 and its application in human plasma samples of
patients with diabetes. *Anal Bioanal Chem* **410**:521–528. doi:10.1007/s00216-
017-0748-1

Olson,' EN, Spizz G, Tainsky2 MA. 1987. The oncogenic forms of N-ras or H-ras
prevent skeletal myoblast differentiation. *Mol Cell Biol* **7**:2104–2111.
doi:10.1128/MCB.7.6.2104-2111.1987

Parrinello S, Coppe JP, Krtolica A, Campisi J. 2005. Stromal-epithelial interactions in
aging and cancer: Senescent fibroblasts alter epithelial cell differentiation. *J Cell
Sci* **118**:485–496. doi:10.1242/jcs.01635

Pylayeva-Gupta Y, Grabocka E, Bar-Sagi D. 2011. RAS oncogenes: weaving a
tumorigenic web. *Nat Rev Cancer 2011 1111* **11**:761–774. doi:10.1038/nrc3106

Robles SJ, Adami GR. 1998. Agents that cause DNA double strand breaks lead to
p16(INK4a) enrichment and the premature senescence of normal fibroblasts.
Oncogene **16**:1113–1123. doi:10.1038/sj.onc.1201862

Rocks O, Peyker A, Kahms M, Verveer PJ, Koerner C, Lumbierres M, Kuhlmann J,
Waldmann H, Wittinghofer A, Bastiaens PIH. 2005. An acylation cycle regulates
localization and activity of palmitoylated ras isoforms. *Science (80-)* **307**:1746–
1752. doi:10.1126/science.1105654

Rommel C, Clarke BA, Zimmermann S, Nuñez L, Rossman R, Reid K, Moelling K,
Yancopoulos GD, Glass DJ. 1999. Differentiation stage-specific inhibition of the
Raf-MEK-ERK pathway by Akt. *Science (80-)* **286**:1738–1741.

doi:10.1126/SCIENCE.286.5445.1738

Santra T, Herrero A, Rodriguez J, von Kriegsheim A, Iglesias-Martinez LF, Schwarzl

T, Higgins D, Aye TT, Heck AJR, Calvo F, Agudo-Ibáñez L, Crespo P,

Matallanas D, Kolch W. 2019. An Integrated Global Analysis of

Compartmentalized HRAS Signaling. *Cell Rep* **26**:3100-3115.e7.

doi:10.1016/J.CELREP.2019.02.038

Scholz ME, Meissner JD, Scheibe RJ, Umeda PK, Chang KC, Gros G, Kubis HP.

2009. Different roles of H-ras for regulation of myosin heavy chain promoters in

satellite cell-derived muscle cell culture during proliferation and differentiation.

Am J Physiol - Cell Physiol **297**:1012–1018.

doi:10.1152/AJPCELL.00567.2008/ASSET/IMAGES/LARGE/ZH001009605700

06.JPEG

Serrano M, Lin AW, McCurrach ME, Beach D, Lowe SW. 1997. Oncogenic ras

provokes premature cell senescence associated with accumulation of p53 and

p16(INK4a). *Cell.* doi:10.1016/S0092-8674(00)81902-9

Shaner NC, Lin MZ, McKeown MR, Steinbach PA, Hazelwood KL, Davidson MW,

Tsien RY. 2008. Improving the photostability of bright monomeric orange and

red fluorescent proteins. *Nat Methods 2008 56* **5**:545–551.

doi:10.1038/nmeth.1209

Shelton DN, Chang E, Whittier PS, Choi D, Funk WD. 1999. Microarray analysis of

replicative senescence. *Curr Biol* **9**:939–945. doi:10.1016/S0960-

9822(99)80420-5

Shibata T, Kondo M, Osawa T, Shibata N, Kobayashi M, Uchida K. 2002. 15-Deoxy-

Δ12,14-prostaglandin J2. *J Biol Chem* **277**:10459–10466.

doi:10.1074/jbc.m110314200

Slanovc J, Mikulčić M, Jahn N, Wizsy NGT, Sattler W, Malle E, Hrzenjak A. 2024.

Prostaglandin 15d-PGJ2 inhibits proliferation of lung adenocarcinoma cells by

inducing ROS production and activation of apoptosis via sirtuin-1. *Biochim*

Biophys acta Mol basis Dis **1870**. doi:10.1016/J.BBADIS.2023.166924

Smuder AJ, Kavazis AN, Min K, Powers SK. 2011. Exercise protects against

doxorubicin-induced oxidative stress and proteolysis in skeletal muscle. *J Appl Physiol* **110**:935–942. doi:10.1152/JAPPLPHYSIOL.00677.2010/ASSET/IMAGES/LARGE/ZDG004119 5300006.JPEG

Stein GH, Beeson M, Gordon L. 1990. Failure to phosphorylate the retinoblastoma gene product in senescent human fibroblasts. *Science (80-)* **249**:666–669. doi:10.1126/science.2166342

Van Der Burgt I, Kupsky W, Stassou S, Nadroo A, Barroso C, Diem A, Kratz CP, Dvorsky R, Ahmadian MR, Zenker M. 2007. Myopathy caused by HRAS germline mutations: implications for disturbed myogenic differentiation in the presence of constitutive HRas activation. *J Med Genet* **44**:459–462. doi:10.1136/JMG.2007.049270

Veliça P, Khanim FL, Bunce CM. 2010. Prostaglandin D2 inhibits C2C12 myogenesis. *Mol Cell Endocrinol* **319**:71–78. doi:10.1016/J.MCE.2010.01.023

Vetter IR, Wittinghofer A. 2001. The guanine nucleotide-binding switch in three dimensions. *Science (80-)* **294**:1299–1304. doi:10.1126/SCIENCE.1062023/ASSET/0AB072BE-F662-4796-8B33- AFF3C6AD829F/ASSETS/GRAPHIC/SE4419929006.JPEG

Wiley CD, Campisi J. 2021. The metabolic roots of senescence: mechanisms and opportunities for intervention. *Nat Metab 2021 310* **3**:1290–1301. doi:10.1038/s42255-021-00483-8

Wiley CD, Campisi J. 2016. From Ancient Pathways to Aging Cells - Connecting Metabolism and Cellular Senescence. *Cell Metab*. doi:10.1016/j.cmet.2016.05.010

Wiley CD, Sharma R, Davis SS, Lopez-Dominguez JA, Mitchell KP, Wiley S, Alimirah F, Kim DE, Payne T, Rosko A, Aimontche E, Deshpande SM, Neri F, Kuehnemann C, Demaria M, Ramanathan A, Campisi J. 2021. Oxylipin biosynthesis reinforces cellular senescence and allows detection of senolysis. *Cell Metab* **33**:1124-1136.e5. doi:10.1016/j.cmet.2021.03.008

Wiley CD, Velarde MC, Lecot P, Liu S, Sarnoski EA, Freund A, Shirakawa K, Lim

HW, Davis SS, Ramanathan A, Gerencser AA, Verdin E, Campisi J. 2016.
Mitochondrial dysfunction induces senescence with a distinct secretory
phenotype. *Cell Metab* **23**:303–314. doi:10.1016/j.cmet.2015.11.011

Woods D, Parry D, Cherwinski H, Bosch E, Lees E, Mcmahon M. 1997. Raf-Induced
Proliferation or Cell Cycle Arrest Is Determined by the Level of Raf Activity with
Arrest Mediated by p21 Cip1. *Mol Cell Biol* **17**:5598–5611.

Yamamoto Y, Takase K, Kishino J, Fujita M, Okamura N, Sakaeda T, Fujimoto M,
Yagami T. 2011. Proteomic Identification of Protein Targets for 15-Deoxy-
Δ12,14-Prostaglandin J2 in Neuronal Plasma Membrane. *PLoS One* **6**:e17552.
doi:10.1371/JOURNAL.PONE.0017552

Yang G, Rosen DG, Zhang Z, Bast RC, Mills GB, Colacino JA, Mercado-Uribe I, Liu
J. 2006. The chemokine growth-regulated oncogene 1 (Gro-1) links RAS
signaling to the senescence of stromal fibroblasts and ovarian tumorigenesis.
Proc Natl Acad Sci U S A **103**:16472–16477. doi:10.1073/pnas.0605752103

Yen CC, Hsiao C Der, Chen WM, Wen YS, Lin YC, Chang TW, Yao FY, Hung SC,
Wang JY, Chiu JH, Wang HW, Lin CH, Chen TH, Chen PCH, Liu CL, Tzeng CH,
Fletcher JA. 2014. Cytotoxic effects of 15d-PGJ2 against osteosarcoma through
ROS-mediated AKT and cell cycle inhibition. *Oncotarget* **5**:716.
doi:10.18632/ONCOTARGET.1704

Zindy F, Williams RT, Baudino TA, Rehg JE, Skapek SX, Cleveland JL, Roussel MF,
Sherr CJ. 2003. Arf tumor suppressor promoter monitors latent oncogenic
signals in vivo. *Proc Natl Acad Sci U S A* **100**:15930–15935.
doi:10.1073/pnas.2536808100

Zwerschke W, Mazurek S, Stöckl P, Hütter E, Eigenbrodt E, Jansen-Dürr P. 2003.
Metabolic analysis of senescent human fibroblasts reveals a role for AMP in
cellular senescence. *Biochem J* **376**:403–411. doi:10.1042/BJ20030816

Materials and Methods

Materials and Methods:

- **Plasmids:**

Unmutated and cysteine mutants of HRas WT [HRas WT, HRas-C181S, and HRas-C184S] and HRas V12 [HRas V12, HRas V12-C181S, HRas V12-C184S] were cloned in the pEGFPC1 vector (Clontech) by restriction digestion-ligation method. Constructs of wild-type HRas were PCR amplified from a previously available HRas construct in the lab with construct-specific primers using Phusion High Fidelity DNA Polymerase (Thermo Scientific) followed by restriction digestion of HRas constructs and the pEGFPC1 vector using Xho1/EcoR1 enzymes (New England Biolabs Inc.). Ligation was set up using T4 DNA ligase (Takara Bio.). proper nucleotide additions were made to the forward primer to maintain the EGFP ORF, marking a 7 amino acid linker between the proteins. The construct sequences were confirmed by Sanger sequencing. GalT – TagRFP construct was a gift from Prof. Satyajit Mayor and was used to mark the Golgi. mCherry-RAF RBD construct was a gift from Prof. Phillipe Bastiens and was used to measure the activity of HRas GTPase using FRET.

- **Cell Culture:**
    - **Cell Maintenance:**

C2C12 mouse myoblasts (CRL – 1772) were obtained from ATCC and were maintained in DMEM complete medium @ 37° C, 5% CO_2. For experiments, the cells were trypsinized with 0.25% trypsin – EDTA (Gibco): DPBS (Gibco) (1:1) @ 37° C, 3 mins. The cells were then resuspended in a triple volume of DMEM complete medium and were counted manually with a hemocytometer after diluting the cells with trypan blue (Gibco) (1:1) and were seeded in required numbers in cell culture dishes. Human Skeletal Muscle Myoblasts P2 (CC-2580) were obtained from Lonza and were maintained in DMEM Skeletal Muscle growth medium @ 37° C, 5% CO_2. For experiments, the cells were trypsinized with 0.25% trypsin – EDTA (Gibco): DPBS (Gibco) (1:1) @ 37° C, 2 mins. The cells were then resuspended in a double volume of DMEM complete medium and were counted manually with a hemocytometer after diluting the cells with trypan blue (Gibco) (1:1) and were seeded in required numbers in cell culture dishes. All cultures tested negative for mycoplasma checked by Mycoalert Mycoplasma Detection Kit (Lonza).

    - **Conditioned medium collection:**

C2C12 cells seeded in 35mm dishes were treated with Doxorubicin (150 nM) for 3 days. The media was then changed to DMEM complete medium without Doxorubicin

for 19 days after treatment with Doxorubicin. The cells were treated with DMSO or AT-56 (30 µM) in the DMEM complete medium for 2 days. On Day 21, the cells were treated with DMSO or At-56 in DMEM Starvation medium for 3 days. The media was then collected, centrifuged @1000g, R.T. for 5 minutes. The media was then stored at -80° C after flashfreezing in liq. N_2 till further requirement.

- **Treatments:**
  - **15d-PGJ$_2$:**

15d-PGJ$_2$ (Cayman Chemical Company) dissolved in methyl acetate was purged with N_2 stream @ R.T. till drying and was then redissolved in DMSO to make a 10 mM stock solution. 15d-PGJ$_2$ (10 mM) stock was then appropriately diluted in an appropriate volume of DMEM media for experiments. DMSO was used as vehicle control. A media change of the same composition was given every 24 hours.

  - **9,10-dihydro-15d-PGJ$_2$:**

9,10-dihydro-15d-PGJ$_2$ (Cayman Chemical Company) dissolved in methyl acetate was purged with N_2 stream @ R.T. till drying and was then redissolved in DMSO to make a 10 mM stock solution. 9,10-dihydro-15d-PGJ$_2$ (10 mM) stock was then appropriately diluted in an appropriate volume of DMEM media for experiments. DMSO was used as a control.

  - **Doxorubicin:**

C2C12 cells with 70-80% confluency in a 10 cm plate were treated with Doxorubicin (Doxo) for 3 days. After 72hr, Doxo was removed from the medium and the cells were kept for 10 more days with media change every 3 days. The cells were harvested on day 13 after Doxo treatment for experiments.

  - **15d-PGJ$_2$-Biotin:**

C2C12 cells transfected with EGFP-HRas WT/C181S/C184S in 35mm dishes were treated with 15d-PGJ$_2$-Biotin in DMEM Hi Glucose medium (Gibco) supplemented with 1% Penicilin-streptomycin-Glutamine (Gibco) without fetal bovine serum for 3 hours. The cells were then harvested for immunoprecipitation.

  - **Conditioned medium:**

Conditioned medium collected from senescent cells was thawed @ 37° C. The medium was then supplemented with 2% heat-inactivated horse serum and 1% penicillin-streptavidin-glutamine. C2C12 myoblasts seeded in 35mm dishes on coverslips were treated with the conditioned medium. A media change was given every 48 hours.

- **Transfections:**

C2C12 cells were seeded in 35 mm dishes at an intermediate density to achieve confluency of ~60 - 70%. For western blot, immunoprecipitation, and differentiation experiments, the cells were transfected with 1.5 µg of EGFP-tagged HRas WT/ HRas-C181S/ HRas-C184S/ HRas V12/ HRas V12-C181S/ HRas V12-C184S using the jetPRIME transfection reagent (Polyplus) using the manufacturer's protocol. Transfection efficiency was confirmed by checking for GFP fluorescence in cells under a Ti2 epifluorescence microscope (Nikon) using appropriate filters. For measuring the intracellular distribution of HRas, the cells were reverse transfected with 1µg each of EGFP-tagged HRas WT/ HRas-C181S/ HRas-C184S/ HRas V12/ HRas V12-C181S/ HRas V12-C184S and GalT - TagRFP, a Golgi apparatus marker protein tagged with red fluorescent TagRFP protein using jetPRIME, where the DNA - jetPRIME mixture was incubated with the cell suspension while seeding the cells after trypsinization, and the transfection efficiency was confirmed by checking GFP and RFP fluorescence after 24 hours of transfection. For measuring the activity of HRas, the cells seeded in imaging dishes (iBidi) were transfected with 0.7 µg EGFP-HRas and 1.4 µg mCherry-RAF RBD using jetPRIME transfection reagent (Polyplus) using the manufacturer's protocol. Transfection efficiency was confirmed by checking for GFP and RFP fluorescence after 24 hours of transfection.

- **Myoblast differentiation:**
  - ➤ **Untransfected cells:**

C2C12 cells were seeded in 35 mm dishes (Corning) at high density to achieve a confluency of ~90 – 95% the next day. The cells were then treated with either 15d-PGJ$_2$ or DMSO in the C2C12 differentiation medium. The cells were given a media change of the same composition every 24 hours. The cells were harvested after 5 days of 15d-PGJ$_2$ treatment for either RNA or protein isolation. For the Immunofluorescence experiment, the experiment was done in 35 mm dishes

(Corning) on glass coverslips (Blue star) coated with 0.2% Gelatin (Porcine, Sigma
Aldrich), and the cells were fixed with the fixative solution at the end of the
experiment and were immunostained. Human Skeletal Muscle Myoblast cells were
seeded in 35 mm dishes in high density to achieve a confluency of ~90 – 95% the
next day. The cells were then treated with DMSO or 15d – PGJ_2 in the Skeletal
Muscle Differentiation medium. A media change of the same composition was given
every 24 hours. The cells were harvested after 5 days of treatment in RIPA - PP.

➤ **Transfected cells:**

C2C12 cells were seeded in 35 mm dishes (Corning) at an intermediate density to
achieve a confluency of ~60 – 70% confluency. The cells were transfected with
EGFP-tagged HRas WT/ HRas-C181S/ HRas-C184S/HRas V12/HRas V12-C181S/
HRas V12-C184S using jetPRIME transfection reagent (Polyplus). After confirming a
transfection efficiency of ~80% the next day, the cells were treated with DMSO
vehicle or 15d – PGJ_2 (4 μM) in the C2C12 differentiation medium. The cells were
given a media change of the same composition every 24 hours. The cells were
harvested after 5 days of 15d – PGJ_2 treatment for either RNA or protein isolation.

▪ **X-Gal staining:**

Proliferative and day-13 dox-treated cells were seeded at a density of $1x10^5$ cells
/well into 35mm dishes. After overnight incubation, SA-β-gal activity was measured
by using Senescence β-Galactosidase Staining Kit (Cell Signalling #9860) per
manufacture's protocol. In brief, the cells were washed with PBS twice and were
fixed for 10-15 min at RT. Following three washes of PBS, cells were incubated
overnight in staining solution at 37℃ in a CO_2–free chamber. Development of blue
colour was examined with Ti2 widefield inverted microscope (Nikon).

• **Immunoprecipitation:**

C2C12 cells transfected with EGFP-HRas and treated with 15d-PGJ_2-Biotin were
harvested and lysed in RIPA-PP buffer and the lysate was centrifuged @15000 rpm,
4^o C, 30 minutes. Protein estimation was done using BCA assay kit (G Biosciences).
100 μg of protein was loaded on 10 μl MyOne Streptavidin C1 dynabeads blocked
with 1% BSA in IP washing buffer. The lysate-streptavidin mix was incubated @4^o C,
10 rpm overnight. The beads were then washed thrice with IP washing buffer and

then boiled in 20 µl Laemmlli buffer. 15 µl of the beads were loaded on 12% SDS –
Polyacrylamide gel for detection of EGFP-HRas by immunoblotting using EGFP
antibody.

- **Gene expression analysis:**
  - **Western blotting:**

For measuring Erk/Akt phosphorylation in C2C12 cells (untransfected/ transfected
with EGFP-tagged HRas WT/ HRas-C181S/ HRas-C184S/ HRas V12/ HRas V12-
C181S/ HRas V12-C184S) were seeded in 35 mm dishes. 1x 35 mm dish was
harvested in RIPA – PP the next day, while the rest were incubated in DMEM
starvation medium @37° C. The cells were treated with 15d - PGJ_2 after 24 hrs of
starvation @37° C. The cells were harvested 1 hr after treatment in RIPA - PP.
Protein quantification was done using BCA assay (G Biosciences) using the
manufacturer's protocol. An equal mass of proteins was loaded onto a 12% SDS
Polyacrylamide gel in Laemmlli buffer. The proteins were transferred onto a PVDF
membrane and were probed with phospho-Erk/Erk antibodies for measuring Erk
phosphorylation and with phospho-Akt/Akt antibodies for measuring Akt
phosphorylation. For measuring the expression of Myosin heavy chain, C2C12 cells
expressing EGFP-tagged HRas WT/ HRas-C181S/ HRas-C184S/ HRas V12/ HRas
V12-C181S/ HRas V12-C184S or Human Skeletal Muscle Myoblasts were seeded in
35 mm dishes and were harvested in RIPA - PP after 5 days of differentiation.
Protein quantification was done using BCA assay (G Biosciences) using the
manufacturer's protocol. An equal mass of proteins was loaded onto an 8% SDS
Polyacrylamide gel in Laemmlli buffer. The proteins were transferred onto a PVDF
membrane and were probed with Myosin Heavy Chain Antibody.

  - **qPCR:**

C2C12 cells, untransfected or expressing EGFP-tagged HRas WT/ HRas-C181S/
HRas-C184S and treated with DMSO/15d - PGJ_2, or expressing EGFP-tagged HRas
V12/ HRas V12-C181S/ HRas V12-C184S were lysed in TRIZol at the end of the
experiment (Invitrogen). RNA was isolated from the lysate by the chloroform-
isopropanol method using the manufacturer's protocol. The RNA was quantified and
1.5 µg of RNA was used to prepare cDNA using PrimeScript 1st strand cDNA
Synthesis Kit (Takara Bio) and random hexamer primer. Gene expression for

differentiation markers was measured by qPCR using PowerUp™ SYBR™ Green Master Mix (Applied Biosystems) and previously reported qPCR primers (p.m. 43). Relative gene expression was quantified using the $\Delta\Delta C_T$ method[44] with 18s rRNA as an internal loading control and DMSO vehicle as an experimental control.

- **Immunofluorescence:**

C2C12 cells were seeded in 35 mm dishes (Corning) on glass coverslips (Blue Star) coated with 0.2% Gelatin (Porcine, Sigma Aldrich) and were fixed with the fixative solution at the end of the experiment. The cells were then permeabilized and blocked with the blocking solution and were then incubated with MyHC antibody in the blocking solution overnight. The cells were then washed with 1x PBS, incubated with fluorophore tagged secondary antibody, and were mounted in Prolong gold antifade medium with DAPI (Invitrogen). The cells were then imaged under FV3000 inverted confocal laser scanning microscope (Olympus – Evident) using appropriate lasers and detectors.

- **HRas distribution between the Golgi and the plasma membrane:**

C2C12 cells expressing EGFP-tagged HRas WT/ HRas-C181S/ HRas-C184S + GalT-TagRFP were starved overnight in DMEM starvation medium and treated with DMSO or 15d – PGJ$_2$ (10 µM) in DMEM complete medium for 24 hrs, with a medium change @ 12 hrs post-treatment. The cells were then fixed with the fixative solution @ R. T., 5 mins, washed with PBS, and stained for plasma membrane with Alexa Fluor 633 conjugated Wheat Germ Agglutinin (WGA - 633) (Invitrogen) (10 µg/ml) @R. T. 15 mins. The cells were washed with PBS and were then mounted on glass slides in ProLong Gold Antifade Mounting medium (Invitrogen). C2C12 cells expressing EGFP-tagged HRas V12/ HRas V12-C181S/ HRas V12-C184S were also fixed with the fixative solution @R. T., 5 mins, washed with PBS, stained with WGA – 633, and mounted on slides in Prolong gold antifade medium (Thermo Scientific). The cells were imaged with FV 3000 inverted confocal laser scanning microscope (Olympus - Evident) using appropriate lasers and detectors. Preliminary image processing was done using ImageJ (NIH), while batch analysis of HRas at the plasma membrane and the Golgi complex was done using a custom MATLAB script, where EGFP-HRas image was overlayed onto the GalT-TagRFP and WGA-633 image to obtain HRas localization at the Golgi complex and the Plasma Membrane

respectively. A ratio of mean HRas intensity at the Golgi complex to that of at the Plasma membrane (R_{mean}) was calculated and was used to compare HRas distribution between treatments.

- **Intracellular activity of HRas:**

C2C12 cells expressing EGFP-tagged HRas WT/HRas C181S/HRas C184S + mCherry-RAF RBD were starved overnight in the DMEM starvation medium. The cells were imaged with FV 3000 inverted confocal laser scanning microscope (Olympus-Evident) using the following lasers and detectors:

1. Donor Channel: 488nm excitation, 510 (+/-) 20nm detection.
2. Acceptor Channel: 561nm excitation, 630 (+/-) 50nm detection.
3. FRET Channel: 488nm excitation, 630 (+/-) 50nm detection.

The cells were then treated with 15d-PGJ$_2$ (10 µM) for 1 hour and were imaged using the same imaging parameters. C2C12 cells expressing EGFP-HRas or mCherry-RAF RBD only were used to calculate the bleed-through corrections (EGFP emission @ 630 (+/-) 50nm, and Excitation of mCherry by 488 nm laser). Preliminary processing was done using ImageJ (NIH). The FRET index was calculated using the FRET and co-localization analyzer plugin(Hachet-Haas et al., 2006). The FRET index was then divided by the intensity of the Acceptor channel to normalize the variation in the expression of mCherry. We used the mean normalized FRET index to compare the activity of HRas before and after treatment with 15d-PGJ$_2$.

- **Animal care:**
  - **Maintenance:**

Mice were maintained at BLiSC Animal Care and Resource Centre (ACRC). All the procedures performed were approved by the Internal Animal Users Committee (IAUC) and the Institutional Animal Ethics Committee (IAEC).

  - **Treatment and tissue collection:**

12–15-week-old C57BL/6J (JAX#000664) mice were injected intraperitoneally (I.P.) with 5 mg/kg Doxorubicin (Doxo) four times, once every three days. Intraperitoneal injection of Saline was used as a control. The mice were sacrificed on Day 11 after the first injection. Hindlimb muscles from 4 animals (control and treated with Doxo

each) were used for qPCR analysis and Hindlimb muscles from 3 animals (control and treated with Doxo each) were used for immunohistochemical analysis.

- **Lipid extraction and detection of 15d-PGJ$_2$ by mass spectrometry:**

For lipid extraction, cell pellets were resuspened in 3ml of a methanol solvent [water: methanol: 2:1, 1% formic acid (FA)] whereas only 1 ml of methanol with 3% FA was added to the 2ml of CM, making a uniform sample volume of 3 ml. Subsequently, 1ml of ethyl acetate was added to each sample and mixed vigorously. Phase separation was done by centrifuging the mixture (12000xg, 4℃ for 10 mins), and the organic phase containing the lipid was collected. This process was repeated thrice in total and all the organic phases were combined and dried under a nitrogen stream at RT. The residues were resuspended in 100 µl of 50% acetonitrile in water with 0.1% FA and were subjected to mass spec analysis using the Waters® Acquity UPLC class I system The detection of 15d-PGJ$_2$ was performed using an electrospray ionisation source (ESI) operating in the negative ion mode and a quadrupole trap mass spectrometer (AB SCIEX QTRAP 6500) connected to a Waters® Acquity UPLC class I system (Waters, Germany) outfitted with a binary solvent delivery system with an online degasser and a column manager with a column oven coupled to a UPLC autosampler. 5 µl samples were injected into the union for analysis. Solvent A consisted of 0.1% ammonium acetate in water and solvent B was 0.1% ammonium acetate in a mixture of acetonitrile/water (95:5). For each run, the LC gradient was: 0 min, 20% B; 0.5 min, 20% B; 1.5 min, 90% B; 2.5 min, 20% B; 3min, 20% B. Analyte detection was performed using multiple reaction monitoring (MRM), 315.100 ➔ 271.100 and 315.100 ➔ 203.100. Source parameters were set as follows: capillary voltage 3.8 kV, desolvation gas flow 25 L/h, source temperature 350 °C, ion source gas 1 flow 40 L/h and ion source gas 2 flow 40 L/h. Acquisition and quantification were completed with Analyst 1.6.3 and Multiquant 3.0.3, respectively (method adopted from (Morgenstern et al., 2018)).

For the standards, 2ml media of different known concentrations (50nM, 100nM, 250nM, and 500nM) of 15d-PGJ$_2$ were prepared and were subjected to the same extraction procedure as that of CM. A standard curve was plotted with the known concentration and the mass spec peak area, and the concentration of the lipid in samples was calculated.

Table 1: List of Primers:

Primer Name	Primer Sequence (5' – 3')
HRas WT Fwd	CCGCTCGAGCTATGACGGAATATAAGCTGG
HRas WT Rev	CCGGAATTCTCAGGAGAGCACACACTTGC
HRas C181S Rev	CCGGAATTCTCAGGAGAGCACACACTTGCAGCTCATGCTGCCGG
HRas C184S Rev	CCGGAATTCTCAGGAGAGCACACACTTGCTGCTCATG
mm 18s qPCR Fwd	CCCGTTGAACCCCATTCGTG
mm 18s qPCR Rev	GGGCCTCACTAAACCATCCA
mm MyoD qPCR Fwd	TCCGCTACATCGAAGGTCTG
mm MyoD qPCR Rev	GTCCAGGTGCGTAGAAGGC
mm MyoG qPCR Fwd	CGATCTCCGCTACAGAGGC
mm MyoG qPCR Rev	GTTGGGACCGAACTCCAGT
mm MyHC qPCR Fwd	TAAACGCAAGTGCCATTCCTG
mm MyHC qPCR Rev	GGGTCCGGGTAATAAGCTGG
mm PTGS1 qPCR Fwd	TTACTATCCGTGCCAGAACCA
mm PTGS1 qPCR Rev	CCCGTGCGAGTACAATCACA
mm PTGS2 qPCR Fwd	TTCCAATCCATGTCAAAACCGT
mm PTGS2 qPCR Rev	AGTCCGGGTACAGTCACACTT
mm PTGDS qPCR Fwd	GAGTACGCTCTGCTATTCAGC
mm PTGDS qPCR Rev	GGTTGGGGCAGGAAAACAATG
mm p16 qPCR Fwd	GCTCAACTACGGTGCAGATTC
mm p16 qPCR Rev	GCACGATGTCTTGATGTCCC
mm p21 qPCR Fwd	CCTGGTGATGTCCGACCTG
mm p21 qPCR Rev	CCATGAGCGCATCGCAATC
mm p53 qPCR Fwd	CCCCTGTCATCTTTTGTCCCT
mm p53 qPCR Rev	AGCTGGCAGAATAGCTTATTGAG
mm CXCL1 qPCR Fwd	ACTGCACCCAAACCGAAGTC
mm CXCL1 qPCR Rev	TGGGGACACCTTTTAGCATCTT
mm CXCL2 qPCR Fwd	CCAACCACCAGGCTACAGG
mm CXCL2 qPCR Rev	GCGTCACACTCAAGCTCTG
mm TNF1α qPCR Fwd	CCTGTAGCCCACGTCGTAG
mm TNF1α qPCR Rev	GGGAGTAGACAAGGTACAACCC

mm IL6 qPCR Fwd	CTGCAAGAGACTTCCATCCAG
mm IL6 qPCR Rev	AGTGGTATAGACAGGTCTGTTGG
mm TGF1β qPCR Fwd	CTTCAATACGTCAGACATTCGGG
mm TGF1β qPCR Rev	GTAACGCCAGGAATTGTTGCTA

1101

1102 **Table 2: List of Reagents**

Item Description	Manufacturer	Catalog No.
pEGFP-C1	Clontech	
Phusion high fidelity DNA polymerase	Thermo Scientific	F 530
Xho1 restriction enzyme	New England Biolabs Inc.	R0146
EcoR1 restriction enzyme	New England Biolabs Inc.	R0101
T4 DNA Ligase	Takara Bio	2011
DMEM High Glucose	Gibco	11995065
FBS, Certified (Origin: United States)	Gibco	16000044
Horse Serum, Heat Inactivated (Origin: New Zealand)	Gibco	26050070
Penicillin – Streptomycin – Glutamine (100x)	Gibco	10378016
Penicillin – Streptomycin (100x)	Gibco	15140163
DPBS, no calcium, no magnesium	Gibco	14190144
0.25% Trypsin - EDTA	Gibco	25200056
jetPRIME transfection reagent	Polyplus	101000046
Doxorubicin	Sigma – Aldrich	D1515
15-deoxy-$\Delta^{12,14}$-Prostaglandin J$_2$	Cayman Chemical Company	18570
9,10-dihydro-15-deoxy-$\Delta^{12,14}$-Prostaglandin J$_2$	Cayman Chemical Company	18590
15-deoxy-$\Delta^{12,14}$-Prostaglandin J$_2$-Biotin	Cayman Chemical Company	10141
Dynabeads™ MyOne™ Streptavidin C1	Invitrogen	65001
cOmplete Protease inhibitor cocktail tablets	Roche	11697498001
Wheat germ agglutinin, Alexa fluor 633 conjugate	Invitrogen	2307289
ProLong Gold Antifade Mounting medium	Invitrogen	P36930
Paraformaldehyde	Sigma – Aldrich	158127

TRIzol reagent	Invitrogen	15596018
PrimeScript™ 1st strand cDNA Synthesis Kit	Takara Bio	6110A
PowerUp™ SYBR™ Green Master Mix	Applied Biosystems	A25742
WesternBrightTM ECL-spray Western blotting detection system	advansta	K-12049-D50
Fetuin (Bovine)	Sigma – Aldrich	F2379
hEGF	Sigma – Aldrich	E9644
N2 Supplement	Thermo Scientific	17502048
Dexamethasone	Sigma – Aldrich	D4902
DMEM Low Glucose	Thermo Scientific	10567014
Bovine Serum Albumin Fraction-V, Cell culture tested	HIMEDIA	9048468
PierceTM Streptavidin Magnetic Beads	Thermo Scientific	88816

1103

1104   **Table 3: List of Antibodies**

Antibody Name	Clonality	Species of Origin	Manufacturer	Catalog No.
Phospho-Erk (Thr202/ Tyr204)	Polyclonal	Rabbit	Cell Signaling Technology	9101S
Erk	Polyclonal	Rabbit	Cell Signaling Technology	9102S
GAPDH	Monoclonal	Mouse	Puregene	PG23002
β-actin	Polyclonal	Rabbit	Cell Signaling Technology	4967S
Phospho-Akt (Ser473)	Monoclonal	Mouse	Cell Signaling Technology	4051S
Akt	Monoclonal	Rabbit	Cell Signaling Technology	4691S
GFP	Monoclonal	Rabbit	Cell Signaling Technology	2956S
Myosin Heavy Chain	Monoclonal	Mouse	Invitrogen	14-6503-82
Myosin Heavy Chain	Monoclonal	Mouse	Developmental Studies Hybridoma Bank (DSHB)	

p21	Monoclonal	Mouse	Santacruz Biotechnology	sc-6246
γH2A.X	Polyclonal	Rabbit	Novus Biologicals	NB100384
Tubulin	Monoclonal	Mouse	Cell Signaling Technology	3873S
HRP - Anti-Mouse		Horse	Cell Signaling Technology	7076S
HRP - Anti-Rabbit		Goat	Cell Signaling Technology	7074P2
Alexa Fluor 568 – Anti-Mouse	Polyclonal	Goat	Invitrogen	A11031

1105

Reagent Composition

• DMEM complete medium: DMEM Hi Glucose medium (Gibco) supplemented
with 1% Penicillin – Streptomycin – Glutamine (Gibco) and heat-inactivated
10% Fetal Bovine Serum (US origin) (Gibco).

• Basal Conditioned medium: DMEM Hi Glucose medium (Gibco)
supplemented with 1% Penicillin – Streptomycin – Glutamine (Gibco) and
heat-inactivated 2% Fetal Bovine Serum (US origin) (Gibco).

• C2C12 differentiation medium: DMEM Hi Glucose medium (Gibco)
supplemented with 2% Horse Serum (Gibco) and 1% Penicillin –
Streptomycin – Glutamine (Gibco).

• DMEM Starvation medium: DMEM Hi Glucose medium (Gibco) supplemented
with 0.2%heat-inactivated fetal bovine serum (US origin) (Gibco) and 1%
Penicillin – Streptomycin – Glutamine (Gibco).

• RIPA – PP buffer: RIPA buffer (Invitrogen) supplemented with protease
inhibitor cocktail (Roche) and 5 mM Sodium Fluoride and 5 mM Sodium
Orthovanadate.

• TBS – T buffer: 50 mM Tris–Cl (pH = 7.5), 150 mM NaCl and 0.1% Tween –
20 in water.

• PBS: 2.67 mM KCl, 1.47 mM KH_2PO_4, 137.93 mM NaCl, 8.06 mM Na_2HPO_4
in water.

- IP Washing Buffer: 150 mM NaCl, 0.1% SDS, 1% NP-40 in 50 mM Tris-Cl (pH=7)
- Fixative Solution: 4% (w/v) Paraformaldehyde (Sigma – Aldrich) in PBS.
- Blocking Solution: 2% Heat Inactivated FBS, 0.2% BSA, 0.2% Triton - X, 0.05% NaN_3 in PBS.
- Skeletal Muscle Growth Medium: DMEM Low Glucose Medium (Gibco), supplemented with 1% Penicillin – Streptomycin – Glutamine (Gibco), heat-inactivated 10% Fetal Bovine Serum (US origin) (Gibco), Bovine Fetuin (50 µg/ml) (Sigma – Aldrich), Dexamethasone (0.4 µg/ml), and hEGF (10 ng/ml).
- Skeletal Muscle Differentiation Medium: DMEM low glucose medium (Gibco) supplemented with 2% Horse Serum, 1% Penicillin – Streptomycin (Gibco), and 1% N2 Supplement.

Supplementary Figures:

Figure S1. Treatment with Doxo induces senescence *in vivo* and *in vitro* and induces release of eicosanoid prostaglandin 15d-PGJ$_2$

A. Schematic Representation of the experimental flow for treatment of B6J mice with Doxo (5 mg/kg) or Saline.

B. Schematic representation of the experimental flow for treatment of senescent C2C12 cells with prostaglandin D synthase (PTGDS) inhibitor AT-56 (30 µM) or DMSO.

C. Widefield micrograph of the morphology of C2C12 cells after treatment with Doxo (150 nM) or DMSO.

D. Widefield micrograph of expression of Senescence Associated β-galactosidase (SA β-gal), measured by X-gal staining, in C2C12 cells after 13 days of treatment with Doxo (150 nM) or DMSO.

E. Representative peaks from Blank sample.

F. Representative peaks from Conditioned medium of C2C12 myoblasts treated with DMSO.

G. Representative peaks from Conditioned medium of C2C12 myoblasts treated with Doxo (150 nM).

H. Standard curve of peak area vs known conc. of 15d-PGJ$_2$ detected using AB SCIEX QTRAP 6500 mass spectrometer.

Sample	15d-PGJ$_2$ (pmol)	Cell Number	15d-PGJ$_2$/Cell (fg)
Intracellular Quiescent C2C12 cells_1	27.087	462500	18.531
Intracellular Quiescent C2C12 cells_2	18.080	650000	8.801
Intracellular Senescent C2C12 cells_1	809.633	287500	891.018
Intracellular Senescent C2C12 cells_2	940.032	300000	991.420
Conditioned medium Quiescent C2C12 cells_1	8.271	462500	5.659
Conditioned medium Quiescent C2C12 cells_2	6.554	650000	3.190
Conditioned medium Senescent C2C12 cells_1	45.814	287500	50.419
Conditioned medium Senescent C2C12 cells_2	38.893	300000	41.019

Supplementary Table 1: 15d-PGJ$_2$ conc. (pmol and fg/cell) detected in samples by mass spectrometry.

Figure S2. Reduced viability of C2C12 myoblasts after treatment with 15d-PGJ$_2$ (5 and 10 µM) in differentiating medium.

A. Representative images taken every 24 hours of C2C12 treated with 15d-PGJ$_2$ (5 and 10 µM) or DMSO in the C2C12 differentiation medium.

B. Representative images taken every 24 hours of C2C12 treated with 15d-PGJ$_2$ (1, 2, and 4 µM) or DMSO in the C2C12 differentiation medium.

Figure S3. Activation of HRas after treatment with 15d-PGJ$_2$.

A. Representative confocal micrograph of C2C12 cells expressing only EGFP-tagged HRas or mCherry-tagged RAF RBD alone for spectral overlap (bleed-through) calculations.

B. Mean bleed-through of EGFP-tagged HRas (Donor) and mCherry-tagged RAF RBD (Acceptor) in the FRET channel.

C. Phosphorylation of Akt (measured by immunoblotting) in C2C12 cells treated with 15d-PGJ$_2$ (5, 10 µM) or DMSO for 1 hr after starving the cells in 0.2% serum medium for 24 hrs. The densitometric ratio of phosphorylated Akt (Ser473) to total Akt was normalized to non-starved C2C12 cells. (Statistical significance tested by two-tailed heteroscedastic student's t-test, N=3).

Figure S4. C-terminal cysteine mediated intracellular distribution of constitutively active HRas (HRas V12) regulates differentiation of C2C12 myoblasts.

A. Distribution of EGFP-tagged HRas WT/ HRas V12/ HRas-C181S/ HRas V12-C181S/ HRas-C184S/ HRas V12-C184S between the Golgi complex and the plasma membrane, as seen by the scatter plot of R_{mean}, the ratio of the mean HRas intensity at the Golgi complex to that at the plasma membrane, in C2C12 cells. (Statistical significance tested by two-tailed heteroscedastic student's t-test, N=3)

B. mRNA levels of MyoD, MyoG, and MyHC relative to 18s rRNA (measured by quantitative PCR) in C2C12 cells expressing EGFP-tagged HRas V12/ HRas V12-C181S/ HRas V12-C184S in C2C12 differentiation medium for 5 days. (Statistical significance tested by two-tailed heteroscedastic student's t-test, N=3)

C. Protein levels of MyHC (measured by immunoblotting) in C2C12 cells expressing EGFP-tagged HRas V12/HRas V12-C181S/HRas V12-C184S in C2C12 differentiation medium for 5 days. The densitometric ratio of levels of MyHC to α-Tubulin was normalized to C2C12 cells expressing HRas V12. (Statistical significance tested by two-tailed heteroscedastic student's t-test, N=3).

D. Brightfield image of C2C12 cells expressing EGFP-HRas V12, V12 C181S, or V12 C184S on Day 0 and Day 5 of differentiation.

E. Brightfield image of C2C12 cells expressing EGFP-HRas WT, C181S, or C184S after 5 days of treatment with 15d-PGJ$_2$ or DMSO.