Distinct ancient structural polymorphisms control heterodichogamy in walnuts and hickories

Jeffrey S. Groh[1,2], Diane C. Vik[1], Kristian A. Stevens[1,3], Patrick J. Brown[4], Charles H. Langley[1,2], and Graham Coop[1,2]

[1] Department of Evolution and Ecology,

[2] Center for Population Biology,

[3] Department of Computer Science,

[4] Department of Plant Sciences,

University of California, Davis.

To whom correspondence should be addressed: `jgroh@ucdavis.edu`

Abstract

The maintenance of stable mating type polymorphisms is a classic example of balancing selection, underlying the nearly ubiquitous 50/50 sex ratio in species with separate sexes. One lesser known but intriguing example a balanced mating polymorphism in angiosperms is heterodichogamy - polymorphism for opposing directions of dichogamy (temporal separation of male and female function in hermaphrodites) within a flowering season. This mating system is common throughout Juglandaceae, the family that includes globally important and iconic nut and timber crops - walnuts (*Juglans*), as well as pecan and other hickories (*Carya*). We show that two distinct genetic systems control heterodichogamous mating systems in these genera. In each genus, we find ancient (>50Mya) structural variants that segregate as trans-species polymorphisms. The *Juglans* locus maps to a ca. 20kb structural variant adjacent to gene with likely function in the trehalose-6-phosphate signaling pathway, which is known to regulate flowering time in Arabidopsis and inflorescence architecture in maize. Notably, an insertion in the dominant (protogynous) haplotype contains 8-12 complex tandem duplicates of the 3' UTR of the gene, and transpecies polymorphism within the gene is restricted to the 3' UTR. In contrast, the *Carya* heterodichogamy locus maps to a ca. 200-450 kb cluster of tightly linked polymorphisms across 20 genes, some of which have known roles in flowering and are differentially expressed between morphs in floral primordia. The dominant (protogynous) haplotype in pecan, which is nearly always heterozygous and appears to rarely recombine, shows reduced genetic diversity and is over twice as long as its recessive counterpart due to proliferation of multiple transposable element families. We did not detect either genetic system in another heterodichogamous genus within the Juglandaceae, suggesting that additional genetic systems for heterodichogamy may yet remain undiscovered.

Keywords: heterodichogamy, walnut, pecan, Juglandaceae, *Juglans*, *Carya*, supergene, trans-species polymorphism, balanced polymorphism, T6P, flowering time, structural variation

Introduction

Flowering plants have evolved a remarkable diversity of sexual heteromorphisms, e.g. dioecy, heterostyly, gynodioecy, mirror-image flowers, that have long fascinated evolutionary biologists (Darwin 1877; Charlesworth and Charlesworth 1979; Barrett 2010). Such systems present key opportunities to test evolutionary theories of sexual reproduction and to understand its ecological and genetic consequences. Heterodichogamy presents an intriguing and relatively unexplored example of a balanced polymorphism in the sexual organization of some angiosperms, in this case not in space but in time. In Darwin's words, heterodichogamous populations

"consist of two bodies of individuals, with their flowers differing in function, though not in structure; for certain individuals mature their pollen before the female flowers on the same plant are ready for fertilisation and are called proterandrous [protandrous]*; whilst conversely other individuals, called proterogynous* [protogynous]*, have their stigmas mature before their pollen is ready"*(1877). This system generates strong disassortative mating between morphs (Bai *et al.* 2007), thus classical sex ratio theory predicts a 50/50 ratio of protandrous and protogynous individuals at equilibrium (Gleeson 1982). Although heterodichogamy has likely evolved at least a dozen times in angiosperms (Renner 2001; Endress 2020), to our knowledge the genetic loci controlling the inherited basis of this mating system have not been described at the molecular level in any species. Thus, it is not known whether similar genetic mechanisms control heterodichogamy in different taxa, nor whether the loci involved experience similar evolutionary dynamics as sex chromosomes and other supergenes that underpin complex mating polymorphisms (Charlesworth 2016; Gutiérrez-Valencia *et al.* 2021).

Heterodichogamy is well-known within Juglandaceae (Delpino 1874; Darwin 1876; Pringle 1879; Stuckey 1915, Fig. 1A,B), the family that includes two major globally important nut and timber-producing groups of trees - the walnuts (*Juglans*) and hickories (*Carya*, includes cultivated pecans). The divergence of these genera roughly 58-72 Mya (Mu *et al.* 2020; Zhou *et al.* 2021) is supported by a rich fossil record for Juglandaceae (Manchester 1987; Manos *et al.* 2007; Zhang *et al.* 2022). In both genera, protogyny is inherited via a dominant Mendelian allele (dominant allele - *G*, recessive allele - *g*, Gleeson 1982; Thompson and Romberg 1985). Furthermore, other genera phylogenetically closer to *Juglans* (*Cyclocarya, Platycarya*) are also known to be heterodichogamous (Fukuhara and Tokumaru 2014; Mao *et al.* 2019). These observations have suggested a single origin of the alleles controlling heterodichogamy in the common ancestor of these taxa. Here, we identify and characterize the evolutionary history of the genetic loci controlling heterodichogamy in both *Juglans* and *Carya*. We find two distinct and substantively different genetic underpinnings for heterodichogamy in these genera, and show that each genetic system is an ancient trans-species balanced polymorphism for a structural variant.

An ancient structural variant controls heterodichogamy across *Juglans*

In a natural population of Northern California black walnut (*J. hindsii*), protandrous and protogynous morphs were found to occur in roughly equal proportions (43 vs. 38, P=0.47 under null hypothesis of equal proportions). A genome-wide association study (GWAS) for dichogamy type in this population recovered a single, strong association peak, consistent with the known Mendelian inheritance of heterodichogamy type in *Juglans* (Fig. 1C)(Gleeson 1982). This region is syntenic with the top GWAS hit location for dichogamy type in *J. regia* (supp Fig. S1, Bernard *et al.* 2020) as well as a broad region associated with dichogamy type in *J. nigra* (Chatwin *et al.* 2023). The association peak extends roughly 20kb (hereafter, the G-locus) across a probable trehalose-6-phosphate phosphatase gene (*TPPD-1*, Fig. 1D, corresponding to LOC108984907 in the Walnut 2.0 assembly). TPPs catalyze the dephosphorylation of trehalose-6-phosphate to trehalose as part of a signaling pathway essential for establishing regular flowering time in Arabidopsis (Wahl *et al.* 2013) and inflorescence architecture in maize (Satoh-Nagasawa *et al.* 2006). Consistent with the width of the GWAS hit, we observed strong linkage disequilibrium (LD) extending across this region, suggesting locally reduced recombination in the genealogical history of the sample for both *J. hindsii* and *J. regia* (Fig. S2).

We identified copy number variation consistent with three indels distinguishing G-locus haplotypes (*GJ1* and *GJ2* in the G haplotype; *gJ3* in the g haplotype, Figs. 1E, S1). Consistent with the known dominance, all the *J. hindsii* protandrous individuals (*gg*) appear to be homozygotes for *gJ3* and protogynous individuals (*G?*) appear to be hemizygous for *GJ1* and *GJ2*. The apparent absence of any homozygotes for the *G* haplotype in our *J. hindsii* sample is consistent with disassortative mating at the G-locus in a natural population (P=0.059). Strikingly similar copy number patterns were seen in *J. regia* (Fig. 1F, S1), with a known homozygote for the protogynous allele ('Sharkey', *GG*) (Gleeson 1982) lending further support for these indels. To look for the presence of this structural variant across *Juglans* species, we generated whole-genome resequencing data from both morphs from 5 additional *Juglans* species (Table S1) combined with existing data from known morphs in two additional species (Stevens *et al.* 2018). Parallel patterns of read depth at the G-locus indicate that the same structural variant segregates in perfect association with dichogamy type across all nine species examined (Fig. 1F, S3). As our species sampling spans the deepest

Figure 1: **(A)** Protandrous and protogynous morphs of *J. hindsii* are readily distinguished by catkin size in the first half of the flowering season. **(B)** A protogynous individual of *J. ailantifolia* with developing fruits and that is currently shedding pollen. Photo by JSG. **(C)** GWAS of flowering type in 44 individuals of *J. hindsii* from a natural population. **(D)** The GWAS peak occurs across and in the 3' region of *TPPD-1*, a probable trehalose-6-phosphate-phosphatase gene (blue rectangle, arrow indicates direction of transcription). Red bar shows the position of a *NDR1/HIN1*-like gene. **(E)** Normalized average read depth in 1kb windows for 44 *J. hindsii* from a natural population against a genome assembly for *J. californica*. Black bars show positions of identified indels. **(F)** Read depth of 26 *J. regia* accessions against an assembly for *J. regia* in the region syntenic with the *J. hindsii* G-locus. Colored bars and arrows indicate positions of orthologous genes bordering the G-locus.

divergence event in the genus (∼47 Mya, Mu *et al.* 2020), this suggests that the G-locus haplotypes may have diverged in the common ancestor of *Juglans*.

Consistent with the *Juglans* G-locus being an ancient balanced polymorphism, nucleotide divergence between G-locus haplotypes is exceptionally high compared to the genome-wide background (falling in the top 1% of values across the entire aligned chromosome in several species, Fig. 2D). Applying a molecular clock, we estimated the age of the G-locus haplotypes to be 68.6 Mya (although there is considerable uncertainty in this date, see methods), approximately 22 million years older than the most recent common ancestor of the genus (Mu *et al.* 2020).

We next examined variation within the *TPPD-1* gene sequence. We identified several transpecific polymorphic SNPs within the 3' UTR of *TPPD-1*, but only two within *TPPD-1* coding sequence, both of these are in the last exon and these do not group by haplotype across all species comparisons. Moving away from the indels, we observed that polymorphism quickly transitions to grouping by species rather than haplotype upstream of the 3' UTR (Fig. S4). These observations imply recombination within the *TPPD-1* transcribed region over phylogenetic timescales, as well as a lack of shared functional divergence between G-locus haplotypes within *TPPD-1* coding sequence. However, we observed 14 fixed differences between G-locus haplotypes in a sample of 113 *J. regia* (Stevens *et al.* 2018; Ji *et al.* 2021; Ding *et al.* 2022) stretching upstream all the way to the 2nd exon, nine of which are nonsynonymous substitutions. Combined with the elevated LD across *TPPD-1* coding sequence seen within both *J. regia* and *J. hindsii*, this raises the intriguing possibility of recruitment and turnover of functional divergence within the *TPPD-1* gene sequence or elsewhere within the G-locus.

Nonetheless, the lack of shared coding sequence divergence suggests that regulatory divergence rather than protein sequence divergence may underlie the differential development of dichogamy types in *Juglans*. We note that in *J. regia*, *TPPD-1* is expressed in a broad range of vegetative and floral tissues (Fig. S7), and we detected full-length transcripts from multiple tissues related to flowering in *J. regia* homozygotes for both G-locus haplotypes (Table S2). The developmental basis of heterodichogamy in *J. regia* is driven by a differential between protogynous and protandrous morphs in the extent of both male and female floral primordia differentiation in the growing season prior to flowering (Luza and Polito 1988; Polito and Pinney 1997). Correspondingly, we found that protandrous and protogynous morphs of *J. regia* differ in the size of dormant catkin buds by mid-summer (Fig. S5). We propose that variation at the G-locus may drive differential gene expression of *TPPD-1* in floral primordia or in leaves, as the T6P signaling pathway has been shown to regulate Arabidopsis flowering time independently in these two tissues (Wahl *et al.* 2013).

We next investigated the three indels between the *G* and *g* haplotypes (Fig 2A). The indels *GJ2* and *gJ3* are not obvious functional candidates. For example, *gJ3* is derived from an insertion of a CACTA-like DNA transposon in *g* haplotypes of *Juglans*, which lacks evidence of gene expression or sequence conservation across species (Fig. S6). Nonetheless, we cannot at present rule out a functional role of these indels in the differential development of alternate dichogamy types.

We next turned to the *GJ1* indel closer to *TPPD-1*. Using pairwise alignment of the haplotypes in *J. regia*, we found that *GJ1* contains a series of ∼1kb tandem repeats with homology to the 3' region of *TPPD-1* (Fig. 2 A,B). We find no evidence for the inclusion of any part of the repeat in full length *TPPD-1* transcripts from a *GG J. regia* individual. Each repeat, ordered *GJ1-1* up to *GJ1-12* moving downstream of *TPPD-1*, consists of three subunits. One subunit (*GJ1a*) is an inverted ∼250-300bp motif homologous with the 3' UTR of *TPPD-1*. The other two subunits (*GJ1b* and *GJ1c*) are non-inverted ∼300bp motifs homologous with regions within 1kb downstream of the 3' UTR. The short tandem duplicate is present in *G* haplotypes across multiple species with long read assemblies, varying from 8-12 copies (Fig. 2B). Consistent with G-locus haplotypes arising in the ancestor of *Juglans*, a maximum-likelihood phylogeny of concatenated aligned sequence from *GJ1* repeats and their homologous sequence immediately downstream of the *TPPD-1* (Table S3) revealed that all *G* haplotype sequences form a sister clade to *g* haplotype sequences (Fig. 2C). Among more closely related species (*J. microcarpa* and *J. californica*, ∼12 Mya species divergence) we found evidence of conservation of specific repeats, with some repeats from the same relative positions in different species clustering together phylogenetically. On the other hand, for deeper divergences (∼ 40-50Mya species divergence), evolution of *GJ1* repeats is characterized by lineage-specific turnover due to expansion and contraction of the repeat array and/or possible gene conversion. For example, the majority of repeats within *J. regia* are most closely related to repeats at other positions within the same species, and likewise for *J. mandshurica*.

Figure 2: **(A)** Schematic of the G-locus structural variant in *Juglans*. Presence/absence of indels are indicated by continuous lines vs. open parentheses. Maroon and blue bars show bordering genes. Black arrow across *TPPD-1* indicates direction of transcription. Colored arrows represent components of the repeated units within the *GJ1* indel and homologous sequence outside of the indel. Each repeat motif (numbered 1,2,...) is comprised of subunits a, b, and c. Subunit a, which is homologous with the 3' UTR of *TPPD-1*, is inverted within the indel. **(B)** Dotplots showing pairwise alignment of alternate haplotypes for three species representing the major clades within the genus. Maroon and blue rectangles indicate the locations of the genes bordering the G-locus as in (A). **(C)** Phylogeny of GJ1 repeats. Sample codes from L-R indicate: haplotype (for *Juglans* only), abbreviated binomial, optional additional identifier, and relative position (as shown in A) (see Table S3 for full taxon list and data sources). Sequences from *g* haplotypes (gold) and *G* haplotypes (blue) form two sister clades (*****, 99% bootstrap support) whose most recent common ancestor predates the radiation of *Juglans*. Daggers and diamonds highlight representative sequences from *J. californica* and *J. regia* that show the trans-species polymorphism. **(D)** Average nucleotide divergence in 500bp windows between alternate G-locus haplotypes for three within-species comparisons. Dotted gray lines indicate the chromosome-wide average, and shaded gray regions indicate 95% quantiles. Maroon and blue bars indicate positions of genes as in (A).

Taken together, multiple independent lines of evidence support that the control of dichogamy type in *Juglans* is governed by structural variation at the G-locus that predates the radiation of the genus. More specifically, the narrow range of conserved trans-specific polymorphisms within the 3' UTR of *TPPD-1* and the conservation of numerous (≥ 8) 3' UTR homologous repeats are evidence for a functional role in the regulation of *TPPD-1*. The known role of the TPPs and their substrate, T6P, in regulating critical and sensitive signaling systems in plants, and specifically developmental transitions in flowering (Ponnu *et al.* 2011), lends further support to this idea.

Finally, we note that while the divergence of G-locus haplotypes predates the *Juglans* radiation, our phylogeny inference suggests it is more recent than the divergence of *Juglans* with its closest cousins, *Pterocarya* and *Cyclocarya*, and the relationships of non-*Juglans* sequences resemble previously estimated species trees for these taxa (Fig. 2C). Consistent with this, we found no evidence of the G-locus structural variant in resequencing data from multiple individuals from *Pterocarya* and *Cyclocarya* (Fig. S8).

The G-locus in *Carya* is a distinct trans-species supergene

While heterodichogamy has been suggested as the ancestral state of all Juglandaceae, our results indicate that the *Juglans* G-locus is specific to walnuts. To identify the basis of the trait in pecan (*C. illinoinensis*), we generated whole-genome resequencing data from 18 pecan varieties of known dichogamy type and combined these with existing data for 12 additional varieties (Xiao *et al.* 2021) to perform a GWAS for dichogamy type. We identified single peak of strong association at a locus on chromosome 4, with many alleles in strong LD (Fig. 3, S13), consistent with previous QTL mapping (Bentley *et al.* 2019). This pecan G-locus region is not syntenic with the *Juglans* G-locus, implying either convergent origins of heterodichogamy within Juglandaceae, or a single origin followed by turnover in the underlying genetic mechanism.

We identified a large segregating structural variant in pecan (445kb in the primary haplotype-resolved assembly of 'Lakota', *Gg* diploid genotype) overlapping the location of the GWAS peak that is perfectly associated with dichogamy type (Fig. 3C), and confirmed the genotype of two known protogynous allele homozygotes, (*GG*) 'Mahan' and 'Apache' (Thompson and Romberg 1985; Bentley *et al.* 2019). Coverage patterns indicate the primary assembly of 'Lakota' is of the *G* haplotype. Consistent with this, coverage patterns against the alternate assembly of 'Lakota' and against the reference assembly of 'Pawnee' (protandrous, *gg*) identified both as *g* haplotype assemblies (Fig. S10).

In contrast to the *Juglans* G-locus, the *Carya* G-locus contains approximately 20 predicted protein-coding genes (Fig. 3B, red boxes, Table S4). Gene-level synteny is highly conserved between the *Carya* G-locus haplotypes (Fig. 3F), with the segregating copy number variation largely localized to intergenic regions. Several of these genes have well-characterized homologs involved in flowering (e.g. *FIL1, EMS1, SLK2, CEN*, see Table S4) with plausible roles in morph-specific development of alternate dichogamy types. We found 237 nonsynonymous fixed differences between the pecan haplotypes across these 20 genes, suggesting many potential functional candidates. However, applying the McDonald-Kreitman test to each gene, the ratio of nonsynonymous to synonymous substitutions and polymorphisms did not identify any signals of adaptive evolution. We note however that two of these genes with annotated functional roles in flowering, *EMS1* and *FIL1*, were previously identified in a small set of highly differentially expressed genes between male catkin buds of protandrous and protogynous cultivars in the season prior to bloom near when a differential in anther development is first established (Rhein *et al.* 2023). Pollen differentiation in *EMS1* mutants of Arabidopsis fail to form pollen tetrads (Zhao *et al.* 2002), and protogynous pecans have a delayed progression from pollen tetrads to mature pollen as compared to protandrous pecans (Stuckey 1915). *FIL1* homologs are expressed uniquely or predominantly in stamens (Nacken *et al.*1991, Fig. S9) and were found to be a downstream target of the B class MADS box transcription factor *DEFICIENS* in Antirrhinum (Nacken *et al.* 1991) (the ortholog of Arabidopsis *AP3*).

We found strongly elevated nucleotide divergence between *Carya* G-locus haplotypes against the genome-wide background, suggesting that they could have been maintained as an ancient balanced polymorphism (Fig. 3D). To search for evidence of trans-species polymorphism of the *Carya* G-locus, we compared the North American pecan to two long-read genome assemblies from species in the East Asian *Carya* clade (*C. sinensis* and *C. cathayensis*, Zhang *et al.* 2023), spanning the deepest split within the genus (see Fig. 3E for species tree). Patterns of read depth of the pecan whole-genome resequencing reads against these two

Figure 3: The G-locus in *Carya*. **A)** GWAS for dichogamy type in 30 pecan varieties identifies a single peak of strong association on chromosome 4. **B)** Zoomed in view of the GWAS peak. Boxes underneath plot indicate positions of predicted genes. Red boxes are those that fall inside the region of strong association. **C)** Normalized average read depth in 1kb windows across the location of the GWAS hit reveals a structural variant segregating in perfect association with dichogamy type. **D)** Nucleotide divergence between G-locus haplotypes within coding sequence is strongly elevated against the genome-wide background. Dotted line shows the genome-wide average, shaded interval shows 95% quantile. Colored lines at bottom show locations where other *Carya* species are heterozygous for SNPs that are fixed between pecan G-locus haplotypes. Blue - North American, red - East Asian. **E)** The phylogeny of the G-locus is discordant with the species tree, reflecting trans-species polymorphism. Divergence of *Carya* G-locus haplotypes occurred after the split with *Juglans* and were present in the most recent common ancestor of *Carya*. **F)** Gene-level synteny for assemblies of both G-locus haplotypes in pecan and East Asian *Carya*, and *J. regia*. *G* haplotypes are longer than *g* haplotypes in both major *Carya* clades due to a derived expansion of transposable elements. Colored boxes show positions of genes in each assembly, with color indicating the strand (blue - plus, green - minus). Red bands connect orthologous genes that fall within the GWAS peak in pecan; gray bands connect genes outside the region. Individual gene trees that support the trans-species polymorphism are indicated with asterisks along the top. Black bands and daggers indicate two genes that appear to be uniquely shared by North American and East Asian *G* haplotypes.

assemblies suggested that the *C. cathayensis* and *C. sinensis* assemblies represent the *g* and *G* haplotypes, respectively (Fig. S11). We next constructed a maximum-likelihood consensus phylogeny from concatenated protein-coding sequence from the 20 G-locus genes, and found that it conflicts with the species tree, with *C. cathayensis* clustering with the pecan *g* haplotype and *C. sinensis* clustering with the pecan *G* haplotype (both 100% bootstrap support). Finally, we used whole-genome resequencing data from 16 individuals (15 from Huang *et al.* 2019, 1 from this study) representing 15 different *Carya* species to examine heterozygosity at the G-locus across the genus. We ascertained in pecan a set of 454 SNPs in coding sequence that were fixed between G-locus haplotypes. We found 8/16 individuals in this sample that are highly heterozygous at these SNPs (25-74%, the other half being 1-5%), consistent with these individuals being heterozygous for the G-locus and with dichogamy types being maintained at 50/50 proportions in species across the entire genus (Fig. S12). We conclude from these multiple lines of evidence that the *Carya* G-locus haplotypes are at least as old as the divergence between Eastern North American and East Asian clades of *Carya*, a depth of ∼25 Mya (Zhang *et al.* 2013; Mu *et al.* 2020; Zhou *et al.* 2021).

If variation at the *Carya* G-locus controlled dichogamy types in the ancestor of *Juglans* and *Carya*, we would expect that nucleotide divergence between the *Carya* G-locus haplotypes should match nucleotide divergence between pecan and the orthologous sequence in *Juglans*. Divergence between *Carya* G-locus haplotypes at four-fold degenerate sites across the 20 *Carya* G-locus genes was significantly less than that between either *Carya* haplotype and *J. regia* (0.05 vs 0.065, P=0.00069), implying that the *Carya* G-locus haplotypes diverged more recently than the *Juglans-Carya* split. Assuming a molecular clock and a divergence time of 72 Mya between *Juglans* and *Carya* (Zhou *et al.* 2021), we estimate the age of the *Carya* G-locus haplotypes to be 59.6 Mya.

The *G* haplotype is rarely found in homozygotes due to strong disassortative mating under heterodichogamy, and it seems to experience little recombination (Fig S13), so it may experience similar evolutionary forces to non-recombination portions of Y chromosomes (Bachtrog 2013). The pecan *G* haplotype is 445kb long, over twice as large as the *g* haplotype (200kb), largely due to a difference in transposable element content within the region of reduced recombination (*g*: 40%, *G*: 65%, genome average: 53%, Fig. S14) . Part of the transposable element expansion in the *G* haplotype may be ancient, given parallel coverage patterns and assembly lengths for the East Asian G-locus haplotypes; the *C. sinensis G* assembly is ∼350kb, while the *C. cathayensis g* assembly is ∼230kb. This proliferation of transposable elements within *G* haplotypes might reflect reduced efficacy of selection on the *G* haplotype due to its lower effective population size and the enhanced effects of linked selection due to its low recombination rate (Charlesworth and Charlesworth 2000). Consistent with *G* haplotypes having a reduced effective population size and/or stronger linked selection, we observe a 2.3-fold reduction in genetic diversity in coding regions for *G* haplotypes compared to *g* haplotypes ($\pi = 0.012$ vs. 0.005, respectively).

Theory and data suggests that supergenes (e.g. sex chromosomes) may progressively assemble over time perhaps through the accumulation of morph-antagonistic variation. While synteny of 20 predicted protein-coding genes across *Carya* G-locus assemblies is conserved as far back as the divergence with oaks (∼90Mya, Larson-Johnson 2016) (Fig. S15), we do note several predicted protein-coding genes that differ among *Carya* G-locus assemblies (Fig. 3F, Table S5). Notably, two of these appear to be derived and shared between North American and East Asian *G* haplotypes and so warrant further attention (Table S5, further details in methods). While the majority of gene trees across the G-locus support the trans-species polymorphism for the four *Carya* assemblies, we also note that the three genes at the left border (including *EMS1*) instead support the species phylogeny for these four taxa (Fig. 3F). Furthermore, while G-locus heterozygotes in other North American *Carya* species are heterozygous at pecan SNPs within these genes, the two G-locus heterozygotes in the East Asian clade are not heterozygous at any of these SNPs (Fig. 3D). These observations suggest the boundary of recombination suppression at the G-locus may have changed since the divergence of the North American and East Asian *Carya*, perhaps after the appearance of genetic variation in neighboring genes with antagonistic fitness effects between the two dichogamy morphs.

Genetic systems for heterodichogamy in other genera   Our estimate of the age of the *Carya* G-locus haplotypes suggests that we do not expect to find these haplotypes segregating in other known heterodichogamous genera within Juglandaceae - *Cyclocarya Platycarya*, and likely *Pterocarya* - which are all more closely related to *Juglans* than to *Carya*. Nonetheless, we considered the possibility that a subset of the pecan G-locus variation could be ancestral across heterodichogamous genera, or that there has been convergent use of

the same region. Therefore, we examined heterozygosity at the region syntenic with the *Carya* G-locus in a sample of 13 *Pterocarya* and 12 diploid *Cyclocarya* (including protandrous and protogynous individuals, Qu *et al.* 2023), but we found no evidence of increased polymorphism in this region comparable to the patterns seen in *Carya* (Fig. S16). As our analyses also indicate that heterodichogamy in *Cyclocarya/Pterocarya* is not controlled by the *Juglans* G-locus (Fig. 2C, S8, S17), this suggests the intriguing possibility that additional genetic systems controlling heterodichogamy in other Juglandaceae genera remain undiscovered.

Discussion

The genetic mechanisms underlying heterodichogamy appear to be more diverse than previously appreciated. Within Juglandaceae, two distinct ancient structural variants underlie this mating polymorphism in different genera. The identification of these loci opens further opportunities to study the ecology and the molecular and cellular mechanisms of this mating system with potential agricultural benefits in these important crop species. Our data cannot rule out the possibility that these genetic systems originated independently through the convergent evolution of heterodichogamy. However, given the clade-wide presence of heterodichogamy Juglandaceae, it seems plausible that a heterodichogamous mating system evolved once in the ancestor and the genetic control has been subject to turnover. Similar dynamics have been discovered in sex determination systems as new data and experiments improve both the phylogenetic scale and resolution of sex-chromosome turnover (Myosho *et al.* 2012; Jeffries *et al.* 2018; Müller *et al.* 2020; Hu *et al.* 2023). As heterodichogamy unites concepts of inbreeding avoidance, sexual interference, and sex-ratio selection, these systems may offer an important complement for testing theories on the evolution of the control and turnover of sexual systems (Sargent *et al.* 2006; Van Doorn and Kirkpatrick 2007; Kozielska *et al.* 2010; Blaser *et al.* 2013; Saunders *et al.* 2018). In summary, the evolution of heterodichogamy within Juglandaceae showcases both dynamic evolution and remarkable stability.

Acknowledgements We are grateful to UC Davis Putah Creek Riparian Reserve, Gene Cripe of Linwood Nursery, CA, USDA Wolfskill Experimental Orchard, Sonoma Botanical Garden, and UC Botanical Garden at Berkeley for access to leaf tissue from natural and cultivated collections of Juglandaceae. We thank Chuck Leslie for providing phenotype records for trees from the USDA germplasm database. We thank Jeff Ross-Ibarra, Quentin Cronk, the Dandekar lab at UC Davis, and the Coop lab for helpful discussions. Funding was provided by the United States Department of Agriculture (NIFA SCRI-Award no. 2012-51181-20027 awarded to PJB, CHL), the National Institutes of Health (NIH R35 GM136290 awarded to GC), and the National Science Foundation (NSF DISES 2307175 to GC, and NSF 1650042 awarded to JSG).

Materials and Methods

Phenotyping We phenotyped 81 individuals of a naturally-occuring population of *J. hindsii* from the UC Davis Putah Creek Riparian Reserve, along a ~2 mile creek-side path, in spring of 2022 and 2023. Dichogamy type was ascertained visually based on the relative developmental stages of male and female flowers. To validate our assessment of flowering phenotype we measured the length and width of 1-3 catkins of each tree on the same day and plotted the size distribution of catkins in the sample (Fig. 1A). We similarly obtained dichogamy phenotypes for individuals of *J. ailantifolia*, *J. californica*, *J. cathayensis*, *J. cinerea*, and *J. major* from trees at USDA Wolfskill Experimental Orchard in spring 2023. Phenotypes of newly sequenced *C. illinoinensis* from UC Davis orchards were also scored. Dichogamy phenotypes were available for 26 previously sequenced individuals of *J. regia* from UC Davis walnut breeding program records (Stevens *et al.* 2018). Flowering phenotypes for *J. microcarpa* and *J. hindsii* trees that were previously sequenced in Stevens *et al.* (2018) were obtained from the USDA Wolfskill Experimental Orchard database (Chuck Leslie, personal communication). Phenotypes of previously sequenced *J. nigra* individuals (Stevens *et al.* 2018) were obtained from Chatwin *et al.* (2023). Phenotypes of previously sequenced varieties of pecan (Xiao *et al.* 2021) were obtained from Bentley *et al.* (2019). Phenotypes of newly sequenced pecan were scored visually in spring of 2023 or otherwise obtained from Bentley *et al.* (2019) and the USDA database (https://cgru.usda.gov/carya/pecans/).

We measured the length of dormant catkin buds in 30 *J. regia* individuals on July 20, 2023. Measurements were made blind with respect to phenotype with the exception of 3 varieties which were known a priori. We took an average measurement of 6-12 of the largest easily accessible catkins using hand calipers. We tested for a difference in length using a linear model with leafing date as a covariate (Fig. S5).

Genomic sequencing and data curation We generated whole-genome resequencing data for 44 *J. hindsii*; 2 individuals each from *J. ailantifolia*, *J. californica*, *J. cathayensis*, *J. cinerea*, *J. major*; 19 *C. illinoinensis*; 1 *C. ovata*; 13 *Pterocarya stenoptera*; 2 *Pterocarya rhoifolia*; 1 *Pterocarya macroptera*; and 2 *Platycarya strobilaceae* (Table S1). Samples of *J. hindsii* were a subset of trees phenotyped from the UC Davis Putah Creek Riparian Reserve, CA. Other sequenced *Juglans* trees were from USDA Wolfskill Experimental Orchard. Samples of *Pterocarya stenoptera* were obtained from Wolfskill and UC Davis campus. Samples of *Carya illinoinensis* were obtained from the UC Davis orchards and the original orchard of Linwood Nursery in Turlock, California. Leaf tissue was flash frozen in liquid nitrogen and preserved at -80 degrees Celsius. DNA was extracted using the Qiagen DNeasy Plant Pro Kit.

We obtained published whole-gene resequencing data from 87 individuals of *J. regia* from Ding *et al.* (2022); Ji *et al.* (2021), as well as 26 *J. regia*, 11 *J. hindsii*, 12 *J. microcarpa*, and 13 *J. nigra* from Stevens *et al.* (2018). Published *Cyclocarya* resequencing data were obtained from Qu *et al.* (2023). We used publicly available long-read genome assemblies of plants within *Fagales* generated by Zhu *et al.* (2019); Marrano *et al.* (2020); Fitz-Gibbon *et al.* (2023); Zhang *et al.* (2021); Lovell *et al.* (2021); Zhang *et al.* (2023); Ding *et al.* (2023); Ning *et al.* (2020); Li *et al.* (2022); Zhang *et al.* (2020); Sork *et al.* (2022); Qu *et al.* (2023); Cao *et al.* (2023); Guzman-Torres *et al.* (2023).

We accessed whole-genome resequence data for 34 varieties of *C. illinoinensis* from Xiao *et al.* (2021). We verified the variety identity for 12 of these samples by comparison of genetic relatedness at overlapping sets of SNPs to reduced representation sequence data from 83 varieties of cultivated pecan from Bentley *et al.* (2019). The remainder of these samples were not used in the analysis, as several were labelled as varieties that did not match their predicted identity using data from Bentley *et al.* (2019). We chose the data from Bentley *et al.* (2019) as the standard as dichogamy type was rigorously documented for the trees in this data set.

Sequence alignment and variant calling We mapped short read sequencing libraries to available long-read reference genomes using *bwa* 0.7.17 (Li and Durbin 2009) with default parameters. To examine structural differences between G-locus haplotyes, we used BLAST (Altschul *et al.* 1990) to identify and align syntenic regions containing *TPPD-1* orthologs. We used both minimap2 (Li 2018) and Anchorwave (Song *et al.* 2022) to align entire chromosomes and G-locus regions for variant calling and divergence calculations. In one analysis, we aligned all available genomes to the Walnut 2.0 (*g* haplotype) (Fig. S6). For divergence calculations, we aligned genomes containing alternate G-locus haplotypes within species for *J. regia*, *J. californica*, and *J. mandschurica* using Anchorwave. As Anchorwave uses a genome annotation in the alignment, we used *liftoff* (Shumate and Salzberg 2021) to port the Walnut 2.0 annotation onto assemblies lacking an annotation. Variant calling was done with bcftools 1.17 (Danecek *et al.* 2021). Variants were filtered in *vcftools* 0.1.16 (Danecek *et al.* 2011) for minimum GQ 30, and minimum depth 10. Additional filters were applied on a per-analysis basis.

GWAS and linkage disequilibrium We performed GWAS for dichogamy type separately in 44 individuals of *J. hindsii*, 26 *J. regia* individuals, and 30 *Carya illinoinensis* individuals. GWAS was done in GEMMA 0.98.3 (Zhou and Stephens 2012), which controls for genome-wide relatedness among samples. We calculated genotypic LD as the r^2 value between genotypes at pairs of loci using *vcftools*, filtering sites for a minimum distance of 100bp.

Phylogenetic analyses and synteny To construct a phylogeny of *GJ1* repeats and their homologs, we extracted the genomic coordinates of individual repeat subunits from pairwise BLAST alignments. We aligned subunits a, b, and c separately using *muscle* (Edgar 2004) and then concatenated alignments. We constructed a maximum-likelihood phylogeny using *IQ-Tree* (Nguyen *et al.* 2015) and obtained node support values using the program's ultrafast bootstrap approximation algorithm. We constructed a species phylogeny

for *Carya* genome assemblies and *J. regia*, from concatenated alignments of 12,101 single copy orthologs identified in OrthoFinder (Emms and Kelly 2019). The phylogeny of the *Carya* G-locus was inferred similarly using just the 20 genes within the G-locus, as well as for each gene individually. We inferred and visualized synteny across the *Carya* G-locus using MCscan (Tang *et al.* 2008) which leverages collinearity of orthologs.

While G-locus haplotypes show strong conservation of synteny, we note a small number of predicted genes that are not shared between haplotypes (see Fig. 3F). None of these was annotated independently in two assemblies. However, we investigated whether homologous sequence in other assemblies may have been missed by the annotation pipeline. We therefore used BLAST to search for sequence homologous to these uniquely annotated genes in other *Carya* G-locus assemblies (Table S5). For two genes that were uniquely annotated in the *C. sinensis* assembly, we found significant BLAST hits within syntenic regions of the 'Lakota' primary assembly (Fig. 3F), but not within G-locus regions of the other assemblies (although we find hits elsewhere in the assemblies). This result suggests that these are ancient duplications, perhaps with conserved function. However, further validation of these predicted genes with gene expression and additional de novo assemblies is needed to fully address the role of gene duplications in the assembly of G-locus haplotypes.

Nucleotide divergence and heterozygosity   We calculated nucleotide divergence across the region encompassing the *Juglans* G-locus in 500bp windows from the genome alignments using custom R scripts. As a estimate of the age haplotype divergence, we took the average of the maximum divergence value within any 500bp window on either side of the G-locus indels within 4kb. We adjusted the values of Dxy for multiple hits using the Jukes and Cantor (1969) distance. We then calculated the divergence time as $T = Dxy/(2\mu)$, using a value of 1.5×10^{-9} per bp per year as an estimate of the substitution rate (Ding *et al.* 2023). We note there is considerable uncertainty about the substitution rate, and hence about this estimated time. Furthermore, we found evidence of recombination between haplotypes within the *TPPD-1* sequence over deep timescales (Fig. S4), suggesting that the genealogies of these linked regions in contemporary G-locus haplotypes may differ from the genealogy of the causal variants. As such we approach this estimate with caution, but note that it is consistent with trans-specific SNPs across *Juglans*, the inferred *GJ1* phylogeny, and read depth analyses in supporting haplotype divergence in the common ancestor of *Juglans*. We used *pixy* (Korunes and Samuk 2021) to calculate divergence between haplotypes from short-read alignments. To calculate divergence between pecan G-locus haplotypes, we calculated Dxy in coding regions between samples of *GG* and *gg* homozygotes, as well as heterozygosity in the same regions for *Gg* heterozygotes, and averaged these values. Pixy was also used to calculate heterozygosity in other *Carya* species at pecan G-locus SNPs, and to calculate heterozygosity for *Pterocarya* and *Cyclocarya* individuals across regions syntenic with *Juglans* and *Carya* G-loci. We used the R package *ape* (Paradis and Schliep 2019) to calculate divergence at fourfold degenerate sites within aligned coding sequences from the *Carya* G-locus. To estimate a divergence time we used the ratio of the Dxy between *Carya* haplotypes and from a *Carya* haplotype to *J. regia*, we adjusted both for multiple hits using Jukes and Cantor (1969).

Polymorphism within haplotypes and tests of selection   To test for an excess of nonsynonymous fixed differences between haplotypes relative to polymorphism (or conversely, an excess of nonsynonymous polymorphism), we performed a McDonald-Kreitman test for *TPPD-1* coding sequence (McDonald and Kreitman 1991) in a sample of 113 *J. regia*. To obtain data partitioned by haplotype, variants were first filtered for a minor allele count of 2 and then phased haplotypes using Beagle 5.4 (Browning *et al.* 2021). Phasing results across *TPPD-1* were manually inspected for errors, and we found that it resulted in haplotypes that produced a bimodal distribution of non-reference alleles with zero overlap. Moreover, phasing of heterozygotes for G-locus haplotypes (assigned on the basis of read depth) resulted in clear separation into haplotypes resembling those homozygotes across *TPPD-1* with no evidence of phase switches. We assigned haplotype identity for phased haplotypes from heterozygotes based on similarity to haplotypes from either homozygote. We used a custom R script to calculate nonsynonymous and synonymous polymorphism and divergence in the sample, and performed the MK test using Fisher's exact test. We applied a similar approach in *Carya*, first phasing all SNPs within coding regions across the *Carya* G-locus genes. Here, we observed erroneous phase switches near the end of the region in a small subset of heterozygotes, which we removed from the analysis. Variants were considered polymorphic within a haplotype group only if they were present within at least 2 copies of that haplotype.

PacBio IsoSeq   Tissues were collected from various locations at the University of California in Davis and the USDA's National Clonal Germplasm Repository in Winters, CA. Collected tissue was wrapped in foil and immediately immersed in liquid nitrogen in the field to preserve RNA quality. Frozen tissues were subsequently pulverized in liquid nitrogen in a mortar and pestle for extraction.

The extraction buffer used was 4M guanidine isothiocyanate, 0.2 M sodium acetate pH 5.0, 2mM EDTA, 2.5% (w/v) PVP-40. To this we added 400 uL Lysis buffer to 100 mg tissue and homogenize with pestle for 30 seconds, then 600 uL Lysis buffer (1 mL total)was added and vortexed for 20 seconds. From this, 500 uL of the homogenate was processed using the RNeasy Plant Mini kit (Qiagen) according to manufacturer's protocol (centrifugations steps were performed at 12,000 rpm for 30 seconds). RNA was eluted with 50 uL nuclease-free water at Step 9 in manufacturer's protocol. This was followed by DNase digestions with Turbo DNA-free kit (Invitrogen) according to manufacturer's protocols. Finally, RNA samples were cleaned up using HighPrep RNA Elite beads (MagBio Genomics) according to the manufacturer's 96 well format protocol for 10 uL reaction volume. Final elution was performed with 20 uL of nuclease-free water that was heated at 60C for 10 minutes before adding to sample.

For quality control and quantitation, samples were subsequently checked for purity on Nanodrop, quantified on Qubit, and checked on Bioanalyzer.

SMRTbell libraries were constructed and sequenced at the University of California Davis Genome center. Sequencing was performed on the Pacbio Sequel II. Demultiplexing and post processing of sequence data to create high quality full length non-chimera consensus transcripts (FLNC) was performed using the PacBio bioinformatics pipeline (ccs v4.2.0, lima v1.11.0, isoseq v3).

Transposable Element Annotation   We performed de-novo whole genome annotation of transposable elements (TEs) for long-read assemblies containing both G-locus haplotypes for *J. regia*, *J. californica*, and *J. mandshurica* using EDTA (Ou *et al.* 2019), and compared coordinates of annotated TEs to the coordinates of the derived *gJ3* insertion. EDTA predicted the presence of a CACTA-like DNA transposon at the coordinates of *gJ3*. We separately identified the presence of terminal inverted repeats near the *gJ3* insertion endpoints and evidence of target site duplication.

We used EDTA to annotate assemblies of 'Pawnee' and 'Lakota' pecan as well as assemblies of *Carya sinensis* and *C. cathayensis*. We computed the proportion of sequence covered by predicted TEs for three categories - (1) within the G-locus (defined by endpoints of genes falling inside the pecan GWAS peak, (2) in 300kb of sequence surrounding the G-locus (150kb on either side), and (3) across the whole genome.

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. Journal of Molecular Biology. 215:403–410.

Bachtrog D. 2013. Y-chromosome evolution: Emerging insights into processes of Y-chromosome degeneration. Nature Reviews Genetics. 14:113–124.

Bai WN, Zeng YF, Zhang DY. 2007. Mating patterns and pollen dispersal in a heterodichogamous tree, *Juglans mandshurica* (Juglandaceae). New Phytologist. 176:699–707.

Barrett SC. 2010. Darwin's legacy: the forms, function and sexual diversity of flowers. Philosophical Transactions of the Royal Society B: Biological Sciences. 365:351–368.

Bentley N, Grauke L, Klein P. 2019. Genotyping by sequencing (GBS) and SNP marker analysis of diverse accessions of pecan (*Carya illinoinensis*). Tree Genetics & Genomes. 15:1–17.

Bernard A, Marrano A, Donkpegan A, Brown PJ, Leslie CA, Neale DB, Lheureux F, Dirlewanger E. 2020. Association and linkage mapping to unravel genetic architecture of phenological traits and lateral bearing in Persian walnut (*Juglans regia* L.). BMC genomics. 21:1–25.

Blaser O, Grossen C, Neuenschwander S, Perrin N. 2013. Sex-chromosome turnovers induced by deleterious mutation load. Evolution. 67:635–645.

Browning BL, Tian X, Zhou Y, Browning SR. 2021. Fast two-stage phasing of large-scale sequence data. The American Journal of Human Genetics. 108:1880–1890.

Cao Y, Almeida-Silva F, Zhang WP, Ding YM, Bai D, Bai WN, Zhang BW, Van de Peer Y, Zhang DY. 2023. Genomic insights into adaptation to karst limestone and incipient speciation in East Asian *Platycarya* spp. (Juglandaceae). Molecular Biology and Evolution. 40:msad121.

Chakraborty S, Britton M, Martínez-García P, Dandekar AM. 2016. Deep RNA-Seq profile reveals biodiversity, plant–microbe interactions and a large family of NBS-LRR resistance genes in walnut (*Juglans regia*) tissues. AMB Express. 6:1–13.

Charlesworth B, Charlesworth D. 2000. The degeneration of Y chromosomes. Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences. 355:1563–1572.

Charlesworth D. 2016. The status of supergenes in the 21st century: recombination suppression in b atesian mimicry and sex chromosomes and other complex adaptations. Evolutionary Applications. 9:74–90.

Charlesworth D, Charlesworth B. 1979. The evolutionary genetics of sexual systems in flowering plants. Proceedings of the Royal Society of London. Series B. Biological Sciences. 205:513–530.

Chatwin W, Shirley D, Lopez J, Sarro J, Carlson J, Devault A, Pfrender M, Revord R, Coggeshall M, Romero-Severson J. 2023. Female flowers first: QTL mapping in eastern black walnut (*Juglans nigra* L.) identifies a dominant locus for heterodichogamy syntenic with that in Persian walnut (*J. regia* L.). Tree Genetics & Genomes. 19:4.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST *et al*. 2011. The variant call format and VCFtools. Bioinformatics. 27:2156–2158.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy SA, Davies RM *et al*. 2021. Twelve years of SAMtools and BCFtools. GigaScience. 10:giab008.

Darwin C. 1876. *The Effects of Cross and Self Fertilisation in the Vegetable Kingdom*. John Murray.

Darwin C. 1877. *The Different Forms of Flowers on Plants of the Same Species*. John Murray.

Delpino F. 1874. Ulteriori osservazioni e considerazioni sulla dicogamia nel regno vegetale. Appendice. Dimorfismo nel noce (*Juglans regia*) e pleiontismo nelle piante. Atti della Societa Italiana Scientia Naturale. 17:402–407.

Ding YM, Cao Y, Zhang WP, Chen J, Liu J, Li P, Renner SS, Zhang DY, Bai WN. 2022. Population-genomic analyses reveal bottlenecks and asymmetric introgression from Persian into iron walnut during domestication. Genome Biology. 23:1–18.

Ding YM, Pang XX, Cao Y, Zhang WP, Renner SS, Zhang DY, Bai WN. 2023. Genome structure-based Juglandaceae phylogenies contradict alignment-based phylogenies and substitution rates vary with DNA repair genes. Nature Communications. 14:617.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Research. 32:1792–1797.

Emms DM, Kelly S. 2019. Orthofinder: phylogenetic orthology inference for comparative genomics. Genome Biology. 20:1–14.

Endress PK. 2020. Structural and temporal modes of heterodichogamy and similar patterns across angiosperms. Botanical Journal of the Linnean Society. 193:5–18.

Fitz-Gibbon S, Mead A, O'Donnell S, Li ZZ, Escalona M, Beraut E, Sacco S, Marimuthu MP, Nguyen O, Sork VL. 2023. Reference genome of California walnut, *Juglans californica*, and resemblance with other genomes in the order Fagales. Journal of Heredity. esad036.

Fukuhara T, Tokumaru Si. 2014. Inflorescence dimorphism, heterodichogamy and thrips pollination in *Platycarya strobilacea* (Juglandaceae). Annals of Botany. 113:467–476.

Gleeson SK. 1982. Heterodichogamy in walnuts: inheritance and stable ratios. Evolution. pp. 892–902.

Gutiérrez-Valencia J, Hughes PW, Berdan EL, Slotte T. 2021. The genomic architecture and evolutionary fates of supergenes. Genome Biology and Evolution. 13:evab057.

Guzman-Torres CR, Trybulec E, LeVasseur H, Akella H, Amee M, Strickland E, Pauloski N, Williams M, Romero-Severson J, Hoban S *et al*. 2023. Conserving a threatened North American walnut: a chromosome-scale reference genome for butternut (*Juglans cinerea*). bioRxiv. doi:10.1101/2023.05.12.539246.

Hu N, Sanderson BJ, Guo M, Feng G, Gambhir D, Hale H, Wang D, Hyden B, Liu J, Smart LB *et al*. 2023. Evolution of a ZW sex chromosome system in willows. Nature Communications. 14:7144.

Huang Y, Xiao L, Zhang Z, Zhang R, Wang Z, Huang C, Huang R, Luan Y, Fan T, Wang J *et al*. 2019. The genomes of pecan and chinese hickory provide insights into carya evolution and nut nutrition. GigaScience. 8:giz036.

Jeffries DL, Lavanchy G, Sermier R, Sredl MJ, Miura I, Borzée A, Barrow LN, Canestrelli D, Crochet PA, Dufresnes C *et al*. 2018. A rapid rate of sex-chromosome turnover and non-random transitions in true frogs. Nature Communications. 9:4088.

Ji F, Ma Q, Zhang W, Liu J, Feng Y, Zhao P, Song X, Chen J, Zhang J, Wei X *et al*. 2021. A genome variation map provides insights into the genetics of walnut adaptation and agronomic traits. Genome Biology. 22:1–22.

Jukes TH, Cantor CR. 1969. Evolution of protein molecules. Mammalian Protein Metabolism. 3:21–132.

Korunes KL, Samuk K. 2021. pixy: Unbiased estimation of nucleotide diversity and divergence in the presence of missing data. Molecular ecology resources. 21:1359–1368.

Kozielska M, Weissing F, Beukeboom L, Pen I. 2010. Segregation distortion and the evolution of sex-determining mechanisms. Heredity. 104:100–112.

Larson-Johnson K. 2016. Phylogenetic investigation of the complex evolutionary history of dispersal mode and diversification rates across living and fossil Fagales. New Phytologist. 209:418–435.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 34:3094–3100.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler transform. Bioinformatics. 25:1754–1760.

Li X, Cai K, Zhang Q, Pei X, Chen S, Jiang L, Han Z, Zhao M, Li Y, Zhang X *et al*. 2022. The Manchurian walnut genome: insights into juglone and lipid biosynthesis. GigaScience. 11:giac057.

Lovell JT, Bentley NB, Bhattarai G, Jenkins JW, Sreedasyam A, Alarcon Y, Bock C, Boston LB, Carlson J, Cervantes K *et al*. 2021. Four chromosome scale genomes and a pan-genome annotation to accelerate pecan tree breeding. Nature Communications. 12:4125.

Luza JG, Polito VS. 1988. Microsporogenesis and anther differentiation in *Juglans regia* L.: A developmental basis for heterodichogamy in walnut. Botanical Gazette. 149:30–36.

Manchester SR. 1987. The fossil history of the Juglandaceae. Monographs in Systematic Botany from the Missouri Botanical Garden. 21:1–137.

Manos PS, Soltis PS, Soltis DE, Manchester SR, Oh SH, Bell CD, Dilcher DL, Stone DE. 2007. Phylogeny of extant and fossil Juglandaceae inferred from the integration of molecular and morphological data sets. Systematic Biology. 56:412–430.

Mao X, Fu XX, Huang P, Chen XL, Qu YQ. 2019. Heterodichogamy, pollen viability, and seed set in a population of polyploidy *Cyclocarya paliurus* (Batal) Iljinskaja (Juglandaceae). Forests. 10:347.

Marrano A, Britton M, Zaini PA, Zimin AV, Workman RE, Puiu D, Bianco L, Pierro EAD, Allen BJ, Chakraborty S *et al*. 2020. High-quality chromosome-scale assembly of the walnut (*Juglans regia* L.) reference genome. GigaScience. 9:giaa050.

McDonald JH, Kreitman M. 1991. Adaptive protein evolution at the Adh locus in Drosophila. Nature. 351:652–654.

Mu XY, Tong L, Sun M, Zhu YX, Wen J, Lin QW, Liu B. 2020. Phylogeny and divergence time estimation of the walnut family (Juglandaceae) based on nuclear RAD-Seq and chloroplast genome data. Molecular Phylogenetics and Evolution. 147:106802.

Müller NA, Kersten B, Leite Montalvão AP, Mähler N, Bernhardsson C, Bräutigam K, Carracedo Lorenzo Z, Hoenicka H, Kumar V, Mader M *et al*. 2020. A single gene underlies the dynamic evolution of poplar sex determination. Nature Plants. 6:630–637.

Myosho T, Otake H, Masuyama H, Matsuda M, Kuroki Y, Fujiyama A, Naruse K, Hamaguchi S, Sakaizumi M. 2012. Tracing the emergence of a novel sex-determining gene in medaka, oryzias luzonensis. Genetics. 191:163–170.

Nacken WK, Huijser P, Beltran JP, Saedler H, Sommer H. 1991. Molecular characterization of two stamen-specific genes, *tap1* and *fil1*, that are expressed in the wild type, but not in the *deficiens* mutant of *Antirrhinum majus*. Molecular and General Genetics MGG. 229:129–136.

Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular Biology and Evolution. 32:268–274.

Ning DL, Wu T, Xiao LJ, Ma T, Fang WL, Dong RQ, Cao FL. 2020. Chromosomal-level assembly of *Juglans sigillata* genome using Nanopore, BioNano, and Hi-C analysis. GigaScience. 9:giaa006.

Ou S, Su W, Liao Y, Chougule K, Agda JR, Hellinga AJ, Lugo CSB, Elliott TA, Ware D, Peterson T *et al*. 2019. Benchmarking transposable element annotation methods for creation of a streamlined, comprehensive pipeline. Genome Biology. 20:1–18.

Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. Bioinformatics. 35:526–528.

Polito V, Pinney K. 1997. The relationship between phenology of pistillate flower organogenesis and mode of heterodichogamy in *Juglans regia* L.(Juglandaceae). Sexual Plant Reproduction. 10:36–39.

Ponnu J, Wahl V, Schmid M. 2011. Trehalose-6-phosphate: connecting plant metabolism and development. Frontiers in Plant Science. 2:70.

Pringle C. 1879. Dimorpho-dichogamy in *Juglans cinerea*, L. Botanical Gazette. 4:237–237.

Qu Y, Shang X, Fang S, Zhang X, Fu X. 2023. Genome assembly of two diploid and one auto-tetraploid *Cyclocarya paliurus* genomes. Scientific Data. 10:507.

Renner SS. 2001. How common is heterodichogamy? Trends in Ecology & Evolution. 16:595–597.

Rhein HS, Sreedasyam A, Cooke P, Velasco-Cruz C, Grimwood J, Schmutz J, Jenkins J, Kumar S, Song M, Heerema RJ *et al*. 2023. Comparative transcriptome analyses reveal insights into catkin bloom patterns in pecan protogynous and protandrous cultivars. PloS one. 18:e0281805.

Sargent RD, Mandegar MA, Otto SP. 2006. A model of the evolution of dichogamy incorporating sex-ratio selection, anther-stigma interference, and inbreeding depression. Evolution. 60:934–944.

Satoh-Nagasawa N, Nagasawa N, Malcomber S, Sakai H, Jackson D. 2006. A trehalose metabolic enzyme controls inflorescence architecture in maize. Nature. 441:227–230.

Saunders PA, Neuenschwander S, Perrin N. 2018. Sex chromosome turnovers and genetic drift: a simulation study. Journal of Evolutionary Biology. 31:1413–1419.

Shumate A, Salzberg SL. 2021. Liftoff: accurate mapping of gene annotations. Bioinformatics. 37:1639–1643.

Song B, Marco-Sola S, Moreto M, Johnson L, Buckler ES, Stitzer MC. 2022. AnchorWave: Sensitive alignment of genomes with high sequence diversity, extensive structural polymorphism, and whole-genome duplication. Proceedings of the National Academy of Sciences. 119:e2113075119.

Sork VL, Cokus SJ, Fitz-Gibbon ST, Zimin AV, Puiu D, Garcia JA, Gugger PF, Henriquez CL, Zhen Y, Lohmueller KE *et al.* 2022. High-quality genome and methylomes illustrate features underlying evolutionary success of oaks. Nature Communications. 13:2047.

Stevens KA, Woeste K, Chakraborty S, Crepeau MW, Leslie CA, Martínez-García PJ, Puiu D, Romero-Severson J, Coggeshall M, Dandekar AM *et al.* 2018. Genomic variation among and within six *Juglans* species. G3: Genes, Genomes, Genetics. 8:2153–2165.

Stuckey HP. 1915. The two groups of varieties of the Hicora pecan and their relation to self-sterility. Georgia Experiment Station. Bulletin No. 124.

Tang H, Wang X, Bowers JE, Ming R, Alam M, Paterson AH. 2008. Unraveling ancient hexaploidy through multiply-aligned angiosperm gene maps. Genome research. 18:1944–1954.

Thompson T, Romberg L. 1985. Inheritance of heterodichogamy in pecan. Journal of Heredity. 76:456–458.

Van Doorn G, Kirkpatrick M. 2007. Turnover of sex chromosomes induced by sexual conflict. Nature. 449:909–912.

Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, Feil R, Lunn JE, Stitt M, Schmid M. 2013. Regulation of flowering by Trehalose-6-Phosphate signaling in *Arabidopsis thaliana*. Science. 339:704–707.

Xiao L, Yu M, Zhang Y, Hu J, Zhang R, Wang J, Guo H, Zhang H, Guo X, Deng T *et al.* 2021. Chromosome-scale assembly reveals asymmetric paleo-subgenome evolution and targets for the acceleration of fungal resistance breeding in the nut crop, pecan. Plant Communications. 2:100247.

Zhang J, Zhang W, Ji F, Qiu J, Song X, Bu D, Pan G, Ma Q, Chen J, Huang R *et al.* 2020. A high-quality walnut genome assembly reveals extensive gene expression divergences after whole-genome duplication. Plant Biotechnology Journal. 18:1848.

Zhang JB, Li RQ, Xiang XG, Manchester SR, Lin L, Wang W, Wen J, Chen ZD. 2013. Integrated fossil and molecular data reveal the biogeographic diversification of the eastern Asian-eastern North American disjunct hickory genus (*Carya* Nutt.). PLoS one. 8:e70449.

Zhang Q, Ree RH, Salamin N, Xing Y, Silvestro D. 2022. Fossil-informed models reveal a boreotropical origin and divergent evolutionary trajectories in the walnut family (Juglandaceae). Systematic Biology. 71:242–258.

Zhang WP, Cao L, Lin XR, Ding YM, Liang Y, Zhang DY, Pang EL, Renner SS, Bai WN. 2021. Dead-end hybridization in walnut trees revealed by large-scale genomic sequence data. Molecular Biology and Evolution. 39:msab308.

Zhang WP, Ding YM, Cao Y, Li P, Yang Y, Pang XX, Bai WN, Zhang DY. 2023. Uncovering ghost introgression through genomic analysis of a distinct East Asian hickory species. bioRxiv. doi:10.1101/2023.06.26.546421.

Zhao DZ, Wang GF, Speal B, Ma H. 2002. The *EXCESS MICROSPOROCYTES1* gene encodes a putative leucine-rich repeat receptor protein kinase that controls somatic and reproductive cell fates in the *Arabidopsis* anther. Genes & Development. 16:2021–2031.

Zhou H, Hu Y, Ebrahimi A, Liu P, Woeste K, Zhao P, Zhang S. 2021. Whole genome based insights into the phylogeny and evolution of the Juglandaceae. BMC Ecology and Evolution. 21:1–16.

Zhou X, Stephens M. 2012. Genome-wide efficient mixed-model analysis for association studies. Nature genetics. 44:821–824.

Zhu T, Wang L, You FM, Rodriguez JC, Deal KR, Chen L, Li J, Chakraborty S, Balan B, Jiang CZ *et al*. 2019. Sequencing a *Juglans regia* × *J. microcarpa* hybrid yields high-quality genome assemblies of parental species. Horticulture Research. 6:55.

 # Supplementary Material

Figure S1: **A)** GWAS for dichogamy type in 26 *J. regia* individuals. **B)** Zoomed in region of strong association. Here, read mapping and variant calling was done using the Walnut 2.0 genome, an assembly from the 'Chandler' variety, a protandrous tree (*gg* genotype.) **C)** Normalized average read depth in 1kb windows for 26 *J. regia* individuals of known genotype across the Walnut 2.0 assembly (*g* haplotype). Black bars indicate the position of the (*gJ3*) indel and *gJ1-0* (See Fig. 2 in main text for notation of structural variants). The spike in coverage at *gJ1-0* seen for protogynous individuals corresponds to sequencing reads originating from repeats within the *GJ1* insertion in the *G* haplotype.

Figure S2: Correlation of diploid genotypes between loci (LD) in the region surrounding the G-locus in two species of *Juglans*. **Left)** LD in a sample of 44 from a natural population of *J. hindsii*. **Right)** LD in a sample of 26 *J. regia* from the UC Davis/USDA walnut breeding program.

Figure S3: Patterns of read depth from multiple species against two pseudohaploid assemblies of *J. californica* at the region syntenic with the GWAS hit in *J. regia* reveal the same segregating structural variant across multiple species in association with heterodichogamy flowering type. The gray bar indicates the location of the TPP ortholog in each assembly, and the arrow indicates direction of transcription. Read depth was normalized by the average coverage across an arbitrarily chosen separate chromosome. Protandrous individuals are known genotype *hh*, while the genotype of protogynous individuals is not known a priori without pedigree information. One homozygote for the protogynous allele is known in *J. regia* (variety 'Sharkey', Gleeson (1982)). Read depth for the *J. nigra* 'Hay' variety also indicate it is homozygous for the protogynous allele.

Figure S4: Genetic variation within the *TPPD-1* gene and its 3' region in long-read assemblies of *G* and *g* haplotypes from three *Juglans* species. The region displayed includes *TPPD-1*, it's 3'UTR (*GJ1a-0*), as well as *GJ1b-0* and *GJ1c-0* (see Fig. 2 in main text). Numbers indicate positions of nucleotides in the Walnut 2.0 reference genome. Singletons are omitted. Diagonal lines connect SNP genotypes to their locations within the gene model of *TPPD-1*. Below, rectangles indicate exons - gray rectangles represent UTRs, and brown rectangles represent CDS. SNPs that consistently group by haplotype are localized to the 3' UTR or further downstream, but within *TPPD-1* CDS SNPs quickly transition to grouping by species. The black vertical line indicates the border between the 3'UTR and the CDS of the last exon in the reference annotation.

Figure S5: Protandrous and protogynous morphs of *J. regia* differ in size of male catkin buds in the season prior to flowering (β = 2 mm, P<0.001). We fit a linear model predicting catkin bud length from dichogamy type with leafing date as a covariate. While leafing date was not a significant predictor, we note that the slope estimate is in the expected direction. Notably, the earliest leafing protogynous varieties have catkin buds roughly of similar size to the latest leafing protandrous varieties.

Figure S6: Nucleotide divergence of Juglandaceae genomes across the *g* haplotype of *J. regia* 'Chandler', calculated in 500bp windows from whole-genome alignments. Dotted gray lines show the average calculated over the entire aligned chromosome. Shaded gray interval indicates the top 95% quantile.

Figure S7: Gene expression levels of *TPPD-1* (LOC108984907) from published short read RNA-seq libraries of *J. regia* 'Chandler' (*gg* genotype). Data from (Chakraborty *et al.* 2016). Y-axis measures Fragments Per Kilobase of transcript per Million mapped reads.

Figure S8: Normalized average read depth in 1kb bp windows for multiple individuals of *Cyclocarya paliurus* and *Pterocarya stenoptera* against an assembly of the *Juglans G* haplotype from *J. californica*. Blue rectangle and arrow indicates the position of *TPPD-1* in the assembly.

Figure S9: Gene expression levels of the *J. regia* ortholog of *FIL1* in 20 transcriptomic libraries from different tissues. Results from the literature indicate that *FIL1*-homologs are expressed uniquely or predominantly in stamens Nacken *et al.* (1991). Consistent with this, we detected abundant expression of *FIL1* in catkins of *J. regia*, and no trace of expression in other tissues. Y-axis measures Fragments Per Kilobase of transcript per Million mapped reads.

Figure S10: Normalized average read depth of pecan varieties in 1kb windows across the pecan G-locus for two assemblies of the *g* haplotype. **Top)** 'Pawnee' is protandrous (*gg*) so the assembly is known as the *g* haplotype a priori. **Bottom)** The alternate assembly of 'Lakota' (*Gg*) shows the same coverage pattern as for 'Pawnee', which is reversed compared to the 'Lakota' primary assembly (see Fig 3C in main text)

.

Figure S11: Read depth of pecan varieties (*C. illinoinensis*) of against two East Asian *Carya* genome assemblies. Top: Patterns of differential read depth by G-locus genotype indicate that the assembly of *C. sinensis* is of the *G* haplotype. The G-locus region in this assembly is roughly 350kb in length, closer in length to the pecan *G* haplotype than to the pecan *g* haplotype. Vertical indicate the outermost G-locus genes as defined in pecan, and these correspond roughly to the span of differential coverage patterns. Bottom: Differential read depth by G-locus genotype against the *C. cathayensis* assembly indicate it is an assembly of the *g* haplotype. The G-locus region in this assembly is roughly 230kb, closer in length to the *g* haplotype of pecan.

.

Figure S12: Individual-level heterozygosity at a set of 454 SNPs in G-locus coding sequence that were ascertained as fixed differences between GG and gg genotypes in pecan. **Left)** Heterozygosity at these SNPs is very close to 100% for heterozygotes for the G-locus haplotypes, confirming that these vast majority of SNPs are indeed fixed differences. **Right)** Heterozygosity at the same set of SNPs for 16 individuals representing 15 species of *Carya* spanning the divergence between North American (black) and East Asian (gray) clades. Error bars show 95% confidence intervals for the proportion of heterozygous SNPs. Note two individuals of *C. ovata*, one of which appears to be a heterozygote for the G-locus haplotypes and the other a homozygote. The putative heterozygote in the East Asian clade, *C. kweichowensis*, is heterozygous at a lower proportion of SNPs, which would be explained by many of these SNPs having arisen after the divergence between East Asian and North American clades. Consistent with this, we note that the individual with the highest proportion of heterozygous SNPs is from *C. aquatica*, which is the sister species to *C. illinoinensis*.

Figure S13: Linkage disequilibrium is exceptionally high across the Carya G-locus, consistent with a lack of recombination between G and g haplotypes. Color indicates the strength of Pearson correlation between diploid genotypes at two loci. Black lines indicate the endpoints of genes that fall within the GWAS peak for dichogamy type in pecan. Shown for variant calls against the 'Pawnee' reference.

.

Figure S14: Transposable element (TE) content within *Carya* G-locus haplotypes. **(Left)** Number of TEs by category within the G-locus, in 150kb of flanking sequence on either end of the H-locus, and across the entire genome for two pecan (*C. illinoinensis*) assemblies containing alternate G-locus haplotypes. **(Right)** TEs within the G-locus for assemblies of haplotype within the East Asian clade. In this case, the assemblies are from different species, so for simplicity we omit showing background TE content outside of the G-locus, but note the relative increase in this region for *C. sinensis* compared to *C. cathayensis* diverges from the genome-wide average pattern.

Figure S15: Gene-level synteny for pecan G-locus haplotypes and *Quercus lobata*. Gray bands connect orthologous genes. Genes are colored by strand (blue - plus, green - minus). Shown for the 20 predicted genes that fall within the pecan GWAS peak for dichogamy type and are present in the annotation of both pecan assemblies.

Figure S16: **Topeft)** Heterozygosity in 1kb windows for pecan varieties of known genotype at the *Carya* G-locus. **Topright)** Heterozygosity in 1kb windows for 12 diploid *Cyclocarya paliurus* individuals across the region syntenic with the *Carya* G-locus. Two of these individuals are of known dichogamy type (Qu *et al.* 2023). **Bottomleft)** Heterozygosity in 1kb windows for 13 diploid *Pterocarya stenoptera* individuals across the region syntenic with the *Carya* G-locus. Lines are LOESS smoothing curves fit for each individual. Dotted lines show the endpoints of the outermost G-locus orthologous genes in each assembly.

.

Figure S17: **Topleft)** Heterozygosity in 500bp windows for Persian walnut varieties of known genotype at the *Juglans* G-locus. Note the clear separation of heterozygotes from homozygotes. Lines are LOESS smoothing curves fit for each individual. Blue rectangle indicates the location of *TPPD-1*, red rectangle indicates the location of a *NDR1/HIN1*-like gene. **Topright)** Heterozygosity in 500bp windows for 12 diploid *Cyclocarya paliurus* individuals across the region syntenic with the *Juglans* G-locus. Two of these individuals represent each dichogamy type (Qu *et al.* 2023). **Bottomleft)** Heterozygosity in 500bp windows for 13 *Pterocarya stenoptera* individuals across the region syntenic with the *Juglans* G-locus. Here, we note two individuals appear highly heterozygous to the left hand side of the *NDR1/HIN1*-like gene, but that individuals also show high heterozygosity in the corresponding region in *J. regia* that is not associated with dichogamy type. We also note that we did not detect G-locus structural variants in *P. stenoptera* and *C. paliurus* (Fig. S8), and our phylogeny inference (Fig. 2C) suggests we should not expect to find the *Juglans* G-locus structural variant segregating in either of these genera.