Multilayer Network Analysis across Cortical Depths in Resting-State 7T fMRI

Parker Kotlarz[1,2], Kaisu Lankinen[1,3], Maria Hakonen[1,3], Tori Turpin[4], Jonathan R. Polimeni[1,3,5],

and Jyrki Ahveninen[1,3]

[1]Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital,

Charlestown, MA, USA

[2]Harvard Medical School, Boston, MA, USA

[3]Department of Radiology, Harvard Medical School, Boston, MA, USA

[4]McLean Hospital, Boston, MA, USA

[5]Division of Health Sciences and Technology, Massachusetts Institute of Technology,

Cambridge, MA, USA

Abstract

In graph theory, "multilayer networks" represent systems involving several interconnected

topological levels. One example in neuroscience is the stratification of connections between

different cortical depths or "laminae", which is becoming non-invasively accessible in humans

using ultra-high-resolution functional MRI (fMRI). Here, we applied multilayer graph theory to

examine functional connectivity across different cortical depths in humans, using 7T fMRI (1-

mm^3 voxels; 30 participants). Blood oxygenation level dependent (BOLD) signals were derived

from five depths between the white matter and pial surface. We compared networks where the

inter-regional connections were limited to a single cortical depth only ("layer-by-layer matrices")

to those considering all possible connections between areas and cortical depths ("multilayer matrix"). We utilized global and local graph theory features that quantitatively characterize network attributes including network composition, nodal centrality, path-based measures, and hub segregation. Detecting functional differences between cortical depths was improved using multilayer connectomics compared to the layer-by-layer versions. Superficial depths of the cortex dominated information transfer and deeper depths drove clustering. These differences were largest in frontotemporal and limbic regions. fMRI functional connectivity across different cortical depths may contain neurophysiologically relevant information; thus, multilayer connectomics could provide a methodological framework for studies on how information flows across this stratification.

Keywords (up to 6): fMRI, laminar, connectomics, graph theory, multilayer network

Introduction

Investigating brain activity and function through network analyses has become an integral methodological foundation of neuroscience. Connectomics has yielded significant advances in understanding brain structure and function (Farahani et al., 2019; Milano et al., 2019). Modeling the brain as a system of nodes (brain regions) connected by edges (mathematical relationships)— often using graph theory—can be used to provide insight into brain characteristics and topological properties (Rubinov & Sporns, 2010). Brain networks can be derived from structural neuroimaging such as MRI or diffusion tensor imaging (DTI) (structural connectomics) (Griffa et al., 2013; Meoded et al., 2020; Yeh et al., 2021) or functional neuroimaging such as fMRI,

EEG, or magnetoencephalography (MEG) (functional connectomics) (De Vico Fallani et al.,

2014; Matthews & Hampshire, 2016; Sadaghiani et al., 2022; Smith et al., 2013; Xia & He,

2017). Both structural and functional connectomics have been used to understand disease models

(Benito-Leon et al., 2019; Fleischer et al., 2019; Jacob et al., 2020; Kotlarz et al., 2022), aid in

surgical mapping (Ahsan et al., 2020; Dadario et al., 2021; Gleichgerrcht et al., 2020; Hart et al.,

2016), and characterize therapeutic effects of neuropsychiatric treatments (Caeyenberghs et al.,

2017; Chen et al., 2020; Filippi et al., 2023; Lei et al., 2021; Tavakol et al., 2019; Yun & Kim,

2021).

An emerging field within connectomics, and more broadly graph theory, is the exploration of

multilayer networks (Boccaletti et al., 2014; Kivela et al., 2014). Multilayer networks are

composed of individual layers of networks with interconnecting edges between different layers.

Connections across layers can be solely between homologous nodes (multiplex) or between

nodes regardless of layer or nodal position (multilayer). Multilayer connectomics enables the

study of multifaceted and multimodal neuroimaging data, with the different groups of data

divided into distinct layers of the connectivity matrix (Betzel & Bassett, 2017; De Domenico,

2017; Vaiana & Muldoon, 2018). For example, multilayer networks can be derived using

correlations between different frequency bands of MEG recordings to identify the interplay

between frequencies (Buldu & Porter, 2018). Additionally, different modalities such as MEG,

fMRI, and diffusion MRI can be combined to identify patterns in brain processing (Breedt et al.,

2023) or pathological dysfunction (Casas-Roma et al., 2022) that were not found in traditional

single-layer analysis. Thus, multilayer connectomics allows for the incorporation of

multidimensional neuroimaging data and can identify relationships between distinct

neuroimaging techniques and analyses.

One potential application of multilayer connectomics is understanding the hierarchical

organization of the cerebral cortex. Neuroanatomical (Felleman & Van Essen, 1991; Rockland &

Pandya, 1979; Zeki, 2018) and electrophysiological (Schroeder & Foxe, 2002; Schroeder et al.,

2001) studies in animal models have identified that laminar input/output patterns can inform

about bottom-up (feedforward) or top-down (feedback) processes between cortical regions.

Despite its high resolution, a limitation of laminar electrophysiological recordings in

comprehensive connectivity analyses is that the coverage area is typically very small. At the

same time, mapping anatomical connections using fiber tracing has limited options for mapping

of the post-synaptic targets (Rockland, 2019). Intracortical analyses of high resolution fMRI data

have the benefit that the coverage can be extended to the entire cerebral cortex. Advancements in

high-resolution fMRI ($\leq 1\,mm^3$ voxel size) have enabled sampling of functional signals from

different depths of the cortical gray matter (Finn et al., 2019; L. Huber et al., 2021; Norris &

Polimeni, 2019; Polimeni et al., 2018). However, there are multiple challenges and unanswered

questions for the feasibility of using cortical depth profiles of fMRI signals (Norris & Polimeni,

2019). Because deoxygenated blood also drains up to the cortical surface through the

intracortical diving venules, fMRI voxels intersecting the superficial layers could also be

affected by deeper neuronal activations (Markuerkiaga et al., 2016b). Despite this limitation,

studies using this emerging methodology have attempted to identify feedback and feedforward

relationships non-invasively in the human brain (Chai et al., 2021; De Martino et al., 2015;

Fracasso et al., 2018; Gau et al., 2020; Klein et al., 2018; Kok et al., 2016; Lankinen et al., 2023;

Lawrence et al., 2019; Moerel et al., 2018; Moerel et al., 2019; Muckli et al., 2015; Wu et al.,

2018), akin to micro-scale recordings in animal models.

Even with the advent of high-resolution functional neuroimaging, multilayer connectomics have

mostly focused on anatomical networks derived from structural MRI and DTI (Shamir & Assaf,

2021; Shamir & Assaf, 2023) due to their direct relationship to cortical architecture. For

example, DTI and histological samples identified that cortical areas with similar laminar

structure were more likely to be connected (Wei et al., 2019). Additionally, even in functional

laminar studies, previous works have predominantly utilized task-based studies in pre-defined

brain regions (Chai et al., 2021; De Martino et al., 2015; Finn et al., 2019; Fracasso et al., 2018;

Gau et al., 2020; Klein et al., 2018; Kok et al., 2016; Lankinen et al., 2023; Lawrence et al.,

2019; Moerel et al., 2018; Moerel et al., 2019; Muckli et al., 2015; Polimeni et al., 2010; Wu et

al., 2018). Consequently, the functional components of the whole-brain cortical depths continue

to be underexplored.

In contrast to task-based studies, which primarily focus on specific cortical areas (Finn et al.,

2021), resting-state analysis enables whole-brain investigation of laminar organization (L. Huber

et al., 2021; L. R. Huber et al., 2021). These resting-state connections have been shown to reflect

anatomical connectivity (Adachi et al., 2012; Honey et al., 2009; Turk et al., 2016; van den

Heuvel et al., 2016) and task-based networks (Di et al., 2013; Hermundstad et al., 2013). Thus,

network differences within laminar resting-state fMRI networks represent functional differences

between cortical depths. This work explores the laminar structure of the cortex using high-

resolution resting-state fMRI and multilayer connectomics. We use a dual-pipeline approach in

comparing the information extracted from layer-by-layer vs. multilayer connectomics to test

whether there are connectivity differences between cortical depths. We demonstrate the validity

of multilayer functional laminar connectomics through showing that cortical depths have distinct

graph theory characteristics that are more clearly identifiable through multilayer connectomics

compared to the traditional single layer methodology.

Methods

Figure 1. Multilayer connectomic pipeline to analyze functional connectivity across different cortical depths. Here, "layers" refer to the dimensions of the connectivity matrices which represent fMRI signals gathered from different "cortical depths." **(A)** The cortex was uniformly divided into five surfaces at different depths, as seen above projected on a 0.75-mm isotropic-resolution anatomical T_1-weighted image. **(B)** The brain was parcellated into 148 regions-of-interest (ROIs) (74 per hemisphere) based on the Destrieux atlas in FreeSurfer (Destrieux et al., 2010; Fischl et al., 2004). The ROIs are shown on an inflated left-hemisphere cortical surface. **(C)** Schematic showing the difference between a layer-by-layer network and a multilayer network. In the layer-by-layer approach, each layer (network) is independent of other layers while in the multilayer approach, the layers are inter-connected. A sparser multilayer network is shown for visualization purposes. **(D)** Example matrix construction from both the layer-by-layer and multilayer approaches. While both approaches use matrices derived from Pearson correlations from the different layers, the multilayer approach generates a supra-adjacency matrix that also has correlations between different layers (shown in grayscale). **(E)** Example matrix construction for within-layer and between-layer matrices. For within-layer matrices, each sub-matrix is extracted individually for analysis. White areas represent connections excluded from the analysis.

Participants

Thirty healthy adults (mean age ± standard deviation = 32.4 ± 10 years, 15 women, all right-

handed) were recruited using an internal online recruiting platform. Participants were screened

for vision problems, hearing problems, cognition-altering medications, and exclusions for MRI

(metal in the body). Twenty-eight of the participants were native English-speakers. Informed

consent was obtained from all participants, and MRI safety screening forms were completed

before each scan. The study design, protocol, and consent were approved by the Mass General

Brigham Institutional Review Board.

Image Acquisition

Participants were measured in sets of 7.9-min resting-state fMRI scans occurring on different

days (3 to 4 sessions per participant). Twenty-three participants were measured in twelve resting-

state scans. Seven participants had between ten to eighteen resting-state scans (10 scans: $n = 1$;

11 scans: $n = 1$; 13 scans: $n = 2$; 14 scans: $n = 2$, 18 scans: $n = 1$) (**Figure S1**). The participants

were instructed to avoid movement during the scans and keep their eyes open and fixated on a

fixation cross projected on a screen viewed through a mirror. The average duration of the

sessions was around two hours. Breathing and heart rate were recorded using the built-in

Siemens system at a sampling rate of 400 Hz. Inhalation and exhalation were measured with the

Siemens respiratory-effort transducer attached to a respiratory belt. The heart rate was recorded

using Siemens photoplethysmogram transducers on the participant's index finger.

The functional and structural neuroimaging data was acquired using a 7T whole-body MRI

scanner (MAGNETOM Terra, Siemens, Erlangen, Germany) with a home-built custom-built 64-

channel array coil (Mareyam et al., 2020). To reduce participant head motion inside the scanner,

MRI-compatible paddings were placed around the head and neck. In each imaging session, T_1-

weighted anatomical images were measured using a 0.75-mm isotropic multi-echo MPRAGE

pulse sequence (van der Kouwe et al., 2008; Zaretskaya et al., 2018) with repetition time (TR) =

2530 ms, four echoes with echo time (TE) of 1.72, 3.53, 5.34, and 7.15 ms, 7° flip angle, 240 $\times$

240 mm^2 field of view (FoV), and 224 sagittal slices. To help with pial surface placement by

avoiding dura mater, T_2-weighted anatomical images (voxel size = 0.83 x 0.83 x 0.80 mm, TR =

9000 ms, TE = 269 ms, flip angle = 120°, FoV = 225 x 225 mm^2, 270 sagittal slices) were

acquired for twenty-eight out of thirty participants in one of the imaging sessions. Resting-state

functional imaging was collected using a T_2*-weighted blipped-CAIPI (Setsompop et al., 2012)

simultaneous multi-slice (SMS) echo planar imaging (EPI) sequence using multi-band RF pulses

(Setsompop et al., 2012) with 4$\times$ acceleration factor in phase-encoding direction, 3$\times$ acceleration

factor in slice-encoding direction, TR = 2800 ms, TE = 27.0 ms, isotropic 1-mm^3 voxels, 78° flip

angle, 192 $\times$ 192 mm^2 FoV, 132 axial slices, anterior-to-posterior phase encoding direction, 1446

Hz/pixel bandwidth, 0.82 ms nominal echo spacing, and fat suppression. In addition, to de-warp

the functional data, an EPI scan was collected with identical parameters but with an opposite

phase-encoding polarity (posterior-to-anterior, PA-EPI) relative to the functional scans. For four

participants with missing PA-EPI scans, the data were de-warped using a gradient-echo field

map (TR = 1040 ms, TE = 4.71 ms and 5.73 ms, isotropic 1.3-mm^3 voxels; flip angle = 75°, FoV

= 240 $\times$ 240 mm^2, 120 slices, bandwidth = 303 Hz/pixel).

MRI Preprocessing

First, SPM12 (http://www.fil.ion.ucl.ac.uk/spm/, [SPM12-spm_preproc_run.m]; bias field

correction, full-width at half-maximum, FWHM: 18 mm, sampling distance: 2 mm, bias

regularization: 1E–4) and customized MATLAB scripts were used to correct the bias field of the

structural T_1 and T_2 images. Next, recon-all of FreeSurfer 6.0 (Fischl, 2012) with an extension

for submillimeter 7 T data (Zaretskaya et al., 2018) was used to automatically create cortical

reconstructions for each participant. An average of multiple T_1-weighted anatomical volumes (3

to 4 per participant) alongside a T_2-weighted volume were used in the reconstruction to enhance

the quality of the cortical surfaces. Nine intermediate surfaces were created between the white

matter and pial surfaces with fixed relative distances, of which five were selected for the laminar

analysis (described below). Lastly, the surfaces generated by recon-all were corrected manually

for inaccuracies with Recon Edit of Freeview.

For the functional data, slice-timing and motion corrections were first implemented in FreeSurfer

7.1 (Fischl, 2012). De-warping was then used to correct for geometric distortions caused by

susceptibility-induced off-resonance fields. In de-warping, the off-resonance distortion field was

estimated using the functional data and the PA-EPI scan collected with reversed a phase-encode

blip; thus, the distortions are reversed in direction in respect to the scans [FreeSurfer: topup,

applytopup] (Andersson et al., 2003; Smith et al., 2004). For four participants that were missing

the PA-EPI scan used above, the distortion field was estimated using the B_0 field map scan in

FreeSurfer 6.0 [FreeSurfer-epidewarp]. The respiratory and heart rate artifacts were corrected

using the RETROspective Image CORrection (RETROICOR) algorithm (3[rd] order heart rate,

respiratory, and multiplicative terms) (Glover et al., 2000). Three participants were missing heart

rate data and, therefore, only respiratory recordings were used in RETROICOR. In addition,

RETROICOR was not applied to five participants with missing respiratory and heart rate data.

Functional data were then co-registered with the structural images using Boundary-Based

Registration in FreeSurfer 6.0 (Greve & Fischl, 2009). By projecting each intersecting voxel onto

the corresponding surface vertex using trilinear interpolation, the fMRI timeseries were then

resampled onto the pial and white matter surfaces, and nine cortical depths between them.

From the nine intracortical surfaces, five alternating depths were selected starting closest to pial

surface (depths 1 to 5, superficial to deep) (**Figure 1A**). The outside surfaces (pial and white

matter) were excluded to avoid partial volume effects from the cerebrospinal fluid and white

matter, respectively. Additionally, depths included were alternated to minimize potential for

partial volume overlap between surfaces that would bias the correlation matrix generation. To

measure this overlap, the distance from each voxel centroid (within cortical volume) to the

cortical surface (white matter/pial surface) was calculated. The relative distance was defined so

that the depth at pial surface was zero, and one at the white matter border. Next, voxels

intersecting each layer were picked and plotted with respect to their relative distances in a

histogram. **Figure S2** illustrates that taking every other layer limits the overlap between the

layers, and thus leaking of information to adjacent layers. Additionally, to explore if tSNR

impacted connectivity matrix generation, the average tSNR per each cortical depth was

calculated for each layer (**Figure S3**).

Matrix Generation and Processing

Two parallel matrix processing pipelines were used to generate individual independent adjacency

matrices for the layer-by-layer approach while creating one supra-adjacency matrix for the

multilayer approach (**Figures 1C and 1D**). The layer-by-layer approach creates an independent

network for each cortical depth while the multilayer approach results in five interconnected

networks that combines all cortical depths.

One important distinction is between the terminology "depth" and "layer." Here, depth refers to

the anatomical depth in the cortex while layer refers to a specific network derived from a cortical

depth. This distinction is critical to avoid equating a network layer with an associated cerebral

cortical layer.

The brain was parcellated into 148 regions-of-interest (ROIs) (74 per hemisphere) based on the

Destrieux atlas in FreeSurfer (Destrieux et al., 2010; Fischl et al., 2004) (**Figure 1B**). A detailed

list of parcellations can be found in the **Table S1**.

Resting-state time series for each participant were concatenated across runs, leading to the

following number of time points: 2028 for $n = 23$ participants; 1690 time points: $n = 1$; 1859

time points: $n = 1$; 2197 time points: $n = 2$; 2366 time points: $n = 2$, and 3042 time points: $n = 1$.

Concatenated time series were detrended and filtered using a second-order Butterworth filter

[high-pass: 0.01 Hz, low-pass: 0.1 Hz, MATLAB-filtfilt].

For the **layer-by-layer approach**, Pearson correlations were derived between ROIs within the

same depth, resulting in 10878 pairwise correlations from 148 nodes (ROIs) after removing 148

self-correlations (number of correlations = $(\text{nodes}^2 - \text{diagonal nodes}) / 2$). Pearson correlation

coefficients were normalized using Fisher's z-transformation resulting in five 148-by-148

symmetric weighted connectivity matrices for each participant, i.e., one matrix for each cortical

depth (**Figure 1D**). Pearson correlation was used as opposed to partial correlations since partial

correlations have been shown to perform poorly in networks with a large number of ROIs (Smith

et al., 2011), the larger impact of noise and time series length on partial correlations (Liegeois et

al., 2020; Matkovic et al., 2023), and the popularity of Pearson correlations in previous studies

(Casas-Roma et al., 2022; Wang et al., 2014).

For the **multilayer approach**, Pearson correlations were derived between ROIs between and

within all depths, resulting in 273430 pairwise correlations from 740 nodes (140 ROIs times 5

depths) after removing 740 self-correlations. Pearson correlation coefficients were then

normalized using Fisher's z-transformation with the final product being a 740 by 740 symmetric

weighted connectivity matrix for each participant (**Figure 1D**).

For both approaches, individual matrices were normalized and thresholded at 2% intervals

ranging from 2 to 40% graph density (ratio of edges present to total number of possible edges) to

understand measure differences over a wide range of thresholds. Thresholding is required to

minimize the effect spurious correlations and consider only positive correlations.

We also examined the account of within- and between-layer connections only in the context of

the complete multilayer matrix. To this end, we draw two additional types of sub-matrices from

the multilayer matrix, selectively concentrating on either their within-layer aspect (here, termed

multilayer within-layer) or the between-layer aspects (termed **multilayer between-layer**) only.

The **multilayer within-layer** matrices were derived by normalizing the supra-adjacency matrix,

thresholding the matrix, and then extracting the nodes included in each individual layer (i.e.

nodes 1 to 148 for layer 1), creating a 148-by-148 weighted connectivity matrix. The **multilayer**

between-layer matrices were, in turn, derived by normalizing the supra-adjacency matrix,

thresholding the matrix, and then zeroing the five diagonal matrices (from each cortical depth)

composing within-layer connections, thus resulting in only between-layer connections. (It is

worth noting that since the within-layer and between-layer connectivity matrices were extracted

after thresholding, analysis that requires normalization, i.e., non-thresholded matrices, could not

be conducted in the context of this analysis; **Figure 1E**.)

Edge Consistency and Variability

Connections (edges) within and between layers were explored to understand edge consistency

and variability between participants. Edge consistency (Finn et al., 2015) was calculated by

selecting the top five percent of edges with the lowest standard deviation in un-thresholded

multilayer networks. In contrast, edge variability (Menon & Krishnamurthy, 2019) was

calculated by selected the top five percent of edges with the highest standard deviation across

participants. In both cases, edges in each layer were then summed and divided by the total

number of significant edges (edges in the top five percent) to identify the percentage of

significant edges in each layer.

Matrix Similarity

Matrix similarity was used to understand how matrices differed across layers. Thresholded (2–

40%) and normalized matrices were compared using cosine similarity,

$$CS_{i,j} = \frac{X_i \cdot X_j}{||X_i||\,||X_j||}$$

where X_i and X_j are vectors of the upper triangular elements of two adjacency matrices

[MATLAB-pdist2], with values ranging from -1 (maximal dissimilarity) to $+1$ (maximal

similarity). Cosine similarity was shown to distinguish between matrices better than traditionally

used Pearson correlation (Cabral et al., 2017; Menon & Krishnamurthy, 2019). Using both the

layer-by-layer approach and within-layer matrices, each layer was compared to the other layers.

Additionally, to understand how matrix generation differs between methods, the same layer was

compared across layer-by-layer and within-layer approaches.

Connectomic Analysis

Global and nodal measures were calculated in MATLAB using the Brain Connectivity Toolbox

(MATLAB Version R2022b) (Rubinov & Sporns, 2010) on the Massachusetts Life Sciences

Center Compute Cluster (DELL R440 servers with two Intel Xeon Silver 4214R twelve core

CPUs). Global measures characterize the entire network while nodal measures characterize

attributes of specific node (ROI). Nodal measures can also be averaged to create a global

measure. Measures can be grouped into four general categories to describe their overall network

characterization: composition, centrality, integration, and segregation. **Composition** measures

describe the topology of the network while **centrality** measures detail specific nodal importance

for network function. **Integration** measures examine how information flows through the network

and **segregation** measures explore how the network is divided into functional components.

Therefore, different measures can be used to understand different characteristics of the network.

For example, decreased nodal and global average strength was found in maltreated children

indicating decreased overall brain connectivity (Puetz et al., 2017) while decreased clustering

coefficient and global efficiency in patients with Parkinson's disease can signify deficits in brain

network integration and segregation (Schill et al., 2023). **Table 1** denotes the measures used in

this work, and detailed explanation of each measure can be found in Rubinov and Sporns

(Rubinov & Sporns, 2010). Additionally, small-worldness, a global quantifier that examines how

"random" a network is organized, was also calculated on layer-by-layer networks since noisier

data will appear more "random" (Humphries & Gurney, 2008). Small-worldness was not

calculated on within-layer and between-layer connectivity matrices since the calculation requires

normalization.

Table 1. List of network measures used in this work organized by functional category. (n) denotes a nodal measures.

Composition	Centrality	Integration	Segregation
- Largest Cluster Size	- Betweenness	- Characteristic Path	- Maximized
- Graph Density	Centrality (n)	Length	Modularity Q
- Degree (n)	- Eigenvector	- Radius	- Transitivity
- Strength (n)	Centrality (n)	- Diameter	- Clustering
	- Participation	- Global Efficiency	Coefficient (n)
	Coefficient (n)	- Assortativity	- Local Efficiency (n)

Statistical Analysis

To avoid the bias of selecting a single threshold, area-under-the-curve (AUC) analysis was

conducted to create a threshold-independent measure. The measures in **Table 1** were calculated

at each threshold (from 2 to 40% graph density in 2% intervals). The measure values at each

threshold were then plotted against their threshold, and the area underneath the generated curve

was calculated using a trapezoidal integration method [MATLAB-trapz].

For each global measure (and averaged nodal measure), the AUC value for each layer for all

participants was compared using a one-way analysis of variance (ANOVA) [MATLAB-anova1].

A one-way ANOVA was also used to compare each nodal measure to find differences at each

specific ROI (node). For the multilayer network and between-layer measures, only nodal values

(both averaged and individual) were compared since global measures for the multilayer network

(and thus between-layer measures) contain effects from every layer. For global values, a False

Discovery Rate (FDR, Benjamini-Hochberg) correction (alpha = 0.05) was applied to account for

multiple comparisons (Benjamini & Hochberg, 1995; Groppe, 2024). Additionally, for nodal

values (non-averaged), a FRD (Bonferroni-Holm method) correction (alpha = 0.01) was applied

to account for multiple comparisons (Groppe, 2023; Holm, 1979).

Cortical Thickness Validation

One potential confounding factor using whole-brain laminar analysis is that different brain

regions have different cortical thicknesses (Barbas, 2015; Ding et al., 2009; Zachlod et al., 2020).

Thus, comparing cortical thickness values of significant brain regions (defined above) can help

evaluate whether our findings may be influenced by cortical thickness. Subsequently, cortical

thickness values for each ROI for each participant were extracted using FreeSurfer and averaged

across all subjects (Fischl, 2012). The distribution of significant nodes versus non-significant

nodes for each nodal measure and pipeline with greater than ten significant nodes were compared

using a t-test [MATLAB-ttest2].

Figure 2. (A) Edge consistency between each participant (multilayer matrix). The edges of

each layer of the multilayer matrix were compared to find the edge strengths that had the

lowest 5% standard deviation between participants. Higher values indicate a higher percentage

of consistent edges, indicating consistent features between participants for those connections. **(B)** Edge variability between each participant (multilayer matrix). The edges of each layer of the multilayer matrix were compared to find the edge strengths that had the highest 5% standard deviation between participants. Higher values indicate a higher percentage of variable edges, indicating variable features between participants for those connections. **(C)** Cosine similarity between layer-by-layer and within-layer matrices (multilayer approach). Within participant, the matrix generation methods were compared using cosine similarity across a range of thresholds at each layer. Cosine similarity values range from -1 (maximal dissimilarity) to $+1$ (maximal similarity). The mean value at each threshold is plotted while the shaded region indicates the standard error. **(D)** Area-under-the-curve (AUC) measure in comparing layer-by-layer and within-layer matrix generation methods. Linear interpolation was used for visualization. The AUC from **(C)** is calculated using trapezoidal approximation. Higher values indicate higher similarity between methods while lower values indicate lower similarity between methods. The mean AUC value at each layer is plotted while the shaded region indicates the standard error.

Results

Matrix Similarity and Edge Comparison

Edges from the multilayer matrix were compared to understand differences and similarities between participants and to see if the laminar connectomic methodology can distinguish different participants. High edge consistency indicates a similar connectivity pattern between participants

while a high edge variability increases the ability to distinguish been participants. **Figure 2A** and

Table 2 shows the percentage of consistent edge strengths between participants that each layer

contains from the multilayer matrix. Layer 1 (derived from the depth closest to pial surface) has

the largest number of consistent edges (29.52%) with connections between layer 1 and layer 5

(7.06%) being the most consistent between participants. In contrast, **Figure 2B** shows the edge

variability between participants with the highest variability found in layer 5 (closest to white

matter) overall (33.65%) and in within layer connections (11.86%). It is important to clarify that

edge consistency and variability are not mutually related, even though they provide

complementary results above.

After edge analysis, connectivity matrices were compared within participants to understand the

two competing matrix generation methodologies: layer-by-layer vs. within-layer (multilayer)

approach. Cosine similarity was used to examine similarities between different connectivity

matrices. In comparing within participant matrices across layers (**Figure S4**), layers were found

to be similar with the most distant layers (layer 1 to layer 5) having the lowest similarity in both

layer-by-layer matrices and within-layer matrices. Matrices within the same layer and within

participant were also compared across the matrix generation methods (**Figure 2C** and **2D**). Layer

3 was the most consistent across the two methodologies while the peripheral layers (layer 1 and

layer 5) differed the most between methods.

Table 2. Edge consistency and variability percentages for each layer derived from the multilayer matrix. Bolded values show the highest percentage for each measure.

Layer	Edge Consistency (%)	Edge Variability (%)

1 (Superficial)	**29.52**	8.25
2	20.73	12.36
3	15.82	19.11
4	16.03	26.60
5 (Deep)	17.89	**33.65**

Figure 3. Area-under-the-curve (AUC) values across different layers for significant global measures ($p \leq 0.05$) for layer-by-layer analysis. Significance was calculated using a one-way ANOVA with an FDR correction (alpha = 0.05). Linear interpolation was used for visualization. The mean value across participants at each layer is plotted while the shaded region indicates the standard error. P values shown are FDR corrected (Benjamini-Hochberg method, alpha = 0.05).

Single Layer Results

Global

Global network measures were calculated for layer-by-layer matrices. AUC values for each

global measure can be found summarized in **Table S2**. Network diameter ($p = 0.049$)

significantly increased from the most superficial layer 1 to the deepest layer 5 (**Figure 3A**).

Largest cluster size ($p = 0.0024$), average betweenness centrality ($p = 0.028$), average local

efficiency ($p = 0.046$), and eigenvector centrality ($p = 0.028$) significantly decreased from layer 1

to layer 5 (**Figure 3B–E**), with a peak in layer 2 for average betweenness centrality and average

local efficiency. There were no significant differences in modularity, transitivity, characteristic

path length, global efficiency, radius, assortativity, average degree centrality, average strength,

average clustering coefficient, and average participation coefficient (**Figure S5–6).** Graph

density and average degree centrality were constant across layers due to both measures being a

direct function of thresholding (**Figure S5–6**). Additionally, small-worldness showed a general

trend of decreasing with depth; however, there was no significant differences between layers ($p =$

0.2651) and small-worldness was greater than one (indicating a small-world network) for all

thresholds except 40% graph density (**Figure S7**).

Nodal

Table 3 shows the number of nodes in brain regions in layer-by-layer matrices with significant

differences between layers (FDR correction with alpha = 0.01) (See **Table S3** for specific values

and regions). Degree centrality, strength, and eigenvector centrality had the greatest number of

significant nodes (4/148) (**Table 3**). The limbic region had more significant nodes than all other

regions for each measure, except for clustering coefficient which was tied with the temporal

region (one significant node for each region). In all measures, the right hemisphere had more

significant nodes than the left hemisphere (**Table 3**). In general, the most superficial layers (1

and 2) had the highest value for significant nodes (**Table 3, TableS3, Figure S8–S9**). Significant

nodes were distributed across node thickness levels (**Figure S10–S11**).

Table 3. Number of significant nodes within each brain region for layer-by-layer analysis. Significance was calculated from the area-under-the curve (AUC) values using a one-way ANOVA with an FDR correction (alpha = 0.01) to account for multiple comparisons (Groppe, 2023; Holm, 1979). Details of nodal mapping to each region can be found in Table S1. Orange: hemisphere with the highest number of nodes; Yellow: measure with the highest number of nodes; Green: region within each measure with the highest number of nodes.

Network Measure	Hemisphere	All Regions	Frontal	Limbic	Occipital	Parietal	Temporal
Degree Centrality	Left:	1/74	0/24	1/14	0/10	0/10	0/16
	Right:	3/74	0/24	2/14	0/10	0/10	1/16
	Both:	4/148	0/48	3/28	0/20	0/20	1/32
Strength	Left:	1/74	0/24	1/14	0/10	0/10	0/16
	Right:	3/74	0/24	2/14	0/10	0/10	1/16
	Both:	4/148	0/48	3/28	0/20	0/20	1/32
Eigenvector Centrality	Left:	1/74	0/24	1/14	0/10	0/10	0/16
	Right:	3/74	0/24	2/14	0/10	0/10	1/16
	Both:	4/148	0/48	3/28	0/20	0/20	1/32

Betweenness Centrality	Left:	0/74	0/24	0/14	0/10	0/10	0/16
	Right:	3/74	1/24	2/14	0/10	0/10	0/16
	Both:	3/148	1/48	2/28	0/20	0/20	0/32
Clustering Coefficient	Left:	0/74	0/24	0/14	0/10	0/10	0/16
	Right:	2/74	0/24	1/14	0/10	0/10	1/16
	Both:	2/148	0/48	1/28	0/20	0/20	1/32
Local Efficiency	Left:	1/74	0/24	1/14	0/10	0/10	0/16
	Right:	3/74	0/24	2/14	0/10	0/10	1/16
	Both:	4/148	0/48	3/28	0/20	0/20	1/32
Participation Coefficient	Left:	1/74	0/24	1/14	0/10	0/10	0/16
	Right:	2/74	0/24	2/14	0/10	0/10	0/16
	Both:	3/148	0/48	3/28	0/20	0/20	0/32

Figure 4. Area-under-the-curve (AUC) values across different layers for significant global measures ($p \leq 0.05$) for within-layer analysis. Significance was calculated using a one-way

ANOVA with an FDR correction (alpha = 0.05). Linear interpolation was used for visualization. The mean value across participants at each layer is plotted while the shaded region indicates the standard error. P values shown are FDR corrected (Benjamini-Hochberg method, alpha = 0.05).

Multilayer Results

Within-layer Global

Figure 4 shows global network measures calculated for within-layer matrices. AUC values for

each global measure can be found summarized in **Table S4**. Characteristic path length ($p =$

0.014) and diameter ($p < 0.001$) all increased from layer 1 to layer 5 (**Figure 4A–B**). Largest

cluster size ($p < 0.001$), graph density ($p < 0.001$), average degree centrality (p < 0.001), average

strength ($p < 0.001$), average eigenvector centrality ($p < 0.001$), and average participation

coefficient ($p = 0.0011$) significantly decreased with cortical depth (layer 1 to 5) (**Figure 4C–H**).

There were no significant differences for modularity, transitivity, global efficiency, radius,

assortativity, average betweenness centrality, average clustering coefficient, and average local

efficiency (**Figure S12–13**). In contrast to layer-by-layer results, graph density and average

degree centrality were different across layers due to the within-layer matrix generation

methodology allowing each individual layer to have a different graph density.

Within-layer Local

The number of nodes in brain regions with significant differences using within-layer matrices

can be found in **Table 4** (FDR correction with alpha = 0.01, see **Table S5** for specific values and

regions). Degree centrality had the largest number of significant nodes (15/148) followed by

strength (9/148), participation coefficient (9/148), and local efficiency (8/148) (**Table 4**). In all

measures except participation coefficient, the limbic region had the most significant nodes; in

participation coefficient, the temporal region had the most significant nodes (4/32). In all

measures, the right hemisphere had more significant nodes than the left hemisphere (**Table 4**).

For significant nodes, either layers 1 or 2 had the highest value (**Table 4**, **Table S5, Figure S14–**

15). Significant nodes were spread across different node thicknesses (**Figure S16–S17**).

Table 4. Number of significant nodes within each brain region for within-layer analysis. Significance was calculated from the area-under-the curve (AUC) values using a one-way ANOVA with an FDR correction (alpha = 0.01) to account for multiple comparisons (Groppe, 2023; Holm, 1979). Details of nodal mapping to each region can be found in Table S1. Orange: hemisphere with the highest number of nodes; Yellow: measure with the highest number of nodes; Green: region within each measure with the highest number of nodes.

Network Measure	Hemisphere	All Regions	Frontal	Limbic	Occipital	Parietal	Temporal
Degree Centrality	Left:	4/74	0/24	4/14	0/10	0/10	0/16
	Right:	11/74	1/24	5/14	0/10	0/10	5/16
	Both:	15/148	1/48	9/28	0/20	0/20	5/32
Strength	Left:	2/74	0/24	2/14	0/10	0/10	0/16
	Right:	7/74	0/24	3/14	0/10	0/10	4/16

	Both:	9/148	0/48	5/28	0/20	0/20	4/32
Eigenvector Centrality	Left:	1/74	0/24	1/14	0/10	0/10	0/16
	Right:	3/74	0/24	2/14	0/10	0/10	1/16
	Both:	4/148	0/48	3/28	0/20	0/20	1/32
Betweenness Centrality	Left:	0/74	0/24	0/14	0/10	0/10	0/16
	Right:	3/74	1/24	2/14	0/10	0/10	0/16
	Both:	3/148	1/48	2/28	0/20	0/20	0/32
Clustering Coefficient	Left:	1/74	0/24	1/14	0/10	0/10	0/16
	Right:	5/74	0/24	3/14	0/10	0/10	2/16
	Both:	6/148	0/48	4/28	0/20	0/20	2/32
Local Efficiency	Left:	1/74	0/24	1/14	0/10	0/10	0/16
	Right:	7/74	0/24	4/14	0/10	0/10	3/16
	Both:	8/148	0/48	5/28	0/20	0/20	3/32
Participation Coefficient	Left:	2/74	0/24	1/14	0/10	0/10	1/16
	Right:	7/74	1/24	2/14	1/10	0/10	3/16
	Both:	9/148	1/48	3/28	1/20	0/20	4/32

Figure 5. Area-under-the-curve (AUC) values across different layers for significant global measures ($p \leq 0.05$) for multilayer analysis. Significance was calculated using a one-way ANOVA with an FDR correction (alpha = 0.05). Linear interpolation was used for visualization. The mean value across participants at each layer is plotted while the shaded region indicates the standard error. P values shown are FDR corrected (Benjamini-Hochberg method, alpha = 0.05).

Multilayer Global

Alongside measures for individual layers, global measures were calculated for the supra-

adjacency matrix created using the multilayer approach. While only global values derived from

nodal averages were statistically compared between layers, AUC values for all global measures

are summarized in **Table S6**. Average degree centrality ($p < 0.001$), average strength ($p =$

0.0019), and average eigenvector centrality ($p < 0.001$) decreased from layer 1 to layer 5 with a

slight peak at layer 2 (**Figure 5A–C**). Average betweenness centrality ($p < 0.001$) and average

participation coefficient ($p = 0.019$) also decreased from layer 1 to layer 5 (**Figure 5D–E**). In

contrast, average clustering coefficient tended to increase from layer 1 to layer 5 ($p = 0.080$)

(**Figure S18–19, Table S6**). Average local efficiency showed no significant difference between

layers (**Figure S18–19, Table S6**). Additionally, layer-wise graph density, graph density derived

from each individual layer within the multilayer network, decreased from layer 1 to layer 5 with

a slight peak at layer 2 (**Figure S20**).

Multilayer Local

Table 5 depicts the number of nodes in specific brain regions that were significantly different

between layers for the multilayer analysis (FDR correction with alpha = 0.01) (See **Table S7** for

specific values and regions). Betweenness centrality identified the most significant nodes

(58/148) with 19/48 frontal nodes, 16/28 limbic nodes, 2/20 occipital nodes, 5/20 parietal nodes,

and 16/32 temporal nodes (**Table 5**). The limbic region had highest percentage of nodes in all

measures except clustering coefficient (4/32 in temporal) and local efficiency (no significant

nodes found). Additionally, using betweenness centrality the frontal region had the highest

number of significant nodes (19/48). Again, in all measures, the right hemisphere had more

nodes with significant differences between layers (**Table 5**). While most significant regions

across measures were highest in the superficial layers (layers 1,2), especially in betweenness

centrality, the deepest layer (layer 5) had the highest values for nodes significant in clustering

coefficient (**Table 5, Table S7, Figure S21**). The thickness of significant nodes was spread

across the spectrum of thickness levels, with a preference towards thicker nodes for betweenness

centrality (**Figure S22–S23**).

Table 5. Number of significant nodes within each brain region for multilayer analysis. Significance was calculated from the area-under-the curve (AUC) values using a one-way ANOVA with an FDR correction (alpha = 0.01) to account for multiple comparisons (Groppe, 2023; Holm, 1979). Details of nodal mapping to each region can be found in Table S1. Orange: hemisphere with the highest number of nodes; Yellow: measure with the highest number of nodes; Green: region within each measure with the highest number of nodes.

Network Measure	Hemisphere	All Regions	Frontal	Limbic	Occipital	Parietal	Temporal
Degree Centrality	Left:	2/74	0/24	2/14	0/10	0/10	0/16
	Right:	4/74	0/24	2/14	0/10	0/10	2/16
	Both:	6/148	0/48	4/28	0/20	0/20	2/32
Strength	Left:	1/74	0/24	1/14	0/10	0/10	0/16
	Right:	4/74	0/24	2/14	0/10	0/10	2/16
	Both:	5/148	0/48	3/28	0/20	0/20	2/32
Eigenvector Centrality	Left:	1/74	0/24	1/14	0/10	0/10	0/16
	Right:	2/74	0/24	2/14	0/10	0/10	0/16
	Both:	3/148	0/48	3/28	0/20	0/20	0/32
Betweenness Centrality	Left:	24/74	8/24	6/14	1/10	2/10	7/16
	Right:	34/74	11/24	10/14	1/10	3/10	9/16
	Both:	58/148	19/48	16/28	2/20	5/20	16/32
Clustering Coefficient	Left:	1/74	0/24	0/14	0/10	0/10	1/16
	Right:	6/74	2/24	1/14	0/10	0/10	3/16

	Both:	7/148	2/48	1/28	0/20	0/20	4/32
Local Efficiency	Left:	0/74	0/24	0/14	0/10	0/10	0/16
	Right:	0/74	0/24	0/14	0/10	0/10	0/16
	Both:	0/148	0/48	0/28	0/20	0/20	0/32
Participation Coefficient	Left:	3/74	0/24	2/14	0/10	0/10	1/16
	Right:	4/74	1/24	2/14	0/10	0/10	1/16
	Both:	7/148	1/48	4/28	0/20	0/20	2/32

Figure 6. Area-under-the-curve (AUC) values across different layers for significant global measures ($p \leq 0.05$) for between-layer analysis. Significance was calculated using a one-way ANOVA with an FDR correction (alpha = 0.05). Linear interpolation was used for visualization. The mean value across participants at each layer is plotted while the shaded region indicates the

standard error. P values shown are FDR corrected (Benjamini-Hochberg method, alpha = 0.05).

Between-layer Global

Significant between-layer global measures can be found in **Figure 6**. AUC values for all global

measures can be found in **Table S8**. Average degree centrality ($p < 0.001$), average strength ($p =$

0.0021), average eigenvector centrality ($p < 0.001$), average betweenness centrality (p < 0.001),

and average participation coefficient ($p = 0.032$) decreased from superficial (layer 1) to deep

(layer 5) with all except average participation coefficient showing a peak in layers 2 and 3

(**Figure 6A–D, F**). Average clustering coefficient ($p = 0.0033$) increased from layer 1 to layer 5

with a slight decrease from layer 1 to layer 2 (**Figure 6E**). Average local efficiency was the only

measure that showed no significant difference between layers (**Figure S24–25, Table S8**).

Between-layer Local

Brain regions with nodes that were significantly different using between-layer matrices can be

found in **Table 6** (FDR correction with alpha = 0.01) (See **Table S9** for specific values and

regions). Betweenness centrality had the largest number of significant nodes (57/148) followed

by clustering coefficient (22/148) (**Table 6**). For betweenness centrality, most significant nodes

had the highest values in the superficial layers (layers 1 and 2) while for clustering coefficient,

all significant nodes were highest in layer 5 (**Figure S26**). In degree centrality, strength,

eigenvector centrality, and betweenness centrality, the limbic region had the highest percentage

of significant nodes within each region (**Table 6**). In betweenness centrality, frontal region nodes

had the highest absolute number of significant nodes (20/48). In clustering coefficient (10/32)

and participation coefficient (4/32), the temporal region had the most significant nodes (**Table**

6). Local efficiency had no significant nodes (**Table 6, Table S9, Figure S26**). The right

hemisphere had more significant nodes than the left hemisphere for all measures (**Table 6**).

Significant nodes were dispersed across different thickness levels, with betweenness centrality

nodes leaning slightly more toward thicker regions (**Figure S27–S28**).

Table 6. Number of significant nodes within each brain region for between analysis. Significance was calculated from the area-under-the curve (AUC) values using a one-way ANOVA with an FDR correction (alpha = 0.01) to account for multiple comparisons (Groppe, 2023; Holm, 1979). Details of nodal mapping to each region can be found in Table S1. Orange: hemisphere with the highest number of nodes; Yellow: measure with the highest number of nodes; Green: region within each measure with the highest number of nodes.

Network Measure	Hemisphere	All Regions	Frontal	Limbic	Occipital	Parietal	Temporal
Degree Centrality	Left:	1/74	0/24	1/14	0/10	0/10	0/16
	Right:	4/74	0/24	2/14	0/10	0/10	2/16
	Both:	5/148	0/48	3/28	0/20	0/20	2/32
Strength	Left:	1/74	0/24	1/14	0/10	0/10	0/16
	Right:	4/74	0/24	2/14	0/10	0/10	2/16
	Both:	5/148	0/48	3/28	0/20	0/20	2/32
Eigenvector Centrality	Left:	1/74	0/24	1/14	0/10	0/10	0/16
	Right:	2/74	0/24	2/14	0/10	0/10	0/16
	Both:	3/148	0/48	3/28	0/20	0/20	0/32
Betweenness	Left:	25/74	10/24	7/14	1/10	1/10	6/16

Centrality	Right:	32/74	10/24	11/14	1/10	2/10	8/16
	Both:	57/148	20/48	18/28	2/20	3/20	14/32
Clustering Coefficient	Left:	6/74	0/24	3/14	0/10	0/10	3/16
	Right:	16/74	3/24	5/14	1/10	0/10	7/16
	Both:	22/148	3/48	8/28	1/20	0/20	10/32
Local Efficiency	Left:	0/74	0/24	0/14	0/10	0/10	0/16
	Right:	0/74	0/24	0/14	0/10	0/10	0/16
	Both:	0/148	0/48	0/28	0/20	0/20	0/32
Participation Coefficient	Left:	2/74	0/24	1/14	0/10	0/10	1/16
	Right:	5/74	1/24	1/14	0/10	0/10	3/16
	Both:	7/148	1/48	2/28	0/20	0/20	4/32

* White sections indicate no significant difference between layers
** Gray sections indicate network measures that cannot separate layers

Figure 7. Layer with the highest area-under-the-curve (AUC) value across different global network measures methods ($p \leq 0.05$). Significance was calculated using a one-way ANOVA with an FDR correction (alpha = 0.05). White sections indicate no significant difference between layers. Gray sections indicate network measures that cannot separate effects of different layers.

Method Comparison

All four network measure methods (layer-by-layer, within-layer, multilayer, between-layer)

identified global differences between layers (**Figure 7**). Layer-by-layer and within-layer methods

showed an increase in an integration-based global measure (diameter) from layer 1 to layer 5;

however, the within-layer approach identified an increase in characteristic path length as well.

Similarly, layer-by-layer and within-layer approaches identified a decrease in largest cluster size

from layer 1 to layer 5, with an additional decrease found in graph density using the within-layer

method.

For nodal-averaged global measures, which can be applied to all four methods, measures

generally decreased from superficial layers (1 and 2) to deeper layers (**Figure 7**). The layer-by-

layer methodology identified significant differences in three measures (average eigenvector

centrality, average betweenness centrality, and average local efficiency) while the within-layer

approach found significant differences in four measures (average degree centrality, average

strength, average eigenvector centrality, and average participation coefficient). The multilayer

approach also found differences in five measures (average degree centrality, average strength,

average eigenvector centrality, average betweenness centrality, and average participation

coefficient). Interestingly, the between-layer method identified the most measures with significant differences between layers using six measures, with two measures (average strength and average eigenvector centrality) peaking in the middle layer. Additionally, the between-layer approach was the only method to identify a difference in average clustering coefficient which was the highest in layer 5.

Figure 8. Nodes with significant differences between layers for each nodal measure pipeline (layer-by-layer, within-layer, multilayer, between-layer) for **(A)** betweenness centrality and **(B)** clustering coefficient. Significance was calculated from the area-under-the curve (AUC) values using a one-way ANOVA with an FDR correction (alpha = 0.01) to account for multiple comparisons (Groppe, 2023; Holm, 1979). The colored section represents the layer with the highest value for the node. The nodes are based on the Destrieux atlas in FreeSurfer (Destrieux et al., 2010; Fischl et al., 2004). LH: left hemisphere; RH: right hemisphere.

Similar to global measures, the multilayer-based approaches (within-layer, multilayer, between-

layer) identified more nodal differences between layers than the layer-by-layer approach (**Table**

3–6). For example, the layer-by-layer method identified a maximum of four nodes with

significant differences per measures compared to fifteen for within-layer (degree centrality),

fifty-eight for multilayer (betweenness centrality), and fifty-seven for between-layer

(betweenness centrality). Despite this, in a majority of measures for all four methods, the limbic

region had the greatest number of nodes with significant differences between layers. The right

hemisphere also had more significant nodes across all methods and measures (**Table 3–6**). The

right (2.4127 ± 0.0306 mm; AVG $\pm$ SE) and left (2.4069 ± 0.0303 mm) hemispheres had

comparable cortical thicknesses overall and across brain regions (**Figure S29**). Additionally,

while the thickness of significant nodes versus non-significant nodes was significantly larger in

betweenness centrality metrics (multilayer, between-layer), the absolute difference between

significant and non-significant nodes was typically less than 1 mm (**Figure S30**).

Two measures that showed a considerable benefit from the multilayer-based approach were

betweenness centrality and clustering coefficient (**Figure 8**). For example, the multilayer and

between-layer methods showed substantial increase in the number of nodes that had significant

differences between layers. Similarly, the number of nodes with significant differences between

layers in clustering coefficient increased using multilayer and between-layer methods. More

importantly, however, is clustering coefficient in multilayer and between-layer approaches is the

only measure to highlight the deepest layer as having the largest value. Likewise, the multilayer

and between-layer methods are the only methods to include nodes that are the highest value in

the middle layer (**Figure S21, S26**).

Discussion

Graph theory analysis of high-resolution (7T) resting-state fMRI revealed global and nodal

network differences between cortical depths. Global integration measures (diameter,

characteristic path length) were higher in deeper layers while composition (largest cluster size,

graph density, degree centrality, strength) and centrality (eigenvector centrality, betweenness

centrality) measures were often higher in superficial layers (**Figure 7**).

Current literature exploring graph theory measures and overall laminar connectivity through

networks in the human brain is very limited. Structural analysis of the human connectome using

diffusion MRI combined with T_1-weighted anatomical imaging found qualitative differences in

degree, strength, and betweenness centrality nodal distributions across cortical depths (Shamir et

al., 2022); however, network-wide global calculations were absent. In a functional approach,

Deshpande et al. found no global differences between layers using mean blind deconvoluted

Pearson correlations from resting-state fMRI (Deshpande et al., 2022). However, no threshold

was used, enabling spurious correlations to impact the mean. Additionally, using the mean across

the whole brain obfuscates any impact of a particular ROI. Our results, thus, significantly extend

global and nodal network analysis of cortical architecture across the entire brain.

Our findings provide evidence of an advantage of applying of multilayer graph theory to

connectomic analysis. While differences between layers were seen across all methodologies, the

multilayer approach provided a greater identification of these differences through identifying

more measures with larger significant differences (**Figure 7**). Previous connectomic studies have

shown a benefit of applying a multilayer network framework (Betzel & Bassett, 2017; De

Domenico, 2017; Vaiana & Muldoon, 2018). For example, multilayer connectomics enables the

integration of complex neuroimaging data (cortical lamina, frequency bands, multi-modal

neuroimaging) (Betzel & Bassett, 2017; Breedt et al., 2023; Buldu & Porter, 2018; Casas-Roma

et al., 2022; De Domenico, 2017; Shamir & Assaf, 2023; Vaiana & Muldoon, 2018) and the

creation of new network features. New network features can be used to explain neuroscientific

findings, as in this work, or even enhance machine learning workflows to better discriminate

between disease states (Zhu et al., 2022). Future connectomics studies with available data should

therefore consider using a multilayer framework to augment brain network modeling and

analysis.

One particular benefit of multilayer analysis in laminar connectomics is the ability to

discriminate between and incorporate the impact of within- and between-layer connections.

While this comparison was limited to nodal averaged global measures and nodal values, there

was still a stark comparison between within- and between-layer connections. When exploring

layers individually (within-layer connections), the most superficial layer had the highest

activation and is densely connected to itself while the deepest layer was relatively sparsely

connected and took longer network paths to transmit information to different brain regions

(**Figure 4,7**). However, in between layer connections, layer 1 becomes less important and the

superficial-middle layers (layers 2 and 3) become integral for cortical connectivity (**Figure 6–7**).

The superficial-middle layers (layers 2 and 3) had the highest between-layer degree, strength,

and eigenvector centrality, indicating both layers are densely connected to other layers.

Additionally, layer 2's significantly larger betweenness centrality demonstrates it is the most

important layer for information flow between layers (**Figure 6–7**). Lastly, the deepest layer had

the highest clustering coefficient meaning it had the highest likelihood of forming local hubs

(triangles) with other layers. Thus, within- and between-layer analysis provides evidence of a

highly active superficial layer that utilizes layers 2 and 3 to transmit information to other cortical

layers.

One important note to contextualize the above findings is that while activity and hubs of

information flow can be identified, the direction of information flow cannot be delineated.

Cortical layer architecture can have diverse connectivity patterns across layers and hierarchal

schemes (Felleman & Van Essen, 1991). Therefore, it is important to recognize the correlation

nature of this work rather than infer causality. Furthermore, while our results primarily focus on

layers with the highest measure/activity, this does not imply that other layers are inactive. This

contextualization is notably important in the context of feedback/feedforward interpretations, a

common framework for laminar analysis. In general, feedback is thought to target superficial and

deep layers, and feedforward targets the middle layer (Barbas, 2015; Felleman & Van Essen,

1991; Rockland & Pandya, 1979). However, the fact that one area might also send information,

makes the interpretation less straightforward. Thus, the information transfer and clustering

processes might not directly reflect the feedback/feedforward processes, but complex

interactions between them. This intricate feedback/feedforward interplay may also explain the

connections between layer 1 and layer 5 (7.06%) being the most consistent between participants

(**Figure 2A**) with both superficial and deep layers are activated on opposite sides of the

feedback/feedforward circuit (Miyashita, 2022). However, this feedback/feedforward interaction

would need to be further validated and studied before drawing concrete conclusions.

The significance of connectivity patterns and characteristics for cortical depths differed across

different brain regions. We identified extensive differences between layers in the frontal, limbic,

and temporal brain regions (**Figure 8, Table 3–6**). Interestingly, the limbic cortex, often with the

most significant regions per measures, typically has less layers than other brain regions (Barbas,

2015). Thus, cellular architecture may play a role in the ability for laminar fMRI, and potentially

multilayer connectomics, to detect differences between cortical layers. Cortical thickness may

also play a role in detecting differences between layers (Fukutomi et al., 2018). The regions

(frontal, limbic, and temporal) with the most differences were often the thickest regions (**Figure**

S29), with our results overlapping with previous studies (Fukutomi et al., 2018). Additionally,

other factors that may impact cortical function and detectability including the neurite density

index, orientation dispersion index, and myelin (Fukutomi et al., 2018). However, Fukutomi's et

al. findings show a varied distribution across regions for these measures (Fukutomi et al., 2018).

Despite this variation, hot-spots in these metrics near the posterior-ventral part of the cingulate

gyrus and transverse temporal sulcus often overlap with significant nodes in our work.

Therefore, our results indicate the need to contextualize layer fMRI results within cortical

metrics while providing a framework for potential regions (frontal, limbic, temporal) that may be

suited for whole-brain laminar analysis.

On a global network level, there were high levels of activation in superficial layers when

compared to deeper layers, in line with previous resting-state fMRI analyses reporting increased

activation patterns in superficial cortical depths (Guidi et al., 2020; L. Huber et al., 2021; L. R.

Huber et al., 2021; Polimeni et al., 2010). This was also reflected in the in higher graph density

in superficial layers (**Figure S20**), despite marginally higher tSNR in deeper layers (**Figure S3**).

Higher composition and centrality measures indicate a more robustly connected network (**Figure**

7). Additionally, at least within the same network (layer-by-layer and within-layer), deeper

layers had significantly longer paths to transmit information, shown by higher global integration

measures. It is, however, important to consider the pial vein bias (Polimeni et al., 2010), which

increases gradient-echo BOLD signals from the deep to the more superficial parts of cortex.

Further studies with alternative contrast mechanisms, which are less affected by the draining vein

effect, are thus needed to determine whether the superficial cortical depths play a more critical

role in the brain at rest, compared to the deeper aspects of the cortex or if this result is a function

of signal strength and variance increasing toward the superficial depth.

Limitations

This study does have some limitations, both in terms of laminar analysis and connectomic

analysis. Regarding our laminar analysis approach, the limitations of this study include biases

associated with our fMRI pulse sequence, signal-to-noise ratio bias, the number of cortical

depths chosen, the depth to cortical layer incongruence, and the impact of resting-state versus

task-based paradigms. This work uses gradient-echo BOLD pulse sequences which may not be

fully optimal for laminar analysis (Bandettini et al., 2021; L. Huber et al., 2021). Gradient-echo

BOLD can be influenced largely by large draining vessels (Markuerkiaga et al., 2016a; Olman et

al., 2007; Polimeni et al., 2010; Turner, 2002), while spin-echo BOLD (Duong et al., 2003;

Uludag et al., 2009; Yacoub et al., 2003; Zhao et al., 2004) and VAscular Space Occupancy

(VASO) (Chai et al., 2020; Chai et al., 2021) have been proposed as alternative fMRI contrasts

for laminar analysis to address this large vein bias (Bandettini et al., 2021; L. Huber et al., 2021).

However, VASO and spin-echo BOLD have lower sensitivity and several practical challenges

(Moerel et al., 2021). Similarly, signal-to-noise ratio (SNR) can vary at different cortical depths.

For example, depths within the middle of the cortex will contain less tissue boundary effects

compared to the depths near the pial and white matter surfaces (Blazejewska et al., 2019). This

difference may be further exacerbated since the thickness and functionality of cortical layers can

change based on brain region (Barbas, 2015; Ding et al., 2009; Zachlod et al., 2020) and cortical

curvature (Fatterpekar et al., 2003; Fischl & Dale, 2000; Hilgetag & Barbas, 2006; Van Essen &

Maunsell, 1980). However, as shown above for nodal analysis, thickness varied across

statistically significant nodes suggesting our results are not purely a function of cortical thickness

since significant differences were identified in "thinner" nodes (**Figure S10–S11, S16–S17,**

S22–S23, S27–S30). However, for multilayer ($p < 0.001$) and between-layer ($p < 0.001$)

betweenness centrality (**Figure S30**), nodes with significant differences between layers had

higher cortical thickness. This finding could either result from methodological constraints

indicating an inability to detect differences at lower cortical thicknesses or a neurophysiological

phenomenon of thicker nodes having larger functional differences between layers. Cortical

curvature was not explored, and future laminar work should include the anatomical constraints of

the cortex to address this. In addition to the location of the cortical depths chosen, the number of

depths can affect the results. Other studies have used a smaller number of depths to ensure

independence between depths (Sharoh et al., 2019), six depths to match the number of cortical

layers (Pais-Roldan et al., 2023), or even a larger amount that showed an improved detection of

cortical responses (Huber et al., 2017). The number of depths chosen should balance

independence, cortical response detection, and computational demands from a higher depth

count. The number of depths can also impact the role of partial volume effects due to voxel

overlap. Furthermore, as mentioned above, the cortical depths do not directly equate to

cytoarchitectural cortical layers. Lastly, this study used resting-state fMRI to study whole-brain

connectivity; however, laminar resting-state fMRI activation patterns may be different than

laminar task-based patterns (Pais-Roldan et al., 2023), limiting the broad applicability to task-

based laminar paradigms. Despite potential activation pattern differences, the underlying

anatomical basis of resting-state connections (Adachi et al., 2012; Honey et al., 2009; Turk et al.,

2016; van den Heuvel et al., 2016) can still inform task-based paradigms. Ideally, a second

dataset would be utilized to validate our results; however, few comparable datasets are available.

Regarding our connectomic analysis, limitations include the parcellation choice, network

construction approach, thresholding methodology, and multilayer measure calculations.

Parcellation choice can impact graph theory results (Albers et al., 2021; Arslan et al., 2018). This

work used the Destrieux atlas in FreeSurfer (Destrieux et al., 2010; Fischl et al., 2004), which is

based on anatomical nomenclature. However, an atlas derived from functional connectivity

(Schaefer et al., 2018) or utilizing functional localizers for specific areas of interest (Nieto-

Castanon & Fedorenko, 2012) may be more appropriate for a functional analysis study.

Additionally, for laminar analysis, a custom atlas using laminar cytoarchitecture and cortical

thickness may improve the accuracy of the results. Another impactful choice in connectomic

methodology is how to construct the network from the fMRI time series. Pearson correlations

perform better for network construction when using a large number of ROIs (Smith et al., 2011)

or in noisier data (Liegeois et al., 2020; Matkovic et al., 2023). However, Pearson correlations

also include indirect effects of ROIs which can alter analysis. A popular alternative is partial

correlation which utilizes the inverse covariance matrix, and, thus, excludes the indirect network

effects. However, partial correlations can also include spurious connections (Berkson's paradox

(Berkson, 1946)) and tend to increase network construction complexity since partial correlations

require regularization that has varying optimization parameters (Kim et al., 2015; Pervaiz et al.,

2020). Thus, partial correlations may have future utility in laminar multilayer analysis; however,

limited ROI number in relation to fMRI time series data points and lack of optimized laminar

multilayer regularization parameters led to Pearson correlation being used in this work. Also, as

mentioned above, graph theory measures are directly impacted by thresholding the network

(Osmanlioglu et al., 2020). AUC analysis attempts to correct for this thresholding bias but still

may be inadequate for eliminating thresholding's effect on network characteristics. Additionally,

network measures may be impacted as a result of SNR and network layer normalization (Mandke

et al., 2018). For example, increased noise will transition network structure from small-world to

more random (Humphries & Gurney, 2008), which may occur as we measure deeper into the

cortex. However, our results showed no significant differences in small-worldness between

layers, indicating that this network structure change is not occurring in our work (**Figure S7**).

Similarly, within multilayer approaches, normalization plays a key role since graph density can

influence network properties. However, even with comparable tSNR (**Figure S3**), the graph

density varied across cortical depths. While we believe this to be an intrinsic property of cortical

connectivity having higher density in superficial cortical depths, as demonstrated by other

studies (Logothetis et al., 2001), future work should explore different multilayer normalization

schemes in laminar connectivity to more thoroughly parse through this effect (Mandke et al.,

2018). Lastly, our statistical analysis of our measures may be limited be the use of ANOVA

since it assumes normality and equal variance which are sometimes violated by network

measures. Additionally, when selecting peak values for each measure, the highest value was

selected instead of using a planned contrast ANOVA.

Increased BOLD signal in superficial vs. deeper layers may be due to vascular-related bias

(Markuerkiaga et al., 2016a; Olman et al., 2007; Pais-Roldan et al., 2023; Polimeni et al., 2010;

Turner, 2002). One might conclude that the present results reflect vascular biases. The most

superficial depth was excluded in this work to reduce this bias; however, the other layers will

still have some effect of vascular draining. Additionally, even with removal of the most

superficial depth, the current most superficial depth may still be including superficial voxels that

are sensitive to vascular-related bias. Despite this limitation, some composition and centrality

measures peaked in layers 2 and 3, notably average strength, suggesting that some observed

effects are not explainable by biases in superficial layers (**Figure 7**, **S10–S11, S16–S17, S22–**

S23, S27–S30). Lastly, even the utility of the multilayer approach to find more significant effects

may be a result of the multilayer model being more sensitive to draining/signal confounds.

Conclusion

Our multilayer connectomics findings demonstrate global and nodal network differences

between cortical depths that can be more aptly identified through the multilayer approach

compared to traditional single layer connectomics. These results demonstrate the validity of the

multilayer connectomic framework on laminar fMRI and provide a methodological foundation

for future multilayer laminar studies. Future work should further explore the intersection of

connectomics and laminar studies and address current methodological constraints.

Supporting Information

Author Contributions

Parker Kotlarz: Conceptualization; Methodology; Software; Validation; Formal analysis;

Investigation; Data curation; Writing – original draft; Writing – review & editing; Visualization.

Kaisu Lankinen: Conceptualization; Methodology; Software; Investigation; Writing – review &

editing; Visualization; Supervision.

Maria Hakonen: Methodology; Software; Investigation; Data curation; Writing – review &

editing.

Tori Turpin: Investigation; Writing – review & editing.

Jonathan R. Polimeni: Methodology; Software; Writing – review & editing.

Jyrki Ahveninen: Conceptualization; Methodology; Investigation; Writing – review & editing;

Supervision; Project administration; Funding acquisition.

Funding Information

Supported by R01DC017991, R01DC016765, R01DC016915 and P41-EB030006 and was made

possible by the resources provided by NIH Shared Instrumentation Grant S10-OD023637.

References

Adachi, Y., Osada, T., Sporns, O., Watanabe, T., Matsui, T., Miyamoto, K., & Miyashita, Y.

(2012). Functional connectivity between anatomically unconnected areas is shaped by

collective network-level effects in the macaque cortex. *Cereb Cortex*, *22*(7), 1586-1592.

https://doi.org/10.1093/cercor/bhr234

Ahsan, S. A., Chendeb, K., Briggs, R. G., Fletcher, L. R., Jones, R. G., Chakraborty, A. R., Nix,

C. E., Jacobs, C. C., Lack, A. M., Griffin, D. T., Teo, C., & Sughrue, M. E. (2020).

Beyond eloquence and onto centrality: a new paradigm in planning supratentorial

neurosurgery. *J Neurooncol*, *146*(2), 229-238. https://doi.org/10.1007/s11060-019-

03327-4

Albers, K. J., Ambrosen, K. S., Liptrot, M. G., Dyrby, T. B., Schmidt, M. N., & Morup, M.

(2021). Using connectomics for predictive assessment of brain parcellations.

Neuroimage, *238*, 118170. https://doi.org/10.1016/j.neuroimage.2021.118170

Andersson, J. L., Skare, S., & Ashburner, J. (2003). How to correct susceptibility distortions in

spin-echo echo-planar images: application to diffusion tensor imaging. *Neuroimage*,

20(2), 870-888. https://doi.org/10.1016/s1053-8119(03)00336-7

Arslan, S., Ktena, S. I., Makropoulos, A., Robinson, E. C., Rueckert, D., & Parisot, S. (2018).

Human brain mapping: A systematic comparison of parcellation methods for the human

cerebral cortex. *Neuroimage*, *170*, 5-30.

https://doi.org/10.1016/j.neuroimage.2017.04.014

Bandettini, P. A., Huber, L., & Finn, E. S. (2021). Challenges and opportunities of mesoscopic

brain mapping with fMRI. *Current Opinion in Behavioral Sciences*, *40*, 189-200.

https://doi.org/10.1016/j.cobeha.2021.06.002

Barbas, H. (2015). General cortical and special prefrontal connections: principles from structure

to function. *Annu Rev Neurosci*, *38*, 269-289. https://doi.org/10.1146/annurev-neuro-

071714-033936

Benito-Leon, J., Sanz-Morales, E., Melero, H., Louis, E. D., Romero, J. P., Rocon, E., &

Malpica, N. (2019). Graph theory analysis of resting-state functional magnetic resonance

imaging in essential tremor. *Hum Brain Mapp*, *40*(16), 4686-4702.

https://doi.org/10.1002/hbm.24730

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*

(Methodological), *57*(1). https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Berkson, J. (1946). Limitations of the application of fourfold table analysis to hospital data.

Biometrics, *2*(3), 47-53. https://www.ncbi.nlm.nih.gov/pubmed/21001024

Betzel, R. F., & Bassett, D. S. (2017). Multi-scale brain networks. *Neuroimage*, *160*, 73-83.

https://doi.org/10.1016/j.neuroimage.2016.11.006

Blazejewska, A. I., Fischl, B., Wald, L. L., & Polimeni, J. R. (2019). Intracortical smoothing of

small-voxel fMRI data can provide increased detection power without spatial resolution

losses compared to conventional large-voxel fMRI data. *Neuroimage*, *189*, 601-614.

https://doi.org/10.1016/j.neuroimage.2019.01.054

Boccaletti, S., Bianconi, G., Criado, R., Del Genio, C. I., Gomez-Gardenes, J., Romance, M.,

Sendina-Nadal, I., Wang, Z., & Zanin, M. (2014). The structure and dynamics of

multilayer networks. *Phys Rep*, *544*(1), 1-122.

https://doi.org/10.1016/j.physrep.2014.07.001

Breedt, L. C., Santos, F. A. N., Hillebrand, A., Reneman, L., van Rootselaar, A.-F., Schoonheim,

M. M., Stam, C. J., Ticheler, A., Tijms, B. M., Veltman, D. J., Vriend, C., Wagenmakers,

M. J., van Wingen, G. A., Geurts, J. J. G., Schrantee, A., & Douw, L. (2023). Multimodal

multilayer network centrality relates to executive functioning. *Network Neuroscience*,

7(1), 299-321. https://doi.org/10.1162/netn_a_00284

Buldu, J. M., & Porter, M. A. (2018). Frequency-based brain networks: From a multiplex

framework to a full multilayer description. *Netw Neurosci*, *2*(4), 418-441.

https://doi.org/10.1162/netn_a_00033

Cabral, J., Vidaurre, D., Marques, P., Magalhaes, R., Silva Moreira, P., Miguel Soares, J., Deco,

G., Sousa, N., & Kringelbach, M. L. (2017). Cognitive performance in healthy older

adults relates to spontaneous switching between states of functional connectivity during

rest. *Sci Rep*, *7*(1), 5135. https://doi.org/10.1038/s41598-017-05425-7

Caeyenberghs, K., Verhelst, H., Clemente, A., & Wilson, P. H. (2017). Mapping the functional

connectome in traumatic brain injury: What can graph metrics tell us? *Neuroimage*, *160*,

113-123. https://doi.org/10.1016/j.neuroimage.2016.12.003

Casas-Roma, J., Martinez-Heras, E., Sole-Ribalta, A., Solana, E., Lopez-Soley, E., Vivo, F.,

Diaz-Hurtado, M., Alba-Arbalat, S., Sepulveda, M., Blanco, Y., Saiz, A., Borge-

Holthoefer, J., Llufriu, S., & Prados, F. (2022). Applying multilayer analysis to

morphological, structural, and functional brain networks to identify relevant dysfunction

patterns. *Netw Neurosci*, *6*(3), 916-933. https://doi.org/10.1162/netn_a_00258

Chai, Y., Li, L., Huber, L., Poser, B. A., & Bandettini, P. A. (2020). Integrated VASO and

perfusion contrast: A new tool for laminar functional MRI. *Neuroimage*, *207*, 116358.

https://doi.org/10.1016/j.neuroimage.2019.116358

Chai, Y., Liu, T. T., Marrett, S., Li, L., Khojandi, A., Handwerker, D. A., Alink, A., Muckli, L.,

& Bandettini, P. A. (2021). Topographical and laminar distribution of audiovisual

processing within human planum temporale. *Prog Neurobiol*, *205*, 102121.

https://doi.org/10.1016/j.pneurobio.2021.102121

Chen, V. C., Lin, K. Y., Tsai, Y. H., & Weng, J. C. (2020). Connectome analysis of brain

functional network alterations in breast cancer survivors with and without chemotherapy.

PLoS One, *15*(5), e0232548. https://doi.org/10.1371/journal.pone.0232548

Dadario, N. B., Brahimaj, B., Yeung, J., & Sughrue, M. E. (2021). Reducing the Cognitive

Footprint of Brain Tumor Surgery. *Front Neurol*, *12*, 711646.

https://doi.org/10.3389/fneur.2021.711646

De Domenico, M. (2017). Multilayer modeling and analysis of human brain networks.

Gigascience, *6*(5), 1-8. https://doi.org/10.1093/gigascience/gix004

De Martino, F., Moerel, M., Ugurbil, K., Goebel, R., Yacoub, E., & Formisano, E. (2015).

Frequency preference and attention effects across cortical depths in the human primary

auditory cortex. *Proc Natl Acad Sci U S A*, *112*(52), 16036-16041.

https://doi.org/10.1073/pnas.1507552112

De Vico Fallani, F., Richiardi, J., Chavez, M., & Achard, S. (2014). Graph analysis of functional

brain networks: practical issues in translational neuroscience. *Philos Trans R Soc Lond B*

Biol Sci, *369*(1653). https://doi.org/10.1098/rstb.2013.0521

Deshpande, G., Wang, Y., & Robinson, J. (2022). Resting state fMRI connectivity is sensitive to

laminar connectional architecture in the human brain. *Brain Inform*, *9*(1), 2.

https://doi.org/10.1186/s40708-021-00150-4

Destrieux, C., Fischl, B., Dale, A., & Halgren, E. (2010). Automatic parcellation of human

cortical gyri and sulci using standard anatomical nomenclature. *Neuroimage*, *53*(1), 1-15.

https://doi.org/10.1016/j.neuroimage.2010.06.010

Di, X., Gohel, S., Kim, E. H., & Biswal, B. B. (2013). Task vs. rest-different network

configurations between the coactivation and the resting-state brain networks. *Front Hum*

Neurosci, *7*, 493. https://doi.org/10.3389/fnhum.2013.00493

Ding, S. L., Van Hoesen, G. W., Cassell, M. D., & Poremba, A. (2009). Parcellation of human

temporal polar cortex: a combined analysis of multiple cytoarchitectonic,

chemoarchitectonic, and pathological markers. *J Comp Neurol*, *514*(6), 595-623.

https://doi.org/10.1002/cne.22053

Duong, T. Q., Yacoub, E., Adriany, G., Hu, X., Ugurbil, K., & Kim, S. G. (2003). Microvascular

BOLD contribution at 4 and 7 T in the human brain: gradient-echo and spin-echo fMRI

with suppression of blood effects. *Magn Reson Med*, *49*(6), 1019-1027.

https://doi.org/10.1002/mrm.10472

Farahani, F. V., Karwowski, W., & Lighthall, N. R. (2019). Application of Graph Theory for

Identifying Connectivity Patterns in Human Brain Networks: A Systematic Review.

Front Neurosci, *13*, 585. https://doi.org/10.3389/fnins.2019.00585

Fatterpekar, G. M., Delman, B. N., Boonn, W. W., Gultekin, S. H., Fayad, Z. A., Hoff, P. R., &

Naidich, T. P. (2003). MR microscopy of normal human brain. *Magn Reson Imaging Clin*

N Am, *11*(4), 641-653. https://doi.org/10.1016/s1064-9689(03)00071-0

Felleman, D. J., & Van Essen, D. C. (1991). Distributed hierarchical processing in the primate

cerebral cortex. *Cereb Cortex*, *1*(1), 1-47. https://doi.org/10.1093/cercor/1.1.1-a

Filippi, M., Spinelli, E. G., Cividini, C., Ghirelli, A., Basaia, S., & Agosta, F. (2023). The human

functional connectome in neurodegenerative diseases: relationship to pathology and

clinical progression. *Expert Rev Neurother*, *23*(1), 59-73.

https://doi.org/10.1080/14737175.2023.2174016

Finn, E. S., Huber, L., & Bandettini, P. A. (2021). Higher and deeper: Bringing layer fMRI to

association cortex. *Prog Neurobiol*, *207*, 101930.

https://doi.org/10.1016/j.pneurobio.2020.101930

Finn, E. S., Huber, L., Jangraw, D. C., Molfese, P. J., & Bandettini, P. A. (2019). Layer-

dependent activity in human prefrontal cortex during working memory. *Nat Neurosci*,

22(10), 1687-1695. https://doi.org/10.1038/s41593-019-0487-z

Finn, E. S., Shen, X., Scheinost, D., Rosenberg, M. D., Huang, J., Chun, M. M., Papademetris,

X., & Constable, R. T. (2015). Functional connectome fingerprinting: identifying

individuals using patterns of brain connectivity. *Nat Neurosci*, *18*(11), 1664-1671.

https://doi.org/10.1038/nn.4135

Fischl, B. (2012). FreeSurfer. *Neuroimage*, *62*(2), 774-781.

https://doi.org/10.1016/j.neuroimage.2012.01.021

Fischl, B., & Dale, A. M. (2000). Measuring the thickness of the human cerebral cortex from

magnetic resonance images. *Proc Natl Acad Sci U S A*, *97*(20), 11050-11055.

https://doi.org/10.1073/pnas.200033797

Fischl, B., van der Kouwe, A., Destrieux, C., Halgren, E., Segonne, F., Salat, D. H., Busa, E.,

Seidman, L. J., Goldstein, J., Kennedy, D., Caviness, V., Makris, N., Rosen, B., & Dale,

A. M. (2004). Automatically parcellating the human cerebral cortex. *Cereb Cortex*, *14*(1),

11-22. https://doi.org/10.1093/cercor/bhg087

Fleischer, V., Radetz, A., Ciolac, D., Muthuraman, M., Gonzalez-Escamilla, G., Zipp, F., &

Groppa, S. (2019). Graph Theoretical Framework of Brain Networks in Multiple

Sclerosis: A Review of Concepts. *Neuroscience*, *403*, 35-53.

https://doi.org/10.1016/j.neuroscience.2017.10.033

Fracasso, A., Luijten, P. R., Dumoulin, S. O., & Petridou, N. (2018). Laminar imaging of

positive and negative BOLD in human visual cortex at 7T. *Neuroimage*, *164*, 100-111.

https://doi.org/10.1016/j.neuroimage.2017.02.038

Fukutomi, H., Glasser, M. F., Zhang, H., Autio, J. A., Coalson, T. S., Okada, T., Togashi, K.,

Van Essen, D. C., & Hayashi, T. (2018). Neurite imaging reveals microstructural

variations in human cerebral cortical gray matter. *Neuroimage*, *182*, 488-499.

https://doi.org/10.1016/j.neuroimage.2018.02.017

Gau, R., Bazin, P. L., Trampel, R., Turner, R., & Noppeney, U. (2020). Resolving multisensory

and attentional influences across cortical depth in sensory cortices. *Elife*, *9*.

https://doi.org/10.7554/eLife.46856

Gleichgerrcht, E., Keller, S. S., Drane, D. L., Munsell, B. C., Davis, K. A., Kaestner, E., Weber,

B., Krantz, S., Vandergrift, W. A., Edwards, J. C., McDonald, C. R., Kuzniecky, R., &

Bonilha, L. (2020). Temporal Lobe Epilepsy Surgical Outcomes Can Be Inferred Based

on Structural Connectome Hubs: A Machine Learning Study. *Ann Neurol*, *88*(5), 970-

983. https://doi.org/10.1002/ana.25888

Glover, G. H., Li, T. Q., & Ress, D. (2000). Image-based method for retrospective correction of

physiological motion effects in fMRI: RETROICOR. *Magn Reson Med*, *44*(1), 162-167.

https://doi.org/10.1002/1522-2594(200007)44:1<162::aid-mrm23>3.0.co;2-e

Greve, D. N., & Fischl, B. (2009). Accurate and robust brain image alignment using boundary-

based registration. *Neuroimage*, *48*(1), 63-72.

https://doi.org/10.1016/j.neuroimage.2009.06.060

Griffa, A., Baumann, P. S., Thiran, J. P., & Hagmann, P. (2013). Structural connectomics in

brain diseases. *Neuroimage*, *80*, 515-526.

https://doi.org/10.1016/j.neuroimage.2013.04.056

Groppe, D. (2023). *Bonferroni-Holm Correction for Multiple Comparisons*. In (Version 1.1.0.0.)

https://www.mathworks.com/matlabcentral/fileexchange/28303-bonferroni-holm-

correction-for-multiple-comparisons

Groppe, D. (2024). *fdr_bh*. In (Version 2.3.0.0.)

Guidi, M., Huber, L., Lampe, L., Merola, A., Ihle, K., & Moller, H. E. (2020). Cortical laminar

resting-state signal fluctuations scale with the hypercapnic blood oxygenation level-

dependent response. *Hum Brain Mapp*, *41*(8), 2014-2027.

https://doi.org/10.1002/hbm.24926

Hart, M. G., Price, S. J., & Suckling, J. (2016). Connectome analysis for pre-operative brain

mapping in neurosurgery. *Br J Neurosurg*, *30*(5), 506-517.

https://doi.org/10.1080/02688697.2016.1208809

Hermundstad, A. M., Bassett, D. S., Brown, K. S., Aminoff, E. M., Clewett, D., Freeman, S.,

Frithsen, A., Johnson, A., Tipper, C. M., Miller, M. B., Grafton, S. T., & Carlson, J. M.

(2013). Structural foundations of resting-state and task-based functional connectivity in

the human brain. *Proc Natl Acad Sci U S A*, *110*(15), 6169-6174.

https://doi.org/10.1073/pnas.1219562110

Hilgetag, C. C., & Barbas, H. (2006). Role of mechanical factors in the morphology of the

primate cerebral cortex. *PLoS Comput Biol*, *2*(3), e22.

https://doi.org/10.1371/journal.pcbi.0020022

Holm, S. (1979). A Simple Sequentially Rejective Multiple Test Procedure. *Scandinavian

Journal of Statistics*, *6*(2), 65-70. https://www.jstor.org/stable/4615733

Honey, C. J., Sporns, O., Cammoun, L., Gigandet, X., Thiran, J. P., Meuli, R., & Hagmann, P.

(2009). Predicting human resting-state functional connectivity from structural

connectivity. *Proc Natl Acad Sci U S A*, *106*(6), 2035-2040.

https://doi.org/10.1073/pnas.0811168106

Huber, L., Finn, E. S., Chai, Y., Goebel, R., Stirnberg, R., Stocker, T., Marrett, S., Uludag, K.,

Kim, S. G., Han, S., Bandettini, P. A., & Poser, B. A. (2021). Layer-dependent functional

connectivity methods. *Prog Neurobiol*, *207*, 101835.

https://doi.org/10.1016/j.pneurobio.2020.101835

Huber, L., Handwerker, D. A., Jangraw, D. C., Chen, G., Hall, A., Stuber, C., Gonzalez-Castillo,

J., Ivanov, D., Marrett, S., Guidi, M., Goense, J., Poser, B. A., & Bandettini, P. A. (2017).

High-Resolution CBV-fMRI Allows Mapping of Laminar Activity and Connectivity of

Cortical Input and Output in Human M1. *Neuron*, *96*(6), 1253-1263 e1257.

https://doi.org/10.1016/j.neuron.2017.11.005

Huber, L. R., Poser, B. A., Bandettini, P. A., Arora, K., Wagstyl, K., Cho, S., Goense, J.,

Nothnagel, N., Morgan, A. T., van den Hurk, J., Muller, A. K., Reynolds, R. C., Glen, D.

R., Goebel, R., & Gulban, O. F. (2021). LayNii: A software suite for layer-fMRI.

Neuroimage, *237*, 118091. https://doi.org/10.1016/j.neuroimage.2021.118091

Humphries, M. D., & Gurney, K. (2008). Network 'small-world-ness': a quantitative method for

determining canonical network equivalence. *PLoS One*, *3*(4), e0002051.

https://doi.org/10.1371/journal.pone.0002051

Jacob, Y., Morris, L. S., Huang, K. H., Schneider, M., Rutter, S., Verma, G., Murrough, J. W., &

Balchandani, P. (2020). Neural correlates of rumination in major depressive disorder: A

brain network analysis. *Neuroimage Clin*, *25*, 102142.

https://doi.org/10.1016/j.nicl.2019.102142

Kim, J., Wozniak, J. R., Mueller, B. A., & Pan, W. (2015). Testing group differences in brain

functional connectivity: using correlations or partial correlations? *Brain Connect*, *5*(4),

214-231. https://doi.org/10.1089/brain.2014.0319

Kivela, M., Arenas, A., Barthelemy, M., Gleeson, J. P., Moreno, Y., & Porter, M. A. (2014).

Multilayer networks. *Journal of Complex Networks*, *2*(3), 203-271.

https://doi.org/10.1093/comnet/cnu016

Klein, B. P., Fracasso, A., van Dijk, J. A., Paffen, C. L. E., Te Pas, S. F., & Dumoulin, S. O.

(2018). Cortical depth dependent population receptive field attraction by spatial attention

in human V1. *Neuroimage*, *176*, 301-312.

https://doi.org/10.1016/j.neuroimage.2018.04.055

Kok, P., Bains, L. J., van Mourik, T., Norris, D. G., & de Lange, F. P. (2016). Selective

Activation of the Deep Layers of the Human Primary Visual Cortex by Top-Down

Feedback. *Curr Biol*, *26*(3), 371-376. https://doi.org/10.1016/j.cub.2015.12.038

Kotlarz, P., Nino, J. C., & Febo, M. (2022). Connectomic analysis of Alzheimer's disease using percolation theory. *Netw Neurosci*, *6*(1), 213-233. https://doi.org/10.1162/netn_a_00221

Lankinen, K., Ahlfors, S. P., Mamashli, F., Blazejewska, A. I., Raij, T., Turpin, T., Polimeni, J. R., & Ahveninen, J. (2023). Cortical depth profiles of auditory and visual 7 T functional MRI responses in human superior temporal areas. *Hum Brain Mapp*, *44*(2), 362-372. https://doi.org/10.1002/hbm.26046

Lawrence, S. J., Norris, D. G., & de Lange, F. P. (2019). Dissociable laminar profiles of concurrent bottom-up and top-down modulation in the human visual cortex. *Elife*, *8*. https://doi.org/10.7554/eLife.44422

Lei, D., Li, W., Tallman, M. J., Patino, L. R., McNamara, R. K., Strawn, J. R., Klein, C. C., Nery, F. G., Fleck, D. E., Qin, K., Ai, Y., Yang, J., Zhang, W., Lui, S., Gong, Q., Adler, C. M., Sweeney, J. A., & DelBello, M. P. (2021). Changes in the brain structural connectome after a prospective randomized clinical trial of lithium and quetiapine treatment in youth with bipolar disorder. *Neuropsychopharmacology*, *46*(7), 1315-1323. https://doi.org/10.1038/s41386-021-00989-5

Liegeois, R., Santos, A., Matta, V., Van De Ville, D., & Sayed, A. H. (2020). Revisiting correlation-based functional connectivity and its relationship with structural connectivity. *Netw Neurosci*, *4*(4), 1235-1251. https://doi.org/10.1162/netn_a_00166

Logothetis, N. K., Pauls, J., Augath, M., Trinath, T., & Oeltermann, A. (2001). Neurophysiological investigation of the basis of the fMRI signal. *Nature*, *412*(6843), 150-157. https://doi.org/10.1038/35084005

Mandke, K., Meier, J., Brookes, M. J., O'Dea, R. D., Van Mieghem, P., Stam, C. J., Hillebrand, A., & Tewarie, P. (2018). Comparing multilayer brain networks between groups:

Introducing graph metrics and recommendations. *Neuroimage*, *166*, 371-384.

https://doi.org/10.1016/j.neuroimage.2017.11.016

Mareyam, A., Kirsch, J. E., Chang, Y., Madan, G., & Wald, L. L. (2020). A 64-Channel 7T array

coil for accelerated brain MRI. Proceedings of the International Society for Magnetic

Resonance in Medicine,

Markuerkiaga, I., Barth, M., & Norris, D. G. (2016a). A cortical vascular model for examining

the specificity of the laminar BOLD signal. *Neuroimage*, *132*, 491-498.

https://doi.org/10.1016/j.neuroimage.2016.02.073

Markuerkiaga, I., Barth, M., & Norris, D. G. (2016b). A cortical vascular model for examining

the specificity of the laminar BOLD signal. *Neuroimage*, *132*, 491-498.

https://doi.org/10.1016/j.neuroimage.2016.02.073

S1053-8119(16)00191-9 [pii]

Matkovic, A., Anticevic, A., Murray, J. D., & Repovs, G. (2023). Static and dynamic fMRI-

derived functional connectomes represent largely similar information. *Netw Neurosci*,

7(4), 1266-1301. https://doi.org/10.1162/netn_a_00325

Matthews, P. M., & Hampshire, A. (2016). Clinical Concepts Emerging from fMRI Functional

Connectomics. *Neuron*, *91*(3), 511-528. https://doi.org/10.1016/j.neuron.2016.07.031

Menon, S. S., & Krishnamurthy, K. (2019). A Comparison of Static and Dynamic Functional

Connectivities for Identifying Subjects and Biological Sex Using Intrinsic Individual

Brain Connectivity. *Sci Rep*, *9*(1), 5729. https://doi.org/10.1038/s41598-019-42090-4

Meoded, A., Goldenberg, N. A., & Huisman, T. (2020). Structural Connectomics: State of the

Art and Applications in Pediatric Neurodevelopmental Disorders, Neuro-Oncology, and

Arterial Ischemic Stroke. *J Pediatr*, *221S*, S37-S42.

https://doi.org/10.1016/j.jpeds.2020.01.056

Milano, M., Guzzi, P. H., & Cannataro, M. (2019). Network building and analysis in

connectomics studies: a review of algorithms, databases and technologies. *Network*

Modeling Analysis in Health Informatics and Bioinformatics, *8*(1).

https://doi.org/10.1007/s13721-019-0192-6

Miyashita, Y. (2022). Operating principles of the cerebral cortex as a six-layered network in

primates: beyond the classic canonical circuit model. *Proc Jpn Acad Ser B Phys Biol Sci*,

98(3), 93-111. https://doi.org/10.2183/pjab.98.007

Moerel, M., De Martino, F., Ugurbil, K., Formisano, E., & Yacoub, E. (2018). Evaluating the

Columnar Stability of Acoustic Processing in the Human Auditory Cortex. *J Neurosci*,

38(36), 7822-7832. https://doi.org/10.1523/JNEUROSCI.3576-17.2018

Moerel, M., De Martino, F., Ugurbil, K., Yacoub, E., & Formisano, E. (2019). Processing

complexity increases in superficial layers of human primary auditory cortex. *Sci Rep*,

9(1), 5502. https://doi.org/10.1038/s41598-019-41965-w

Moerel, M., Yacoub, E., Gulban, O. F., Lage-Castellanos, A., & De Martino, F. (2021). Using

high spatial resolution fMRI to understand representation in the auditory network. *Prog*

Neurobiol, *207*, 101887. https://doi.org/10.1016/j.pneurobio.2020.101887

Muckli, L., De Martino, F., Vizioli, L., Petro, L. S., Smith, F. W., Ugurbil, K., Goebel, R., &

Yacoub, E. (2015). Contextual Feedback to Superficial Layers of V1. *Curr Biol*, *25*(20),

2690-2695. https://doi.org/10.1016/j.cub.2015.08.057

Nieto-Castanon, A., & Fedorenko, E. (2012). Subject-specific functional localizers increase

sensitivity and functional resolution of multi-subject analyses. *Neuroimage*, *63*(3), 1646-

1669. https://doi.org/10.1016/j.neuroimage.2012.06.065

Norris, D. G., & Polimeni, J. R. (2019). Laminar (f)MRI: A short history and future prospects.

Neuroimage, *197*, 643-649. https://doi.org/10.1016/j.neuroimage.2019.04.082

Olman, C. A., Inati, S., & Heeger, D. J. (2007). The effect of large veins on spatial localization

with GE BOLD at 3 T: Displacement, not blurring. *Neuroimage*, *34*(3), 1126-1135.

https://doi.org/10.1016/j.neuroimage.2006.08.045

Osmanlioglu, Y., Alappatt, J. A., Parker, D., & Verma, R. (2020). Connectomic consistency: a

systematic stability analysis of structural and functional connectivity. *J Neural Eng*,

17(4), 045004. https://doi.org/10.1088/1741-2552/ab947b

Pais-Roldan, P., Yun, S. D., Palomero-Gallagher, N., & Shah, N. J. (2023). Cortical depth-

dependent human fMRI of resting-state networks using EPIK. *Front Neurosci*, *17*,

1151544. https://doi.org/10.3389/fnins.2023.1151544

Pervaiz, U., Vidaurre, D., Woolrich, M. W., & Smith, S. M. (2020). Optimising network

modelling methods for fMRI. *Neuroimage*, *211*, 116604.

https://doi.org/10.1016/j.neuroimage.2020.116604

Polimeni, J. R., Fischl, B., Greve, D. N., & Wald, L. L. (2010). Laminar analysis of 7T BOLD

using an imposed spatial activation pattern in human V1. *Neuroimage*, *52*(4), 1334-1346.

https://doi.org/10.1016/j.neuroimage.2010.05.005

Polimeni, J. R., Renvall, V., Zaretskaya, N., & Fischl, B. (2018). Analysis strategies for high-

resolution UHF-fMRI data. *Neuroimage*, *168*, 296-320.

https://doi.org/10.1016/j.neuroimage.2017.04.053

Puetz, V. B., Parker, D., Kohn, N., Dahmen, B., Verma, R., & Konrad, K. (2017). Altered brain network integrity after childhood maltreatment: A structural connectomic DTI-study. *Hum Brain Mapp*, *38*(2), 855-868. https://doi.org/10.1002/hbm.23423

Rockland, K. S. (2019). What do we know about laminar connectivity? *Neuroimage*, *197*, 772-784. https://doi.org/10.1016/j.neuroimage.2017.07.032

Rockland, K. S., & Pandya, D. N. (1979). Laminar origins and terminations of cortical connections of the occipital lobe in the rhesus monkey. *Brain Res*, *179*(1), 3-20. https://doi.org/10.1016/0006-8993(79)90485-2

Rubinov, M., & Sporns, O. (2010). Complex network measures of brain connectivity: uses and interpretations. *Neuroimage*, *52*(3), 1059-1069. https://doi.org/10.1016/j.neuroimage.2009.10.003

Sadaghiani, S., Brookes, M. J., & Baillet, S. (2022). Connectomics of human electrophysiology. *Neuroimage*, *247*, 118788. https://doi.org/10.1016/j.neuroimage.2021.118788

Schaefer, A., Kong, R., Gordon, E. M., Laumann, T. O., Zuo, X. N., Holmes, A. J., Eickhoff, S. B., & Yeo, B. T. T. (2018). Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. *Cereb Cortex*, *28*(9), 3095-3114. https://doi.org/10.1093/cercor/bhx179

Schill, J., Simonyan, K., Lang, S., Mathys, C., Thiel, C., & Witt, K. (2023). Parkinson's disease speech production network as determined by graph-theoretical network analysis. *Network Neuroscience*, 1-19. https://doi.org/10.1162/netn_a_00310

Schroeder, C. E., & Foxe, J. J. (2002). The timing and laminar profile of converging inputs to multisensory areas of the macaque neocortex. *Brain Res Cogn Brain Res*, *14*(1), 187-198. https://doi.org/10.1016/s0926-6410(02)00073-3

Schroeder, C. E., Lindsley, R. W., Specht, C., Marcovici, A., Smiley, J. F., & Javitt, D. C.

(2001). Somatosensory input to auditory association cortex in the macaque monkey. *J*

Neurophysiol, *85*(3), 1322-1327. https://doi.org/10.1152/jn.2001.85.3.1322

Setsompop, K., Gagoski, B. A., Polimeni, J. R., Witzel, T., Wedeen, V. J., & Wald, L. L. (2012).

Blipped-controlled aliasing in parallel imaging for simultaneous multislice echo planar

imaging with reduced g-factor penalty. *Magn Reson Med*, *67*(5), 1210-1224.

https://doi.org/10.1002/mrm.23097

Shamir, I., & Assaf, Y. (2021). An MRI-Based, Data-Driven Model of Cortical Laminar

Connectivity. *Neuroinformatics*, *19*(2), 205-218. https://doi.org/10.1007/s12021-020-

09491-7

Shamir, I., & Assaf, Y. (2023). Expanding connectomics to the laminar level: A perspective.

Network Neuroscience, 1-12. https://doi.org/10.1162/netn_a_00304

Shamir, I., Tomer, O., Krupnik, R., & Assaf, Y. (2022). Modelling the laminar connectome of

the human brain. *Brain Struct Funct*, *227*(6), 2153-2165. https://doi.org/10.1007/s00429-

022-02513-x

Sharoh, D., van Mourik, T., Bains, L. J., Segaert, K., Weber, K., Hagoort, P., & Norris, D. G.

(2019). Laminar specific fMRI reveals directed interactions in distributed networks

during language processing. *Proc Natl Acad Sci U S A*, *116*(42), 21185-21190.

https://doi.org/10.1073/pnas.1907858116

Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E., Johansen-Berg,

H., Bannister, P. R., De Luca, M., Drobnjak, I., Flitney, D. E., Niazy, R. K., Saunders, J.,

Vickers, J., Zhang, Y., De Stefano, N., Brady, J. M., & Matthews, P. M. (2004).

Advances in functional and structural MR image analysis and implementation as FSL.

Neuroimage, *23 Suppl 1*, S208-219. https://doi.org/10.1016/j.neuroimage.2004.07.051

Smith, S. M., Miller, K. L., Salimi-Khorshidi, G., Webster, M., Beckmann, C. F., Nichols, T. E.,

Ramsey, J. D., & Woolrich, M. W. (2011). Network modelling methods for FMRI.

Neuroimage, *54*(2), 875-891. https://doi.org/10.1016/j.neuroimage.2010.08.063

Smith, S. M., Vidaurre, D., Beckmann, C. F., Glasser, M. F., Jenkinson, M., Miller, K. L.,

Nichols, T. E., Robinson, E. C., Salimi-Khorshidi, G., Woolrich, M. W., Barch, D. M.,

Ugurbil, K., & Van Essen, D. C. (2013). Functional connectomics from resting-state

fMRI. *Trends Cogn Sci*, *17*(12), 666-682. https://doi.org/10.1016/j.tics.2013.09.016

Tavakol, S., Royer, J., Lowe, A. J., Bonilha, L., Tracy, J. I., Jackson, G. D., Duncan, J. S.,

Bernasconi, A., Bernasconi, N., & Bernhardt, B. C. (2019). Neuroimaging and

connectomics of drug-resistant epilepsy at multiple scales: From focal lesions to

macroscale networks. *Epilepsia*, *60*(4), 593-604. https://doi.org/10.1111/epi.14688

Turk, E., Scholtens, L. H., & van den Heuvel, M. P. (2016). Cortical chemoarchitecture shapes

macroscale effective functional connectivity patterns in macaque cerebral cortex. *Hum*

Brain Mapp, *37*(5), 1856-1865. https://doi.org/10.1002/hbm.23141

Turner, R. (2002). How much cortex can a vein drain? Downstream dilution of activation-related

cerebral blood oxygenation changes. *Neuroimage*, *16*(4), 1062-1067.

https://doi.org/10.1006/nimg.2002.1082

Uludag, K., Muller-Bierl, B., & Ugurbil, K. (2009). An integrative model for neuronal activity-

induced signal changes for gradient and spin echo functional imaging. *Neuroimage*,

48(1), 150-165. https://doi.org/10.1016/j.neuroimage.2009.05.051

Vaiana, M., & Muldoon, S. F. (2018). Multilayer Brain Networks. *Journal of Nonlinear Science*,

30(5), 2147-2169. https://doi.org/10.1007/s00332-017-9436-8

van den Heuvel, M. P., Scholtens, L. H., Turk, E., Mantini, D., Vanduffel, W., & Feldman

Barrett, L. (2016). Multimodal analysis of cortical chemoarchitecture and macroscale

fMRI resting-state functional connectivity. *Hum Brain Mapp*, *37*(9), 3103-3113.

https://doi.org/10.1002/hbm.23229

van der Kouwe, A. J. W., Benner, T., Salat, D. H., & Fischl, B. (2008). Brain morphometry with

multiecho MPRAGE. *Neuroimage*, *40*(2), 559-569.

https://doi.org/10.1016/j.neuroimage.2007.12.025

Van Essen, D. C., & Maunsell, J. H. (1980). Two-dimensional maps of the cerebral cortex. *J

Comp Neurol*, *191*(2), 255-281. https://doi.org/10.1002/cne.901910208

Wang, H. E., Benar, C. G., Quilichini, P. P., Friston, K. J., Jirsa, V. K., & Bernard, C. (2014). A

systematic framework for functional connectivity measures. *Front Neurosci*, *8*, 405.

https://doi.org/10.3389/fnins.2014.00405

Wei, Y., Scholtens, L. H., Turk, E., & van den Heuvel, M. P. (2019). Multiscale examination of

cytoarchitectonic similarity and human brain connectivity. *Netw Neurosci*, *3*(1), 124-137.

https://doi.org/10.1162/netn_a_00057

Wu, P. Y., Chu, Y. H., Lin, J. L., Kuo, W. J., & Lin, F. H. (2018). Feature-dependent intrinsic

functional connectivity across cortical depths in the human auditory cortex. *Sci Rep*, *8*(1),

13287. https://doi.org/10.1038/s41598-018-31292-x

Xia, M., & He, Y. (2017). Functional connectomics from a "big data" perspective. *Neuroimage*,

160, 152-167. https://doi.org/10.1016/j.neuroimage.2017.02.031

Yacoub, E., Duong, T. Q., Van De Moortele, P. F., Lindquist, M., Adriany, G., Kim, S. G., Ugurbil, K., & Hu, X. (2003). Spin-echo fMRI in humans using high spatial resolutions and high magnetic fields. *Magn Reson Med*, *49*(4), 655-664. https://doi.org/10.1002/mrm.10433

Yeh, C. H., Jones, D. K., Liang, X., Descoteaux, M., & Connelly, A. (2021). Mapping Structural Connectivity Using Diffusion MRI: Challenges and Opportunities. *J Magn Reson Imaging*, *53*(6), 1666-1682. https://doi.org/10.1002/jmri.27188

Yun, J. Y., & Kim, Y. K. (2021). Graph theory approach for the structural-functional brain connectome of depression. *Prog Neuropsychopharmacol Biol Psychiatry*, *111*, 110401. https://doi.org/10.1016/j.pnpbp.2021.110401

Zachlod, D., Ruttgers, B., Bludau, S., Mohlberg, H., Langner, R., Zilles, K., & Amunts, K. (2020). Four new cytoarchitectonic areas surrounding the primary and early auditory cortex in human brains. *Cortex*, *128*, 1-21. https://doi.org/10.1016/j.cortex.2020.02.021

Zaretskaya, N., Fischl, B., Reuter, M., Renvall, V., & Polimeni, J. R. (2018). Advantages of cortical surface reconstruction using submillimeter 7 T MEMPRAGE. *Neuroimage*, *165*, 11-26. https://doi.org/10.1016/j.neuroimage.2017.09.060

Zeki, S. (2018). The Rough Seas of Cortical Cartography. *Trends Neurosci*, *41*(5), 242-244. https://doi.org/10.1016/j.tins.2018.03.005

Zhao, F., Wang, P., & Kim, S. G. (2004). Cortical depth-dependent gradient-echo and spin-echo BOLD fMRI at 9.4T. *Magn Reson Med*, *51*(3), 518-524. https://doi.org/10.1002/mrm.10720

1217   Zhu, Y., Zhu, G., Li, B., Yang, Y., Zheng, X., Xu, Q., & Li, X. (2022). Abnormality of

1218   Functional Connections in the Resting State Brains of Schizophrenics. *Front Hum*

1219   *Neurosci*, *16*, 799881. https://doi.org/10.3389/fnhum.2022.799881

1220

1221