Dual-specificity phosphatases 13 and 27 as key switches in muscle stem cell transition from proliferation to differentiation

Takuto Hayashi[1], Shunya Sadaki[1,2], Ryosuke Tsuji[1,2], Risa Okada[3], Sayaka Fuseya[1,4], Maho Kanai[1], Ayano Nakamura[1,5], Yui Okamura[1,5], Masafumi Muratani[6], Gu Wenchao[7], Takehito Sugasawa[8], Seiya Mizuno[9], Eiji Warabi[1], Takashi Kudo[1], Satoru Takahashi[1*], Ryo Fujita[10, 11*]

[1] Laboratory Animal Resource Center in Transborder Medical Research Center, and Department of Anatomy and Embryology, Institute of Medicine, University of Tsukuba, Ibaraki 305-8575, Japan

[2] Ph.D. Program in Humanics, School of Integrative and Global Majors, University of Tsukuba, Ibaraki 305-8575, Japan

[3] JEM Utilization Center, Human Spaceflight Technology Directorate, Japan Aerospace Exploration Agency (JAXA), Ibaraki 305-8505, Japan

[4] Cellular and Molecular Biotechnology Research Institute, National Institute of Advanced Industrial Science and Technology, Ibaraki 305-8565, Japan

[5] College of Medicine, School of Medicine and Health Sciences, University of Tsukuba, Ibaraki 305-8575, Japan

[6] Department of Genome Biology, Transborder Medical Research Center, Institute of Medicine, University of Tsukuba, Ibaraki 305-8575, Japan

[7] Department of Diagnostic and Interventional Radiology, Institute of Medicine, University of Tsukuba, Ibaraki 305-8575, Japan

[8] Laboratory of Clinical Examination and Sports Medicine, Department of Clinical Medicine, Institute of Medicine, University of Tsukuba, Ibaraki 305-8575, Japan

[9] Laboratory Animal Resource Center in Transborder Medical Research Center, and Department of Laboratory Animal Science, Institute of Medicine, University of Tsukuba, Ibaraki 305-8575, Japan.

[10] Division of Regenerative Medicine, Transborder Medical Research Center, Institute of Medicine, University of Tsukuba, Ibaraki 305-8575, Japan

[11] Lead contact

*Correspondence:

Ryo Fujita, Ph.D.

Division of Regenerative Medicine, Transborder Medical Research Center, Institute of Medicine, University of Tsukuba, Ibaraki 305-8575, Japan.

E-mail: fujiryo@md.tsukuba.ac.jp

Satoru Takahashi, M.D., Ph.D.

37   Laboratory Animal Resource Center in Transborder Medical Research Center, and Department of

38   Anatomy and Embryology, Institute of Medicine, University of Tsukuba, Ibaraki 305-8575, Japan

39   E-mail: satoruta@md.tsukuba.ac.jp

Summary

Muscle regeneration depends on muscle stem cell (MuSCs) activity. These cells are typically quiescent but become activated upon muscle injury, transitioning from proliferation to differentiation. Myogenic regulatory factors, including myoblast determination protein 1 (MyoD), regulate the fate transition of MuSCs. However, the role of MYOD in this process is not yet clear. Using previously established MyoD knock-in (MyoD-KI) mice, we revealed that MyoD targets dual-specificity phosphatase (Dusp) 13 and Dusp27. In *Dusp13*:*Dusp27* double knock-out (DKO) mice, the ability for muscle regeneration after injury was found to be reduced. Moreover, single-cell RNA sequencing of MyoD-high expressing MuSCs from MyoD-KI mice revealed that *Dusp13* and *Dusp27* are only expressed in specific populations within MyoD-high MuSCs, which also express *Myogenin*. Overexpressing *Dusp13* in MuSCs causes premature muscle differentiation. Thus, we propose a model where DUSP13 and DUSP27 contribute to the fate transition of MuSCs from proliferation to differentiation during myogenesis.

Keywords: Muscle stem cell, MyoD, Dual specificity phosphatase 13:27, Regeneration, Single-cell RNA-sequencing

INTRODUCTION

Skeletal muscle stem cells (MuSCs), also called satellite cells (Mauro, 1961), play a vital role in postnatal muscle development and the regeneration of skeletal muscles (Murphy et al., 2011). MuSCs are located near the myofibers beneath the basal lamina and exist in a quiescent state. However, in response to injury, they become activated, differentiate, and generate new myofibers (Collins et al., 2005; van Velthoven and Rando, 2019). Some of the progeny of the activated MuSCs undergo self-renewal to perpetually sustain the MuSC pool throughout a lifespan (Kuang et al., 2007; Sénéchal et al., 2022; Zammit et al., 2006).

The dynamics of MuSCs are intricately regulated by a series of myogenic regulatory factors (MRFs) that are sequentially expressed during myogenesis. Quiescent MuSCs express the paired homeodomain family of transcription factors PAX7 (Relaix et al., 2005; Seale et al., 2000). The MRFs, including MyoD, Myf5, myogenin, and MRF4, act as basic helix–loop–helix (bHLH) transcription factors that bind to E-box sites. While MYOD and MYF5 have overlapping functions as essential transcription factors for muscle lineage specification (Rudnicki et al., 1993; Yamamoto et al., 2018), MYOGENIN and MRF4 are indispensable for muscle differentiation (Kassar-Duchossoy et al., 2004; Nabeshima et al., 1993). Additionally, MRFs are induced in activated MuSCs during muscle regeneration in adult mice. Interestingly, quiescent MuSCs repress the translation of specific transcripts that keep them primed to rapidly activate the myogenic program in case of injury (Zismanov et al., 2016). MicroRNAs and RNA-binding proteins primarily mediate the repression of *Myf5* and *MyoD* transcripts (Crist et al., 2012; Fujita et al., 2017; Hausburg et al., 2015; de Morrée et al., 2017). Upon injury, the block on translation is lifted, leading to the rapid translation of *MyoD* and *Myf5*, initiating the myogenic program for regeneration.

MyoD knockout (KO) mice exhibited delayed myogenic differentiation and accelerated self-renewal ability in MuSCs. Furthermore, *MyoD* KO mice on an mdx background, a model for Duchenne muscular dystrophy, have a low survival rate (Megeney et al., 1996; Yablonka-Reuveni et al., 1999) . These reports suggest that MYOD plays a critical role in determining the fate of activated MuSCs by balancing self-renewal and differentiation during myogenesis. However, the specific mechanisms by which MYOD influences the fate of activated MuSCs remain unclear.

In a prior study, we created a MyoD knock-in (KI) reporter mouse line to recapitulate the expression levels of endogenous MyoD using tdTomato fluorescence (Fujita et al., 2023). This fluorescence allowed us to identify the state of MuSCs during myogenesis. We found that the MyoD-low population, characterized by low tdTomato fluorescence, consisted mostly of undifferentiated stem-like cells. On the other hand, the MyoD-high population, with high tdTomato fluorescence, showed features of committed and differentiating MuSC (Fujita et al., 2023). To investigate the molecular mechanisms governing the stemness and commitment/differentiation of MuSCs, we collected whole transcriptome data from MyoD-low MuSCs, MyoD-high MuSCs, and quiescent MuSCs isolated from Pax7$^{YFP/YFP}$

(Pax7-YFP) mice (Fujita et al., 2023; Kitajima and Ono, 2018). The establishment of MyoD-KI mice

and the availability of whole transcriptome data at different activation stages of MuSCs based on

MYOD levels provide a valuable platform for understanding the fate determination mediated by

MYOD.

The DUSP family comprises members of the protein tyrosine phosphatase superfamily involved in

the dephosphorylation of phosphotyrosine and phosphoserine/threonine residues (Alonso and Pulido,

2016; Denu and Dixon, 1995, 1998). They interact directly with MAPK effectors, including

extracellular signal-regulated kinases 1/2 (ERK1/2), leading to their dephosphorylation and

subsequent inactivation. Therefore, DUSPs play regulatory roles in cell growth, cell differentiation,

and apoptosis (Bermudez et al., 2010; Patterson et al., 2009; Tonks, 2006). The phosphatase domain

common to DUSPs is known as protein tyrosine phosphatases (PTP). DUSPs can be classified into

two subfamilies based on the presence or absence of the MAP kinase-binding (MKB) motif or the

kinase-interacting motif (KIM). The DUSPs with the KIM domain are classified as typical DUSPs,

whereas those lacking the KIM domain are classified as atypical DUSPs (Aoki et al., 2001; Huang

and Tan, 2012).

Interestingly, the functional regulation of substrates by DUSPs does not always require phosphatase

activity. DUSPs can control the function of MAPKs by sequestering them in the cytoplasm or nucleus.

It has also been observed that some DUSPs are catalytically inactive (Reiterer et al., 2020). For

example, DUSP27 lacks phosphatase activity due to the absence of a highly conserved cysteine residue

essential for enzyme activity (Fero et al., 2014). Nevertheless, the loss of *Dusp27* in zebrafish leads to

severe disorganization of the sarcomere structure, underscoring the significance of its

pseudophosphatase function in muscle development (Fero et al., 2014). Within skeletal muscles,

DUSP5 and DUSP6, classified as typical DUSPs, regulate muscle hypertrophy and myofiber types by

inhibiting ERK1/2 signaling (Boyer et al., 2019). However, our understanding of the function of

atypical DUSPs in skeletal muscle in mammals is limited.

In the present study, we used our MyoD-KI transcriptome and various datasets to identify *Dusp13*

as a direct target for MYOD. *Dusp13* is located on chromosome 10a22.2 and encodes two distinct

DUSPs, *Dusp13a* and *Dusp13b*. Although DUSP13a is predominantly expressed in murine skeletal

muscles (Chen et al., 2004), its specific roles in skeletal muscle remain unclear. Single KO of *Dusp13*

did not show any regenerative defects, but there was compensatory upregulation of *Dusp27* in skeletal

muscles lacking *Dusp13*. To further investigate, we generated a double KO (DKO) of *Dusp13* and

Dusp27 and observed diminished muscle regeneration activity after acute muscle injury. Furthermore,

through single-cell RNA-seq (scRNA-seq) analysis, we found that the expressions of *Dusp13* and

Dusp27 are limited to specific subsets of MyoD-high expressing MuSCs that also show the expression

of late myogenic markers. Based on these findings, we propose a model in which DUSP13 and

DUSP27 instigate the transition of MuSCs from an activated/proliferative state to a differentiation

state during regeneration.

RESULTS

Dusp13 is directly regulated by MyoD activity.

We previously generated MyoD-KI mice that allow us to assess MYOD expression through tdTomato

fluorescence intensity (Fujita et al., 2023). To identify new regulators directly controlled by MYOD

activity, we utilized RNA-seq data from MyoD-low and MyoD-high expressing MuSCs from the

MyoD-KI mice, as well as freshly isolated quiescent MuSCs from Pax7$^{\text{YFP/YFP}}$ (Pax7-YFP) mice

(Figure 1A). Initially, we visualized the differentially expressed genes (FDR < 0.05) between the

MyoD-low and MyoD-high populations using a heatmap (Figure 1A). We found that 966 genes were

significantly upregulated as *MyoD* levels increased (MyoD-KI Cluster2). To further narrow down the

targets, we consulted a gene expression database of proliferating and differentiating MuSCs from

MyoD KO mice (Wang et al., 2022). We identified 1797 genes that were not upregulated in *MyoD* KO

mice (MKO Cluster2 in Figure 1B). Through a Venn diagram comparison, we combined 505 genes

from the MyoD-KI and MKO clusters. Additionally, when we included *MyoD* ChIP-seq data from

ChIP-Atlas, we found that 35 genes were enriched (Figure 1C) (Oki et al., 2018; Zou et al., 2022).

Finally, out of these 35 genes, six genes (*Ank1*, *Dmpk*, *Dusp13*, *Jsrp1*, *Rapsn*, and *Stac3*) were induced

by *MyoD* overexpression in mouse embryonic fibroblasts (MEFs) (Conerly et al., 2016) (Figure 1C

and 1D; Figure S1A and S1B). We also confirmed that *Dusp13* mRNA was predominantly enriched

in the quadriceps muscles compared to other tissues (Figure S2A).

Due to the modulation of *Dusp13* expression by *MyoD* (Figure 1E–1H), we assessed the E-box site

on *Dusp13*. We identified a highly conserved E-box site 150 bp upstream of the transcription initiation

site of *Dusp13*, according to the UCSC Genome Browser (Figure 1I). To determine if MYOD binds

to the E-box of *Dusp13*, we ligated the putative E-box site of *Dusp13* to a pGL4.10 vector to generate

a luciferase-*Dusp13*-E-box vector. We then co-transfected the plasmid with a *MyoD* expression vector

into the HEK293T cell line. The luciferase activity was significantly upregulated in the presence of

MyoD when the *Dusp13*-Ebox was present (Figure 1J). These results suggest that DUSP13 is a new

target of MYOD during myogenesis.

Single Dusp13 or Dusp27 KO mice exhibits no abnormality in muscle development or regeneration.

To investigate the role of DUSP13 in skeletal muscle, we generated *Dusp13* KO mice using CRISPR-

Cas9 technology. We confirmed that *Dusp13* expression in skeletal muscle was depleted in these

Dusp13 KO mice (Figure S2C). The *Dusp13* KO mice were born according to Mendelian laws and

grew normally, similar to the litter mate wild type (WT) control mice, with respect to body weight and

muscle cross-sectional area (CSA) of the soleus (SOL) and plantaris (PLA) muscles (Figure S3A–

S3C). We then investigated the regenerative ability in the *Dusp13* KO mice. We injected cardiotoxin

(CTX) into the tibialis anterior (TA) muscles of both WT and *Dusp13* KO mice and found no

significant histological differences in the CSA of the damaged myofibers with central nuclei at day 7

after the injury (Figure S3D and S3E). Furthermore, immunohistochemical analysis for PAX7 and

MYOGENIN showed that the number of PAX7 or MYOGENIN (+) nuclei was the same between WT

and *Dusp13* KO mice (Figure S3F–S3I). These findings indicate that there is no significant impairment

of muscle regeneration in the absence of *Dusp13*.

We hypothesized that other members of the *Dusp* family expressed in skeletal muscles could

compensate for the function of DUSP13 upon *Dusp13* deletion. Within overlapping genes between

the MyoD-KI Cluster 2 and MKO Cluster 2 in Figure 1A and 1B, we identified *Dusp27* as another

member of the *Dusp* family that is abundantly expressed in the quadriceps muscles and the heart

(Figure S2B). *Dusp27* expression was significantly higher in *Dusp13* KO muscles compared to WT

muscles, suggesting compensatory induction following the loss of *Dusp13* (Figure S4A). We also

observed a significant increase in *Dusp27* expression when *MyoD* expression increased (Figure S4B),

similar to *Dusp13* (Figure 1F). Interestingly, we found a potential Ebox site located 10 bp upstream of

the transcription initiation site of *Dusp27*, which is highly conserved among multiple mammalian

species (Figure S4C). The luciferase activity of luciferase-Dusp27-Ebox was significantly upregulated

in the presence of *MyoD* (Figure S4D), suggesting that both *Dusp13* and *Dusp27* are directly regulated

by MYOD activity. To test our hypothesis that DUSP27 compensates for the loss of *Dusp13*, we first

generated *Dusp27* KO mice using CRISPR-Cas9. We confirmed the deletion of the gene through

qPCR analysis (Figure S2D). Similar to the *Dusp13* KO mice, the *Dusp27* KO mice did not show any

differences in body weight (data not shown), muscle CSA before and after muscle CTX injury (Figure

S4E–S4J), or the number of myogenic populations labeled by either PAX7 or MYOGENIN after CTX

injury (Figure S4K–S4N). These results demonstrate that neither DUSP13 nor DUSP27 deficiency

influences muscle development and acute muscle regeneration.

Muscle regeneration is delayed in Dusp13:Dusp27 DKO mice.

To investigate the role of DUSP13 and DUSP27 in skeletal muscles, we utilized DKO mice. The body

weights and muscle weights of DKO mice were found to be similar to age-matched WT mice (data

not shown). Furthermore, both the SOL and PLA muscles exhibited normal myofiber size (Figure

S5A–S5C), and the number of MuSCs was also normal (Figure S5D–S5G). However, histological

analysis on days 3, 7, and 28 after CTX injury clearly demonstrated the essential role of DUSP13 and

DUSP27 in muscle regeneration (Figure 2A–2D). Specifically, seven days post-injury, we observed a

notable increase in PAX7 (+) cells and a decrease in MYOGENIN (+) cells (Figure 2E–2H) in DKO

muscles compared to those in WT mice. At 28 days post-injury, when MuSCs are returning to their

niche after completing the muscle regeneration process, we observed a trend towards a reduction in

the number of PAX7 (+) cells (Figure S5H and S5I), accompanied by a notable increase in the number

of perduring MYOGENIN (+) cells undergoing regeneration (Figure S5J and S5K).

To confirm our *in vivo* findings indicating a severe delay in muscle regeneration in DKO mice, we

isolated MuSCs from DKO and WT mice using magnetic cell sorting (MACS) with α7-integrin

microbeads (Fujita et al., 2023; Sénéchal et al., 2022) and cultured them for three days (Figure 2I, 2J).

Immunofluorescent analysis using antibodies against PAX7 and MYOD showed a lower proportion

of differentiating PAX7 (-) and MYOD (+) cells in the DKO mice (Figure 2I, 2J)., indicating that

DUSP13/DUSP27 are necessary for the transition from activation to differentiation during myogenesis.

Since the lower differencing PAX7 (-), MYOD (+) cells in DKO mice, we observed a higher number

of proliferating Ki67 (+) MuSCs in DKO mice (Figure S6A–S6C). As expected, both myotube width

and fusion index exhibited a significant reduction in DKO mice relative to WT mice after six days of

MuSCs differentiation (Figure 2K–2M). These findings suggest that DUSP13 and DUSP27 dictate the

balance between the proliferation and differentiation of MuSCs during myogenesis.

Single-cell RNA sequencing revealed that Dusp13 and Dusp27 are highly expressed in the late

differentiation stage of MuSCs.

While MYOD activates *Dusp13* and *Dusp27* through their E-box, and *Dusp13:Dusp27* deficient mice

exhibit diminished muscle regeneration activity, the molecular mechanisms by which DUSP13 and

DUSP27 regulate myogenesis remain elusive. Additionally, the expression patterns of *Dusp13* and

Dusp27 in different stages of myogenic progenitors within the highly heterogeneous MuSC

populations are not well-defined. In order to gain a more comprehensive understanding of the

molecular signatures of *Dusp13* and *Dusp27*-expressing myogenic cells, we conducted single-cell

RNA sequencing (scRNA-seq) on MyoD-high expressing cells isolated from cultured MuSCs

obtained from MyoD-KI mice (Figure 3A). This approach allowed us to enrich activated/differentiated

MyoD (+) cells (Fujita et al., 2023). MuSCs from MyoD-KI mice were cultured for five days in a

growth medium, and only live MyoD-high MuSCs with PI (-) negative populations were sorted by

fluorescence-activated cell sorter (FACS) for subsequent scRNA-seq library preparation (Figure 3A).

Utilizing the Seurat single-cell analysis package, we identified five distinct clusters of cells (C0–C4)

presented in the uniform manifold approximation and projection (UMAP) (Figure 3B). Each cluster

exhibited a unique transcriptomic fingerprint characterized by the top 10 differentially expressed genes

(Figure S7A).

To further understand the characteristics of these clusters, we conducted a Gene Ontology (GO)

analysis on differentially expressed genes (DEGs) among the clusters (Figure S7B–S7F). C0 and C4

showed upregulation of genes associated with chromosome segregation and DNA replication (Figure

S7B and S7F). C1 and C3 displayed strong enrichment for terms related to muscle differentiation,

maturation, and contraction (Figure S7C and S7E). C2 was characterized by genes associated with

ameboidal-type cell migration and extracellular organization (Figure S7D), akin to the previously

reported MyoD-low MuSC signatures (Fujita et al., 2023). Next, we investigated cell cycle and

myogenic markers, including *MKi67, E2f1, Ccnd1, Ccnd3, Pax7, Myf5, Myod1, Myogenin, Cdkn1c*

(p21), Myh3, Dusp13, and Dusup27 on UMAP (Figure 3C). *MyoD* is ubiquitously expressed across

all clusters but with variations between the clusters (Figure 3C). *Mki67* and *E2f1*, which are

proliferative markers, were highly expressed in C0 and C4 but not in C1, C2, and C3 (Figure 3C).

Undifferentiated MuSC markers, *Pax7* and *Myf5*, were exclusively enriched in C0 and C2 (Figure 3C).

Notably, *Dusp13* and *Dusp27* exhibited pronounced enrichment in C1 and C3, characterized by

upregulation of genes associated with myogenic differentiation and maturation (Figure 3C).

Considering the GO term and UMAP, C0 and C4 represent proliferative states, while C1 and C3

contain the differentiation and maturation state of MuSCs. On the other hand, C2 reflects the self-

renewing MuSCs returning to the undifferentiated state. Furthermore, we used Monocle to analyze

these MYOD-high expressing MuSC clusters and infer MuSC state trajectories (Figure 3D–3F). The

cell state trajectory model revealed three branches for the MyoD-high-expressing MuSCs. These

branches suggested that some clusters were transitioning towards further proliferation (C1, C4),

myogenic differentiation/maturation (C1, C3), or self-renewal (C2) to return to an undifferentiated

quiescent state (Figure 3D). We then focused the Monocle analysis on selected genes related to

myogenic development/regeneration and cell cycle (*Pax7, Myf5, Myod1, Myogenin, Myf6, Myh3,*

Myh4, Hes6, Acta1, ccnd1, ccnd3, E2f1, Mki67, Cdkn1c (p21), Dusp13, and Dusp27). Cluster 3

contained *Dusp13* and *Dusp27*, along with *Myogenin, Myh3, Hes6, Acta1,* and *Cdkn1c (p21)* (Figure

3E–3F). When visualizing the pseudotime progression axis of these selected genes, we observed that

cells from C0 and C4, which expressed genes associated with the cell cycle (*Mki67* and *E2f1*) but had

lower expression of *Pax7* and *Myf5*, were located at the initiation point of pseudotime (Figure 3E–3F).

Cells from C2, which showed lower levels of *Mki67* and *E2f1* but higher levels of *Pax7, Myf5*, and

Ccnd1, were mainly found in the middle point of pseudotime, suggesting that C2 represents a self-

renewing MuSCs state (Brett et al., 2020; Fujita et al., 2023). In contrast, cells from C1 and C3, which

highly expressed genes associated with myogenic differentiation and maturation, were located distally

at the endpoint of pseudotime (Figure 3E and 3F), indicating myogenic differentiation and maturation

states (Figure 3E and 3F). *Dusp13* and *Dusp27*, enriched in C1 and C3, progressed alongside myogenic

differentiation markers and *Cdkn1c (p21)* (Figure 3E and 3F). Overall, the clustering and cell state

trajectory analysis based on the scRNA-seq data of MyoD-high expressing MuSCs suggest that

Dusp13 and *Dusp27* are enriched only in the late stage of MyoD-positive MuSCs co-expressing

Myogenin, Myh3, and *Cdkn1c*. They may regulate the switch from a proliferative state to a myogenic

differentiation state.

Dusp13 drives precocious muscle differentiation through phosphatase activity in an independent
manner.

Since our scRNA-seq analysis of MyoD-high expressing cells indicates that *Dusp13* and *Dusp27* are
expressed in the subset of cells expressing late myogenic markers, such as myogenin. To investigate
whether DUSP13 plays a role in initiating the differentiation process during myogenesis, we
transduced FACS-isolated MuSCs from adult Pax7-YFP mice with an adenovirus vector expressing
Dusp13 or GFP under the CMV promoter (Figure 4A and Figure S8A and S8B). Two days post-
infection, we observed a significant reduction in the number of MuSCs transduced with *Dusp13*
compared to the GFP control (Figure 4B). Notably, immunofluorescent analysis using antibodies
against PAX7 and MYOGENIN showed that *Dusp13* overexpression led to an induction of
MYOGENIN expression in PAX7 (+) cells, which are typically not detected under normal culture
conditions (Figure 4C and 4D). Furthermore, when we cultured MuSCs overexpressing *Dusp13* for
seven days and assessed their myotube formation ability, we found that both the myotube width and
fusion index were lower compared to the GFP control (Figure 4E–4G).

To investigate whether the phosphatase activity of DUSP13 is involved in myogenic induction, we
transduced MuSCs from adult PAX7-YFP mice with pAd-Dusp13_Mut, which lacks enzymatic
activity. Surprisingly, we discovered that DUSP13 without phosphatase activity still enhanced the
population of PAX7 (+) and MYOGENIN (+) cells to the same extent as normal DUSP13 (Figure
S8C and S8D). Taken together, these results suggest that DUSP13 acts as a driver for myogenin during
myogenic progression, which is mediated by phosphatase activity-independent mechanisms.

DISCUSSION

Skeletal muscles have a remarkable capacity for regeneration, primarily relying on the activity of
MuSCs. The regulation of MuSC dynamics involves the sequential expression of multiple
transcription factors during myogenesis. MYOD plays a crucial role in determining the myogenic
lineage specification during embryogenesis (Davis et al., 1987; Rudnicki et al., 1992, 1993). In adult
mice regeneration, MYOD is induced in activated MuSCs, modulating the balance between self-
renewal and differentiation (Megeney et al., 1996; Sabourin et al., 1999; White et al., 2000; Wood et
al., 2013; Yablonka-Reuveni et al., 1999). However, the molecular mechanisms by which MYOD
governs fate determination in MuSCs remain poorly understood. The diverse and highly
heterogeneous nature of MuSCs, particularly during myogenesis, has posed a challenge in deciphering
molecular signatures in MYOD-expressing MuSCs due to a lack of appropriate tools. This obstacle
has hindered the elucidation of MYOD's role in fate determination. To address this, we previously
engineered MyoD-KI mice, providing a platform to distinguish and isolate undifferentiated (MyoD-
low) and committed MuSCs (MyoD-high).

In this study, we integrated our previously established whole transcriptome data encompassing
quiescent, undifferentiated (MyoD-low), and committed (MyoD-high) MuSCs from MyoD-KI
reporter mice (Fujita et al., 2023) with publicly available databases from *MyoD* KO mice (Wang et al.,
2022), Chip-atlas (Oki et al., 2018; Zou et al., 2022), and *MyoD* overexpression (Conerly et al., 2016).
Through this integration, we identified *Dusp13* as a novel molecular switch regulated by MyoD,
stimulating the initiation of myogenic differentiation. To understand the function of DUSP13, we
generated *Dusp13* KO mice using CRISPR-Cas9. Surprisingly, muscle development and regeneration
abilities in *Dusp13* KO mice mirrored those of their WT counterparts. This led us to investigate the
compensatory role of other members of the DUSP family. We found that *Dusp27* was also directly
regulated by MyoD. Although *Dusp27* KO mice generated in this study did not exhibit marked
differences in muscle development and regenerative ability following CTX injury, the combined
deficiency of *Dusp13* and *Dusp27* in mice resulted in substantial delays in muscle regeneration after
CTX injury. These findings suggest a compensatory interplay between DUSP13 and DUSP27 during
muscle regeneration. The quantities of PAX7 (+) quiescent MuSCs were identical between WT and
DKO mice before injury. However, at 28 days post-CTX injury, when muscle regeneration was almost
complete, DKO mice had lower levels of PAX7 (+) cells compared to WT mice. Conversely, DKO
mice had more MYOGENIN (+) cells remaining at day 28 post-injury compared to WT mice. At day
7 post-injury, DKO mice had higher numbers of PAX7 (+) cells and fewer MYOGENIN (+) cells.
These results suggest that the absence of *Dusp13* and *Dusp27* promotes the imbalance of the fate
determination processes in MuSCs following muscle injury. *In vitro* analyses employing isolated
MuSCs from DKO mice supported these findings, showing that the absence of *Dusp13* and *Dusp27*
promotes proliferation rather than differentiation. Consequently, the delay/failure in transitioning from
the activation/proliferation phases to differentiation, attributed to the deficiency of *Dusp13* and
Dusp27, disrupts the muscle repair process in adult mice.
To further investigate the regulatory dynamics of *Dusp13* and *Dusp27* in fate determination, we
conducted scRNA-seq on MyoD-high expressing MuSCs isolated from MyoD-KI mice. Intriguingly,
both *Dusp13* and *Dusp27* expression were enriched in specific clusters (C1 and C3) within the MyoD-
high expressing MuSC populations. These clusters were characterized by increased expression of late
myogenic markers, such as *Myogenin, Myh3,* and *Cdkn1a*, and decreased expression of *Pax7*, *MKi67*,
and *Myf5*. Pseudotime cell state trajectory analysis was used to distinguish the fate determination of
MuSCs during myogenesis. Cluster C0 represented a proliferative state with high expression of
undifferentiated markers (*Pax7* and *Myf5*) and cell cycle markers (*MKi67* and *E2f1*), which projected
to further proliferate (C4), differentiate/mature (C1 and C3), or self-renew to return to an
undifferentiated quiescent state (C2). We believe that the scRNA-seq data from our MyoD-KI mice
will provide a valuable resource for uncovering new molecular mechanisms involved in myogenic fate

transition (proliferation, differentiation, or self-renewal/quiescence) of activated MuSCs during

myogenesis.

Given the limited expression of *Dusp13* and *Dusp27* in C1 and C3, we hypothesized that these two

genes play a crucial role in the transition from proliferation to differentiation. In our study, we

overexpressed *Dusp13* in MuSCs using an adenovirus vector. While the expression of PAX7 and

MYOGENIN expression is typically mutually exclusive during myogenesis (Olguin et al., 2007), a

small subset of cells does exhibit positivity for both PAX7 and MYOGENIN, representing an

intermediate population initiating differentiation (Halevy et al., 2004). In our model, a limited number

of PAX7 (+) and MYOGENIN (+) cells were also identified. Notably, *Dusp13* overexpression

significantly augmented the population of PAX7 (+) and MYOGENIN (+) cells. Collectively, our

findings align with the scRNA-seq analysis results, supporting the notion that the MYOD-DUSP13

and DUSP27-MYOGENIN regulatory network governs the initiation of differentiation for muscle

regeneration in adult mice.

There are limited studies on the regulatory roles of the DUSP family in myogenesis. Therefore,

the molecular targets modulated by DUSP13 and DUSP27 in MuSCs are currently unknown. The

DUSPs family possesses the dual ability to dephosphorylate phosphor-serine/threonine and phospho-

tryosoine residues. Currently, 25 different DUSPs are identified and categorized into six subfamilies

(Alonso and Pulido, 2016). DUSP13, an atypical DUSP, has versatile dephosphorylation substrates,

including mitogen-activated protein kinases (MAPKs). Forced expression of pAd-Dusp13_Mut vector,

without phosphatase activity, into MuSCs led to the emergence of PAX7 (+) and MYOGENIN (+)

cells *in vitro*, indicating phosphatase-independent mechanisms mediated by DUSP13 for myogenic

differentiation. Interestingly, DUSP27 is recognized as a pseudo-phosphatase, characterized by

catalytically inactive phosphatase activity. Various pseudo-phosphatases are reported to possess

pivotal functions in different organisms and pathological conditions (Reiterer et al., 2020). Inhibition

of ERK1/2 promotes myogenic differentiation and fusion in both mice and avian models (Eigler et al.,

2021). The DUSP family, lacking a kinetic-interacting motif, inhibits ERK1/2 signaling (Hu and

Mivechi, 2006; Huang et al., 2000; Marti et al., 2001), suggesting that the phosphatase-independent

activities of DUSP13 and DUSP27 potentially regulate ERK signaling in MuSCs. Future

investigations are expected to unravel the detailed molecular mechanisms by which DUSP13 and

DUSP27 orchestrate fate switching in activated MuSCs during myogenesis.

In summary, using our previously established MyoD-KI mice model, we identified *Dusp13* and

Dusp27 as novel MYOD target genes that play a pivotal role in the fate switching of MuSCs.

Furthermore, we demonstrated that *Dusp13* and *Dusp27* expressions are highly enriched in

differentiation clusters revealed by scRNA-seq of MyoD-high expressing MuSCs. Beyond both being

new determinants for the initiation of myogenic differentiation, further analysis of scRNA-seq data

within each cluster will yield valuable insights into identifying more potent fate determinants in

MuSCs during myogenesis.

Limitation of the study

Although our study shows that DKO mice have reduced muscle regeneration ability, there are certain

limitations that need to be taken into account in the interpretation of our results, given that *Dusp13*

and *Dusp27* deletion is systemically induced. To further investigate muscle regeneration, we isolated

MuSCs from DKO mice and found that they exhibited impaired myotube formation compared to WT

mice. However, optimal regeneration requires coordination among multiple cell types, including

inflammatory cells (Arnold et al., 2007; Fujita et al., 2014) and fibro-adipogenic progenitors (Joe et

al., 2010; Uezumi et al., 2010). The expression of *Dusp13* and *Dusp27* is comparatively lower in other

tissues than in skeletal muscles (Figure S2A–S2B). Nevertheless, further investigations are necessary

to reveal the inflammatory status in DKO mice both before and after injury. Under normal conditions,

the size, weight, and histology of skeletal muscles in DKO mice are indistinguishable from those in

WT mice. The reason why the loss of *Dusp13* and *Dusp27* does not affect muscle development during

embryonic development is unclear. It is possible that other members of the DUSP family might

compensate for their roles in embryonic and postnatal muscle formation, or the downstream targets of

DUSP13 and DUSP27 may diverge between muscle regeneration and embryonic myogenesis. Our

demonstration of *Dusp13* overexpression promoting MYOGENIN in PAX7(+) MuSCs indicates

precocious differentiation. Moreover, our scRNA-seq data of MyoD-high MuSCs clearly show that

the late myogenic stage MyoD-positive MuSCs express *Dusp13* and *Dusp27*. Expanding on the

molecular functions of the genes within these clusters will contribute to deciphering the molecular

targets of DUSP13 and DUSP27.

Acknowledgments

We would like to thank Satoshi Yamazaki and Yuji Yamazaki from the Division of Stem Cell Therapy, University of Tsukuba, for their valuable technical assistance in cell sorting using flow cytometry. This work was supported by a grant from the MEXT Leading Initiative for Excellent Young Researchers (R.F.), and the Grant-in-Aid for Early-Career Scientists (21K17679; R.F.). This work was also supported, in part, by grants to R.F. from the Takeda Science Foundation and Mochida Memorial Foundation for Medical and Pharmaceutical Research. T.H. and S.S were funded by JSPS Research Fellowships for Young Scientists (22J10954 (T.H.) and 23KJ0287 (S.S.)). Furthermore, this work received support from the Grant-in-Aid for the Japan Aerospace Exploration Agency (14YPTK-407 005512; S.T.) and the Grant-in-Aid for Scientific Research on Innovative Area from MEXT (18H04965; S.T.).

Author contributions

T.H., S.M., S.T., and R.F. conceived the study and designed the experiments. T.H., S.S., R.T., R.O., S.F., M.K., A.N., Y.O., M.M., S.M., D.S., E.W., T.K., and R.F. performed the experiments and bioinformatic analysis. T.H., G.W., T.S., and R.F. performed single-cell RNA-seq analysis. T.H., S.T., and R.F. wrote the manuscript. All authors contributed to the analysis and interpretation of results and reviewed the manuscript.

Competing interests

The authors declare no competing financial interests.

Additional information

Supplementary information

Supplementary Information is available for this paper.

Corresponding authors

Correspondence and requests for materials should be addressed to Ryo Fujita.

 STAR METHODS

RESOURCE AVAILABILITY

Lead contact

Further information and request for reagents may be directed to and will be fulfilled by the lead contact,

Ryo Fujita (fujiryo@md.tsukuba.ac.jp).

Materials availability

All unique reagents generated in this study are available from the lead contact in accordance with the

relevant material transfer agreements.

Data and code availability

● Single-cell RNA-seq data have been deposited in the NCBI Gene Expression Omnibus (GEO).

The accession number is listed in the key resources table.

● This paper does not report original code.

● Any additional information required to reanalyze the data reported in this paper is available from

the lead contact upon request.

EXPERIMENTAL MODELS AND SUBJECT DETAILS

Mice

Mice were kept in a specific pathogen-free environment at the Laboratory Animal Resource Center at

the University of Tsukuba, Ibaraki, Japan. All experiments were performed in compliance with the

relevant Japanese and institutional laws and guidelines and were approved by the University of

Tsukuba Animal Ethics Committee (authorization number: 23-046). The study used male and female

WT, *Dusp13* KO, *Dusp27* KO, *Dusp13*/*Dusp27* DKO mice, unless otherwise stated. Quiescent

MuSCs were directly isolated from the leg muscles of 15–16-week-old male Pax7-YFP ($Pax7^{YFP/YFP}$)

(Kitajima and Ono, 2018) (Fujita et al., 2023). The TA muscle was acutely injured by injecting 50 μL

of 10 μM cardiotoxin (Funakoshi Co., Ltd., Tokyo, Japan) under anesthesia using isoflurane inhalation.

METHOD DETAILS

Generation of *Dusp13* and *Dusp27* KO mice

The *Dusp13* and *Dusp27* KO mice were generated using the CRISPR/Cas9 system. Two target

sgRNAs were designed as follows; 5′-GTAGGGTTCACACAGAACGG-3′ and 5′-

ATGTCACAGTAAGCGTACGG-3′ for targeting all exons of the *Dusp13* gene on mouse

chromosome 14, and 5′-GCATCCTTCCACATCAACCC-3′ and 5′-

GGTAAGGGTGCTCTCGACAA-3′ for targeting regions located in introns 2 and 3, respectively, of

the *Dusp27* gene on mouse chromosome 1. Each target sgRNA was cloned into the guide RNA-Cas

co-expression plasmid px330-U6-Chimeric_BB-CBh-hSpCas9 vector (Addgene, Watertown, MA,

USA). A mixture of the four cloned plasmids was microinjected into the pronuclei of C57BL/6 J

fertilized eggs, and living embryos were transferred to pseudo-pregnant ICR female mice. Genomic

DNA screening was performed using PCR with the primers and protocols listed in Table S1.

Muscle stem cell isolation and culture

MuSCs were isolated from the hindlimb muscles of 11–13-week-old mice following MACS using a

Satellite Cell Isolation Kit (Miltenyi Biotec, Bergisch Gladbach, Germany) and anti-integrin α7

microbeads (Miltenyi Biotec), as previously described (Fujita et al., 2023). The adult hindlimb

muscles were minced in ice-cold Ham's F12 medium. The minced muscles were then incubated in

0.1% collagenase D (Roche, Basel, Switzerland) and 0.1% trypsin in Ham's F12 (Gibco, Waltham,

MA, USA) with 1% penicillin-streptomycin for 45 min at 37 °C on a shaker for three rounds of

digestion. After each round, the supernatant was collected in a 50 mL tube containing 8 mL of fetal

bovine serum (FBS) (Gibco) on ice, and fresh 0.1% trypsin/0.1% collagenase digestion buffer was

added to the tube for the next round of digestion. The collected supernatant was filtered through a

sequence of 100 and 40 μm nylon mesh strainers and then centrifuged at $400 \times g$ for 10 min at 4 °C.

The pellets were resuspended in PBS containing 2% FBS and processed for MACS or flow cytometry.

The MuSCs were cultured in a growth medium containing 39% DMEM (Gibco), 39% Ham's F12,

20% FBS (Gibco), and 1% UltroserG (Pall Life Sciences, Port Washington, NY, USA). Differentiation

media consisted of 2% horse serum in Dulbecco's modified Eagle's medium (DMEM). Gelatin (0.2%)-

coated dishes were used for MuSC culture unless otherwise indicated.

Immunofluorescence and histological analysis

Cultured cells were washed twice with PBS after removing the media and fixed with 4%

paraformaldehyde (PFA) for 15 minutes. The fixed cells were blocked with 2% BSA/PBS for 1 hour

at room temperature and then washed three times with PBS. Isolated single EDL myofibers (Sénéchal

et al., 2022) were fixed using 2% PFA for 15 min, permeabilized with 0.1% Triton in PBS, and blocked

with 5% goat serum in 0.1% Triton/PBS for 30 min on a shaker. The cultured cells and single EDL

myofibers were incubated with primary antibodies at 4 °C overnight. The myotube diameter and

nucleus number were measured using the ImageJ software (NIH, Bethesda, MI, USA). The myotube

diameter was defined as the average diameter at the three ends (both ends and the center) for each

MYHC-positive and multinucleated myotube. TA muscles were neatly stacked on top of each other.

The Achilles tendon side was vertically mounted on tragacanth gum on a cork disc and quickly frozen

in isopentane cooled in liquid nitrogen. Thin frozen muscle sections (8 μm) were subjected to

immunohistochemical analyses. The sections were fixed with 4% PFA for 10 min at room temperature.

For PAX7 staining, sections underwent antigen retrieval with DAKO buffer (Agilent, Santa Clara, CA,

USA) for 10 min at 110 °C, followed by blocking with M.O.M. blocking reagent (Vector Laboratories,

Newark, CA, USA). The sections were then incubated with primary antibodies overnight at 4 °C.

Tissue sections were mounted with VECTASHIELD Vibrance Antifade Mounting Medium with

DAPI (Vector Laboratories). Primary antibodies were against PAX7 (Developmental Studies

Hybridoma Bank (DSHB)), MyoD (sc-377460; SantaCruz), LAMININ (sc-59854; SantaCruz),

MYOGENIN (124800; Abcam, Cambridge, UK), MYHC (DSHB; MF20), and KI67 (MA5-14520;

Invitrogen, Waltham, MA, USA). For immunofluorescence, secondary antibodies conjugated with

Alexa Fluor-488, Alexa Fluor-546, Alexa Fluor-555, Alexa Fluor-633, and Alexa Fluor-647 were used.

The secondary antibodies included anti-mouse IgG1, anti-mouse IgG2b, anti-rat IgG, and anti-rabbit

antibodies (Thermo Fisher Scientific, Waltham, MA, USA). For HE staining, the SOL, PLA, and TA

muscle sections were stained with Mayer's hematoxylin (Wako, Richmond, VA, USA) for 10 min.

After washing in warm running tap water (50 ± 5 °C) for 3 minutes, eosin (Wako) was applied for 1

min. The sections were then dehydrated using two changes (5 s each) of 95% ethanol and two changes

(5 s each) of 100% ethanol (Wako), followed by three changes of xylene (Wako) for 5 min each.

Finally, the sections were mounted with DPX mounting medium (Sigma-Aldrich, St. Louis, MA,

USA) and observed with a BIOREVO BZ-X800 microscope (Keyence, Osaka, Japan). The CSA of

myofibers was assessed using the same microscope system and a hybrid cell count application

(Keyence).

Flow cytometry analysis and cell sorting

MACS-isolated MuSCs were cultured for 5 days. They were detached using 0.25% trypsin/EDTA,

then centrifuged and resuspended in 2% FBS/PBS. Propidium iodide (PI) was added to the cells at a

1:500 (vol/vol) ratio and incubated on ice for 10 min. The cells were either analyzed using a flow

cytometer (CytoFLEX S; Beckman Coulter, Brea, CA, USA) or sorted using a MoFlo XDP flow

cytometer (Beckman Coulter). Debris and dead cells were excluded using forward scatter, side scatter,

and PI gating. Gates were defined based on WT control fluorescence. The data were analyzed using

FlowJo^{TM} software ver. 10.7.1 (BD Biosciences, Franklin Lakes, NJ, USA).

DNA construct

The *MyoD*, *Dusp13*, and *Dusp27* genes were amplified using PrimeStar GXL DNA Polymerase

(Takara Bio, Shiga, Japan) with soleus cDNA as the template. After confirming the size of the PCR

products using 1% agarose gel electrophoresis, the target PCR product was purified from the gel. The

purified DNA fragment of *Dusp13* was digested with EcoRI. The digested fragments were then ligated

to pENTR3c-IRES-EGFP, which was also digested with the same restriction enzymes, using Ligation

High (TOYOBO, Osaka, Japan). DUSP13 deficient in phosphatase activity (=Dusp13_Mut) was

generated using the PrimeSTAR® Mutagenesis Basal Kit (Takara Bio). The Dusp family contains the

amino acid cysteine, which is essential for enzyme activity (Alonso et al., 2004). The present mutant

was generated using Dusp13_D97A and Dusp13_C128S primers (Table S1) to replace aspartic acid

at position 97 and cysteine at position 128 with alanine and serine, respectively. The reaction was then

transformed into One shot TOP10 Competent cells (Thermo Fisher Scientific) and seeded on

kanamycin-LB agar plates. A single colony was grown in kanamycin-LB medium, and after

confirming each insert by colony PCR and DNA sequencing, plasmid purification was performed

using the FastGene Plasmid Mini Kit. Clones without any mutations in the product were selected. The

LR reaction of the Gateway system was performed to recombine pENTR3c-Dusp13-IRES-EGFP into

pAd/CMV/V5-DEST (Thermo Fisher Scientific). In total, 100 ng of the purified plasmid and 300 ng

of pAd/CMV/V5-DEST were mixed at room temperature and reacted according to the manual for

Gateway LR Clonase II Enzyme Mix (Thermo Fisher Scientific). The 5 μL reaction was subsequently

transformed into 50 μL of competent cells and seeded on ampicillin-LB agar plates. Single colonies

were cultivated in LB medium, and confirmation of the insert was achieved through colony PCR. The

confirmed plasmid was prepared in large quantities using the PureLink HiPure Plasmid Midiprep Kit

(Thermo Fisher Scientific). The plasmid concentration was measured using a NanoDrop 2000

Spectrophotometer (Thermo Fisher Scientific), and 20 μg of the plasmid was digested with the

restriction enzyme PacI. Verification of complete digestion by PacI was performed through 1%

agarose gel electrophoresis, and the linearized plasmid was purified via phenol:chloroform extraction

and ethanol precipitation. To generate the luciferase vector, the purified DNA fragment of the *MyoD*

gene was cloned into a pAAV-CMV vector using the In-Fusion Cloning HD Kit (Takara Bio). The

DNA fragment corresponding to the 300 bp 5′ upstream region (−250 to +50) of *Dusp13* or the 1,200

bp upstream region (−1,000 to +200) of *Dusp27* from EDL muscle DNA of C57BL/6 mice was

inserted into a pGL4.10 promoter-less luciferase vector (Promega, Madison, WI, USA) digested with

XhoI. The resulting constructs were named pGL4-Dusp13 or pGL4-Dusp27. The primer sequences

for the construction of vectors are provided in Table S1.

Adenovirus production and infection

293A cells were seeded on BD BioCoat Collagen I 60 mm Culture Dishes (IWAKI) using the medium

(DMEM and 10% FBS). Purified 5 μg of pAd/CMV/V5-DEST plasmid was transfected using Fugene

HD (Promega) when the 293A cells were 80% confluent. When the cells began to float, the bottom

surface of the dish was scraped off using a scraper and collected in a 15 mL High-Clarity

Polypropylene Conical Tube 17 × 120 mm style (FALCON) (primary virus). The primary virus

underwent repeated freeze-thaw cycles in a water bath at 37 °C and in liquid nitrogen. This freeze-

thaw process was repeated three times. Finally, it was frozen again and stored at −80 °C. Next, 2 mL

of the primary virus was thawed at 37 °C and centrifuged for 15 min at 3,000 rpm (KUBOTA 2800).

The supernatant was added to 293A cells (80% confluent) cultured in a 100 mm Collagen-Coated Dish

Collagen I (IWAKI) and infected for 1 hour at 37 °C in a CO_2 incubator. Then, 8 mL of medium was

added. When the cells floated, they were scraped off using a scraper and collected in 15-mL tubes.

The freeze-thaw process was repeated three times. Finally, it was frozen again and stored at −80 °C.

This virus was created as the 2nd virus. This procedure was repeated five times, and the 5th virus was

stored at −80 °C. Next, to purify the virus, the fifth virus was freeze-thawed three times and

ultracentrifuged at 23 k rpm at 4 °C for 1.5 hours in the Super-centrifuge Optima XE-90

Ultracentrifuge (Beckman Coulter) using Ultra-Clear Centrifuge Tubes (Beckman Coulter). After

ultracentrifugation, recombinant adenovirus was extracted from the second viral layer from the bottom.

The dialysis buffer was prepared in three heat-resistant plastic bottles by adding 1976 mL of MilliQ,

20 mL of 1 M Tris HCl (pH 8.0), and 4 mL of 1 M $MgCl_2$, and the bottles were then autoclaved. For

dialysis, the recombinant adenovirus extract was injected into Slide-A-Lyzer Dialysis Cassettes

(Thermo Fisher Scientific) and dialyzed in a dialysis buffer at 4 °C with stirring. The dialysis buffer

was changed three times every 12 hours. The purified recombinant adenovirus was stored at −80 °C.

Infectious titers were measured using the Adeno-X Rapid Titer Kit (Takara Bio). MuSCs were

cultured in a growth medium (GM) composed of DMEM (Gibco, high glucose, 11960-044), 10% FBS,

1% Pen Strep (Gibco), and 2 mM L-glucose (Gibco) at 37 °C under 5% CO_2. To differentiate into

myotube cells, cells were seeded at 1×10^4 cells/well (12-well plate). Differentiation was induced by

replacing the growth medium with a differentiation medium (DM) consisting of DMEM, 2% horse

serum (Gibco), 1% Pen Strep (Gibco), and 2 mM L-glucose (Gibco). After 48 hours from the start of

differentiation, the medium was replaced with a fresh DM and incubated with adenovirus at a

multiplicity of infection (MOI) of 250 in a differentiation medium. Two days after the virus

transduction, the cells were fixed with 4% PFA/PBS and permeabilized with 0.1% Triton X-100/PBS.

The fixed cells were blocked with 2% BSA/PBS for 1 h at room temperature and washed three times

with PBS. The myotube cells were then reacted with monoclonal anti-PAX7 antibody (DSHB) (1:100)

and anti-MYOGENIN (ab124800; Abcam) (1:100) in 2% BSA/PBS at 4 °C overnight. After washing

with PBS, they were incubated with appropriate Alexa Fluor-conjugated secondary antibodies

(Thermo Fisher Scientific) for 1 hour at room temperature and mounted with 20 mM Tris and 90%

glycerol. The detection was photographed with a fluorescence microscope BIOREVO (BZ-X800;

Keyence).

Luciferase assay

To determine if MyoD could activate the *Dusp13* or *Dusp27* promoter, we co-transfected 100 ng of

pGL4-Dusp13 or pGL4-Dusp27 with 50 ng of pRL-TK vectors (Promega) and varying amounts (0,

10, 20, 100, 200 ng) of pAAV MyoD expression vectors into AAVpro 293T cells (Takara Bio) using

the FuGENE HD transfection reagent (Promega). Twenty-four hours after transfection, the cells were

harvested and assayed using the Dual-Luciferase Reporter Assay System (Promega). Luciferase activity was measured using the GloMax® 20/20 Luminometer (Promega) and normalized based on Renilla luciferase activity from the pRL-TK vector.

Motif enrichment analysis

The presence of Ebox sites in the promoter regions of genes was predicted using the Find Individual Motif Occurrences (FIMO) algorithm from the MEME Suite (Grant et al., 2011), utilizing the position weight matrix available from JASPAR (https://jaspar.genereg.net/). This analysis was conducted from 1 kbp upstream to 500 bp downstream of the transcriptional start site of *Dusp13*/*Dusp27*.

RNA isolation and RT-qPCR

Total RNA was extracted from various tissues and MuSCs using ISOGEN (NIPPON GENE, Chiyoda, Tokyo, Japan). We synthesized first-strand cDNA using the QuantiTect Reverse Transcription Kit (Qiagen, Hilden, Germany). Polymerase chain reaction (PCR) was then performed using a THUNDERBIRD SYBR qPCR system (TOYOBO) and a TP850 Thermal Cycler Dice Real Time System (Takara Bio, Shiga, Japan). The primer sequences are listed in Table S1. The relative amount of each transcript was normalized to the number of beta-actin transcripts.

Single-cell RNA-seq analysis

MACS-isolated MuSCs from MyoD-KI mice (Fujita et al., 2023) were cultured for five days and then detached using 0.25% trypsin/EDTA. The cells were centrifuged and resuspended in 2% FBS/PBS. The cells were incubated with PI at a 1:500 (v/v) ratio. Debris and dead cells were excluded using forward scatter, side scatter, and PI gating. Live MyoD-tdTomato-high MuSCs were sorted using a MoFlo XDP flow cytometer (Beckman Coulter) based on our previous study's gate of MyoD-tdTomato-high population. Single RNA-seq libraries from MyoD-tdTomato-high MuSCs were generated using a chromium controller (10× Genomics, Pleasanton, CA, USA) following the manufacturer's instructions. Approximately 10,000 cells were processed using the Chromium Next GEN Single Cell 3'Reagent Kits v3.1 (10× Genomics). The sequencing was performed on a NovaSeq 6000 (Illumina, San Diego, CA, USA) operated by the Rhelixa Corporation (Tokyo, Japan). The fastq files were processed using default parameters for adaptor sequence filtering, and low-quality reads were removed to obtain clean data. Feature-barcode matrices were then generated by aligning the reads to the mouse genome (mm10 Ensemble: version 92) with CellRanger v3.1.0. The scRNA-seq data was processed using the standard Seurat pipeline. For quality control, the range of gene reads per cell was set between 300 and 6000. Additionally, the filtering threshold for mitochondrial gene proportion was established at 15%. The "FindVariableFeature" function was used to determine the highly variable genes. Cell clustering was determined using the "FindNeighbors" and "FindCluster" functions in the

Seurat package. To investigate the proliferation of genes involved in the development of muscle stem

cells, the "Monocle2" package was applied for conducting pseudotime analysis. The biological

function analysis was performed using GO analysis.

Quantification and statistical analysis

All data are represented as the mean of biological replicates, with individual sample data shown as

dots. Values are reported as mean ± standard error of the mean (SEM). Statistical analyses were

performed using unpaired Student's t-tests or non-repeated measures analysis of variance (ANOVA),

followed by Tukey's test, using GraphPad Prism (GraphPad Software, Inc., San Diego, CA, USA). P

values for statistical significance are indicated as * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, not detected

(n.d.), or not significant (n.s.).

References

References

Alonso, A., and Pulido, R. (2016). The extended human PTPome: a growing tyrosine

phosphatase family. FEBS J *283*, 1404–1429. https://doi.org/10.1111/febs.13600.

Alonso, A., Sasin, J., Bottini, N., Friedberg, I., Friedberg, I., Osterman, A., Godzik, A.,

Hunter, T., Dixon, J., and Mustelin, T. (2004). Protein tyrosine phosphatases in the

human genome. Cell *117*, 699–711. https://doi.org/10.1016/j.cell.2004.05.018.

Aoki, N., Aoyama, K., Nagata, M., and Matsuda, T. (2001). A growing family of dual

specificity phosphatases with low molecular masses. J Biochem *130*, 133–140.

https://doi.org/10.1093/oxfordjournals.jbchem.a002952.

Arnold, L., Henry, A., Poron, F., Baba-Amer, Y., van Rooijen, N., Plonquet, A.,

Gherardi, R.K., and Chazaud, B. (2007). Inflammatory monocytes recruited after

skeletal muscle injury switch into antiinflammatory macrophages to support

myogenesis. Journal of Experimental Medicine *204*, 1057–1069.

https://doi.org/10.1084/jem.20070075.

Bermudez, O., Pagès, G., and Gimond, C. (2010). The dual-specificity MAP kinase

phosphatases: critical roles in development and cancer. Am J Physiol Cell Physiol *299*,

C189-202. https://doi.org/10.1152/ajpcell.00347.2009.

Boyer, J.G., Prasad, V., Song, T., Lee, D., Fu, X., Grimes, K.M., Sargent, M.A.,

Sadayappan, S., and Molkentin, J.D. (2019). ERK1/2 signaling induces skeletal muscle

slow fiber-type switching and reduces muscular dystrophy disease severity. JCI Insight

4, e127356. https://doi.org/10.1172/jci.insight.127356.

Brett, J.O., Arjona, M., Ikeda, M., Quarta, M., De Morrée, A., Egner, I.M., Perandini,

L.A., Ishak, H.D., Goshayeshi, A., Benjamin, D.I., et al. (2020). Exercise rejuvenates

quiescent skeletal muscle stem cells in old mice through restoration of Cyclin D1. Nat

Metab *2*, 307–317. https://doi.org/10.1038/s42255-020-0190-0.

Chen, H.-H., Luche, R., Wei, B., and Tonks, N.K. (2004). Characterization of two

distinct dual specificity phosphatases encoded in alternative open reading frames of a

single gene located on human chromosome 10q22.2. J Biol Chem *279*, 41404–41413.

https://doi.org/10.1074/jbc.M405286200.

Collins, C.A., Olsen, I., Zammit, P.S., Heslop, L., Petrie, A., Partridge, T.A., and

Morgan, J.E. (2005). Stem cell function, self-renewal, and behavioral heterogeneity of

cells from the adult muscle satellite cell niche. Cell *122*, 289–301.

https://doi.org/10.1016/j.cell.2005.05.010.

Conerly, M.L., Yao, Z., Zhong, J.W., Groudine, M., and Tapscott, S.J. (2016). Distinct

Activities of Myf5 and MyoD Indicate Separate Roles in Skeletal Muscle Lineage

Specification and Differentiation. Dev Cell *36*, 375–385.

https://doi.org/10.1016/j.devcel.2016.01.021.

Crist, C.G., Montarras, D., and Buckingham, M. (2012). Muscle satellite cells are

primed for myogenesis but maintain quiescence with sequestration of Myf5 mRNA

targeted by microRNA-31 in mRNP granules. Cell Stem Cell *11*, 118–126.

https://doi.org/10.1016/j.stem.2012.03.011.

Davis, R.L., Weintraub, H., and Lassar, A.B. (1987). Expression of a single transfected

cDNA converts fibroblasts to myoblasts. Cell *51*, 987–1000.

https://doi.org/10.1016/0092-8674(87)90585-X.

Denu, J.M., and Dixon, J.E. (1995). A catalytic mechanism for the dual-specific

phosphatases. Proc Natl Acad Sci U S A *92*, 5910–5914.

https://doi.org/10.1073/pnas.92.13.5910.

Denu, J.M., and Dixon, J.E. (1998). Protein tyrosine phosphatases: mechanisms of

catalysis and regulation. Curr Opin Chem Biol *2*, 633–641.

https://doi.org/10.1016/s1367-5931(98)80095-1.

Eigler, T., Zarfati, G., Amzallag, E., Sinha, S., Segev, N., Zabary, Y., Zaritsky, A.,

Shakked, A., Umansky, K.-B., Schejter, E.D., et al. (2021). ERK1/2 inhibition promotes

robust myotube growth via CaMKII activation resulting in myoblast-to-myotube fusion.

Developmental Cell *56*, 3349-3363.e6. https://doi.org/10.1016/j.devcel.2021.11.022.

Fero, K., Bergeron, S.A., Horstick, E.J., Codore, H., Li, G.H., Ono, F., Dowling, J.J.,

and Burgess, H.A. (2014). Impaired embryonic motility in dusp27 mutants reveals a

developmental defect in myofibril structure. Dis Model Mech *7*, 289–298.

https://doi.org/10.1242/dmm.013235.

Fujita, R., Kawano, F., Ohira, T., Nakai, N., Shibaguchi, T., Nishimoto, N., and Ohira,

Y. (2014). Anti-interleukin-6 receptor antibody (MR16-1) promotes muscle

regeneration via modulation of gene expressions in infiltrated macrophages. Biochimica

et Biophysica Acta (BBA) - General Subjects *1840*, 3170–3180.

https://doi.org/10.1016/j.bbagen.2014.01.014.

Fujita, R., Zismanov, V., Jacob, J.-M., Jamet, S., Asiev, K., and Crist, C. (2017). Fragile

X mental retardation protein regulates skeletal muscle stem cell activity by regulating

the stability of Myf5 mRNA. Skeletal Muscle *7*, 18. https://doi.org/10.1186/s13395-

017-0136-8.

Fujita, R., Mizuno, S., Sadahiro, T., Hayashi, T., Sugasawa, T., Sugiyama, F., Ono, Y.,

Takahashi, S., and Ieda, M. (2023). Generation of a MyoD knock-in reporter mouse line

to study muscle stem cell dynamics and heterogeneity. iScience *26*, 106592.

https://doi.org/10.1016/j.isci.2023.106592.

Grant, C.E., Bailey, T.L., and Noble, W.S. (2011). FIMO: scanning for occurrences of a

given motif. Bioinformatics *27*, 1017–1018.

https://doi.org/10.1093/bioinformatics/btr064.

Halevy, O., Piestun, Y., Allouh, M.Z., Rosser, B.W.C., Rinkevich, Y., Reshef, R.,

Rozenboim, I., Wleklinski-Lee, M., and Yablonka-Reuveni, Z. (2004). Pattern of Pax7

expression during myogenesis in the posthatch chicken establishes a model for satellite

cell differentiation and renewal. Developmental Dynamics *231*, 489–502.

https://doi.org/10.1002/dvdy.20151.

Hausburg, M.A., Doles, J.D., Clement, S.L., Cadwallader, A.B., Hall, M.N.,

Blackshear, P.J., Lykke-Andersen, J., and Olwin, B.B. (2015). Post-transcriptional

regulation of satellite cell quiescence by TTP-mediated mRNA decay. eLife *4*, e03390.

https://doi.org/10.7554/eLife.03390.

Hu, Y., and Mivechi, N.F. (2006). Association and regulation of heat shock

transcription factor 4b with both extracellular signal-regulated kinase mitogen-activated

protein kinase and dual-specificity tyrosine phosphatase DUSP26. Mol Cell Biol *26*,

3282–3294. https://doi.org/10.1128/MCB.26.8.3282-3294.2006.

Huang, C.-Y., and Tan, T.-H. (2012). DUSPs, to MAP kinases and beyond. Cell Biosci

2, 24. https://doi.org/10.1186/2045-3701-2-24.

Huang, K., Serria, M.S., Nakabayashi, H., Nishi, S., and Sakai, M. (2000). Molecular

cloning and functional characterization of the mouse mafB gene. Gene *242*, 419–426. .

Joe, A.W.B., Yi, L., Natarajan, A., Le Grand, F., So, L., Wang, J., Rudnicki, M.A., and

Rossi, F.M.V. (2010). Muscle injury activates resident fibro/adipogenic progenitors that

facilitate myogenesis. Nat Cell Biol *12*, 153–163. https://doi.org/10.1038/ncb2015.

Kassar-Duchossoy, L., Gayraud-Morel, B., Gomès, D., Rocancourt, D., Buckingham,

M., Shinin, V., and Tajbakhsh, S. (2004). Mrf4 determines skeletal muscle identity in

Myf5:Myod double-mutant mice. Nature *431*, 466–471.

https://doi.org/10.1038/nature02876.

Kitajima, Y., and Ono, Y. (2018). Visualization of PAX7 protein dynamics in muscle

satellite cells in a YFP knock-in-mouse line. Skelet Muscle *8*, 26.

https://doi.org/10.1186/s13395-018-0174-x.

Kuang, S., Kuroda, K., Le Grand, F., and Rudnicki, M.A. (2007). Asymmetric self-

renewal and commitment of satellite stem cells in muscle. Cell *129*, 999–1010.

https://doi.org/10.1016/j.cell.2007.03.044.

Marti, F., Krause, A., Post, N.H., Lyddane, C., Dupont, B., Sadelain, M., and King, P.D.

(2001). Negative-feedback regulation of CD28 costimulation by a novel mitogen-

activated protein kinase phosphatase, MKP6. J Immunol *166*, 197–206.

https://doi.org/10.4049/jimmunol.166.1.197.

Mauro, A. (1961). Satellite cell of skeletal muscle fibers. J Biophys Biochem Cytol *9*,

493–495. https://doi.org/10.1083/jcb.9.2.493.

Megeney, L.A., Kablar, B., Garrett, K., Anderson, J.E., and Rudnicki, M.A. (1996).

MyoD is required for myogenic stem cell function in adult skeletal muscle. Genes Dev.

10, 1173–1183. https://doi.org/10.1101/gad.10.10.1173.

de Morrée, A., van Velthoven, C.T.J., Gan, Q., Salvi, J.S., Klein, J.D.D., Akimenko, I.,

Quarta, M., Biressi, S., and Rando, T.A. (2017). Staufen1 inhibits MyoD translation to

actively maintain muscle stem cell quiescence. Proc Natl Acad Sci USA *114*, E8996–

E9005. https://doi.org/10.1073/pnas.1708725114.

Murphy, M.M., Lawson, J.A., Mathew, S.J., Hutcheson, D.A., and Kardon, G. (2011).

Satellite cells, connective tissue fibroblasts and their interactions are crucial for muscle

regeneration. Development *138*, 3625–3637. https://doi.org/10.1242/dev.064162.

Nabeshima, Y., Hanaoka, K., Hayasaka, M., Esuml, E., Li, S., Nonaka, I., and

Nabeshima, Y. (1993). Myogenin gene disruption results in perinatal lethality because

of severe muscle defect. Nature *364*, 532–535. https://doi.org/10.1038/364532a0.

Oki, S., Ohta, T., Shioi, G., Hatanaka, H., Ogasawara, O., Okuda, Y., Kawaji, H.,

Nakaki, R., Sese, J., and Meno, C. (2018). ChIP-Atlas: a data-mining suite powered by

full integration of public ChIP-seq data. EMBO Rep *19*, e46255.

https://doi.org/10.15252/embr.201846255.

Olguin, H.C., Yang, Z., Tapscott, S.J., and Olwin, B.B. (2007). Reciprocal inhibition

between Pax7 and muscle regulatory factors modulates myogenic cell fate

determination. J Cell Biol *177*, 769–779. https://doi.org/10.1083/jcb.200608122.

Patterson, K.I., Brummer, T., O'Brien, P.M., and Daly, R.J. (2009). Dual-specificity

phosphatases: critical regulators with diverse cellular targets. Biochem J *418*, 475–489.

https://doi.org/10.1042/bj20082234.

Reiterer, V., Pawłowski, K., Desrochers, G., Pause, A., Sharpe, H.J., and Farhan, H.

(2020). The dead phosphatases society: a review of the emerging roles of

pseudophosphatases. FEBS J *287*, 4198–4220. https://doi.org/10.1111/febs.15431.

Relaix, F., Rocancourt, D., Mansouri, A., and Buckingham, M. (2005). A Pax3/Pax7-

dependent population of skeletal muscle progenitor cells. Nature *435*, 948–953.

https://doi.org/10.1038/nature03594.

Rudnicki, M.A., Braun, T., Hinuma, S., and Jaenisch, R. (1992). Inactivation of MyoD

in mice leads to up-regulation of the myogenic HLH gene Myf-5 and results in

apparently normal muscle development. Cell *71*, 383–390.

https://doi.org/10.1016/0092-8674(92)90508-A.

Rudnicki, M.A., Schnegelsberg, P.N.J., Stead, R.H., Braun, T., Arnold, H.-H., and

Jaenisch, R. (1993). MyoD or Myf-5 is required for the formation of skeletal muscle.

Cell *75*, 1351–1359. https://doi.org/10.1016/0092-8674(93)90621-V.

Sabourin, L.A., Girgis-Gabardo, A., Seale, P., Asakura, A., and Rudnicki, M.A. (1999).

Reduced Differentiation Potential of Primary MyoD−/− Myogenic Cells Derived from

Adult Skeletal Muscle. Journal of Cell Biology *144*, 631–643.

https://doi.org/10.1083/jcb.144.4.631.

Seale, P., Sabourin, L.A., Girgis-Gabardo, A., Mansouri, A., Gruss, P., and Rudnicki,

M.A. (2000). Pax7 Is Required for the Specification of Myogenic Satellite Cells. Cell

102, 777–786. https://doi.org/10.1016/S0092-8674(00)00066-0.

Sénéchal, C., Fujita, R., Jamet, S., Maiga, A., Dort, J., Orfi, Z., Dumont, N.A., Bouvier,

M., and Crist, C. (2022). The adhesion G-protein-coupled receptor Gpr116 is essential

to maintain the skeletal muscle stem cell pool. Cell Reports *41*, 111645.

https://doi.org/10.1016/j.celrep.2022.111645.

Tonks, N.K. (2006). Protein tyrosine phosphatases: from genes, to function, to disease.

Nat Rev Mol Cell Biol *7*, 833–846. https://doi.org/10.1038/nrm2039.

Uezumi, A., Fukada, S., Yamamoto, N., Takeda, S., and Tsuchida, K. (2010).

Mesenchymal progenitors distinct from satellite cells contribute to ectopic fat cell

formation in skeletal muscle. Nat Cell Biol *12*, 143–152.

https://doi.org/10.1038/ncb2014.

van Velthoven, C.T.J., and Rando, T.A. (2019). Stem Cell Quiescence: Dynamism,

Restraint, and Cellular Idling. Cell Stem Cell *24*, 213–225.

https://doi.org/10.1016/j.stem.2019.01.001.

Wang, R., Chen, F., Chen, Q., Wan, X., Shi, M., Chen, A.K., Ma, Z., Li, G., Wang, M.,

Ying, Y., et al. (2022). MyoD is a 3D genome structure organizer for muscle cell

identity. Nat Commun *13*, 205. https://doi.org/10.1038/s41467-021-27865-6.

White, J.D., Scaffidi, A., Davies, M., McGeachie, J., Rudnicki, M.A., and Grounds,

M.D. (2000). Myotube formation is delayed but not prevented in MyoD-deficient

skeletal muscle: studies in regenerating whole muscle grafts of adult mice. J Histochem

Cytochem *48*, 1531–1544. https://doi.org/10.1177/002215540004801110.

Wood, W.M., Etemad, S., Yamamoto, M., and Goldhamer, D.J. (2013). MyoD-

expressing progenitors are essential for skeletal myogenesis and satellite cell

development. Developmental Biology *384*, 114–127.

https://doi.org/10.1016/j.ydbio.2013.09.012.

Yablonka-Reuveni, Z., Rudnicki, M.A., Rivera, A.J., Primig, M., Anderson, J.E., and

Natanson, P. (1999). The Transition from Proliferation to Differentiation Is Delayed in

Satellite Cells from Mice Lacking MyoD. Developmental Biology *210*, 440–455.

https://doi.org/10.1006/dbio.1999.9284.

Yamamoto, M., Legendre, N.P., Biswas, A.A., Lawton, A., Yamamoto, S., Tajbakhsh,

S., Kardon, G., and Goldhamer, D.J. (2018). Loss of MyoD and Myf5 in Skeletal

Muscle Stem Cells Results in Altered Myogenic Programming and Failed Regeneration.

Stem Cell Reports *10*, 956–969. https://doi.org/10.1016/j.stemcr.2018.01.027.

Zammit, P.S., Partridge, T.A., and Yablonka-Reuveni, Z. (2006). The Skeletal Muscle

Satellite Cell: The Stem Cell That Came in From the Cold. J Histochem Cytochem. *54*,

1177–1191. https://doi.org/10.1369/jhc.6R6995.2006.

Zismanov, V., Chichkov, V., Colangelo, V., Jamet, S., Wang, S., Syme, A., Koromilas,

A.E., and Crist, C. (2016). Phosphorylation of eIF2α Is a Translational Control

Mechanism Regulating Muscle Stem Cell Quiescence and Self-Renewal. Cell Stem Cell

18, 79–90. https://doi.org/10.1016/j.stem.2015.09.020.

Zou, Z., Ohta, T., Miura, F., and Oki, S. (2022). ChIP-Atlas 2021 update: a data-mining

suite for exploring epigenomic landscapes by fully integrating ChIP-seq, ATAC-

seq and Bisulfite-seq data. Nucleic Acids Res *50*, W175–W182.

https://doi.org/10.1093/nar/gkac199.

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
anti-PAX7	Santa Cruz	Cat# sc-81648; RRID: AB_2159836
anti-MYOD	Santa Cruz	Cat# sc-377460, RRID: AB_2813894
anti-MYOGENIN	abcam	Cat# ab124800, RRID: AB_10971849
Anti-KI67	Invitrogen	Cat# MA5-14520, RRID:AB_10979488
Anti-LAMININ2 ALPHA	abcam	Cat# ab11576, RRID:AB_298180
Anti-MYH1E	DSHB	MF20, RRID:AB_2147781
Alexa Fluor 488 goat anti-mouse IgG1	Thermo Fisher Scientific	Cat# A-21121, RRID: AB_2535764
Alexa Fluor 555 goat anti-mouse IgG1	Thermo Fisher Scientific	Cat# A-21127, RRID:AB_2535769
Alexa Fluor 546 goat anti-rabbit IgG	Thermo Fisher Scientific	Cat# A-11010, RRID: AB_2534077
Alexa Fluor 546 goat anti-mouse IgG2b	Thermo Fisher Scientific	Cat# A-21143, RRID:AB_2535779
Alexa Fluor 633 goat anti-rabbit IgG	Thermo Fisher Scientific	Cat# A-21070, RRID:AB_2535731
Alexa Flour 647 donkey anti-rat IgG	abcam	Cat# ab150155, RRID:AB_2813835
Satellite Cell Isolation Kit	Miltenyi Biotec	Cat# 130-104-268
anti-mouse integrin α-7 MicroBeads	Miltenyi Biotec	Cat# 130-104-261
Bacterial and Virus Strains		
pAd-CMV-EDFP	This study	N/A
pAd-CMV-Dusp13-IRES-EDFP	This study	N/A
pAd-CMV-Dusp13_Mut-IRES-EDFP	This study	N/A
Chemicals, Peptides, and Recombinant Proteins		
DMEM	Thermo Fisher Scientific	Cat# 11965-092
F12	Thermo Fisher Scientific	Cat# 11765-054
Collagenase D	Roche	Cat# 11088866001
D-PBS (-)	FUJIFILM	Cat# 045-29795
OPTI-MEM	Thermo Fisher Scientific	Cat# 31985062
M.O.M. Blocking Reagent	Vector	Cat# MKB-2213-1
Cardiotoxin	LATOXIAN	Cat# 56574-47-1
TRIzol	Thermo Fisher Scientific	Cat# 15596018
Gelatin from porcine skin	MERK	Cat# G1890

Ultroser G	PALL	Cat# 15950-017
Fetal bovine serum	Thermo Fisher Scientific	Cat# 10270-106
Horse serum	Thermo Fisher Scientific	Cat# 26050088
SsoAdvanced^{TM} Universal SYBR Green Supermix	Bio-Rad	Cat# 1725271
4% Paraformaldehyde Fixative	MUTO PURE CHEMICALS CO., LTD.	Cat# 33111
FuGENE 6 Transfection	Promega	Cat# E2692
PrimeSTAR Max DNA	TAKARA BIO Inc.	Cat# R045A
PrimeSTAR Mutagenesis Basal Kit	TAKARA BIO Inc.	Cat#R046A
Normal Goat Serum	Vector	Cat# S-1000
VECTASHIELD Vibrance Antifade Mounting Medium with DAPI	Vector	Cat# H-1800
Glycerol Gelatin	Sigma-Aldrich	Cat# GG1-15ML
Mayer's Hematoxylin Solution	FUJIFILM Wako Pure Chemical Corporation	Cat# 131-09665
Bovine Serum Albumin	Sigma-Aldrich	Cat# A7906
Tragacanth Gum, Powder	FUJIFILM Wako Pure Chemical Corporation	Cat# 200-02245
DPX. Neutral mounting medium	Sigma-Aldrich	Cat# 317616
RNase-free Water, non DEPC	Biodynamics Laboratory Inc.	Cat# DR127
Ethanol (99.5)	FUJIFILM Wako Pure Chemical Corporation	Cat# 057-00456
Ethanol (99)	FUJIFILM Wako Pure Chemical Corporation	Cat# 056-06967
Xylene	FUJIFILM Wako Pure Chemical Corporation	Cat# 242-00087
Acetone	FUJIFILM Wako Pure Chemical Corporation	Cat# 016-00346
Critical Commercial Assays		
Satellite cell isolation Kit, mouse	Miltenyi Biotec	Cat# 130-104-268
ReverTra Ace^{TM} qPCR RT Master Mix with gDNA Remover	TOYOBO	Cat# FSQ-301

Wizard Plus SV Minipreps DNA Purification System	Promega	Cat# A1460
QIAquick PCR Purification Kit	QIAGEN	Cat# 28104
In-Fusion® HD Cloning Kit	TAKARA BIO Inc.	Cat# Z9649N
QuikChange Site-Directed Mutagenesis Kit	Agilent	Cat# 200519
Dual-Luciferase® Reporter Assay System	Promega	Cat# E1910
Deposited Data		
Single RNA-seq data	This paper	Accession number: GSE250228
Experimental Models: Cell Lines		
AAVpro 293T Cell Line	Takara Bio	Cat# 632273
Experimental Models: Organisms/Strains		
Mouse: C57BL/6J	The Jackson Laboratory	RRID:IMSR_JAX:000664
Mouse: Dusp13 KO	This paper	N/A
Mouse: Dusp27 KO	This paper	N/A
Mouse: Pax7-YFP	Kitajima et al.	PMID: 30139390
Mouse: *C57BL/6J-Myod1$^{em1(tdTomato)Utr}$ (MyoD^{KI})*	Fujita et al.	PMID: 37250337
Oligonucleotides		
See Table S1 for genotyping primers	This paper	N/A
See Table S1 for RT-qPCR primers	This paper	N/A
See Table S1 for cloning primers	This paper	N/A
Recombinant DNA		
pGL4-Dusp13	This study	N/A
pGL4-Dusp27	This study	N/A
pAAV-MyoD	This study	N/A
Software and Algorithms		
ImageJ	NIH	https://imagej.net/software/fiji/
Illustrator	Adobe	N/A
Photoshop	Adobe	N/A
Chromium Next GEN Single Cell 3'Reagent Kits v3.1	10× Genomics	Cat# 1000268
Chromium controller	10× Genomics	https://www.10xgenomics.com/
NovaSeq 6000	Ilumina	https://jp.illumina.com/
MEME FIMO v5.5.0	Grant et al.[51]	RRID:SCR_001783
Prism v10	GraphPad	https://www.graphpad.com/
Other		
Thermal Cycler Dice® Real Time System Single	Takara Bio	Cat# TP850

| GloMax® 20/20 Luminometer | Promega | Cat# E5311 |
| NanoDrop™ 2000/2000c | Thermo-Fisher Scientific | Cat# ND-2000 |

Figures

Figure 1. *Dusp13* **is a novel target regulated by MyoD**

(A) Clustering heatmap showing differentially expressed genes in RNA-seq data from Pax7-YFP

(Quiescent MuSCs; QSCs), MyoD-low (MyoD-tdTomato low), and MyoD-high (MyoD-tdTomato

high) (n = 3 per group).

(B) Clustering heatmap of differentially expressed genes in RNA-seq between wild-type (WT**)** growth

medium (GM) (WT_GM), *MyoD* knock-out GM (MKO_GM), WT differentiation medium (WT_DM),

and MKO_DM (n = 3 per group).

(C) Venn diagrams of indicated clusters from **(A, B)** and ChIP atlas top 100 genes potentially regulated

by *MyoD* (Left). Venn diagrams of overlapped 35 genes (left) and 80 genes upregulated by *MyoD*

overexpression in MEFs (Right).

(D) The list of six genes from **(C)**.

(E, F) Gene expression of *MyoD* **(E)** or *Dusp13* **(F)** analyzed by RNA-seq in the Pax7-YFP, MyoD-

low, and MyoD-high MuSCs (n = 3 per group).

(G, H) Gene expression of *MyoD* **(G)** or *Dusp13* **(H)** analyzed by RNA-seq in the WT_GM, WT_DM,

MKO_GM, and MKO_DM (n = 3 per group).

(I) E box-like sequences in the promoter region of *Dusp13* in mammalian species.

(J) Luciferase reporter constructs of *Dusp13* upstream region (-150 to $+50$) containing an E box-like

sequence (Top). Relative *Dusp13*-luciferase reporter activities in HEK293T cells overexpressing

MyoD (Bottom) (n = 3 per group).

All data are represented as the mean ± standard error of the mean (SEM)

P values calculated using edge R **(E, F)** (FDR-corrected), DEseq2 **(G, H)** (FDR-corrected), or Tukey's

test **(J)**; **$P < 0.01$, ***$P < 0.001$, n.s., not significant.

Figure 2. Loss of both *Dusp13* and *Dusp27* impairs muscle regeneration after acute muscle injury.

(A) Hematoxylin and eosin (H&E) staining of tibialis anterior (TA) muscle cross-sections of male *Dusp13* and *Dusp27* double knock-out (DKO) (n = 3) and WT (n = 3) mice at 12–15 weeks of age, 3, 7, and 28 days post-cardiotoxin (CTX) injury. Scale bar: 100 μm.

(B–D) Cross-sectional area (CSA) of damaged TA muscles at indicated time points.

(E, F) Immunofluorescence of PAX7 (**E**) or MYOGENIN (**F**) with LAMININ on sections of TA

muscles from WT and DKO mice, 7 days post-CTX injury. Nuclei are stained with DAPI. Scale bar:

100 μm. The area in white dotted boxes is shown at a higher magnification. Scale bar: 25 μm.

(G, H) Average number of PAX7 (+) or MYOGENIN (+) nuclei in **(E)** and **(F)** are quantified (n =

5–8 independent experiments).

(I) Immunofluorescence of PAX7 and MYOD on cultured MuSCs from WT and DKO mice. Nuclei

are stained with DAPI. Scale bar: 100 μm.

(J) Quantification of the proportion of MuSCs undergoing self-renewal (PAX7 (+), MYOD (-)),

activation (PAX7 (+), MYOD (+)), and differentiation [PAX7 (+), MYOD (-)] in **(I)**. (n = 6

independent experiments)

(K) Immunofluorescence with an antibody against myosin heavy chain (MyHC) on MuSCs

differentiated for six days in DM. Nuclei are stained with DAPI. Scale bar: 100 μm.

(L, M) Myotube width **(L)** and Fusion index (numbers of myonuclei per MyHC (+) myotube) **(M)**

in **(K)** are quantified (WT, n = 50**;** DKO, n = 53).

All data are represented as the mean ± standard error of the mean (SEM)

P values calculated using Student's t-test; *P < 0.05, **P < 0.01, ***P < 0.001, n.s., not significant.

901
Figure 3. Single-cell RNA-sequencing analysis of MyoD-high MuSCs from MyoD-KI mice
reveals *Dusp13* and *Dusp27* expression in the mature stage of MuSCs.
(A) Schematic diagram illustrating the scRNA-seq analysis of MyoD-high MuSCs from MyoD-KI
mice.
(B) Transcriptomic atlas and clustering of the subset of MyoD-high MuSCs from MyoD-KI mice.
(C) Single-cell expression pattern of the target genes.
(D) Pseudotime analysis performed using the Monocle package. The plot depicts the hierarchy of
MyoD-high MuSCs in the pseudotime trajectory.

(E) A heatmap showing the expression of representative genes in the pseudotime analysis. Each

column represents a cell, and each row represents a gene.

(F) Pseudotime-ordered single-cell expression of representative genes.

Figure 4. Overexpression of *Dusp13* induces precocious muscle differentiation independent of phosphatase activity.

(A) Schematic diagram illustrating adenovirus infection into MuSCs from Pax7-YFP mice on day 4 of growth medium (GM) culture.

(B) Number of cultured MuSCs after transduction with adenovirus vectors overexpressing GFP (pAd-EGFP) or Dusp13 (pAd-Dusp13-EGFP).

(C) Immunofluorescence staining of PAX7 and MYOGENIN in cultured MuSCs after transduction with adenovirus vectors overexpressing GFP (pAd-EGFP) or Dusp13 (pAd-Dusp13-EGFP) on GM day 4. Nuclei are stained with DAPI. The yellow arrow indicates PAX7 (+)/MYOGENIN (+) cells. (n = 6 per group). Scale bar: 100 μm (Upper); 25 μm (Bottom).

(D) The proportion of each population in (**C**).

(E) Schematic diagram illustrating adenovirus infection into MuSCs from Pax7-YFP mice on GM day 9.

(F) Immunofluorescence staining of MYOGENIN in cultured MuSCs after transduction with adenovirus vectors overexpressing GFP (pAd-EGFP) or Dusp13 (pAd-Dusp13-EGFP) on GM day 9. Nuclei are stained with DAPI. Scale bar: 100 μm

(G, H) Myotube width **(G)** and Fusion index (numbers of myonuclei per myotube) **(H)** in **(F)** are quantified (WT, n = 66, DKO, n = 104)

All data are represented as the mean ± standard error of the mean (SEM)

P values calculated using Student's t-test; *P < 0.05, **P < 0.01, ***P < 0.001, n.s., not significant.

Supplementary Figure 1 (Hayashi et al.)

Figure S1. Expression changes of candidate genes that are potentially MyoD-regulated.

(A) Gene expression of *Rapsn*, *Jsrp1*, *Dmpk*, *Ank1*, and *Stac3* in RNA-seq data from Pax7-YFP (Quiescent MuSCs; QSCs), MyoD-low (MyoD-tdTomato low), and MyoD-high (MyoD-tdTomato high) (n = 3 per group).

(B) Gene expression of *Rapsn*, *Jsrp1*, *Dmpk*, *Ank1*, and *Stac3* in RNA-seq between wild-type (WT)_growth medium (GM) (WT_GM), *MyoD* knock-out_GM (MKO_GM), WT_differentiation medium (WT_DM), and MKO_DM (n = 3 per group).

All data are represented as the mean ± standard error of the mean (SEM)

P values calculated using edge R **(A)** (FDR-corrected), DEseq2 **(B)** (FDR-corrected); *P < 0.05, ***P < 0.001 among all groups.

Figure S2. Expression pattern of *Dusp13/Dusp27* in a collection of tissues.

(A, B) qPCR analysis of *Dusp13* (A) and *Dusp27* (B) mRNA levels in the tissues (brain, heart, kidney, liver, lung, pancreas, muscle, and spleen) of WT mice at 13–14 weeks (n = 4 per tissue).

(C) qPCR analysis of *Dusp13* mRNA levels in the gastrocnemius muscles of WT and *Dusp13* KO (13 KO) mice at 8–12 weeks (n = 4–6 independent experiments).

(D) qPCR analysis of *Dusp27* mRNA levels in the gastrocnemius muscles of WT and *Dusp27* KO (27 KO) mice at 8–12 weeks (n = 3 per group).

All data are represented as the mean ± standard error of the mean (SEM)

P values calculated using Student's t-test; *P < 0.05, n.d., not detected.

Figure S3. Absence of skeletal muscle phenotype in *Dusp13* knock-out (13 KO) mice during development and regeneration.

(A) Hematoxylin and eosin (H&E) staining of Soleus (SOL) and Plantaris (PLA) muscle cross-sections of male *Dusp13* knock-out (13 KO) and WT mice at 10 weeks. Scale bar: 100 μm.

(B, C) Cross-sectional area (CSA) of SOL (B) or PLA (C) (n = 3 per group).

(D) H&E staining of tibialis anterior (TA) muscle cross-sections of male 13 KO and WT mice at 12 weeks, 7 days post cardiotoxin (CTX) injury. Scale bar: 100 μm.

(E) CSA of damaged TA muscles (n = 3 per group).

(F) Immunofluorescence of PAX7 with LAMININ on sections of TA muscles from WT and *Dusp13* KO mice, 7 days post-CTX injury. Nuclei are stained with DAPI. Scale bar: 100 μm. The area in the white dotted boxes is shown at a higher magnification. Scale bar: 25 μm.

(G) Average number of PAX7 (+) nuclei in **(F)** are quantified (n = 3–4 independent experiments).

(H) Immunofluorescence of MYOGENIN with LAMININ on sections of TA muscle from WT and 13 KO mice, 7 days post-CTX injury. Nuclei are stained with DAPI. Scale bar: 100 μm. The area in the white dotted boxes is shown at a higher magnification. Scale bar: 25 μm.

(I) Average number of MYOGENIN (+) nuclei in **(H)** are quantified (n = 3–4 independent experiments).

All data are represented as the mean ± standard error of the mean (SEM)

P values calculated using Student's t-test; n.s., not significant.

Figure S4. *Dusp27* **KO mice exhibit comparable muscle development and regeneration to WT mice.**

(A) qPCR analysis of *Dusp27* mRNA levels in gastrocnemius muscles of WT and 13 KO mice at 12 weeks (n = 3 per group).

(B) Gene expression of *Dusp27* in RNA-seq data from Pax7-YFP, MYOD-low, and MYOD-high (n = 3 per group).

(C) Ebox-like sequences in the promoter region of *Dusp27* in mammalian species.

(D) Luciferase reporter constructs of *Dusp27* upstream region (−1,000 to +200) containing an Ebox-like sequence (Top). Relative *Dusp27*-luciferase reporter activities in HEK293T cells overexpressing MyoD (Bottom) (n = 3 per group).

(E, F) H&E staining of SOL and PLA muscle cross-sections. Scale bar: 100 μm.

(G, H) CSA of SOL **(G)** or PLA **(H)** muscles of male *Dusp27* KO (27 KO) and WT (n = 3–6 independent experiments) mice at 12 weeks.

(I) H&E staining of TA muscle cross-sections of male 27 KO and WT mice at 12 weeks (n = 5–6 independent experiments), 7 days post-cardiotoxin injury. Scale bar: 100 μm.

(J) CSA of damaged TA muscles.

(K, L) Immunofluorescence of PAX7 **(K)** or MYOGENIN **(L)** with LAMININ on sections of TA muscles from WT and 27 KO mice, 7 days post-CTX injury. Nuclei are stained with DAPI. Scale bar: 100 μm. The area in the white dotted boxes is shown at a higher magnification. Scale bar: 25 μm.

(M, N) Average number of PAX7 (+) or MYOGENIN (+) nuclei in (M) and (N) are quantified (n = 5–8 independent experiments).

All data are represented as the mean ± standard error of the mean (SEM)

P values calculated using Student's t-test, and edge R; *P < 0.05, **P < 0.01, ***P < 0.001, n.s., not significant.

Figure S5. Comparative analysis of intact muscles in *Dusp13:Dusp27* DKO mice and WT mice.

(A) H&E staining of SOL and PLA muscle cross-sections. Scale bar: 100 μm.

(B, C) CSA of SOL **(B)** or PLA **(C)** muscles of male *Dusp13:Dusp27* double knock-out (DKO) and WT mice at 13 weeks (n = 5–7 independent experiments).

(D) Immunoflurescence of PAX7 and LAMININ in intact TA muscles. Scale bar: 100 μm. The area in the white dotted boxes is shown at a higher magnification. Scale bar: 25 μm. **(E)** Average number of PAX7 (+) nuclei in DAPI (+) in **(D)**.

(F) Immunofluorescence of PAX7 on freshly isolated single extensor digitorum longus (EDL) myofibers from DKO and WT mice at 13 weeks (n = 3 per group). Nuclei stained with DAPI. White arrows indicate PAX7 (+) cells on EDL myofibers. Scale bar: 100 μm.

(G) Number of PAX7 (+) cells EDL myofibers in **(F)**.

(H) Immunofluorescence of PAX7 with LAMININ on sections of TA muscles from WT and DKO mice, 28 days post-CTX injury. Nuclei are stained with DAPI. Scale bar: 100 μm. The area in the white dotted boxes is shown at a higher magnification. Scale bar: 25 μm.

(I) Average number of PAX7 (+) nuclei in **(H)** are quantified (n = 6 per group).

(J) Immunofluorescence of MYOGENIN with LAMININ on sections of TA muscles from WT and DKO mice, 28 days post-CTX injury. Nuclei are stained with DAPI. Scale bar: 100 μm. The area in the white dotted boxes is shown at a higher magnification. Scale bar: 25 μm.

(K) Average number of MYOGENIN (+) nuclei in **(J)** are quantified (n = 4–6 independent experiments).

All data are represented as the mean ± standard error of the mean (SEM)

P values calculated using Student's t-test; **P < 0.01, n.s., not significant.

Figure S6. MuSCs derived from *Dusp13*:*Dusp27* DKO mice stay in the proliferation stage.

(A) Immunofluorescence of PAX7 and Ki67 on MuSCs cultured for three days in a growth medium (GM).

(B) Average number of Ki67 (+) nuclei in PAX7 (+) in **(A)** are quantified (n = 6 per group).

(C) Cell numbers of MuSCs derived from WT or DKO at GM day 4 (n = 3 per group).

All data are represented as the mean ± standard error of the mean (SEM)

P values calculated using Student's t-test; **P < 0.01.

Figure S7. Pathway analysis in single-cell RNA-sequencing data of MYOD-high MuSCs from MyoD-KI mice.

(A) Dot plot representing the expression of the top 10 differentially expressed genes for each cluster.

(B–F) A Gene Ontology analysis for cluster 0 **(B)**, cluster 1 **(C)**, cluster 2 **(D)**, cluster 3 **(E)**, and cluster 4 **(F)**.

Figure S8. Expression levels of *Dusp13* and *Dusp27* in a *Dusp13* overexpression experiment.

(A, B) qPCR analysis of *Dusp13* **(A)** and *Dusp27* **(B)** mRNA levels in MuSCs which was overexpressed EGFP or *Dusp13*-EGFP (n = 3 per group).

(C) Immunofluorescence of PAX7 and MYOGENIN on cultured MuSCs after transduction with adenovirus vectors overexpressing Dusp13 (pAd-Dusp13-EGFP) or Dusp13 without phosphatase activity (pAd-DUSP13_Mut-EGFP). Nuclei are stained with DAPI (n = 3 per group).

(D) The proportion of each population in **(C)**.

All data are represented as the mean ± standard error of the mean (SEM)

P values calculated using Student's t-test; *P < 0.05, ***P < 0.001.

Table S1. List of oligonucleotide sequences used as primers

Gene (genotyping)	Orientation	Sequence
Dusp13 (WT)	Fw	5′-GAATGAGGCTATTACCCGGC-3′
	Rv	5′-ACATAACCCACTCCTTCCGG-3′
Dusp13 (KO)	Fw	5′-CCCACATACATATTTGTAGGCACT-3′
	Rv	5′-GCCCCAATGCTAGGTTTACA-3′
Dusp27 (WT)	Fw	5′-AACCGAAAGCATCTTCATGG-3′
	Rv	5′-GTGGGCTAGAGAAGCATTCG-3′
Dusp27 (KO)	Fw	5′-TTGACCTGCCCTACATCTCC-3′
	Rv	5′-CTCACCCCAGGACAAGATGT-3′
Gene (RT-qPCR)	**Orientation**	**Sequence**
b-actin	Fw	5′-CATTGCTGACAGGATGCAGAAGG-3′
	Rv	5′-TGCTGGAAGGTGGACAGTGAGG-3′
Dusp13	Fw	5′-AAGTCTGGCCCAACCTTTTC-3′
	Rv	5′-CACTGCTGCCGTAGAAGTCA-3′
Dusp27	Fw	5′-AACCGAAAGCATCTTCATGG-3′
	Rv	5′-AGCCACGCTTTTCTCAGCTA-3′
Gene (Cloning)	**Orientation**	**Sequence**
Dusp13 for Adenovirus	Fw	5′-CCCAGGATCCATGTCACAGACC-3′
	Rv	5′-GGGTCTCGAGCTAGTGCTGCTGGCTC-3′
Dusp13 for Luciferase	Fw	5′-GCTCGCTAGCCTCGATAACCCTTGCAACAACTCC-3′
	Rv	5′-TCTTGATATCCTCGAACTTCATCCACTCTACTACA-3′
Dusp27 for Luciferase	Fw	5′-GCTCGCTAGCCTCGAATGTCTCCCACGAAGCCT-3′
	Rv	5′-TCTTGATATCCTCGAACACTCAGGGCAGACTGG-3′
MyoD	Fw	5′-GGATTCGCGAGAATTATGGAGCTTCTACGCCG-3′
	Rv	5′-CGACTCTAGAGAATTTCAAAGCACCTGATAAATCG-3′
Gene (Mutation)	**Orientation**	**Sequence**
Dusp13 D97A	Fw	5′-GACCCCTAGGGTCGGGTGCGGGAGGGA-3′
	Rv	5′-CTATAACTTTAGTCCCTCCCGCACCCG-3′
Dusp13 C128S	Fw	5′-CGATTCCACGACCACGTGAGGCACCAC-3′
	Rv	5′-AGCCGAGTGTGGGTGGTGCCTCACGTG-3′