Dining in the daylight: Nocturnal rhinoceros beetles extend feeding periods on host

trees with reduced sap exudation

Asahi Kanda[1], Ryo Shibata[2], Takahiro Ueno[1], Namifu Yamashita[1], Wataru Kojima[1]

[1]Graduate School of Sciences and Technology for Innovation, Yamaguchi University,

1677-1 Yoshida, Yamaguchi 7538511, Japan

[2]Miyashiro-Town Maehara Junior High School, 461 Naka Miyashiro Town,

Minamisaitama, Saitama 3450815, Japan

Correspondence: Asahi Kanda, Graduate School of Sciences and Technology for

Innovation, Yamaguchi University, 1677-1 Yoshida, Yamaguchi City, Yamaguchi

7538511, Japan.

Email: asahikanda083@gmail.com

Telephone: +81-83-933-5767

This MS consists of 16 pages, 3299 words and 4 figures.

Abstract

While the Japanese rhinoceros beetle *Trypoxylus dichotomus* typically feeds on the sap of the oak *Quercus acutissima* and the crape myrtle *Lagerstroemia subcostata* during the night, it exhibits feeding activity both during the day and night when it utilizes the ash tree *Fraxinus griffithii*. However, the mechanisms underlying the variations in temporal activity patterns remain unknown. We compared feeding rates (measured as body mass increments) and sap exudation rates among *F. griffithii*, *Q. acutissima*, and *L. subcostata*. We found that beetles feeding on *L. subcostata* and *Q. acutissima* exhibited significantly higher feeding rates than those feeding on *F. griffithii*. No significant differences in feeding rates were observed between *L. subcostata* and *Q. acutissima*. The sap exudation rate was significantly higher for *Q. acutissima* than for *F. griffithii*. However, there were no significant differences in the sap exudation rates between *F. griffithii* and *L. subcostata* or between *Q. acutissima* and *L. subcostata*. These findings suggest that lower feeding rates on *F. griffithii* prolong the feeding duration, resulting in daytime activity. While the low sap exudation in *L. subcostata* seems inconsistent with high feeding rates on this host, this apparent contradiction could be related to the extended duration of sap exudation.

Keywords; Diel rhythm, Coleoptera, Daily activity pattern, Foraging behavior, Plasticity, Sap-feeding insect, Scarab beetle, Scarabaeidae

Introduction

The daily activity patterns of animals have evolved through natural selection to maximize their fitness. Although activity patterns are typically fixed within species and are classified as diurnal, nocturnal, or crepuscular, some species exhibit adaptive changes in their activity patterns in response to various factors (Levy et al. 2018), including developmental stage (Paulissen 1988), predation pressure (Fenn and Macdonald 1995; Shiojiri et al. 2006), competitor (Siva-Jothy 1987; Levy et al. 2007), season (Stiles et al. 2017), temperature (Nisimura et al. 2005) and physiological state. Hunger is a primary physiological factor influencing animal activity patterns (Lockard 1978; Kramer et al. 2001; Pereira 2010; Hut et al. 2011). Starvation compels individuals to engage in foraging activities to secure the necessary energy, causing them to deviate from their original temporal niches, even if such behavior incurs some costs.

The Japanese rhinoceros beetle *Trypoxylus dichotomus* (Coleoptera, Scarabaeidae, Dynastinae) feeds on tree sap exudates. In Japan, beetles utilize various host tree species across different taxonomic groups, such as Fagaceae, Ulmaceae, and Oleaceae (Siva-Jothy 1987; Hongo 2006; Moriue 2009; Yagihashi et al. 2014; Del sol et al. 2021). Although the rhinoceros beetle is typically nocturnal, its activity patterns vary plastically depending on the host species. Temporal activity patterns were quantitatively investigated in two host species: the oak *Quercus acutissima* and the ash tree *Fraxinus griffithii*. In *Q. acutissima*, *T. dichotomus* feeds on sap from wounds formed by wood-boring insects, such as the carpenter worm *Cossus jezoensis* (Yoshimoto et al. 2005; Ichikawa and Ueda 2010), the bark beetle *Platypus quercivorus* (Akaishi et al. 2006), and the white stripe long-horned beetle *Batocera lineolate* (Takakuwa 2007). *T. dichotomus* flies to sap sites

63 on *Q. acutissima* after sunset and leaves its roosts before sunrise; therefore, beetles are

64 rarely seen during the daytime (Siva-Jothy 1987). In contrast, *T. dichotomus* carves the

65 bark of *F. griffithii*, using its mandible to feed on the sap with head-scooping movements

66 (Hongo 2006; Ichiishi et al. 2019). *T. dichotomus* individuals fly to *F. griffithii* trees at

67 night and remain at their feeding sites for extended periods compared to their feeding

68 behavior on *Q. acutissima*. Consequently, more than half of them continue to feed during

69 the day (Shibata and Kojima 2021). Despite the distinct contrast in activity patterns

70 among host species, the underlying mechanisms remain unclear.

72 Since wounds on *F. griffithii* formed by *T. dichotomus* exuded less sap than those on *Q.*

73 *acutissima* (R. Shibata, personal observation), we predicted that lower feeding rates on *F.*

74 *griffithii* would prolong feeding duration, resulting in daytime activity. To confirm this

75 prediction, we compared the temporal activity patterns, sap exudation rates, and feeding

76 rates of beetles among the host species. In addition to *F. griffithii* and *Q. acutissima*, we

77 dealt with the crape myrtle *Lagerstroemia subcostata*, which is taxonomically distinct

78 from the other two hosts.

80 **Materials and Methods**

81 *Activity Patterns on Lagerstroemia subcostata*

82 Although the activity patterns of *T. dichotomus* have been previously reported for two

83 host species, *Q. acutissima* and *F. griffithii* (see Introduction), no such information is

84 available for *L. subcostata*. Therefore, we assessed the activity patterns of beetles feeding

85 on *L. subcostata*. The study was conducted from August 6 to August 10, 2021, along a

86 path (total length of approximately 300 m) in Anbo, Yakushima-cho, Kagoshima

Prefecture, Japan. Twenty-five trees were selected for observation based on preliminary

surveys, and the number of beetles on these trees was counted hourly. Surveys were not

conducted during rainy weather, and there were 83 counting events over the course of the

study. If the beetles were high up in the trees, they were observed using binoculars (Atrek

II HR 8 × 32 WP; Vixen). For night-time observations, a headlight (MM285-h; GENTOS)

was used; however, direct illumination of the beetles was avoided.

Feeding Rate

The weight gain of beetles per hour was measured as an indicator of feeding rate. The

experiments were conducted in Sugito Town, Saitama Prefecture, in August 2021 for *F.*

griffithii (n = 20 beetles); Yamaguchi City, Yamaguchi Prefecture, in August 2021 for *Q.*

acutissima (n = 26 beetles); and Yakushima-cho in August 2022 for *L. subcostata* (n = 36

beetles). All measurements were conducted at night (20:00 ~ 26:00) when *T. dichotomus*

is most active. Prior to the experiments, field-collected beetles were fasted for 3 days to

stimulate their feeding behavior. Beetles were weighed using a portable digital scale (SF-

700; TUO) at an accuracy of 0.01 g. They were then immediately released into the trees.

They typically initiated sap feeding within approximately 10 min. After an hour of feeding,

beetles were collected and weighed. Individuals who did not continuously feed were

excluded from the analysis. Beetles defecated during feeding, potentially influencing the

accuracy of the feeding rate estimate. However, our experimental setup did not allow us

to measure the weight loss caused by defecation.

Sap Exudation Rate

The sap exudation rate at the beetle feeding sites was measured following the method

described by Yoshimoto (2007). The experiments were conducted in Mitaka City, Tokyo,

in July 2023 for *F. griffithii*; Atsugi City, Kanagawa Prefecture, in July 2023 for *Q.*

acutissima; and Yakushima-cho in July 2023 for *L. subcostata*. As our aim was to test the

relationship between activity time of beetles and sap exudation rates, we measured these

rates during the beetles' activity periods on the focal host species. Specifically,

measurements were conducted at night (20:00 ~ 24:00) for *Q. acutissima* (n = 5 trees and

10 wounds) and *L. subcostata* (n = 2 trees and 8 wounds), and both during the daytime

(10:00 ~ 14:00, n = 5 trees and 10 wounds) and at night (20:00 ~ 24:00, n = 5 trees and

10 wounds) for *F. griffithii*. The beetles were removed from the feeding sites, and solids

and liquids were wiped away with a paper towel. An unused paper towel (approximately

1.5 g), the weight of which was measured beforehand, was pressed against the wounds to

absorb the sap. The paper towel was covered with cling wrap to prevent evaporation.

After 20 min, the towel was collected and weighed again. The increase in weight was

used as an indicator of the sap exudation rate.

Statistical Analyses

Tukey's multiple comparison method (HSD) was used to determine variation in feeding

rates among the three host species. As our preliminary analyses showed that pronotum

width (an indicator of body size) and sex did not significantly affect the feeding rate, these

variables were not included in the model. Comparisons of sap exudation rates among host

species were conducted using TukeyHSD. The R software package (version 4.0.4; R

Foundation for Statistical Computing, Vienna, Austria) was used for statistical analysis.

Results

The number of *T. dichotomus* on *L. subcostata* started increasing around 8 pm, reaching

a maximum at midnight, and most individuals left the tree by 5 am. The number of beetles

during the daytime was approximately one-fifth of the nighttime peak. Most of the

individuals on the trees were engaged in activities such as feeding and mating. These

results indicated that the beetles feeding on *L. subcostata* were nocturnal.

The weight gain of the beetles feeding on *L. subcostata* and *Q. acutissima* was

significantly greater than those feeding on *F. griffithii*. There was no significant difference

in the weight gain between those feeding on *Q. acutissima* and *L. subcostata*. The body

weights of some individuals decreased after 1-hour feeding, probably because of

defecation.

Since the sap exudation rate of *F. griffithii* did not vary between day and night (Welch's

t-test: t = -1.2358, df = 17.89, p = 0.2325), the data were combined for comparison with

other tree species. The sap exudation rate of *Q. acutissima* was significantly higher than

that of *F. griffithii*. There were no significant differences in the sap exudation rates

between *F. griffithii* and *L. subcostata* or *Q. acutissima* and *L. subcostata*.

Discussion

This study revealed that the feeding rate of *T. dichotomus* on *F. griffithii* was lower than

that on other host species. In addition, the sap exudation rate of *F. griffithii* was lower

than that of *Q. acutissima*. Furthermore, we found that the beetle activity patterns on *L.*

subcostata were similar to those on *Q. acutissima*. Based on these findings, when beetles

utilize host trees with low sap exudation, their activity periods are likely to be extended.

Feeding behavior on *Q. acutissima* and *L. subcostata* is probably completed during the

night, whereas beetles feeding on *F. griffithii* face the challenge of inadequate sap intake

at night, compelling them to continue feeding until daytime.

Sap quantity in *Q. acutissima* was greater than that of the other tree species. This may be

due to the sap exudation mechanism of *Q. acutissima* was different from that of the other

tree species. In *Q. acutissima*, the interior of the trunk was damaged by wood-boring

insects, whereas in the other two tree species, only the surface of the bark was carved by

T. dichotomus. There was no significant difference in sap exudation rates between *F.*

griffithii and *L. subcostata*. Nevertheless, the feeding rates were significantly higher in *L.*

subcostata. These results are seemingly contradictory but could be explained by the

difference in the duration of sap exudation. *L. subcostata* may continue to produce sap

longer with a single excavation than *F. griffithii*.

Diurnal activities are likely to impose costs on *T. dichotomus*. The large black body of *T.*

dichotomus makes it more conspicuous during the day, increasing its vulnerability to

visual predators such as crows (Kojima et al. 2014). Furthermore, the daytime ambient

temperature in central Japan reaches 35–40 °C, which may cause physiological stress to

T. dichotomus, as has been reported in other insect species (Chen et al. 2018). Despite

these potential costs, *T. dichotomus* might gain net benefits from the extended activity

period of *F. griffithii*.

Although this study suggests that feeding rate affects the activity patterns of rhinoceros

beetles, multiple unexplored factors could also be attributable to host-dependent activity

patterns. For example, insufficient nutritional content (e.g., sugar and protein) in the sap

may lead to prolonged feeding times. Phloem sap consists mainly of sugars, with trace

amounts of amino acids, organic acids, and proteins (Jensen et al. 2016). For example,

the sap from *Q. acutissima* contains fructose, glucose, ethanol, acetic acid, glycerol, lactic

acid, and 2,3-butanediol (Yoshimoto 2008). However, due to the absence of sap

composition data for the other two host species, we cannot examine or test this hypothesis.

Interspecific competition for sap sites is also one of the potential factors that might affect

the activity patterns of rhinoceros beetles. Kojima (2023) reported that the giant hornet

Vespa mandarinia visited sap sites on *Q. acutissima* at approximately 5:00 am and

physically excluded *T. dichotomus*. After the hornets were experimentally removed, the

beetles continued to occupy the sap sites until noon. Thus, the activity duration of

rhinoceros beetles may be extended into the daytime even on *Q. acutissima,* if they cannot

obtain adequate sap due to factors such as lower conditions of sap sites or intense

intraspecific competition. Additionally, male beetles are active at different times of the

day depending on their body size (Siva-Jothy 1987). As the body size of individuals at

the sap sites was not measured in this or previous studies (Shibata and Kojima 2021;

Kojima 2023), further investigation of individual-level activity patterns is required.

Host-dependent variation in activity patterns has been observed in the sap-feeding stag

beetle *Prosopocoilis dissimilis okinawanus* (Hongo 2005). While some individuals of this

stag beetle exhibit daytime activity on *Citrus trichotomum* var. esculentum and *Persea*

thunbergii, they become exclusively nocturnal with an adequate food in laboratory.

Although the study did not examine the sap feeding rates and the sap exudation rates,

similar mechanisms to those observed in *T. dichotomus* might underlie the activity

patterns of the stag beetle. This example in the stag beetle and our study in *T. dichotomus*

also highlight the possibility that host-dependent variation in activity patterns may be

widespread across sap-feeding insects.

Acknowledgements

We thank Kazuya Shibata and Mariko Shibata for assistance with field observation, and

Yoshiaki Kobayashi for the permission to perform fieldwork and provision of information.

Funding

This study was partly supported by grants from Yamaguchi University.

Declarations

Conflict of interest

The authors declare that they have no conflict of interest.

Ethical approval

No approval of research ethics committees was required for this research.

References

Akaishi D, Kamata N, Nakamura K (2006) Initial stage of an infestation of *Platypus*
quercivorus (Coleoptera: Platypodidae) in a secondary forest dominated by *Quercus*
serrata and *Quercus variabilis* (in Japanese). J Jpn For Soc 88:274–278

Chen H, Zheng X, Luo M, Guo J, Solangi G, Wan F, Zhou Z (2018) Effect of short-term
high-temperature exposure on the life history parameters of *Ophraella communa*. Sci Rep
8:13969

Del Sol JF, Hongo Y, Boisseau RP, Berman GH, Allen CE, Emlen DJ (2021) Population
differences in the strength of sexual selection match relative weapon size in the Japanese
rhinoceros beetle, *Trypoxylus dichotomus* (Coleoptera: Scarabaeidae). Evolution 75:394–
413

Fenn MG, Macdonald DW (1995) Use of middens by red foxes: risk reverses rhythms of
rats. J Mammal 76:130–136

Hut RA, Pilorz V, Boerema AS, Strijkstra AM, Daan S (2011) Working for food shifts
nocturnal mouse activity into the day. PloS One 6:e17527

Hongo Y (2005) Diurnal activity of the Japanese stag beetle *Prosopocoilus dissimilis*
okinawanus NOMURA (Coleoptera, Lucanidae). Elytra 33:245–257

Hongo Y (2006) Bark-carving behavior of the Japanese horned beetle *Trypoxylus*
dichotomus septentrionalis (Coleoptera: Scarabaeidae). J Ethol 24:201–204

Ichiishi W, Shimada S, Motobayashi T, Abe H (2019) Completely engaged three-
dimensional mandibular gear-like structures in the adult horned beetles: reconsideration
of bark-carving behaviors (Coleoptera, Scarabaeidae, Dynastinae). ZooKeys 813:89–110

Ichikawa T, Ueda K (2010) Predation on exuded sap-dependant arthropods by the larvae
of oriental carpenter moth, *Cossus jezoensis* (Matsumura)(Lepidoptera, Cossidae):
preliminary observations (in Japanese). Tech Bull Fac Agr Kagawa Univ 62:39–58

Jensen KH, Berg-Sørensen K, Bruus H, Holbrook NM, Liesche J, Schulz A, Zwieniecki
MA, Bohr T (2016) Sap flow and sugar transport in plants. Rev Mod Phys 88:035007

Kojima W, Sugiura S, Makihara H, Ishikawa Y, Takanashi T (2014) Rhinoceros beetles
suffer male-biased predation by mammalian and avian predators. Zool Sci 31:109–115

Kojima W (2023) Temporal niche shifts driven by interference competition: Giant hornets
exclude rhinoceros beetles at sap sites at dawn. Ecology 104:e3914

Kramer K, Birney E (2001) Effect of light intensity on activity patterns of Patagonian
leaf-eared mice, *Phyllotis xanthopygus*. J Mammal 82:535–544

Levy O, Dayan T, Kronfeld‑Schor N (2007) The relationship between the golden spiny
mouse circadian system and its diurnal activity: an experimental field enclosures and
laboratory study. Chronobiol Int 24:599–613.

Levy O, Dayan T, Porter WP, Kronfeld-Schor N (2019) Time and ecological resilience:
can diurnal animals compensate for climate change by shifting to nocturnal activity? Ecol
Monogr 89:e01334

Lockard RB (1978) Seasonal change in the activity pattern of *Dipodomys spectabilis*. J
Mammal 59:563–568

Moriue N (2009) The handbook of the insects which sap attracts (in Japanese). Bunichi-
Sogo Syuppan, Tokyo

Nisimura T, Numata H, Yoshioka E (2005) Effect of temperature on circadian rhythm
controlling the crepuscular activity of the burying beetle *Nicrophorus quadripunctatus*
Kraatz (Coleoptera: Silphidae). Entomol Sci 8:331–338

Paulissen MA (1988) Ontogenetic and seasonal comparisons of daily activity patterns of
the six-lined racerunner, *Cnemidophorus sexlineatus* (Sauria: Teiidae). Am Midl Nat
120:355–361

Pereira JA (2010) Activity pattern of Geoffroy's cats (*Leopardus geoffroyi*) during a
period of food shortage. J Arid Environ 74:1106–1109

Schlaepfer MA, Sherman PW, Blossey B, Runge MC (2005) Introduced species as
evolutionary traps. Ecol Lett 8:241–246

Stiles RM, Halliday TR, Engbrecht NJ, Swan JW, Lannoo MJ (2017) Wildlife cameras
reveal high resolution activity patterns in threatened crawfish frogs (*Lithobates*
areolatus). Herpetol Conserv Biol 12:160–170

Shibata R, Kojima W (2021) An introduced host plant alters circadian activity patterns of
a rhinoceros beetle. Ecology 102:e03366

Shiojiri K, Ozawa R, Takabayashi J (2006) Plant volatiles, rather than light, determine
the nocturnal behavior of a caterpillar. PLoS Biol 4:e164

Siva-Jothy MT (1987) Mate securing tactics and the cost of fighting in the Japanese
Horned Beetle. J Ethol 5:165–172

Takakuwa M (2007) A Longihorn Beetle *Batocera lineolata* Chevrolat and the Sap Insects
in the Secondary Forests, Why Have Both Remarkably Been Declined ?. Bull Kanagawa
prefect Mus Nat Sci 36:75–90

Yagihashi T, Saitoh T, Maehara N, Noguchi M (2014) Bark-carving and sap-obtaining

behavior of *Trypoxylus dichotomus septentrionalis* on *Fraxinus japonica* (in Japanese).

Tohoku Soc Forest Sci 19:63–65

Yoshimoto J, Kakutani T, Nishida T (2005) Influence of resource abundance on the

structure of the insect community attracted to fermented tree sap. Ecol Res 20: 405–414

Yoshimoto J, Nishida T (2007) Boring effect of carpenterworms (Lepidoptera: Cossidae)

on sap exudation of the oak, *Quercus acutissima*. Appl Entomol Zool 42:403–410

Yoshimoto J (2008). Ecological studies on the insect community attracted to sap (in

Japanese). PhD thesis,Kyoto University.

Figures

Figure 1. *Trypoxylus dichotomus* feeding on the sap of *Quercus acutissima* (a), *Fraxinus*

griffithii (b), and *Lagerstroemia subcostata* (c). In (b) and (c), *T. dichotomus* is engaging

in bark carving, with the scars indicated by black arrows. In (c), the smaller beetle

above the female *T. dichotomus* is a female stag beetle *Prosopocoilus inclinatus*.

Figure 2. Experimental setup for measuring sap exuding rate on *Quercus acutissima*.

The sap is absorbed by the paper towel at the center, and covered with cling wrap. Tape

was applied to secure the cling wrap.

Figure 3. Daily activity pattern of *Trypoxylus dichotomus* on *Lagerstroemia subcostata*.

The number of beetles is the total number of individuals in all surveyed trees. Different

symbols represent the data of three days. The gray area indicates the night.

Figure 4. Feeding rates of *Trypoxylus dichotomus* (a) and sap exudation rates (b) on

three host species. Feeding rates were assessed using hourly weight gain of the beetle.

Sap exudation rates were assessed by measuring the weight gain during the 20-minute

period that the paper towel was pressed against sap sites. Three asterisks indicate P <

0.001, two asterisks indicate P < 0.01, and 'N.S.' indicates no significant difference.