Aberrant connective tissue sheath contraction drives premature hair follicle regression by inducing progenitor cell depletion in androgenetic alopecia

Zhili Deng[1, 2, 3, #], Li Yang[1, 2, 3, #], Shixin Duan[1, 2, 3], Mengting Chen[1, 2, 3], Yujin Zhang[4], Fangfen Liu[1, 2, 3], Yan Tang[1, 2, 3], Yunying Wang[1, 2, 3, 5], Jiayun Li[1, 2, 3], San Xu[1, 2, 3], Zheng Wu[1, 2, 3], Ben Wang[1, 2, 3], Zhixiang Zhao[1, 2, 3], Wei Shi[1, 2, 3], Hongfu Xie[1, 2, 3], Guo Li[1, 2, 3, *], Ji Li[1 ,2, 3, *]

[1]Department of Dermatology, Xiangya Hospital, Central South University, Changsha, Hunan, China.

[2]Hunan Key Laboratory of Aging Biology, Xiangya Hospital, Central South University, Changsha, Hunan, China.

[3]National Clinical Research Center for Geriatric Disorders, Xiangya Hospital, Central South University, Changsha, Hunan, China.

[4]Department of Dermatology, The Second Hospital of Hunan University of Chinese Medicine, Changsha, Hunan, China.

[5]Department of Dermatology, Second Affiliated Hospital of Xinjiang Medical University, Urumqi, China.

[#]These authors contributed equally: Zhili Deng, Li Yang.

[*]Corresponding authors: Ji Li, E-mail: liji_xy@csu.edu.cn; Guo Li, E-mail: liguo237@csu.edu.cn.

Abstract

Androgenetic alopecia (AGA) is characterized by progressive miniaturization of hair, forming a distinctive patterned baldness in the scalp; yet, the mechanisms for hair miniaturization in this disease remain largely unknown. Here by single-cell transcriptome analysis, we describe a concise single-cell atlas, and identify the early changes in cell subpopulations, hair follicle (HF) stem cell fate determination and cell-cell communications in AGA anagen HF units. Thereinto, apoptotic loss of HF progenitor cells is significantly increased, correlated with HF miniaturization in AGA. Mechanistically, enhanced contraction of connective tissue sheath (CTS) activates the mechanosensitive channel PIEZO1, which triggers ectopic apoptosis of progenitor cells in human anagen HFs. Continuous CTS contraction during AGA causes long-term loss of progenitor cells via inducing persistent ectopic apoptosis through PIEZO1, eventually leading to premature hair regression. Most importantly, we show that targeting CTS contraction by ML-7, a selective myosin light chain kinase (MLCK) inhibitor, can obviously improve the growth of HFs from balding scalps of AGA patients. Our study reveals the cellular hierarchies and identifies CTS with increased muscle contraction activity as the key determinant of premature hair regression in AGA, highlighting CTS surrounding human HF as therapeutic targets for treating this disorder.

Introduction

Androgenetic alopecia (AGA), also known as male pattern baldness, is the most common type of baldness, affecting both men and women. Testosterone and androgen receptor (AR) signaling pathway are considered to be necessary for AGA development(Ceruti et al., 2018; Inui and Itami, 2013). Genetic susceptibility loci mapping to AR gene and its neighboring ectodysplasin A2 receptor gene (EDA2R) have also been found in AGA(Liu et al., 2016). Current recognized therapies for AGA include finasteride, minoxidil, platelet-rich plasma (PRP) and low-level laser light therapy (LLLT)(Adil and Godwin, 2017; Katzer

et al., 2019). Among these, finasteride suppresses the activity of 5-a reductase 2 (SRD5A2), which transforms testosterone to dihydrotestosterone (DHT), a more potent AR agonist(Ceruti et al., 2018). Although the potential targets of other therapies remain elusive, additional factors are suggested to be involved in the development of AGA besides AR-related pathways; however, specific instances have not been conclusively identified.

AGA is featured by the progressive and patterned conversion of large terminal hair follicles (HFs) into small vellus HFs (termed as HF miniaturization), which eventually results in balding(Deng et al., 2022). All HFs undergo periodic cycles of growth (anagen), regression (catagen), and rest (telogen) stages throughout the lifetime(Deng et al., 2015). Normally, most human hair follicles will remain in anagen for 2-6 years to produce large terminal hair. But in AGA, the growth phase decreases to only days or weeks, followed by premature entry into hair regression; this leads to an increase in the percentage of resting HFs, generating microscopic hairs in the balding scalp as the HF cycles(Garza et al., 2012; Gu et al., 2022). In HF cycle, the growth of HFs relies on the proliferation and differentiation of HF stem/progenitor cells, whose activities are mainly regulated by the components of HF niche (including dermal papilla (DP), dermal sheath (DS), adipocytes and blood vessels *etc.*)(Deng et al., 2022; Festa et al., 2011; Heitman et al., 2020; Morgan, 2014; Plikus and Chuong, 2014). In balding scalp of AGA, HF stem cells (HFSCs), also known as bulge cells, are basically retained, whereas the number of progenitor cells, a cluster of more actively proliferating cells, is decreased(Garza et al., 2011). This suggests that HFs in AGA balding scalp lack fuels for growth, but the details and underlying mechanism are unclear.

To date, the pathogenetic study of AGA has been scant and limited to histomorphology, global gene analysis and immunohistochemistry due to the fact that most laboratory models are not fully representative of AGA(Lolli et al., 2017; Ntshingila et al., 2021). Single-cell RNA-sequencing (scRNA-seq) has emerged as a powerful tool for dissecting the transcriptome dynamics, cell

heterogeneity, cell lineage tracing and cell-cell communications in complex

tissues at uniquely high resolution(Deutsch et al., 2020; Jin et al., 2021; Mereu

et al., 2020; Papalexi and Satija, 2018; Potter, 2018). However, to our

knowledge, a study regarding scRNA-seq application for exploring the

pathogenesis of AGA has never been described.

In this study, we report a comprehensive scRNA-seq analysis on human

anagen HF units from 3 conditions: balding frontal and non-balding occipital

scalps of patients with AGA, and normal frontal scalps of healthy individuals.

With these data, we generated a concise single-cell atlas and elucidated the

early cellular and molecular changes in AGA anagen HF units. Importantly, via

combining human HF organ culture system, we revealed an unexpected role of

connective tissue sheath (CTS), the analogous structure of murine DS in

human, in regulating HF progenitor cell survival in the development of AGA. In

AGA, CTS contraction is significantly enhanced, resulting in mechanical

compression of HF progenitor cells; increased mechanical force activates the

mechanosensitive calcium channel PIEZO1, thus induces apoptotic loss of

progenitor cells, which eventually leads to premature hair regression. Finally,

we demonstrated that targeting CTS contraction by ML-7 can promote the

growth of HFs from the balding scalps of AGA patients, even better than

minoxidil. These findings illustrate the cellular hierarchies in AGA and

emphasize relaxing hair follicle by targeting CTS as a promising therapy for its

treatment.

Results

Single-cell RNA-seq reveals cell heterogeneity of androgenetic alopecia

and healthy anagen HF units

To dissect the cellular heterogeneity and explore the early changes in AGA

development, we obtained anagen HFs from frontal balding (B) scalps and

occipital non-balding (NB) scalps of AGA patients, and frontal normal scalps of

healthy individuals (HS), to perform scRNA-seq (Fig. 1A). Among these,

anagen HFs of balding scalps were at the early stage of hair miniaturization,

isolated from the edge of frontal hairline of AGA patients. First, to identify the

hair cycle phase of human HFs, we established an identification system by

integrating multiple methods reported previously(Langan et al., 2015; Oh et al.,

2016) (Fig. S1). Before conducting scRNA-seq, the HFs were confirmed in

anagen phase. After sequencing and stringent cell filtration, 76368 cells were

retained for subsequent analyses. Following a standard scRNA-seq data

analysis procedure, totally 28 clusters (C0-C27) were retained, which were

sufficiently separated from each other (Fig. S2A), and these clusters could be

reproduced using cells from each of the 12 samples, suggesting that they are

robustly present across different samples and conditions (including HS, NB and

B) (Fig. S2B, C).

To dissect the cell types, we first defined the expression pattern of SOX9 (Fig.

S2D), a reported marker for hair follicle cells(Zou et al., 2021). By

immunohistochemistry, we confirmed that SOX9 is expressed in most cells of

hair follicle and almost none in other cells, distinguishing hair follicle cells from

other epidermal and dermal cells (Fig. S2E). Based on the expression of SOX9,

the above clusters were further identified as 20 different cell types, according

to the expression of known lineage markers, including Bulge (KRT15$^+$,

CXCL14$^+$), DP (VCAN$^+$, CFD$^+$), DS (ACTA2$^+$, CCN2$^+$), outer root sheat basal

(ORS, KRT14$^+$, PTHLH$^+$), ORS suprabasal (ORS SB, KRT17$^+$), inner root sheat

(KRT6A$^+$), medulla (KRT85$^+$, KRT31$^+$), proliferating progenitors (PR, MKI67$^+$),

endothelial cells (Endo, PECAM1$^+$, VWF$^+$), immune cells (CD3D$^+$), Langerhans

(CD207$^+$, CD1A$^+$), Melanocytes (DCT$^+$, MLANA$^+$), Vascular smooth muscle

cells/pericytes (vSMC, ACTA2$^+$, RGS5$^+$, RERGL$^+$), Sebaceous gland cells

(DCD$^+$, SCD$^+$), sweet gland cells (SwG, KRT8$^+$, KRT18$^+$), interfollicular

epidermis spinous (IFE spinous, KRT1$^+$, KRT10$^+$), IFE granular (FLG$^+$, KRT2$^+$);

cells highly expressing KRT14 but not SOX9 were defined as IFE basal; Inner

II (KRT79$^+$) is the outer layer of Infundibulum and Isthmus which highly

expressed KRT14 and KRT6A in the upper HF region; in contrast, Outer II has

a relatively low expression of KRT6A(Cui and Schlessinger, 2015; He et al.,

2020; Hughes et al., 2020; Joost et al., 2020; Joost et al., 2018; Joost et al.,

2016; Liu et al., 2022; Rahmani et al., 2014; Takahashi et al., 2020; Wu et al.,

2022; Yang et al., 2017) (Fig. 1B-D; Supplementary Table 1; Fig. S2F-K).

To investigate the trajectories among all 20 cell types, monocle3(Cao et al.,

2019) was employed to perform the pseudotime analysis with the root node

setting at start point of bulge on the SOX9$^+$ HF branch and start point of IFE

basal on the SOX9$^-$ IFE branch separately. Our data demonstrate that bulge

cells first differentiate into ORS and IRS cells, which then further branch into

SOX9$^+$ PR, also known as matrix cells, and finally contribute to medulla (HF)

formation. For the SOX9$^-$ IFE branch, IFE basal cells can directly differentiate

into SOX9$^-$ PR and IFE spinous cells which finally develop into IFE granular

cells (Fig. 1E). These patterns are consistent with the knowledge from previous

studies(Wu et al., 2022; Yang et al., 2017).

Consistent with the scRNA-seq data, immunostaining analysis showed that

KRT15 was highly expressed in bulge HFSCs and moderately expressed in IFE

basal cells (Fig.1F), similar to previous observations(Garza et al., 2011; Wu et

al., 2022). As an IRS marker, KRT6A was exactly expressed mainly in the IRS

and some of the inner layer cells of the infundibulum and isthmus, which are

the upward continuation of IRS (Fig.1G and Fig. S2L). Moreover, we verified

IFE spinous cells with KRT10 and DS cells with α-SMA (Fig. 1H, I). We

determined the relative proportion of each cell type in all samples, showing

increased DS and sebaceous gland cells in B anagen HF units compared with

NB or HS (Fig. S2M). We next explored the number of differentially expressed

genes (DEGs) between B and NB/HS. The results showed that all cell types

had differences in various degree (Fig. S2N, Supplementary Table 2),

suggesting that anagen HFs undergo significant changes at the early stage of

hair miniaturization in AGA.

To further explore the potential heterogeneity, bulge cells were subclustered

into three distinct subclusters (Fig. 2A), in which the percentage of subcluster

was reduced in the balding bulge of AGA patients (Fig. 2B). Wilcoxon rank-sum test was used to characterize the marker genes specifically enriched in each subpopulation versus other subpopulations. Top marker genes with avg_log2FC>0.25, pct (minimum fraction in either of the two populations)>0.25 and P value<0.05 of each subcluster were showed in Fig. S3A and Supplementary Table 3. Among these, KRT15 was commonly highly expressed (Fig. 2C); CD200 was visibly enriched in subcluster 0 (Fig. 2D), while CD34 was mainly enriched in subcluster 1 (Fig. 2E). Consistently, CD34$^+$ bulge cells were indeed decreased in the balding bulge (Fig. 2F). By immunostaining, we defined that CD200$^+$ cells are located in the upper bulge with high stemness (known as HFSCs), while CD34 is mainly expressed in the lower bulge cells referred to as HF progenitor cells, which are differentiated from the upper bulge HFSCs and give rise to matrix cells and produce the new hair shaft (Fig. 2G, H). Via RNA Velocyto analysis, we further confirmed that CD200$^+$ bulge cells are the HFSCs, which have the ability to differentiate into CD34$^+$ progenitor cells (Fig. 2I).

PR cells are the proliferating progenitors which highly expressed MKI67. Further subclustering analysis separated PR cells into 2 subclusters (Fig. S3B). Cell composition analysis showed the subcluster 1decreased in B anagen HFs of AGA (Fig. S3C). Among these two subclusters, one was enriched for epithelial markers (KRT1 and KRT10), representing the IFE proliferating progenitor (ePR) cells; another one (subcluster 1) was specifically enriched for HF cell marker SOX9, suggesting that it represented the HF proliferating progenitors also named as matrix cells (Fig. S3D, E, and Supplementary Table 4).

Subclustering of DP cells, the signaling center for hair growth(Morgan, 2014), revealed 5 subclusters (Fig. 2J, Fig. S3F and Supplementary Table 5). The percentage and distribution of subclusters were comparable across HS, NB and B, suggesting that DP cell identity may not alter in AGA anagen HFs (Fig. 2K and Supplementary Table 6). Gene ontology (GO) analysis showed that

angiogenesis-related pathway was downregulated (Fig. 2L), consistent with our

previous findings from bulk RNA sequencing(Deng et al., 2022).

DS cells were divided into 4 subclusters (Fig. 2M, Fig. S3G and

Supplementary Table 7), which showed no significant changes in the balding

anagen HFs of AGA patients (Fig. 2N and Supplementary Table 8). Pathway

enrichment analysis of differential genes revealed muscle contraction was

obviously upregulated in balding anagen HFs compared to non-balding or

healthy groups (Fig. 2O).

In human hair follicle, the DS is tightly surrounded by a network of blood

vessels, composed of vSMCs and endothelial cells(Martino et al., 2021).

Among them, vSMCs were separated into 4 subclusters, and the percentage

and distribution of subclusters were comparable across three groups (Fig. 2P,

O; Fig. S3H; Supplementary Table 9 and Supplementary Table 10). Pathway

enrichment analysis showed that smooth muscle contraction and multiple

pathways involved in inflammatory regulation (such as IL-17 signaling pathway)

were upregulated in vSMCs of balding anagen HFs (Fig. 2R). Endothelial cells

were divided into 4 subclusters, which exhibited no obvious alterations (Fig.

S3I-K; Supplementary Table 11 and Supplementary Table 12), and IL-17

signaling pathway was also upregulated in endothelial cells of balding anagen

HFs (Fig. S3L).

Immune cells expressing CD3D were further separated into 6 subclusters,

including Th17 cells (IL17A$^+$, IL17F$^+$ and IL22$^+$), naive CD4$^+$ T cells (IL7R$^+$),

CD8$^+$ T cells (GZMK$^+$ and CD8A$^+$), natural killer T (NKT) cells (ZNF683$^+$) and

two unknown subclusters (Fig. S3M-O; Supplementary Table 13). Cell

composition analysis showed significant increase of Th17 cells in balding HF

units (Fig. S3P, Q), which provides evidence for inflammatory infiltration in AGA.

Identification of early changes in HFSC fate determination and cell-cell

communications in AGA

Epidermal and sebaceous differentiation of HFSCs has been reported to be

associated with HF miniaturization during ageing-induced and obesity-induced

hair loss in mice(Matsumura et al., 2016; Morinaga et al., 2021), but it was unclear whether there existed HFSC fate determination alterations in AGA development. To address this question, we conducted cell fate analysis among potential cell types: bulge, ORS, sebaceous gland and IFE basal cells. As expected, more HFSCs showed a pronounced tendency to differentiate into sebaceous gland and IFE basal cells in balding HFs compared with non-balding and healthy HFs, and obviously decreased HFSCs differentiated into ORS cells (Fig. S4A, D). To explore the underlying mechanisms, branched expression analysis modeling (BAEM) was performed, and identified three common cell fate determination genes, COL17A1, PPP1R1C and SGK1 (Fig. S4B, C, E, F).

To further characterize the cell-cell communications between different cell types in HF units. Cellchat(Jin et al., 2021) was employed to compare the cell-cell communications between balding and non-balding/healthy HFs. Although the interaction strength showed that most cell types had no obvious changes in incoming and outgoing interaction strength, Th17 cells and langerhans exhibited increased incoming interaction strength in balding and non-balding HFs compared to healthy HFs, while ORS cells displayed decreased outgoing interaction strength in balding anagen HFs in AGA (Fig. S4G).

Apoptotic loss of HF progenitor cells is correlated with HF miniaturization in AGA

AGA is characterized by the progressive conversion of the terminal (t) scalp HFs into intermediate/miniaturized (i/m) HFs, which we called the process of HF miniaturization and may be attributed to the deficiency of progenitor cells based on our scRNA-seq data as mentioned above and the early study(Garza et al., 2011). To verify changes in HFSC and progenitor cell numbers, we performed immunostaining of CD200 and CD34 on anagen HFs from frontal balding (B) and occipital non-balding scalps (NB) of AGA patients, and frontal normal scalps of healthy individuals (HS). Surprisingly, HFs retain CD200$^+$ HFSCs with the same percentage in AGA, despite a decrease in cell numbers in B scalps (Fig. S5A-C). In contrast, CD34$^+$ progenitor cells were gradually

diminished during the process of HF miniaturization (Fig. 3A-C). Further analysis indicated a strong positive correlation between the number/ proportion of CD34[+] progenitor cells and the hair follicle diameter in balding scalps of AGA patients (Fig. 3D-G). Collectively, these observations implicated an intriguing role of progenitor cells in HF miniaturization.

Next, investigation into the causes of the progenitor cell loss via signaling pathway enrichment analysis revealed upregulated apoptotic signals in progenitor cells of balding anagen HFs (Fig. 3H). A significant increase in apoptosis was observed in the CD34[+] progenitor cells of balding anagen HFs according to TUNEL staining (Fig. 3I, J). These results support the notion that an apoptosis-induced gradual loss of CD34[+] progenitor cells may be involved in the HF miniaturization in AGA.

Hyperactive CTS contraction drives premature hair regression via inducing apoptosis of progenitor cells in AGA

Intimate crosstalk between stem cells and the surrounding microenvironment is vital for stem cell fate decisions. On the outside of the progenitor cells is a layer of elastic DS in murine pelage HF, while in human scalp HF is the multi-layer of connective tissue sheath (CTS) mainly composed of DS cells, collagens and vasculature (including vSMCs and endothelial cells) (Martino et al., 2021)(Fig. S6A; Fig. 4A, B). Pathway enrichment analysis revealed increased contractile activity in both DS cells and vSMCs of balding anagen HFs compared to non-balding or healthy groups (Fig. 2O and R), which can be confirmed by the immunostaining of phosphorylated myosin light chain 2 (p-MLC2), a marker of myosin active state(Nayak et al., 2020) (Fig. 4C, D; Fig. S6B, C). Collectively, these results suggest that CTS contraction is enhanced in anagen HFs of balding scalps in AGA patients.

Given that the contractile force of DS is a key driver of HF regression during hair regeneration in mice(Heitman et al., 2020), we wondered whether excessive contraction of the progenitor cell-encapsulating CTS, the analogous structure of murine DS in human, might affect hair follicle by modulating

progenitor cell fate in AGA. To this end, we utilized methacholine (MCH)(Huang

et al., 2012; Kudo et al., 2012) and norepinephrine (NE)(Cairns and Borrani,

2015; Lynch and Ryall, 2008) to activate CTS contraction in the in vitro cultured

human anagen HF organs. The efficiency of MCH and NE on contraction

activation in this model was evaluated and verified by p-MLC2 immunostaining

(Fig. S6D). Strikingly, HF growth was significantly inhibited (Fig. S6E-G) and

premature catagen induction was promoted (Fig. S6H, I) in HFs treated with

MCH or NE. Furthermore, the observed activation of CTS contraction by MCH

resulted in a dose-dependent negative effect on hair growth and anagen

maintenance (Fig. 4E-H, and Fig. S7A, B). To figure out whether progenitor cells

act as the targets of CTS contraction, we examined the effect of MCH on the

number and apoptosis of HF progenitor cells. The immunostaining of CD34 and

TUNEL showed that activated CTS contraction by MCH promoted the loss and

apoptosis of CD34[+] progenitor cells in a concentration-dependent manner. Of

note, TUNEL[+]CD34[+] cells are barely detectable under 100 μM MCH treatment

due to the fact that CD34+ progenitor cells were almost depleted under this

condition (Fig. 4I-L).

Notably, blockade of CTS contraction with ML-7 (Fig. S7C), a specific myosin

activation inhibitor(Colgren and Nichols, 2022), not only improved the growth

retardation and catagen premature induction (Fig. 4M-O, and Fig. S7D), but

also restored the apoptotic loss of progenitor cells in MCH-treated human HFs

(Fig. 4P, Q), confirming the critical role of CTS contraction in MCH effects.

Consistently, we demonstrated that the percentage of catagen HFs is indeed

markedly increased in the B scalps of AGA patients compared to NB or HS

scalps (Fig. 4R-T). Collectively, these findings

significantly strengthen the link between enhanced CTS contraction and AGA

pathogenesis.

Subsequently, to determine whether apoptosis is necessary, we applied Z-

VAD-FMK to block cell apoptosis(Li et al., 2019). Our results showed that the

observed phenotypes caused by MCH, including HF growth inhibition (Fig. 5A-

D), premature catagen induction (Fig. 5E) and depletion of CD34$^+$ progenitor cells (Fig. 5F-H), were counteracted after Z-VAD-FMK administration. Taken together, these data suggest that over-activated CTS contraction compresses hair growth and drives premature hair regression via promoting the apoptosis of progenitor cells.

CTS contraction induces progenitor cell apoptosis and premature hair regression via PIEZO1 signaling in AGA

To pinpoint the specific molecular mechanisms by which the contractility of CTS regulates progenitor cells, we first examined the expression levels of the known mechanosensitive ion channel protein PIEZO1, which has been found to be highly expressed in HFSCs in mice(Xie et al., 2022). Our results showed that PIEZO1 was abundantly expressed in CD34$^+$ progenitor cells (Fig. S8A) and was expressed at much higher levels in balding anagen HFs compared to non-balding group (Fig. 6A, B). We then used intracellular Ca^{2+} as a downstream indicator of PIEZO1 signaling(Kefauver et al., 2020; Ranade et al., 2015). As quantified by Fluo-4 (a Calcium influx indicator) RFI, there was a significant increase in intracellular Ca^{2+} concentration in CD34$^+$ progenitor cells of balding anagen HFs (Fig. 6C, D). These results suggest that PIEZO1 signaling is activated in progenitor cells of balding anagen HFs.

We next investigated whether activation of PIEZO1 is sufficient to induce HF growth retardation and premature regression. To this end, we applied YODA1, a PIEZO1-specific activator to cultured anagen HFs (Fig. S8B, C). Intriguingly, anagen HFs treated with YODA1 displayed a phenotype similar to MCH (Fig. 6E-I and Fig. S8D). In addition, the Ca^{2+} concentration in MCH-treated HF progenitor cells was elevated and could be reversed by the specific PIEZO1 inhibitor GsMTX4 and the myosin activation inhibitor ML-7 (Fig. 6J, K), indicating the activation of PIEZO1 signaling by CTS contraction. Not surprisingly, inhibition of PIEZO1 signaling by GsMTX4 rescued the phenotypes induced by MCH (Fig. 6L-P).

These results prompt us to speculate that CTS contraction induced in balding

anagen HFs activates the PIEZO1 signaling pathway in progenitor cells via mechano-transduction, thereby inducing progenitor cell apoptosis and thus promoting premature HF regression.

Targeting CTS contraction by ML-7 improves the growth of HFs from the balding scalps of AGA patients

All the above results inspired us to wonder whether targeting CTS contraction is a promising therapy for AGA. We first obtained anagen HFs from the balding scalps of AGA patients, which had been experiencing the process of miniaturization. As expected, the balding anagen HFs exhibited obvious growth retardation compared with those from occipital non-balding scalps of the same AGA patients in ex vivo organ culture system (Fig. S9A-C). We then treated the balding HFs with ML-7, or minoxidil (MNX), the FDA-approved drug for AGA treatment. Immunostaining of p-MLC2 confirmed that ML-7 abolished muscle contraction activity in CTS, but MNX did not (Fig. S9D). Excitingly, ML-7 could significantly improve the growth and premature regression of balding HFs of AGA patients, even better than MNX (Fig. 7A-D; Fig. S9E). Consistently, ML-7 showed a better effect on decreasing the apoptotic loss of progenitor cells compared with MNX (Fig. 7E, F). Taken together, these findings suggest that relaxing HFs via targeting CTS contraction is a hopeful strategy for AGA therapy.

Discussion

Here we have presented what is, to the best of our knowledge, the first and currently the only comprehensive single-cell transcriptome profiling of all cell types within HF units from balding and non-balding scalps of AGA patients compared with healthy individuals. Considering that HFs of balding scalps used in the present study were in anagen phase and at the early stage of hair miniaturization, isolated from the edge of frontal hairline of AGA patients, our high-quality data allowed us to illustrate the early changes in cell subpopulations, the HF cell lineage trajectory and cell-cell communications in AGA in unprecedented detail. Our atlas uncovers at least 16 different broad cell

types in human HF unit, which are more than those described in previous

scRNA-seq studies(Takahashi et al., 2020; Wu et al., 2022).

The reason for this obvious difference could be likely attributed to the

optimization of human HF single cell isolation by combining mechanical

separation with multienzyme digestion. Based on this, we obtained previously

undescribed DP cells and major cell types of CTS (including DS, vSMC and

endothelial cells), which are reported to be essential for regulating the growth

of hair follicles(Driskell et al., 2011; Heitman et al., 2020; Martino et al., 2021).

Delineation of DP and CTS cells at single-cell level provides an opportunity to

determine the precise roles of these cells in the pathogenesis of AGA.

The hair growth is powered by HFSCs (termed as bulge stem cells) that have

the ability to give rise to all epithelial components of the hair follicle(Chen et al.,

2020). However, current understanding of the cellular and molecular changes

in the bulge in AGA development is limited. Our scRNA-seq data show that

bulge stem cells contain two major subclusters: $CD200^+$ bulge stem cells and

$CD34^+$ progenitor cells. Thereinto, the number and percentage of progenitor

cells are significantly decreased in the balding anagen HFs of AGA patients.

These results are consistent with the previous study showing that balding scalp

in men with AGA retains HFSCs but lacks HF progenitor cells(Garza et al.,

2011). Besides, we demonstrate that HFSCs are more likely to differentiate into

interfollicular epidermal and sebaceous lineages, rather than HF lineage cells,

in the balding scalps, which may be responsible for the progressive hair follicle

miniaturization and hair thinning in AGA. Similarly, in the case of ageing- and

obesity-induced hair loss in mice, those aberrant fate changes also occur in a

fraction of HFSCs, which prefer to undergo epidermal and sebocyte

differentiation regulated by COL17A1 proteolysis and inflammatory signals

respectively, thereby decreasing the HFSC pool and leading to hair follicle

miniaturization in a stepwise manner(Matsumura et al., 2016; Morinaga et al.,

2021). Here, our data also suggest an important role of COL17A1 in the

aberrant differentiation of HFSCs in AGA development. During normal hair

growth, HF stem/progenitor cells proliferate and differentiate to generate hair

matrix cells, which rapidly proliferate to produce the hair shaft of growing hair

in the end(Martel et al., 2023). Here, we show that the proliferating matrix cells

are also decreased in balding HFs in AGA. It is tempting to speculate that the

apoptotic loss of progenitor cells contributes to a decrease in matrix cells, which

may cause the premature induction of catagen (namely shortening of anagen),

and eventually weakens the generation of terminal hair shaft in AGA

development.

DP acts as the signaling center to govern the proliferation and differentiation

of HF stem/progenitor cells in hair growth(Morgan, 2014; Plikus, 2012). Our

previous study demonstrated that androgen receptor (AR) is upregulated in DP

cells, which induce apoptosis of microvascular endothelial cells via paracrine

signaling in the DP of balding HFs in AGA(Deng et al., 2022). However, in this

study, except for the downregulated angiogenesis, our scRNA-seq analysis

identifies no other obvious changes in cell subpopulations and signaling

pathways, which may be involved in the regulation of hair growth, in DP cells of

balding HFs from AGA patients. The reasons for this discrepancy are currently

not clear, and further analysis and experimentation will be required to address

this point. Although microinflammation has been commonly reported in

AGA(Jaworsky et al., 1992; Magro et al., 2011; Martinez-Jacobo et al., 2018;

Peyravian et al., 2020), the details remain largely undefined. Here, we identity

that T cells are the main lymphocytes in human HF units, and AGA is featured

by increased Th17 cells.

Although a previous study has identified that progenitor cells are significantly

diminished by flow cytometry in the bald scalp of AGA patients(Garza et al.,

2011), the details in this aspect are obscure. In this study, we further confirm

that CD34[+] progenitor cells are decreased in the balding anagen HFs in AGA,

and determine that these progenitor cells are derived from CD200[+] bulge stem

cells via scRNA-seq analysis and immunostaining. Importantly, we also find that

the reduction in HF progenitor cells is attributed to the increased apoptosis, and

the number and percentage of these cells is positively correlated with the hair follicle size in the balding scalp in AGA. These findings suggest a pivotal role for progenitor cell depletion in the pathogenesis of AGA.

HF growth is tightly spatiotemporally regulated by the crosstalk between stem/progenitor cells and the microenvironment, mainly the mesenchymal niche(Plikus and Chuong, 2014). In addition to DP whose role is well recognized, DS is also an important component of the mesenchymal niche. DS is continuous with the DP and generates an envelope surrounding the hair follicle, whose role has been far less examined than the DP yet may be equivalently important for the regulation of hair growth(Martino et al., 2021). Until recently, DS has been identified as smooth muscle that drives hair follicle regression for reuniting HFSCs and niche to regenerate tissue structure during hair regeneration in mice(Heitman et al., 2020). Further study demonstrates that progenitor cell-derived endothelin governs DS contraction in mice(Martino et al., 2023). However, the role of CTS (the analogous structure of murine DS in human), in human hair follicle, especially in hair disorders, remains undefined. In this study, via single-cell sequencing analysis we identify that smooth muscle contraction activity is markedly increased in the major cell types of CTS, including DS and vSMC cells in balding anagen HFs of AGA patients. We further, by co-immunostaining, confirm this conclusion, and demonstrate that muscle contraction-related pathways are hyperactivated in the CTS of balding anagen HFs; In human hair follicle organ culture system, activation of muscle contraction in CTS drives premature hair follicle regression via inducing progenitor cell apoptosis. Surprisingly, we find that pharmacological inhibition of muscle contraction can obviously improve the growth and premature catagen induction of hair follicles from the balding scalps of AGA patients, even better than the FDA-approved drug minoxidil, strongly suggesting that targeting muscle contraction in CTS is a very promising therapy for AGA. It will also be interesting in future studies to determine the mechanism that hyperactivates the muscle contraction of CTS cells in the development of AGA.

Our results suggest PIEZO1, a mechanosensitive ion channel, as a regulator

in HF progenitor cells that senses the mechanical force generated by CTS

contraction. We show that PIEZO1 is expressed in HF progenitor cells, and it

becomes markedly upregulated and activated in the balding anagen HFs of

AGA patients. Inhibition of PIEZO1 signaling evidently alleviates progenitor cell

apoptosis and premature hair follicle regression induced by CTS contraction; in

contrast, PIEZO1 hyperactivation induces apoptosis of progenitor cells and

premature entry into regression in human anagen hair follicles. Similarly,

PIEZO1 has been reported to be abundantly expressed in HFSCs, and is

responsible for niche-size-decrease-triggered ectopic HFSC apoptosis in

mice(Xie et al., 2022). We conclude that the role of PIEZO1 in sensing the

abnormal mechanical force from the microenvironment to regulate the HFSC

apoptosis is conserved across species, and the mechanosensitive channel

might be a therapeutic target for hair loss. However, the detailed molecular

mechanisms by which aberrant PIEZO1 activation induces progenitor cell

apoptosis in AGA need further study to clarify.

In summary, the present study reveals the cellular hierarchies via scRNA-seq

analysis and demonstrates hyperactive CTS contraction as a critical cause of

hair miniaturization in AGA, strongly suggesting that relaxing hair follicle by

targeting CTS might be a promising therapy for this disorder.

Methods

Human scalp and HF samples

Human scalp skin biopsies and HF units were collected from non-balding

occipital and balding frontal sites of male AGA patients undergoing hair

transplantation surgery, and normal frontal sites of male healthy volunteers from

the Department of Dermatology in Xiangya Hospital, Central South University.

The procedures were in accordance with protocols set out in the WMA

Declaration of Helsinki and the Department of Health and Human Services

Belmont Report and approved by the ethical committee of the Xiangya Hospital

of Central South University and written informed consent was acquired from all
participants.

Identification of hair cycle phases in human HFs

Hair cycle phases of human HFs were determined by morphological analysis,
HE staining, co-immunostaining of Ki67 and TUNEL, and immunostaining of C-
Caspase-3 as previously described(Langan et al., 2015; Oh et al., 2016). With
these identification methods (Fig. S1), all human HFs used in this study
(including scRNA-seq, staining and in vitro organ culture) had been confirmed
in the typical anagen phase before experimentation unless otherwise stated.

Preparation of single-cell suspensions of HF units

To obtain single-cell suspension of entire HF units, DPs were isolated first by
micro-dissection from HFs and were chopped. The remaining hair follicle tissue
samples were cut with scissors and digested with the chopped DPs in 4mg/ml
Dispase II and Collagenase IV for 40 min at 37℃, centrifuged at 200g for 5min.
The pellet was resuspended and digested in 0.25% trypsin for 10 min at 37℃,
washed in PBS+5%FBS and filtered through a 70 µm cell strainer. The Dead
Cell Removal Kit (Miltenyi Biotec) was used in accordance with the
manufacturer's instruction to obtain live cells. 10 µL of the cell suspension were
mixed with the same volume of Trypan blue for cell counting. The loading
volume for sequencing was verified by the cell concentration

ScRNA-seq

scRNA-seq mainly includes GEM (gel bead-in-emulsion) generation, barcoding,
cDNA amplification, library construction and sequencing. These steps were
completed according to the user's instructions of Chromium Single Cell 3'
Reagent Kits v3.1 (10× Genomics, product code: 1000268, 1000215, 1000120)
(https://www.10xgenomics.com/support/single-cell-gene-expression). Libraries
were sequenced by an Illumina NovaSeq6000 System. Approximately 10,000
cells (targeting 5000–12,000) per sample were used for single-cell RNA
sequencing.

ScRNA-seq data processing and cell type identification

We used the default parameters of Cell Ranger software suite (v6.0.2) to align

and quantify the raw reads data from 10x Genomics. After the initial Cell Ranger

metric assessment, Seurat (v4.3.0)(Hao et al., 2021) in R version 4.3.1 was

used to exclude cells with fewer than 200 genes or more than 6000 genes

detected, and more than 20% mitochondrial reads from downstream analysis.

After quality control, 76368 cells remained and were used for downstream

bioinformatic analysis. The "NormalizeData" function was used to normalize the

feature expression measurements for each cell by the total expression. The

"ScaleData" function was used to scale and center the expression of each gene

for dimensional reduction. To avoid batch effects among samples and

experiments, we integrated data from all samples using harmony (v0.1.1). Total

cell clustering was performed by "FindClusters"function with Louvain algorithm

at a resolution of 0.2. Non-linear dimensional reduction was performed by

"RunUMAP" function and visualized by Uniform Manifold Approximation and

Projection (UMAP). Markers genes of each cell cluster were determined by

"FindAllMarkers" function with Wilcoxon rank-sum test. Only those with

|'avg_logFC'| ≥0.25 and 'adjusted P value'≤0.05 were considered as marker

genes.

For subgroup cell clustering, cells of each cell type were extracted separately

and clustered by their first 20 PCs and appropriate resolution. To further

approximate the low-dimensional data manifold representing the differentiation

trajectory for Bulge cells, spliced status of mRNAs in single-cell RNA

sequencing data were used to estimations of RNA velocities through

velocyto(La Manno et al., 2018). Markers genes of each subcluster were

identified by "FindAllMarkers" function with the default parameters.

"DoHeatmap" function were used to show top marker genes in heatmap.

Identification of balding-associated DEGs

566　We used the function of "FindMakers" in Seurat to identify balding-vs-

567　nonbalding and balding-vs-healthy differentially expressed genes (DEGs) for

568　each cell type and sub-cell type. The adjusted P values of each DEG were

calculated by on-parametric two-sided Wilcoxon rank-sum test and only those with |'avg_logFC'| > 0.25 and 'adjusted P value' < 0.05 were considered to be balding-associated DEGs.

Function enrichment analysis

Gene Set Enrichment Analysis (GSEA) of GO and KEGG were performed by clusterProfiler R package (v4.5.1.902) and visualized with ggplot2 R package (v3.4.1)(Bechhofer et al., 1986). Representative terms selected from the top ranked GO terms and KEGG pathways from MsigDB were displayed.

Pseudotime analysis

Monocle3 was used to reconstruct the differentiation trajectory for all cell types(Cao et al., 2019). The expression matrix of all cells was used as input and the UMAP embedding coordinates were import from Seurat result. The start points of bulge and IFE basal cells were specified as the root nodes separately for the SOX9$^+$ HF branch and SOX9$^-$ IFE branch of the trajectory graph by "order_cells" function. "Plot_cells" function was used to visualize each cell type along the pseudotime trajectory. To further reconstruct the differentiation trajectory and do "branched expression analysis modeling" (BAEM) for bulge cell fate cell determination related cell types, such as bulge to ORS and IFE, Monocle2(Qiu et al., 2017) was used to reconstruct the differentiation trajectory for small set of cell types and "BAEM" function were performed to identify cell determination related molecules.

Cell-cell communication analysis

To assess cell-cell communications among different cell types, we used cellchat (v1.1.3) to infer the intercellular communication network from single-cell RNA-seq data. Only cell types with more than 10 cells were considered in the analysis. The "trimean" method is used for calculating the average gene expression per cell group in "computeCommunProb" function. Pairwise comparison and visualization are performed by cellchat build-in function, only interactions with P value lower than 0.05 are considered to be real.

HF organ culture

Human HFs were obtained from the non-balding occipital sites (unless stated otherwise) of male AGA patients undergoing hair transplantation surgery. After microdissection, the HFs were first incubated in William'E medium (WEM, ThermoFisher) for 24 hours to re-equilibrate. HFs in anagen VI phase were randomly assigned to the different experimental groups. HFs were cultured at 37℃ with 5% CO_2 in WEM supplemented with 2 mM of L-glutamine, 10 ng/ml hydrocortisone, 10 ug/ml insulin and 1% penicillin/streptomycin mixure, and photographed under a stereoscope every two days to count hair shaft length and hair cycle stage. The hair cycle stage (anagen/early-catagen/mid-catagen) of HFs was determined according to the identification methods described above(Langan et al., 2015; Oh et al., 2016).

Chemical stimulation of HFs

All the chemicals were purchased from Selleck. After 24 hours, medium was replaced and HFs were treated with vehicle (0.1% DMSO), MCH (0.1,1 10,100 µM as indicated), NE (0.1 µM), ML-7(0.3 µM), Z-VAD-FMK (10 µM), YODA1(15 µM), or GsMTX4(1 µM) according to the design of different experiments and the medium was changed daily with the indicated treatment. HFs were cultured for 2-10 days for analysis of hair growth and immunostaining.

Immunostaining and TUNEL staining

Immunostaining was performed as previously described(Deng et al., 2021). Briefly, OCT-embedded samples were sectioned (10 µm) with a Leica cryostat. The sections were fixed for 10 min with 4% PFA, and washed with PBS, then blocked for 60 min with blocking buffer (5% NDS, 1% BSA, 0.3% Triton X-100). Primary antibodies were incubated overnight at 4℃. Secondary antibody was incubated for 60 min at room temperature. After washed, sections were counterstained with DAPI. TUENL staining was performed with TUNEL assay kit (Roche, USA) according to manufacturer's instructions, and then stained with CD34 following the immunostaining protocol. For the co-immunostaining of two antibodies from the same host species, double immunohistochemistry for paraffin sections (8 µm) was conducted by using the Opal 4 color manual

immunohistochemistry (IHC) kit (NEL810001KT, PerkinElmer). The signal of p-MLC2 was amplified by the Opal 4 color manual IHC kit (NEL810001KT, PerkinElmer). All pictures were taken with a Zeiss Axioplan 2 microscope. The fluorescence intensity was evaluated with ImageJ. The primary antibodies were used in this study: Rabbit anti-SOX9 (1:100, Cell Signaling), Mouse anti-K15 (1:2000, Lab vision), Mouse anti-K14 (1:4000, Abcam), Rabbit anti-KRT6A (1:500, Atlas Antibodies), Rabbit anti-KRT10 (1:1000, Abcam), Mouse anti-α-SMA (1:4000, Abcam), Rabbit anti-CD200 (1:200, Cell Signaling), Rabbit anti-CD34 (1:250, Abcam), Rabbit anti-p-MLC2 (1:100, Cell Signaling), Rabbit anti-Ki67 (1:500, Abcam), Rabbit anti-PIEZO1 (1:100, Proteintech). Staining intensity was evaluated in well-defined reference by quantitative (immuno-) histomorphometry using NIH ImageJ software.

In vitro Ca^{2+} imaging

HFs were incubated with Fluo-4 staining solution (Beyotime) in culture medium at 37℃ for 1 hour. Afterward HFs were washed with DPBS and cultured DPBS with 1mM calcium. For different treatments, MCH, ML-7, YODA1 or GsMTX4 was added to the culture medium for 30 min. The HFs were embedded in OCT and were sectioned with a Leica cryostat. Images were taken immediately to record the fluorescence intensity to indicate Ca^{2+} concentration and the adjacent section was used for CD34 staining to indicate progenitor cells.

Statistical analysis

All data were expressed as mean±SEM and were analyzed by ANOVA or Kruskall-Wallis test when more than two groups were compared or Student's t-test when only two groups were compared. Graphpad Prism 8, R (v4.1.3) and Excel (Microsoft) were used to assess statistical significance. Statistical significance was set at a $P < 0.05$. *, ** and *** indicate $P < 0.05$, $P < 0.01$ and $P < 0.001$, respectively.

Data availability

All data needed to assess the conclusions in the present study are provided in the manuscript and/or the Supplementary Materials. The scRNA-seq datasets

are available from the genome sequence archive under accession number
HRA005629 (http://bigd.big.ac.cn/gsa-human/). Any other data supporting the
findings of this study are available from the corresponding author upon
reasonable request.

Acknowledgements
This work was supported by the National Key Research and Development
Program of China (No. 2021YFF1201205), the National Natural Science Funds
for Distinguished Young Scholars (No. 82225039), the National Natural Science
Foundation of China (No. 82304057, No. 81874251, No. 82073457, No.
82173448, No. 82373508, No. 82303992), the Natural Science Funds of Hunan
province for excellent Young Scholars (No. 2023JJ20094). We thank our
colleagues (Department of Dermatology, Xiangya Hospital, Central South
University, China) for their generous support throughout this work.

Author contributions
J.L., Z.D., G.L. and L.Y. designed and conceived the study. Z.D., G.L. and L.Y.
performed data analyses. L.Y. and Z.D. performed scRNA-seq analysis. G.L.,
Z.D., S.D. and M.C. performed most experiments. Y.Z., F.L., Y.T., Y.W., S.X.,
Z.W., B.W., Z.Z. and W.S. contributed to sample collection. Z.D. and M.C. help
to generate sequencing libraries. H.X. and Y.Z. provide critical discussion and
suggestion. Z.D., G.L., L.Y. and J.L. prepared the manuscript with input from
coauthors.

Competing interests
The authors declare no competing interests.

References

Adil, A., and Godwin, M. (2017). The effectiveness of treatments for androgenetic alopecia: A systematic
review and meta-analysis. J Am Acad Dermatol *77*, 136-141 e135.

Bechhofer, D.H., Kornacki, J.A., Firshein, W., and Figurski, D.H. (1986). Gene control in broad host range

plasmid RK2: expression, polypeptide product, and multiple regulatory functions of korB. Proc Natl Acad

Sci U S A *83*, 394-398.

Cairns, S.P., and Borrani, F. (2015). beta-Adrenergic modulation of skeletal muscle contraction: key role

of excitation-contraction coupling. J Physiol *593*, 4713-4727.

Cao, J., Spielmann, M., Qiu, X., Huang, X., Ibrahim, D.M., Hill, A.J., Zhang, F., Mundlos, S., Christiansen,

L., Steemers, F.J.*, et al.* (2019). The single-cell transcriptional landscape of mammalian organogenesis.

Nature *566*, 496-502.

Ceruti, J.M., Leiros, G.J., and Balana, M.E. (2018). Androgens and androgen receptor action in skin and

hair follicles. Mol Cell Endocrinol *465*, 122-133.

Chen, C.L., Huang, W.Y., Wang, E.H.C., Tai, K.Y., and Lin, S.J. (2020). Functional complexity of hair follicle

stem cell niche and therapeutic targeting of niche dysfunction for hair regeneration. J Biomed Sci *27*,

43.

Colgren, J., and Nichols, S.A. (2022). MRTF specifies a muscle-like contractile module in Porifera. Nat

Commun *13*, 4134.

Cui, C.Y., and Schlessinger, D. (2015). Eccrine sweat gland development and sweat secretion. Exp

Dermatol *24*, 644-650.

Deng, Z., Chen, M., Liu, F., Wang, Y., Xu, S., Sha, K., Peng, Q., Wu, Z., Xiao, W., Liu, T.*, et al.* (2022).

Androgen Receptor-Mediated Paracrine Signaling Induces Regression of Blood Vessels in the Dermal

Papilla in Androgenetic Alopecia. J Invest Dermatol *142*, 2088-2099 e2089.

Deng, Z., Chen, M., Liu, Y., Xu, S., Ouyang, Y., Shi, W., Jian, D., Wang, B., Liu, F., Li, J.*, et al.* (2021). A

positive feedback loop between mTORC1 and cathelicidin promotes skin inflammation in rosacea.

EMBO Mol Med *13*, e13560.

Deng, Z., Lei, X., Zhang, X., Zhang, H., Liu, S., Chen, Q., Hu, H., Wang, X., Ning, L., Cao, Y.*, et al.* (2015).

mTOR signaling promotes stem cell activation via counterbalancing BMP-mediated suppression during

hair regeneration. J Mol Cell Biol *7*, 62-72.

Deutsch, A., McLellan, B.N., and Shinoda, K. (2020). Single-cell transcriptomics in dermatology. JAAD Int

1, 182-188.

Driskell, R.R., Clavel, C., Rendl, M., and Watt, F.M. (2011). Hair follicle dermal papilla cells at a glance. J

Cell Sci *124*, 1179-1182.

Festa, E., Fretz, J., Berry, R., Schmidt, B., Rodeheffer, M., Horowitz, M., and Horsley, V. (2011). Adipocyte

lineage cells contribute to the skin stem cell niche to drive hair cycling. Cell *146*, 761-771.

Garza, L.A., Liu, Y., Yang, Z., Alagesan, B., Lawson, J.A., Norberg, S.M., Loy, D.E., Zhao, T., Blatt, H.B.,

Stanton, D.C.*, et al.* (2012). Prostaglandin D2 inhibits hair growth and is elevated in bald scalp of men

with androgenetic alopecia. Sci Transl Med *4*, 126ra134.

Garza, L.A., Yang, C.C., Zhao, T., Blatt, H.B., Lee, M., He, H., Stanton, D.C., Carrasco, L., Spiegel, J.H., Tobias,

J.W.*, et al.* (2011). Bald scalp in men with androgenetic alopecia retains hair follicle stem cells but lacks

CD200-rich and CD34-positive hair follicle progenitor cells. J Clin Invest *121*, 613-622.

Gu, Y., Bian, Q., Zhou, Y., Huang, Q., and Gao, J. (2022). Hair follicle-targeting drug delivery strategies for

the management of hair follicle-associated disorders. Asian J Pharm Sci *17*, 333-352.

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., 3rd, Zheng, S., Butler, A., Lee, M.J., Wilk, A.J., Darby,

C., Zager, M.*, et al.* (2021). Integrated analysis of multimodal single-cell data. Cell *184*, 3573-3587 e3529.

He, H., Suryawanshi, H., Morozov, P., Gay-Mimbrera, J., Del Duca, E., Kim, H.J., Kameyama, N., Estrada,

Y., Der, E., Krueger, J.G.*, et al.* (2020). Single-cell transcriptome analysis of human skin identifies novel

fibroblast subpopulation and enrichment of immune subsets in atopic dermatitis. J Allergy Clin Immunol *145*, 1615-1628.

Heitman, N., Sennett, R., Mok, K.W., Saxena, N., Srivastava, D., Martino, P., Grisanti, L., Wang, Z., Ma'ayan, A., Rompolas, P.*, et al.* (2020). Dermal sheath contraction powers stem cell niche relocation during hair cycle regression. Science *367*, 161-166.

Huang, F., Zhang, H., Wu, M., Yang, H., Kudo, M., Peters, C.J., Woodruff, P.G., Solberg, O.D., Donne, M.L., Huang, X.*, et al.* (2012). Calcium-activated chloride channel TMEM16A modulates mucin secretion and airway smooth muscle contraction. Proc Natl Acad Sci U S A *109*, 16354-16359.

Hughes, T.K., Wadsworth, M.H., 2nd, Gierahn, T.M., Do, T., Weiss, D., Andrade, P.R., Ma, F., de Andrade Silva, B.J., Shao, S., Tsoi, L.C.*, et al.* (2020). Second-Strand Synthesis-Based Massively Parallel scRNA-Seq Reveals Cellular States and Molecular Features of Human Inflammatory Skin Pathologies. Immunity *53*, 878-894 e877.

Inui, S., and Itami, S. (2013). Androgen actions on the human hair follicle: perspectives. Exp Dermatol *22*, 168-171.

Jaworsky, C., Kligman, A.M., and Murphy, G.F. (1992). Characterization of inflammatory infiltrates in male pattern alopecia: implications for pathogenesis. Br J Dermatol *127*, 239-246.

Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.H., Myung, P., Plikus, M.V., and Nie, Q. (2021). Inference and analysis of cell-cell communication using CellChat. Nat Commun *12*, 1088.

Joost, S., Annusver, K., Jacob, T., Sun, X., Dalessandri, T., Sivan, U., Sequeira, I., Sandberg, R., and Kasper, M. (2020). The Molecular Anatomy of Mouse Skin during Hair Growth and Rest. Cell Stem Cell *26*, 441-457 e447.

Joost, S., Jacob, T., Sun, X., Annusver, K., La Manno, G., Sur, I., and Kasper, M. (2018). Single-Cell Transcriptomics of Traced Epidermal and Hair Follicle Stem Cells Reveals Rapid Adaptations during Wound Healing. Cell Rep *25*, 585-597 e587.

Joost, S., Zeisel, A., Jacob, T., Sun, X., La Manno, G., Lonnerberg, P., Linnarsson, S., and Kasper, M. (2016). Single-Cell Transcriptomics Reveals that Differentiation and Spatial Signatures Shape Epidermal and Hair Follicle Heterogeneity. Cell Syst *3*, 221-237 e229.

Katzer, T., Leite Junior, A., Beck, R., and da Silva, C. (2019). Physiopathology and current treatments of androgenetic alopecia: Going beyond androgens and anti-androgens. Dermatol Ther *32*, e13059.

Kefauver, J.M., Ward, A.B., and Patapoutian, A. (2020). Discoveries in structure and physiology of mechanically activated ion channels. Nature *587*, 567-576.

Kudo, M., Melton, A.C., Chen, C., Engler, M.B., Huang, K.E., Ren, X., Wang, Y., Bernstein, X., Li, J.T., Atabai, K.*, et al.* (2012). IL-17A produced by alphabeta T cells drives airway hyper-responsiveness in mice and enhances mouse and human airway smooth muscle contraction. Nat Med *18*, 547-554.

La Manno, G., Soldatov, R., Zeisel, A., Braun, E., Hochgerner, H., Petukhov, V., Lidschreiber, K., Kastriti, M.E., Lonnerberg, P., Furlan, A.*, et al.* (2018). RNA velocity of single cells. Nature *560*, 494-498.

Langan, E.A., Philpott, M.P., Kloepper, J.E., and Paus, R. (2015). Human hair follicle organ culture: theory, application and perspectives. Exp Dermatol *24*, 903-911.

Li, X., Yao, X., Zhu, Y., Zhang, H., Wang, H., Ma, Q., Yan, F., Yang, Y., Zhang, J., Shi, H.*, et al.* (2019). The Caspase Inhibitor Z-VAD-FMK Alleviates Endotoxic Shock via Inducing Macrophages Necroptosis and Promoting MDSCs-Mediated Inhibition of Macrophages Activation. Front Immunol *10*, 1824.

Liu, F., Hamer, M.A., Heilmann, S., Herold, C., Moebus, S., Hofman, A., Uitterlinden, A.G., Nothen, M.M., van Duijn, C.M., Nijsten, T.E.*, et al.* (2016). Prediction of male-pattern baldness from genotypes. Eur J Hum Genet *24*, 895-902.

Liu, X., Chen, W., Zeng, Q., Ma, B., Li, Z., Meng, T., Chen, J., Yu, N., Zhou, Z., and Long, X. (2022). Single-
Cell RNA-Sequencing Reveals Lineage-Specific Regulatory Changes of Fibroblasts and Vascular
Endothelial Cells in Keloids. J Invest Dermatol *142*, 124-135 e111.

Lolli, F., Pallotti, F., Rossi, A., Fortuna, M.C., Caro, G., Lenzi, A., Sansone, A., and Lombardo, F. (2017).
Androgenetic alopecia: a review. Endocrine *57*, 9-17.

Lynch, G.S., and Ryall, J.G. (2008). Role of beta-adrenoceptor signaling in skeletal muscle: implications
for muscle wasting and disease. Physiol Rev *88*, 729-767.

Magro, C.M., Rossi, A., Poe, J., Manhas-Bhutani, S., and Sadick, N. (2011). The role of inflammation and
immunity in the pathogenesis of androgenetic alopecia. J Drugs Dermatol *10*, 1404-1411.

Martel, J.L., Miao, J.H., and Badri, T. (2023). Anatomy, Hair Follicle. In StatPearls (Treasure Island (FL)).

Martinez-Jacobo, L., Villarreal-Villarreal, C.D., Ortiz-Lopez, R., Ocampo-Candiani, J., and Rojas-Martinez,
A. (2018). Genetic and molecular aspects of androgenetic alopecia. Indian J Dermatol Venereol Leprol
84, 263-268.

Martino, P., Sunkara, R., Heitman, N., Rangl, M., Brown, A., Saxena, N., Grisanti, L., Kohan, D., Yanagisawa,
M., and Rendl, M. (2023). Progenitor-derived endothelin controls dermal sheath contraction for hair
follicle regression. Nat Cell Biol *25*, 222-234.

Martino, P.A., Heitman, N., and Rendl, M. (2021). The dermal sheath: An emerging component of the
hair follicle stem cell niche. Exp Dermatol *30*, 512-521.

Matsumura, H., Mohri, Y., Binh, N.T., Morinaga, H., Fukuda, M., Ito, M., Kurata, S., Hoeijmakers, J., and
Nishimura, E.K. (2016). Hair follicle aging is driven by transepidermal elimination of stem cells via
COL17A1 proteolysis. Science *351*, aad4395.

Mereu, E., Lafzi, A., Moutinho, C., Ziegenhain, C., McCarthy, D.J., Alvarez-Varela, A., Batlle, E., Sagar,
Grun, D., Lau, J.K.*, et al.* (2020). Benchmarking single-cell RNA-sequencing protocols for cell atlas
projects. Nat Biotechnol *38*, 747-755.

Morgan, B.A. (2014). The dermal papilla: an instructive niche for epithelial stem and progenitor cells in
development and regeneration of the hair follicle. Cold Spring Harb Perspect Med *4*, a015180.

Morinaga, H., Mohri, Y., Grachtchouk, M., Asakawa, K., Matsumura, H., Oshima, M., Takayama, N., Kato,
T., Nishimori, Y., Sorimachi, Y.*, et al.* (2021). Obesity accelerates hair thinning by stem cell-centric
converging mechanisms. Nature *595*, 266-271.

Nayak, A., Wang, T., Franz, P., Steffen, W., Chizhov, I., Tsiavaliaris, G., and Amrute-Nayak, M. (2020).
Single-molecule analysis reveals that regulatory light chains fine-tune skeletal myosin II function. J Biol
Chem *295*, 7046-7059.

Ntshingila, S., Khumalo, N.P., Engel, M., and Arowolo, A.T. (2021). An appraisal of laboratory models of
androgenetic alopecia: A systematic review. Skin Health Dis *1*, e15.

Oh, J.W., Kloepper, J., Langan, E.A., Kim, Y., Yeo, J., Kim, M.J., Hsi, T.C., Rose, C., Yoon, G.S., Lee, S.J.*, et al.*
(2016). A Guide to Studying Human Hair Follicle Cycling In Vivo. J Invest Dermatol *136*, 34-44.

Papalexi, E., and Satija, R. (2018). Single-cell RNA sequencing to explore immune cell heterogeneity. Nat
Rev Immunol *18*, 35-45.

Peyravian, N., Deo, S., Daunert, S., and Jimenez, J.J. (2020). The Inflammatory Aspect of Male and
Female Pattern Hair Loss. J Inflamm Res *13*, 879-881.

Plikus, M.V. (2012). New activators and inhibitors in the hair cycle clock: targeting stem cells' state of
competence. J Invest Dermatol *132*, 1321-1324.

Plikus, M.V., and Chuong, C.M. (2014). Macroenvironmental regulation of hair cycling and collective
regenerative behavior. Cold Spring Harb Perspect Med *4*, a015198.

Potter, S.S. (2018). Single-cell RNA sequencing for the study of development, physiology and disease.
Nat Rev Nephrol *14*, 479-492.

Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H.A., and Trapnell, C. (2017). Reversed graph
embedding resolves complex single-cell trajectories. Nat Methods *14*, 979-982.

Rahmani, W., Abbasi, S., Hagner, A., Raharjo, E., Kumar, R., Hotta, A., Magness, S., Metzger, D., and
Biernaskie, J. (2014). Hair follicle dermal stem cells regenerate the dermal sheath, repopulate the
dermal papilla, and modulate hair type. Dev Cell *31*, 543-558.

Ranade, S.S., Syeda, R., and Patapoutian, A. (2015). Mechanically Activated Ion Channels. Neuron *87*,
1162-1179.

Takahashi, R., Grzenda, A., Allison, T.F., Rawnsley, J., Balin, S.J., Sabri, S., Plath, K., and Lowry, W.E. (2020).
Defining Transcriptional Signatures of Human Hair Follicle Cell States. J Invest Dermatol *140*, 764-773
e764.

Wu, S., Yu, Y., Liu, C., Zhang, X., Zhu, P., Peng, Y., Yan, X., Li, Y., Hua, P., Li, Q.*, et al.* (2022). Single-cell
transcriptomics reveals lineage trajectory of human scalp hair follicle and informs mechanisms of hair
graying. Cell Discov *8*, 49.

Xie, Y., Chen, D., Jiang, K., Song, L., Qian, N., Du, Y., Yang, Y., Wang, F., and Chen, T. (2022). Hair shaft
miniaturization causes stem cell depletion through mechanosensory signals mediated by a Piezo1-
calcium-TNF-alpha axis. Cell Stem Cell *29*, 70-85 e76.

Yang, H., Adam, R.C., Ge, Y., Hua, Z.L., and Fuchs, E. (2017). Epithelial-Mesenchymal Micro-niches
Govern Stem Cell Lineage Choices. Cell *169*, 483-496 e413.

Zou, Z., Long, X., Zhao, Q., Zheng, Y., Song, M., Ma, S., Jing, Y., Wang, S., He, Y., Esteban, C.R.*, et al.* (2021).
A Single-Cell Transcriptomic Atlas of Human Skin Aging. Dev Cell *56*, 383-397 e388.

Figures and Figure legends

Fig. 1: scRNA-seq reveals cell type composition in anagen HF units of AGA and healthy scalps.

A Flowchart overview of scRNA-seq, and subsequent validation and functional experiments with anagen HFs from balding frontal and non-balding occipital scalps of AGA patients, and normal scalps of healthy individuals. **B** Uniform manifold approximation and projection (UMAP) plot showing the twenty cell

types of human anagen HFs. DP, dermal papilla; DS, dermal sheath; Endo, endothelial cells; IFE, interfollicular epidermis; Inner II, inner layer of infundibulum and isthmus; IRS, inner root sheath; ORS, outer root sheath basal; ORS SB, outer root sheath suprabasal; Outer II, outer layer of infundibulum and isthmus; PR, proliferating progenitors; SwG, sweat gland; vSMC, vascular smooth muscle cells. **C** Heatmap of top 20 marker genes of each HF cell type. **D** Feature plots showing expression of marker genes for the Bulge (KRT15, KRT14), DP (VCAN, CFD), DS (ACTA2, CCN2), ORS (PTHLH), IRS (KRT6A). **E** Monocle3 pseudotime of all HF cell types. **F-I** Representative immunostaining images of KRT15/KRT14 (**F**), KRT6A/KRT15 (**G**), KRT15/KRT10 (**H**) and α-SMA (**I**) in human HF sections and the corresponding magnified views of boxed areas. (n=5 HFs from 3 individuals. The same 3 individuals' samples were used in all following HF stainings unless otherwise noted). Scale bar: 50 μm.

Fig. 2: Heterogeneity of cell subpopulations in AGA and healthy scalps.

A UMAP plots showing subclusters and sample conditions of bulge cells. **B** Bar

graph representing the percentage of subpopulations of bulge cells. **C-E**

Feature plots showing the expression of the KRT15 (**C**), CD200 (**D**) and CD34

(**E**) in bulge cells. **F** The ratio of CD34+ bulge cells in Healthy (HS), non-balding

(NB) and balding (B) conditions. **G, H** Representative images showing CD200

protein expression in upper bulge stem cells (**G**) and CD34 in progenitor cells

(**H**). Right panels, magnified images of boxed areas. Bu, bulge. Red, KRT15

(K15) staining. DAPI staining (blue) indicates nuclear localization. ORS–DS

boundary is indicated by dashed lines. Scale bar: 50 μm. **I** velocyto showing the single-cell trajectories of bulge subclusters in UMAP. **J** UMAP plots showing subclusters and sample conditions of DP cells. **K** Bar graph representing the percentage of subpopulations of DP cells. **L** Representative GO terms of upregulated (left) and downregulated (right) DEGs between balding and healthy groups (B vs HS), balding and non-balding groups (B vs NB) in DP cells. NES stands for enrichment scores. The color keys from red to blue indicate the range of P value. **M** UMAP plot showing subclusters and sample conditions of DS cells. **N** Bar graph representing the percentage of subpopulations of DS cells. **O** Representative GO terms enriched in balding versus non-balding or healthy DS cells. NES stands for enrichment scores. The color keys from red to blue indicate the range of p value. e. **P** UMAP plot showing subclusters and sample conditions of vSMC cells. **Q** Bar graph representing the percentage of subpopulations of vSMC cells. **R** Representative GO terms enriched in balding versus non-balding or healthy vSMC cells. NES stands for enrichment scores. The color keys from red to blue indicate the range of p value. The data represent the means±SEM. P values were determined by unpaired (B vs HS) or paired (B vs NB) student's t-test. ns, not significant.

Fig. 3: Apoptotic loss of HF progenitor cells is associated with HF miniaturization in AGA.

908
909

A Immunostaining images show progenitor cells in terminal HFs of occipital
910

scalps from healthy individuals (n=5 anagen HFs from 3 individuals), terminal
911

HFs of non-balding occipital scalps (n=8 anagen HFs) and intermediate/mini
912

HFs of balding frontal scalps (n=11 anagen HFs) from 3 AGA patients. Bulge
913

HFSCs and progenitor cells were labelled with K15 (green) and CD34 (red)
914

respectively and indicated by the dotted line. Right panels, magnified images
915

of boxed areas. Bu, bulge. Types of HFs were defined by their diameter.
916

Terminal: diameter>200 μm; Intermediate: 150 μm <diameter<200 μm; mini: diameter<150 μm. **B, C** Quantification of the number (**B**) of CD34$^+$ progenitor cells in different groups and their proportion in the total outer layer of HFs (**C**). **D-G** Scatter plots show the correlation between HF diameters and the number or percentage of CD34$^+$ progenitor cells in AGA patients. Spearman's correlation coefficient was used for the correlation analysis (two-tailed). **H** Top10 GO terms upregulated or downregulated in B versus in NB CD34$^+$ progenitor cells. NES stands for enrichment scores. The color keys from red to blue indicate the range of P value. **I, J** Representative images of TUNEL staining on anagen HFs from non-balding occipital scalps and balding frontal scalps of AGA patients (n=6 HFs from 3 donors for each group); Right panels, magnified images of boxed areas (**I**). The quantification of TUNEL$^+$CD34$^+$ cell numbers in two groups (**J**). Progenitor cells were indicated with dotted line. Scale bar: 50μm. The data represent the means±SEM. *p<0.05, **p<0.01, ***p<0.001, determined by one-way ANOVA with Tukey's post hoc test (**B, C**) or unpaired Student's t-test (**J**). ns, not significant.

Fig. 4: Hyperactive CTS contraction induces premature hair regression in

AGA.

A Dermal sheath (DS)/ connective tissue sheath (CTS) in murine pelage follicle/

human scalp follicle. In murine pelage follicle (left), the DS is the outer layer of

the ORS progenitors. In human scalp follicle, the outer layer is the CTS (right),

which is mainly composed of DS cells, vasculature and multiple types of

collagens. **B** Immunostaining of CD31 (an endothelial cell marker) and α-SMA

(a marker for both DS and vSMC cells) in human anagen HFs. CTS is located

between the two dotted lines. White arrows indicate DS. Yellow arrows indicate blood vessels. Right panels, magnified images of boxed areas. **C** The activation of DS muscle contraction was evaluated with p-MLC2 immunostaining on anagen HFs from non-balding occipital scalps (n=5 HFs) and balding frontal scalps (n=6 HFs) of 3 AGA patients. Right panels, magnified images of boxed areas. **D** Quantification of relative mean fluorescent intensity (RFI) of p-MLC2 in α-SMA positive cells in CTS in different groups. **E** Representative images of hair shaft elongation of human HFs ex vivo with vehicle (CON) or MCH treatment at concentration of 0.1 μM, 1 μM, 10 μM, 100 μM as indicated (n=21-32 HFs from 3 donors for each group). The same individuals' HF samples were used in the following staining of HFs with the same treatments. **F** Quantification of hair shaft elongation on day 2/4/6 with indicated treatments (n=32/24/29/21 in different groups respectively). **G** Macroscopic quantification of hair cycle stage of HFs with vehicle or MCH treatments on day 6. **H** Percentages of HFs in anagen on day 2/4/6 with indicated treatments. **I** Representative images of co-immunostaining of CD34 (green) and K15 (red) in HFs with vehicle or MCH treatments. Progenitor cells and bulge cells were indicated by dotted lines on green and red single-channel IF images, respectively. **J** Quantification of CD34$^+$ progenitor cells numbers and proportions in the total outer layer of HFs with vehicle or MCH treatments. **K** Apoptosis of progenitor cells with vehicle or MCH treatment were indicated with co-immunostaining of CD34 and TUNEL. Progenitor cells were indicated with dotted line. Right panels, magnified images of boxed areas. **L** Quantification of TUNEL$^+$CD34$^+$ progenitor cells in HFs treated with vehicle or MCH. To be mentioned, TUNEL$^+$CD34$^+$ progenitor cells are barely detectable under 100 μM MCH administration because CD34+ progenitor cells were almost vanished under this condition. **M** Representative images of hair shaft elongation of human HFs ex vivo with vehicle or MCH (10 μM) and ML-7 (0.3 μM) treatment as indicated (n=23-26 HFs from 3 donors for each group, independent experiments). **N** Quantification of hair shaft elongation on day 2/4 with vehicle or MCH and ML-7 treatments (n=24/26/23 in different

groups respectively). **O** Macroscopic quantification of hair cycle stage of HFs with vehicle or MCH and ML-7 treatments on day 4. **P** Co-immunostaining of CD34 and TUNEL showed progenitor cell apoptosis with indicated treatments (n=6 HFs from 3 donors for each group). Right panels, magnified images of boxed areas. **Q** Quantification of total and apoptotic CD34$^+$ progenitor cells numbers in HFs with vehicle or MCH and ML-7 treatments. **R** Representative images of HE staining with HS frontal scalps, NB occipital and B frontal scalps from AGA patients. A, anagen HF. C, catagen HF. T, telogen HF. **S** Macroscopic quantification of hair cycle stage of HFs in HE staining (n=197/489/228 HFs from 5 HS and 5 AGA donors). **T** Percentage of anagen, catagen and telogen HFs in HS, NB and B scalps (n=197/489/228 HFs from 5 HS and 5 AGA donors). Scale bar, 50 μm. Data are expressed as mean±SEM and were analyzed by one-way ANOVA with Tukey's post hoc test (J, L, Q, T) and two-way ANOVA with a post hoc Holm–Sidak's multiple comparisons test (F, N), or unpaired Student's t-test (D). *p<0.05, **p<0.01, ***p<0.001. ns, not significant. NA, not applicable.

Fig. 5: CTS contraction drives premature hair regression in an apoptosis-dependent manner.

A Representative images of hair shaft elongation of human HFs ex vivo with or without MCH (10 μM) and Z-VAD-FMK (10 μM) as indicated (n=24-30 anagen HFs from 3 donors for each group). The same individuals' HF samples were used in the following staining of HFs with the same treatments. **B** Quantification of hair shaft elongation on day 2/4/6 with indicated treatments. **C, D** Scatter plots show the elongation length of hair shaft with indicated treatments on day 2 (**C**) and day 4 (**D**). **E** Macroscopic quantification of hair cycle stage of HFs with indicated treatments on day 4. **F** Co-immunostaining of CD34 and TUNEL showed apoptosis of progenitor cells with indicated treatments (n=5 HFs for each group from 3 individuals). Progenitor cells were indicated with dotted line.

Right panels, magnified images of boxed areas. **G, H** Quantification of total (G)

and apoptosis (H) CD34+ progenitor cell numbers in HFs with indicated

treatments. Scale bar: 50 µm. The data represent the means±SEM. *p<0.05,

**p<0.01, ***p<0.001, determined by one-way ANOVA with Tukey's post hoc

test (C, D, G, H) and two-way ANOVA with a post hoc Holm–Sidak's multiple

comparisons test (B).

1007

1008 **Fig. 6: PIEZO1 signaling is necessary for CTS contraction-induced**

1009 **progenitor cell apoptosis and premature hair regression.**

1010 **A, B** (**A**) Representative immunostaining of PIEZO1 on skin sections from

1011 balding frontal and non-balding occipital scalps of AGA patients (n=3) and (**B**)

1012 the quantification of mean fluorescent intensity in outermost layer of mid-HF

cells (n=70-90 cells from 3 patients). **C, D** (**C**) CD34 and Fluo-4 staining were performed on serial sections from balding frontal and non-balding occipital scalps of AGA patients (n=3) and (**D**) the Fluo-4 mean fluorescent intensity in CD34$^+$ progenitor cells was quantified (n=100 cells in non-balding HFs and 80 cells in balding HFs from 3 patients). Fluo-4 indicates the cellular Ca^{2+} concentration. **E** Representative images of hair shaft elongation of human HFs ex vivo with or without YODA1 (15 μM) as indicated (n=22-26 anagen HFs from 3 individuals for each group). The same individuals' HF samples were used in the following staining of HFs with the same treatments. **F** Quantification of hair shaft elongation on day 2/4 with vehicle or YODA1 treatments. **G** Macroscopic quantification of hair cycle stage of HFs with YODA1 treatment on day 4. **H** Co-immunostaining of CD34 and TUNEL showed apoptosis of progenitor cells with YODA1 treatment. 5 HFs from 3 individuals were performed with immunostaining. **I** Quantification of total and apoptotic CD34$^+$ progenitor cell numbers in HFs treated with YODA1. **J, K** (**J**) CD34 and Fluo-4 staining were performed on serial sections of HFs with indicated treatments (n=3 HFs for each group) and (**K**) the Fluo-4 mean fluorescent intensity in CD34$^+$ progenitor cells was quantified (n=80-100 cells from 3 different HFs). **L** Representative images of hair shaft elongation of human anagen HFs ex vivo treated with MCH (10 μM) and GsMTX4(1 μM) (n=25 HFs for CON, 28 HFs for MCH and 26 HFs for MCH+GsMTX4; from 3 individuals). The same individuals' HF samples were used in the following staining of HFs with the same treatments. **M** Quantification of hair shaft elongation on day 2/4 with MCH and GsMTX4 treatments. **N** Macroscopic quantification of hair cycle stage of HFs with MCH and GsMTX4 treatments on day 4. **O** Co-immunostaining of CD34 and TUNEL showed apoptosis of progenitor cells with MCH and GsMTX4 treatments. 5 HFs from 3 individuals were performed with staining. **P** Quantification of total and apoptotic CD34$^+$ progenitor cell numbers in HFs treated with MCH and GsMTX4. Right panels, magnified images of boxed areas. Scale bar: 50 μm. Data are expressed as mean±SEM and were analyzed by one-way ANOVA with Tukey's

 post hoc test (K, P), two-way ANOVA with a post hoc Holm–Sidak's multiple

 comparisons test (F, M), and two-tailed unpaired Student's t-test (B, D, I).

 *p<0.05, **p<0.01, ***p<0.001.

1046

1047

1048 Fig. 7: ML-7 exhibits a better effect than MNX on promoting the growth of

1049 HFs from the balding scalps of AGA patients.

A Representative images of hair shaft elongation of anagen HFs from balding

scalps of AGA patients ex vivo treated with vehicle (CON), ML-7 (0.3 µM) or

minoxidil (MNX) ((100 µM)) as indicated (n=31-37 balding HFs from 5 AGA

patients for each group). 3 patients' HF samples were used in the following

staining. **B** Quantification of hair shaft elongation on day 2/4/6 with indicated

treatments. **C** Scatter plots show the elongation length of hair shaft on day 6. **D**

Macroscopic quantification of hair cycle stage of HFs with indicated treatments on day 6. **E** Co-immunostaining of CD34 and TUNEL showed apoptosis of progenitor cells with indicated treatments (n=7 balding HFs from 3 AGA patients for each group). Right panels, magnified images of boxed areas. **F** Quantification of total and apoptosis CD34$^+$ progenitor cell numbers in HFs with indicated treatments. Scale bar: 50 µm. The data represent the means±SEM. *$p<0.05$, **$p<0.01$, ***$p<0.001$, determined by one-way ANOVA with Tukey's post hoc test (C, F) and two-way ANOVA with a post hoc Holm–Sidak's multiple comparisons test (B).