1

TITLE: Machine Learning Based Modelling of Human and Insect Olfaction Screens Millions of compounds to Identify Pleasant Smelling Insect Repellents

AUTHORS: Joel Kowalewski[1], Sean M. Boyle[2], Ryan Arvidson[2], Jadrian Ejercito[2], and Anandasankar Ray[1,2]

AFFILIATION:

[1]Interdepartmental Neuroscience Program

[2]Department of Molecular, Cell and Systems Biology

University of California, Riverside, CA 92521, USA

CORRESPONDING AUTHOR: Anandasankar Ray

Department of Molecular Cell and Systems Biology, University of California Riverside, 3401 Watkins Drive, Riverside, CA 92521, USA, Phone:+1(951) 827-5998, Email: anand.ray@ucr.edu

Abstract

The rational discovery of behaviorally active odorants is impeded by a lack of understanding on how the olfactory system generates percept or valence for a volatile chemical. In previous studies we showed that chemical informatics could be used to model prediction of ligands for a large repertoire of odorant receptors in *Drosophila* (Boyle et al., 2013). However, it remained difficult to predict behavioral valence of volatiles since the activities of a large ensembles of odor receptors encode odor information, and little is known of the complex information processing circuitry. This is a systems-level challenge well-suited for Machine-learning approaches which we have used to model olfaction in two organisms with completely unrelated olfactory receptor proteins: humans (~400 GPCRs) and insects (~100 ion-channels). We use chemical structure-based Machine Learning models for prediction of valence in insects and for 146 human odor characters. Using these predictive models, we evaluate a vast chemical space of >10 million compounds *in silico.* Validations of human and insect behaviors yield very high success rates. The discovery of desirable fragrances for humans that are highly repulsive to insects offers a powerful integrated approach to discover new insect repellents.

Introduction

Animals can detect volatile molecules in their environment using specialized protein receptors in their olfactory organs. Mammalian odorant receptors were discovered in 1991 and are G-protein coupled receptors (GPCRs) numbering >400 in humans (Buck L and Axel, 1991). Several years later starting in 1999 three families of distinct insect odorant receptors were discovered from the *Drosophila* genome which are ligand-gated ion channels and number ~100 (Clyne et al., 1999, 2000). Over the subsequent 20-30 years, studies in mice and insects focusing on odor detection, odorant receptors, odorant receptor

neurons, and higher-level circuitry, have revealed important principles. A single volatile is usually

detected by a combination of receptors, and the activity from the neural circuit processing the

information is extremely complex. Previously, we successfully developed a chemical informatics

approach to discover ligands for a large number of odorant receptors in *Drosophila* (Boyle et al., 2013).

However, the mechanisms connecting olfactory perception (e.g., rose, lemon, pleasant, unpleasant, etc.)

to olfactory receptor activity are incompletely understood and it is therefore challenging to predict

behavior from odorant receptor activity alone. There is also often a lack of correspondence between

simple, 2D representations of odorants and their perceptual descriptions, and the molecular-cellular

organization of the olfactory system lacks a simple discernible coding strategy that enables predictions,

unlike auditory or visual processing (Buck L and Axel, 1991; Grabe & Sachse, 2018; P Mombaerts, 1999;

Peter Mombaerts, 2001; Sachse & Beshel, 2016; Vassar et al., 1993).

Yet Machine Learning algorithms have shown promise in accurately predicting human odor

perception (Keller et al., 2017; Kowalewski et al., 2021; Kowalewski & Ray, 2020b). Recently, we have

shown that the volatile structure-to-percept prediction can be improved by using the activities of a small

number of human odorant receptors and applying machine learning algorithms to identify highly

predictive combinations (Kowalewski & Ray, 2020b). In insects, we have limited success modeling the

behavioral valence of volatiles from olfactory receptor neuron activity in the fruit fly, *Drosophila*

melanogaster. Since receptor activity depends on chemical structure, this suggests machine learning

algorithms can potentially detect a consistent physicochemical basis for simple behavioral or complex

perceptual responses (Kowalewski & Ray, 2020b; MacWilliam et al., 2018).

To test whether Machine learning based modeling can lead to identification of strong

behaviorally active volatiles, we selected a challenge that has a very high potential for easing human

suffering and yet had little innovation: pleasant smelling fragrances that are also repulsive to harmful

insects that transmit diseases to humans or cause agricultural loss. Annually, mosquitos account for

~500,000-700,000 deaths and a billion infections due to vector-borne illnesses like malaria, Dengue,

Yellow Fever. Insects also destroy ~30% of the global agricultural produce causing additional suffering.

Insects are currently managed through widespread application of insecticides, but mutations are

emerging that confer insecticide resistance and affect insect olfactory responses in both human disease

vectors and plant pests (Deletre et al., 2019; Duneau et al., 2018).

The most widely used topical repellent for mosquitoes is DEET (N,N-Diethyl-m-toluamide), a

chemical with undesirable chemical and cosmetic properties. DEET (N,N-Diethyl-m-toluamide) was

discovered >75 years ago through empirical testing. It is not widely used outside the developed world,

where vector borne diseases are present globally, and are also use restrictions due to its strong solvent

properties, even dissolving plastics and nylons. Recent studies have also raised human health concerns ,

as the chemical interferes with inhibitory neurotransmission in mammals via $GABA_A$ and Glycine

receptors (Grant & Hall, 2019). Unfortunately, not enough is understood about the odorant receptors

for aversion in mosquitoes as a result there has been limited innovation in the search for repellents even

a quarter century after discovery of insect odorant receptors. The last repellent registered by the US EPA

for commercial use was Picaridin, over 20 years ago.

Volatile compounds that have pleasant fragrances are desired by humans in cosmetics and are

therefore excellent candidates for novel mosquito repellents. In order to identify fragrances that would

be aversive to insects, we assumed machine learning (ML) could be used to identify structural features

enriched among repellents and then detect these features within any arbitrary, large chemical space

(Maldonado et al., 2006). We trained a support vector machine (SVM) on a comprehensive set of

chemicals with known repellency from literature and identified predictive physicochemical features. In

parallel, we adapted a collection of recently developed ML models to predict 146 different human odor

percepts, helping us organize the candidate repellents by their potential fragrances (Kowalewski et al.,

2021; Kowalewski & Ray, 2020b). We then validated the accuracy of behavioral predictions for insects

and humans by randomly testing 50 predicted chemicals for repellency in mosquitoes and odor

perception in humans. The validation success was >75% in each assay, repellent and perceptual, with

~95% of the chemicals tested for repellency in round 1 showing aversion, which left ~5% as neutral or

slightly attractive to mosquitoes. Collectively, this suggests a strong physicochemical basis for odor

valence and percept in insects and humans. In addition, we also built *in silico* toxicity models, filtering

the compounds across multiple parameters for human use. A small number of the fragrances were

tested on evolutionarily distinct insects, finding that they are repellent to mosquitoes and the fruit fly,

Drosophila.

Tens of millions of dollars and several years may be required for identification and adequate

human-safety analysis for registration of new repellent chemicals by the EPA (Gupta & Bhattacharjee,

2006). Machine learning approaches for modeling toxicity as well as other physical and cosmetic

attributes are steadily advancing, suggesting it is possible to accelerate this search and significantly

reduce costs. Accordingly, we included additional computational filters to select predicted repellents

such as predicted fragrant odor perceptual qualities. Because fragrant molecules that are also insect

repellents may be rare to find, we applied this approach to a large library (>10 million) of purchasable

chemicals from the ZINC 15 database. These efforts represent a dramatic expansion of the insect

repellent chemical space and provides a theory-based high-throughput method to drive rational

discovery of novel insect repellents.

Results

Machine learning can learn to predict aversive valence in insects from chemical structure

To predict insect behavior, we developed a novel chemical informatics approach using Machine Learning

(ML). We assembled a training set of volatile chemicals known in the literature to have a negative

behavioral valence including eucalyptol, linalool, geraniol, citronella, alpha-thujone, beta-thujone (Kline

et al., 2003; Klocke et al., 1987; Syed & Leal, 2008), DEET, picaridin, 34 N-acyl piperidines that are

structurally related to picaridin (Katritzky et al., 2008) and structurally diverse odorants from fruits and

food sources that are not expected to elicit aversion (Carey et al., 2010; Hallem & Carlson, 2006). We

generated energy-minimized 3D structures for each and calculated 3,242 physicochemical features for

201 training chemicals. From it we identified 18 features that were highly correlated with aversive

valence (correlation of 0.912). We then fit a support vector machine (SVM) with these features to learn

to predict the repellency of test chemicals (Figure 1).

The trained model successfully predicted the valence of the test chemicals (avg AUC = 0.994),

indicating that it could distinguish volatiles with aversive valence from non-repellents (Figure 1).

Evaluation of the predictive chemical features indicated six membered rings, carbon-nitrogen distances,

tertiary amides, and the positioning of oxygen atoms in the molecule are associated with repellency

(Figure 1).

We subsequently used this repellency model to evaluate a >440,000 small molecules assembled

from databases which includes ~3,000 chemicals from natural sources like plants, insects, vertebrates,

and compounds that are in food, beverages, and cosmetics. The top 500 predicted compounds (top

~0.1% of hits) represent a structurally diverse group of chemicals, with some showing similarities to

volatiles in the training set (Figure 1). Several naturally occurring compounds were also in the list of

precited repellents. Since the ML model successfully classified known repellents versus non-repellents,

we next experimentally tested the candidate repellents.

Machine learning can prioritize candidate repellents by modeling their odor qualities to humans

Importantly, repellents must meet many physical and cosmetic demands for human use. However,

evaluating the scent characteristics of candidate repellents with human subjects is a laborious,

expensive, and time-consuming task. In order to create a method to bypass this bottleneck, we adapted

our recent perceptual modeling approach involving the prediction of 146 odor perceptual descriptors

from chemical structure; example perceptual descriptors include "fruity other than citrus", "lemon",

"orange", "cinnamon", "perfumery" and unpleasant ones such as "sickening", "rancid", "animal", and

"dirty linen" to name a few (Kowalewski et al., 2021; Kowalewski & Ray, 2020b). The models were

trained with data from prior human behavior studies in which participants evaluated ~150 odorants and

mixtures according to the 146 descriptors. Using these models, we predicted odor descriptors for the

candidate insect repellents that were discovered using the model described in Figure 1. The compounds

had a wide variety of predicted odor characteristics covering both pleasant and unpleasant fragrance

profiles. Together with the insect valence predictions, this gives us the foundations for selecting

desirable fragrances that can protect humans and plants from pests.

Validation of computational behavioral predictions of human smell and insect aversion

To validate the behavioral predictions of scent or repellency, on humans and insects,

respectively, we selected 47 compounds based on their availability and affordability. First, we compared

them to reports from human testing of smell properties. While odor qualities remain poorly

characterized, 37 of the 47 tested had odor and taste annotation in a sensorial databases at the The

Good Scents Company. We could then rigorously evaluate the accuracy of the trained perceptual

models by ROC analysis. The perceptual models predicted the observed descriptors for most volatiles

with a high success rate (Avg. AUC = 0.77) (Figure 2A,B). Interestingly, many chemicals in the predicted

repellents have pleasant odor characteristics, including highly desirable ones such as fruity, floral or

perfume-like, such as seen in the word cloud of enriched predicted characters for the predicted insect

repellents (Figure 2C).

We next tested the compounds against *Aedes aegypti* mosquitoes. We developed a high

throughput assay utilizing the innate attraction of the female mosquitoes to a heat block approximating

human skin temperature (37 °C) covered with a filter paper (Figure 3A). Each of the test compounds

were individually evaluated by applying on the filter paper (3% in acetone) and counting the numbers of

mosquitoes landing on the surface with solvent control before and after applying the compound. A

repellency index indicated that most of the predicted compounds (~90%) showed >50% repellency to

female *A. aegypti* mosquitoes (Figure 3B). These results suggest that the machine learning (ML)

algorithm was capable o predicting a chemical's perceptual qualities for humans and its valence for

insects. This set of chemicals represents a very large increase in new insect repellents, comprising

diverse chemical structures and smell characteristics that are desirable for human use.

Since we do not know the molecular receptor targets for these diverse repellents, and also know

that different insect species have both highly diverse and conserved olfactory receptors, we cannot tell

a priori whether the repellency is narrowly effective in mosquitoes or conserved across diverse insects.

To determine broad-spectrum aversive valence among insects, we tested another distantly related

dipteran, *Drosophila melanogaster*, which is >250 MY apart evolutionarily. All of the predicted

compounds displayed concentration-dependent aversive valence to *Drosophila* in a trap-based 2-choice

assay (Reeder et al., 2009; Syed et al., 2011). Most of the tested compounds showed strong repellency

(Figure 3C). The simplest interpretation is that the machine learning algorithm was able to identify

chemical features that predict negative valence across multiple species of dipteran insects.

Repellents show calcium mobilization in mammalian cells

An alternative approach to identifying insect repellents would be to computationally model a known

cellular or molecular activity relevant to repellency. Interestingly, one of the aversive compounds used

in the repellency training set is the commercial repellent DEET, had previously been seen to cause

calcium mobilization in human HEK293 cells at high concentrations (Dennis et al., 2018, Supplement). In

order to further evaluate if such Ca^{2+} mobilization is a common feature for repellency, we performed

additional experiments.

When HEK293 cells were exposed to DEET, a dose-dependent calcium transient was observed,

with an EC50 of 56.6 mM (~1% vol/vol), below the concentrations used on skin applied formulations (7-

98% vol/vol) of which up to 60% is absorbed into the skin (Schoenig et al., 1996; Selim et al., 1995).

(Figure 4A). Additional cell lines from different organisms were tested, including an insect S2 cell line,

and all of them were sensitive to DEET at similar EC50s (Figure 4A). The DEET-induced calcium transients

were similar in magnitude to the positive control, a 100 µM ATP induced signal (e.g., GPCR mediated)

(Figure 4B). To assess the source of DEET induced calcium flux, cells were depleted of intercellular

calcium with treatment with thapsigargin (a SERCA antagonist) 10 minutes before the assay. These

calcium-depleted cells did not show increased Ca transients when treated with DEET. This indicates that

DEET is likely inducing release from intercellular calcium stores, in a dose-dependent manner (Figure

4C). Interestingly, two additional commercially used repellents, picaridin and diethyl benzamide (a

structural analog of DEET), also evoked a strong intracellular Ca^{2+} signal (Figure 4D). Because these three

compounds were part of the training data to build our machine learning insect repellent models, we

tested whether the newly identified aversive fragrances also showed similar activity. We tested a

sampling of the verified aversive fragrances, and a few traditionally known repellents, at two

concentrations (5% and 0.5%) in the HEK293 cell line and found that there were indeed some that

showed Ca^{2+} release (Figure 4E). However, there appeared to be no correlation between the repellency

index and the Ca^{2+} activation of the fragrant repellents (R^2=0.0093), suggesting that this conserved effect

is unlikely to contribute to aversive behavior.

Nevertheless, the data provided an opportunity to develop a better understanding of the

physicochemical basis of the Ca^{2+} mobilization seen *in vitro* and we trained machine learning models to

predict the calcium activity (Figure S1A). Upon analyses, the top chemical features for predicting the

calcium activity were different from the features selected by the model to predict repellency. This too

suggest that while there is a structural basis for basis for calcium activity which does not overlap with

shared features for detection as a repellent. The features fell into 3 broad categories including 2D

autocorrelation, RDF, and 3D MoRSE features. Autocorrelations are distance-based metrics, with the 2D

autocorrelation referring to the number of bonds between a pair of atoms in the molecule, while radial

distribution function (RDF) features encode dependencies between atoms that give rise to molecules'

characteristic infrared spectra (Hemmer et al., 1999). The 3D MoRSE features are based on theoretical

electron diffraction patterns and are therefore sensitive to differences in 3D geometries (Devinyak et al.,

2014). In previous work, these features have been shown to be predictive of ligand bonding and

biological activity (Beglari et al., 2020; Cheng et al., 2010; González et al., 2006).

Evaluation of models trained with these features on test data showed a high correlation

between the predicted and observed Ca^{2+} activities (Figure S1). Using another approach, where we

classified the chemicals as broadly falling into one of two categories (high versus low activity), the

predictions were also accurate, further emphasizing that the Ca^{2+} signal does indeed have a strong

physicochemical basis. Collectively, this suggests the unique 3D shape, rather than generic, possibly

indiscriminate information, underlies the observed Ca^{2+} activity, supporting our contention that the

activity may be a persistent off target effect among certain chemical repellents and should therefore be

further investigated. While the physiological consequences of this effect have not been studied, we

incorporated this activity as a parameter into our machine learning pipeline, allowing us to further triage

repellents for future use by low estimated Ca^{2+} activity, in addition to other desirable characteristics for

human or animal use.

Mining massive commercially available chemical spaces for novel insect repellents

The size and diversity of the chemical library we initially screened (~0.45M) is in the range of high-

throughput wet lab screening that pesticide discovery companies can undertake. The Machine learning

approach allows us to screen far more without the limitations of cost of synthesis and screening time. To

demonstrate the massive throughput capacity of this approach, we increased our search space to >10

million commercially available chemicals (from the ZINC 15 database), canvasing far more structural

diversity than the ~450,000 chemicals we previously screened. We built a new training set including the

chemical repellents we experimentally verified, more than doubling the number of the actives while

adding greater structural diversity. The top features fell mostly into 2D and 3D autocorrelations and 3D

MoRSE categories, suggesting that certain types of features became increasingly important following the

addition of more repellent activities. Notably, some features changed from this vastly improved training

set; that is, moving away from simple features that account for amides, or more generally, chemical

bonds including nitrogen (N). This possibly reflects an initial bias toward DEET, an amide, that was

resolved when including more complex feature selection methods and increasing the size and diversity

of the training set.

There was a strong correlation between the predicted and observed repellency when validating

these updated models on test data (R = 0.73). Then, by applying the filters that we progressively

incorporated into the pipeline, and described throughout (e.g., insect valence, odor perceptual qualities,

and intracellular calcium release), we identified a set of candidates still numbering in the hundreds. We

next predicted the acute oral mammalian toxicity (Rat) using a computational model from our earlier

work (Kowalewski & Ray, 2020a) and eliminated all compounds falling into EPA Tox categories I and II.

The top predictions contain many aminobenzoates comprised of benzene, amino, and ester groups.

Some chemicals notably differ in that they resemble amide repellents like diethyltoluamide (DEET)

(Figure 5A). The top hits were predicted to have a variety of different human odor characters, many of

which are perceptually pleasant or are useful in agriculture (Figure 5B). This approach accelerates the

identification of novel, fragrance-based, broad spectrum insect repellents.

Discussion

Here, we have developed a comprehensive machine learning pipeline to screen small or large

chemical spaces for suitable alternatives and verified numerous predicted chemicals for repellency in

various behavioral assays and across species, indicating the effectiveness of the pipeline. The rapid

identification of fragrances that have excellent behavioral activity, pleasant smell and safety greatly

improves the likelihood of regulatory approval and potential for use.

Interestingly, we identified a strong intracellular Ca^{2+} signal in response to DEET and picaridin as

well as some anthranilates. While the mechanism remains unclear, it has been previously observed in

cultured cell lines after DEET exposure (Dennis et al., 2018; Swale et al., 2014). One possibility proposed

in the literature is that the elevated intracellular Ca^{2+} is due to activation of octopamine receptors.

Octopamine plays a modulatory role, with targets including OctαR and OctαβR in insects and TAAR1 in

mammals (Kleinau et al., 2011; Swale et al., 2014), and while this could explain the many complex

physiological responses that have reported in the DEET literature (Briassoulis, 2001; Clem et al., 1993;

Corbel et al., 2009; Tavares et al., 2019), an unknown but conserved receptor subtype is still possible.

Additional research will be needed to determine whether this phenomenon is important for off-target

mammalian toxicity. In either case, the model we developed to predict the Ca^{2+} mobilization may help

isolate a set of key physicochemical features, enabling identification of compounds that have this effect.

However, since the calcium activity does not explain the repellency, additional electrophysiological and

genetic experiments are needed to further clarify the mechanism of action.

Although many repellents have been studied, DEET remains the most widely used commercially

available chemical repellent. Yet research continues to emerge about off-target toxicities in humans

who are repeatedly exposed to DEET; specifically, accounts suggest adverse skin reactions and activities

in human or broadly mammalian nervous systems (Briassoulis, 2001; Grant & Hall, 2019; Osimitz et al.,

2010). This study demonstrates the successful development and application of a machine learning

pipeline that accelerates research into insect repellency and its physicochemical basis. The methods and

data presented provide a high-throughput and rational way to identify additional novel repellent

chemicals. Here, we emphasized screening candidate chemicals that are most likely to fit multiple

requirements (smell and repellency) rather than simply repellency. As these requirements steadily

increase, it is obvious that massive, unexplored chemical spaces offer the most promising leads. By

predicting and filtering 10+ million purchasable chemicals according to their desirability for human use,

we further illustrate an essential role for computational approaches (e.g., machine learning) in future

repellency studies, accelerating the discovery of safe, ecologically friendly, and effective repellent

chemicals.

Material and Methods

Behavior testing

Repellency Assay in Drosophila

Repellency was tested using a *Drosophila melanogaster* 2-choice trap assay. A chamber was built from

DRAM vials, with four air holes made in the plastic lid and the bottom of chamber is coated with 5 ml of

1% agarose to maintain moisture. Traps are made from 1.5 ml Natural Microcentrifuge tubes with

bottom 3 mm cut to form an opening and a 1000 ul blue graduated pipette tip catalogue number 1111-

2023. To create a plastic funnel, a pipette tip was cut on both ends; cut to 1.0 cm from tip end to expand

the opening of the funnel and then cut on the other end of plastic to create a final height of 2.5 cm for

funnel. A BioRad PCR 0.2 ml Tube cap was cut from the strip and was inserted into the lid of

microcentrifuge tube. Test odor (35 ul) was placed in the PCR tube cap and lure (90 ul) was placed in

outer ring between the PCR cap and the microcentrifuge cap. In the dark, 4–7-day old, starved flies (10

males, 10 females) were released per trap and followed for 2 days. Counts were taken manually, and the

Preference Index (PI) was computed using the formula below.

$$Preference\ Index\ (PI) = \frac{(\#\ flies\ in\ treated\ trap) - (\#\ flies\ in\ control\ trap)}{(\#\ files\ in\ test\ trap) + (\#\ flies\ in\ control\ trap)}$$

Repellency Assay in Mosquitoes

A non-contact heat attraction assay was performed by simultaneously offering mosquitoes a

test chemical on a filter paper coupled with an attractive heat source. Heat for the assay was provided

by a Hot hands® Hand warmers HH2 (Heat Max, Dalton, GA) which was activated by shaking in gloved

hands. Temperature (~37^{o}C) was verified using IR thermometers. The hand warmer was fitted snugly

into a 100 mm x 15 mm petri dish base and covered with 15 cm x 15 cm piece of polyester net secured

round the petri dish by a pair or 8-inch plastic cable ties (Gardner Bender, Milwaukee, WI). The petri

dishe with hand warmer was covered with a larger 150 mm X 15 mm petri dish with a 10 cm x 7.5 cm

rectangular window cut out. This netted window aligned with the hand warmers exposing a heated

surface.

A polyester net (measuring 9 cm x 8 cm) was treated with 500 µl of either acetone (solvent) or

3% (v/v or w/v) test chemical and allowed to air dry for 30 minutes to evaporate solvent. The test

compound ($\sim 0.2\mu g/cm^2$) treated net was secured by stretching between a frame made using two 10 cm

x 7.5 cm flexible magnetic frames with 7.5 cm x 6.2 cm cut out windows. This dose is 40% lower than the

WHO, WHOPES "Guidelines for Efficacy Testing of Mosquito Repellents for Human Skin" recommended

dose for the positive control stimulus. Three more flexible magnets were added on top as spacers from

the heat source. The net-magnet assembly was placed directly over the 10 cm x 7.5 cm cut out openings

of the larger 150 mm X 15 mm petri dishes holding the heat pads. The stimulus was placed on the top

net of the frames. Temperature was monitored by a laser Infra-red thermometer and assay started at

37^oC. A puff of 5% CO_2 was blown into the cage to behaviorally activate the mosquitoes and initiate the

assay. A webcam timelapse is used to record mosquitoes landing on the net for automated counting

offline using ImageJ. A custom Macro is used in ImageJ to count the mosquitoes in the selected area (the

net) across the timepoints. The counts from each frame are automatically tabulated.

Macro:

run("8-bit");

setAutoThreshold("Default");

//run("Threshold...");

setThreshold(0, 80);

run("Analyze Particles...", "size=2-70 show=[Overlay Outlines] clear summarize stack");

```
328        dir = getDirectory("image");

329         name = getTitle;

330         index = lastIndexOf(name, ".");

331         if (index!=-1) name = substring(name, 0, index);

332         name = name + ".csv";

333         selectWindow("Summary of "+getTitle);

334         saveAs("Measurements", dir+name);
```

Since this is a 1-choice assay, the solvent alone controls are done before and after the test material with

at least 20 minutes between the trials. Any significant changes in behavior between the control solvent

should indicate if mosquito behavior is acceptable.

Repellency index = 100 x [1 - (mean cumulative number of mosquitoes on the window of treatment for 5

seconds at time points 2,3,4,5 min/ mean cumulative number of mosquitoes that remained on window

of solvent treatment for 5 seconds at time points 2,3,4,5 min)].

Each time point has N=5 trials, 40 mosquitoes per trial, Except for EA, where N=4.

In vitro assay of Ca^{2+} mobilization

All mammalian cell lines were maintained at 37 C, 5% CO_2, and 99 percent relative humidity. S2 Cells

were maintained at 26 C and 99% relative humidity. HEK293, CHOk1 and NG108 were grown in DMEM

media (Gibco), supplemented with 10% FBS (Gibco), and antibiotic-antimycotic supplement (Gibco). S2

cells were cultured in Shields and Sang M3 Insect Medium (Sigma), supplemented with Insect Medium

Supplement (Sigma). Cells were transfected with the luminescent calcium reporter aequorin 24 hours

before experiment with X-tremeGENE 9 transfection reagent, in the case of HEK293 and CHOK1, and X-

tremeGENE HP, in the case of NG108 and S2 cells. Transfected cells were harvested with enzyme-free

cell dissociation buffer (Gibco), centrifuged, and suspended in assay media (DMEM with 0.1% BSA for

mammalian cells, unsupplemented SFX-100 insect medium for S2). The aequorin cofactor

coelenterazine-F (Nanolight) was added (5 µM final concentration) to the cell suspension, in the dark,

with gentle stirring for 2 to 3 hours before assay. DEET emulsion was prepared by adding a percentage

of DEET indicated, vol/vol to assay media, vortexed, and bath-sonicated for at least 15 min, but up to

one hour. Cells (100 µl, at approx. 50,000 cells) were injected into a 96 well plate containing 100 DEET µl

emulsion, and resulting luminescence recorded on a BMG Labtech Lumistar Omega plate reading

luminometer. Thapsigargin was added to the cell suspension were indicated to a final concentration of 1

µM, 10 minutes before assay.

Chemical Informatics

Machine learning predictions of repellency

Physicochemical features for figures 1-2 were selected using Sequential Forward Selection (SFS)

(Haddad et al., 2008) from 3,224 using Dragon software (Talete). This algorithm is an iterative search

over the chemical feature pool that seeks to maximize the correlation between chemical feature values

and, here, the % repellency. These chemical features were subsequently used to fit a support vector

machine (SVM) with a radial basis function kernel (RBF). The implementation of the support vector

machine algorithm is from the e1071 R package. We performed 20 independent 5-fold cross-validations

followed by a receiver operating characteristics (ROC) analysis to analyze the performance of the

computational repellency prediction method. The trained SVM then ranked both a library obtained from

eMolecules (http://www.emolecules.com) (>400,000) and natural odor libraries (Supplementary

Material) (>3,000) by repellency using the optimized descriptor set to identify the best candidate

repellent compounds. The EPI Suite (http://www.epa.gov/oppt/exposure/pubs/episuite.htm) was used

to calculate predicted LogP and Vapour Pressure values.

For Figures 3, 5, S1-S2, we updated the methods and algorithms to further optimize the pipeline.

Physicochemical features were selected using AlvaDesc software, which offers > 5000 chemical features.

Optimal chemical features were then selected using cross-validation and recursive feature elimination

algorithm (CV-RFE). The RFE CV algorithm involves iterative partitioning of the data and the selection of

feature subsets (n_1, n_2, …, n_{i+1}; where "n" is the size of the feature subset and "i" is an arbitrary indexer).

Given a data partition and n features, an arbitrary ML algorithm is fit, importance is assigned to the

feature set and the model of n features predicts a test partition, that is, data excluded from training,

differentiating the CV-RFE from RFE alone. In this study, the ML algorithm was a support vector machine

(SVM) or a random forest. Over 100s of iterations the algorithm converges on the optimal (n) number of

features to maximize predictive success. Additionally, for every round the contribution of each chemical

feature is assessed, giving rise to an aggregate rank; namely, the frequency at which a feature has been

identified as important to the predictive success. Again, distinguishing the standard RFE algorithm from

CV-based adaptation. As for computing importance, the random forest algorithm internally tests the

contribution of the features. It accomplished this by fitting multiple decision trees on bootstrap samples

of the data. Consequently, a percentage of data has been excluded and serves as a test set. By

predicting the test set with and without permuting or shuffling the values of a given feature, the

algorithm can then assign an importance according to the % increase in error that is observed after

permuting the feature.

The support vector machine (SVM) algorithm does not natively perform a comparable

bootstrapping procedure as random forest, and therefore does not generate a test set to assess feature

importance. In this case, an external metric must be supplied to assign the importance. Depending on

the type of fit, classification or regression, this is a pseudo R^2, based on non-linear regression, or the

AUC for the feature and the outcome that is being predicted (e.g., % repellency). For additional details,

refer to the documentation provided with caret package in R.

Machine learning predictions of Ca^{2+} mobilization

Due to the smaller number of observations (datapoints or chemicals) we opted for an ensemble

method that could take advantage of more of the available data. Random forest fits multiple decision

trees on bootstrap samples of the data, thereby reducing the tendency to overfit to one or few

chemicals in the dataset. As compared to the SVM, it has two simple parameters (mtry and ntrees) and

the tuning grid (or number combinations of these parameters that need to be tested to optimize

performance tends to be small). Accordingly, it is an ideal approach when little is known about the

activity being predicted.

To further reduce overfitting, we identified a small set of physicochemical features on different

subsets of the data and then randomly sampled from this pool. By doing so, the features are slightly

underfit, which is the better approach, again, when less is known about the activity being predicted.

Additionally, we used two different methods and metrics to independently verify that

physicochemical features were indeed predictive of the activity. First, a regression-based approached

predicting the raw assay activities. Second, a classification-based approach where the top ~40% of the

activity distribution was converted into an "Active" label, assessing predictive success by ROC analysis.

Finally, we fit classification models that were otherwise identical but trained on shuffled labels. This

ensured that the chance level-performances significantly differed.

Machine learning predictions of odor perceptual qualities

Perceptual training data is from a study published in a reference book: ATLAS of Odor Character Profiles

(Dravnieks, 1985). In the study, panellists supplied ratings for ~150 odorants and mixtures across 146

possible descriptors; the ratings are reported as the % usage. The % usage refers to the portion of

panellists that rated an odorant (1-5) using a particular descriptor (1-146), indicating its relevance to the

odorant. It is on the scale 0-100, with the maximum of 100% suggesting every participant found the

descriptor to be relevant.

The top perceptual descriptors were selected by thresholding the predictions based on the

mean value in the training data. For each descriptor in the training data, thresholds were computed as

2x the mean. These thresholds were then applied to the predicted repellents to generate a set of

exemplar descriptors. Subsequently, these exemplars were converted into a descriptor frequency table,

where the frequency is the total number of occurrences of a descriptor. The data and word cloud plot

were generated using the R programming language, with additional functionality provided by the

Wordcloud package in R.

References

Beglari, M., Goudarzi, N., Shahsavani, D., Arab Chamjangali, M., & Mozafari, Z. (2020). Combination of

radial distribution functions as structural descriptors with ligand-receptor interaction information

in the QSAR study of some 4-anilinoquinazoline derivatives as potent EGFR inhibitors. *Structural*

Chemistry, *31*(4), 1481–1491. https://doi.org/10.1007/s11224-020-01505-z

Boyle, S. M., McInally, S., & Ray, A. (2013). Expanding the olfactory code by in silico decoding of odor-

receptor chemical space. *ELife*, *2013*(2). https://doi.org/10.7554/eLife.01120.001

Briassoulis, G. (2001). Toxic encephalopathy associated with use of DEET insect repellents: A case

analysis of its toxicity in children. *Human and Experimental Toxicology*.

https://doi.org/10.1191/096032701676731093

Buck L and Axel, R. (1991). A Novel Multigene Family May Encode Odorant Receptors: A Molecular Basis.

Cell, *65*, 175–187.

Carey, A. F., Wang, G., Su, C. Y., Zwiebel, L. J., & Carlson, J. R. (2010). Odorant reception in the malaria

mosquito Anopheles gambiae. *Nature*. https://doi.org/10.1038/nature08834

Cheng, Z., Zhang, Y., & Fu, W. (2010). QSAR study of carboxylic acid derivatives as HIV-1 Integrase

inhibitors. *European Journal of Medicinal Chemistry*. https://doi.org/10.1016/j.ejmech.2010.05.052

Clem, J. R., Havemann, D. F., & Raebel, M. A. (1993). Insect Repellent (N,N-Diethyl-m-Toluamide)

Cardiovascular Toxicity in an Adult. *Annals of Pharmacotherapy*.

https://doi.org/10.1177/106002809302700305

Clyne, P. J., Warr, C. G., & Carlson, J. R. (2000). Candidate taste receptors in Drosophila. *Science*.

https://doi.org/10.1126/science.287.5459.1830

Clyne, P. J., Warr, C. G., Freeman, M. R., Lessing, D., Kim, J., & Carlson, J. R. (1999). A Novel Family of

Divergent   Seven-Transmembrane   Proteins: Candidate   Odorant   Receptors   in

Drosophila. *Neuron*, *22*(2), 327–338. https://doi.org/10.1016/S0896-6273(00)81093-4

Corbel, V., Stankiewicz, M., Pennetier, C., Fournier, D., Stojan, J., Girard, E., Dimitrov, M., Molgó, J.,

Hougard, J. M., & Lapied, B. (2009). Evidence for inhibition of cholinesterases in insect and

mammalian   nervous   systems   by   the   insect   repellent   deet.   *BMC   Biology*.

https://doi.org/10.1186/1741-7007-7-47

Deletre, E., Martin, T., Duménil, C., & Chandre, F. (2019). Insecticide resistance modifies mosquito

response   to   DEET   and   natural   repellents.   *Parasites   &   Vectors*,   *12*(1),   89.

https://doi.org/10.1186/s13071-019-3343-9

Dennis, E. J., Dobosiewicz, M., Jin, X., Duvall, L. B., Hartman, P. S., Bargmann, C. I., & Vosshall, L. B.

(2018). A natural variant and engineered mutation in a GPCR promote DEET resistance in C.

elegans. *Nature*. https://doi.org/10.1038/s41586-018-0546-8

Devinyak, O., Havrylyuk, D., & Lesyk, R. (2014). 3D-MoRSE descriptors explained. *Journal of Molecular*

Graphics and Modelling. https://doi.org/10.1016/j.jmgm.2014.10.006

Dravnieks, A. (1985). *Atlas of Odor Character Profiles*. ASTM. https://doi.org/10.1520/DS61-EB

Duneau, D., Sun, H., Revah, J., San Miguel, K., Kunerth, H. D., Caldas, I. V, Messer, P. W., Scott, J. G., &

Buchon, N. (2018). Signatures of Insecticide Selection in the Genome of Drosophila melanogaster.

G3 (Bethesda, Md.), *8*(11), 3469–3480. https://doi.org/10.1534/g3.118.200537

González, M. P., Terán, C., Teijeira, M., & Helguera, A. M. (2006). QSAR Studies Using Radial Distribution

Function for Predicting A1 Adenosine Receptors Agonists. *Bulletin of Mathematical Biology*, *69*(1),

347. https://doi.org/10.1007/s11538-006-9127-3

Grabe, V., & Sachse, S. (2018). Fundamental principles of the olfactory code. In *BioSystems*.

https://doi.org/10.1016/j.biosystems.2017.10.010

Grant, G. G., & Hall, A. C. (2019). Interactions of N, N- diethyl-meta-toluamide (DEET) and Novel Insect

Repellents with Mammalian GABAA and Glycine Receptors. *The FASEB Journal*, *33*(1_supplement),

813.13-813.13. https://doi.org/10.1096/fasebj.2019.33.1_supplement.813.13

Gupta, R. K., & Bhattacharjee, A. (2006). *Discovery and Design of New Arthropod/Insect Repellents by*

Computer-Aided   Molecular   Modeling   (pp.   195   –   228).

https://doi.org/10.1201/9781420006650.ch10

Haddad, R., Khan, R., Takahashi, Y. K., Mori, K., Harel, D., & Sobel, N. (2008). A metric for odorant

comparison. *Nature Methods*. https://doi.org/10.1038/nmeth.1197

Hallem, E. A., & Carlson, J. R. (2006). Coding of Odors by a Receptor Repertoire. *Cell*, *125*(1), 143–160.

https://doi.org/http://dx.doi.org/10.1016/j.cell.2006.01.050

Hemmer, M. C., Steinhauer, V., & Gasteiger, J. (1999). Deriving the 3D structure of organic molecules

from   their   infrared   spectra.   *Vibrational   Spectroscopy*.   https://doi.org/10.1016/s0924-

2031(99)00014-4

Katritzky, A. R., Wang, Z., Slavov, S., Tsikolia, M., Dobchev, D., Akhmedov, N. G., Hall, C. D., Bernier, U. R.,

Clark, G. G., & Linthicum, K. J. (2008). Synthesis and bioassay of improved mosquito repellents

predicted from chemical structure. *Proceedings of the National Academy of Sciences of the United*

States of America. https://doi.org/10.1073/pnas.0800571105

Keller, A., Gerkin, R. C., Guan, Y., Dhurandhar, A., Turu, G., Szalai, B., Mainland, J. D., Ihara, Y., Yu, C. W.,

Wolfinger, R., Vens, C., Schietgat, L., De Grave, K., Norel, R., Stolovitzky, G., Cecchi, G. A., Vosshall,

L. B., & Meyer, P. (2017). Predicting human olfactory perception from chemical features of odor

494  molecules. *Science*, *355*(6327), 820–826. https://doi.org/10.1126/science.aal2014

495 Kleinau, G., Pratzka, J., Nürnberg, D., Grüters, A., Führer-Sakel, D., Krude, H., Köhrle, J., Schöneberg, T., &

496  Biebermann, H. (2011). Differential modulation of beta-adrenergic receptor signaling by trace

497  amine-associated receptor 1 agonists. *PLoS ONE*. https://doi.org/10.1371/journal.pone.0027073

498 Kline, D. L., Bernier, U. R., Posey, K. H., & Barnard, D. R. (2003). Olfactometric Evaluation of Spatial

499  Repellents for Aedes aegypti . *Journal of Medical Entomology*. https://doi.org/10.1603/0022-2585-

500  40.4.463

501 Klocke, J. A., Darlington, M. V., & Balandrin, M. F. (1987). 1,8-Cineole (Eucalyptol), a mosquito feeding

502  and ovipositional repellent from volatile oil of Hemizonia fitchii (Asteraceae). *Journal of Chemical*

503  *Ecology*. https://doi.org/10.1007/BF01012562

504 Kowalewski, J., Huynh, B., & Ray, A. (2021). A systems-wide understanding of the human olfactory

505  percept chemical space. *Chemical Senses*. https://doi.org/10.1093/chemse/bjab007

506 Kowalewski, J., & Ray, A. (2020a). Predicting novel drugs for SARS-CoV-2 using machine learning from a

507  >10 million chemical space. *Heliyon*. https://doi.org/10.1016/j.heliyon.2020.e04639

508 Kowalewski, J., & Ray, A. (2020b). Predicting Human Olfactory Perception from Activities of Odorant

509  Receptors. *IScience*, *23*(8), 101361. https://doi.org/10.1016/j.isci.2020.101361

510 MacWilliam, D., Kowalewski, J., Kumar, A., Pontrello, C., & Ray, A. (2018). Signaling Mode of the Broad-

511  Spectrum Conserved CO_2 Receptor Is One of the Important Determinants of Odor Valence in

512  Drosophila. *Neuron*. https://doi.org/10.1016/j.neuron.2018.01.028

513 Maldonado, A. G., Doucet, J. P., Petitjean, M., & Fan, B. T. (2006). Molecular similarity and diversity in

514  chemoinformatics: From theory to applications. In *Molecular Diversity*.

515  https://doi.org/10.1007/s11030-006-8697-1

Mombaerts, P. (1999). Molecular biology of odorant receptors in vertebrates. *Annual Review of*

Neuroscience, *22*, 487–509. https://doi.org/10.1146/annurev.neuro.22.1.487

Mombaerts, Peter. (2001). THE HUMAN REPERTOIRE OF ODORANT RECEPTOR GENES AND

PSEUDOGENES. *Annual Review of Genomics and Human Genetics*, *2*(1), 493–510.

https://doi.org/10.1146/annurev.genom.2.1.493

Osimitz, T. G., Murphy, J. V., Fell, L. A., & Page, B. (2010). Adverse events associated with the use of

insect repellents containing N,N-diethyl-m-toluamide (DEET). *Regulatory Toxicology and*

Pharmacology. https://doi.org/10.1016/j.yrtph.2009.09.004

Reeder, N. L., Ganz, P. J., Carlson, J. R., & Saunders, C. W. (2009). Isolation of a Deet-Insensitive Mutant

of <I>Drosophila melanogaster</I> (Diptera: Drosophilidae). *Journal of Economic Entomology*.

https://doi.org/10.1603/0022-0493-94.6.1584

Sachse, S., & Beshel, J. (2016). The good, the bad, and the hungry: How the central brain codes odor

valence to facilitate food approach in Drosophila. In *Current Opinion in Neurobiology*.

https://doi.org/10.1016/j.conb.2016.06.012

Schoenig, G. P., Hartnagel, R. E., Osimitz, T. G., & Llanso, S. (1996). Absorption, distribution, metabolism,

and excretion of N,N-diethyl-M-toluamide in the rat. *Drug Metabolism and Disposition*.

Selim, S., Hartnagel, R. E., Osimitz, T. G., Gabriel, K. L., & Schoenig, G. P. (1995). Absorption, metabolism,

and excretion of N,N-diethyl-m-toluamide following dermal application to human volunteers.

Toxicological Sciences. https://doi.org/10.1093/toxsci/25.1.95

Swale, D. R., Sun, B., Tong, F., & Bloomquist, J. R. (2014). Neurotoxicity and mode of action of N, N-

diethyl-Meta-toluamide (DEET). *PLoS ONE*. https://doi.org/10.1371/journal.pone.0103713

Syed, Z., & Leal, W. S. (2008). Mosquitoes smell and avoid the insect repellent DEET. *Proceedings of the*

National   Academy   of   Sciences   of   the   United   States   of   America.

https://doi.org/10.1073/pnas.0805312105

Syed, Z., Pelletier, J., Flounders, E., Chitolina, R. F., & Leal, W. S. (2011). Generic insect repellent detector

from   the   fruit   fly   Drosophila   melanogaster.   *PLoS   ONE*.

https://doi.org/10.1371/journal.pone.0017705

Tavares, E. M., Judge, B. S., & Jones, J. S. (2019). Bug off! Severe toxicity following inhalational exposure

to   N,   N-diethyl-meta-toluamide   (DEET).   *American   Journal   of   Emergency   Medicine*.

https://doi.org/10.1016/j.ajem.2019.04.019

Vassar, R., Ngai, J., & Axel, R. (1993). Spatial segregation of odorant receptor expression in the

mammalian olfactory epithelium. *Cell*, *74*(2), 309–318. https://doi.org/0092-8674(93)90422-M [pii]

    FIGURE LEGENDS

Figure 1. A chemical informatics pipeline to screen for pleasant smelling insect repellents

A, Overview of the cheminformatics pipeline used to identify novel DEET-like ligands from a larger

chemical space. Summary table of top 10 physicochemical features that optimally corelate with

repellency. Receiver-operating-characteristic curve (ROC), which indicates the success in predicting test

repellents. The performance is the average over 100 80/20 splits, where 80% of the data is for training

and 20% for testing. The mean area under the curve (AUC) is provided. Representative structures from

the top 150 predicted repellent compounds.

Figure 2. Models successfully predict odor perceptual qualities of repellents in humans

A, Validation of models predicting the odor perceptual qualities (146 perceptual descriptors) of a set of

compounds that were screened for repellency in Figure 3A. The performance metric for the validation is

the area under the ROC curve, which assesses the rate of correctly predicted perceptual descriptors

relative the rate of incorrectly predicted descriptors. Dashed line is the average Area Under the Curve

(AUC). **B,** Sample ROC curves for select compounds as well as the ROC curve across all compounds (e.g.,

shown as separate curves in the ROC plot as well as those not shown but whose identities appear in

Figure 2A); the respective AUC values are reported inside the plot. **C,** The top 25 perceptual descriptors

for the predicted repellents that underwent experimental validation are displayed as a word cloud. The

font size of the descriptor is scaled relative to frequency.

Figure 3. Experimental validation of predicted insect repellents show a high success rate several of

which are pleasant smelling

A, Representative still photograph of landing assay of female *Aedes aegypti* mosquitoes on filter paper

covered heat block over 5 mins (overlay of all frames), and **B,** mean repellency index for each of the

indicated compounds tested at ~0.2μg/cm^2. The repellency coefficient = number mosquitoes attracted

to solvent treated control netting, minus number of mosquitoes on treated netting compared to solvent,

with 37C heated rectangle as an attractant. Mosquito numbers are counted for the 5 min duration, over

multiple frames, as described in the methods. N=2-4 trials, ~20 mosquitoes/trial. For DEET, N=23 trials.

C, Mean Preference index of *Drosophila* adults to repellents at three different concentrations dissolved

in acetone in a Two-choice trap assay baited with 10% Apple Cider Vinegar measured after 48 hrs. N = 7-

10 trials each treatment at 48 hrs, 10 flies/trial, error bars = s.e.m.

Figure 4. Some, but not all insect repellents increase Ca^{2+} mobilization in a human cell line

DEET induces a non-specific calcium repose *in vitro*. **A.** Dose response curves for three mammalian cell

lines (HEK293. CHOK1, NG108) and one insect cell line (S2). EC50 and Hill coefficients for each cell line

are given below. **B.** Representative DEET induced calcium kinetics in HEK293 cells. **C.** HEK293 cells

treated with thapsigargin 10 min before assay to deplete intracellular calcium stores no longer respond

to DEET, suggesting DEET response is dependent on intracellular calcium. Error bars are standard

deviation.**D. F,** Luminescence of the Ca^{2+} indicator at two different concentrations for predicted and

previously known repellent chemicals, scaled relative to DEET.

Figure 5. Screen of 10+ million chemicals predicts several insect repellents with different smells

A, Tabulated predicted % repellency (relative to DEET) from a library of 10+ million chemicals, filtered to

the top values. The best matching known repellent is displayed alongside the Euclidean distance, the

predicted LD50 and Ca^{2+} mobilization. **B,** heatmap organizing top predictions according to estimated

perceptual qualities.

Supplemental Figure S1

A, Overview of the approach to generate and validate the machine learning models. Here, a model-

averaged prediction is made, where each model has access to different physicochemical features and is

trained on different combinations of training set chemicals. **B,** The R^2 values over the 100 train/test

splits, averaged into 10 bins. **C,** The average classification success is reported over 100 train/test splits,

assessed by the area under the ROC curve. Active labels (positive cases) were assigned as chemical

scoring in the top ~40% of Ca^{2+} values. This is compared to shuffling the labels before training, reported

as "Shuffle."

Descriptor Symbol	Descriptor Definition
B05[C-N]	Presence/absence of C-N at topological distance 5
piID	conventional bond order ID number
N-072	RCO-N< />N-X=X
nRCONR2	number of tertiary amides (aliphatic)
RDF035v	Radial Distribution Function - 035 / weighted by van der Waals volume
EEig08r	Eigenvalue 08 from edge adj. matrix weighted by resonance integrals
D-Dr06	distance/detour ring index of order 6
Mor10p	3D-MoRSE - signal 10 / weighted by atomic polarizables
BAC	Balaban centric index
EEig14d	Eigenvalue 14 from edge adj. matrix weighted by dipole movements

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5