Evaluating HIV-1 Infectivity and Virion Maturation Across Varied Producer Cells with a Novel FRET-Based Detection and Quantification Assay

Aidan McGraw[1†], Grace Hillmer[1†], Jeongpill Choi[1†], Kedhar Narayan[1†], Dacia Marquez[1], Hasset Tibebe[1], Taisuke Izumi[1, 2*]

[1]Department of Biology, College of Arts and Sciences, American University, Washington D.C., USA

[2] District of Columbia Center for AIDS Research, Washington D.C., USA

[†]These authors contributed equally to this work

*Author to whom correspondence should be addressed

Abstract

The maturation of HIV-1 virions is a crucial process in viral replication. Although T cells are a primary source of virus production, much of our understanding of virion maturation comes from studies using the HEK293T human embryonic kidney cell line. Notably, there is a lack of comparative analyses between T cells and HEK293T cells in terms of virion maturation efficiency in existing literature. We previously developed an advanced virion visualization system based on the FRET principle, enabling the effective distinction between immature and mature virions via fluorescence microscopy. In this study, we utilized pseudotyped, single-round infectious viruses tagged with FRET labels (HIV-1 Gag-iFRETΔEnv) derived from Jurkat (a human T lymphocyte cell line) and HEK293T cells to evaluate their virion maturation rates. HEK293T-derived virions demonstrated a maturity rate of 81.79%, consistent with other studies and our previous findings. However, virions originating from Jurkat cells demonstrated a significantly reduced maturation rate of 68.67% ($p < 0.0001$). Correspondingly, viruses produced from Jurkat cells exhibited significantly reduced infectivity compared to those derived from HEK293T cells, with the relative infectivity measured at 65.3%. This finding is consistent with the observed relative maturation rate of viruses produced by Jurkat cells. These findings suggest that initiation of virion maturation directly correlates with viral infectivity. Our observation highlights the dynamic nature of virus-host interactions and their implications for virion production and infectivity.

Keywords:

36   Human Immunodeficiency Virus type I (HIV-1), Forster Resonance Energy Transfer

37   (FRET), Virion Maturation, Comparative Viral Analysis, Fluorescent Microscopic

38   Imaging,

Introduction

Human Immunodeficiency Virus-1 (HIV-1) continues to pose a significant global health challenge. The advancement of combination antivirals has dramatically improved the management of HIV-1, targeting various stages of the virus lifecycle, including entry, reverse transcription, integration, and protease activity [1-6]. Recently, the development of capsid inhibitors has emerged as a promising strategy for long-acting antiviral treatments against HIV-1 [7-12]. These novel inhibitors, targeting critical steps in both the early and late stages of the virus lifecycle, notably disrupt the formation of infectious particles. This innovation underscores the need for a comprehensive understanding of the virion maturation process, a key phase in the HIV-1 life cycle where non-infectious particles are transformed into infectious virions. Our previous research developed a unique tool for detecting virus maturation utilizing a fluorescence resonance energy transfer (FRET)-based technique [13]. This approach enables the semi-automatic evaluation of virion maturation through fluorescent microscopy, effectively minimizing human bias. It offers crucial insights into the structural changes that are vital for virus infectivity. Building upon this, our current study explores virion maturation in different cellular environments. Although T cells are recognized as the primary source for HIV-1 replication [14-20], most existing research assessing viral maturation predominantly utilizes the HEK293T human embryonic kidney cell line [21-24]. Utilizing the FRET-based system, we generated infectious FRET-labeled viruses in both HEK293T and Jurkat cells, an immortal human T-cell leukemia cell line. Surprisingly, we observed that viruses produced by Jurkat cells exhibited a significantly lower maturation efficiency (approximately 69%) compared to those from HEK293T cells (approximately 82%). In alignment with the observed differences in maturation rates, the infectivity of

viruses from Jurkat cells was approximately half that observed in viruses from

HEK293T cells. Given that CD4+ T cells are the principal producers of HIV-1, these

cells have evolved multiple antiretroviral mechanisms. For example, the HIV-1 Viral

Infectivity Factor (Vif) is a crucial accessory molecule for HIV-1 replication in vivo.

However, it is not necessary for HEK293T cells, as these kidney-derived cells do not

express the potent antiviral host factor APOBEC3G [13, 25, 26]. It is highly plausible

that kidney cells, having not been exposed to lentiviruses throughout evolutionary

history, exhibit greater susceptibility to these viruses compared to T cells, which have

been concurrently evolving to resist viral infections. This study is at the forefront of

comparing virion maturation and infectivity among cell lines from different origins,

highlighting a novel approach in this field of HIV-1 research.

Results

Virion Maturation Efficiency in Different Producer Cells

In this study, we quantified the proportion of mature and immature virions in FRET-

labeled viruses produced by HEK293T (kidney) and Jurkat (T-cell leukemia) cell

lines. In each sample, we captured a total of 21 images, from which we created

binary images based on the emitted and excited images of YFP (Fig. 1B). We then

calculated the FRET ratio for each virion by dividing the intensity of YFP (the FRET

acceptor) by the total CFP intensity, which serves as the FRET donor. After

calculating the FRET ratio for each virion, based on the extracted signal intensities of

the FRET donor and acceptor, we applied kernel density estimation to the

histograms of FRET efficiencies. The area where the HIV-1 Gag-iFRETΔEnv curve

overlapped with the HIV-1 Gag-iFRETΔPRΔEnv curve was identified as the

proportion of immature virions, as per our established methodology (Fig. 1C) [13]. In

our analysis, we examined 14,021 and 22,779 particles, as well as 20,457 and

23,181 particles of HIV-1 Gag-iFRETΔEnv and HIV-1 Gag-iFRETΔPRΔEnv labeled

virions produced by HEK293T and Jurkat cells, respectively, across three

independent experiments. The mature virion proportion in the HIV-1 Gag-iFRETΔEnv

population from HEK293T cells was 81.79% ± 0.06 (Fig. 1D), consistent with our

previous findings and electron microscopic counts reported in other studies [13, 21-

24]. In contrast, Jurkat cells produced mature virions at a lower rate of 68.67% ±

0.04 (Fig. 1D), which is significantly different from that of HEK293T cells (p-value <

0.0001). Furthermore, we quantified the Mean Fluorescent Intensity (MFI) of the YFP

signal in both mature and immature virions. The results confirmed that the average

incorporation of fluorescent proteins, indicated by the MFI of YFP (Fig. 1E), was

similar in both mature and immature virions from HEK293T and Jurkat cells

(26,691.31 ± 2,733.73 vs. 23,309.51 ± 766.93 in mature virions and 24,619.37 ±

2,857.71 vs. 24,536.10 ± 2,046.71 in immature virions, respectively, and 24,619.37 ±

2,857.71 vs. 24,536.10 ± 2,036.70 in ΔPR). These findings indicate that the number

of incorporated fluorescent proteins is comparable, ensuring that the FRET energy

transfer efficiency is neither under nor overestimated in virions produced by either

HEK293T or Jurkat cells.

Virus Infectivity in Different Producer Cells

To further examine the characteristic differences between viruses produced by

HEK293T and Jurkat cells, we performed a single-round infection assay using TZM-

bl cells to assess the infectivity of FRET-labeled viruses originating from these cell

lines. The infectivity of viruses from Jurkat cells was significantly lower compared to

those from HEK293T cells, measured at 65.3% ± 5.6 (Fig. 2). This discrepancy is

consistent with the observed differences in the maturation rates of viruses produced

by Jurkat cells. (Fig. 1).

Discussion

The findings of this study provide groundbreaking insights into the maturation

process of HIV-1 virions and their impact on viral infectivity, challenging several

established concepts in the field. Our utilization of an advanced FRET-based

visualization system has allowed for a detailed comparative analysis of virion

maturation in two distinct cellular environments: HEK293T and Jurkat cell lines. One

of the most notable findings of our study is that virus maturation and subsequent

infectivity in T cell lines, which are derived from the natural targets for HIV-1, are

lower compared to kidney-derived cell lines. HEK293T cells exhibited a reduced rate

of immature virions (18.21%), whereas virions derived from Jurkat cells displayed a

higher rate of immaturity (31.33%), correlating with its respective infectivity (65.3% ±

5.6 for Jurkat cell-derived virions compared to the infectivity observed in virions from

HEK293T cells). Recent studies have provided deeper insights into the role of

inositol hexakisphosphate (IP6) in HIV-1 capsid assembly, which is a critical factor in

virus infectivity [27]. These studies reveal that the immature Gag lattice of HIV-1

virions enriches IP6, facilitating capsid maturation. This process is characterized by

proteolysis of the Gag polyprotein by the viral protease, which induces vital

conformational changes in the capsid protein (CA). These changes led to the

formation of a conical, cone-shaped mature capsid consisting of over 1,000 CA

copies and forming approximately 200 hexamers and 12 pentamers, collectively

known as capsomers. Sowd et al. showed that IP6-depleted T-cell lines produce

virion particles with incompletely cleaved Gag proteins [28]. This study indicated that

virions from IP6-deficient cells exhibited lower quantities of mature MA and CA

cleavage products and higher proportions of MA-CA-SP1/MA-CA cleavage

intermediates than virions from wild-type cells. In the context of our FRET-based

assay, which specifically detects cleavage of the Gag polyprotein between MA-CA,

our FRET detection system can potentially monitor the IP6-dependent efficiency of

Gag polyprotein cleavage. Furthermore, they characterized virus particle morphology

using electron microscopy and calculated that mature particles in another T-cell

leukemia cell line, MT-4 cells, comprised 62% of the total virions, which aligns with

the results from our FRET detection system in Jurkat cells (Fig. 1C [II], and 1D).

Given that the majority of inositol synthesis in humans occurs in kidney tissue [29], it

is reasonable to infer that HEK293T cells would exhibit abundant IP6 expression.

This is likely to contribute to enhanced virion maturation efficiency. It has also been

suggested that IP6 plays a crucial role in facilitating the assembly of the immature

HIV-1 Gag lattice [30]. Consequently, the elevated expression of IP6 in HEK293T

cells might enhance the assembly of Gag precursors at the budding site prior to

virion release, potentially initiating the trimerization of the MA domain. This, in turn,

could enhance the efficient incorporation of Env proteins into progeny virions [31, 32]

Another study employing a similar concept of FRET-based assay with different FRET

pair proteins revealed that the activation of HIV-1 viral protease occurs during

assembly and budding before the release of particles in HEK293T cells [33]. The

early maturation observed in HEK293T cells could lead to an increase in progeny

virion maturation. This effect likely accounts for the elevated maturation ratio

observed in HEK293T cells in our study (Fig. 1). This early maturation could also

result in enhanced viral infectivity compared to virions originating from Jurkat cells

(Fig. 2)

Several accessory molecules, such as Vif, are not required to produce infectious

viruses from HEK293T cells [13, 25, 26]. This is due to the absence of antiviral host

factors in HEK293T cells that would normally be counteracted by viral accessory

proteins. This discrepancy may hint at the heightened susceptibility of kidney cells to

retrovirus infection, potentially attributable to their lack of prior exposure to these viral

components. Thus, the environment in HEK293T cells is optimized not only by the

absence of antiviral host protein expression but also by the enrichment of host

cofactors, which collectively contribute to increased virus infectivity.

Our research addresses a critical gap in the existing literature by presenting a

comparative analysis of the efficiency of Gag polyprotein cleavage in virions

produced by kidney cells (HEK293T) and T cells (Jurkat). This comparison is vital,

considering T cells are the primary source of HIV-1 replication *in vivo*. The observed

differences in maturation efficiencies and infectivity rates between HEK293T and

Jurkat cells highlight the importance of considering the cellular context in HIV-1

biology and pathogenesis studies. Furthermore, our findings have significant

implications for the evaluation of antiviral effects, particularly in the context of long-

acting treatments with maturation and capsid inhibitors. These inhibitors target

various stages of virus infection, including viral budding and virion maturation [31,

34, 35]. The substantial disparities in viral maturation rates between T-cells and

HEK293T cells, the latter of which is a standard cellular resource for *in-vitro* virus

production, suggest that relying solely on HEK293T cells could lead to misleading

assessments of antiviral efficacy. Therefore, our study indicates the necessity of

including T-cell-based assessments to more accurately determine the antiviral

efficacy of maturation and capsid inhibitors in virus maturation.

In conclusion, the results of our study not only enhance our understanding of HIV-1

pathogenesis but also emphasize the complexity of the virus life cycle. The unique

insights gained into the relationship between virion maturation and infectivity open

new pathways for future research, potentially leading to novel therapeutic

approaches in the fight against HIV-1.

Materials and Methods

Plasmid DNA Construction

The plasmid DNA utilized for HIV-1 Gag-iFRET labeling virus production incorporates

an efficient intramolecular FRET pair, ECFPΔC11 and cp173Venus [36]. These

genes are positioned between the MA and CA regions of the HIV-1 Gag protein

based on the HIV-1 Gag-iGFP construct [37]. The FRET pair genes, along with the

junction with the MA and CA regions, are flanked by HIV-1 protease cleavage sites

(SQNYPIVQ). This design enables the HIV-1 Protease to specifically cleave at these

sites, resulting in a modification of the FRET signal during the maturation of the

virion. Additional details are provided in our previous publication [13].

Virus Production and Cell Culture

HIV-1 Gag-iFRET plasmid DNA was transfected into HEK293T cells using a PEI

transfection method (Polysciences) and into Jurkat cells using Lipofectamine LTX

transfection reagent (Invitrogen). HEK293T and TZM-bl cells were cultured in

Dulbecco's Modified Eagle's Medium (Cytiva) supplemented with 10% Fetal Bovine

Serum (Gibco), 1% Penicillin-Streptomycin-Glutamine (Invitrogen), and 1%

GlutaMax (Gibco)(D10) at 37°C in a 5% CO_2 environment. Suspended Jurkat cells

were cultured in RPMI 1640 media (Cytiva) with the same supplements (R10). To

produce FRET-labelled virions, HEK293T cells (7.0×10^6 cells/10 cm dish) and

Jurkat cells (10×10^6 cells/10 cm dish) were co-transfected with pHIV-1 Gag-

iFRETΔEnv or iFRETΔPRΔEnv, along with the parental plasmids pNL4-3ΔEnv or

pNL43ΔPRΔEnv, and the pSVIII-92HT593.1 dual-tropic HIV-1 Env expression

plasmid, kindly provided by Dr. Viviana Simon at Mount Sinai, using a transfection

ratio of 1:10:2.5. The culture medium was replaced with fresh D10 medium after 3.5

hours for PEI transfection. Virus-containing supernatants were collected 48 hours

post-transfection. The supernatants were then filtered through a 0.45 μm sterile

polyvinylidene difluoride (PVDF, Millipore) membrane and concentrated up to 10-fold

using Lenti-X Concentrator (TaKaRa). The virus pellet was resuspended in 100 μL of

Phosphate Buffered Saline (PBS) without phenol red (Cytiva).

FRET-Based Virion Visualization

To visualize HIV-1 Gag-iFRETΔEnv/iFRETΔPRΔEnv labeled virions, dilute the

concentrated virus supernatant 800-fold in 0.22 μm PVDF-filtered Hank's Balanced

Salt Solution (HBSS) without Phenol Red (Cytiva) and load 350 μL onto a glass-

bottomed 8-chamber slide (ibidi). The slide was then incubated overnight at 4°C.

Single-virion images were captured using an A1R MP+ Multiphoton Confocal

Microscope (Nikon). Two sets of 21 images were automatically acquired for each

sample under optimal focus conditions. The first set employed a 457.9 nm laser for

CFP excitation, capturing emissions through 482 nm/35 nm and 540 nm/30 nm filter

cubes for CFP and YFP signals, respectively (FRET images). The second set used a

514.5 nm laser for Venus excitation, with emissions read through a 540 nm/30 nm

filter cube to detect the YFP signal. The maturation status was quantified as FRET

efficiency relative to signals detected in HIV-1 Gag-iFRETΔ1PRΔEnv labeled virions.

Images were captured in RAW ND2 format and converted to TIFF files using Fiji, an

image processing package based on ImageJ2. Binary images, generated based on

the YFP signal, provided XY coordinates for each particle. The centroid of the XY

position for each virion particle was determined, and a circle with a 2-pixel radius

centered on the calculated centroid was drawn to represent the virion map. Using

this virion map, the FRET signal intensity for each virion was extracted from the raw

data, and the FRET ratio (YFP/CFP) was calculated for each particle. Additionally,

the mean fluorescent intensity (MFI) of YFP within the circle of the virion map was

also extracted from the raw data. Histograms displaying the distribution of FRET

ratio values were generated, with 100 bins division, and kernel density estimation

curves were plotted alongside. The proportion of the total kernel density estimation

area overlapping with the HIV-1 Gag-iFRETΔPRΔEnv area was used to determine

the proportion of immature virions. Image data analysis and subsequent calculations

of mature and immature virion proportions were performed using an updated in-

house MATLAB program [13].

Infectivity Assays

Viral titers were determined using the HIV-1 Gag p24 DuoSet ELISA (R&D Systems).

After seeding TZM-bl cells at 1×10^4 in flat-bottom 96-well plates (CELLTREAT),

they were exposed to an equivalent amount of virus (5 ng of HIV-1 p24 total) in the

following day. The cells were incubated at 37°C for 48 hours in a CO_2 incubator.

Subsequently, luciferase activity in the infected cells was measured with the

Luciferase Assay System (Promega) using a Molecular Devices SpectraMax

Microplate Reader.

Statistical Analysis

Data were analyzed using standard statistical methods. The significance of

differences in virion maturation and infectivity between the two cell types was

determined using the Mann-Whitney U test with a *p*-value of less than 0.05, which is

considered statistically significant. The statistical differences in YFP MFI between the

two cell types were calculated using the Wilcoxon matched-pairs signed-rank test.

Acknowledgments

We extend our thanks to Dr. Viviana Simon at Mount Sinai for generously providing

the pSVIII-92HT593.1 plasmid DNA. In addition, Dr. Luca Sardo at ViiV Healthcare

kindly reviewed and provided invaluable comments that enriched this research.

Author's contributions

TI conceptualized and designed the study. Experiments were carried out by AM, JC,

GH, DM, HT, and TI. The development of the in-house MATLAB script for automated

image data analysis was a collaborative effort between KN and TI. Data analysis was

conducted by KN and TI. The manuscript was written by TI. TI provides financial

support for the research. All authors have made substantial contributions to the work

and have given their approval for the final version of the article to be published.

Funding

This study is partially supported by a research start-up fund and the Spring 2024

CAS Faculty Mellon Fund from American University to TI. Additional funding was

provided by the National Institutes of Health, specifically through a grant from the

National Institute of Allergy and Infectious Diseases (Grant Number: 7R15AI172610-

02 to TI).

Figure Legend

Figure 1. Quantitative Analysis of Mature Virions from HEK293T and Jurkat

Cells.

(A) Schematic representation of the HIV-1 Gag iFRET construct in the Gag region.

ECFPΔC11 and cp173Venus are used for efficient single-molecule CFP and YFP

FRET pairing. (B) Representative FRET-labeled HIV-1 Gag-iFRETΔEnv virion

images from Jurkat cells displayed in a single field. The left panel shows an image

captured through the YFP excitation (515 nm) and emission (540 nm) channels (YFP

channel). The second image shows the binary image, derived from the YFP channel,

identifying virus particle locations as regions of interest (ROI). The third and fourth

images are captured using CFP (FRET donor) excitation (454 nm), with CFP

emission (483 nm, third) and YFP emission (540 nm, fourth) channels. The right

panel shows the FRET efficiency ratio view, computationally constructed from FRET

donor (CFP excitation/CFP emission) and acceptor (CFP excitation/YFP emission)

images. The color spectrum indicates FRET signal strength, with high FRET

appearing in red (yellow arrows) and decreasing FRET shifting toward blue (white

arrows). (C) Kernel Density Estimation Curves of FRET Intensity in HIV-1 Gag-

iFRETΔEnv Virions Produced by HEK293T and Jurkat Cells. The curves depict the

distribution histograms of FRET intensity for virions labeled with HIV-1 Gag-

iFRETΔEnv from (I) HEK293T and (II) Jurkat cells. The x-axis represents the range

of FRET intensity. The y-axis shows the adjusted density of curves for both HIV-1

Gag-iFRETΔEnv and HIV-1 Gag-iFRETΔPRΔEnv virions. The area under the curve

for HIV-1 Gag-iFRETΔEnv overlapping with that of HIV-1 Gag-iFRETΔPRΔEnv is

calculated and considered indicative of the proportion of immature virions within the

total population of HIV-1 Gag-iFRETΔEnv virions. (D) Scatter Plot of Mature Virion

Quantification displays the median percentage of mature virions across 21 images

derived from three independent experiments, culminating in a total of 63 data points.

Statistical significance was assessed using the Mann-Whitney U test. (E)

Comparison of Mean Fluorescent Intensity (MFI) of YFP Signals in Mature and

Immature HIV-1 Gag-iFRETΔEnv labeling Virions. The levels of fluorescent protein

incorporation, as indicated by the YFP signal, were not significantly different between

mature and immature virions in either the wild-type or ΔPR viruses. The error bars

represent the standard error from three independent experiments. Statistical

significance was assessed using the Wilcoxon matched-pairs signed rank test.

Figure 2. Comparing Virus Infectivity Between HEK293T and Jurkat Cells.

Single-round infectivity assays using TZM-bl cells were performed to evaluate the

differences in virus infectivity produced by HEK293T and Jurkat cells. The bar plot

illustrates the relative infectivity of the FRET-labeled virus produced by Jurkat cells in

comparison to its infectivity when produced by HEK293T cells. Error bars represent

the standard deviation from four independent experiments. Statistical significance

was determined using a Mann-Whitney U test.

References

References

1.   Ghosh, A. K., Four decades of continuing innovations in the development of
antiretroviral therapy for HIV/AIDS: Progress to date and future challenges. *Glob*
Health Med **2023,** 5, (4), 194-198.
2.   Landovitz, R. J.; Scott, H.; Deeks, S. G., Prevention, treatment and cure of HIV
infection. *Nat Rev Microbiol* **2023,** 21, (10), 657-670.
3.   Arts, E. J.; Hazuda, D. J., HIV-1 antiretroviral drug therapy. *Cold Spring Harb*
Perspect Med **2012,** 2, (4), a007161.
4.   Magden, J.; Kaariainen, L.; Ahola, T., Inhibitors of virus replication: recent
developments and prospects. *Appl Microbiol Biotechnol* **2005,** 66, (6), 612-21.
5.   Adamson, C. S.; Freed, E. O., Novel approaches to inhibiting HIV-1 replication.
Antiviral Res **2010,** 85, (1), 119-41.
6.   Wiebe, L. I.; Knaus, E. E., Concepts for the design of anti-HIV nucleoside prodrugs
for treating cephalic HIV infection. *Adv Drug Deliv Rev* **1999,** 39, (1-3), 63-80.
7.   Thenin-Houssier, S.; Valente, S. T., HIV-1 Capsid Inhibitors as Antiretroviral Agents.
Curr HIV Res **2016,** 14, (3), 270-82.
8.   Yant, S. R.; Mulato, A.; Hansen, D.; Tse, W. C.; Niedziela-Majka, A.; Zhang, J. R.;
Stepan, G. J.; Jin, D.; Wong, M. H.; Perreira, J. M.; Singer, E.; Papalia, G. A.; Hu, E.
Y.; Zheng, J.; Lu, B.; Schroeder, S. D.; Chou, K.; Ahmadyar, S.; Liclican, A.; Yu, H.;
Novikov, N.; Paoli, E.; Gonik, D.; Ram, R. R.; Hung, M.; McDougall, W. M.; Brass,
A. L.; Sundquist, W. I.; Cihlar, T.; Link, J. O., A highly potent long-acting small-
molecule HIV-1 capsid inhibitor with efficacy in a humanized mouse model. *Nat Med*
2019, 25, (9), 1377-1384.
9.   Singh, K.; Gallazzi, F.; Hill, K. J.; Burke, D. H.; Lange, M. J.; Quinn, T. P.; Neogi, U.;
Sonnerborg, A., GS-CA Compounds: First-In-Class HIV-1 Capsid Inhibitors Covering
Multiple Grounds. *Front Microbiol* **2019,** 10, 1227.
10.   Sun, L.; Zhang, X.; Xu, S.; Huang, T.; Song, S.; Cherukupalli, S.; Zhan, P.; Liu, X.,
An insight on medicinal aspects of novel HIV-1 capsid protein inhibitors. *Eur J Med*
Chem **2021,** 217, 113380.
11.   Flexner, C.; Owen, A.; Siccardi, M.; Swindells, S., Long-acting drugs and
formulations for the treatment and prevention of HIV infection. *Int J Antimicrob*
Agents **2021,** 57, (1), 106220.
12.   Gillis, E. P.; Parcella, K.; Bowsher, M.; Cook, J. H.; Iwuagwu, C.; Naidu, B. N.; Patel,
M.; Peese, K.; Huang, H.; Valera, L.; Wang, C.; Kieltyka, K.; Parker, D. D.;
Simmermacher, J.; Arnoult, E.; Nolte, R. T.; Wang, L.; Bender, J. A.; Frennesson, D.
B.; Saulnier, M.; Wang, A. X.; Meanwell, N. A.; Belema, M.; Hanumegowda, U.;
Jenkins, S.; Krystal, M.; Kadow, J. F.; Cockett, M.; Fridell, R., Potent Long-Acting
Inhibitors Targeting the HIV-1 Capsid Based on a Versatile Quinazolin-4-one
Scaffold. *J Med Chem* **2023,** 66, (3), 1941-1954.
13.   Sarca, A. D.; Sardo, L.; Fukuda, H.; Matsui, H.; Shirakawa, K.; Horikawa, K.;
Takaori-Kondo, A.; Izumi, T., FRET-Based Detection and Quantification of HIV-1
Virion Maturation. *Front Microbiol* **2021,** 12, 647452.
14.   Chen, B., Molecular Mechanism of HIV-1 Entry. *Trends Microbiol* **2019,** 27, (10),
878-891.
15.   Pan, X.; Baldauf, H. M.; Keppler, O. T.; Fackler, O. T., Restrictions to HIV-1
replication in resting CD4+ T lymphocytes. *Cell Res* **2013,** 23, (7), 876-85.
16.   Banga, R.; Procopio, F. A.; Noto, A.; Pollakis, G.; Cavassini, M.; Ohmiti, K.;
Corpataux, J. M.; de Leval, L.; Pantaleo, G.; Perreau, M., PD-1(+) and follicular

383   helper T cells are responsible for persistent HIV-1 transcription in treated aviremic
384   individuals. *Nat Med* **2016,** 22, (7), 754-61.

17. Berg, R. K.; Rahbek, S. H.; Kofod-Olsen, E.; Holm, C. K.; Melchjorsen, J.; Jensen, D. G.; Hansen, A. L.; Jorgensen, L. B.; Ostergaard, L.; Tolstrup, M.; Larsen, C. S.; Paludan, S. R.; Jakobsen, M. R.; Mogensen, T. H., T cells detect intracellular DNA but fail to induce type I IFN responses: implications for restriction of HIV replication. *PLoS One* **2014,** 9, (1), e84513.

18. Jones, R. B.; Ndhlovu, L. C.; Barbour, J. D.; Sheth, P. M.; Jha, A. R.; Long, B. R.; Wong, J. C.; Satkunarajah, M.; Schweneker, M.; Chapman, J. M.; Gyenes, G.; Vali, B.; Hyrcza, M. D.; Yue, F. Y.; Kovacs, C.; Sassi, A.; Loutfy, M.; Halpenny, R.; Persad, D.; Spotts, G.; Hecht, F. M.; Chun, T. W.; McCune, J. M.; Kaul, R.; Rini, J. M.; Nixon, D. F.; Ostrowski, M. A., Tim-3 expression defines a novel population of dysfunctional T cells with highly elevated frequencies in progressive HIV-1 infection. *J Exp Med* **2008,** 205, (12), 2763-79.

19. Zhang, Z.; Schuler, T.; Zupancic, M.; Wietgrefe, S.; Staskus, K. A.; Reimann, K. A.; Reinhart, T. A.; Rogan, M.; Cavert, W.; Miller, C. J.; Veazey, R. S.; Notermans, D.; Little, S.; Danner, S. A.; Richman, D. D.; Havlir, D.; Wong, J.; Jordan, H. L.; Schacker, T. W.; Racz, P.; Tenner-Racz, K.; Letvin, N. L.; Wolinsky, S.; Haase, A. T., Sexual transmission and propagation of SIV and HIV in resting and activated CD4+ T cells. *Science* **1999,** 286, (5443), 1353-7.

20. Groot, F.; van Capel, T. M.; Schuitemaker, J.; Berkhout, B.; de Jong, E. C., Differential susceptibility of naive, central memory and effector memory T cells to dendritic cell-mediated HIV-1 transmission. *Retrovirology* **2006,** 3, 52.

21. Mattei, S.; Flemming, A.; Anders-Osswein, M.; Krausslich, H. G.; Briggs, J. A.; Muller, B., RNA and Nucleocapsid Are Dispensable for Mature HIV-1 Capsid Assembly. *J Virol* **2015,** 89, (19), 9739-47.

22. Keller, P. W.; Huang, R. K.; England, M. R.; Waki, K.; Cheng, N.; Heymann, J. B.; Craven, R. C.; Freed, E. O.; Steven, A. C., A two-pronged structural analysis of retroviral maturation indicates that core formation proceeds by a disassembly-reassembly pathway rather than a displacive transition. *J Virol* **2013,** 87, (24), 13655-64.

23. Fontana, J.; Jurado, K. A.; Cheng, N.; Ly, N. L.; Fuchs, J. R.; Gorelick, R. J.; Engelman, A. N.; Steven, A. C., Distribution and Redistribution of HIV-1 Nucleocapsid Protein in Immature, Mature, and Integrase-Inhibited Virions: a Role for Integrase in Maturation. *J Virol* **2015,** 89, (19), 9765-80.

24. Hanne, J.; Gottfert, F.; Schimer, J.; Anders-Osswein, M.; Konvalinka, J.; Engelhardt, J.; Muller, B.; Hell, S. W.; Krausslich, H. G., Stimulated Emission Depletion Nanoscopy Reveals Time-Course of Human Immunodeficiency Virus Proteolytic Maturation. *ACS Nano* **2016,** 10, (9), 8215-22.

25. Sheehy, A. M.; Gaddis, N. C.; Choi, J. D.; Malim, M. H., Isolation of a human gene that inhibits HIV-1 infection and is suppressed by the viral Vif protein. *Nature* **2002,** 418, (6898), 646-50.

26. Izumi, T.; Shirakawa, K.; Takaori-Kondo, A., Cytidine deaminases as a weapon against retroviruses and a new target for antiviral therapy. *Mini Rev Med Chem* **2008,** 8, (3), 231-8.

27. Renner, N.; Kleinpeter, A.; Mallery, D. L.; Albecka, A.; Rifat Faysal, K. M.; Bocking, T.; Saiardi, A.; Freed, E. O.; James, L. C., HIV-1 is dependent on its immature lattice to recruit IP6 for mature capsid assembly. *Nat Struct Mol Biol* **2023,** 30, (3), 370-382.

431   28.   Sowd, G. A.; Aiken, C., Correction: Inositol phosphates promote HIV-1 assembly and
432   maturation to facilitate viral spread in human CD4+ T cells. *PLoS Pathog* **2021,** 17,
433   (3), e1009389.
434   29.   Parthasarathy, L. K.; Seelan, R. S.; Tobias, C.; Casanova, M. F.; Parthasarathy, R. N.,
435   Mammalian inositol 3-phosphate synthase: its role in the biosynthesis of brain inositol
436   and its clinical use as a psychoactive agent. *Subcell Biochem* **2006,** 39, 293-314.
437   30.   Dick, R. A.; Zadrozny, K. K.; Xu, C.; Schur, F. K. M.; Lyddon, T. D.; Ricana, C. L.;
438   Wagner, J. M.; Perilla, J. R.; Ganser-Pornillos, B. K.; Johnson, M. C.; Pornillos, O.;
439   Vogt, V. M., Inositol phosphates are assembly co-factors for HIV-1. *Nature* **2018,** 560,
440   (7719), 509-512.
441   31.   Freed, E. O., HIV-1 assembly, release and maturation. *Nat Rev Microbiol* **2015,** 13,
442   (8), 484-96.
443   32.   Tedbury, P. R.; Novikova, M.; Alfadhli, A.; Hikichi, Y.; Kagiampakis, I.;
444   KewalRamani, V. N.; Barklis, E.; Freed, E. O., HIV-1 Matrix Trimerization-Impaired
445   Mutants Are Rescued by Matrix Substitutions That Enhance Envelope Glycoprotein
446   Incorporation. *J Virol* **2019,** 94, (1).
447   33.   Tabler, C. O.; Wegman, S. J.; Chen, J.; Shroff, H.; Alhusaini, N.; Tilton, J. C., The
448   HIV-1 Viral Protease Is Activated during Assembly and Budding Prior to Particle
449   Release. *J Virol* **2022,** 96, (9), e0219821.
450   34.   Wang, D.; Lu, W.; Li, F., Pharmacological intervention of HIV-1 maturation. *Acta*
451   *Pharm Sin B* **2015,** 5, (6), 493-9.
452   35.   Kleinpeter, A. B.; Freed, E. O., HIV-1 Maturation: Lessons Learned from Inhibitors.
453   *Viruses* **2020,** 12, (9).
454   36.   Nagai, T.; Miyawaki, A., A high-throughput method for development of FRET-based
455   indicators for proteolysis. *Biochem Biophys Res Commun* **2004,** 319, (1), 72-7.
456   37.   Hubner, W.; Chen, P.; Del Portillo, A.; Liu, Y.; Gordon, R. E.; Chen, B. K., Sequence
457   of human immunodeficiency virus type 1 (HIV-1) Gag localization and
458   oligomerization monitored with live confocal imaging of a replication-competent,
459   fluorescently tagged HIV-1. *J Virol* **2007,** 81, (22), 12596-607.

Figure 1

Figure 2