Generating single-cell gene expression profiles for

high-resolution spatial transcriptomics based on cell boundary

images

Bohan Zhang[1,2,†], Mei Li[1,3,†], Qiang Kang[1,†], Zhonghan Deng[1], Hua Qin[2], Kui Su[1], Xiuwen Feng[1],

Lichuan Chen[1], Huanlin Liu[1], Shuangsang Fang[2], Yong Zhang[1], Yuxiang Li[1], Susanne Brix[3,*], Xun

Xu[1,*]

1 BGI Research, Shenzhen 518083, China

2 BGI Research, Beijing 102601, China

3 Department of Biotechnology and Biomedicine, Technical University of Denmark, 2800 Kgs.

Lyngby, Denmark

† Contributed equally.

* Corresponding author. E-mail: sbrix@dtu.dk, xuxun@genomics.cn.

ABSTRACT

Stereo-seq is a cutting-edge technique for spatially resolved transcriptomics that combines subcellular resolution with centimeter-level field-of-view, serving as a technical foundation for analyzing large tissues at the single-cell level. Our previous work presents the first one-stop software that utilizes cell nuclei staining images and statistical methods to generate high-confidence single-cell spatial gene expression profiles for Stereo-seq data. With recent advancements in Stereo-seq technology, it is possible to acquire cell boundary information, such as cell membrane/wall staining images. To take advantage of this progress, we update our software to a new version, named STCellbin, which utilizes the cell nuclei staining images as a bridge to align cell membrane/wall staining images with spatial gene expression maps. By employing an advanced cell segmentation technique, accurate cell boundaries can be obtained, leading to more reliable single-cell spatial gene expression profiles. Experimental results verify that STCellbin can be applied on the mouse liver (cell membranes) and *Arabidopsis* seed (cell walls) datasets and outperforms other competitive methods. The improved capability of capturing single cell gene expression profiles by this update results in a deeper understanding of the contribution of single cell phenotypes to tissue biology.

Availability & Implementation: The source code of STCellbin is available at https://github.com/STOmics/STCellbin.

STATEMENT OF NEED

Spatially resolved single cell transcriptomics enables the generation of comprehensive molecular maps that provide insights into the spatial distribution of molecules within the single cells that make up tissues. This groundbreaking technology offers insights into the location and function of cells in various tissues, enhancing our knowledge of organ development [1], tumor heterogeneity [2], cancer evolution [3], and other biological mechanisms. Resolution and field-of-view are two critical parameters in spatial transcriptomics. High resolution enables detailed molecular information at the single-cell level, and large field-of-view facilitates the creation of complete 3D maps that represent biological functions at the organ level. Stereo-seq simultaneously achieves subcellular resolution and a centimeter-level field-of-view, providing a technical foundation for obtaining comprehensive spatial gene expression profiles of whole tissues at single-cell level [4]. Our previous work offers the one-stop software StereoCell for acquiring high signal-to-noise ratio single-cell spatial gene expression profiles from Stereo-seq data [5]. StereoCell takes the cell nuclei staining image tiles and its corresponding spatial gene expression data as input, and its main process includes image stitching, image registration, tissue segmentation, cell nuclei segmentation and molecule labeling steps. The image data generated by Stereo-seq used for StereoCell are cell nuclei staining images. However, there is a big difference between cell nuclei and cell boundary staining images, based on cell membrane/wall staining, in terms of the ability to capture robust and precise cell specific gene expression profiles. Despite the widespread use of spatial techniques, such as MERFISH [6], CosMx [7], and Xenium [8], several of these techniques still struggle to achieve accurate cell boundary information, as they are based on cell nuclei staining images that can be generated using stains such as 4,6-diamidino-2-phenylindole (DAPI). Hematoxylin-eosin

(H&E) and single strand DNA fluorescence (ssDNA) nuclei staining images are also commonly

used and readily obtainable data. The updated Stereo-seq technology implement a procedure based

on simultaneous cell membrane/wall and cell nuclei staining by adding multiplex

immunofluorescence (mIF) and calcofluor white (CFW) staining [9,10], which can automatically

acquire more accurate cell boundary information and thereby obtain more reliable single-cell

spatial gene expression profiles.

We update StereoCell to a new version, named STCellbin. We retain the image stitching,

tissue segmentation and molecule labeling steps from StereoCell and improve the image

registration and cell segmentation steps. The "track line" is a crossed linear marker embedded on

the Stereo-seq chip, which is the key information for the image registration step of StereoCell [5].

As the cell membrane/wall staining images miss the "track line" information, we utilize the cell

nuclei staining images as a bridge to align the cell membrane/wall staining images with the spatial

gene expression maps, upon which we obtain the registered cell boundary information in the cell

segmentation step. Based on the cell boundary information, we directly assign the molecules to

their corresponding cells, obtaining single-cell spatial gene expression profiles. We apply

STCellbin on mouse liver (cell membrane) and *Arabidopsis* seed (cell wall) datasets and confirm

the accuracy of cell segmentation. This update offers a comprehensive workflow to obtain more

reliable single-cell spatial gene expression profiles based on cell membrane/wall information,

providing support and guidance for related scientific investigations, particularly those based on

Stereo-seq data.

IMPLEMENTATION

Overview of STCellbin

The process of STCellbin includes image stitching, image registration, cell segmentation and molecule labeling (Fig. 1). The Stereo-seq spatial gene expression data, cell nuclei and cell membrane/wall staining image tiles are input into STCellbin. The stitched cell nuclei and cell membrane/wall staining images are obtained through the MFWS algorithm [5]. These two stitched staining images are registered using the Fast Fourier Transform (FFT) algorithm [11]. The spatial gene expression data is transformed into a map, this map and a stitched cell nuclei staining image are registered based on "track lines". Thus, the registration of the gene expression map and cell membrane/wall staining image is implemented. Cell segmentation is performed on the registered cell membrane/wall staining image by the adjusted Cellpose 2.0 [12] to obtain the cell mask. The molecules are assigned to their corresponding cells according to the cell mask to obtain the single-cell spatial gene expression profile as the output. The tissue segmentation step based on Bi-Directional ConvLSTM U-Net [13] is set as optional, which can generate a tissue mask to assist in filtering out impurities outside the tissue.

Image stitching

The image stitching step in STCellbin is consistent with it in StereoCell. The MFWS algorithm [5] is adopted, which calculates the offsets of two adjacent tiles with overlapping areas using FFT [11] to stitch these two tiles and extends this process to all tiles. The relative error, absolute error and computational efficiency of MFWS have been verified in our previous work [5].

Image registration

The registration of STCellbin includes three stages. The first stage is the registration of the

stitched cell nuclei and stitched cell membrane/wall staining images. These two staining images

have similar sizes and no large difference in rotation because the chip does not move when they

are photographed. The key to this registration is to calculate their offset. The size of the cell

membrane/wall staining image is adjusted to be consistent with the cell nuclei staining image by

cutting and zero-padding (Fig. 2A). The two staining images are mean-based subsampled [14]

(Fig. 2B). The offset of the subsampled images is calculated through FFT [11], similar to MFWS

[5] (Fig. 2C). The calculated offset is restored to the scale of the original images (Fig. 2D). Thus,

these two staining images can be registered. The second stage is the registration of the stitched

nuclei staining image and spatial gene expression map. This registration is the same as it in

StereoCell [5]. The spatial gene expression data is transformed into a map. The stitched cell nuclei

staining image is registered with the map based on "track lines" by performing scaling, rotating,

flipping and translating on the stitched cell nuclei staining image. The third stage is the

registration of the stitched cell membrane/wall staining image and spatial gene expression map.

Since the cell nuclei and cell membrane/wall staining images have been registered in the first

stage, the same operations in the second stage, include scaling, rotating, flipping and translating,

are applied to the cell membrane/wall staining image (Fig. 2E). Then, the cell membrane/wall

staining image and spatial gene expression map can be registered. Moreover, when utilizing

staining images produced with a multi-channel microscope, STCellbin can omit the registration

among these images. STCellbin can also process the case of multiple mIF staining images taken

from identical tissues using the same microscope when there is only a difference in offsets among

these images.

Cell segmentation

The cell segmentation step of STCellbin is performed using Cellpose 2.0 [12] with some

adjustments. The model architecture of Cellpose 2.0 and its weight files "cyto2" are downloaded.

Due to the large size of staining images derived from Stereo-seq data, Cellpose 2.0 cannot be

executed smoothly using normal hardware configurations. To circumvent this issue, the staining

images are therefore cropped into multiple tiles with overlapping areas to perform cell

segmentation and record the coordinates of tiles. The overlapping areas rescue cells at the border

of the tiles from being cropped. To obtain the best results, segmentations with different values of

the cell diameter parameter are performed independently, and the result with the largest sum of

cell areas is retained. All the segmented tiles are assembled into the final segmented result

according to the recorded coordinates. Moreover, when selecting the tissue segmentation option,

an additional step is executed to apply a filter on the cell mask using the tissue mask, resulting in a

filtered segmented outcome.

Molecule labeling

The molecule labeling of STCellbin is the same as the one used in StereoCell in principle.

StereoCell assigns molecules in the cell nuclei to the cell by using the cell nuclei mask, and then

assigns molecules outside the cell nuclei to the cells with the highest probability density using

Gaussian Mixture Model [15]. STCellbin assigns molecules to the cells directly based on the cell

mask, while the process of assigning molecules outside the cell is included as an option. The latter

decision was made as the cell membranes/walls are usually tightly packed, with only a few

molecules outside the cells, and the assignment of these molecules takes a lot of time. Thus, we

generally do not recommend this option, and the users can use it according to actual requirements.

RESULTS

Datasets and computing resource

We adopt two datasets acquired via Stereo-seq technology [4]. One is a mouse liver dataset, a

tissue that offers cell boundary information via cell membranes, as in all mammalian tissues. The

other dataset is derived from seeds of the plant *Arabidopsis*, a tissue that provides cell boundary

information based on rigid cell walls. More details of the two datasets are shown in Table 1.

The experiment for image segmentation is implemented on the STOmics cloud platform [16]

with settings of 32 CPUs, 32 GB memory and "ALL" resource type. Except for the watershed

method [17], which is implemented using ImageJ on a computer with a 16-core CPU and 16GB of

RAM. The experiment for downstream analysis is implemented on a server with 40-core CPU,

128 GB of RAM and 24 GB of GPU.

Evaluation criteria for cell segmentation performance

In a cell mask image, the gray value of a pixel is set to 255 in the cell area and 0 in the background.

True positive (TP, the number of pixels with gray value of 255 in both ground truth and segmented

result), true negative (TN, the number of pixels with gray value of 0 in both ground truth and

segmented result), false positive (FP, the number of pixels with gray value of 0 in ground truth and

255 in segmented result) and false negative (FN, the number of pixels with gray value of 255 in

ground truth and 0 in segmented result) are calculated. The number of cells segmented by a

method is ns. For each segmented cell ($cell_i$), there should be a corresponding area in ground truth

($area_i$), where i is the cell index ($i = 1, 2, …, ns$). The intersection over union metric (IoU) [18] is

set as:

$$IoU_i = ao_i \big/ au_i \tag{1}$$

163 where ao_i is the overlap area between $cell_i$ and $area_i$, and au_i is the union area of $cell_i$ and $area_i$.

164   Then the precision (*Pre*), recall (*Rec*), F1 score (*F1_s*), Dice coefficient (*Dc*) and average Jaccard

165   index (*Avg_J*) are calculated as:

$$Pre = \mathrm{TP}/(\mathrm{TP} + \mathrm{FP}) \tag{2}$$

$$Rec = \mathrm{TP}/(\mathrm{TP} + \mathrm{FN}) \tag{3}$$

$$F1\_s = 2 \times (Pre \times Rec)/(Pre + Rec) \tag{4}$$

$$Dc = 2 \times \mathrm{TP}/(2 \times \mathrm{TP} + \mathrm{FP} + \mathrm{FN}) \tag{5}$$

$$Avg\_J = \sum_{i=1}^{ns} IoU_i \Big/ ns \tag{6}$$

Process and evaluation of downstream analysis

The generated single-cell spatial gene expression profiles are input into Stereopy (v0.6.0) [19].

The cells with fewer than 10 expressed genes, fewer than 3 expression counts and more than 3%

mitochondrial genes are removed, and genes present in less than 3 cells are also removed. After

normalization, the differentially expressed genes are summarized by Principal Component

Analysis to reduce the data dimensionality, in which the number of features after reducing is 10.

The Leiden algorithm [20] is used for clustering and the Uniform Manifold Approximation and

Projection (UMAP) algorithm [21] is used to obtain 2D data projection. The Silhouette coefficient

(*Sc*) and Moran's I (*MI*) are used to evaluate the effect of clustering and the spatial self-correlation

of each cluster, respectively. *Sc* is calculated as:

$$Sc = (b_j - a_j)/\max\{a_j, b_j\} \tag{7}$$

176   where a_j is the average distance between the *j*-th sample and other samples in its cluster, and b_j is

177   the average distance between the *j*-th sample and the samples in other clusters. *MI* is calculated as:

$$MI = (n/W_0)\left(\sum_{k=1}^{n}\sum_{l=1}^{n}\omega_{k,l}(y_k - \bar{y})(y_l - \bar{y})\Big/\sum_{k=1}^{n}(y_k - \bar{y})^2\right) \tag{8}$$

178   where *n* is the number of clusters, y_k and y_l are the attribute value of the *k*-th and *l*-th clusters,

179   respectively, $y\square$ is the mean of all cluster attributes, $\omega_{k,l}$ is the spatial weight between k-th and l-th

180   clusters, and W_0 is the aggregation of all spatial weights as:

$$W_0 = \sum_{k=1}^{n} \sum_{l=1}^{n} \omega_{k,l} \tag{9}$$

STCellbin more accurately segments cells based on cell membrane/wall staining

images

We crop two areas with higher image quality from the two datasets and design their ground truths

based on the manual markup of the cells according to the cell membranes/walls in the staining

images. The cell segmentation method of STCellbin is compared with the original Cellpose [18], a

state-of-the-art method DeepCell [22] and a traditional watershed method [17].

On the mouse liver dataset, STCellbin effectively identifies cell membranes for segmentation,

yielding cell masks that exhibit acceptable agreement with the stating image and ground truth (Fig.

3A, upper). Among all cell mask images, STCellbin provides the best description of the cell

boundaries, outperforming other methods that tend to miss quite a few cells (Fig. 3A, lower). We

observe a similar situation on the *Arabidopsis* seed dataset, which indicates that STCellbin can

also effectively identify cell walls for segmentation (Fig. 3B). Compared with other methods,

STCellbin obtains higher values in most indicators on two datasets (Fig. 3C). The comparison

with the original Cellpose validates the effectiveness of STCellbin in adjusting segmentation.

While DeepCell is a powerful method, it is primarily designed for segmenting cell nuclei, which

involves identifying highlighted areas in the nuclei staining images. This strategy is not suitable

for cell membrane/wall staining images, resulting in less desirable results. Similarly, the

traditional watershed method performs poorly on cell membrane/wall staining images. In general,

STCellbin's cell segmentation is the most feasible method.

STCellbin generates more reliable single-cell spatial gene expression profile for downstream analysis

Currently, there is a lack of image-based one-stop software like STCellbin for Stereo-seq data. Therefore, we compare STCellbin with Baysor (v0.6.2) [23], a tool that generates the spatial gene expression profile without relying on image. However, Baysor cannot output the results on the complete mouse liver and *Arabidopsis* seed datasets in an acceptable time or a given computational resource. We run Baysor on a smaller *Arabidopsis* seed dataset, which is the cropped area in cell segmentation experiment and contains two complete seed data.

The cell area, number of unique genes per cell and number of gene counts per cell are statistically calculated from the results of STCellbin (Fig. 4A). The clustering results of STCellbin are obtained by utilizing the generated single-cell spatial gene expression profiles. The clusters of cells are spatially mapped within the tissue (Fig. 4B, left hand side for each tissue), allowing for the observation of their specific positions. From the UMAPs, it is apparent that the different cell types are effectively distinguished (Fig. 4B, right hand side for each tissue). The spatial location of the different cell types will positively influence a series of downstream analyzes such as cellular annotation in less well-studied tissues. The cells are clustered into 7 clusters on the profile of STCellbin, while 14 clusters on the profile of Baysor (Fig. 4C, the 1st subfigure from the left). We observe that the number of cells segmented by Baysor is significantly higher than that segmented by STCellbin, and it does not align with the cell count observed in the ground truth image. This may be the main reason for the higher number of clusters of Baysor. The silhouette coefficient and Moran's I obtained by both STCellbin and Baysor are not very satisfactory (Fig. 4C, the 2nd and 3rd subfigures from the left), possibly due to the limited information from a small dataset.

Nevertheless, the values of STCellbin are higher than those of Baysor. Moreover, STCellbin has

significant advantages in terms of computing resource usage and running time (Fig. 4C, the 4th

and 5th subfigures from the left), which also explains why Baysor cannot process the complete

mouse liver and Arabidopsis seed datasets. It should be noted that the stitching and registration

steps of STCellbin cannot be performed on the cropped dataset, and then, the related

computational resource usage and running time cannot be recorded. Thus, the resource usage and

time of STCellbin for comparison are obtained on the complete *Arabidopsis* seed dataset. That is,

STCellbin can process a larger dataset with fewer computational resources and less time compared

to Baysor. Overall, STCellbin is a more reliable method, particularly for analyzing high-resolution

and large-field-of-view spatial transcriptomic data.

Discussion

Accurate identification of cell boundaries plays a crucial role in generating single-cell resolution

in spatial omics applications. Based on previous work in StereoCell using cell nuclei staining

images to generate single-cell spatial gene expression profiles, this STCellbin update extends the

capability to automatically process Stereo-seq cell membrane/wall staining images for

identification of cell boundaries that facilitates downstream analyses. We also showcase a few

examples of the performance of cell membrane/wall segmentation in STCellbin. Currently, the

tools for cell nuclei and cell membrane/wall segmentation can be independently executed,

allowing users to choose the more suitable solution for their specific applications. In future work,

these two techniques can be combined by training a deep learning model that is compatible with

any staining image type, thereby achieving more accurate results.

AVAILABILITY OF SOURCE CODE AND REQUIREMENTS

- Project name: STCellbin

- Project home page: https://github.com/STOmics/STCellbin

- Operating system(s): Platform independent

- Programming language: Python

- Other requirements: Python 3.8

- License: MIT License

- RRID: SCR_024438

DATA AVAILABILITY

The data that support the findings of this study have been deposited into Spatial Transcript Omics DataBase (STOmics DB) of China National GeneBank DataBase (CNGBdb) with accession number STT0000048: https://db.cngb.org/stomics/project/STT0000048. A backup link for the data is also provided at Github link of STCellbin: https://github.com/STOmics/STCellbin.

LIST OF ABBREVIATIONS

DAPI: 4,6-diamidino-2-phenylindole; H&E: hematoxylin-eosin; ssDNA: single strand DNA fluorescence; mIF: multiplex immunofluorescence; CFW: calcofluor white; FFT: Fast Fourier Transform.

DECLARATIONS

Ethics Approval and Consent to Participate

Not applicable.

Competing Interests

The authors declare that they have no competing interests.

Funding

This work was supported by the National Key R&D Program of China (2022YFC3400400).

Authors' Contributions

Conceptualization: BZ and ML; Project administration and supervision: SB and XX; Software

implementation: ZD, HQ, KS and HL; Data collection and processing: QK, XF and LC;

Validation: QK and ZD. Project coordination: BZ and ML; Manuscript writing and figure

generation: BZ, ML and QK; Manuscript review: ML, SF, YZ, YL and SB.

Acknowledgements

We thank China National GeneBank for providing technical support.

REFERENCES

1.   Fang S, Chen B, Zhang Y, Sun H, Liu L, Liu S, et al. Computational Approaches and Challenges

in Spatial Transcriptomics. *Genom Proteom Bioinf*. 2023;21:24–47.

2.   Lu T, Ang CE, Zhuang X. Spatially resolved epigenomic profiling of single cells in complex

tissues. *Cell*. 2022;185:4448–4464.

3.   Erickson A, He M, Berglund E, Marklund M, Mirzazadeh R, Schultz N, et al. Spatially resolved

clonal copy number alterations in benign and malignant tissue. *Nature*. 2022;608:360–367.

4.   Chen A, Liao S, Cheng M, Ma K, Wu L, Lai Y, et al. Spatiotemporal transcriptomic atlas of mouse

organogenesis using DNA nanoball-patterned arrays. *Cell*. 2022;185:1777-1792.

283 5. Li M, Liu H, Li M, Fang S, Kang Q, Zhang J, et al. StereoCell enables highly accurate single-cell

284   segmentation for spatial transcriptomics. *bioRxiv*. 2023.

285 6. Chen KH, Boettiger AN, Moffitt JR, Wang S, Zhuang X. Spatially resolved, highly multiplexed

286   RNA profiling in single cells. *Science*. 2015;348:aaa6090.

287 7. He S, Bhatt R, Brown C, Brown EA, Buhr DL, Chantranuvatana K, et al. High-plex imaging of

288   RNA and proteins at subcellular resolution in fixed tissue by spatial molecular imaging. *Nat*

289   *Biotechnol*. 2022;40:1794-1806.

290 8. Janesick A, Shelansky R, Gottscho AD, Wagner F, Rouault M, Beliakoff G, et al. High resolution

291   mapping of the breast cancer tumor microenvironment using integrated single cell, spatial and in

292   situ analysis of FFPE tissue. *bioRxiv*. 2022.

293 9. Liao S, Heng Y, Liu W, Xiang J, Ma Y, Chen L, et al. Integrated Spatial Transcriptomic and

294   Proteomic Analysis of Fresh Frozen Tissue Based on Stereo-seq. *bioRxiv*. 2023.

295 10. STOmics Decuments. https://en.stomics.tech/.

296 11. Duhamel P, Vetterli M. Fast fourier transforms: A tutorial review and a state of the art. *Signal*

297   *Process*. 1990;19(4):259-299.

298 12. Pachitariu M, Stringer C. Cellpose 2.0: how to train your own model. *Nat Methods*.

299   2022;19:1634-1641.

300 13. Azad R, Asadi-Aghbolaghi M, Fathy M, Escalera S. Bi-Directional ConvLSTM U-Net with

301   densley connected convolutions. *arXiv*. 2019.

302 14. Levina A, Priesemann V. Subsampling scaling. *Nat Commun*. 2017;8:15140.

303 15. Reynolds D. Gaussian Mixture Models. *Encycl Biom*. 2009;741:659-663.

304 16. https://cloud.stomics.tech/

17. Wen T, Tong B, Liu Y, Pan T, Du Y, Chen Y, et al. Review of research on the instance

segmentation of cell images. *Comput Meth Prog Bio*. 2022;227:107211.

18. Stringer C, Wang T, Michaelos M, Pachitariu M. Cellpose: a generalist algorithm for cellular

Segmentation. *Nat Methods*. 2021;18:100-106.

19. Fang S, Xu M, Cao L, Liu X, Bezulj M, Tan L, et al. Stereopy: modeling comparative and

spatiotemporal cellular heterogeneity via multi-sample spatial transcriptomics. bioRxiv. 2023.

20. Traag VA, Waltman L, Van Eck NJ. From Louvain to Leiden: guaranteeing well-connected

communities. *Sci Rep*. 2019;9:5233.

21. Becht E, McInnes L, Healy J, Dutertre C, Kwok IWH, Ng LG, et al. Dimensionality reduction for

visualizing single-cell data using UMAP. *Nat Biotechnol*. 2019;37:38-44.

22. Greenwald NF, Miller G, Moen E, Kong A, Kagel A, Dougherty T, et al. Whole-cell segmentation

of tissue images with human-level performance using large-scale data annotation and deep

learning. *Nat Biotechnol*. 2022;40:555-565.

23. Petukhov V, Xu RJ, Soldatov RA, Cadinu P, Khodosevich K, Moffitt JR, et al. Cell segmentation

in imaging-based spatial transcriptomics. *Nat Biotechnol*. 2022;10:345-354.

Figure legends

Figure 1. Overview of STCellbin. The cell nuclei and cell membrane/wall staining image tiles are stitched into

individual large images respectively. The spatial gene expression map and stitched cell membrane/wall staining

image are registered with the stitched cell nuclei staining image as a bridge. The cell mask is directly obtained

from the registered cell membrane/wall staining image by cell segmentation. The single-cell spatial gene

expression profile is obtained by overlaying the generated cell mask and the gene expression map.

Figure 2. Registration of the cell membrane/wall staining image and spatial gene expression map using the cell

nuclei staining image as a bridge. **A**. Size of the cell membrane/wall staining image is adjusted to be consistent

with the cell nuclei staining image. **B**. Cell nuclei and cell membrane/wall staining images are subsampled. **C**.

Calculating the offsets of the subsampled images. **D**. Restoring the offsets to the scale of original images for

registration. E. Registering the spatial gene expression map and cell nuclei staining image by performing scaling,

rotating, flipping and translating, and registering the spatial gene expression map and cell membrane/wall staining

image by performing the same operations.

Figure 3. Comparison of cell segmentation performance. **A**. Cell segmentation results on the cropped area from

mouse liver dataset, where in the merged images, cell masks are set in yellow, staining images are set in cyan, and

ground truths are set in red. **B**. Cell segmentation results on the cropped area from *Arabidopsis* seed dataset, where

in the merged images, cell masks are set in yellow, staining images are set in cyan, and ground truths are set in red.

C. Indicator comparison of cell segmentation results on the cropped areas from mouse liver and *Arabidopsis* seed

datasets.

Figure 4. Downstream analysis results and comparison. **A**. Statistical results of cell area, gene number per cell and

gene expression per cell of STCellbin on the mouse liver dataset and cropped area from *Arabidopsis* seed dataset.

B. Clustering results and UMAPs of STCellbin on the mouse liver dataset and cropped area from *Arabidopsis* seed

dataset. **C**. Indicator comparison of downstream analysis results on the cropped area from *Arabidopsis* seed dataset,

where *Sc* and *MI* are taken from mean of the values of all clusters.

Table

Table 1. Details of two datasets used for evaluation of cell boundary information

Detail	Mouse liver dataset	*Arabidopsis* seed dataset

Data source	A slice of liver	Slices of multiple seeds
Cell nuclei dye	DAPI	ssDNA
Cell membrane/wall dye	mIF	CFW
Number of molecules	16,177,288	62,884,637

tissue

microscope

cell membrane/wall
staining tiles

image stitching

stitched cell membrane/wall
staining image

image stitching

cell nuclei
staining tiles

stitched cell nuclei
staining image

sequenator

spatial gene
expression map

Image registration

molecule labeling

cell segmentation

single-cell spatial gene
expression profile

cell mask

registered images

A

cell nuclei
staining image

cell membrane/wall
staining image

zero-padding

cutting

B

subsampling

subsampled cell
nuclei staining image

subsampled cell
membrane/wall staining image

C

x'

y'

**offset calculation
on subsampled
images using FFT**

D

x

y

**offset restoration to
the scale of original
images for registration**

E

cell membrane/wall
staining image

**registration by performing
same operations**

spatial gene
expression map

cell nuclei
staining image

scaling

rotating

translating

flipping