Scalable ultra-high-throughput single-cell chromatin and RNA sequencing reveals gene regulatory dynamics linking macrophage polarization to autoimmune disease

Author list

Sara Lobato-Moreno[*,1,2,5], Umut Yildiz[*,2,5], Annique Claringbould[*1,2], Nila H. Servaas[1], Evi P. Vlachou[1], Christian Arnold[1], Hanke Gwendolyn Bauersachs[1], Víctor Campos-Fornés[2,5], Karin D. Prummel[1,2], Kyung Min Noh[+,2], Mikael Marttinen[+,1,2,3], Judith B. Zaugg[+,1,2,4]

Affiliations

1) European Molecular Biology Laboratory, Structural Computational Biology Unit
2) European Molecular Biology Laboratory, Genome Biology Unit
3) current address: Faculty of Medicine and Health Technology, Tampere University
4) Molecular Medicine Partnership Unit (MMPU), Heidelberg, Germany
5) Faculty of Biosciences, Collaboration for Joint PhD Degree between EMBL and Heidelberg University, Heidelberg, Germany
*) these authors contributed equally
+) Co-correspondence: mikael.marttinen@embl.de, kyung.min.noh@embl.de, judith.zaugg@embl.de

Abstract

Enhancers and transcription factors (TFs) are crucial in regulating cellular processes, including disease-associated cell states. Current multiomic technologies to study these elements in gene regulatory mechanisms lack multiplexing capability and scalability. Here, we present SUM-seq, a cost-effective, scalable **S**ingle-cell **U**ltra-high-throughput **M**ultiomic sequencing method for co-assaying chromatin accessibility and gene expression in single nuclei. SUM-seq enables profiling hundreds of samples at the million cell scale and outperforms current high-throughput single-cell methods. We applied SUM-seq to dissect the gene regulatory mechanisms governing macrophage polarization and explored their link to traits from genome-wide association studies (GWAS). Our analyses confirmed known TFs orchestrating M1 and M2 macrophage programs, unveiled key regulators, and demonstrated extensive enhancer rewiring. Integration with GWAS data further pinpointed the impact of specific TFs on a set of immune traits. Notably, inferred enhancers regulated by the STAT1/STAT2/IRF9 (ISGF3) complex were enriched for genetic variants associated with Crohn's disease, ulcerative colitis and multiple sclerosis, and their target genes included known drug targets. This highlights the potential of SUM-seq for dissecting molecular disease mechanisms. SUM-seq offers a cost-effective, scalable solution for ultra-high-throughput single-cell multiomic sequencing, excelling in unraveling complex gene regulatory networks in cell differentiation, responses to perturbations, and disease studies.

Introduction

Gene regulatory elements in enhancer regions play a major role in disease development, as evidenced by the fact that most genome-wide association studies (GWAS) single nucleotide polymorphisms (SNPs) fall in non-coding gene regulatory regions (reviewed in[1]). Moreover, many diseases manifest through an imbalance of cellular differentiation, highlighting the importance of studying enhancer dynamics in the context of cellular differentiation. For example, many autoimmune diseases are characterized by an imbalance of pro- and anti-inflammatory immune cells[2–4]. Thus, studying such disease mechanisms requires a technology that allows the joint analysis of enhancer and transcription factor (TF) dynamics along a differentiation time course at single-cell resolution.

Recent developments in scalability of single-cell omics technologies, specifically single-cell RNA-seq (scRNA-seq) and single-nucleus (sn)ATAC-seq have revolutionized our understanding of the diversity of cell states and cellular responses to perturbations (reviewed in[5]). Particularly, multimodal profiling has enhanced our ability to unravel gene regulatory dynamics governing fundamental biological processes[6–8]. However, while scalability for individual modalities has become available (e.g., scifi-RNA-seq[9], sci-RNA-seq[10], dsci-ATAC-seq[11], sci-ATAC-seq[12]), the majority of current multiomic methods are limited in scalability, multiplexing capability, or cost effectiveness (e.g., 10x multiome, ISSAAC-seq[13]), with a handful providing scalability in terms of number of cells assayed (e.g., SHARE-seq[14], Paired-seq[15]), but at the expense of complexity (**Table 1**).

Here, we present SUM-seq (**S**ingle-cell **U**ltra-high-throughput **M**ultiomic sequencing): a cost effective and scalable sequencing technique for multiplexed multiomics profiling. SUM-seq enables simultaneous profiling of chromatin accessibility and gene expression in single nuclei at ultra-high-throughput scale (up to millions of cells and hundreds of samples[9]). To this end, we refined the two-step combinatorial indexing approach, originally introduced by Datlinger et al.[9] for snRNA-seq, tailoring it to the multiomic setup. In brief, accessible chromatin and nuclear mRNA first receive a sample-specific index by transposition and reverse transcription, respectively. Subsequently, a second index is introduced using a droplet-based microfluidic platform (e.g. 10x Chromium). The dual indexing allows overloading of microfluidic droplets while retaining the capacity to assign matched RNA- and ATAC-seq reads to the same individual cell.

We demonstrate the application of SUM-seq in two independent experimental setups, a species-mixing benchmarking experiment and a macrophage M1 and M2 polarization time course experiment, exemplifying its flexibility to accommodate complex experimental setups. We use both modalities to resolve temporal patterns of gene regulation, and integrate with genetic evidence to link regulatory networks to disease. Additionally, we validate the feasibility of decoupling sample collection from library generation, rendering the method well-suited for extensive atlas projects, time course experiments, and other experimental designs that involve prolonged sample collection periods.

Main

Overview of the SUM-seq method

SUM-seq is built upon the concept of a recently developed combinatorial fluidic indexing scRNA-seq approach, scifi-RNA-seq[9], and a low-throughput multiomic assay, ISSAAC-seq[13]. SUM-seq has 5 key steps: In the first step, nuclei are isolated and fixed with glyoxal, then split into equal bulk aliquots in a multi-well format. In the second and third steps, unique sample indices (one per well) are introduced for the ATAC and RNA modalities. For ATAC, accessible genomic regions are indexed by Tn5 loaded with barcoded oligos. For RNA, the mRNA molecules are indexed with barcoded oligo-dT primers via reverse transcription (**Figure 1a, detailed library structures in Extended Data Figure 1a**,**b, Supplementary Table 1**). In step four, samples are pooled for Tn5-based tagmentation of the cDNA-mRNA hybrids to introduce a primer binding site necessary for binding of the barcoded oligonucleotides in the microfluidic system. In step five, nuclei are overloaded onto the microfluidic system (e.g., 10x Chromium), facilitating the encapsulation of multiple nuclei in a single droplet. Within these droplets, fragments are barcoded with the microfluidic barcode, resulting in dual barcoding of fragments with both a sample index and a droplet barcode enabling downstream demultiplexing and assignment of sequencing reads to individual nuclei. This is followed by an initial amplification of both modalities after which the library is split into two equal proportions and subjected to modality-specific amplification. At this stage, a library index can be introduced to enable multiplexing libraries from independent experiments on the sequencer. snATAC and snRNA libraries are sequenced using standard Illumina sequencing primers. Additionally, we have developed a scalable and reproducible Snakemake pipeline for processing SUM-seq data (https://git.embl.de/grp-zaugg/SUMseq; available upon publication), in which reads are assigned to sample indices and further demultiplexed to single-cell resolution by the droplet barcode (**Figure 1b**). From here, reads are mapped and a gene expression matrix and tile matrix are generated for RNA and ATAC, respectively. Modalities can be matched based on the assigned sample index-cell barcode combinations. The pipeline provides reproducibility, high scalability and flexibility for execution on any system (local, cluster, cloud). In brief, SUM-seq offers a user-friendly and rapid workflow for ultra-high-throughput single-cell multiomic sequencing that doesn't require specialized instrumentation and is compatible with a widely available microfluidic platform.

SUM-seq enables efficient and scalable single-cell multiomics analysis

To evaluate the performance of SUM-seq and the data quality, we performed SUM-seq on an equal mixture of human leukemia (K562) and mouse fibroblast (NIH-3T3) cell lines. For this species-mixing benchmarking experiment, we loaded 100,000 nuclei into a single channel of the 10x Chromium system which equates to ~7-fold overloading compared to the standard 10x workflow. To limit the sequencing requirements, we processed only 20 % (20,000 nuclei) of the generated microfluidic droplets for final library preparation. After combinatorial index demultiplexing, human and mouse reads were well-separated with a collision rate of 0.1 % (UMIs) and 3.8 % (ATAC fragments), resulting in 6,215 human cells and 7,607 mouse cells with data from both modalities (**Figure 1c**,**d**). This translates to a ~70 % recovery rate of input loaded to the 10x Chromium system and represents a ~7-fold increase in throughput at low collision rates compared to the standard 10x workflow (**Extended Data Figure 1d**). Key performance metrics of snRNA (UMIs and genes per cell) and snATAC (fragments in peaks

per cell, TSS enrichment score, fragment size distribution) were consistently of very high
quality for both modalities in SUM-seq and outperformed other high-throughput assays for
scRNA, snATAC, and multiomic approaches (**Figure 1e, Extended Data Figure 2c-g**).
Notably, there was no reduction in data quality for nuclei in overloaded droplets compared to
droplets containing a single nucleus (**Figure 1f**). Lastly, the aggregate of snRNA and snATAC
data resembled the published bulk RNA-seq and ATAC-seq in the K562 benchmarks from
ENCODE data (**Figure 1g;** GSE86660[16] (RNA), GSE170214[16] (ATAC)).
131
During protocol optimization, we evaluated the effects of three factors on data quality: I) adding
polyethylene glycol (PEG), II) methods to mitigate barcode hopping, and III) freezing of
samples. (I) We found that, consistent with previous studies[17,18], the addition of 12 % PEG to
the reverse transcription reaction increased the number of UMIs and genes detected per cell
(~2.5- and ~2-fold, respectively) with minor impact on the quality metrics of the ATAC modality
(**Extended Data Figure 2h,i**). (II) We observed that barcode hopping can occur within
multinucleated droplets[19,20], primarily affecting the ATAC modality. However, we were able to
mitigate it by using two complementary strategies: adding a blocking oligonucleotide[20] in
excess to the droplet barcoding step and reducing the number of linear amplification cycles
during droplet barcoding from twelve to four (**Methods**) (**Figure 1b,c** and **Extended Data
Figure 2a**). (III) To assess the suitability of SUM-seq for asynchronous sample collection (e.g.
time course data or clinical samples), we tested glycerol-based cryopreservation following
glyoxal fixation and found that it had minimal impact on the performance metrics of the assay
(**Extended Data Figure 2j**). Thus, SUM-seq can be employed for gathering, fixing, and
cryopreserving samples prior to library construction, which enables multiplexing samples in
one experiment and thereby reducing costs and batch effects.
148

149
Figure 1. SUM-seq allows simultaneous profiling of chromatin accessibility and gene expression
in single cells at ultra-high-throughput scale
a, Schematic depiction of the SUM-seq workflow. Key steps and detailed structures are described in
the main text and **Extended Data Figure 1**. **b,** Schematic overview of the computational analysis
pipeline. **c, d,** Species-mixing plots for the ATAC (**c**) and the RNA (**d**) modality, indicating the fraction
of reads from singlets and multiplets assigned to the mouse genome (mm10, y-axes) and the human
genome (hg38, x-axes). Genomic DNA fragments (**c**) and transcripts (**d**) are demultiplexed based on
the combination of the sample index and the droplet barcode. The pie-chart shows the fraction of human
and mouse cells as well as the frequency of multiplets and singlets. Collision rates are highlighted in
red. **e,** Number of accessible DNA fragments in peaks (FRiP = fragments in peaks; upper panel) and
genes detected (lower panel) per cell are shown as violin plots for SUM-seq and other methods (dsci-
ATAC-seq[11]: GSM3507387; SHARE-seq[14]: GSM4156590 (ATAC), GSM4156602 and GSM4156603
(RNA); Paired-seq[15]: GSM3737488 (ATAC), GSM3737489 (RNA); scifi-RNA-seq[9]: GSM5151362). The

median values are shown on top. **f,** Distribution of accessible DNA fragments in peaks per cell and the
TSS score per cell (upper panels) as well as UMIs and genes detected per cell (lower panels) are shown
as violin plots, split by the number of nuclei in a droplet. The number of droplets with N nuclei
encapsulated is shown on the top. **g,** A representative genome browser view (*chr12*: *52,334,173 -*
55,349,410) of SUM-seq data is shown for K562 as aggregated transcript and ATAC fragment
mappings. For comparison, K562 ATAC-seq and RNA-seq data tracks were retrieved from the
ENCODE database (tracks labeled in green; GSE86660[16] (RNA), GSE170214[16] (ATAC)). The bottom
panel shows binarised accessibility for 200 randomly selected K562 nuclei at single-cell resolution.
Boxplots in **e**,**f**: center line represents median; lower and upper hinges represent the 25th and 75th
quartiles respectively.

SUM-seq recapitulates regulatory dynamics during M1/M2 macrophage
polarization

SUM-seq is a suitable method for unraveling gene regulatory networks during cellular
differentiation. As a model system for cellular polarization, we used macrophages, innate
immune cells that can polarize towards a pro-inflammatory M1 or an anti-inflammatory M2
state depending on microenvironmental signaling. Despite the inherent complexity and
heterogeneity observed *in vivo*, the M1/2 dichotomy serves as a useful framework for
elucidating major transcriptional events and regulatory elements governing macrophage
polarization. Previous research has determined TFs including STAT1, IRF5, and NF-κB as
key factors involved in M1 polarization, and STAT6, PPARγ, and IRF4 in M2 polarization[21-23].
However, the complex regulatory networks orchestrating macrophage polarization over time
remain elusive.
185
To unravel these intricate regulatory networks, we applied SUM-seq to profile hiPSC-derived
macrophages across a polarization time course from the naive M0-state to the M1- and M2-
states. We stimulated the M0 macrophages with LPS and IFN-γ to induce M1 polarization or
IL-4 to induce M2 polarization. To discern early and sustained responses at chromatin
accessibility and gene expression levels, we collected samples at five time points along the
two polarization trajectories; prior to stimulation (M0) and at 1-hour, 6-hour, 10-hour, and 24-
hour intervals, each sampled in duplicates totaling 18 samples **(Figure 2a)**. Leveraging the
overloading capacity of SUM-seq, we loaded 150,000 nuclei into a single microfluidic channel
of the 10x Chromium system, resulting in multiomic data from 51,750 high-quality nuclei evenly
distributed across samples (**Figure 2b; Extended Data Figure 3**). For the snATAC, nuclei
passing the QC filters (**Methods**) exhibited an average of 11,900 unique fragments, a TSS
score of 8, and 40 % of reads in peaks. For the snRNA-seq readout, nuclei displayed on
average 407 UMIs and 342 genes (**Extended Data Figure 3a-c**). The lower average UMI and
gene counts per nucleus in the macrophage experiment, in comparison to the mixed-species
experiment, are a consequence of the omission of PEG during reverse transcription due to a
technical issue (**Methods**). However, the high number of nuclei that passed the quality filter
enabled effective downstream analyses.
203

204
Figure 2. Integrated chromatin accessibility and gene expression data at single-nucleus
resolution via SUM-seq characterizes differentiation trajectories along M1/M2 macrophage
polarizations
a, Schematic overview of the macrophage polarization experiment. hiPSC-derived macrophages were
stimulated with LPS and IFN-γ (M1) or IL-4 (M2). Nuclei were fixed with glyoxal and collected for SUM-
seq at 0h, 1h, 6h, 10h, and 24h (*n*=2 per time point). **b,c,** Weighted nearest neighbor (WNN) UMAP
projection of integrated SUM-seq data of macrophage polarization. Cells are annotated and labeled
according to their sample index (**b**), AUC score of M1 signature genes (**c** left panel) and AUC score of
M2 signature genes (**c** right panel; Supplementary Table 2). **d,e,** Distributions of cells from each time
point across three MOFA factor weights associated with M1 polarization (**d;** early response, sustained
response, late response) and M2 polarization (**e**; early response, sustained response I and II). Dotted
arrows indicate the direction of the time-resolved response. **f,g,** Gene set enrichment analysis (GSEA)
(Methods) for M1 (**f**) and M2 (**g**) polarization factors are shown as heatmaps. **h,** Motif activity for TFs
associated with M1 early response (Methods) across M0 and M1-1h cells sorted by M1 early response
factor weights (top). Motif activity for TFs associated with M1 sustained response across all M0 and M1
cells sorted by sustained factor weights (bottom). STAT1 and STAT1.1 represent the homodimer and
heterodimer motifs of STAT1 respectively. **i,** Motif activity for TFs associated with M2 early response
across M0 and M2-1h cells sorted by M2 early response factor weights (top). TF motif activity for TFs

associated with M2 sustained response II across all M0 and M2 cells sorted by sustained factor II
weights (bottom).
225

After preprocessing the data (**Methods**), we obtained a joint visualization of the two modalities
by constructing a weighted nearest-neighbor (WNN) graph, which was further used to
determine a uniform manifold approximation and projection (UMAP) (**Methods**). The resulting
UMAP separated cells along the M1 and M2 polarization trajectories (**Figure 2b**). By
calculating M1 and M2 scores from the expression of literature-based marker genes
(**Supplementary Table 2; Methods**), and mapping them onto the UMAP, we observed
increased M1 and M2 scores during the respective polarization trajectories (**Figure 2c**),
confirming the validity of our experimental setup. M2 cells were not clearly distinguished from
the M0 state and underwent only gradual changes throughout the time course. In contrast,
during M1 polarization, we observed a clear separation of cells from the M0 state, with a
substantial shift between the 1-hour and 6-hour time points. This separation of cells from
different timepoints along the M1 and M2 trajectories underscores the capacity of our system
to effectively capture distinct early and late polarization stages.
239

To delineate the key features that influence gene expression and chromatin accessibility
during macrophage M1/M2 polarization, we performed Multi-Omics Factor Analysis (MOFA)[24]
to establish a shared low-dimensional representation for the snATAC- and snRNA-seq data
(**Extended Data Figure 3d,e**). Investigating the association of the MOFA factors with
metadata, we pinpointed three factors associated with M1 polarization, explaining the early
response (Factor 4), sustained response (Factor 1), and late response (Factor 7) (**Figure 2d,**
Extended Data Figure 3d). Notably, the early response was predominantly driven by
chromatin accessibility, while the sustained and late responses were influenced by both gene
expression and chromatin accessibility (**Extended Data Figure 3d**). For M2 polarization, we
identified an early response (Factor 7) and two distinct sustained response factors (Factors 3
and 5) (**Figure 2e**). For functional interpretation of the MOFA factor associated genes, we
performed gene set enrichment analysis (**Supplementary Table 3, Methods**). Genes
associated with the early M1 response were enriched in IFN-γ and cytokine signaling (**Figure**
2f). IFN signaling was also enriched in genes associated with the sustained response (Factor
1), along with general immune system processes, IL-1 signaling, and metabolic terms, in line
with metabolic reprogramming previously observed in M1 macrophages[25]. Genes associated
with the M1 late response were predominantly enriched for terms related to antigen (cross-)
presentation and RHO signaling (**Figure 2f**). The early M2 polarization factor was enriched for
IL-4 signaling, while the two M2 sustained factors were enriched in biological oxidation - likely
signifying the switch in metabolism of M2 macrophages[26], and non-canonical NF-κB-signaling
and Notch signaling (**Figure 2g**).

SUM-seq uncovers key TF motif activity dynamics along the M1/M2 polarization

Next, we investigated which TFs underlie the chromatin remodeling during M1 polarization.
Using chromVAR, we quantified TF motif accessibility variation[27] and use this as a proxy for
the activity of a TF binding a particular motif (here referred to as TF motif activity). We selected
the top 10 % most variable TF motifs and filtered for those enriched in peaksets associated
with either the early, the sustained, or the late response of M1 (**Extended Data Figure 4a**),
resulting in 103 unique TFs (139 motifs; **Supplementary Table 4**).
268

When plotting the chromatin-based TF motif activity in cells from M0 and M1-1h along the
early response (Factor 4) trajectory, we observed increasing motif activity for prototypical M1
TFs, such as STAT1 and IRF5[22,23], followed by NF-κB (NFKB1, NFKB2, RELA, REL, TF65;
Figure 2h). Furthermore, we observed a transient upregulation of the AP-1 complex motifs
(JUN, FOS, etc.), indicative of a cellular activation state[28]. Meanwhile, the myeloid
differentiation factors of the ETS family, including SPI1 (PU.1)[29], were active in M0
macrophages and slightly decreased their activity upon polarization. To investigate the M1
response at later time points, we ordered all M1 cells along the sustained response factor
(Factor 1; **Figure 2h**). This revealed a switch-like increase in motif activity for many
prototypical M1 TFs, including STATs and IRFs, such as IRF5. STAT1 can function as a
homodimer (also called IFN-γ activated factor (GAF)) or can form, together with STAT2 and
IRF9, the interferon-stimulated gene factor-3 (ISGF3) complex[30]. STAT1 has two motifs in the
HOCOMOCO database: one motif is bound by the STAT1 homodimer
(STAT1.H12INVIVO.0.P.B), and the other motif (STAT1.H12INVIVO.1.P.B) is highly similar to
the motif of STAT2. In the JASPAR database[31], this second motif is classified as the interaction
of STAT1 and STAT2. As such, herein, we refer to the second motif as the STAT1 heterodimer
motif. Notably, we observed a steady drop in STAT1 homodimer motif activity over time, which
coincided with a marked increase of its heterodimer motif activity as well as the motifs of
STAT2 and IRF9 (**Figure 2h**). These observations highlight the role of STAT1 homodimer in
initiating chromatin remodeling in response to IFN-γ. Meanwhile, the sustained response to
IFN-γ is maintained by partnering with STAT2 and IRF9 in the ISGF3 complex, the driver of
type-I IFN response[30]. Additional factors like AP-1 and ETS family members also exhibited
switch-like activity patterns, increasing and decreasing with time, respectively. In contrast to
the switch-like dynamic of the IRFs, STATs, AP-1 and ETS, the NFKB2 motif activity presented
a more gradual increase along the M1 polarization. In the late response phase we observed
only subtle changes in TF motif activity, including a slight increase in NFY motifs and a slight
decrease in CEBP motifs (**Extended Data Figure 4c**). This pattern suggests that TF-driven
gene regulation is stable between 6 and 24 hours of M1 polarization.
297
A similar analysis of TFs along the M2 polarization focused on 93 TFs (121 motifs) (**Extended
Data Figure 4b, Supplementary Table 4**). Overall, we observed less striking dynamics along
the M2 polarization trajectories compared to the M1 response. The most notable pattern was
observed for the IL-4-responsive STAT6[32], which showed increasing motif activity along the
early M2 response trajectory (**Figure 2i**). The sustained response patterns were mostly
characterized by a steady increase in motif activity for ETS family members, most notably
SPIB[33], ETV transcription factors, such as ETV3 and ETV6[34], and increased activity of NFY
factors (**Figure 2i, Extended Data Figure 4d**). Conversely, SNAIL family member motif
activity decreased along the sustained response. Likewise, accessibility of CTCF, a protein
shaping chromatin structure[35], decreased upon M2 polarization, suggesting a global rewiring
of chromatin architecture.
309
310   Overall, we observed strong remodeling of the chromatin landscape along the macrophage
311   polarization trajectories, much of it driven by specific sets of TFs.

Gene regulatory network analysis reveals sustained response by STAT1/STAT2/IRF9 in M1 macrophages

We next aimed to understand how the TF-driven chromatin remodeling leads to activation of specific gene expression programs. To that end, we constructed an enhancer-mediated gene regulatory network (eGRN) using GRaNIE[36] (**Supplementary Table 5; Methods**). Out of the 103 TFs identified in association with M1 polarization, we retrieved 44 within the eGRN, either as a TF itself or as a target gene of the M1-TFs (**Figure 3a; Extended Data Figure 5b**). The eGRN shows that the most connected TFs share many of their target genes. For example, IRF1 and IRF8, connected to 1002 and 804 genes respectively, share 728 of their regulon genes.

We further resolved the hierarchy of a TF's activity on chromatin accessibility and their effect on inducing transcription by comparing motif and regulon activity similarity along the M1 immediate response factor (**Methods**). For the majority of TFs, we found a positive correlation between their motif activity and the expression of their target genes derived from the eGRN (**Extended Data Figure 6a**). Some evident exceptions to this were STAT1, STAT3, NFKB1, NFKB2 and RELB. For NFKB1, we observed a slightly delayed expression of its regulon despite the gradual increase in NFKB1 motif activity from an early time point (**Figure 3b**). Additionally, we found that STAT1 regulon activity was discordant with the chromatin activity for the STAT1 homodimer motif, but it closely tracked the STAT1 heterodimer motif activity (**Figure 3b,c**).

333
Figure 3. Gene regulatory networks inferred from SUM-seq data in macrophages coupled with
genetic evidence reveals TF and regulon hierarchy, linking TFs to immune traits. a, Gene
regulatory network visualization of TFs and their assigned target genes, including only TFs with at least
4 target genes. **b,** TF motif activity (chromVAR Z-score) and regulon activity (AUC cell score) for STAT1
(top) and NFKB1 (bottom) motifs along the M1 sustained response factor. Cells are ranked by their
respective value for activity measures, lines show best fit (generalized additive model). **c,** Scatter plots
of the correlation between the STAT1 regulon activity and STAT1 homodimer (left) and STAT1
heterodimer (right) TF motif activity (chromVAR Z-score). Points represent cells and are coloured by
their experimental time point. **d,** Scatter plots of each cell across the M1 macrophage polarization
showing the correlation between STAT1 and STAT2 regulons (left) and STAT1 homodimer and STAT2
TF motif activity (right). Points represent cells and are coloured by their experimental time point. **e,**
Schematic of the type II IFN-associated STAT1 homodimer (known as GAF) response, representing
the early response, and the type I IFN-associated STAT1/STAT2/IRF9 (known as ISGF3 complex),

representing the sustained response. IFN-γ binding to its receptor initiates an early response
characterized by the phosphorylation and subsequent dimerization of STAT1, forming GAF. The GAF-
homodimer then translocates into the nucleus and binds the GAS motif, activating the expression of 8
downstream targets identified in our eGRN. As a secondary response, STAT1, STAT2 and IRF9
undergo phosphorylation and activation, forming the ISGF3 complex. The ISGF3 complex binds the
ISRE motif, potentially activating the expression of 1008 downstream genes identified in our eGRN. **f,**
Identification of diseases and traits associated with genetic variants enriched in open regions of all GRN
peaks and a highly connected regulon in the eGRN (the STAT1/STAT2/IRF9 - putative ISGF3 complex)
using linkage disequilibrium score regression (LDSC). All open regions in macrophages are used as
background. **g,** AUC cell scores for disease modules derived by intersection of putative genome-wide
SNPs for the top enriched diseases with the putative ISGF3 complex regulon (union of STAT1, STAT2,
IRF9 gene targets) (**Methods**), compared to AUC cell scores for all ISGF3 regulon genes along the M1
sustained response factor. **h,** eGRN peak-gene interactions for the CD40 intronic peak intersecting RA-
associated SNP rs4810485 (top), zooming in to the CD40 region with the TFs binding to this peak
(insert). M0 and M1 cell aggregate ATAC-seq tracks split by timepoint highlighting the CD40 intronic
peak (bottom).

While the STAT1 homodimer motif activity was strongest at 1h post-stimulation and decreased
from 6h onwards, the STAT1 heterodimer motif, along with STAT2 and IRF9 motif activities,
exhibited a gradual increase over time (**Figure 3d, Extended Data Figure 6b,c**). This pattern
was also reflected by the STAT1, STAT2 and IRF9 regulons. These three regulons shared
601 genes and their activities were highly correlated, providing evidence that the ISGF3
complex was captured by the eGRN (**Figure 3d**; **Extended Data Figure 6d-f**). In contrast, we
identified only 8 target genes connected to the homodimer motif, all of which were also
connected with the heterodimer motif (**Supplementary Table 6**). After IFN-γ/LPS stimulation,
we first observe increased activity for the STAT1 homodimer and other early response TFs
(**Figure 2h**). This early response potentially leads to endogenous type I IFN production, which
in turn activates the ISGF3 complex, explaining the delayed response of ISGF3 as compared
to the STAT1 homodimer activity (**Figure 3e**).

Given that the activity of many TFs, including STAT1, is regulated by phosphorylation, we
made use of published phosphoproteomic data on IFN-γ/LPS and IL-4 stimulated THP1-
derived macrophages to validate our findings[37]. We observed that upon IFN-γ/LPS stimulation,
STAT1 becomes phosphorylated at sites T699 and Y701 almost immediately, followed by a
gradual decrease over time (**Extended Data Figure 6g**). Phosphorylation of these sites is
known to be induced by IFN signaling and triggers STAT1 accumulation in the nucleus and
activation of DNA binding activity[38,39]. This pattern thus recapitulates the early STAT1
homodimer response we observed on chromatin. Moreover, the observed gradual increase in
STAT2 and IRF9 activity based on ATAC-seq and RNA-seq data is corroborated by a delayed
but sustained increase in the phosphorylation of IRF9 at sites S131 and S253 (**Extended Data**
Figure 6h). S253 is known to be induced by IFN-β (a type I IFN) and may play a role in
regulating the expression of interferon-stimulated genes and interactions with STAT2[40].

In summary, integrating chromatin accessibility and gene expression data at single-cell
resolution over a time course, we characterized the M1 and M2 polarization trajectories. This
highlights SUM-seq's capability to provide detailed insights into cellular states and TF
activities. Using both ATAC and RNA modalities along a time course enabled us to dissect the
regulatory layers of M1 polarization, leading to the identification of a putative hierarchy of IFN-
driven responses. Specifically, we observed a marked shift in STAT1-mediated regulation,

from its homodimer-driven chromatin remodeling during the early M1 polarization, to an
ISGF3-driven response reflected in both accessibility and transcription at later time points.
This switch is in line with previous reports on the timing of IFN stimulations[41,42] (**Figure 3e**).

GWAS enrichment analyses of the SUM-seq identified eGRN

To further investigate the role of the macrophage regulatory network in disease, we tested
whether the chromatin accessible regions linked to regulon genes in the network were
enriched for heritability of specific diseases using linkage disequilibrium score regression
(LDSC)[43].
404
Overall, we found all open regions in our macrophage dataset strongly enriched for white blood
cell count traits and autoimmune diseases (**Extended Data Figure 7a**). Therefore, we used
all open regions as background, to ensure the disease enrichments are not driven by the cell
type only. The eGRN peaks were enriched for specific immune-related traits including
inflammatory bowel disease (IBD), two subtypes of IBD (ulcerative colitis (UC) and Crohn's
disease (CD)), hayfever and multiple sclerosis (**Figure 3f**). Mapping the putative genome-wide
SNPs (p < 10^{-6}) for the top heritability-enriched diseases and traits (p < 0.1), we found 112
unique SNPs that directly overlap with the peaks in the eGRN (**Supplementary Table 7**). For
UC, the most significant overlapping SNP was rs153109, located in an intron of the Interleukin-
27 (IL-27) gene (**Extended Data Figure 7b**). This SNP has also been associated with CD and
with IBD. These diseases are characterized by chronic inflammation of the digestive tract. IL-
27 is mainly produced by antigen presenting cells including macrophages, and has been
implicated in the regulation of the immunological response in IBD[44]. As such, it has been
proposed as a potential new drug target for IBD[45]. In our eGRN, this IL-27 peak is connected
to and putatively regulated by NFKB1. Interestingly, a NFKB1 knockout in a THP1 monocyte
cell line showed downregulation of IL-27, providing further evidence for its regulatory
function[46].
422
To zoom in further, we tested heritability enrichment for the peaks in the putative ISGF3
regulon (union of peaks linked to STAT1, STAT2 and IRF9). Although this subset consisted of
only half as many peaks as the full eGRN (1673 vs. 3217), the enrichments for the autoimmune
diseases remained significant (**Figure 3f**). This suggests that STAT1, STAT2 and IRF9 play
an important role in autoimmune diseases involving macrophages. Indeed, disease SNPs
mapping specifically to the ISGF3 regulon resulted in prioritizing 19 genes for IBD, 16 for CD,
and 37 for eosinophil counts (**Supplementary Table 8; Methods**). We then calculated the
expression activity for these disease genes (**Methods**) and projected them along the M1
sustained response factor. The disease gene sets (IBD, CD) showed a stronger increase in
activity over time than the full ISGF3 regulon and "eosinophil count" gene set, suggesting the
polarized M1 state is particularly important for these diseases and traits (**Figure 3g**). Another
example bridging the M1 polarization and genetics is rs4810485, which overlaps with the
ISGF3 regulon. This SNP, located in the intron of CD40, is associated with IBD, CD and UC
and the CD40 intronic peak is linked to CD40, PLTP, NEURL2 and SLC35C2 in our eGRN
(**Figure 3h**). Previous work has linked this SNP to CD40 as an expression, protein and splice
quantitative trait locus (e-, p-, sQTL) in blood and monocyte-specific datasets, as well as to
PLTP as a blood eQTL[47] (**Supplementary Table 9)**. CD40 is a cell surface receptor expressed
by antigen presenting cells including macrophages, that contributes to the activation of T-cells
through its interaction with CD40 ligand (CD40-L). Given this role, CD40 is likely involved in

the pathogenesis of IBD and other autoimmune diseases[48]. CD40 expression can be induced
by IFN-γ stimulation in macrophages, a process thought to be involved in the initiation of
chronic inflammation in IBD[49,50].
445
In summary, the integration of our high-resolution single-cell multimodal time course data on
M1 polarization with genetic disease evidence revealed molecular disease mechanisms and
suggests novel candidate pathways that will require validation in future studies.

Discussion

SUM-seq is a single-cell multiomic method that allows highly multiplexed and scalable
experimental setups to investigate the molecular underpinnings of gene regulation via profiling
of chromatin accessibility and gene expression from the same cell. In this study, we applied
SUM-seq to examine macrophage polarization, demonstrating how the multimodal data from
a single multiplexed experiment can unravel the molecular underpinnings of TF-driven disease
mechanisms, integrating TF response dynamics with genetic evidence from GWAS.
456
We benchmarked SUM-seq, showcasing its ability to effectively capture both chromatin
accessibility and gene expression at the singel-cell level, outperforming previously published
scalable high-throughput methods. Notably, we demonstrated its capability to generate high-
quality data from both fresh and cryopreserved samples, making it adaptable for extensive
atlas studies, where sample collection can span a long time period and requires multi-center
efforts. While SUM-seq does not reach the gene expression complexity of lower throughput
droplet-based single-cell multiomic methods, it provides matching chromatin accessibility
complexity paired with significant improvement in cost-efficient scalability. We anticipate that
through minor optimisations 1) gene expression complexity will further improve and 2) SUM-
seq can expand to incorporate surface proteomic readouts by use of single cells instead of
single nuclei[51]. Furthermore, it should be noted that although the current format measures
chromatin accessibility and gene expression, the strategy can be adopted for the
measurement of other omic layers such as TF binding and histone modifications (Cut&Tag[52]),
DNA methylation (sci-MET[53]), and whole genome sequencing (s3-WGS[54]).
471
We applied SUM-seq to hiPSC-derived macrophages in a time-course of M1 and M2
polarization, as a prototypical model of cellular polarization. By analyzing the activity of TF
motifs on the chromatin versus the expression of associated TF target genes, we observed a
hierarchical activation of a TF cascade upon IFN-γ/LPS stimulation of M0 macrophages.
Specifically, we identified TFs that clearly act in a stepwise manner (i.e. STAT1), whereas
others exhibited a continuous activation profile (i.e. NFKB1). We could recapitulate the
hierarchy of STAT1 homodimer activity preceding STAT1/STAT2/IRF9 activity. A potential
explanation for the delay in the activity of the NFKB regulon with respect to its motif activity is
the previously reported observation that enhancers of NF-κB response genes are primed by
NF-κB but their expression additionally requires activity of the IFN-I-induced ISGF3 complex[55].
482
Genes with genetic links to disease have been shown to be better drug targets[56]. Yet for the
majority of disease-associated genetic variants, we lack molecular mechanisms and it is often
unclear in which cell type or disease state a variant exerts its effect. Most disease-associated
common genetic variants lie in non-coding regulatory elements that are presumably regulated
by TFs. Indeed, TFs are often identified as main drivers of disease-relevant phenotypes in

genetic screens[57]. Our study emphasizes that understanding disease mechanisms requires
integrating genetic evidence with disease-specific gene-regulatory dynamics. SUM-seq was
instrumental in pinpointing gene-regulatory dynamics and linking them to genetic disease
evidence in a single experiment. Specifically, profiling the macrophage polarization response
in a single SUM-seq experiment, followed by integrative analysis of the eGRN and publicly
available summary statistics from GWAS, identified several known disease mechanisms, as
well as novel associations. For example, our findings reinforce the link between the ISGF3
complex, CD40 expression, and IBD. Indeed, the role of macrophages in maintaining intestinal
immune homeostasis makes them an appealing target to develop new therapeutic strategies
for IBD[58].
498
In conclusion, our study showcases the effective implementation of SUM-seq in the context of
a time course experiment. Importantly, our findings suggest broad applicability of this method
to diverse experimental settings that demand gene regulatory analyses at single-cell
resolution. Specifically, SUM-seq is a promising tool for arrayed CRISPR, drug or perturbation
screens, and large-scale atlas projects, underscoring its potential as a versatile and powerful
technique across a spectrum of biological studies. Overall, we envision SUM-seq as an easily
adoptable, scalable method for projects requiring multiomic single-cell profiling up to millions
of cells from hundreds of samples.

Methods

Cell culture

Human myelogenous leukemia cells (K562) were cultured in RPMI 1640 medium (Gibco,
catalog no. 11875093) supplemented with 10 % FBS (Gibco, catalog no. 10270-106), 100
U/ml penicillin/streptomycin (PenStrep, Gibco, catalog no. 15140122) and 1x non-essential
amino acids (Gibco, catalog no. 11140050) at 37°C with 5 % CO_2. Mouse fibroblast cells (NIH-
3T3) were cultured in DMEM medium (high glucose, Gibco, catalog no. 11965092)
supplemented with 10 % FBS, 100 U/ml penicillin/streptomycin and 1x non-essential amino
acids at 37 °C with 5 % CO_2.

Macrophage differentiation

hiPSCs were cultured in Essential 8 medium (Gibco, catalog no. A1517001) in dishes coated
with vitronectin (ThermoFisher, catalog no. A14700). hiPSCs were differentiated to
macrophages as previously described by van Wilgenburg et al[59]. In brief, 4 million hiPSCs
were resuspended in EB medium (Essential 8 medium (Gibco, catalog no. A1517001), 50
ng/ml Recombinant Human BMP4 (Peprotech, catalog no. 120-05ET), 50 ng/ml Recombinant
Human VEGF (Peprotech, catalog no. 100-20), 50 ng/ml Recombinant Human SCF
(Peprotech, catalog no. 300-07)) with 10 µM Y-27632 (AbCam Biochemicals, catalog no.
ab120129), seeded in 400-microwell Aggrewells (Stemcell Technologies, catalog no. 34450)
and centrifuged at 100 x g for 3 min to evenly distribute the cells in microwells. 75 % of the
medium was replaced by fresh EB medium on the next 2 consecutive days. On the third day,
EBs were transferred to a low-attachment plate (Sigma-Aldrich, catalog no. CLS3471-24EA)
and cultured for two additional days with minimal disruption. On the fifth day, EBs were
transferred to the final format of choice and left undisturbed for 1 week in factory media (X-

VIVO^{TM}-15 (Lonza, catalog no. BE02-060F) 1 % GlutaMax (Thermo Scientific, catalog no.
35050061), 1 % PenStrep (Gibco, catalog no. 15140122), 50 µM ß-mercaptoethanol, 50 µg/ml
Normocin (Invivogen, catalog no. ant-nr-05) and 100 ng/ml Recombinant Human M-CSF
(Peprotech, catalog no. 300-25), 25 ng/ml Recombinant Human IL-3 (Peprotech, catalog no.
200-03)) to allow them to attach. Fresh media was added weekly until macrophage precursor
production started around week 4, as identified by the presence of large suspension cells with
spherical morphology. From that point on, factories were maintained for a total of 8-10 weeks.
Precursors were harvested weekly, taking approximately 1/4th of the total volume containing
precursors and replacing it with fresh media. For the terminal differentiation of precursors to
macrophages, harvested precursors were resuspended in macrophage media (*X-VIVO*^{TM}-15,
1 % GlutaMax, 1 % PenStrep, Recombinant Human M-CSF) and plated in the desired format
at a density of 160.000 cells/cm^2 for 7 days.

Polarization to M1 or M2 macrophages

At day 7, mature macrophages were polarized to M1 or M2. For M1 polarization, macrophages
were cultured in macrophage media supplemented with 20 ng/ml human recombinant IFN-γ
(PeproTech, catalog no. 300-02) and 25 ng/ml LPS (Invivogen, catalog no. tlrl-3pelps). For
M2 polarization macrophages were cultured in macrophage media supplemented with 20
ng/ml recombinant human IL-4 (PeproTech, catalog no. 200-04). Cells were harvested after
24, 10, 6 and 1h, including an undifferentiated control (M0) in which medium was replaced by
fresh macrophage medium.

Nuclei preparation

For the species mixing experiments, mouse NIH-3T3 cells were washed once with PBS and
dissociated with Accutase (Stem Cell Technologies, catalog no. 07920). Two million NIH-3T3
and K562 cells were collected and fixed in a 3 % glyoxal solution (40 % glyoxal (Merck, catalog
no. 128465), 0.75 % acetic acid; adjusted to pH 5 by addition of 1 M NaOH) for 7 min at room
temperature. For cryopreservation, cells were washed after fixation with RSB-1%BSA-RI (10
mM Tris-HCl pH 7.5, 10 mM NaCl, 3 mM MgCl$_2$, 1 mM DTT, 1 % BSA, 20 µg/ml in-house
produced RNasin (referred to RNasin hereafter)) and slowly frozen (Freezing buffer: 50 mM
Tris-HCl pH 7.5, 5 mM MgAc, 0.1 mM EDTA and 25 % glycerol). On the experiment day, fresh
cells were processed as described above, while cryopreserved cells were thawed slowly on
ice and washed with RSB-1%BSA-RI. Nuclei were extracted with 1x lysis buffer (10 mM Tris-
HCl pH 7.5, 10 mM NaCl, 3 mM MgCl2, 0.1 % Tween20, 20 µg/ml RNasin, 1 mM DTT, 1 %
BSA, 0.025 % IGEPAL CA-630 and 0.01 % Digitonin) incubating samples for 5 min on ice.
Next, nuclei were washed with wash buffer (10 mM Tris-HCl pH 7.5, 10 mM NaCl, 3 mM MgCl$_2$,
0.1 % Tween20, 20 µg/ml RNasin, 1 mM DTT, 1 % BSA) and filtered through a 40 µm cell
strainer (Falcon, catalog no. 352340).
566
567 After the desired polarization time, macrophages were harvested by replacing the medium
568 with ice-cold PBS-1%BSA and 2 mM EDTA and pipetting. Nuclei were extracted by
569 resuspending the cell pellet in 1x lysis buffer for 4 min on ice and washed with wash buffer.
570 To maintain nuclei integrity, nuclei were fixed with a 3 % glyoxal solution for 7 min at room
571 temperature and washed with RSB-1%BSA-RI.

Transposome generation

Tn5 was produced in-house according to a previously described protocol with minor modifications. In short, the pETM11-Sumo3-Tn5(E54K,L372P) plasmid was transformed into *Escherichia coli* BL21(DE3) codon + RIL cells (Stratagene). Cells were grown in TB-FB media (1.2 % Bacto-Tryptone, 2.4 % yeast-extract, 0.4 % glycerol, 17 mM KH_2PO_4, 72 mM K_2HPO_4, 1.5 % Lactose, 0.05 % Glucose, 2 mM $MgSO_4$) supplemented with kanamycin and chloramphenicol at 37 °C until OD600 ~0.5. The temperature was then lowered to 18 °C and cells were grown overnight, wherafter cells were harvested by centrifugation. The cell pellet was resuspended in running buffer (20 mM HEPES-NaOH pH 7.2, 800 mM NaCl, 5 mM imidazole, 1 mM EDTA, 2 mM DTT, and 10 % glycerol) supplemented with cOmplete protease inhibitors (Roche) and lysed using a Microfluidizer. Polyethyleneimine (PEI) pH 7.2 was added dropwise to a final concentration of 0.5 % to remove nucleic acids. The cleared lysate was loaded onto a cOmplete His-tag purification column (Roche) and the His6-Sumo3-Tn5 was eluted with a running buffer containing 300 mM imidazole. To remove the fusion tag, His6-tagged SenP2 protease was added to the elution fractions. The sample was digested overnight at 4 °C while being dialyzed back to the running buffer. The next morning, the dialyzed sample was loaded again onto a cOmplete His-tag purification column (Roche), and the untagged Tn5 was collected in the flow-through. The concentrated flow through was then loaded onto a HiLoad Superdex200 16/600 pg column (GE Healthcare) equilibrated with 50 mM Tris pH 7.5, 800 mM NaCl, 0.2 mM EDTA, 2 mM DTT, and 10 % glycerol. Elution fractions corresponding to the Tn5 dimer peak were pooled, aliquoted, snap frozen in liquid nitrogen and stored at -80 °C. For transposome assembly, 100 µM of oligonucleotides with the Nextera Read1 sequence and 100 µM of oligonucleotides with a Read2-sample_index-spacer structure were annealed with 100 µM mosaic end-complement oligonucleotides with a 3' dideoxynucleotide end (ddc) at a 1:1:2 ratio by heating for 95 °C for 3 minutes followed by cooling down to 25 °C at a ramp rate of -1 °C/min. Annealed oligos were mixed with an equal volume of 100 % glycerol and stored at -20 °C until use. Finally, the annealed oligos were mixed with the in-house produced Tn5 (at 1 mg/ml in 50 mM Tris, 100 mM NaCl, 0.1 mM EDTA, 1 mM DTT, 0.1 % NP-40, and 50 % glycerol), at a 1:1 ratio and incubated for 30-60 min at room temperature. For the species-mixing experiment, annealed oligos were first diluted with H_2O at a 1:1 ratio before generating the transposomes. Assembled Tn5 was stored at -20 °C until use.

SUM-seq procedure

DNA transposition

For accessible DNA transposition, isolated and fixed nuclei were resuspended in a transposition mix containing 38.8 mM Tris-acetate, 77.6 mM potassium acetate, 11.8 mM magnesium acetate, and 18.8 % dimethylformamide supplemented with 0.005x protease inhibitor cocktail (Roche, catalog no. 11697498001), 0.4 U/µl SUPERaseIn (Thermo Fisher, catalog no. AM2694), 1.2 U/µl RiboLock (Thermo Fisher, catalog no. EO0382) and the barcoded Tn5 transposomes (1:10 (v/v); 938 µM) and incubated at 30 °C for 30 min with shaking at 400 rpm. The reaction was terminated by adding an equal volume of 2x stop buffer (10 mM Tris-HCl pH 7.5, 20 mM EDTA pH 8.0, 2 % BSA).

In situ reverse transcription

Transposed nuclei were washed 3 times with RSB-1%BSA-RI and resuspended in reverse
transcription mix containing 1x RT buffer (50 mM Tris-HCl pH 8.0, 75 mM NaCl, 3 mM $MgCl_2$,
10 mM DTT), 0.5 mM dNTPs, 1 U/µl Protector RNase inhibitor (Roche, catalog no.
3335402001), 0.2 U/µl SUPERaseIn (Thermo Fisher, catalog no. AM2694), 10 U/µl Maxima
H Minus RT (Thermo Fisher catalog no. EP0752) and 5 µM of the respective barcoded oligo-
dT primer. In the NIH-3T3/K562 species mix experiment, 12 % PEG 8000 (w/v; Jena
Bioscience, CSS-256) was included in the RT mix as a crowding agent. For the macrophage
experiment, crowding agent PEG was omitted from the reverse transcription reaction as we
observed formation of white precipitates, which hampered downstream steps of library
preparation. This issue was however later resolved by omitting BSA from pre- and post reverse
transcription washes. The samples were incubated at 50 °C for 10 min, followed by 3 thermal
cycles (8 °C for 12 sec, 15 °C for 45 sec, 20 °C for 45 sec, 30 °C for 30 sec, 42 °C for 2 min
and 50 °C for 3 min), and finally incubated at 50 °C for 5 min. Following the RT reaction, all
samples were pooled in RSB-RI and washed twice with RSB-1%BSA-RI.

cDNA/mRNA hybrid tagmentation

For transposome preparation, 100 µM of oligonucleotides with a Nextera Read1 sequence
overhang were annealed with 100 µM of the Tn5-mosaic end oligonucleotides at a 1:1 (v/v)
ratio by heating for 95 °C for 3 min followed by cooling down to 25 °C at a ramp rate of -1
°C/min. Annealed oligos were mixed with an equal volume of 100 % glycerol and stored at -
20 °C until use. For transposome assembly, annealed oligos were mixed with the in-house
produced Tn5 at a 1:1 ratio for 30 min at RT and stored at -20 °C until use.
636
The collected nuclei pool was counted, and the reaction volume was adjusted according to
the total number of nuclei (100 µl reaction for each 100,000 nuclei). Nuclei were resuspended
in the transposition mix containing 1x transposition buffer (38.8 mM Tris-acetate, 77.6 mM
potassium acetate, 11.8 mM magnesium acetate, and 18.8 % dimethylformamide and 0.005x
protease inhibitor cocktail and a 1:100 ratio (v/v) of assembled Tn5 (93.8 µM). The
tagmentation was performed on a shaking thermoblock at 37 °C for 30 min at 400 rpm. The
reaction was terminated by adding an equal volume of 2x stop buffer. Next, the sample pool
was washed twice with RSB-1%BSA.

Gap fill and ExoI

Sample pool was resuspended a mix containing 1x RT buffer (50 mM Tris-HCl pH 8.0, 75 mM
NaCl, 3 mM $MgCl_2$, 10 mM DTT), 0.5 mM dNTPs, 8 U/µl Maxima H Minus RT (Thermo Fisher
catalog no. EP0752) and 2 U/µl Thermolabile Exonuclease I (New England Biolabs, catalog
no. M0568S) and incubated for 15 min at 37 °C on a shaking thermoblock at 400 rpm.
Afterwards the sample was washed twice with RSB-1%BSA filtered through a 40 µm cell
strainer and count.

GEM generation

Complex samples with a high proportion of debris were cleaned prior loading of the 10x
Genomics chip using a glycerol-based buffer (50 mM Tris-HCl pH 7.5, 5 mM Mg-acetate, 0.1
mM EDTA, 25 % glycerol). For this, nuclei were resuspended in a glycerol-based buffer,
layered on top of an additional 500 µl glycerol-based buffer and centrifuged at 800 x g for 15

minutes. The supernatant was discarded, and the pellet was washed twice with RSB-1%BSA
and filtered through a 40 µM Flowmi cell strainer (Sigma-Aldrich, catalog no. BAH136800040).
The nuclei were counted and loaded into the Chromium Controller (10x Genomics) according
to 10x Genomics single-cell-ATAC standard protocol. 100,000 nuclei were loaded for the NIH-
3T3/K562 species-mixing experiment and 150,000 for the macrophages experiment. In brief,
nuclei in 1x Nuclei buffer (10x Genomics) were mixed with barcoding reagents (Barcoding
Reagent B, Reducing Reagent B and Barcoding Enzyme). For the NIH-3T3/K562 species mix
experiment, this reaction mix was supplemented with ~500 nM of a blocking oligonucleotide
to prevent Tn5-barcode hopping in multinucleated droplets. In brief, barcode hopping may
occur when residual barcoded Tn5 molecules, which contain the ATAC sample index, are
carried over to the droplets. During droplet barcoding, these residual transposomes are
disassembled, and barcoded oligonucleotides released. These oligonucleotides have
complementarity to sequence on ATAC fragments downstream of the sample index, which
can lead to fragment swapping within multinucleated droplets. To circumvent this, we
employed two complementary strategies: first, we added a blocking oligonucleotide in excess
to the droplet barcoding step with a 3'-inverted dT-end. These blocking oligonucleotides
compete with the carry-over Tn5-barcodes for binding of ATAC fragments. Their 3'-inverted
dT-modification prevents further amplification of these fragments, preserving the balance of
ATAC and RNA fragments and the correct sample barcodes in the library. Second, we
decreased the number of thermal amplification cycles during droplet barcoding from twelve to
four. By reducing the number of linear amplification cycles, the probability of a carryover Tn5-
barcode binding a fragment and erroneously barcoding and exponentially amplifying it
decreases. For Chip loading, the standard 10x workflow was followed. In brief, 70 µl of the
sample, 50 µl of Barcoding Gel Beads, 40 µl Partitioning Oil were loaded into the respective
channels. After the Chromium run, the droplet emulsion was transferred to 10x verified PCR
tubes (Eppendorf, catalog no. EP0030124359) and the cell barcoding reaction was run in the
thermocycler as follows: 5 min at 72 °C, 30 seconds at 98 °C, 12 cycles (98 °C 10 seconds,
59 °C 30 seconds and 72 °C 1 min). In the NIH-3T3/K562 species mix experiment, the droplet
emulsion was split into two fractions to test 12 versus 4 cycles of linear amplification, and 20
% of each fraction was further processed to limit the sequencing requirements. Sample
cleanup was performed following the standard 10x Genomics protocol with minor
modifications: 125 µl Recovery Reagent was added to sample and inverted 10-15 times. After
brief centrifugation, 125 µl of the pink oil phase was discarded and the aqueous phase was
mixed with a dynabead cleanup mastermix (182 µl Cleanup Buffer, 13 µl Dynabeads MyOne
SILANE and 5 µl Reducing Agent B (10x Genomics)) and incubated for 10 min at RT. Samples
were placed on a magnetic separator and washed 2 times with 80 % ethanol before eluting
them in 42 µl of elution buffer. Finally, samples were subjected to a 1.4x SPRI cleanup
(Beckman Coulter, catalog no. B23318).

Pre- and final library amplification

Following GEM generation, cell barcoding and cleanup, samples containing a mixture of
single-cell ATAC and RNA libraries were resuspended in the pre-amplification mastermix
containing 1x NEBNext HF 2x PCR Master Mix (NEB, catalog no. M0541L) and a combination
of three primers: partial-P5-PTO primer (common for both modalities), TruseqR2 (specific for
the RNA modality) and an ATAC-spacer primer (specific for ATAC modality) each at a final
concentration of 500 nM. Samples were incubated in the thermal cycle as follows: 98 °C 1 mi,
5 cycles of 98 °C for 20 sec, 63 °C for 20 sec and 72 °C for 20 sec), 72 °C for 1 min and then

subjected to a 1.2x SPRI cleanup eluting in 42 µl EB buffer (Qiagen). Each sample was split
into two fractions and each of them was further amplified with modality-specific primers: a
partial-P5-PTO/P7-index-spacer primer pair for the ATAC libraries and a partial-P5-PTO/P7-
index-TruseqR2 primer pair for the gene expression libraries. The PCR programme for both
library types was as follows: 98 °C for 1 min, 5 cycles of 98 °C for 20 sec, 63 °C for 20 sec
and 72 °C for 20 sec followed by a final extension step at 72 °C for 1 min. To estimate the
number of endpoint PCR cycles required for each library, 1/20th of the total library volume was
used to perform a qPCR reaction using library-specific primers, and the number of cycles
required to reach 50 % of saturation was calculated. Finalized ATAC libraries were subjected
to a 1.2x SPRI cleanup and RNA libraries to 0.72x SPRI cleanup. The concentration of the
libraries was measured using Qubit 1x dsDNA-HS (Thermo Scientific, catalog no. Q33230)
and the library fragment distribution was assessed using Bioanalyzer High Sensitivity DNA kit
(Agilent, catalog no. 5067-4626).

Sequencing

717   Libraries were sequenced on the NovaSeq6000 (Illumina) using a 100-cycle S2 Kit or
718   NextSeq2000 (Illumina) using a 100-cycle P3 kit (ATAC: Read 1: 55 cycles, Index 1: 11 cycles,
719   Index 2: 16 cycles, Read 2: 55 cycles, RNA: Read 1: 95 cycles, Index 1: 6 cycles, Index 2: 16
720   cycles, Read 2: 21 cycles).

Data analysis procedures

ATAC sequencing data preprocessing

Base calls were converted to fastq format and demultiplexed by i7, allowing for one mismatch,
using bcl-convert (v4.0.3). Demultiplexed reads were aligned to the hg38 genome and
fragment file generation was performed with chromap (v0.2.3)[60]. For the species mixture
experiment to evaluate collision rates, a combined hg38/mm10 reference was used and
fragments aligning to either genome were counted. Using the R package ArchR (v1.02)[61],
generated fragment counts for each cell were computed in 500 base genome bins. The basic
preprocessing pipeline for data processing until the ArchR object generation is available at:
https://git.embl.de/grp-zaugg/SUMseq (available upon publication).
731
Cell barcode filtering was performed based on number of unique fragments, TSS enrichment
score, and fraction of reads in promoter regions for each sample. Peak calling was performed
for each sample separately. In brief, MACS2 peak caller is used to identify peaks for each
group of cells, whereafter an iterative overlap peak merging procedure is done to derive a
consensus peakset between all samples. BigWig files for trace plots were generated with
ArchR::plotBrowserTrack normalizing for reads in TSS.

M1 and M2 signature score generation

To determine M1 and M2 signature genes, we obtained publicly available datasets and used
DESeq2[62] to identify genes that were differentially expressed in M1 and M2 macrophages
compared to M0 (adjusted p-value < 0.05 and fold change > 0). The following datasets were
utilized: GSE159112[63], where THP1-derived macrophages were stimulated with LPS and IFN-
γ or IL-4 and IL-13 for 24 hours to obtain M1 and M2 macrophages, respectively; and
GSE55536[64], involving iPSC-derived macrophages and human monocyte-derived

macrophages stimulated with IFN-γ and LPS for M1 polarization, or IL-4 for M2 polarization.
Genes identified as differentially expressed in both datasets were used as genesets for input
to the ranking based scoring-approach AUCell[65].

Transcription factor motif accessibility

749 Position weight matrices (PWMs) of TF-binding motifs were obtained from HOCOMOCO v12
750 (*in vivo* subcollection)[66]. Motif positions in the accessible chromatin regions were determined
751 using the function ArchR::addMotifAnnotations. Z-score of bias-corrected (GC content and
752 mean peak accessibility) per-cell motif accessibility was calculated with the function
753 ArchR::addDeviationsMatrix, which leverages the chromVAR framework[27].

RNA sequencing data preprocessing

Base calls were converted to fastq format using bcl2fastq (v2.20.0) or in the case of multiple
pooled libraries base calls were converted to fastq format and demultiplexed by library index
(i7) using bcl-convert (v4.0.3). The cell barcode (i5) was concatenated to the sample index
and UMI (Read2), whereafter reads were demultiplexed by sample index using Je (v2.0.RC),
allowing for two mismatches. Read alignment to Hg38 and gene expression matrix generation
for each demultiplexed sample was conducted with STARsolo. For the species mixing
experiment, a combined Hg38/mm10 reference was used and number of UMIs aligning to
each genome was determined for each cell to estimate collision rates. For each sample, cell
calling was either performed with EmptyDrops (v.1.16) or by determining inflection points on
a rank vs UMI plot. Additionally, cells were filtered by mitochondrial and ribosomal read
percentage. Feature by cell matrices were merged between samples, and finally features
found in a minimum of 10-25 cells were retained.

Comparison to other technologies

We compare the performance of SUM-seq to dsci-ATAC-seq[11]GSM3507387), SHARE-
seq[14]GSM4156590, RNA: GSM4156602 and GSM4156603), Paired-seq[15] (ATAC:
GSM3737488, RNA: GSM3737489), and scifi-RNA-seq[9] (GSM5151362) using cell line data.
We used NIH-3T3 data for SUM-seq, sequenced to approximately 30.000 reads/cell for each
modality. For other methods, we use authors' count matrices provided in the indicated
repositories.

Dimensionality reduction and modality co-embedding

775 The peak matrix was normalized using TF-IDF, whereafter singular value decomposition was
776 applied to derive a low-dimensional representation. The number of components considered
777 was based on the proportion of variance explained (range 30-50), discarding the first
778 component from downstream analysis due to high correlation with number of fragments per
779 cell.
780
781 The gene expression matrix was normalized by proportional fitting (cell depth normalization to
782 the mean cell depth), followed by logarithmic transformation (log(x + 0.5)), and another round
783 of proportional fitting[67]. Next, a low dimensional representation of the normalized gene
784 expression matrix was determined with PCA. The number of components considered was
785 based on the proportion of variance explained (range 30-50).
786

787 Derived low-dimensional representations for each modality was used as input to learn cell
788 modality weights, from which a WNN graph was constructed [68]. The WNN graph was further
789 used to define a common UMAP visualization of the data modalities.

Multiomic factor analysis

To generate a common low-dimensional latent space for ATAC and RNA in the M1 and M2
polarizations, we utilized the multiomics integration platform MOFA[24]. First, ATAC data was
collapsed from peaks to cis-regulatory topics with cisTopic (v0.3)[69], determining a suitable
number of topics based on the maximum on the second-derivative of likelihood curve and
minimum on the model perplexity curve. All topics were used as input for MOFA. For RNA,
top 4000 (default) most variable genes were used as input.
797
798 To determine polarization-associated latent factors, latent factors were correlated with
799 biological metadata (state (M0/M1/M2), timepoint) as well as technical metadata (UMI/cell,
800 fragments/cell). Factors with a high correlation against biological metadata, but not technical
801 metadata were chosen as factors for downstream analysis.

Gene set enrichment analysis

803 Gene set enrichment analysis was performed for Reactome database genesets (v59). For
804 every gene set G, significance is evaluated via a parametric *t*-test, where weights of the
805 foreground set (features in set G) are contrasted against a background set (weights of features
806 not in set G). *P* values were adjusted for multiple testing using the Benjamini–Hochberg
807 procedure. Enrichments with false discovery rate 5 % were considered significant.

Motif enrichment analysis

809 Motif enrichment analysis was performed with R package monaLisa (v1.8) against
810 HOCOMOCO v12 (*in vivo* subcollection) PWMs. Peaks within topics in the top and bottom 3
811 % quantile of feature weights for MOFA factors of interest were considered for analysis. TFs
812 with FDR 0.1 % and log2 enrichment in the top or bottom 10 % quantiles were considered
813 significant.

Gene regulatory network inference and transcription factor prioritization

We inferred an enhancer-based gene regulatory network (eGRN) using GRaNIE[36] v1.5.3 with
TFBS predictions based on HOCOMOCO v12[66] database (*in vivo* subset) that were generated
using PWMScan (see[70] for methodological details how these were produced). The single-cell
data from all time points and stimulations was first clustered using the smart local moving
(SLM) algorithm with a resolution of 1, giving rise to 31 clusters that have at least 25 cells.
These clusters largely align with the stimulations and time points (**Extended Data Figure 5a**).
We then calculated the mean RNA and ATAC values for each of these clusters as a
pseudobulk, and input these clusters as 'samples' to create a eGRN following the GRaNIE
single   cell   vignette   (https://grp-zaugg.embl-
community.io/GRaNIE/articles/GRaNIE_singleCell_eGRNs.html). We kept links if the TF-
peak connection had an FDR < 0.2 and the peak-gene connection had an FDR < 0.1, the
defaults in the GRaNIE package. We collapsed TF motif connections to the level of TFs to
construct TF regulons.

Residual analysis

To identify discrepancy and concordance between TF motif accessibility and regulon activity
scores, values were rank min-max normalized and the difference between the two was
computed, and is referred to as the residual value. AUCell was used to infer single-cell activity
scores for regulon geneset

GWAS integration

LDSC

To integrate GWAS with our eGRN, we performed stratified linkage disequilibrium score
regression (S-LDSC[43]). We added all open regions in all macrophage cells as background, on
top of the default baseline model that includes genic regions, enhancer regions, and
conserved regions. We then tested enriched heritability for all peaks that were part of the
STAT1/STAT2/IRF9 eGRN, all peaks in the eGRN, and all peaks in M1 and M2 cells only. We
included 8890 traits from UK biobank, FinnGen and several GWAS repositories as included in
the 22.10 release of Open Targets Genetics (ref). SNP heritability (h2) was calculated with
defaults from the LDSC codebase. We tested S-LDSC only if h2 was > 0.05, resulting in a list
of 1230 traits. We corrected the enrichment p-values for the number of peaksets tested in
each trait.

Gene modules and SNP overlap

For those traits that were enriched for heritability (adjusted p < 0.1), the SNPs associated to
the particular trait (suggestive GWAS p-value < 5*10^-6) were intersected with the peaks in
the STAT1/STAT2/IRF9, and full eGRNs (Supplementary Tables 7-8). We selected the genes
connected (in the STAT1/STAT2/IRF9 eGRN) to the peaks that overlapped a SNP to obtain
disease gene modules for CD, IBD and eosinophil counts.

Code availability

A Snakemake preprocessing pipeline for SUM-seq data is available in a public Git repository
(https://git.embl.de/grp-zaugg/SUMseq, available upon publication). All analysis source code
will be available on a separate Git repository.

Data availability

All data will be deposited to the NCBI GEO (GSE253165) or EGA databases.
The provided links can be utilized to privately access the datasets before their public release.
858
859 Data related to the species mixing experiment is available at GEO with accession number
860 GSE253165 (available upon publication)

862 Processed data related to the iPSC-derived macrophage polarization experiment is available
863 upon publication.

Acknowledgements

We thank the EMBL Genomic Core facility for help with sequencing, Protein Purification Core facility for Tn5 and RNasin production, Michael Snyder for providing hiPSCs, and Yakov Tsepilov and Xiangyu (Jack) Ge for providing access to the cleaned and harmonized database of GWAS summary statistics. We further thank members of the Zaugg and the Noh group for extensive discussions, Nadine Fernandez-Novel Marx for experimental support, and Josephine Brysting for providing cell lines.

A.C. was supported by an EIPOD4 Fellowship from the EC Horizon2020 MSCA (grant agreement number 847543). N.S. and J.B.Z. acknowledge funding from GSK through the EMBL-GSK collaboration framework (3000032294). N.H.S. and J.B.Z. acknowledge funding from the EMBL Infection Biology Transversal theme. K.D.P. was supported by SNSF (P2ZHP3_199669) and EMBO (ALTF538) Postdoctoral Fellowships. This work was supported by the Cariplo foundation grant, the GSK basic research fund, and the EMBL research fund (to K.M.N.). M.M. was supported by the Research council of Finland (grant number 347543), Sigrid Jusélius foundation, and Instrumentarium Science foundation. This project is co-funded by the European Union (ERC, EpiNicheAML,101044873) to J.B.Z.. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

Author Information

These authors contributed equally: Sara Lobato-Moreno, Umut Yildiz, and Annique Claringbould; position of S.L. and U.Y. was determined by coin flipping

Contributions
M.M. conceived the project with input from U.Y., K.M.N. and J.B.Z. M.M., S.L. and U.Y. established and implemented the protocol and conducted experiments with the assistance of H.G.B., K.D.P. and V.C. M.M. and A.C. performed computational analyses. C.A. and E.P.V. constructed the pre-processing pipeline. N.H.S. assisted in data interpretation and manuscript writing. S.L., U.Y., A.C., M.M., J.B.Z. and K.M.N wrote the manuscript with input from all co-authors. M.M., J.B.Z. and K.M.N. supervised and directed the project. J.B.Z., K.M.N and M.M. secured financial support.

Corresponding authors
Correspondence to Kyung-Min Noh, Mikael Marttinen or Judith B. Zaugg

Competing interests

The authors declare no competing interest.

Ethics declarations

hiPSCs used for macrophage generation were derived from peripheral blood mononuclear cells with institutional review board approval (Stanford University, reference numbers 29904, 30064). The use of hiPSCs was approved by the EMBL Research Ethics Committee.

906
907
908 **Table 1: Comparison between SUM-seq and other ATAC+RNA single cell methods**
909

Method	Microfluidics/ plate-based	Modalities	Multiplexing	Costs	No° cells	Sensitivity
SUM-seq	mf	ATAC/RNA	unlimited	$	ultra-high	high
10x multiome	mf	ATAC/RNA	no	$$$$	high	high
SNARE-seq2	mf	ATAC/RNA	no	$$	low	high
Paired-seq	mf	ATAC/RNA	no	$$	ultra-high	low
ASTAR-seq	mf	ATAC/RNA	no	$$	low	high
ISSAAC-seq	mf	ATAC/RNA	no	$$$	high	high
Smart3-ATAC	plate	RNA	yes-plate	$$	low	high
SHARE-seq	plate	ATAC/RNA	yes, but limited to first index	$$$	ultra-high	low
scCAT-seq	plate	ATAC/RNA	no	$$$	low	high

910 mf = microfluidics
911

Bibliography

1. Claringbould, A. & Zaugg, J. B. Enhancers in disease: molecular basis and emerging
treatment strategies. *Trends Mol. Med.* **27**, 1060–1073 (2021).

2. Lazarov, T., Juarez-Carreño, S., Cox, N. & Geissmann, F. Physiology and diseases of
tissue-resident macrophages. *Nature* **618**, 698–707 (2023).

3. Dey, K. K. *et al.* SNP-to-gene linking strategies reveal contributions of enhancer-related
and candidate master-regulator genes to autoimmune disease. *Cell Genomics* **2**,
(2022).

4. Tang, P. M.-K., Nikolic-Paterson, D. J. & Lan, H.-Y. Macrophages: versatile players in
renal inflammation and fibrosis. *Nat. Rev. Nephrol.* **15**, 144–158 (2019).

5. Vandereyken, K., Sifrim, A., Thienpont, B. & Voet, T. Methods and applications for
single-cell and spatial multi-omics. *Nat. Rev. Genet.* **24**, 494–515 (2023).

6. Sun, Z. *et al.* Joint single-cell multiomic analysis in Wnt3a induced asymmetric stem cell

division. *Nat. Commun.* **12**, 5941 (2021).

7. Wang, Y. *et al.* Single-cell multiomics sequencing reveals the functional regulatory

landscape of early embryos. *Nat. Commun.* **12**, 1247 (2021).

8. Bian, S. *et al.* Single-cell multiomics sequencing and analyses of human colorectal

cancer. *Science* **362**, 1060–1063 (2018).

9. Datlinger, P. *et al.* Ultra-high-throughput single-cell RNA sequencing and perturbation

screening with combinatorial fluidic indexing. *Nat. Methods* **18**, 635–642 (2021).

10. Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis.

Nature **566**, 496–502 (2019).

11. Lareau, C. A. *et al.* Droplet-based combinatorial indexing for massive-scale single-cell

chromatin accessibility. *Nat. Biotechnol.* **37**, 916–924 (2019).

12. Domcke, S. *et al.* A human cell atlas of fetal chromatin accessibility. *Science* **370**,

eaba7612 (2020).

13. Xu, W. *et al.* ISSAAC-seq enables sensitive and flexible multimodal profiling of

chromatin accessibility and gene expression in single cells. *Nat. Methods* **19**, 1243–

1249 (2022).

14. Ma, S. *et al.* Chromatin Potential Identified by Shared Single-Cell Profiling of RNA and

Chromatin. *Cell* **183**, 1103-1116.e20 (2020).

15. Zhu, C. *et al.* An ultra high-throughput method for single-cell joint analysis of open

chromatin and transcriptome. *Nat. Struct. Mol. Biol.* **26**, 1063–1070 (2019).

16. ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the

human genome. *Nature* **489**, 57–74 (2012).

17. Hagemann-Jensen, M. *et al.* Single-cell RNA counting at allele and isoform resolution

using Smart-seq3. *Nat. Biotechnol.* **38**, 708–714 (2020).

18. Bagnoli, J. W. *et al.* Sensitive and powerful single-cell RNA sequencing using mcSCRB-

seq. *Nat. Commun.* **9**, 2937 (2018).

19. Zhang, X., Marand, A. P., Yan, H. & Schmitz, R. J. Massive-scale single-cell chromatin

accessibility sequencing using combinatorial fluidic indexing. *BioRxiv* (2023)

doi:10.1101/2023.09.17.558155.

20. Zhang, H. *et al.* txci-ATAC-seq, a massive-scale single-cell technique to profile

chromatin accessibility. *BioRxiv* (2023) doi:10.1101/2023.05.11.540245.

21. Satoh, T. *et al.* The Jmjd3-Irf4 axis regulates M2 macrophage polarization and host

responses against helminth infection. *Nat. Immunol.* **11**, 936–944 (2010).

22. Krausgruber, T. *et al.* IRF5 promotes inflammatory macrophage polarization and TH1-

TH17 responses. *Nat. Immunol.* **12**, 231–238 (2011).

23. Kawashima, T. *et al.* STAT5 induces macrophage differentiation of M1 leukemia cells

through activation of IL-6 production mediated by NF-kappaB p65. *J. Immunol.* **167**,

3652–3660 (2001).

24. Argelaguet, R. *et al.* MOFA+: a statistical framework for comprehensive integration of

multi-modal single-cell data. *Genome Biol.* **21**, 111 (2020).

25. Kelly, B. & O'Neill, L. A. J. Metabolic reprogramming in macrophages and dendritic cells

in innate immunity. *Cell Res.* **25**, 771–784 (2015).

26. Huang, S. C.-C. *et al.* Metabolic Reprogramming Mediated by the mTORC2-IRF4

Signaling Axis Is Essential for Macrophage Alternative Activation. *Immunity* **45**, 817–830

(2016).

27. Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR: inferring

transcription-factor-associated accessibility from single-cell epigenomic data. *Nat.*

Methods **14**, 975–978 (2017).

28. Iles, K. E., Dickinson, D. A., Watanabe, N., Iwamoto, T. & Forman, H. J. AP-1 activation

through endogenous H(2)O(2) generation by alveolar macrophages. *Free Radic. Biol.*

Med. **32**, 1304–1313 (2002).

29. Karpurapu, M. *et al.* Functional PU.1 in macrophages has a pivotal role in NF-κB

activation and neutrophilic lung inflammation during endotoxemia. *Blood* **118**, 5255–

5266 (2011).

30. Au-Yeung, N., Mandhana, R. & Horvath, C. M. Transcriptional regulation by STAT1 and

STAT2 in the interferon JAK-STAT pathway. *JAKSTAT* **2**, e23931 (2013).

31. Rauluseviciute, I. *et al.* JASPAR 2024: 20th anniversary of the open-access database of

transcription factor binding profiles. *Nucleic Acids Res.* **52**, D174–D182 (2024).

32. Czimmerer, Z. *et al.* The transcription factor STAT6 mediates direct repression of

inflammatory enhancers and limits activation of alternatively polarized macrophages.

Immunity **48**, 75-90.e6 (2018).

33. Huang, Q. *et al.* Spi-B Promotes the Recruitment of Tumor-Associated Macrophages via

Enhancing CCL4 Expression in Lung Cancer. *Front. Oncol.* **11**, 659131 (2021).

34. Villar, J. *et al.* ETV3 and ETV6 enable monocyte differentiation into dendritic cells by

repressing macrophage fate commitment. *Nat. Immunol.* **24**, 84–95 (2023).

35. Ong, C.-T. & Corces, V. G. CTCF: an architectural protein bridging genome topology

and function. *Nat. Rev. Genet.* **15**, 234–246 (2014).

36. Kamal, A. *et al.* GRaNIE and GRaNPA: inference and evaluation of enhancer-mediated

gene regulatory networks. *Mol. Syst. Biol.* **19**, e11627 (2023).

37. He, L. *et al.* Global characterization of macrophage polarization mechanisms and

identification of M2-type polarization inhibitors. *Cell Rep.* **37**, 109955 (2021).

38. Viengkhou, B., White, M. Y., Cordwell, S. J., Campbell, I. L. & Hofer, M. J. A novel

phosphoproteomic landscape evoked in response to type I interferon in the brain and in

glial cells. *J. Neuroinflammation* **18**, 237 (2021).

39. Sadzak, I. *et al.* Recruitment of Stat1 to chromatin is required for interferon-induced

serine phosphorylation of Stat1 transactivation domain. *Proc Natl Acad Sci USA* **105**,

8944–8949 (2008).

40. Paul, A., Ismail, M. N., Tang, T. H. & Ng, S. K. Phosphorylation of interferon regulatory

factor 9 (IRF9). *Mol. Biol. Rep.* **50**, 3909–3917 (2023).

41. Platanias, L. C. Mechanisms of type-I- and type-II-interferon-mediated signalling. *Nat.*

Rev. Immunol. **5**, 375–386 (2005).

42. Sekrecka, A. *et al.* Time-dependent recruitment of GAF, ISGF3 and IRF1 complexes

shapes IFNα and IFNγ-activated transcriptional responses and explains mechanistic

and functional overlap. *Cell. Mol. Life Sci.* **80**, 187 (2023).

43. Finucane, H. K. *et al.* Heritability enrichment of specifically expressed genes identifies

disease-relevant tissues and cell types. *Nat. Genet.* **50**, 621–629 (2018).

44. Wang, Q. *et al.* Research Progress of Interleukin-27 in Inflammatory Bowel Disease.

Inflamm. Bowel Dis. (2023) doi:10.1093/ibd/izad153.

45. Andrews, C., McLean, M. H. & Durum, S. K. Interleukin-27 as a Novel Therapy for

Inflammatory Bowel Disease: A Critical Review of the Literature. *Inflamm. Bowel Dis.*

22, 2255–2264 (2016).

46. Somma, D., Kok, F. O., Kerrigan, D., Wells, C. A. & Carmody, R. J. Defining the Role of

Nuclear Factor (NF)-κB p105 Subunit in Human Macrophage by Transcriptomic

Analysis of NFKB1 Knockout THP1 Cells. *Front. Immunol.* **12**, 669906 (2021).

47. Ghoussaini, M. *et al.* Open Targets Genetics: systematic identification of trait-associated

genes using large-scale genetics and functional genomics. *Nucleic Acids Res.* **49**,

D1311–D1320 (2021).

48. Peters, A. L., Stunz, L. L. & Bishop, G. A. CD40 and autoimmunity: the dark side of a

great activator. *Semin. Immunol.* **21**, 293–300 (2009).

49. Steinbach, E. C. & Plevy, S. E. The role of macrophages and dendritic cells in the

initiation of inflammation in IBD. *Inflamm. Bowel Dis.* **20**, 166–175 (2014).

50. Nguyen, V. T. & Benveniste, E. N. Involvement of STAT-1 and ets family members in

interferon-gamma induction of CD40 transcription in microglia/macrophages. *J. Biol.*

Chem. **275**, 23674–23684 (2000).

51. Swanson, E. *et al.* Simultaneous trimodal single-cell measurement of transcripts,

epitopes, and chromatin accessibility using TEA-seq. *eLife* **10**, (2021).

52. Bartosovic, M., Kabbe, M. & Castelo-Branco, G. Single-cell CUT&Tag profiles histone

modifications and transcription factors in complex tissues. *Nat. Biotechnol.* **39**, 825–835

(2021).

53. Nichols, R. V. *et al.* High-throughput robust single-cell DNA methylation profiling with

sciMETv2. *Nat. Commun.* **13**, 7627 (2022).

54. Mulqueen, R. M. *et al.* High-content single-cell combinatorial indexing. *Nat. Biotechnol.*

39, 1574–1580 (2021).

55. Wienerroither, S. *et al.* Cooperative Transcriptional Activation of Antimicrobial Genes by

STAT and NF-κB Pathways by Concerted Recruitment of the Mediator Complex. *Cell*

Rep. **12**, 300–312 (2015).

56. Nelson, M. R. *et al.* The support of human genetic evidence for approved drug

indications. *Nat. Genet.* **47**, 856–860 (2015).

57. Freimer, J. W. *et al.* Systematic discovery and perturbation of regulatory genes in

human T cells reveals the architecture of immune networks. *Nat. Genet.* **54**, 1133–1144

(2022).

58. Pan, X., Zhu, Q., Pan, L.-L. & Sun, J. Macrophage immunometabolism in inflammatory

bowel diseases: From pathogenesis to therapy. *Pharmacol. Ther.* **238**, 108176 (2022).

59. van Wilgenburg, B., Browne, C., Vowles, J. & Cowley, S. A. Efficient, long term

production of monocyte-derived macrophages from human pluripotent stem cells under

partly-defined and fully-defined conditions. *PLoS ONE* **8**, e71098 (2013).

60. Zhang, H. *et al.* Fast alignment and preprocessing of chromatin profiles with Chromap.

Nat. Commun. **12**, 6566 (2021).

61. Unable to find information for 10550671.

62. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

63. Xu, M. *et al.* Arachidonic acid metabolism controls macrophage alternative activation

through regulating oxidative phosphorylation in pparγ dependent manner. *Front.*

Immunol. **12**, 618501 (2021).

64. Zhang, H. *et al.* Functional analysis and transcriptomic profiling of iPSC-derived

macrophages and their application in modeling Mendelian disease. *Circ. Res.* **117**, 17–

28 (2015).

65. Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering. *Nat.*

Methods **14**, 1083–1086 (2017).

66. Vorontsov, I. E. *et al.* HOCOMOCO in 2024: a rebuild of the curated collection of

binding models for human and mouse transcription factors. *Nucleic Acids Res.* (2023)

doi:10.1093/nar/gkad1077.

67. Booeshaghi, A. S., Hallgrímsdóttir, I. B., Gálvez-Merchán, Á. & Pachter, L. Depth

normalization for single-cell genomics count data. *BioRxiv* (2022)

doi:10.1101/2022.05.06.490859.

68. Unable to find information for 11129215.

69. Bravo González-Blas, C. *et al.* cisTopic: cis-regulatory topic modeling on single-cell

ATAC-seq data. *Nat. Methods* **16**, 397–400 (2019).

70. Berest, I. *et al.* Quantification of Differential Transcription Factor Activity and Multiomics-

Based Classification into Activators and Repressors: diffTF. *Cell Rep.* **29**, 3147-

3159.e12 (2019).