Feeding- state dependent modulation of reciprocally interconnected inhibitory neurons biases sensorimotor decisions in *Drosophila*

Eloïse de Tredern[1,2], Dylan Manceau[1,2], Alexandre Blanc[3,4], Panagiotis Sakagiannis[5], Chloe Barre[3,4], Victoria Sus[2], Francesca Viscido[2], Md Amit Hasan[2], Sandra Autran[2], Martin Nawrot[5], Jean-Baptiste Masson[3,4], Tihana Jovanic[2]*

[1]These authors contributed equally

[2]Université Paris-Saclay, CNRS, Institut des neurosciences Paris-Saclay, 91400, Saclay, France

[3]Institut Pasteur, Université Paris Cité, CNRS UMR 3571, Decision and Bayesian Computation, 75015 Paris, France

[4]Epiméthée, INRIA

[5]Computational Systems Neuroscience, Institute of Zoology, University of Cologne, 50674 Cologne, Germany

*corresponding author

Abstract

The feeding state of an animal changes behavioral priorities and thus influences even non-feeding related decisions. How is the influence of the feeding state transmitted to non-feeding related circuits and what are the circuit mechanisms involved in biasing non-feeding related decisions remains an open question. By combining calcium imaging, neuronal manipulations, behavioral analysis and computational modeling we determined that the competitive interactions between different behavioral responses to a mechanical stimulus are biased by the feeding state and found that this is achieved by differentially modulating two different types of reciprocally connected inhibitory neurons promoting opposing actions. The modulation of these inhibitory neurons influences the activity in the output layer of the network towards encoding more frequently an active type of response and less frequently a protective type of response in larvae fed on sugar compared to those fed on a balanced diet. The information about the internal state is conveyed to the inhibitory neurons through homologues of the vertebrate neuropeptide Y known to be involved in regulating feeding behavior.

Introduction

Physiological states like hunger and thirst are powerful regulators of behavior across the animal kingdom due to strong homeostatic drives that are critical for survival[1–7]. For example, across model systems, food-deprivation was shown to modulate responsiveness to stimuli by influencing sensory neurons and sensory pathways[8–11], suggesting that food-deprivation can alter the perceived value of a stimulus, which in turn affects the behavioral decisions. Various studies have implicated changes in central processing[12–15] which lead to changes in behavioral decisions in hungry animals. However, the detailed neural circuit mechanisms of this state-dependent flexibility of behaviors remain largely unknown.

Internal drives (e.g., hunger and thirst) need to be balanced by environmental demands such as the need to avoid dangers. Avoiding dangers is a critical instinctive behavior that needs to be balanced with finding and consuming food in order to ensure survival. Avoidance behaviors tend to be robust, which makes them excellent systems for studying neural bases of behavior[16–19]; yet they also need to be flexible in order for animals to adapt their behavioral strategies to different contexts and according to different internal states[17,20–22]. Feeding states and contexts can for example influence both the tolerance to the level of threat and action selection during threat avoidance[16–18,23].

At the neural circuit level, such behavioral flexibility is thought to be implemented by neuromodulation (modulation of existing synaptic connections by neuropeptides, for instance) that could bias the outcome of competition between diverse behaviors. The outcome could in that case differ depending on the neuropeptide released[24–30] . Alternatively, information rerouting (using alternative circuit pathways)[12,13,31] where information is processed differently depending on the context or state, could alter behavior choice in a context-dependent manner. The types of circuit motifs underlying competitive selection must allow for such flexible processing of information. However, the detailed neural circuit mechanisms underlying the state-dependent modulation of behavior, the neuromodulators involved, and their mechanism of actions on specific circuits, especially those that pertain directly to non-feeding or non-water seeking behavior, are not well understood.

This paper addresses the understudied aspect of the influence of physiological drives on competitive selection during avoidance behaviors and investigates the neural circuit mechanisms involved in a powerful model organism for neural circuit analysis: the *Drosophila* larva[32]. *Drosophila* larvae are ideally suited for combining comprehensive, synaptic-resolution circuit mapping in electron microscopy (EM) across the nervous system[33–36] with targeted manipulation of uniquely identified circuit motifs at the individual neuron level, which makes it possible to establish causal relationships between circuit structure and function in a brain-wide manner. In addition, evolutionarily conserved neuropeptidergic and hormonal pathways in *Drosophila*, have been shown to regulate its diverse behaviors[11,18,27,37,38].

Previous studies have described larval avoidance response to a mechanical stimulus (air puff) and detailed the neural circuit underlying the competition between "Hunching", a protective or "startle-like" type of behavior, and "Head Casting", an "active exploratory type of behavior that can

lead to escape[35,39–43]. We identified circuit motifs underlying competitive interactions between behavioral actions (reciprocal inhibition of inhibition) and sequence transitions (lateral and feedback disinhibition) between the two behaviors. These types of motifs based on disinhibition would allow for flexible behavioral selection in a context/state dependent manner. By combining calcium imaging, neuronal and neuropeptide manipulation at the single cell level with automated tracking, behavior classification and computational modeling this work shows that larval responses to the air puff are biased towards less protective actions and towards more active, exploratory actions upon changes in their feeding condition. We determine that this bias is due to the differential modulation of two reciprocally connected inhibitory neurons that drive competing behaviors (Hunching and Head Casting): the activity of the neuron that promotes the protective Hunching is decreased and the activity of the neuron that inhibits Hunching is increased. We also show that the modulation at the level of reciprocally interconnected inhibitory neurons results in a bias at the level of the network output, towards a state that will lead to Head Casting at the expense of Hunching upon feeding on sucrose. Finally, we determine that NPF and sNPF modulate the activity of the reciprocally interconnected inhibitory neurons to bias the behavioral output in a feeding state-dependent manner.

RESULTS

Changes in feeding conditions for short periods of time affect larval feeding and locomotion

To study the effect of diet on behavioral decisions as a result of cognitive control and prioritization of needs and motivations rather than to long-term physiological changes that would affect circuit properties, we sought to establish a food-deprivation protocol of short duration. We determined the shortest duration of food deprivation that was sufficient to induce quantifiable changes in behavior. Depriving larvae of food completely (by putting them on a water-soaked filter paper) or feeding them on 20% sucrose only (and therefore deprived of proteins and increasing their sugar intake) for 90 mins was sufficient to alter larval locomotion. Starved larvae and larvae fed on sucrose moved at a higher speed and spent more time Crawling than normally-fed larvae eventually dispersing faster in less-curved trajectories (Figure 1D, Figure S1A-B). These changes in locomotion are consistent with increased exploration and are likely due to an increased drive to find nutrients caused by deprivation. In order to determine that the 90 minutes feeding deprivation protocol induced changes in behavior that were reversible we monitored locomotion in these larvae upon refeeding them for 15 minutes. Upon refeeding, larvae were slower and their locomotion phenotype was similar to the one of larvae that were constantly fed (Figure S1C).

To determine whether the larval need for nutrients was affected by the feeding protocols that they were subjected to, the consumption rate of different foods was quantified in different feeding conditions. Starved larvae significantly increased their intake of standard *Drosophila* food and yeast (rich in amino acids) compared to both normally-fed larvae and larvae that were fed on sucrose (Figure 1A), which is consistent with a deficit in nutrients in starved larvae. Larvae fed on sucrose significantly increased their intake of yeast, which is consistent with a deficit in amino acids (Figure 1A). These larvae, however, did not increase their intake of standard *Drosophila* food, suggesting that the increased sugar consumption might suppress their intake of

carbohydrate-rich food. To test whether larvae fed on sucrose were repulsed by sugar, a choice assay was performed: larvae were added to the middle of an agar plate where half of the plate was covered in agar and the other half in agar mixed with sucrose. The larvae were then monitored for 15 minutes (Figure 1C). After the 15 minutes, more larvae were found on the agar+sucrose half of the arena in normally-fed and starved larvae, with a preference index increasing over time, while larvae fed on sucrose showed the opposite effect and a decrease in preference for agar supplemented with sucrose. Larvae fed on only 20% sucrose for 90 minutes significantly decreased their sucrose intake compared to fed and starved larvae, and their sucralose intake was similar to that of normally-fed and starved larvae (Figure 1A4), suggesting that their avoidance of sugar was mediated by energy sensing pathways and not taste.

To determine whether the 90-minute food-deprivation or sucrose diet protocols caused changes in glucose levels, the hemolymph glucose levels were quantified in the different feeding conditions. Glucose levels were slightly but not significantly decreased in starved larvae compared to the fed larvae. In larvae that were fed on 20% sucrose, the glucose level was significantly increased compared to both normally-fed and starved larvae (Figure 1B). This high glucose level suggests that larvae fed on sucrose suppress consumption of carbohydrates due to high circulating levels of glucose. Finally, the water consumption of larvae in different feeding conditions was quantified. Larvae fed on 20% sucrose increased their water consumption significantly compared to both normally-fed and starved larvae (Fig 1A3), likely due to changes in extracellular osmolality as a result of high sugar intake[6]. We therefore tested whether rehydration would reduce the increase in exploration and locomotion observed in larvae fed on sucrose. Indeed, after larvae fed on sucrose only were put on water for 15 minutes, their locomotion was similar to the ones of normally fed larvae (Figure S1D)

Altogether, these results suggest that depriving larvae of nutrients for 90 minutes is sufficient to alter feeding and larval locomotion and due to lack of nutrients and lower energy levels in starved larvae and protein hunger and thirst in larvae fed on sucrose. The observed increase in locomotion larvae deprived of nutrients would thus likely result from an increase in motivational drive to find the missing nutrients.

Changes in physiological states affect sensorimotor decisions in response to an air puff

To determine whether starvation and a sucrose-only diet can affect non-feeding-related behaviors, we monitored *Drosophila* larva sensorimotor decisions in response to an aversive mechanical stimulus, the air puff. In response to an air puff, larvae perform probabilistic sequences of five mutually exclusive actions that we have characterized in detail in the past[35,44]: Hunch, Bend, Stop, Back-up, and Crawl. We have also identified circuit motifs and characterized the neural circuit mechanisms underlying the competitive interactions between the two most prominent actions (i.e. the Hunch and the Bend) that occur in response to an air puff[35]. The model in that study predicts that the different activation levels of inhibitory neurons determine which actions will take place: the Hunch, Bend, or the Hunch Bend sequence.

With characterized neurons and synaptic connectivity between the neurons, as well as the availability of driver lines that label neurons of interest, this circuit provides an excellent system to study whether the activity of the neurons is modulated by the changes in an animal's physiological state.

To determine whether the feeding state modulates larval behavior in response to the mechanosensory stimulus and, by extension, whether sensorimotor decision-making in response to an air puff is an adequate behavioral paradigm to study the effects of food deprivation on neural circuit activity, we compared larval behavioral responses to an air puff after subjecting them to the different feeding protocols (as described in the first section of the Results). For this purpose we used automated tracking[45] to monitor larva behavior in response to an air puff and updated the machine-learning-based classification method developed in our previous work[44] to compute probabilities of the different actions that occur in response to an air puff. The new classifiers were trained on larvae fed on different diets, thus taking into account a broader range of behavioral dynamics and separated the bend behavioral category into two types of bending behavior: Static Bends, a form of protective action where the larva responds to the stimulus by immobilizing in a curved position for a period of time, and the exploratory active Head Casts that can lead to escape[39].

Our analysis showed that larvae fed on sucrose Hunch less (and perform less of Static Bends) and Head Cast more (Figure 1E1). Similarly, starved larvae performed less Static Bends and more Head Casts (Figure 1E2). Transitions to Hunching were decreased in larvae fed on sucrose and starved larvae, while transitions to Head Cast were increased (from Hunching to Head Casting in larvae fed on sucrose only fed larvae and from Crawl to Head Casting in both starved larvae and larvae fed on sucrose) (Figure 1). Increasing duration of sucrose feeding and starvation to 5h resulted in similar phenotypes (Figure S1E, F).

Using our machine-learning based classification to monitor internal state induced behavioral changes revealed that changing feeding conditions for short periods of time alters sensorimotor decisions in response to an aversive mechanical cue and results in less protective type of actions, and more actions consistent with active exploration and escape.

The feeding state does not affect the responses of chordotonal sensory neurons to a mechanical stimulus

To test whether changes in feeding conditions alter the perceived value of a stimulus which in turn would affect the behavioral decisions, calcium responses in sensory neurons that sense the air puff, the chordotonal sensory neurons[35,41,44,45] were monitored in the three different feeding conditions: normally fed, fed on sucrose and starved to a moderately strong mechanical stimulus (Figure S2B). We found no significant differences in chordotonal calcium responses in the different feeding states (Figure 2B, Figure S2A), although the responses were slightly higher in starved larvae compared to the larvae fed on standard food and on sucrose only. To ensure that the changes in feeding conditions did not influence protein expression and, by extension, GCAMP expression levels, GFP fluorescence was compared in larvae in all three feeding conditions. No differences were found in GFP fluorescence in the three feeding states (Figure S2C). The results therefore suggest that chordotonal responses are not significantly affected by starvation or sucrose feeding and that the behavioral changes observed in the different feeding conditions are not due to the changes in stimulus sensitivity at the level of sensory neurons. We further confirmed this by optogenetically activating chordotonal sensory neurons in larvae in all three feeding states using a driver that labels all eight subtypes of chordotonal R61D08. We found that although the activation of chordotonal sensory neurons in all three feeding states was the same due to

optogenetic stimulation, larvae performed fewer Hunches and more Head Casts when they were fed only on sucrose than when they were normally fed (Figure 2C). Similarly, the modulation of behavior when they were starved could be observed upon chordotonal optogenetic activation (Figure S2D). Altogether, these results show that the feeding state-dependent modulation could target neurons downstream of chordotonals.

Feeding states modulates the output of the circuit for the choice between Hunch and Bend

To determine whether the changes in feeding state influence the output of the network for the selection between the Hunch and the Bend, characterized in previous work[35], calcium responses of projection neurons Basin-1 and Basin-2 were monitored at different intensities of mechanical stimulation in the three feeding states. Basin-1 and Basin-2 are differentially involved in Hunching and Bending, as shown in Jovanic *et al.*, 2016: Basin-1 is required for both Hunching and Bending, while Basin-2 promotes Bending and inhibits Hunching. Responses in Basin-2 neurons can thus be used as a read-out of the Bend state: if Basin-2 is ON, the larvae will Bend, and if it is OFF, the larvae will Hunch. We found that for most stimulus intensities, the Basin-1 responses remain only mildly affected by the sucrose-only diet (Figure 3B). Only for the very weak stimulus intensity do we observe a decrease in Basin-1 response (Figure S3C). In the starved condition, the effect on Basin-1 neurons was similar to the sucrose-only feeding condition (Figure S3A, C). The activity of Basin-2, on the other hand, was moderately increased in larvae fed only sucrose at all intensities except at the lowest stimulus intensity, where there was no difference between Basin-2 responses in different feeding conditions (Figure 3C, D, Figure S3D). However, the activity of Basin-2 in starved larvae was slightly (but not significantly) decreased compared to fed larvae at lower intensities of stimulation, whereas it was increased at higher intensities of stimulation (Figure S3B, D).

We then compared the distributions of Basin-2 responses in larvae fed on either standard food or sucrose to a moderately strong mechanical stimulus (5V) and observed a higher probability of weak response in normally-fed larvae (Figure S3E). The previous study has recorded depolarization responses simultaneously from Basin-1 and Basin-2 to a mechanical stimulus and has shown that Basin-1 neurons always respond while Basin-2 responses are probabilistic[35]. In this study, we computed the probability of Basin-2 responses to the mechanical stimulus and observed a significantly higher probability of responses that cross the threshold in larvae fed on sucrose, which is consistent with higher Head Casting and lower Hunching probabilities in these larvae. In starved larvae, similar trends were observed at higher stimulus intensities, while at medium intensity (5V), there was no difference between Basin-2 response probability in fed and starved larvae. This difference in Basin-2 responses between larvae fed only sucrose and starved larvae could explain why Hunching is not consistently significantly reduced in starved larvae.

Reciprocally connected Inhibitory interneurons in the decision circuit are differentially modulated by the feeding state.

Based on the simulation of the model of the circuit for a choice between two air puff-induced actions (Hunch and Bend)[35], the prediction was that the relative level of activation of the reciprocally connected feedforward inhibitory interneurons determines the outcome of the competition: Basin-1 only state (Hunch) or Co-active state (Bend). In this study, we investigated

if changes in the feeding state could affect the level of activation of two different classes of the inhibitory neurons: feedforward and feedback inhibitory neurons.

To address this, we started by building a simple rate model, that as in the previous study[35] is similarly based on the observed connectivity between neuron classes from the EM reconstruction data, and where weights of excitatory or inhibitory connections between neurons were proportional to the number of synapses (Figure 3C-E). As previously described the model predicts that the level of activation of the two classes of feedforward inhibitory neurons will determine the output state of the network at the level of the Basin-1 and Basin-2 neurons (Figure S4A). The "sucrose state" was modeled in one version of the model as the decreased maximum intensity of LNa neurons to explore the role of the feedforward inhibitory neurons in the feeding state modulation (Figure 3C). In order to explore the contribution of the feedback inhibitory neurons, in another version of the model, the sucrose state was represented by adding an input to the Handle-b neurons (Figure 3D). Finally, we made a combined model where both the LnA and Handle-b were modulated in the sucrose state (Figure 3E, Figure S3A-D). In all of the versions of the model the Hunching decreased and Head Casting increased in the "sucrose" state compared to normal fed state (Figure 3C3, D3, E3).

These model results suggest that modulating both feedforward and feedback inhibitory neurons of the circuit could result in the observed behavioral changes upon sucrose feeding. To test this experimentally, calcium transients in response to a mechanical stimulus were imaged in the two different types of inhibitory neurons, to which we had genetic access. An LnA neuron type, Griddle-2 (G2), that promotes Hunching[35], showed significantly lower responses to a mechanical stimulus in larvae fed on sucrose compared to the normally fed larvae (Figure 4B). The decrease in response was observed at different intensities of mechanical stimulation (Figure S4C). Similarly, in starved larvae, the responses of Griddle-2 neurons were also decreased across different intensities of stimulation (Figure S4A, C).

Based on the model, the decrease in the activity of LNa would result in less Hunching and more Head Casting which is consistent with the behavioral changes we observed after feeding on sucrose. The model has also shown that another motif in the circuit, feedback disinhibition, promotes Bending by amplifying the activity of Basin-2. We recorded Handle-b responses to the mechanical stimulus in the different feeding states and found that the Handle-b neurons that inhibits Hunching[35] show stronger responses to a mechanical stimulus in larvae fed on sucrose and in starved larvae compared to normally fed larvae (Figure 4C, Figure S4B,D). This was true for all the different intensities of stimulation we tested (Figure S5D). Therefore, the decreased response of Griddle 2 neurons (Figure 4B) and increased responses of Handle-b neurons (Figure 4C) in larvae fed on sucrose could bias the behavioral outcome, consistent with the behavioral changes observed in these larvae (less Hunching and more Head Casting), (Figure 1E1) upon changes in the feeding state.

EM analysis shows that Griddle-2 and Handle-b neurons are reciprocally interconnected (Figure 4A)[35].The circuit and the model predict that silencing Handle-b results in an increase in Griddle-2 activity and vice-versa (Figure 4C, D). Further, the reciprocal connections between Griddle-2 and Handle-b were probed functionally and the model predictions tested experimentally; calcium responses to a mechanical stimulus were monitored in Griddle-2 neurons while inactivating

Handle-b using tetanus-toxin in both larvae fed on standard food and on sucrose only. The responses were increased compared to control larvae in both feeding states (Figure 4C1-C2, Figure S4), which is consistent with the Handle-b inhibition of Griddle-2. However, the increase in Griddle-2 responses upon Handle-b inactivation in larvae fed on sucrose did not reach the levels of Griddle-2 responses upon inactivation in larvae fed on standard food (Figure 4C2, Figure S4E). This could suggest that the increase in Handle-b activity after sucrose feeding is insufficient to reduce the Griddle-2 responses in larvae fed on sucrose and that both Handle-b and Griddle-2 neurons could be independently modulated (Figure S4). Indeed, this is in line with model 1 and 3 predictions where Griddle-2 is modulated extrinsically (Figure 3CE, Figure 4C, Figure S5E).

The converse experiments were also performed, i.e., inactivating Griddle-2 and imaging the activity in Handle-b (Figure 4C4, C5). Handle-b responses to the mechanical stimulus were increased in normally fed larvae upon Griddle-2 inactivation (Figure 4C5, Figure S4F), which is consistent with Griddle-2 inhibition of Handle-b. In larvae fed on sucrose, the activity of the Handle-b neuron was not significantly different and was similar to the activity upon Griddle-2 inactivation in larvae fed on standard food (Figure 4C5, Figure S4F). This lack of increase in Handle-b activity upon Griddle-2 inactivation in larvae fed on sucrose could be due to the different state of the network in the larvae fed only sucrose: the activity of Griddle-2 is already low in these larvae (Figure 4B, Figure S4C), and removing it may not impact its inhibition of Handle-b. Moreover, when silencing the LNa type Griddle-2, other two LNa type neurons remain intact and may continue to inhibit Handle-b. In addition, removing Griddle-2 in this condition could impact the disinhibition of Basin-1 and/or the breaking of the positive feedback Basin-2-Handle-b-Basin-2. Indeed, removing Griddle-2 abolishes the difference between the Handle-b activity in larvae fed on standard food and larvae fed on sucrose only (Figure 4C6).

The feeding state-dependent increase in Basin-2 responses depends on the changes in activity of the inhibitory interneurons

Our previous work has shown that Basin-2 (B2) neurons receive input directly from the inhibitory neuron LNa and are disinhibited by the feedback inhibitory neuron Handle-b that makes inhibitory connections onto LNa and LNb neurons; Handle-b receives input primarily from Basin-2 neurons. The feedback disinhibition of Basin-2 creates positive feedback that stabilizes the Basin-2 ON state[35]. To determine whether the increase in Basin-2 responses in larvae fed on sucrose depended on feeding-state-dependent changes in the activity of the inhibitory neurons, Handle-b neurons were optogenetically activated, and the activity of B2 neurons were recorded in response to a mechanical stimulus. We found that activating Handle-b in normally fed larvae increases the levels of Basin-2 responses to a level similar to that observed in larvae fed on sucrose (Figure 5B, Figure S5B). We further silenced Handle-b neurons and monitored calcium responses in Basin-2 neurons in different feeding states (Figure 5A, Figure S5A). Silencing Handle-b neurons in larvae fed on sucrose decreased Basin-2 responses and erased the difference in Basin-2 response levels between larvae fed on standard food and ON sucrose (Figure 5A, Figure S5A). This result is consistent with previous characterization of the Handle-b disinhibition of Basin-2. Because Handle-b disinhibits Basin-2, silencing Handle-b therefore increases the inhibition of Basin-2 by LNa, for example, and thus reduces its responses to mechanical stimuli. In addition, in normally fed larvae where the activity of LNa is high, Basin-2 responses were almost completely abolished (Figure S5A).

Griddle-2 neurons were then inactivated and the activity in Basin-2 was imaged in the two feeding states. As expected, silencing Griddle-2 abolished the difference in Basin-2 responses in the two states. This result is consistent with the effect of Griddle-2 on Handle-b activity (Figure 5). Silencing Griddle-2 in fed larvae resulted in small (but not significant) increase in Basin-2 responses (Figure 5C, Figure S5C). However, the mean responses of Basin-2 in larvae fed only sucrose unexpectedly and significantly decreased upon Griddle-2 silencing. We also computed Basin-2 response probabilities. While inactivating Griddle-2 did not affect Basin-2 response probabilities in normally fed larvae, it did so in larvae fed on sucrose; the non-response probabilities are significantly increased upon Griddle-2 inactivation in sucrose-fed larvae. (Figure 5C, Figure S5C). The mean responses of trials with only Basin-2 ON responses reveal that the level of responses of Basin-2 was increased in normally fed larvae to the level found in larvae fed on sucrose in the control. The mean responses of trials with only Basin-2 ON responses also show that the level of Basin-2 response was decreased in larvae fed on sucrose (Figure S5C) and similar to the level of activity in fed control larvae

These findings suggest that there are two mechanisms that control Basin-2 response probabilities and amplitudes, respectively. The two mechanisms are decoupled upon Griddle-2 inactivation. Inactivating Griddle-2 may decrease Basin-2 response probabilities by decreasing Basin-1 responses (that then disinhibits less Basin-2). At the same time, inactivating Griddle-2 increases Handle-b responses and thus amplifies Basin-2 responses more upon Griddle-2 activation (when Basin-2 responses are triggered).
Griddle-2 is thus required for state-dependent modulation of Basin-2, possibly by mediating the disinhibition of Basin-2 by Handle-b. Removing Griddle-2 in normally fed larvae disinhibits Basin-2. On the other hand, removing Griddle-2 in larvae fed on sucrose when the activity of Griddle-2 is already low (and Griddle-2 is not inhibiting Basin-2 strongly) may thus favor other inhibitory pathways and result in lower Basin-2 responses in larvae fed on sucrose upon Griddle-2 inactivation.

These results show that two types of reciprocally connected inhibitory neurons that promote competing actions inhibit each other during responses to a mechanosensory cue and are differentially modulated by the change in feeding states (starvation and feeding on sucrose). The activity of the Griddle-2 that promotes a Hunch is decreased, while the activity of Handle-b that promotes a Bend is increased. This differential modulation of the inhibitory neurons, in turn, has an impact on the state of the network in a feeding state-dependent manner and biases the behavioral responses towards less Hunching and more Head Casting by increasing the activity of the Basin-2 neurons when larvae are fed on sucrose only. In addition, it reveals that two different layers of the network are modulated by the changes of the feeding state: the reciprocally connected feedforward inhibitory neurons as well as the feedback inhibitory neurons that stabilizes the state of the network resulting in the co-activation of both Basin-1 and Basin-2.

The feedback inhibitory neuron receives input from an NPF-releasing neuron, that senses the changes in the feeding state

In order to determine the sources of feeding state-dependent modulation, we looked in the connectome for upstream partners of Griddle-2 and Handle-b neurons. Previous work has shown that the different neurons in the circuit receive input from long-range projection neurons that, as

suggested by their morphology and connectivity, could bias the output of the network based on contextual/internal state information[35]. Interestingly, by matching light microscopy images to electron microscopy reconstruction images, we found one of these neurons is an NPF-expressing neuron that synapses on Handle-b. NPF (neuropeptide F), a homolog of the mammalian neuropeptide Y[46] is a hunger signal, and its expression promotes feeding in both adult and larval Drosophila[27]. We, therefore, sought to investigate the implication of NPF neurons in modulating the activity of the inhibitory neurons in a state-dependent manner. In the larva, there are two pairs of NPF-expressing neurons. Both of them have cell bodies in the brain; the dorsolateral (DL-NPF) projects ipsilaterally within the brain lobes and the dorsomedial (DM) sends descending projections via the suboesophageal zone (SEZ) through the ventral nerve cord (VNC).

To study the influence of NPF neurons on Handle-b activity, the activity of the DM-NPF descending neuron was monitored in the different feeding states. To this end, we drove GCAMP6s in the NPF descending neurons with NPF-GAL4and measured the intensity of fluorescence in the projections of the NPF descending neuron in the VNC of larvae fed either on standard food (fed), on 20% sucrose (sucrose fed) or completely starved (given only water for 90 minutes). We showed that starvation or feeding on sucrose both increase the activity of the NPF descending neuron (Figure 6B).

NPF was shown to be involved in promoting feeding[47]. To investigate whether the increase in activity of NPF descending neurons upon 90-minute sucrose feeding or starvation had an influence on NPF release and behavior NPF was knockdown selectively in the NPF descending neuron using a split-GAL4 driver that labels only the descending pair of NPF neurons and larval locomotion monitored in larvae fed on standard food, larvae feed on sucrose and starved larvae (Figure 6C). Downregulating NPF in the DM-NPF neurons descending into VNC impaired the increase in exploration observed in larvae fed on sucrose and starved larvae. it abolishes the difference in tortuosity over 20 s, time fraction in runs, mean scaled speed and maximum dispersal over 60 s between larvae fed on standard food and starved larvae and abolishes the difference between larvae fed on standard food and larvae fed on sucrose or even decrease exploration in larvae fed on sucrose compared to larvae fed on standard food.

To understand the influence of NPF neurons on Handle-b activity, the interactions between the two neurons were tested functionally by hyperpolarizing the NPF neurons with the inwardly rectifying potassium channel Kir2.1 using an NPF-LexA driver line that labels both pairs of NPF neurons and monitored calcium responses of Handle-b to a mechanical stimulation. Upon silencing the NPF neurons, we observed a decrease in Handle-b responses to mechanical stimulations (Figure 6D). The NPF descending neuron could thus facilitate responses of Handle-b neurons. Upon changes in the feeding state, the NPF neurons whose activity increases upon starvation and sucrose feeding, could enhance Handle-b responses to a mechanical stimulus, resulting in higher Handle-b responses (Figure 4B, D) that biases behavioral choice towards less Hunching and more Head Casting.

In the connectome we found large dense core vesicles on the NPF neurons in the proximity of the Handle-b neuron. Moreover, we showed that Handle-b expresses the NPFR1 using T2A-LexA method[48]where we drove one reporter in all NPFR1 neurons using the NPFR1 T2A-Lexa and another reporter with a Handle-b specific split-GAL4 driver. The intersection of the two expression patterns revealed that the Handle-b neurons express the NPFR (Figure 6E). Having established

that Handle-b and Griddle-2 are instrumental in the feeding state-dependent modulation of the behavioral choice of larvae in response to mechanical stimulation and that NPF modulates Handle-b, given that the model and functional connectivity experiment predict that the Handle-b and Griddle-2 could be independently modulated (Figure 4C, Figure S5E-F) we investigated whether Griddle-2 could be also influenced by NPF. We found that Griddle-2 does not receive synaptic input from the NPF descending neurons. Since neuropeptides can act outside of synaptic sites and we found dense core vesicles characteristic for neuropeptide release along the NPF axon in the proximity of Griddle-2 neurons, we also examined the NPFR1 expression in Griddle-2 neurons, as well as Basin-1 and Basin-2 expressed the NPFR using a similar approach as for Handle-b and didn't find any NPFR1 positive cells in these neurons (Figure S7D).

Altogether these results show that NPF senses the changes in feeding state and its activity increases upon starvation or sucrose only feeding. The changes in NPF activity will influence Handle-b activity that will increase upon starvation or sucrose only feeding due to a facilitating effect of NPF. This increase in activity will then bias the sensorimotor decisions towards less Hunching and more Head Casting.

sNPFR in the interconnected inhibitory interneurons mediates the state-dependent modulation of larval behavioral responses to a mechanical stimulus.

The model and calcium imaging predict that Griddle-2 and Handle-b could be independently modulated (Figure 4). Since Griddle-2 does not express the NPFR receptor, another neuropeptide could influence Griddle-2 and the circuit. We therefore investigated the effect of another neuropeptide on the circuit: sNPF (the short neuropeptide F). sNPF is a second homolog of the mammalian NPY whose receptor is widely distributed in the larval nervous system including the VNC[49] and was shown to be involved in regulating hunger-driven behaviors[27] and in facilitating mechano-nociceptive responses[37]. Therefore, we sought to examine whether the interneurons in the circuits express the receptor for sNPF. By genetically co-expressing a red fluorescent reporter (jRGeco) under the control of LexA drivers that label either Handle-b, Griddle-2, Basin-1 or Basin-2 neurons we identified in the literature and existing Gal4 databases (see Methods for details) and a green fluorescent reporter (GCamP6s) under the control of the sNPFR1 promoter with a T2A-Gal4 construct, we showed that both Griddle-2 (Figure 7A) and Handle-b (Figure 7D) express the sNPFR, while the two Basin neurons do not (Figure S7E).

To determine whether sNPF signaling was involved in responses to a mechanical stimulus in these two neurons, their responses were monitored upon sNPFR1 downregulation. Calcium imaging recordings show that genetically downregulating sNPFR1 expression in Handle-b increases its responses to mechanical stimulation (Figure 7B), suggesting that sNPFR1 has an inhibitory effect on Handle-b. On the other hand, responses of Griddle-2 were decreased upon sNPFR1 downregulation (Figure 7E), suggesting that sNPF facilitates Griddle-2 responses to a mechanical stimulus. Since Handle-b inhibits Hunching and promotes Bending (Head Casting), while Griddle-2 promotes Hunching and inhibits Bending (Head Casting) downregulating the sNPFR1 in these neurons would inhibit Hunching and promote Head Casting. Indeed, the experiments showed that sNPFR1 knockdown in either of these two neurons leads to less Hunching and more transitions to Head Casting (Figure 7C, F) in response to an air puff. The sNPFR1 knockdown in Handle-b and Griddle-2 leads to a modulation of their activity similar to that caused by feeding on only sucrose (lower Griddle-2 and higher Handle-b activity) and yields the same behavioral changes. This suggests that sNPF signaling on these neurons could be

downregulated in larvae fed on sucrose and at least partially responsible for the modulation of the behavioral choice of the larvae in response to a mechanical stimulation. Giving stock to this hypothesis, the downregulation of sNPFR1 in Handle-b or Griddle-2 did not impact their calcium responses to mechanical stimulation in larvae fed on sucrose or starved (Figure S8).

Altogether these results show that Handle-b and Griddle-2, reciprocally interconnected inhibitory neurons that drive opposing actions, are differentially modulated by the sNPF signaling pathway to bias the response to air puff towards less Hunching and more Head Casting

Discussion

Whereas many studies showed that various behaviors are affected by an animal's physiological state, especially hunger, this work goes further by describing the neural circuit mechanisms by which transient changes in the physiological status impact behaviors that are not directly related to this state. To determine whether and how changes in the feeding state affect behaviors unrelated to feeding, we took advantage of a well characterized behavioral response of *Drosophila* larva to a mechanical stimulus for which we had in the past dissected the circuit mechanisms underlying the selection between two main types of responses: Hunch and Bend. The current study reveals the neural circuit mechanisms of modulation of these sensorimotor decisions by the feeding conditions. Slight changes in feeding conditions affect an animal's motivational state and bias responses to an air puff towards less Hunching and more Head Casting. Reciprocally interconnected inhibitory neurons that drive competing actions are differentially modulated by the changes in the feeding state: the activity of the neuron that promotes a Hunch is decreased and the activity of the neuron that inhibits a Hunch is increased. This modulation at the level of inhibitory neurons involving NPF and sNPF signaling systems biases the output of the network towards promoting Head Casting and inhibiting Hunching, a modulation consistent with the state-dependent behavioral changes we observed.

Changes in internal physiology can affect unrelated behaviors

Hungry animals behaving differently in non-feeding related contexts have been reported in various studies and across the animal kingdom[7,51]. However, the underlying logic and neural mechanisms have not been well understood. These changes could be linked to overall changes in behavioral strategies in hungry individuals[7] to ensure survival. Risk-taking has been shown to be increased by hunger[50–52], even in social-related decisions[7,53], further raising questions about how food-deprivation signals are integrated into the neural computations underlying non-feeding related behaviors. Animals need to be able to make any decisions flexibly depending on the need that is most critical at a given moment. Our study shows that a circuit underlying sensorimotor decisions in response to a mechanical stimulus located at the early stages of sensory processing is influenced by the changes in diet. This finding supports the idea that state dependent flexibility of behavior could be achieved by the physiological state acting on circuits throughout the nervous system and thus reorganizing its activity in a distributed manner. In various ecological contexts, survival often entails a trade-off between avoiding dangers and pursuing food and water-seeking behaviors[51]. For example, increased exploration will increase the likelihood of finding food but also of encountering a dangerous situation[51]. If animals' need for food or water increases either due to food deprivation or thirst, they might be more likely to explore more despite an increasing risk of threat. Similarly, food-deprived and thirsty animals might take more risks and ignore

aversive cues to increase their chances of getting near food or water sources. Hungry animals use different strategies when escaping predators compared to satiated ones[17,23].

Feeding state does not alter the behavioral responses to mechanical stimuli by acting on the sensory neurons

While various studies have shown that hunger affects behaviors by altering the responses of sensory neurons[9,11,54,55], others have implicated central mechanisms in feeding state dependent behavioral flexibility[13,31,56]. We found that the activity of chordotonal mechanosensory neurons that sense the air puff is not significantly altered upon feeding on sucrose or starvation, while the activity of the downstream neurons is. The changes at the level of sensory pathways tunes animals' perception in order to increase their likelihood of finding food and feeding by increasing responsiveness to appetitive and decreasing responsiveness to aversive food-related stimuli. The implication of central mechanisms, on the other hand, may suggest that hunger acts as a global regulator of behavior, i.e. hunger may change brain activity in such a way to reevaluate goals and behavioral strategies in order to increase animals' chances of survival[7].

Our experiment monitors calcium responses of all the chordotonal sensory neurons together. The different inhibitory and projection neurons in the circuit receive inputs from different chordotonal subtypes. This could also explain the differential modulation of the different neuron subtypes in the circuit if some subtypes of chordotonal are modulated by the changes in the internal state while others are not. However, optogenetic activation of all the chordotonal neurons in the different feeding conditions still resulted in decreased Hunching and increased Head Casting upon starvation or sucrose diet, strongly suggesting that downstream neurons are involved. Even if chordotonals are involved in altering behavior in a state- dependent manner, their modulation is not required to alter the behavior in response to the mechanical stimulus due to the contribution of the downstream circuitry.

Reciprocal interconnected inhibitory neurons are targeted by the modulation

Calcium imaging combined with neuronal manipulations revealed that inhibitory neurons (and not the projection neurons) in the circuit for selecting between the Hunch and the Bend are the target of modulation by the changes in the feeding conditions. Inhibitory neurons were shown to be the target of (contextual) modulation in other systems[57–59].
Previous work has identified reciprocal inhibition of inhibition as a motif underlying the competition between a startle-type action and an exploratory action[35]. Such a motif was shown to underlie similar computations in different species and brain areas[35,60–65] and was proposed to provide flexibility to the selection process[35,63,65]. The current study determines that one of the reciprocally connected inhibitory neurons within this motif, LNa type Griddle-2, is modulated by the changes in the feeding conditions and that this modulation contributes to biasing sensorimotor decisions. These results confirm theoretical predictions that such a motif confers the sensorimotor circuit the capacity to be tuned to other types of information, in this case, internal state information (starvation and thirst). Moreover, these findings are in line with predictions that shaping the output of the network through disinhibition by reciprocally connected inhibited neurons allows for flexible, competitive selection[35,63,65].

Feedback disinhibition may act as a handle for neuromodulation of the circuit.

This work shows that another type of inhibitory neuron, Handle-b, which is a feedback inhibitory neuron that provides positive feedback to the Hunch inhibiting Basin-2 neuron through feedback disinhibition, is also modulated by changes in internal physiology. Handle-b is also reciprocally connected to the LNa neurons that participate in the reciprocal inhibition of inhibition motifs. This connectivity pattern suggests that, in addition to competition within the reciprocal inhibition of inhibition motifs, the competition between the two layers of the circuits is also modulated by the changes in internal physiology. Similar to recurrent excitation, the feedback disinhibition motif provides positive feedback that stabilizes the selected output[35]. Using inhibitory rather than excitatory connection may have the advantage of allowing the decisions to be influenced by contextual and state information. In addition, in this circuit architecture, the feedback inhibitory neuron Handle-b contacts both reciprocally connected inhibitory neurons in the circuit and can thus shape the circuit activity at the level of the site of competition. It is thus well suited to be the site of integration of mechanosensory information and information about an animal's state.

The inhibitory neurons shape the activity of projection neurons to bias behavioral decisions

The behavioral response of the larvae to a mechanical cue depends on the state of the circuit at the level of reciprocally interconnected inhibitory neurons, which will shape the activity of the Basin projection neurons to either give rise to a state where Basin-1 only is active or a state where Basin-1 and -2 are co-active[35]. We showed that the feeding state-dependent modulation of the Basin-2 neuron is dependent on the modulation of the activity of Handle-b and Griddle-2. Accordingly, genetic co-labeling of Basin-1 or Basin-2 neurons with either sNPFR1 or NPFR1 using the T2A GAL4/LexA technology showed no expression of these neuropeptide receptors in the two Basin neurons. This is in line with the finding that the state-dependent changes in the air puff induced sensorimotor decisions are caused by the modulation that acts on the reciprocally interconnected inhibitory neurons that integrate mechanosensory information with current internal state needs.

Feedback inhibitory neurons receive input from an NPF expressing neuron

Previous work revealed that the inhibitory neurons in the mechanosensory circuit are contacted by long-range projection neurons[35]. These long-range projection neurons may carry contextual or state information to the mechanosensory circuit. We found that among these long-range projection neurons is a pair of NPF-releasing neurons that contacts the Handle-b. NPF is a homologue of the mammalian neuropeptide Y that is a hunger signal. NPF neurons have indeed been shown to be involved in hunger dependent behaviors in both adult flies and larvae[27,47]. We found that the descending pair of NPF neuron activity changes as a function of the feeding state. These neurons could thus convey the information about the satiation state to the mechanosensory circuit and bias the sensorimotor decisions to mechanosensory cues by modulating the activity of the Handle-b. Additionally, the release of NPF at the proximity of the circuit by the NPF neurons (as revealed by the existence of dense core vesicles) could modulate Handle-b that expresses the NPFR (while other neurons in the circuit do not). The fact that the feedback inhibitory neuron

is directly influenced by the NPF neurons supports the idea that it could gate circuit activity in a state dependent manner.

Since Handle-b expresses the NPFR1 and also receives direct input from the NPF descending neurons, the NPF neurons could facilitate Handle-b activity either by direct synaptic transmission, by neuropeptide modulation or both. Our results indicate that the feeding state induced changes in motivation locomotor state and exploration persistence are dependent on the NPF from a pair of descending neurons while the acute response to transient environmental stimuli could rely on direct synaptic transmission (by the same neurons). Thus, this neuron could couple various sensorimotor behaviors to the motivational/exploratory state of the animal in a state dependent manner and thus regulate the different behaviors and different timescales

Thus, in the larva, the descending NPF neurons could convey information about the internal state to diverse neuronal populations in the VNC to regulate exploratory locomotion and stimulus-dependent motor responses, thus adjusting behavioral interactions with the environment according to the current animal needs.

Material and methods

Drosophila rearing and handling

Flies (*Drosophila Melanogaster*) were raised on a standard food medium (ethanol 2%, methylhydroxybenzoate 0.4%, yeast 8%, cornmeal 8%, and agar 1%) at 18°C. Third instar larvae were collected as follows: male and female flies from the appropriate genotypes were placed together for mating, then transferred at 25°C for 12-16 h on a petri dish containing a fresh food medium for egg laying. The petri dish was then placed at 25°C for 72 h. Foraging third instar larvae were collected from the food medium by using a denser solution of 20% sucrose, scooped with a paint brush into a sieve and gently and quickly washed with water. Larvae used for optogenetic experiments were raised at 25°C in complete darkness, on standard food supplemented with all-trans retinal at 0.25 mM (R240000, Toronto Research Chemicals). The full list of genotypes used in the study can be found in the Supplementary Material 1 Resource table.

Dietary treatments

For dietary treatments, larvae were placed in 60x15 mm circular petri dishes that contained a 45 mm circular Whatman paper. Larvae were subjected to different diets: standard food without agar for "fed" larvae (as described in the Drosophila rearing section), 20% sucrose solution for "protein deprived" larvae, and water for "starved" larvae. The Whatman paper was soaked with 0.6 mL MilliQ water ("starved"), sucrose solution ("fed on sucrose"), or soaked with 0.6 mL water and 1 mL of standard food medium was added on top ("fed"). Larvae were collected after the appropriate amount of time (90 or 300 min) and rinsed in water before behavioral, imaging, or biochemistry experiments. For the behavioral experiments with rehydration, larvae were collected after 90 minutes, rinsed in water, and placed in a petri dish with a Whatman paper soaked with 0.6 mL water for 15 minutes. Likewise, for the refeeding experiments, starved larvae were collected after 300 minutes, rinsed, and placed for 15 minutes in a petri dish with standard food medium and

water as in the "fed" condition. After the treatment, larvae were once more collected and rinsed before the experiment.

Behavioral tracking

We used an apparatus previously described[44,45]. Briefly, the apparatus comprises a video camera (Basler ace acA2040-90 **µ**m) for monitoring larvae, a ring light illuminator (Cree C503B-RCS-CW0Z0AA1 at 624 nm in the red), a computer and a hardware module for controlling air puff, controlled through multi worm tracker (MWT) software (http://sourceforge.net/projects/mwt)[45,66]. The arena consisted of a 25625 cm2 square of 3% Bacto agar gel (CONDALAB 1804-5) with charcoal (Herboristerie Moderne, 66000 Perpignan) in a plastic dish, and was changed for each experiment. For optogenetic experiments, plates without charcoal were used, and larvae were tracked thanks to IR light. Collected third instar larvae were washed with water, moderately dried and spread on the agar starting from the center of the arena. We tested approximately 30–100 larvae at once during each experiment. The temperature of the behavioral room was kept at 25˚C.

Locomotion analysis

To assess larval locomotion in the absence of stimulation, larvae were placed on top of the agar in the arena inside the tracker, and either tracked for 5 min continuously (wild type larvae) or for 60s. The coordinates of the 11 points along the central spine and the outline of each larva were computed as described previously[44,45]. These 2D X-Y coordinates, structured as time series of irregular framerate, labeled by the instantaneous tracking time of the recording, comprise the raw datasets which have been subsequently analyzed to derive all secondary metrics.

Analysis was performed in python using the *larvaworld* behavioral analysis and simulation platform (https://pypi.org/project/larvaworld/). The 3-step analysis pipeline included preprocessing, computation of secondary angular, translational and temporal metrics and behavioral epoch detection to annotate strides, runs and pauses, as described previously[67]. During preprocessing the raw time series were adjusted to a 10 Hz constant framerate by interpolating them at a 0.1 second timestep. Noise reduction was achieved by applying a low-pass filter with a 2Hz cut-off frequency, a threshold high enough not to alter the crawling-related dynamics around the dominant ~1.4Hz crawling frequency.

For trajectory-based spatial metrics such as pathlength and dispersal, to avoid the cumulative effect of body micromovements, the position of the 9th point along the midline was used as a proxy for the larva's position, a relatively stable rear point unaffected by lateral and translational jitter. To correct for different larval sizes, any metric measured in absolute spatial units (m or mm) can be scaled to body-length, measured in dimensionless body-length units. As the instantaneous body-length of an individual larva fluctuates during crawling due to subsequent stretching and contraction, individual larva length is defined as the median of the midline length across time (total length of the line connecting all 11 midline points). A trajectory's pathlength is the cumulative displacement of the larva during the entire track. Dispersal is the instantaneous straight-line distance relative to its initial position. Track tortuosity was quantified by the straightness index (S.I), computed by advancing a fixed time window (20 seconds in this study) along the track and

calculating at each point the ratio of the dispersal to the actual distance traveled. This index, which varies from 0 (no movement) to 1 (straight line movement), can capture very well the complexity of the movement at various scales (set by the window time frame) throughout the track.

For the detection of peristaltic strides and crawl-pauses, the scaled crawling speed time series were used. To this end the dominant crawling frequency for each track was extracted by applying a Fourier analysis and its inverse was used as the expected duration of a peristaltic cycle. A stride was therefore defined as the epoch between two local speed minima, that included a local maximum of at least 0.3 body-lengths/s and lasted between 0.7 and 1.5 times the expected cycle duration. A run was defined as an uninterrupted sequence of consecutive strides and a crawl-pause as an epoch lacking any strides during which the scaled speed was constantly below the 0.3 body-lengths/s threshold.

Air puff stimulation during behavioral tracking

Air puff was delivered as described previously[35,44,45] to the 25625 cm2 arena at a pressure of 1.1 MPa through a 3D-printed flare nozzle placed above the arena, with a 16 cm x 0.17 cm opening, connected through a tubing system to plant supplied compressed air. The strength of the airflow was controlled through a regulator downstream from the air amplifier and turned on and off with a solenoid valve (Parker Skinner 71215SN2GN00). Air-flow rates at 9 different positions in the arena were measured with a hot-wire anemometer to ensure consistent coverage of the arena across experimental days. The air-current relay was triggered through TTL pulses delivered by a Measurement Computing PCI-CTR05 5-channel, counter/timer board at the direction of the MWT. The onset and duration of the stimulus were also controlled through the MWT. Larvae were left to crawl freely on the agar plate for 60 seconds prior to stimulus delivery. Air puff was delivered at the 60th second and applied for 30 seconds.

Behavior classification and analysis

Behaviors were detected thanks to a custom-made machine learning algorithm that was previously described[44]. Behaviors were defined as mutually exclusive actions. Larvae were tracked using MWT software, all the time series of the contours and the spine of individual larvae are obtained using Choreography. From these times series some features are computed, center of the larva, velocities, etc., all key features are presented in Masson *et al.*, 2020. Behavior classification consists of a hierarchical procedure that were trained separately based on a limited amount of manually annotated data. Here, we required a more detailed definition of behavior. Hence, we extended the hierarchy with another layer to separate some Bends and Hunches, between different behaviors. Bends were separated into Static Bend, Head-Cast (see the description of each behavior below). We take all the Bends, Hunches and Back-ups obtained by the first classification algorithm, to reclassify with new annotated data on new lines.

Action definition
Head Cast: dynamic bends in which the head moves laterally from one side to the other. There are two exits from Head-Cast in both the head moves strongly, but one that is slower and the tail

moves at the same speed as the center of mass, and on the contrary a second one where the tail moves a lot (fast Head-Cast).

Static Bend: Low speed turning movement, and where the head moves little, and the angle between the segment between the center of mass and the head and the center of mass and the tail remains constant.

New annotated data.
We required new annotated data to ensure the classifier matched the phenotype of the larva used in these experiments. The sets of, Hunch, Bends, Backs and Crawls were manually tagged from actions selected using the Masson *et al, 2020* old behavior classification pipeline. A few numbers of tags are used for the model.

New features.
To train the new layer of the pipeline, we combined features that were previously evaluated in the classifier's preceding layer with new features. Each characteristic is calculated for the time step we are examining and the three time steps before and after.

- The three velocities include the head velocity, the motion velocity, and the tail velocity, all normalized by the length of the larva. The motion velocity is the velocity of center of mass return by the MWT software. Head and tail are the terminal point of the spine; The averaging along the spine curve and its derivative, $S = \frac{1}{2}(3 \langle cos^2\theta \rangle - 1)$ with $cos\,\theta$ the scalar product between normalized vectors associated to a segment of the spine and the direction of the larva body.
- The shape factor $\lambda = \frac{\lambda_1 - \lambda_2}{\lambda_1 + \lambda_2}$ with λ_i the eigenvalues of the mean covariance matrix of movement which characterizes the shape of the larva and takes value between 0 and 1.

We have also introduced new features:

- The ratio between the length of the head-center of mass and the tail-center of mass $\frac{||\overrightarrow{HG}||}{||\overrightarrow{TG}||}$ with H, T and G respectively coordinate points of the head, the tail and the center of the masse
- The projection of the head and tail velocity on the spine of the larva. If we note the velocity vector of the head HV_h with H coordinate point of the head and V_h the coordinate point at the end of the velocity vector, the projection point satisfies the basic relationship: $||V_h - V'_h|| = min\,||V_h - x||$ with $x \in \overrightarrow{r_{si}}$.
- The cosine of the angle between the vector of the head (tail) velocity and the first (last) segment of the larva. $cos(\theta) = \frac{\overrightarrow{r_{s1}} \cdot \overrightarrow{v_{head}}}{||\overrightarrow{r_{s1}}||\,||\overrightarrow{v_{head}}||}$ with $\overrightarrow{v_{head}}$ the vector velocity of the head and $\overrightarrow{r_{s1}}$ the first vector of the spine.

All features are normalized by the length of the larva to ensure scale-free properties.

New classifications

We employed a Random Forest algorithm for classification, distinguishing each time step of original Bends, Hunches, and Backs into new categories. We conducted ten random forests on all Hunch and back tags, along with a random selection of a thousand bends, utilizing balanced weights. The predicted behavior represents the most probable outcome, with each random forest's confusion matrix exceeding 80% accuracy for each behavior.

To address the issue of having a behavior span only two time steps, which is not biologically plausible, we introduced a preventive measure by adding five time steps before and after the behavior. Additionally, to mitigate noisy results, we implemented smoothing. This involved excluding behaviors with fewer than three time steps, logically categorizing them as the behavior before or after (based on the length of the behavior and behaviors N+2 and N-2). According to our knowledge, Hunch behavior initiates at the beginning of stimulation, around 60 second[35,44]. Larvae typically do not exhibit multiple Hunches. In cases where they do, we classify the second Hunch as a cast, a classification verified through ground truthing. We applied the same threshold as outlined in Masson *et al.,* 2020 to the effective length change during the behavior. If a Hunch fails to surpass this threshold, the time window is assigned to the small behavior. The threshold is not the same depending on the line, some lines were slower or smaller than others (threshold between 0.6 and 0.3). The validation of these thresholds was performed through ground truthing, contributing to the enhancement of classification, particularly in cases where performance may be suboptimal for certain lines.

The distinction between Static Bends and Head-Cast is determined by applying a threshold to the head velocity. If bending occurs over 'n' time steps, the motion velocity normalized by the length of 'P%' of those 'n' time steps must be below 'p' times the mean head velocity of the larva before the stimulation. The values of the two thresholds, 'P' and 'p', are line-specific, contingent upon the statistical characteristics of the velocity for that particular line; some lines are slower than others.

We computed the cumulative probabilities of actions (Stop, Hunch, Back-up) during five seconds after stimulus onset (as described in Masson *et al,* 2020), only in larvae that were tracked during the entire time window. For actions that occur during baseline locomotion and at high frequency (Crawl and Head Cast), we computed the mean probability over five seconds after stimulus onset and over three seconds starting one second after stimulus onset. We corrected the mean probabilities for these actions it by the mean probability computed over twenty seconds of recording prior to stimulus onset. For optogenetic activation of the mechanosensory neurons, because the dynamic of the response was different to that of air puff experiments, the time window used for computing the cumulative probability of Hunching was two seconds after the stimulus onset and that for bending probability was ten seconds after stimulus onset, by which time bending probability reached baseline levels in larvae fed on standard food. Transition probabilities were computed as the frequency of transition from one action to another over five seconds after stimulus onset, for larvae that were tracked throughout this entire duration.

Food intake quantification

In order to measure food intake, we quantified the amount of fluorescent food inside the digestive tract of larvae that were allowed to feed ad libitum for 15 min on fluorescent feeding media. To

this end, Rhodamine B (Sigma R6626-25G) was diluted in different feeding media (water, yeast extract, sucrose solution or normal food medium, see in media section) to a final concentration of 0.20 µmol/L. 0.8 ml L of each food medium was poured on top of a circular Whatman paper (Fisher scientific, Cytiva 1001-045) placed into a petri dish. 10 larvae were placed into each petri dish. After 15 min of ad libitum feeding on the fluorescent medium, larvae were collected, rinsed in ethanol and in water, and immediately mounted under a coverslip for imaging. Intact larvae were imaged thanks to a fluorescent binocular Zeiss Discovery.V12. The surface of the digestive tract stained by the fluorescent dye was quantified for each larva thanks to a custom-made Fiji script. This script quantified the number of pixels whose intensity was above background.

Sucrose preference

To measure the preference of larvae between a sucrose-containing agar and a water-containing agar, we performed a place-preference assay. To this end, we prepared petri dishes filled with 0.3% agar diluted in water, divided each agar in two halves, and transferred one half into a new petri dish. Then, we filled the missing half in each petri dish with 0.3% agar diluted in a 20% sucrose solution. Therefore, each petri dish finally contained one side with 20% sucrose agar and one side with agar only.
After cooling down, about 20 third instar larvae were put on the midline of a petri dish and the dish was imaged every 30 s in a behavioral tracker. The number of larvae on each side was then counted over time, excluding the larvae touching the limit between the two media. The preference index was then calculated as : PI = (Ns - Na)/Ntot, where Ns is the number of larvae on the sucrose side, Na the number of larvae on the agar side, and Ntot the total number of larvae.

Carbohydrate measurements

To measure carbohydrates concentrations inside larval hemolymph, we combined and adapted different methods already published[68,69]. Glucose measurements - Groups of 5 third instar larvae were rinsed in water and placed on a parafilm layer and their cuticle was cut with forceps. 2 µL of the bleeding hemolymph was collected from each group, and 1 µL of 0.05 g/L N-phenylthiourea diluted in PBS was added to avoid darkening of the samples. Samples were heat-inactivated by a 10 min incubation at 90°C and centrifuged 10 min at 10 000 rpm. 1 µL of the supernatant was then mixed with 4 µL of glucose assay kit (Sigma GAHK20-1KT) and incubated for 1 h at 37°C. Absorbance at 340 nm was measured against the blank thanks to a NanoDrop following the manufacturer's instructions. Hemolymph glucose concentration was finally calculated thanks to a standard curve of glucose concentration.
Trehalose measurements - Groups of 10 third instar larvae were rinsed in water and placed on a parafilm layer and their cuticle was cut with forceps. 3 µL of the bleeding hemolymph was collected from each group, and 97 µL of 0.05 g/L N-phenylthiourea diluted in trehalase buffer (Tris pH 5.5 5 mM, NaCl 137 mM, KCl 2.7 mM) was added to avoid darkening of the samples. Samples were heat-inactivated at 70°C for 10 min, centrifuged for 10 min at 10 000 rpm, and 5 µL of the supernatant were either mixed to 5 µL of trehalase (Sigma T8778-1UN) diluted in trehalase buffer (described above) to a 500 dilution. Samples were incubated at 37°C overnight. 1 µL of each sample was then mixed with 4 µL of glucose assay kit (Sigma GAHK20-1KT) and incubated for 1

h at 37°C. Absorbance at 340 nm was measured against the appropriate blank thanks to a NanoDrop following the manufacturer's instructions. Trehalose concentration was finally calculated thanks to a standard curve of glucose and trehalose concentrations and by additionally subtracting the concentration of glucose in the sample without trehalase.

Histochemistry labeling

To determine the neurotransmitter identification in the interneurons, immuno-labeling was performed from the split lines or Gal4 lines crossed to UAS-myr::GFP, or LexA lines crossed to LexAop-myr::GFP. The VNC was dissected out from 3rd instar larvae and fixed with 4% PFA for 45 min at room temperature. After rinsing in PBS, ten minutes permeabilization in PBS-T and two hours blocking in PBS-T-BSA 1%, the CNS preparations were incubated at 4°C (one to three nights) in the first antibodies raised against neurotransmitter and GFP in PBS-T. Then they were incubated at 4°C (one to two nights) in fluorophore-coupled secondary antibodies in PBS-T raised against species of the first antibodies. After rinsing, the preparations were mounted in an anti-bleaching mounting medium (SlowFade Gold, ThermoFisher S36939) under a cover slip. The confocal images were captured with a Leica SP8 confocal laser microscope. Alexa Fluor 488 was excited with a laser light of 488 nm, Cy3 with a laser light of 561 nm, Alexa Fluor 647 with a light of 633 nm wavelength.

Neuropeptide receptor characterisation

In order to characterize the expression pattern of sNPFR and NPFR in the circuit, expression of two genetically encoded reporter proteins of two different colors was targeted to two different subsets of neurons. To this aim, a T2A Gal-4[48] or LexA (for sNPFR and NPFR respectively) was used to express LexAop-jRGeco1a or UAS-GCaMP6s (for sNPFR and NPFR respectively) under the control of the promoter of the gene coding for that receptor, thus tagging all neurons which express the receptor transcript. A second genetic driver (LexA for sNPFR and Gal4 for NPFR) was used to individually label target neurons of the circuit with a second reporter protein (UAS-GCaMP6s for sNPFR and LexAop-jRGeco1a for NPFR). The VNC was then dissected and stained with antibodies as described in the previous section.

To the best of our knowledge, no clean or sparse LexA line exists to selectively target the Handle-B and Griddle-2 inhibitory interneuron. For sNPFR expression in Griddle-2, the LexA line L55C05 used targets many neurons in addition to Griddle-2. Griddle-2 could nevertheless be identified by comparing the cell body position and projections in the cross-section of the anterior abdominal segments of the CNS of the R55C05 LexA line and the sparse R55C05 GAL4 line that selectively labels griddle-2 in the VNC (see Figure S7).

For Handle-B we used a L60E02-LexA line for which a neuron with a cell boy in the midline resembling Handle-B is part of a very dense expression pattern. In order to confirm that the candidate enron was indeed Handle-B, we used the selective split-Gal4 line GMR_SS00888 that specifically labels only Handle-B neurons. We expressed two reporters of different colors (LexAop-GCaMP6s and UAS-Chrimson-mCherry under the control of GMR_SS00888 and L60E02 in the same larva. The colocalization confirmed the line 60E02-LexA to target Handle-B (Figure S7).

In vivo imaging of intact larvae

Because opening the cuticle might affect the larval internal state, we developed a simple preparation for the imaging of intact larvae. For this purpose, third instar larvae were rinsed in water, and mounted between a 2 cm circular coverslip and a custom-made device that delivers mechanical stimulations in low melting point agarose 4% (melted in phosphate buffer saline), ventral side facing up. Larvae were gently squeezed in this position until agar cooled down, so that the ventral nerve cord could be imaged through the cuticle.

All Gal4 and LexA drivers used for *in vivo* imaging are listed in the figure legends. The imaging plane was restricted to the location of the projections of neurons of interest, in particular when sparse lines that lacked specificity towards a unique neuronal type were used (R22E09).

Mechanical stimulations were generated by a waveform generator (Siglent sdg1032x) connected to a quick-mount extension actuator (Piezo Systems, Inc.), which was embedded in the sylgard-coated recording chamber (Sylgard Silicone Elastomer, WPI). The stimulation was set at 1000 Hz, with an intensity of 1 to 20 V applied to the actuator. The amplitude of the acceleration produced by the actuator was measured thanks to a triple axis accelerometer (Sparkfun electronics ADXL313) connected to a RedBoard (Sparkfun electronics) and bound to the sylgard surface thanks to high vacuum grease. Acceleration was 1.14 $m.s^{-2}$ at 20 V, and 0.61 $m.s^{-2}$ at 10 V. Mechanical stimulations were precisely triggered by the Leica SP8 software thanks to the Leica "Live Data Mode" and to a trigger box branched to the scanning head of the microscope. A typical stimulation experiment consisted in 5 s of recording without stimulation, then 5 s of stimulation, and 5 s of recording in the absence of stimulation.

For optogenetic activation during in vivo imaging, larvae were mounted in the dark, with the least intensity of light possible in the room, to avoid nonspecific activation of the targeted neurons. Optogenetic stimulation of CsChrimson was achieved by a 617-nm wavelength LED (Thorlabs, M617F2), controlled by a LED driver (Thorlabs, LEDD1B) connected to the waveform generator, and conveyed through a Ø 400 µm Core Patch Cable (Thorlabs) to the imaging field. Optogenetics stimulations were triggered at 50 Hz, 50% duty during 1 s, concomitantly to mechanical stimulations thanks to the waveform generator. Irradiance was measured at the level of the imaging field at 500 µW using a PM16-130 THORLABS photometer.

Imaging was achieved with 1-photon or 2-photon scanning Leica SP8 microscope, at 200 Hz, with a resolution of 512 x 256 pixels or 512 x 190 pixels. The rate of acquisition was 1 frame/s or 2 frames/s depending on the experiment. For Basin-2 recordings, the stimulation was repeated 5 times with resting intervals of 60 s in order to calculate a frequency of response. Optogenetic experiments were conducted with 2-photon imaging.

Neuronal processes were imaged in the VNC at the axonal level and fluorescence intensity was measured by manually drawing a region of interest (ROI) in the relevant areas using custom Fiji macros. Data were further analyzed using customized MATLAB scripts. F0 was defined as the mean fluorescence in the ROI during baseline recording, in the absence of mechanical stimulus or optogenetic activation. ΔF/F0 was defined at each time step t in the ROI as: ΔF/F0 = (F(t) - F0)/F0.

For recording the baseline activity level of NPF neurons, one frame was recorded each second for 20 seconds and, for each larva, the frame showing the most intense fluorescence in the neuronal projections was used to evaluate its raw fluorescence level.

For imaging neurons in different feeding conditions, the effect of food treatment on the expression level of fluorescent reporter proteins was assessed by expressing GFP in the chordotonal neurons. Fluorescence was measured after exposing larvae to different food treatments.

Statistical analysis

Locomotion analysis

For all boxplots and histograms pairwise Mann-Whitney tests were used to evaluate significant differences between larva groups, with Bonferroni correction for multiple comparisons. Significance was illustrated according to the p-value by asterisks in histograms (*:<0.05, **:<0.01, ***:<0.001, ****:<0.0001) and by pairs of colored semicircles in boxplots, the left always corresponds to the group with highest mean value. Non-significant tests were omitted for visual clarity.

Behavior probabilities

Chi² tests were used for statistical analysis of behavioral and transition probabilities.

To assess the effects of different states/neuronal manipulations on Head Casting in response to air puff, we calculated an estimator designed to identify the emergence of behaviors at the population scale. We aim to determine the probability induced by the stimulation, so we need to subtract the stationary probability without the stimulus. We calculate the probability of larvae bending 5 seconds after the stimulus, on tracking larvae throughout this time window (denoted as p_A). For the probability of bending before the stimulus, we consider all larvae tracked continuously for 20 seconds prior to the stimulus, between 30 and 50 seconds (denoted as p_B). A probability is defined as $p_k = N_k/N_{k,all}$ with $k \in \{A, B\}$ and $N_{k,all}$ the total number of larva taking to compute probabilities. In order to quantify the effect of the stimulus we defined $\chi = p_A - p_B$ as the difference in the ratio after and before the stimulus.

In order to compare test line our estimator was defined as

$\Theta(p, q) = \chi(p) - \chi(q)$ with p and q the ratios of the lines and the control respectively. $\Theta(p, q)$ takes value in $[-1,1]$. The null hypothesis is $\Theta(p, q) = 0$, if there are no differences between the line tested and the control. Positive or negative values indicate an effect of neuron silencing when compared to the control.

We use numerical simulations to conduct a statistical test where $\{p_k, q_k\}$ are generated from a hypergeometric distribution, $X \sim \boldsymbol{Hypergeometric}(\boldsymbol{N_{k,all}, N_k, 1})$. We perform $N_{sim} = 10^3$ repetitions, computing $\Theta(p, q)$ each time. The p-value is determined by the number of instances when the hypothesis is not verified, divided by the total number of repetitions, (pseudo-code in Jovanic *et al*, 2016[35]).

Note that this estimator, $\Theta(p, q)$, has the advantage of being able to detect the non-synchronous emergence of a behavior at a population scale. For example, head casting can either emerge as an immediate response to the puff or as the second response after Hunching. The statistics of start time of Head Casting is thus widely distributed at the population scale. Time evolution of the

instantaneous ratio of larva performing bending would not exhibit a strong increase after stimuli because larvae are not all going to bend immediately after stimuli. $\Theta(p, q)$ by accumulating events during a time window allows efficient detection of a behavior even if it is widely distributed in time.

Calcium imaging

All data in line plots are presented as mean ± SEM. Violin plots show the first and third quartiles, the average of all recordings as a white line and the median as a white dot. Comparisons of the data series between two conditions were achieved by a two-tailed unpaired t-test. Comparisons between more than two distinct groups were made using a one-way ANOVA test, followed by Bonferroni pairwise comparisons between the experimental groups and their controls.

Mathematical model

We reproduced and extended the rate-based system model of the circuit that was published in a previous publication[35]. The circuit is described as a rate model with a connection matrix derived from the larva connectome. Each neuron population (mechano-ch, iLNa, iLNb, fbLN-Ha and fLN-Hb) was modeled by a single node (Figure 3C-E). The dynamics read:

$$\tau \frac{d\mathbf{r}}{dt} = -\mathbf{V_0} - \mathbf{r} + \mathbf{s} + i + k_{ex} \left(\mathbf{r} - \mathbf{r}^{\max} \right) \left(\underline{A}^{ex}\mathbf{r} - \underline{A}^{in}\mathbf{r} \right) \quad (1)$$

with $\boldsymbol{\tau}$ representing the vector of the characteristic time constants of the neurons, **r** the rate vector, $\mathbf{V_0}$ the threshold vector, **s** the sensory stimulus input vector, **i** the vector of inputs from other brain regions, k_{ex} a sensitivity factor to overall input, $\mathbf{r}^{max}$ the maximal rate vector, $\underline{A}^{ex}$ and $\underline{A}^{in}$ respectively the excitatory and inhibitory coupling matrices.
Values in $\underline{A}^{in}$ and $\underline{A}^{ex}$ were directly extracted from synaptic counts (see Jovanic *et al.,* 2016).
In order to represent the variety of stimuli larvae are subjected to, and thus the variety of behavior they elicit, we follow the approach in Jovanic *et al.*, 2016 and vary the connection strength between Mch and iLNa populations. In the original paper, the connection strength between Mch and iLNb populations is also varied. We decided to fix the value of this connection strength at 2 in order to reduce the number of parameters to explore.

We used the *solve_ivp* routine of the integration package from SciPy, which internally calls the LSODA solver, able to switch between the Adams method and the BDF method, based on the stiffness of the equation. We used relative and absolute tolerances of 10^{-3}. Additionally, the solution vector is constrained to stay positive. This is obtained by replacing r by max(r, 0), and the components of r' by those of max(r', 0) whenever the corresponding component of r is smaller than 10^{-9}.

The behavior is defined based on the end state of the network and, more specifically, the rates of the neurons B1 and B2. We used a k-means clustering with k=2 to separate the values of the output neurons for an ensemble of simulations corresponding to connection strengths spanning

[0.5, 1.5] between MCh and iLNa, and [1.5, 2.5] between iLNa and iLNb. The output activations cluster strongly in a coactive state (rate(B1) > 0, rate(B2) > 0) corresponding to bends, and a monoactive state (rate(B1) > 0, rate(B2) = 0) corresponding to Hunches. The two behaviors can also be distinguished by the single scalar rate(B2)/rate(B1), which is positive for bends and zero for Hunches. This ratio is sometimes plotted instead of the discrete category.

We explore two models for neuromodulation, which modify the behavior output of the network without altering its connectivity.

The first model hypothesizes that in the sucrose state, Hb receives an additional input current, modeled by a nonzero entry to the vector **i**. We show that as this input current increases, the range of stimuli evoking bends increases, allowing us to claim that an additional input current to Hb increases the likelihood of bends. We further fix the value of the input current to 10 a.u. in the sucrose state and 0 a.u. in the fed state, to perform silencing analyses.

The second model hypothesizes that in the sucrose state, the maximum rate for the iLNa neuron population is decreased, representing a saturation of the response to external stimuli. We show that as the r_{max} parameter for iLNa decreases, the range of stimuli evoking bends increases, consistent with experimental observations and once again despite the use of arbitrary units. For silencing analyses, we define the sucrose state as r_{max} = 18 a.u. and the fed state as r_{max} = 20 a.u. Finally, we also consider a model combining both hypotheses. In this model, every combination of a decrease in r_{max} for iLNa and an increase in input current to Hb results in more bend. For silencing analyses, we fix the exact values of those parameters. In the combined model, we define the sucrose set by setting each parameter to the value defining the sucrose state in the single hypothesis models.

We provide here the list of parameters used in the simulations.

k_{ex}	2.5
i	[0, 0, 0, 0, 0, 0, 0] in fed state [0, 10, 0, 0, 0, 0, 0] in sucrose state for the combined model
r_{max}	[20, 20, 20, 20, 20, 20, 20] in fed state [20, 20, 20, 20, 20, 20, 18] in sucrose state for the combined model
V_0	[0, 20, 20, 20, 20, 20, 20]
τ	[1, 35, 35, 35, 35, 35, 35]

The excitatory and inhibitory matrices read

$$A_{ex} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1.5 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0.75 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 0 & 0 & 0 & 0 & 0 & 0 \\ W_{iLNa} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0.2 & 0.2 & 0 & 0 & 0 & 0 \\ 0.4 & 0 & 0.5 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$A_{in} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1.7648 & 1.3841 & 0 & 0 \\ 0 & 0 & 0 & 1 & 5.9167 & 0 & 0 \\ 0 & 0 & 0 & 0 & 3.3744 & 1.6659 & 2.191 \\ 0 & 0 & 0 & 2.7133 & 0 & 1.1010 & 3.3031 \\ 0 & 0 & 0 & 1.8411 & 1.1158 & 0 & 0 \\ 0 & 0 & 0 & 1.7331 & 2.2145 & 0 & 0 \end{bmatrix}$$

Author contributions, E.T. behavioral and physiology experiments and analysis, Calcium-imaging experiments and analysis; Writing: figures, edits and revisions; D. M. behavioral experiments and analysis, Calcium-imaging experiments and analysis, immunohistochemistry experiments, Writing: original draft, figures; A.B. modeling; P.S. Data analysis; C.B. behavioral classification and analysis, statistical analysis, writing, methods; F. V. behavioral experiments; V. S. behavioral and physiology experiments; A.H. and S. A immunohistochemistry experiments; M.N. supervision, funding acquisition; J.B. Methodology, supervision, funding acquisition, writing-edits and revision; T.J. Conceptualization, analysis, supervision, funding acquisition and project administration; writing: original draft, figures, edits and revision

Acknowledgments

Funding. This work was supported by ANR PIA funding: ANR-20-IDEES-0002 (T.J), Agence Nationale de la Recherche: ANR-17-CE37-0019-01 (T.J.), ANR-NEUROMOD (ANR-22-CE37-0027) (T.J.), ANR-NEUC-0002-01 (T.J), Fondation de recherche sur le cerveau (FRC) (T.J.). Fondation de Treille (E.T.), D. M. received a PhD fellowship from the Paris-Saclay University; Tramway, ANR-17-CE23-0016 (J.B.M), the inception Project PIA/ANR-16-CONV-0005,OG (J.B.M), Investissement d'avenir programme under the management of ANR, ANR-19-P3iA-0001 (PRAIRIE 3IA Institute (J.B.M), German Federal Ministry of Education and Research (BMBF DrosoExpect, 01GQ2103A, M.N.), Ministry of Culture and Science of the State of Northrhine

References

1. Allen, W.E., Chen, M.Z., Pichamoorthy, N., Tien, R.H., Pachitariu, M., Luo, L., and Deisseroth, K. (2019). Thirst regulates motivated behavior through modulation of brainwide neural population dynamics. Science *364*, 253. 10.1126/science.aav3932.

2. Anderberg, R.H., Hansson, C., Fenander, M., Richard, J.E., Dickson, S.L., Nissbrandt, H., Bergquist, F., and Skibicka, K.P. (2016). The Stomach-Derived Hormone Ghrelin Increases Impulsive Behavior. Neuropsychopharmacol. Off. Publ. Am. Coll. Neuropsychopharmacol. *41*, 1199–1209. 10.1038/npp.2015.297.

3. Burnett, C.J., Li, C., Webber, E., Tsaousidou, E., Xue, S.Y., Brüning, J.C., and Krashes, M.J. (2016). Hunger-Driven Motivational State Competition. Neuron *92*, 187–201. 10.1016/j.neuron.2016.08.032.

4. Eiselt, A.-K., Chen, S., Chen, J., Arnold, J., Kim, T., Pachitariu, M., and Sternson, S.M. (2021). Hunger or thirst state uncertainty is resolved by outcome evaluation in medial prefrontal cortex to guide decision-making. Nat. Neurosci. *24*, 907–912. 10.1038/s41593-021-00850-4.

5. Jourjine, N. (2017). Hunger and thirst interact to regulate ingestive behavior in flies and mammals. BioEssays *39*, 1600261. 10.1002/bies.201600261.

6. Jourjine, N., Mullaney, B.C., Mann, K., and Scott, K. (2016). Coupled Sensing of Hunger and Thirst Signals Balances Sugar and Water Consumption. Cell *166*, 855–866. 10.1016/j.cell.2016.06.046.

7. Smith, N.K., and Grueter, B.A. (2021). Hunger-driven adaptive prioritization of behavior. FEBS J. 10.1111/febs.15791.

8. Inagaki, H.K., Panse, K.M., and Anderson, D.J. (2014). Independent, reciprocal neuromodulatory control of sweet and bitter taste sensitivity during starvation in Drosophila. Neuron *84*, 806–820. 10.1016/j.neuron.2014.09.032.

9. Ko, K.I., Root, C.M., Lindsay, S.A., Zaninovich, O.A., Shepherd, A.K., Wasserman, S.A., Kim, S.M., and Wang, J.W. (2015). Starvation promotes concerted modulation of appetitive olfactory behavior via parallel neuromodulatory circuits. eLife *4*, 691. 10.7554/eLife.08298.

10. Loch, D., Breer, H., and Strotmann, J. (2015). Endocrine Modulation of Olfactory Responsiveness: Effects of the Orexigenic Hormone Ghrelin. Chem. Senses *40*, 469–479. 10.1093/chemse/bjv028.

11. Sayin, S., Boehm, A.C., Kobler, J.M., De Backer, J.-F., and Grunwald Kadow, I.C. (2018). Internal State Dependent Odor Processing and Perception—The Role of Neuromodulation in the Fly Olfactory System. Front. Cell. Neurosci. *12*. 10.3389/fncel.2018.00011.

12. Bräcker, L.B., Siju, K.P., Varela, N., Aso, Y., Zhang, M., Hein, I., Vasconcelos, M.L., and Grunwald Kadow, I.C. (2013). Essential Role of the Mushroom Body in Context-Dependent CO_2 Avoidance in Drosophila. Curr. Biol. *23*, 1228–1234. 10.1016/j.cub.2013.05.029.

13. Crossley, M., Staras, K., and Kemenes, G. (2018). A central control circuit for encoding perceived food value. Sci. Adv. *4*, eaau9180. 10.1126/sciadv.aau9180.

14. Filosa, A., Barker, A.J., Dal Maschio, M., and Baier, H. (2016). Feeding State Modulates Behavioral Choice and Processing of Prey Stimuli in the Zebrafish Tectum. Neuron *90*, 596–608. 10.1016/j.neuron.2016.03.014.

15. Padilla, S.L., Qiu, J., Soden, M.E., Sanz, E., Nestor, C.C., Barker, F.D., Quintana, A., Zweifel, L.S., Rønnekleiv, O.K., Kelly, M.J., et al. (2016). Agouti-related peptide neural circuits mediate adaptive behaviors in the starved state. Nat. Neurosci. *19*, 734–741. 10.1038/nn.4274.

16. Card, G.M. (2012). Escape behaviors in insects. Curr. Opin. Neurobiol. *22*, 180–186. 10.1016/j.conb.2011.12.009.

17. Evans, D.A., Stempel, A.V., Vale, R., and Branco, T. (2019). Cognitive Control of Escape Behaviour. Trends Cogn. Sci. *23*, 334–348. 10.1016/j.tics.2019.01.012.

18. Imambocus, B.N., Zhou, F., Formozov, A., Wittich, A., Tenedini, F.M., Hu, C., Sauter, K., Varela, E.M., Herédia, F., Casimiro, A.P., et al. (2022). A neuropeptidergic circuit gates selective escape behavior of Drosophila larvae. Curr. Biol. *32*, 149-163.e8. 10.1016/j.cub.2021.10.069.

19. Peek, M.Y., and Card, G.M. (2016). Comparative approaches to escape. Curr. Opin. Neurobiol. *41*, 167–173. 10.1016/j.conb.2016.09.012.

20. Ache, J.M., Namiki, S., Lee, A., Branson, K., and Card, G.M. (2019). State-dependent decoupling of sensory and motor circuits underlies behavioral flexibility in Drosophila. Nat. Neurosci. *22*, 1132–1139. 10.1038/s41593-019-0413-4.

21. Devineni, A.V., and Scaplen, K.M. (2022). Neural Circuits Underlying Behavioral Flexibility: Insights From Drosophila. Front. Behav. Neurosci. *15*, 350. 10.3389/fnbeh.2021.821680.

22. Oram, T.B., and Card, G.M. (2022). Context-dependent control of behavior in Drosophila. Curr. Opin. Neurobiol. *73*, 102523. 10.1016/j.conb.2022.02.003.

23. Schadegg, A.C., and Herberholz, J. (2017). Satiation level affects anti-predatory decisions in foraging juvenile crayfish. J. Comp. Physiol. A Neuroethol. Sens. Neural. Behav. Physiol. *203*, 223–232. 10.1007/s00359-017-1158-8.

24. Dickinson, P.S. (2006). Neuromodulation of central pattern generators in invertebrates and vertebrates. Curr. Opin. Neurobiol. *16*, 604–614. 10.1016/j.conb.2006.10.007.

25. Eriksson, A., Raczkowska, M., Navawongse, R., Choudhury, D., Stewart, J.C., Tang, Y.L., Wang, Z., and Claridge-Chang, A. (2017). Neuromodulatory circuit effects on Drosophila feeding behaviour and metabolism. Sci. Rep. *7*, 8839. 10.1038/s41598-017-08466-0.

26. Krasne, F.B., and Edwards, D.H. (2002). Modulation of the crayfish escape reflex--physiology and neuroethology. Integr. Comp. Biol. *42*, 705–715. 10.1093/icb/42.4.705.

27. Lin, S., Senapati, B., and Tsao, C.-H. (2019). Neural basis of hunger-driven behaviour in Drosophila. Open Biol. *9*, 180259. 10.1098/rsob.180259.

28. Marder, E. (2012). Neuromodulation of neuronal circuits: back to the future. Neuron *76*, 1–11. 10.1016/j.neuron.2012.09.010.

29. Taghert, P.H., and Nitabach, M.N. (2012). Peptide neuromodulation in invertebrate model systems. Neuron *76*, 82–97. 10.1016/j.neuron.2012.08.035.

30. Tierney, A.J. (2020). Feeding, hunger, satiety and serotonin in invertebrates. Proc. R. Soc. B Biol. Sci. *287*. 10.1098/rspb.2020.1386.

31. Lewis, L.P.C., Siju, K.P., Aso, Y., Friedrich, A.B., Bulteel, A.J.B., Rubin, G.M., and Grunwald Kadow, I.C. (2015). A Higher Brain Circuit for Immediate Integration of Conflicting Sensory Information in Drosophila. Curr. Biol. CB *25*, 2203–2214. 10.1016/j.cub.2015.07.015.

32. Jovanic, T. (2020). Studying neural circuit of decision-making in Drosophilalarva. J. Neurogenet. *55*, 1–9. 10.1080/01677063.2020.1719407.

33. Eschbach, C., and Zlatic, M. (2020). Useful road maps: studying Drosophila larva's central nervous system with the help of connectomics. Curr. Opin. Neurobiol. *65*, 129–137. 10.1016/j.conb.2020.09.008.

34. Ohyama, T., Schneider-Mizell, C.M., Fetter, R.D., Aleman, J.V., Franconville, R., Rivera-Alba, M., Mensh, B.D., Branson, K.M., Simpson, J.H., Truman, J.W., et al. (2015). A multilevel multimodal circuit enhances action selection in Drosophila. Nature *520*, 633–639. 10.1038/nature14297.

35. Jovanic, T., Schneider-Mizell, C.M., Shao, M., Masson, J.-B., Denisov, G., Fetter, R.D., Mensh, B.D., Truman, J.W., Cardona, A., and Zlatic, M. (2016). Competitive Disinhibition Mediates Behavioral Choice and Sequences in Drosophila. Cell *167*, 858-870.e19. 10.1016/j.cell.2016.09.009.

36. Schneider-Mizell, C.M., Gerhard, S., Longair, M., Kazimiers, T., Li, F., Zwart, M.F., Champion, A., Midgley, F.M., Fetter, R.D., Saalfeld, S., et al. (2016). Quantitative neuroanatomy for connectomics in Drosophila. eLife *5*, 1133. 10.7554/eLife.12059.

37. Hu, C., Petersen, M., Hoyer, N., Spitzweck, B., Tenedini, F., Wang, D., Gruschka, A., Burchardt, L.S., Szpotowicz, E., Schweizer, M., et al. (2017). Sensory integration and neuromodulatory feedback facilitate Drosophila mechanonociceptive behavior. Nat. Publ. Group *20*, 1085–1095. 10.1038/nn.4580.

38. Nässel, D.R., and Zandawala, M. (2019). Recent advances in neuropeptide signaling in Drosophila, from genes to physiology and behavior. Prog. Neurobiol. *179*, 101607. 10.1016/j.pneurobio.2019.02.003.

39. Lehman, M., Barre, C., Hasan, M.A., Flament, B., Autran, S., Dhiman, N., Soba, P., Masson, J.-B., and Jovanic, T. (2023). Neural circuits underlying context-dependent

competition between defensive actions in Drosophila larva. Preprint at bioRxiv, 10.1101/2023.12.24.573276 10.1101/2023.12.24.573276.

40. Gershow, M., Berck, M., Mathew, D., Luo, L., Kane, E.A., Carlson, J.R., and Samuel, A.D.T. (2012). Controlling airborne cues to study small animal navigation. Nat. Methods *9*, 290–296. 10.1038/nmeth.1853.

41. Jovanic, T., Winding, M., Cardona, A., Truman, J.W., Gershow, M., and Zlatic, M. (2019). Neural Substrates of Drosophila Larval Anemotaxis. Curr. Biol. CB *29*, 554-566.e4. 10.1016/j.cub.2019.01.009.

42. Gomez-Marin, A., and Louis, M. (2014). Multilevel control of run orientation in Drosophila larval chemotaxis. Front. Behav. Neurosci. *8*, 38. 10.3389/fnbeh.2014.00038.

43. Klein, M., Afonso, B., Vonner, A.J., Hernandez-Nunez, L., Berck, M., Tabone, C.J., Kane, E.A., Pieribone, V.A., Nitabach, M.N., Cardona, A., et al. (2015). Sensory determinants of behavioral dynamics in Drosophila thermotaxis. Proc. Natl. Acad. Sci. U. S. A. *112*, E220-9. 10.1073/pnas.1416212112.

44. Masson, J.-B., Laurent, F., Cardona, A., Barre, C., Skatchkovsky, N., Zlatic, M., and Jovanic, T. (2020). Identifying neural substrates of competitive interactions and sequence transitions during mechanosensory responses in Drosophila. Plos Genet. *16*, e1008589. 10.1371/journal.pgen.1008589.

45. Ohyama, T., Jovanic, T., Denisov, G., Dang, T.C., Hoffmann, D., Kerr, R.A., and Zlatic, M. (2013). High-Throughput Analysis of Stimulus-Evoked Behaviors in Drosophila Larva Reveals Multiple Modality-Specific Escape Strategies. PloS One *8*, e71706. 10.1371/journal.pone.0071706.

46. Brown, M.R., Crim, J.W., Arata, R.C., Cai, H.N., Chun, C., and Shen, P. (1999). Identification of a Drosophila brain-gut peptide related to the neuropeptide Y family. Peptides *20*, 1035–1042. 10.1016/s0196-9781(99)00097-2.

47. Wu, Q., Wen, T., Lee, G., Park, J.H., Cai, H.N., and Shen, P. (2003). Developmental Control of Foraging and Social Behavior by the Drosophila Neuropeptide Y-like System. Neuron *39*, 147–161. 10.1016/S0896-6273(03)00396-9.

48. Lee, P.-T., Zirin, J., Kanca, O., Lin, W.-W., Schulze, K.L., Li-Kroeger, D., Tao, R., Devereaux, C., Hu, Y., Chung, V., et al. A gene-specific T2A-GAL4 library for Drosophila. eLife *7*, e35574. 10.7554/eLife.35574.

49. Carlsson, M.A., Enell, L.E., and Nässel, D.R. (2013). Distribution of short neuropeptide F and its receptor in neuronal circuits related to feeding in larval Drosophila. Cell Tissue Res. *353*, 511–523. 10.1007/s00441-013-1660-4.

50. Hunter, P. (2013). Brain drain, brain gain or brain sharing? EMBO Rep. *14*, 315–318. 10.1038/embor.2013.33.

51. Moran, N.P., Sánchez-Tójar, A., Schielzeth, H., and Reinhold, K. (2021). Poor nutritional condition promotes high-risk behaviours: a systematic review and meta-analysis. Biol. Rev. *96*, 269–288. https://doi.org/10.1111/brv.12655.

52. Sandhu, P., Shura, O., Murray, R.L., and Guy, C. (2018). Worms make risky choices too: the effect of starvation on foraging in the common earthworm (Lumbricus terrestris). Can. J. Zool. 10.1139/cjz-2018-0006.

53. Levy, D.J., Thavikulwat, A.C., and Glimcher, P.W. (2013). State dependent valuation: the effect of deprivation on risk preferences. PloS One *8*, e53978. 10.1371/journal.pone.0053978.

54. Inagaki, H.K., Ben-Tabou de-Leon, S., Wong, A.M., Jagadish, S., Ishimoto, H., Barnea, G., Kitamoto, T., Axel, R., and Anderson, D.J. (2012). Visualizing neuromodulation in vivo: TANGO-mapping of dopamine signaling reveals appetite control of sugar sensing. Cell *148*, 583–595. 10.1016/j.cell.2011.12.022.

55. Root, C.M., Ko, K.I., Jafari, A., and Wang, J.W. (2011). Presynaptic facilitation by neuropeptide signaling mediates odor-driven food search. Cell *145*, 133–144. 10.1016/j.cell.2011.02.008.

56. Petzold, A., van den Munkhof, H.E., Figge-Schlensok, R., and Korotkova, T. (2023). Complementary lateral hypothalamic populations resist hunger pressure to balance nutritional and social needs. Cell Metab. *35*, 456-471.e6. 10.1016/j.cmet.2023.02.008.

57. Hsu, W.-M.M., Kastner, D.B., Baccus, S.A., and Sharpee, T.O. (2021). How inhibitory neurons increase information transmission under threshold modulation. Cell Rep. *35*, 109158. 10.1016/j.celrep.2021.109158.

58. Morozova, E., Newstein, P., and Marder, E. (2022). Reciprocally inhibitory circuits operating with distinct mechanisms are differently robust to perturbation and modulation. eLife. 10.7554/eLife.74363.

59. Pakan, J.M., Lowe, S.C., Dylda, E., Keemink, S.W., Currie, S.P., Coutts, C.A., and Rochefort, N.L. (2016). Behavioral-state modulation of inhibition is context-dependent and cell type specific in mouse visual cortex. eLife *5*, e14985. 10.7554/eLife.14985.

60. Chen, Q., Pei, Z., Koren, D., and Wei, W. (2016). Stimulus-dependent recruitment of lateral inhibition underlies retinal direction selectivity. eLife *5*, 4481. 10.7554/eLife.21053.

61. Fadok, J.P., Krabbe, S., Markovic, M., Courtin, J., Xu, C., Massi, L., Botta, P., Bylund, K., Müller, C., Kovacevic, A., et al. (2017). A competitive inhibitory circuit for selection of active and passive fear responses. Nature *542*, 96–100. 10.1038/nature21047.

62. Koyama, M., Minale, F., Shum, J., Nishimura, N., Schaffer, C.B., and Fetcho, J.R. (2016). A circuit motif in the zebrafish hindbrain for a two alternative behavioral choice to turn left or right. eLife *5*, 451. 10.7554/eLife.16808.

63. Koyama, M., and Pujala, A. (2018). Mutual inhibition of lateral inhibition: a network motif for an elementary computation in the brain. Curr. Opin. Neurobiol. *49*, 69–74. 10.1016/j.conb.2017.12.019.

64. Mysore, S.P., and Knudsen, E.I. (2012). Reciprocal Inhibition of Inhibition: A Circuit Motif for Flexible Categorization in Stimulus Selection. Neuron *73*, 193–205. 10.1016/j.neuron.2011.10.037.

65. Mysore, S.P., and Kothari, N.B. (2020). Mechanisms of competitive selection: A canonical neural circuit framework. eLife *9*, e51473. 10.7554/eLife.51473.

66. Swierczek, N.A., Giles, A.C., Rankin, C.H., and Kerr, R.A. (2011). High-throughput behavioral analysis in C. elegans. Nat. Methods *8*, 592–598. 10.1038/nmeth.1625.

67. Sakagiannis, P., Jürgensen, A.-M., and Nawrot, M.P. (2021). A realistic locomotory model of Drosophila larva for behavioral simulations. Preprint at bioRxiv, 10.1101/2021.07.07.451470 10.1101/2021.07.07.451470.

68. Matsuda, H., Yamada, T., Yoshida, M., and Nishimura, T. (2015). Flies without Trehalose *. J. Biol. Chem. *290*, 1244–1255. 10.1074/jbc.M114.619411.

69. Matsushita, R., and Nishimura, T. (2020). Trehalose metabolism confers developmental robustness and stability in Drosophila by regulating glucose homeostasis. Commun. Biol. *3*, 1–12. 10.1038/s42003-020-0889-1.

A Feeding behavior

Legend:
- Fed (black)
- Sucrose fed (red)
- Starved (purple)
- +/UAS-TNT (green)
- Gr43a-Gal4/UAS-TNT (orange)

A1 Normal food
A2 Yeast
A3 Water
A4 Sucralose
A5 Sucrose 20%

B Circulating carbohydrates in hemolymph
B1 Hemolymph glucose
B2 Hemolymph trehalose

C Preference for sucrose
- Fed (black)
- Protein-deprived (red)
- Starved (purple)

D Baseline locomotion, 1h30 of treatment

Maximum dispersion over 60s (mm); Scaled speed (mm.s⁻¹); Mean tortuosity over 20 s; Time fraction in runs

E Stimulus induced behavioral responses

Legend:
- Fed (black)
- Sucrose fed (red)
- Starved (purple)

E1
Hunch (%); Static-Bend (%); Stimulus induced Head Cast (%)

Crawl — Hunch 0.69***, 4.05***; Crawl — Head-Cast 1.29*, 0.86; Hunch — Head-Cast 0.72**, 1.16***

E2
Hunch (%); Static-Bend (%); Stimulus induced Head Cast (%)

Crawl — Hunch 0.71***, 4.00***; Crawl — Head-Cast 1.55***, 1.89***; Hunch — Head-Cast 0.62***, 0.99

Figure 1. Physiological state of food-deprived larvae. A: Larval feeding on different substrates was quantified, in control animals fed ad-libitum, in animals fed on sucrose or in animals subjected to complete starvation during 90 min. A1: Starved animals increased their feeding on a standard food medium as compared to fed animals, while animals fed on sucrose only did not (ANOVA, n = 35-41 larvae, Tukey post-hoc test *** : p < 0.001). A2 : Both animals fed on sucrose only and starved animals increase their yeast feeding as compared to normally fed larvae (ANOVA, n = 51-59 larvae, Tukey post-hoc test *** : p < 0.001, ** : p < 0.01). A3 : While starved animals consume a similar amount of water as fed ones, animals fed on sucrose only double their water consumption as compared to fed and starved larvae (ANOVA, n = 52-54 larvae, Tukey post-hoc test *** : p < 0.001, ** : p < 0.01). A4 Consumption of the non-energetic sweetener sucralose was similar among conditions (n = 25-32 larvae). A5 : Larvae that had fed on a 20% sucrose solution for 90 min decrease their sucrose intake as compared to fed and starved larvae (ANOVA, n = 29-33 larvae, Tukey post-hoc test *** : p < 0.001, * : p < 0.05). Gustatory neurons inhibition (Gr43a-Gal4>TNT) decreases sucrose intake in fed larvae to the level of larvae fed on sucrose, while the consumption of starved larvae remains higher (ANOVA, n = 24-33 larvae, Tukey post-hoc test ** : p < 0.01, * : p < 0.05). B: Concentration of glucose and trehalose in larval hemolymph. B1: Glucose concentration is increased in the hemolymph of larvae that were fed on 20% sucrose solution for 90 min compared to larvae fed on a standard diet and to starved animals (ANOVA, N = 3 samples from 5 larvae each, Tukey post-hoc test ** : p < 0.01, * : p < 0.05). B2: Trehalose concentration is similar in the hemolymph of larvae fed on the different diets (N = 6 samples from 10 larvae each). C: Place preference assay for sucrose. Larvae that fed on a 20% sucrose solution for 90 min exhibit a decreased glucose preference compared to normally fed and starved larvae (n = 79-133 larvae, Chi-2 test ** : p < 0.01, * : p < 0.05). D: Analysis of larval locomotion in the absence of sensory stimulus. After 90 min of sucrose feeding or complete starvation, larval dispersion, time spent crawling and speed are significantly increased, while the tortuosity of the trajectory is decreased compared to larvae fed on a standard diet (Mann-Whitney test, **** : p < 0.0001, *** : p < 0.001, ** : p < 0.01, * : p < 0.05). E: Manipulating the feeding state modulates behavioral responses to mechanosensory stimuli. Behavior in response to air puff during the first five seconds uoin stimulus onset, for, E1 in larvae fed on sucrose only (n = 592-629 larvae) and E2 starved larvae compared to larvae fed on standard food (n =704-771 larvae, *** : p < 0.001, ** : p < 0.01, * : p < 0.05). See also Figure S1 and Supplementary material 2,3 and 4.

A   Structure of the mechano-sensory circuit

- mechanosensory neurons
- feedback inhibitory neurons
- feedforward inhibitory neurons
- projection neurons

B   Calcium imaging in chordotonal mechanosensory neurons

C   Behavioral response to optogenetic activation of chordotonal mechanosensory neurons

Figure 2: Bias in behavioral strategy does not come from the modulation of sensory neurons. A: Organization of the circuit. B: Calcium responses to mechanical stimulations in chordotonal neurons larvae fed on sucrose only and on standard food (R61D08-Gal4/UAS-GCaMP6s). Left panel : calcium responses of chordotonal neuron projections in the VNC from different individuals fed on different diets. Lower right panel : mean calcium response of chordotonal neurons over time. The green dashed line corresponds to stimulus duration. Upper right panel: calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. Stimulus-induced activity of chordotonal mechanosensory neurons is unchanged in animals fed on sucrose as compared to larvae fed on standard food (n = 10 larvae, 3 trials per larva, t-test p = 0.249). C: Optogenetic activation of cho in larvae fed only on sucrose. Hunch probability is computed during the first 2 seconds from stimulus onset. Bend is the mean probability during the first 10 seconds from stimulus onset, corrected by the baseline recording prior to the stimulus. (n = 192 larvae for fed, 168 for sucrose fed, p = 0.011 for Hunch, 0.028 for Bend). See also Figure S2

A Calcium imaging in Basin-1

Fed · Sucrose fed

B Calcium imaging in Basin-2

Fed · Sucrose fed

C Model 1 (LnA RMAX)

C1 · C2 · C3

D Model 2 (Hb input current)

D1 · D2 · D3

E Model Combined

E1 · E2 · E3

Figure 3. The feeding status affects the output of the sensorimotor circuit. A: Basin-1 calcium responses to mechanical stimulations in larvae fed on standard food and on sucrose only larvae (20B01-lexA; LexAop-GCaMP6s, UAS-CsChrimson-mCherry). Left panel: calcium responses of Basin-1 from different individuals fed on different diets. Lower right panel: mean calcium response of Basin-1 over time. The green dashed line corresponds to stimulus onset. Upper right panel: mean calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. Stimulus-induced activity of Basin-1 neurons is similar in larvae fed on sucrose only compared to larvae fed on a standard food (n = 17/20 larvae, 1 trial per larva, t-test p = 0.2577). B: Basin-2 calcium responses to mechanical stimulations in different states (SS00739/UAS-GCaMP6s). Left panel : calcium responses of Basin-2 from different individuals fed on different diets. Lower right panel : mean calcium response of Basin-2 over time. The green dashed line corresponds to stimulus onset. Upper right panel : mean calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. Low panel : percentage of trials that failed to elicit a calcium response in Basin-2. Stimulus-induced activity of Basin-2 neurons is enhanced in animals fed on sucrose only compared to larvae fed on a standard food (n = 12 larvae, 5 trials per larva, mean response comparisons t-test p = 0.0118, percentage of failed responses comparison Chi-2 test p = 0.0027).

C-E: Simple connectome based rate model of the decision circuit for the selection Hunch and Bend/Hedcast C: Model 1: version of the model where the sucrose state is modeled by a decreased maximum rate of the LNa node. C1. schematic of the circuit where the influence of LNa on downstream targets is weaker C2. state space trajectories of Basin-1 (B1) and Basin-2 Activity as a function of Lna and LNa activities for model 1. C3. phase ratio phase diagram for model 1 D: Model 2: version of the model where the sucrose state is modeled by an input current to Hb) Handle B neuron D1. schematic of the circuit where the influence of Hb on downstream targets is stronger D2. state space trajectories of Basin-1 (B1) and Basin-2 Activity as a function of Lna and LNa activities for model 2. D3. phase ratio phase diagram for model 2

E: Combined model : version of the model where the sucrose state is modeled by both an input current to Handle B (Hb) neuron and decreased Maximum rate of LNa E1. schematic of the circuit where the influence of Hb on downstream targets is stronger and the one of Lna weaker E2. state space trajectories of Basin-1 (B1) and Basin-2 Activity as a function of Lna and LNa activities for the combined model. E3. phase ratio phase diagram for the combined model. See also Figures S4 and S5

A Calcium imaging in Griddle-2

Fed

Sucrose fed

Fed
Sucrose fed

Mean ΔF/F$_0$ (%)

Individual

Time (s)

ΔF/F$_0$ (%)

Stimulus

B Calcium imaging in Handle-B

Fed

Sucrose fed

Fed
Sucrose fed

Mean ΔF/F$_0$ (%)

Individual

Time (s)

ΔF/F$_0$ (%)

Stimulus

C Reciprocal inhibition of G2 and Hb

C1 iLNa activity upon silencing Hb

In Fed condition

silenced
not silenced

iLNa activity (AU)

time (AU)

w_iLNa

In sucrose condition

iLNa activity (AU)

time (AU)

w_iLNa

C2 Hb inactivation and Ca imaging in G2

Fed, no inactivation
Fed, Handle-B inactivation

Stimulus

Sucrose fed no inactivation
Sucrose fed, Handle-B inactivation

Stimulus

Fed
Sucrose fed
No inactivation
Inactivation

C3

Mean ΔF/F$_0$ (%)

C4 Hb activity upon iLNa silencing

In Fed condition

silenced
not silenced

fbLNHb activity (AU)

time (AU)

w_iLNa

In sucrose condition

fbLNHb activity (AU)

time (AU)

w_iLNa

C5 G2 inactivation and calcium imaging in Hb

Fed, no inactivation
Fed, Griddle-2 inactivation

Stimulus

Sucrose fed, no inactivation
Sucrose fed, Griddle-2 inactivation

Stimulus

Fed
Sucrose fed
No inactivation
Inactivation

C6

Mean ΔF/F$_0$ (%)

Figure 4. Two interneuron subtypes are oppositely modulated by the internal state. A : Griddle-2 calcium responses to mechanical stimulations in larvae fed on standard food and sucrose only (SS00918/UAS-GCaMP6s). Left panel: calcium responses of Griddle-2 from different individuals, fed on standard food or on sucrose only. Lower right panel: mean calcium response of Griddle-2 over time. The green dashed line corresponds to stimulus onset. Upper right panel: mean calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. Stimulus-induced activity of Griddle-2 is decreased in animals fed on sucrose only compared to larvae fed on a standard food (n = 10 larvae, 1 trial per larva, t-test p = 0.0068). B: Handle-b calcium responses to mechanical stimulations in larvae fed on standard food or on sucrose only (SS00888/UAS-GCaMP6s). Left panel: calcium responses of Handle-b from different individuals, fed on standard food or on sucrose only. Lower right panel: mean calcium response of Handle-b over time. The green dashed line corresponds to stimulus onset. Upper right panel: mean calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. Stimulus-induced activity of Handle-b is increased in animals fed on sucrose only compared to larvae fed on a standard food (n = 23/25 larvae, 1 trial per larva, t-test p = < 0.0001). C: Reciprocal inhibition between Handle-b and Griddle-2. C1: model simulation, trajectories of LNa activity upon Hb inactivation in fed (top panel) and sucrose condition (bottom panel) C2: Left panel: mean calcium response of Griddle-2 over time, with (SS0888-Gal4>UAS-TNT 55C05-LexA>LexAop-GCaMP6s) or without Handle-b inactivation (+/UAS-TNT 55C05-LexA>LexAop-GCaMP6s), in larvae fed on standard food or on sucrose only. The green dashed line corresponds to stimulus onset. C3:: mean calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. Griddle-2 responses to mechanical stimuli are increased after Handle-b inactivation, both in larvae fed on standard food (top panel) or on sucrose (bottom panel) (n = 7-9 larvae, 4 trials per larva, t-test after inactivation in larvae fed on standard food p = 0.009, fed on sucrose only p = 0.002). C1. model simulation, trajectories of Hb activity upon iLna inactivation in fed (top panel) and sucrose condition (bottom panel) C5. mean calcium response of Handle-b over time, with (55C05-LexA>LexAop-TNT 22E09-Gal4>UAS-GCaMP6s) or without Handle-b inactivation (+/LexAop-TNT 22E09-Gal4>UAS-GCaMP6s), in larvae larvae fed on standard food or on sucrose. The green dashed line corresponds to stimulus onset. C6: mean calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. Handle-b responses to mechanical stimuli are increased after Griddle-2 inactivation only in fed animals (n = 6-10 larvae, 5 trials per larva, t-test after inactivation in fed p = 0.0027, sucrose condition p = 0.065). See also Figure S4

A Handle-B inactivation and calcium imaging in Basin-2

Fed, no inactivation
Sucrose fed, no inactivation
Sucrose fed, Handle-B inactivation

calcium activity

genetic inactivation

Stimulus
1 s
40 % ΔF/F₀

Fed
Sucrose fed
No inactivation
Inactivation

Mean ΔF/F₀ (%)

Failed responses (%)

B Handle-B activation and calcium imaging in Basin-2

calcium activity

Fed, no activation
Fed, Handle-B activation
Protein-deprived, no activation

optogenetic activation

40 % ΔF/F₀
1 s
Stimulus

Fed
Sucrose fed
No activation
Activation

Mean ΔF/F₀ (%)
NS

C Griddle-2 inactivation and calcium imaging in Basin-2

genetic inactivation

calcium activity

Fed, no inactivation
Fed, Gridle-2 inactivation
Sucrose fed, no inactivation
Sucrose fed, Griddle-2 inactivation

100 % ΔF/F₀
1 s
Stimulus

Fed
Sucrose fed
No inactivation
Inactivation

Mean ΔF/F₀ (%)

Failed responses (%)

Figure 5. The modulation of the circuit output is a consequence of the modulation of inhibitory interneurons. A: Left panel: mean calcium responses of Basin-2 over time from individuals fed on different diets, with (38H09-LexA>LexAop-GCaMP6s 22E09-Gal4>UAS-TNT) or without (38H09-LexA>LexAop-GCaMP6s +/>UAS-TNT) Handle-b inhibition. The green dashed line corresponds to stimulus onset. Middle panel: calcium response of Basin-2 averaged during the stimulus. White line represents the mean, white dot represents the median. Right panel: percentage of trials that failed to elicit a calcium response in Basin-2. Inactivating Handle-b in larvae fed on sucrose only prevents the effect of sucrose feeding on Basin-2 activity (n = 12-14 larvae, 5 trials per larva ; ANOVA with post-hoc tests: *** : $p < 0.001$; chi-2 test *** : $p < 0.001$, ** : $p < 0.01$). B: Left panel: mean calcium responses of Basin-2 over time from individuals fed on different diets, with or without optogenetic activation of Handle-b (22E09-Gal4>UAS-CsChrimson::tdTomato 38H09-LexA>LexAop-GCaMP6s) during the first second of mechanical stimulus (red box). The green dashed line corresponds to stimulus onset. Right panel : calcium response of Basin-2 averaged during the first second of the stimulus. White line represents the mean, white dot represents the median. Activating Handle-b in fed larvae phenocopies the effect of sucrose feeding on Basin-2 activity (n = 8 larvae, 5 trials per larva; ANOVA with post-hoc tests: *** : $p < 0.001$, * : $p < 0.05$). C: Left panel: mean calcium responses of Basin-2 over time from individuals fed on different diets, with (38H09-LexA>LexAop-GCaMP6s 55C05-Gal4>UAS-TNT) or without (38H09-LexA>LexAop-GCaMP6s +/>UAS-TNT) Griddle-2 inhibition. The green dashed line corresponds to stimulus onset. Middle panel: calcium response of Basin-2 averaged during the stimulus. White line represents the mean, white dot represents the median (ANOVA with post-hoc tests: *** : $p < 0.001$). Right panel: percentage of trials that failed to elicit a calcium response in Basin-2 (chi-2 test *** : $p < 0.001$, * : $p < 0.05$). See also Figure S6

A. NPF neuron connects to Handle-B

B. NPF activity in different feeding states

- Fed
- Sucrose fed
- Starved

Hb
NPF Neuron

90 min | 90 min | 5 hours

C. Locomotion of larvae upon NPF knockdown

C1. control

maximum dispersal over 60s (mm) | scaled speed (1/s) | tortuosity over 20s | time fraction in runs

C2 NPF knockdown in DM-NPF neuron

maximum dispersal over 60s (mm) | scaled speed (1/s) | tortuosity over 20s | time fraction in runs

D. Ca-imaging in Hb upon silencing of NPF neurons

- Control
- KIR

Control | KIR

Recordings
Time (s) | Time (s)

ΔF/F₀ (%)

Mean ΔF/F₀ (%)

50% ΔF/F₀
1 s
Stimulus

E. Immunolabeling SS00888-NPFR T2A-LexA

15 μm | 5 μm SS00888 | 5 μm NPF-R | 5 μm MERGE

Figure 6 A: Expression profile characterization of GMR_SS01635>UAS-GFP stained with a GFP-specific antibody. GMR_SS01635 split-Gal4 line selectively targets one pair of NPF neurons. Comparison of light microscopy images with electron microscopy (EM) reconstruction images indicates that a descending neuron synapses on Handle-b (Jovanic *et al*, 2016) is the DM-NPF descending neuron. EM image shows neuropeptide-containing dense core vesicles in the NPF descending neuron near one of its synapses with Handle-b. B: Baseline calcium fluorescence measured in the projections of NPF descending neurons in the VNC (ventral nerve cord) of larvae fed on sucrose only for 90 minutes (N = 10 larvae), starved for 90 minutes (N = 10 larvae) or 5 hours (N = 8-9 larvae). White line represents the mean, white dot represents the median. Starvation or feeding only on sucrose only increase NPF descending neuron activity (two-tailed T-test: * : $p < 0.05$). C : Analysis of larval locomotion in the absence of sensory stimulus. Top row shows control larvae. Bottom row shows larvae with NPF knockdown in DM- NPF descending neurons. NPF knockdown in DM-NPF descending neuron abolishes differences in locomotion mean between larvae in different states; it abolishes the difference in trajectory tortuosity, time allocation in exploration (time fraction in runs), average movement speed and dispersal between larvae fed on standard food and starved larvae. In larvae fed on sucrose the increase in exploration compared to larvae fed on standard food was also abolished (Mann-Whitney test, ****: $p<0.0001$, *** : $p < 0.001$, ** : $p < 0.01$, * : $p < 0.05$) D: Calcium responses in Handle-b with (UAS-GCamP6s; 22E09-Gal4, NPF-LexA>LexAop-KIR) or without (UAS-GCamP6s; 22E09-Gal4, NPF-LexA;+) NPF neurons silencing. Left panel: calcium responses of Handle-b, in each trial of mechanosensory stimulation. Right lower panel: mean calcium response of Handle-b over time. The green dashed line corresponds to the stimulus Top right panell: mean calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. Stimulus-induced activity of Handle-b is decreased upon Coaster silencing (n = 10 larvae, 3 trials per larva, t-test $p = 0.045$). E: Immunohistochemical labeling for NPFR in Handle_B UAS-GCamP6s is expressed in Handle-b using the GMR_SS00888 split-Gal4 line (green) and LexAop-jRGeco1a is expressed under the control of the NPFR promoter using a T2A-LexA construct (magenta). Antibodies against GFP and dsRed were used to increase detection sensitivity. Co-localization of LexAop-jRGeco1a and GCamP6s show that Handle-b expresses the NPFR. See also Figure S7

A sNPFR expression in Handle-b

sNPF-R   60E02-LexA   MERGE

B Ca-imaging in Hb upon sNPFR knockdown

Control   sNPFR-RNAi

Control   sNPFR-RNAi

Mean ΔF/F₀ (%)

Stimulus

C

Hunch (%)   Static-Bend (%)

Stimulus induced Head Cast (%)

Crawl   0.90*   Hunch
         0.82

0.98   0.58***
5.69***   1.31***
         Head-Cast

D sNPFR expression in Griddle-2

sNPF-R   55C05-LexA   MERGE

E Ca-imaging in Hb upon sNPFR knockdown

Control   sNPFR-RNAi

Control   sNPFR-RNAi

Mean ΔF/F₀ (%)

Stimulus

F

Hunch (%)   Static-Bend (%)

Stimulus induced Head Cast (%)

Crawl   0.76***   Hunch
         1.18

1.48*   0.96
1.90**   1.14*
         Head-Cast

Figure 7. sNPFR is required for the modulation. A: Immunohistochemical labeling for sNPFR in Handle-b. LexAop-jRGeco1a is expressed in Handle-b using the 60E02-LexA line (magenta) and UAS-GCamP6s is expressed under the control of the sNPFR promoter using a T2A-Gal4 construct (green). Antibodies against GFP and dsRed were used to increase detection sensitivity. Co-localization of antibodies against jRGeco1a and GCamP6s shows that Handle-b expresses the sNPFR. B: Calcium responses in Handle-b upon sNPFR knockdown (GMR_SS00888>UAS-GCamP6s; UAS-sNPFR-RNAi) compared to a control (GMR_SS00888>UAS-GCamP6s) in Handle-b. Left panel: calcium responses of Handle-b from different individuals and trials. One line is a trial. Lower right panel: mean calcium response of Handle-b over time. The green dashed line corresponds to the stimulus Top right panel: mean calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. Stimulus-induced activity of Handle-b is increased by sNPFR knockdown (n = 15 larvae, 4 trials per larva, t-test p = 0.035). C: Behavior in response to air puff during the first five seconds upon stimulus onset,for larvae in which sNPFR was knocked down in Handle-b neurons (SS00888>sNPFR-RNAi) compared to the control (n = 185-260 larvae) (*** : $p < 0.001$, ** : $p < 0.01$, * : $p < 0.05$). D: Griddle-2 immunostaining for sNPFR expression. LexAop-jRGeco1a is expressed in Griddle-2 using the 55C05-LexA line (magenta) and UAS-GCamP6s is expressed s expressed under the control of the sNPFR promoter using a T2A-Gal4 construct (green). Antibodies against GFP and dsRed were used to increase detection sensitivity. Co-localization of antibodies against jRGeco1a and GCamP6s show that Griddle-2 expresses sNPFR. E: Calcium responses in Griddle-2 upon sNPFR knockdown (SS_TJ001>UAS-GCamP6s; UAS-sNPFR-RNAi) compared to the control (SS_TJ001>UAS-GCamP6s). Left panel: calcium responses of Griddle-2 from different individuals and trials. One line is a trial. Lower right panel: mean calcium response of Griddle-2 over time. The green dashed line corresponds to the stimulus Top right panel. mean calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. Stimulus-induced activity of Griddle-2 is decreased by sNPFR knockdown (n = 10 larvae, 3 trials per larva, t-test p = 0.003). F: Behavior in response to air puff during the first five seconds upon stimulus onset, for larvae in which sNPFR was knocked down in Griddle-2 neurons (SS_TJ001>sNPFR-RNAi) compared to the control (n = 243-438 larvae, *** : $p < 0.001$, ** : $p < 0.01$, * : $p < 0.05$). See also Figure S7, S8 and Supplementary material 3 and 4

Supplementary information

Feeding- state dependent modulation of reciprocally interconnected inhibitory neurons biases sensorimotor decisions in *Drosophila*

Eloïse de Tredern[1,2], Dylan Manceau[1,2], Alexandre Blanc[3,4], Panagiotis Sakagiannis[5], Chloe Barre[3,4], Victoria Sus[2], Francesca Viscido[2], Md Amit Hasan[2], Sandra Autran[2], Martin Nawrot[5], Jean-Baptiste Masson[3,4], Tihana Jovanic[2]*

[1]These authors contributed equally

[2]Université Paris-Saclay, CNRS, Institut des neurosciences Paris-Saclay, 91400, Saclay, France

[3]Institut Pasteur, Université Paris Cité, CNRS UMR 3571, Decision and Bayesian Computation, 75015 Paris, France

[4]Epiméthée, INRIA

[5]Computational Systems Neuroscience, Institute of Zoology, University of Cologne, 50674 Cologne, Germany

*corresponding author

A Baseline locomotion, 5h of treatment

B Baseline locomotion, 18h of treatment

C Baseline locomotion, 5h of treatment +refeeding 15 minutes

D Baseline locomotion, 5h of treatment +refeeding 15 minutes

E Stimulus induced behavioral responses, 5h of treatment

E1 feeding on sucrose only

E2 starvation

Figure S1. A-D: Analysis of larval locomotion in different feeding states. A-B: Larvae starved for 5h (A, n = 283, 211, 309 larvae) or 18h (B, n = 173, 141, 139 larvae) disperse further, at a higher movement speed, allocating more time to crawling (time fraction in runs) with lower-tortuosity trajectories compared to larvae fed on a standard diet. C: Refeeding larvae on standard food for 15 min after a period of 5h starvation rescues the normal locomotion as dispersion distance, time allocation to crawling and crawling speed are no longer significantly increased, nor the trajectories' tortuosity decreased compared to larvae fed on a standard diet (n= 244, 160, 223 larvae) D: Rehydration by putting larvae for 15 minutes on water after 90 minutes of feeding of sucrose restores normal locomotion similar to larvae fed on standard food, since dispersion distance, time allocation to crawling and movement speed are no longer significantly increased, nor the trajectories' tortuosity decreased. (Mann-Whitney test, ****: $p<0.0001$, *** : $p < 0.001$, ** : $p < 0.01$, * : $p < 0.05$) (n = 102, 109,123 larvae) E-F: Behavior in response to air puff during the first five seconds upon stimulus onset, Behavioral and transition probabilities upon 5h of sucrose feeding (E) and starvation (F) (n=233-301 larvae, *** : $p < 0.001$, ** : $p < 0.01$, * : $p < 0.05$).

A Calcium imaging in chordotonal neurons, starved

B Calcium imaging in chordotonal neurons

C GFP expression in chordotonal neurons

Fed
Sucrose fed
Starved

D Behavioral response to optogenetic activation of chordotonal mechanosensory neurons

Fed
Starved

0-10 s
**

Figure S2: A: Calcium responses to mechanical stimulations at 5V in chordotonal neurons in fed and starved larvae (R61D08-Gal4/UAS-GCaMP6s). Left panel: calcium responses of chordotonal neurons from different individuals fed on different diets. Lower right panel: mean calcium response of chordotonal neurons over time. The green dashed line corresponds to stimulus duration. Upper right panel: calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. Stimulus-induced activity of chordotonal mechanosensory neurons is not significantly increased in starved animals as compared to larvae fed on a standard food (n = 10 larvae, 1 trial per larva, t-test p = 0.092). B: Calcium responses of chordotonal neurons to different intensities of mechanical stimulations, in larvae fed on standard food medium (n = 5 larvae). C: GFP expression levels in chordotonal neurons R61D08>GFP larvae fed on different food media: standard food, 20% sucrose and water (n = 4 larvae per condition). D: Optogenetic activation of cho in starved larvae. Hunch is cumulative probability during the first 2 seconds from stimulus onset. Bend is the mean probability during the first 10 seconds from stimulus onset, corrected by 40 seconds of recording prior to the stimulus. Hunching and stimulus-induced Bending (n = 192 larvae for fed, 157 for starved, p = 0.080 for hunch, 0.003 for bend).

A Calcium imaging in Basin-1

Fed | Starved

Time (s) | Time (s)

ΔF/F$_0$ (%)

Mean ΔF/F$_0$ (%)

Fed
Starved

50 % ΔF/F$_0$

1 s | Stimulus

B Calcium imaging in Basin-2

Fed | Starved

Time (s) | Time (s)

ΔF/F$_0$ (%)

Mean ΔF/F$_0$ (%)

Fed
Starved

50 % ΔF/F$_0$

1 s | Stimulus

C Basin-1 activity for different intensities of stimulation

Fed
Sucrose fed

Mean ΔF/F$_0$ (%)
Intensity of stimulation (V)

Fed
Starved

Mean ΔF/F$_0$ (%)
Intensity of stimulation (V)

D Basin-2 activity for different intensities of stimulation

Fed
Sucrose fed

Mean ΔF/F$_0$ (%)
Intensity of stimulation (V)

Fed
Starved

Mean ΔF/F$_0$ (%)
Intensity of stimulation (V)

E Basin-2 probability of response

Fed
Sucrose fed

Failed responses (%)
Intensity of stimulation

Fed
Starved

Failed responses (%)
Intensity of stimulation

Figure S3: A: Basin-1 calcium responses to mechanical stimulations in fed and starved larvae (20B01-lexA; LexAop-GCaMP6s, UAS-CsChrimson-mCherry). Left panel: calcium responses of Basin-1 from different individuals fed on different diets. Lower right panel: mean calcium response of Basin-1 over time. The green dashed line corresponds to stimulus onset. Upper right panel: mean calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. Stimulus-induced activity of Basin-1 neurons is similar in starved animals compared to larvae fed on a standard food (n = 11/9 larvae, 1 trial per larva, t-test p = 0.2892). B: Basin-2 calcium responses to mechanical stimulations in different states (SS00739/UAS-GCaMP6s). Left panel: calcium responses of Basin-2 from fed and starved animals. Lower right panel: mean calcium response of Basin-2 over time. The green dashed line corresponds to stimulus onset. Upper right panel: mean calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. Stimulus-induced activity of Basin-2 is not significantly decreased in starved animals compared to larvae fed on a standard food (n = 12/13 larvae, 1 trial per larva, t-test p = 0.1804). C: Basin-1 calcium responses for different intensities of stimulation, in larvae fed on standard food, on sucrose only or completely starved larvae (n = 8-20 larvae, 1 trial per larva). Average neuronal activities over time for different intensities of stimulation are plotted. D: Basin-2 calcium responses for different intensities of stimulation, in larvae fed on standard food, on sucrose only or completely starved larvae (n = 11-13 larvae, 1 trial per larva, t-test ** : p < 0.01). Averaged neuronal activity over time for different intensities of stimulation are plotted. E. percentage of failed responses comparison Chi-2 test p = 0.0027).

A Calcium imaging in Griddle-2

B Calcium imaging in Handle-B

C Griddle-2 activity for different intensities of stimulation

D Handle-B activity for different intensities of stimulation

E Handle-B inactivation and calcium imaging in Griddle-2

F Griddle-2 inactivation and calcium imaging in Handle-B

Figure S4: A: Griddle-2 calcium responses to mechanical stimulations in fed and starved larvae (SS00918/UAS-GCaMP6s). Left panel: calcium responses of Griddle-2 from different individuals fed on different diets. Lower right panel: mean calcium response of Griddle-2 over time. The green dashed line corresponds to the stimulus Upper right panel: mean calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. Stimulus-induced activity of Griddle-2 neurons is not significantly decreased in starved animals compared to larvae fed on a standard food (n = 10/9 larvae, 1 trial per larva, t-test p = 0.0979). B: Handle-b calcium responses to mechanical stimulations in different states (SS00888/UAS-GCaMP6s). Left panel: calcium responses of Handle-b from fed and starved animals. Lower right panel: mean calcium response of Handle-b over time. The green dashed line corresponds to stimulus onset. Upper right panel: mean calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. Stimulus-induced activity of Handle-b is significantly increased in starved animals compared to larvae fed on a standard food (n = 10/13 larvae, 1 trial per larva, t-test p = 0.019). C: Griddle-2 calcium responses for different intensities of stimulation, in larvae fed on standard food, on sucrose only .and starved larvae (n = 9-10 larvae, 1 trial per larva, t-test ** : p < 0.01, * : p < 0.05). Average neuronal activities over time for different intensities of stimulation are plotted D: Handle-b calcium responses for different intensities of stimulation, in larvae fed on standard food or on sucrose only.and starved larvae (n = 9-25 larvae, 1 trial per larva, t-test *** : p < 0.001, ** : p < 0.01, * : p < 0.05). Responses averaged during the stimulus for different intensities of stimulation are plotted E. Calcium responses in Griddle-2 with (55C05-LexA>LexAop-GCaMP6s SS00888-Gal4>UAS-TNT) or without Handle-b inactivation (55C05-LexA>LexAop-GCaMP6s +/UAS-TNT), in each trial of mechanosensory stimulation (n = 7-9 larvae, 4 trials per larva). F: Calcium responses in Handle with (55C05-LexA>LexAop-TNT 22E09-Gal4>UAS-GCaMP6s) or without Handle-b inactivation (+/LexAop-TNT 22E09-Gal4>UAS-GCaMP6s), in each trial of mechanosensory stimulation (n = 6-10 larvae, 5 trials per larva).

Figure S5 A: State space trajectories of B1 and B2 Dynamics as a function of inputs to LNa and LNb neurons in a simple rate model (as in Jovanic *et al.,* 2016). B: Dynamics of a single model neuron with self inhibition shows two equilibria, only one of which is stable C: Level-set for threshold in combined model. C1. Points on lines at the intersection of the surface and horizontal planes represent sets of parameters with identical behavior response. C2. State-space trajectories of the models parameterized by the three points in C1 do not differ qualitatively D: Combined model D1. Points on a vertical line correspond to the same model parameter, with different inputs to iLNa. Points above the surface converge to the monoactive state, while points below the surface converge to the coactive state. D2. Trajectories corresponding to the points in D1, color-coded. E: Silencing Handle-B in model 1, where the sucrose state is modeled as a decrease in LnA max. F/ Silencing Handle-b in model 2 where the sucrose state is modeled as increase input in Handle-b

A Handle-B inactivation and calcium imaging in Basin-2

A1

Fed, no activation | Fed, inactivation | Sucrose, no inactivation | Sucrose, Hb inactivation

A2

B Handle-B activation and calcium imaging in Basin-2

Fed, no activation | Fed, Hb activation | Sucrose, no activation

C Griddle-2 inactivation and calcium imaging in Basin-2   **C1**

Fed, no inactivation | Fed, G2 inactivation

Sucrose fed | Sucrose fed G2 inactivation

Fed, no inactivation
Fed, Gridle-2 inactivation
Sucrose fed, no inactivation
Sucrose fed, Griddle-2 inactivation

C2

Fed
Sucrose fed
No inactivation
Inactivation

Figure S6: A: Calcium responses in Basin-2 with (38H09-LexA>LexAop-GCaMP6s 22E09-Gal4>UAS-TNT) or without (38H09-LexA>LexAop-GCaMP6s +/>UAS-TNT) Handle-b inhibition, in each trial of mechanosensory stimulation, in larvae fed on different diets (n = 12-14 larvae, 5 trials per larva). B: Individual calcium responses in Basin-2 from individuals fed on different diets, with or without optogenetic activation of Handle-b (22E09-Gal4>UAS-CsChrimson::tdTomato 38H09-LexA>LexAop-GCaMP6s) during the first second of mechanical stimulus (n = 8 larvae, 5 trials per larva). In all plots, the green dashed line corresponds to stimulus onset. C: Calcium responses in Basin-2 with (38H09-LexA>LexAop-GCaMP6s 55C05-Gal4>UAS-TNT) or without (38H09-LexA>LexAop-GCaMP6s +/>UAS-TNT) Griddle-2 inhibition, in each trial of mechanosensory stimulation, in larvae fed on different diets (n = 6-10 larvae, 5 trials per larva).

A Neurotransmitter expression in NPF neurons

A1 ChAT expression

A2 VGlut expression

A3 GABA expression

B Identifying Griddle-2 in L55C05

C Identifying Handle-b in L60E02

D NPFR expression

D1 NPFR expression in Griddle-2

D2 NPFR expression in Basin-1

D3 NPFR expression in Basin-2

E sNPFR expression

E1 sNPFR expression in Basin-1

E2 sNPFR expression in Basin-2

Figure S7. A.: Immunohistochemical labeling of NPF neuron neurotransmitters (NPF-Gal4>UAS-GFP) labels both pairs of NPF expressing neurons: dorsomedial (DM) and dorsolateral (DL). Brains were then stained with antibodies against either ChAT as a proxy of acetylcholine (A1), V-Glut as a proxy of glutamate (A2) and GABA (A3). Co-localization shows expression of acetylcholine in the DL NPF neurons and glutamate in both pairs of NPF descending neurons. Both pairs were negative for GABA antibody labeling. B: identifying griddle-2 in 55C05. The position of cell bodies and projections was compared between neurons in R55C05 (GAL4) (previously identified as Griddle-2) and L55C05 (LexA) that had a much broader expression pattern. C: Identification of Handle-b in line L60E02. L60E02 is driving the expression of LexAop-GCamP6s and GMR_SS00888 the expression of UAS-Chrimson-mCherry (60E02-LexA; UAS-Chrimson-mCherry, LexAop GCamP6s>GMR_SS00888). Specific antibodies against GFP and mCherry were used to increase detection sensitivity. Co-localization of jRGeco1a and GCamP6s show that the neuron with the cell body in the midline labeled by the 60E02-LexA is Handle-b. D1. Immunohistochemical labeling for NPFR in Griddle-2. UAS-GCamP6s is expressed in Griddle-2 using the SS_TJ001 split-Gal4 line (green) and LexAop-jRGeco1a is expressed under the control of the NPFR promoter using a T2A-LexA construct (magenta). Antibodies against GFP and dsRed were used to increase detection sensitivity. No expression of NPFR could be detected in Griddle-2. D2-D3: Basin-1 and -2 immunostaining for NPFR expression. UAS-GCamP6s is expressed in Basin-1 or Basin-2 using the R20B01 or GMR_SS00739 driver lines (green) respectively and LexAop-jRGeco1a is expressed instead under the control of the NPFR transcript using a T2A-LexA construct (magenta). Antibodies against GFP and dsRed were used to increase detection sensitivity. No expression of NPFR could be detected in Basin-1 or -2. E: Basin-1 and -2 immunostaining for sNPFR expression. LexAop-jRGeco1a is expressed in Basin-1 or Basin-2 respectively using the L20B01 or L38H09 lines (magenta) and UAS-GCamP6s is expressed under the control of the sNPFR promoter using a T2A-Gal4 construct (green). Specific antibodies against GFP and dsRed were used to increase detection sensitivity. No expression of NPFR could be detected in Basin-1 or -2.

A. Ca-imaging in Hb upon sNPFR knockdown in larvae fed on sucrose only

B. Ca-imaging in Hb upon sNPFR knockdown in starved larvae

C. Ca-imaging in G2 upon sNPFR knockdown in larvae fed on sucrose only

D. Ca-imaging in G2 upon sNPFR knockdown in starved larvae

Figure S8: Handle-b and Griddle-2 calcium response upon sNPFR knockdown in larvae fed on sucrose only and in starved larvae

A: Calcium responses in Handle-b sNPFR knockdown in Handle-b (GMR_SS00888>UAS-GCamP6s; UAS-sNPFR-RNAi) compared to the control (GMR_SS00888>UAS-GCamP6s) in larvae fed on sucrose only (top row) or starved (bottom row). Left panel: calcium responses of Griddle-2, in each trial of mechanosensory stimulation. Lower right panel: mean calcium response of Handle-b over time. The green dashed line corresponds to stimulus onset. Top right panel: response averaged during the stimulus. White line represents the mean, white dot represents the median. sNPFR knockdown does not influence the stimulus-induced activity of Handle-b in larvae fed on sucrose only or starved (n = 9-15 larvae, 4 trials per larva) .B: Calcium responses in Griddle-2 with (SS_TJ001>UAS-GCamP6s; UAS-sNPFR-RNAi) or without (SS_TJ001>UAS-GCamP6s) sNPFR knockdown in Griddle-2 in larvae fed on sucrose only (top row) or starved (bottom row). Left panel calcium responses of Griddle-2, for each trial Lower right panel: mean calcium response of Griddle-2 over time. The green dashed line corresponds to stimulus onset. Lower top panel: mean calcium response averaged during the stimulus. White line represents the mean, white dot represents the median. sNPFR knockdown does not influence the stimulus-induced activity of Griddle-2 in larvae fed on sucrose only or starved (n = 10 larvae, 3 trials per larva).