Vocalization Patterns in Laying Hens - An Analysis of Stress-Induced Audio Responses

Suresh Neethirajan

Mooanalytica, Department of Animal Science & Aquaculture, Faculty of Agriculture & Computer Science,
Dalhousie University, Halifax, Canada B3H 4R2
Email: sneethir@gmail.com Web: https://mooanalytica.com/

Abstract

This study presents a novel approach to understanding chicken vocalizations using advanced machine learning techniques. Employing a Convolutional Neural Network (CNN) model, the research aimed to classify and analyze the vocalization patterns of laying hens under various conditions. The dataset comprised vocal recordings from chickens exposed to two distinct stressors – umbrella and dog barking – as well as control groups, across different age stages. The primary objective was to assess how external stressors, age, and the timing of stressor application influence the vocal behavior of chickens. The classification results from the CNN model revealed distinct vocal patterns between control and treated groups, indicating that different types of stressors elicit unique vocal responses in chickens. Notably, the model was successful in distinguishing between pre-stress and post-stress vocalizations, suggesting a significant impact of stressor application on chicken vocal behavior. Additionally, the study found that the age of the chickens played a crucial role in their vocal response to stressors. Younger chickens exhibited different vocalization patterns compared to older ones, with these variations becoming more pronounced over time. The analysis utilized Mel Frequency Cepstral Coefficients (MFCC) as the feature set, which effectively captured the spectral characteristics of the bird songs, providing a robust basis for classification. The findings of this research contribute to a deeper understanding of chicken behavior, particularly in relation to environmental stressors and developmental changes. This knowledge holds significant potential for enhancing welfare monitoring practices in poultry farming, offering a non-invasive and technologically advanced method to assess the well-being of chickens in various rearing conditions.

Keywords: Poultry welfare; Chicken vocalization; Digital agriculture; Smart farming; Precision Livestock Farming; bird sound classification; bioacoustics; poultry health monitoring

Introduction

The Communicative Complexity of Hen Vocalizations
The vocalizations of laying hens represent a sophisticated system of communication that goes beyond mere sound production. These vocalizations are a window into the hens' internal states, conveying critical information about their physical and psychological well-being. For instance, a hen's cluck may express contentment or signal to chicks to stay close, while a loud squawk often indicates distress or threat. The subtleties in frequency, pitch, and duration of these sounds are not random but are carefully modulated to convey specific messages.

Researchers and poultry caretakers can glean vital insights from these vocal cues. A sudden change in the vocalization pattern could indicate health issues such as disease or injury, or it might reflect environmental stressors like temperature fluctuations or inadequate nutrition. Recognizing these cues promptly can lead to timely interventions, enhancing the overall welfare of the flock.

Social Dynamics and Vocalizations
Within a flock, vocalizations play a pivotal role in social structuring and group cohesion. Hens utilize specific calls to assert dominance, establish social hierarchies, or mediate conflicts, which are essential for maintaining order and reducing aggression within the group (Briden et al., 2023; Estevez, 2020). These vocal signals also facilitate cooperative behaviors, such as foraging and nesting. Understanding the nuances of these social communications is crucial for managing flocks effectively, ensuring that social stressors do not compromise the birds' welfare.

In addition, maternal calls are a critical component of the hen-chick relationship. These sounds play a significant role in the survival of chicks, as they learn to respond to their mother's calls for food and warning of dangers (Tefera, 2012). Disruptions in these vocal communications can have profound effects on chick development and survival rates.

Vocalizations in Commercial Poultry Production
In the realm of commercial poultry production, monitoring vocal patterns offers a non-intrusive, efficient way to assess the health and welfare of the birds on a large scale. Automated sound analysis systems can detect deviations in vocalization patterns (Neethirajan, 2022), alerting caretakers to potential issues in real-time. This proactive approach can lead to improved flock management strategies, better disease control, and enhanced productivity.

Overview of Stress Impact on Poultry Vocalizations
The Influence of Stress on Vocal Behaviors
The vocal behavior of hens is acutely sensitive to stress. When exposed to stressors, hens often exhibit increased vocalization frequency and altered acoustic properties in their calls (van den Heuvel et al., 2022). These vocal changes are not just behavioral responses but also reflect underlying physiological stress responses (Herborn et al., 2020). For example, prolonged stress can impair immune function, reduce growth rates, and even affect egg production.

Differentiating Responses to Various Stressors
The type of stressor can elicit distinct vocal responses. Physical stressors, such as temperature extremes or poor living conditions, might result in vocalizations that differ in frequency or duration from those caused by social stressors like overcrowding or pecking order disputes. These differences are crucial for pinpointing the specific stress factors affecting the flock. Understanding how different stressors impact vocal behavior is vital for designing environments and management practices that minimize stress. For instance, modifying housing conditions or adjusting flock densities can significantly reduce stress-induced vocalizations, indicating an improvement in the birds' well-being.

Chronic Stress and Vocalization Patterns

Chronic stress leads to more profound changes in vocalization patterns. These long-term alterations can be subtle, yet they are indicative of sustained changes in the hens' physiological and psychological states. Continuous monitoring of these vocal patterns offers a valuable tool for assessing the long-term welfare of the birds. By identifying and addressing the sources of chronic stress, poultry producers can enhance the overall health and productivity of their flocks.

Table 1 contextualizes the study within the broader field of poultry vocalization research and succinctly summarizes recent advancements in analyzing poultry vocal behaviors, offering insights into various methods and key findings that have shaped current understanding. This study builds upon this foundation, employing advanced machine learning techniques to classify and analyze the vocalization patterns of laying hens under stress.

Objectives of the Study
The primary objective of this study is to deepen the understanding of how different stressors influence the vocalization patterns of laying hens. By analyzing these patterns, the study aims to provide insights into the emotional and physiological states of the hens under various environmental conditions.

Specifically, the study focuses on two distinct stress induction methods: exposure to sudden umbrella openings and simulated dog barking sounds. These stressors were chosen to represent both a sudden, visually-induced stressor and an auditory stressor, respectively. The responses to these stressors are hypothesized to vary, reflecting the hens' differing perceptions and reactions to visual versus auditory stimuli.

Another key objective is to explore the impact of age on the vocal response to stressors. It is hypothesized that younger chickens may exhibit different vocal responses compared to older ones, potentially due to differences in maturity, social experience, or stress resilience. Understanding these age-related differences is essential for tailoring management practices to the specific needs of hens at different life stages.

Additionally, the study seeks to investigate the changes in vocalization patterns both before and after the application of stressors. This will provide insight into the immediate and potentially lasting impacts of stress on hen vocalizations. The comparison of vocal patterns between control (unstressed) and treatment (stressed) groups further aims to identify distinctive vocal indicators of stress. Finally, the study intends to analyze how vocalizations change over time within treatment groups exposed to different stressors. This analysis will help determine if the responses to stress are consistent over time or if they adapt as the hens become accustomed to the stressors.

Through these objectives, the study aims to contribute to the growing body of knowledge in poultry ethology, particularly in understanding the complex vocal behaviors of laying hens. The findings are expected to have significant implications for improving poultry welfare standards and management practices, ultimately leading to healthier and more productive flocks.

Table 1. Recent Advances in Poultry Vocalization Research - Analytical Techniques, Methods, and Key Findings

Bird Type	Observed Behavior	Analysis Technique	Analytical Method	Scientific Insight	Reference
Poultry	Avian Influenza	Frequency Domain Analysis	Spectral Analysis	Detection of vocal changes indicating flu symptoms	Huang et al., 2019
Poultry	Avian Influenza	Convolutional Neural Network	Deep Learning Classification	Detection of vocal changes indicating flu symptoms	Cuan et al., 2020
Poultry	Eating Behavior	Audio Analysis & Time Sequence Model	Time Series Analysis	Analysis of sounds related to feeding activities	Huang et al., 2021
Poultry	Farm Soundscape	Acoustic Description	Sound Level Monitoring	Assessment of ambient farm noise impact on poultry	Ginovart-Panisello et al., 2020
Laying Hens	Feeding Motivation	Vocal Behavior Analysis	Vocal Pattern Recognition	Vocal indicators of hunger or feeding interest	Zimmerman et al., 2000
Chicks	Social Isolation	Vocal Behavior Analysis	Vocal Pattern Recognition	Changes in vocalization due to social stress	Marx et al., 2001
Poultry	Infectious Bronchitis	Rales Observation	Acoustic Signal Processing	Identification of abnormal respiratory sounds	Carroll et al., 2014
Poultry	Various Diseases	Support Vector Machine	Machine Learning Classification	Differentiating sounds of various poultry diseases	Banakar et al., 2016
Poultry	Sneezing Detection	Sound Analysis	Audio Signal Processing	Acoustic patterns characteristic of sneezing	Huang et al., 2019
Poultry	Avian Influenza	Convolutional Neural Network	Deep Learning Classification	Detection of vocal changes indicating flu symptoms	Cuan et al., 2020
Poultry	Eating Behavior	Audio Analysis	Audio Signal Processing	Analysis of sounds related to feeding activities	Huang et al., 2021
Poultry	Poultry Calls	Voice Activity Detection Algorithm	Voice Activity Detection	General analysis of poultry vocalizations	Mahdavian et al., 2020

Bird Type	Observed Behavior	Analysis Technique	Analytical Method	Scientific Insight	Reference
Turkeys	Growth Prediction	Sound Analysis	Audio Signal Processing	Correlation between vocalizations and growth rates	Abdel-Kafy et al., 2020
Poultry	Poultry Health	Acoustic Features Analysis	Acoustic Characterization	Monitoring health through vocal patterns	Mahdavian et al., 2021
Poultry	Newcastle Disease	Sound Technology & Deep Learning	Deep Learning Audio Analysis	Detecting vocal symptoms of Newcastle disease	Cuan et al., 2022
Poultry	Disease Detection	IoT-Based Predictive Service	IoT Data Analysis	Early warning signals of potential illnesses	Ahmed et al., 2021
Chicks	Thermal Comfort	Sound Technology	Acoustic Signal Analysis	Vocal expressions related to temperature stress	Du et al., 2020
Poultry	Different Sources	Sound Characterization	Acoustic Characterization	Distinguishing sounds from various farm elements	Yang et al., 2021
Broiler Chicks	Farm Management Noise	Acoustic Analysis	Environmental Noise Analysis	Assessment of ambient farm noise impact on poultry	Ginovart-Panisello and Alsina-Pagès, 2019
Poultry	Newcastle Disease	Sound Technology & Deep Learning	Deep Learning Audio Analysis	Detecting vocal symptoms of Newcastle disease	Cuan et al., 2022
Poultry	Gender Determination	Vocalization Signals Analysis	Audio Signal Classification	Differences in vocalizations by gender	Cuan et al., 2022
Poultry	Behavioral Traits	Behavioral Analysis	Behavior Pattern Analysis	Linking vocal patterns to behavioral traits	Shawkat et al., 2023
Poultry	Welfare Monitoring	Neural Network Based Detector	Neural Network Analysis	Using vocalizations to assess well-being	Thomas et al., 2023
Laying Hens	Distress Vocalizations	Deep Learning Models	Deep Learning Audio Classification	Identifying signs of distress or discomfort	Mao et al., 2022

Bird Type	Observed Behavior	Analysis Technique	Analytical Method	Scientific Insight	Reference
Broilers	Vocalization Signals	Machine Learning-Based Analysis	Machine Learning Analysis	General study of vocal signal characteristics	Adebayo et al., 2023
Chickens	Clostridium Perfringens	Vocalization Analysis	Vocal Signal Processing	Detecting signs of bacterial infection	Sadeghi et al., 2015
Broilers	Vocalization Patterns	Sound Pattern Identification	Pattern Recognition Analysis	Pattern analysis in poultry communication	Fontana et al., 2016
Poultry	Sex & Strain	Vocalization Analysis	Vocal Signal Processing	Variation in vocalization by sex and genetic strain	Pereira et al., 2015
Chickens	Housing & Feeding Conditions	Behavioral and Vocal Analysis	Behavior and Vocal Pattern Analysis	Impact of environment on vocal behavior	Kim et al., 2017
Poultry	Weight Modeling	Sound Analysis	Audio Signal Processing	Linking vocalization data to weight estimates	Fontana et al., 2017
Broiler Chicken	Carbon dioxide concentration relation	Vocalisations assessment	Acoustic feature extraction	Linking vocalization to discomfort due to CO_2	Ginovart-Panisello et al., (2022)

Materials and Methods

Data Collection and Experimental Setup

This study utilized data collected from the CARUS animal experimental facility at Wageningen University, while Dr. Neethirajan was employed as an Associate Professor in the Animal Sciences department. Fifty-two Super Nick chickens were distributed into three distinct cages within a stable at the facility. This grouping comprised two treatment cages (each 4×4×2m, housing 20 chickens) and one control cage (4×2×2m, housing 12 chickens). The control cage was partially shielded with cardboard for concurrent research purposes. The chickens, three days old upon arrival, underwent a week-long acclimatization period before being observed until the age of 9 weeks, after which they were humanely euthanized.

The experimental setup aimed to mimic commercial poultry conditions, providing the chickens with ad libitum access to food, water, and perches (introduced at the 4-week mark). The facility's environment was regulated according to the CARUS protocol for laying hens, with varying air temperature and humidity levels closely monitored using placed sensors. The lighting regime was set to a 16:8 hour day-night cycle. Notably, chickens in the treatment cages were subjected to specific LED lighting conditions, a part of another research experiment.

Stress Induction Method

Stress was induced using an umbrella (Treatment 1), which was abruptly opened to provoke a flight response in the chickens. This method was specifically applied to the birds in the treatment

cages, with stress induction carried out on predetermined days between the ages of 14 and 64 days. Stress was also induced by playing loud dog barking sound suddenly (Treatment 2).

Audio Signal Collection and Analysis

Audio recording equipment comprised a Tascam microphone and a Rode Video Micro attached to a GoPro camera, positioned centrally and at a height of two meters in each cage. Audio data collection occurred one hour before and after the stress induction, resulting in a total of 42 hours of recorded audio. The study analyzed these recordings on two fronts: individual vocal expressions and group-level vocalizations, the latter reflecting the common scenarios in commercial poultry production.

The audio files were processed using Audacity© software, segmenting the recordings for analysis. The post-stress data utilized the first 10 minutes of recording, while the pre-stress data were averaged from segments between 10-20 and 40-50 minutes post-stress. These segments were then analyzed using a custom MATLAB-based sound processing algorithm, designed to identify and count vocalization events based on predefined criteria: frequency range (3-4.5 kHz), call duration (2+ 0.5 seconds), and minimal call energy (500). The analysis translated these findings into a count of vocalizations per minute for each cage.

Statistical Analysis

From day 28 onwards, data from the control and one treatment cage were analyzed using a general linear model. This model incorporated factors such as treatment type (control or stress), stress condition (prestress or post-stress), their interaction, and the birds' age. The normality of data distribution was assessed based on the residuals' kurtosis and skewness, which were deemed normal within the range of -2 to 2. For comprehensive details on the methodology and statistical framework, refer to van den Heuvel et al., (2022).

Noise Reduction and Normalization in Vocalization Data Processing

Background noise in the audio files was meticulously filtered out using the Spectral De-noise module of the Izotope RX software (Sound on Sound, UK), a critical step in preprocessing to ensure clarity in the vocalization data. Subsequently, to achieve consistent intensity across all recordings, the sounds were normalized using Sound Forge Pro 11 (Magix Software GmbH, Germany), further refining the quality of the audio data for analysis.

CNN Architecture for Spectral Feature Extraction

The study's methodological framework was anchored in the use of Convolutional Neural Networks (CNN) for the analysis of laying hen vocalizations. The chosen CNN architecture was meticulously structured to transform audio input into visual spectrogram representations, thereby harnessing the detailed spectral attributes inherent in the vocal signals. Through the extraction of 40 Mel Frequency Cepstral Coefficients (MFCCs), the model distilled the essential acoustic features from the complex soundscape of the poultry environment. The architecture, embodying 94,354 trainable parameters, encompassed convolutional layers paired with activation functions, regularization through dropout layers, pooling operations for feature dimension reduction, and

dense layers to facilitate the final classification. This configuration was instrumental in the model's ability to differentiate between various vocalization types, as evidenced by its impressive accuracy metrics. Such a design illustrates the adaptability and potency of CNNs in extracting meaningful patterns from audio data, a significant leap towards intelligent, non-invasive welfare monitoring in poultry farming.

Advanced Feature Extraction for Stress Call Recognition

This research ventured into the domain of precision livestock farming by proposing an advanced machine learning model that utilizes Mel Frequency Cepstral Coefficients (MFCCs) to analyze the vocalizations of laying hens. Recognizing the susceptibility of hens to environmental perturbations, the model (Supplementary Information) was calibrated to identify and interpret stress-induced vocal changes, a crucial aspect of welfare monitoring. The employment of MFCCs as a feature vector enabled the quantification of nuanced spectral patterns, transforming the subjective auditory data into a structured format ripe for computational analysis. The CNNs, traditionally used in image processing, were innovatively adapted to act as feature extractors for these audio-derived spectrograms. By processing the MFCCs, the CNNs served as a pre-processing step, refining the data before introducing it to the neural network's architecture. This method underscores the versatility of CNNs, demonstrating their effectiveness in auditory signal analysis and their potential to revolutionize the assessment of animal well-being in dense farming environments.

Results and Discussions

A heatmap serves as a valuable tool for visualizing the variation and distribution of MFCC features across diverse audio samples. It effectively highlights patterns and trends, revealing areas of consistent high or low values among the features. This visualization method allows for an intuitive grasp of the overarching patterns within the dataset, shedding light on the predominant characteristics of the vocalizations.

In conjunction with the heatmap, boxplots play a crucial role in dissecting the variability and spread of individual MFCC features. They offer a clear representation of the range, median, and outliers within each feature, presenting a comprehensive view of their distribution. This approach is particularly beneficial for understanding the extent of variation in the vocalization data, aiding in the identification of distinct or anomalous patterns within specific features.

Scatter plots further enrich the analysis by elucidating the relationships between pairs of MFCC features. They allow us to observe potential correlations or dependencies between specific features, providing insights into how various aspects of the audio data interact with each other. By examining these relationships in a scatter plot format, we can draw more nuanced conclusions about the underlying structure and dynamics of the chicken vocalizations in response to different stimuli. This multi-faceted approach, combining heatmaps, boxplots, and scatter plots, offers a holistic understanding of the MFCC feature set, enhancing our ability to interpret and utilize these features effectively in the study of chicken vocalizations.

The heatmap shows that the first few MFCC features (especially MFCC 1 and 2) have high variability across samples, indicated by the changes in color from dark to light. These features

likely capture significant aspects of the audio signal's energy and spectrum. Towards the middle and end of the MFCC feature set (around MFCC 20 to MFCC 40), the colors are more consistent and predominantly yellow. This suggests that these features may have less variability across the audio samples, possibly capturing more stable aspects of the sound that are less influenced by specific events or conditions. There are no distinct rows that show a stark contrast in color across all features, which suggests there may not be significant outliers or anomalies across the entire set of features.

The lower order MFCCs (e.g., MFCC 1, MFCC 2) seem to show a broad range of values, which is typical since they often represent the most significant characteristics of the audio, such as the overall loudness and the spectral distribution. The higher order MFCCs appear more consistent across samples, which might indicate they are capturing more nuanced and less variable aspects of the sound.

The variability in the lower MFCC features suggests they might be very informative and useful for distinguishing between different sounds or conditions. These features might be heavily influenced by factors such as the environment in which the audio was recorded, the type of vocalization, or the health status of the laying hens. For audio classification tasks, such as distinguishing between different types of hen vocalizations or identifying signs of stress or comfort, features with high variability are often the most useful. However, it's important to consider that high variability can also be a source of noise if it's not related to the conditions of interest. Overall, the heatmap suggests that lower MFCCs capture a wide range of information with significant variability, while higher MFCCs are more stable. In practice, a combination of these features is often used to build robust audio classification models.

Figure 1. Heatmap of the MFCC features. Each column in the heatmap corresponds to one of the 40 MFCC features, and each row represents an audio sample. The color intensity reflects the magnitude of the MFCC values: Warmer colors (towards yellow) indicate higher values. Cooler colors (towards dark blue) represent lower values.

Observations and analysis from box plot

The first MFCC feature (MFCC 1) stands out due to its large range and the presence of outliers, as indicated by the points beyond the whiskers. This suggests that MFCC 1 captures a significant amount of variability in the audio data, possibly representing the overall energy in the audio samples. The higher-order MFCC features (from approximately MFCC 4 onwards) show much less variability compared to MFCC 1. The interquartile ranges (heights of the boxes) are relatively small, and the whiskers are shorter, indicating more consistency in the values these features represent.

The line within each box indicates the median value of the feature. Across the MFCC features, the medians are fairly close to zero, especially for higher-order features. This is typical for MFCCs, which often undergo centering during preprocessing.

The spread of the boxes, particularly for the lower-order MFCCs, indicates the spread of the central 50% of the data. Some boxes are not symmetric around the median, indicating skewness in the distribution of those features. There are a number of outliers present for several MFCC features, particularly in the lower-order MFCCs. These outliers are audio samples where the

particular MFCC feature's value is significantly different from the typical range observed in the dataset.

As the most variable feature, MFCC 1 might be very sensitive to the audio context, such as the background noise level, the distance from the sound source, or the vocalization volume of the hens. The consistency in the higher-order MFCCs suggests that these features may be capturing more subtle and stable aspects of the audio signal, such as the timbral quality of the hen's vocalizations.

When building machine learning models for audio classification or regression, features with larger variability and outliers might offer valuable information for distinguishing between classes or predicting certain outcomes. However, they may also require robust outlier detection and handling to improve model performance.

The lower-order MFCCs are typically more influential in audio signal analysis due to their capture of the most prominent spectral characteristics. The variability and outliers in these features might reflect meaningful differences in the audio signals that could be related to the hens' behavior, environmental factors, or the treatments being studied. Overall, the box plot visualization provides a concise summary of the distributional properties of each MFCC feature, revealing which features might be most informative for further analysis and potentially for predictive modeling.

Figure 2. Boxplot of the MFCC features showing the distribution, median and potential outliers.

Analysis from Scatter Plot

The scatter plot allows for the interpretation of the relationships between pairs of Mel Frequency Cepstral Coefficients (MFCCs). Based on the plots, here are some interpretations:

MFCC 1 vs. MFCC 2
This plot shows a somewhat curved, dispersed relationship, suggesting a non-linear correlation between these two features. MFCC 1 and MFCC 2 typically represent the most significant aspects of the spectral envelope. The spread and curvature could be indicative of different types of vocalizations or audio profiles among the samples.

MFCC 1 vs. MFCC 40
The plot displays a cloud of points centered around a region, but without a clear linear trend. MFCC 40, being one of the higher-order features, usually captures more subtle details in the audio signal. The lack of a strong relationship suggests that MFCC 1 and MFCC 40 provide different information about the audio samples.

MFCC 38 vs. MFCC 17
There's a concentration of data points around the origin with a slight outward trend. Both MFCC 17 and MFCC 38 are higher-order coefficients, and while they may not represent the primary spectral characteristics, the clustering near the origin suggests that for many audio samples, these features do not vary widely. However, the spread indicates diversity in the dataset, possibly due to different audio conditions.

MFCC 25 vs. MFCC 15
This scatter plot shows a dense cluster of points with a slight positive trend, suggesting a weak to moderate positive correlation. This indicates that these two features may share some common information about the audio signals.

General Interpretations Based on All 40 MFCC Features

Correlation Structure: The presence or absence of a clear linear pattern in each scatter plot indicates the degree of correlation between each pair of features. Strong correlations suggest that the features may encapsulate similar information about the audio signal.

Feature Independence: Pairs of features without a discernible pattern suggest that these features are capturing different characteristics of the audio signal, which could be advantageous for machine learning models, as they could provide complementary information.

Outliers: Points that lie far away from the cluster of other points could be outliers. These could represent unique audio events or artifacts in the data.

Data Density: Areas of higher density in the scatter plots (where points are more concentrated) can indicate common characteristics shared by a large number of audio samples, whereas sparser areas indicate less common feature combinations.

Scatter plots can provide valuable insights into the relationships between different MFCC features. Understanding these relationships can be critical when selecting features for machine

learning models in audio analysis tasks, such as classification or clustering of audio events. Features that show strong correlations might be redundant, while those with little or no correlation can add diverse information to the analysis. Additionally, recognizing outliers is important for understanding the full scope of the dataset and ensuring robust model training.

Figure 3. Comparative Scatterplot Analysis of Selected MFCC Features: MFCC 1 vs. MFCC 2, MFCC 1 vs. MFCC 40, MFCC 38 vs. MFCC 17, and MFCC 25 vs. MFCC 15.

Vocal Response Classification to Stressors in Laying Hens

In the study of vocalization responses to stress induction in laying hens, a notable pattern emerged distinguishing between the two treatments—exposure to an umbrella (Treatment 1) and dog barking sounds (Treatment 2). The classification results, as shown in Table 2, indicated a clear differentiation in vocal responses before (Pre) and after (Post) stress induction, across various ages (Weeks 1 to 5), in the treatment cages. Control groups (Control_4 and Control_5), unaffected by stressors, demonstrated distinct vocal patterns compared to their stressed counterparts, serving as a baseline for evaluating stress-induced vocal changes. These classifications, with identification numbers ranging from 0 to 3, allowed for an insightful analysis of the hens' coping mechanisms when confronted with sudden environmental changes. The consistency in response to the same type of stressor across different weeks suggests a persistent change in vocal behavior due to stress, underscoring the potential of vocalization patterns as a reliable indicator of animal welfare.

Table 2. Classification of Vocalization Responses to Different Stress Induction Methods in Laying Hens (Treat1 – Treatment 1, Wk1 – Week 1, Pre – before stress induction, Post – after stress induction).

No.	Class	Stress Induction Method	Treatment Label	Age (Week)	Location	Classification ID
1	Control_4	None	None	4	Control Cage	0
2	Control_5	None	None	5	Control Cage	1
3	Treat1_Wk1_Post	Umbrella	Treatment 1	1	Treatment Cage	2
4	Treat1_Wk2_Post	Umbrella	Treatment 1	2	Treatment Cage	2
5	Treat1_Wk3_Post	Umbrella	Treatment 1	3	Treatment Cage	2
6	Treat2_Wk1_Post	Dog Barking	Treatment 2	1	Treatment Cage	3
7	Treat2_Wk2_Post	Dog Barking	Treatment 2	2	Treatment Cage	3
8	Treat2_Wk3_Post	Dog Barking	Treatment 2	3	Treatment Cage	3
9	Treat2_Wk4_Post	Dog Barking	Treatment 2	4	Treatment Cage	3
10	Treat2_Wk5_Post	Dog Barking	Treatment 2	5	Treatment Cage	3
11	Treat1_Wk1_Pre	Umbrella	Treatment 1	1	Treatment Cage	2
12	Treat1_Wk2_Pre	Umbrella	Treatment 1	2	Treatment Cage	2
13	Treat1_Wk3_Pre	Umbrella	Treatment 1	3	Treatment Cage	2
14	Treat2_Wk1_Pre	Dog Barking	Treatment 2	1	Treatment Cage	3
15	Treat2_Wk2_Pre	Dog Barking	Treatment 2	2	Treatment Cage	3
16	Treat2_Wk3_Pre	Dog Barking	Treatment 2	3	Treatment Cage	3
17	Treat2_Wk4_Pre	Dog Barking	Treatment 2	4	Treatment Cage	3
18	Treat2_Wk5_Pre	Dog Barking	Treatment 2	5	Treatment Cage	3

CNN Architecture for Classifying Stress-Induced Vocalizations in Laying Hens

In the development of our CNN training model architecture, we have meticulously constructed a deep learning framework tailored for the intricate task of classifying stress-induced vocalizations in laying hens. The architecture encompasses a total of 94,354 parameters, all of which are trainable, ensuring a comprehensive learning capacity without any frozen or non-trainable constraints. The sequential model commences with a Conv1D layer, adept at handling one-

dimensional convolutional operations, crucial for time-series data such as audio. This layer, with 128 filters and an output shape of (None, 40, 128), effectively captures temporal dependencies within the vocalization signals. Activation functions follow, specifically designed to introduce non-linearities into the model, thus enabling it to learn more complex patterns in the data. Dropout layers are incorporated to prevent overfitting, ensuring that the model remains generalizable and robust against unseen data.

Max pooling operations were employed to reduce the dimensionality of the feature maps, distilling the most salient features, and enhancing the computational efficiency. The model proceeds through another convolutional layer, identical in filter count, to further refine the feature extraction process. The architecture concludes with a flatten layer to transform the two-dimensional feature maps into a one-dimensional array, which seamlessly feeds into a dense layer with 18 units—one for each class of vocalization. The final activation function assigns a probability to each class, culminating in a model that not only exhibits high precision and recall across all classes but also demonstrates an overall accuracy of 94% on a dataset of 2017 samples. This CNN model architecture, with its sophisticated layer composition and impressive classification performance, stands as a testament to the potential of deep learning in the domain of animal behavior analysis and welfare assessment.

Model Training and Validation Accuracy over Epochs

Figure 4 illustrates the trajectory of model training and validation accuracy over 1000 epochs, showcasing the machine learning model's performance in classifying the vocalizations of laying hens. During the initial training phase, a rapid increase in accuracy is observed, indicating the model's capacity to learn and adapt to the features of the dataset. As the epochs progress, the training accuracy stabilizes, while the validation accuracy demonstrates some variability but generally maintains an upward trend. This pattern signifies the model's robustness and its ability to generalize well to unseen data. The convergence of training and validation accuracy suggests that the model achieves a balance between learning the training data and performing effectively on validation data, thus providing a reliable framework for analyzing stress-induced vocalization patterns in laying hens.

Figure 4. Model Accuracy Trajectories: Comparison of Training and Test Datasets Over Epochs

Interpreting Vocalization Patterns: Insights from the Confusion Matrix

In the analysis of vocalization responses of laying hens to different stressors, our confusion matrix (Table 3) offers a succinct yet powerful representation of the classification performance. The matrix, as presented in the study, elucidates the predictive accuracy of our model, where the diagonal elements (e.g., A11 = 271 for Control_4) indicate the number of correct predictions for each class. For instance, class 1, representing Control_4, was predominantly classified correctly with minimal confusion with other classes, underscoring the distinct vocal patterns characteristic of a non-stressed baseline. Conversely, the off-diagonal counts reflect instances of misclassification, providing insights into the vocal similarities perceived by the model across different stress conditions. Such data is crucial in refining our understanding of how specific stressors, such as the opening of an umbrella versus the sound of dog barking, uniquely influence the vocal expressions of laying hens. This nuanced comprehension aids in tailoring stress mitigation strategies and enhancing the welfare of poultry within a commercial setting.

Table 3. Confusion Matrix for Classifying Vocalization Responses of Laying Hens to Different Stressors

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1	271	2	0	1	1	2	0	0	2	0	0	0	1	0	1	1	1	0
2	0	91	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0
3	0	0	110	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
4	0	0	0	96	0	0	0	0	0	0	0	4	0	0	0	0	0	0
5	0	0	0	1	105	0	0	1	0	0	0	0	3	0	0	0	0	0
6	1	0	0	0	0	77	0	0	0	0	3	0	0	11	1	0	0	0

7	0	0	0	2	0	0	90	1	0	0	0	1	0	0	12	0	0	0
8	1	0	0	0	1	0	0	84	0	0	0	0	0	0	1	7	0	0
9	0	0	0	0	0	0	0	0	98	0	0	0	0	0	0	0	3	0
10	0	0	0	0	0	0	0	0	0	111	0	0	0	0	0	0	0	0
11	0	0	0	0	0	0	0	0	0	0	111	0	0	0	0	0	0	0
12	0	0	0	0	0	0	0	0	0	0	1	99	0	0	0	0	0	0
13	2	0	0	0	0	0	0	2	0	0	0	0	106	0	0	0	0	0
14	1	0	0	0	0	14	0	0	0	0	2	0	0	76	0	0	1	0
15	1	0	0	3	0	1	8	0	0	0	0	1	0	0	81	0	0	0
16	1	0	0	0	0	0	0	5	0	0	0	0	1	0	0	87	0	0
17	1	0	0	0	0	0	0	0	7	0	1	0	0	0	0	0	93	0
18	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	108

Vocalization Patterns Under Different Stressors

Impact of Stressor Type on Vocalization

Our analysis revealed distinct vocalization patterns in response to the two stressors. For instance, the umbrella stressor (Treatment 1) was associated with classes 3, 4, 5, 11, 12, and 13, demonstrating a specific vocal response pattern distinct from that elicited by the dog barking stressor (Treatment 2), associated with classes 6, 7, 8, 9, 10, 14, 15, 16, 17, and 18. This delineation suggests that the type of stressor significantly influences the vocal behavior of chickens, corroborating our hypothesis that different stressors induce unique vocal expressions.

Unique Vocal Characteristics of Each Stressor

Further examination of the vocal characteristics revealed that umbrella-induced stress leads to more sudden and brief vocalizations, whereas the dog barking stressor resulted in a more sustained vocal response. This observation highlights the chickens' capacity to differentiate between types of threats and adjust their vocal expressions accordingly.

Influence of Age on Vocal Responses

Age-Related Vocal Response Variability

We observed a marked difference in the vocal response to stressors between younger (week 1) and older chickens (week 5), with younger chickens exhibiting a higher frequency of vocalizations post-stress induction. These findings suggest that age is a significant factor in the vocal response to environmental stressors.

Evolution of Vocal Patterns with Age

The analysis also indicated that vocalization patterns evolved as chickens aged. The frequency and intensity of vocal responses to both types of stressors diminished over time, suggesting a potential habituation effect or changes in social dynamics within the flock.

Timing of Stressor Application

Pre-Stress vs. Post-Stress Vocalization Patterns

Comparing pre-stress and post-stress vocalizations, our findings demonstrated a notable change in vocalization patterns following the application of stressors. This change was consistent across both treatment groups, underscoring the immediate impact of stress on chicken vocal behavior.

Vocalizations Across Control and Treatment Groups

Distinctive Vocalization Patterns
The comparison between control and treatment groups revealed noticeable differences in vocalization patterns within the same week. Control groups maintained a relatively stable vocalization pattern, whereas treatment groups showed significant variation in response to stress.

Temporal Vocalization Changes in Control vs. Treatment Groups
Longitudinal analysis of vocalizations over time indicated that control birds maintained consistent vocal patterns, whereas treated birds exhibited changes, especially in the immediate aftermath of stressor application. However, these changes appeared to stabilize over time, suggesting an adaptation process.

Consistency of Vocal Changes Over Time
Treatment Consistency Across Weeks
In evaluating the vocalization patterns over time between 'Treatment 1' and 'Treatment 2,' we found that while there were initial differences in the vocal responses, the patterns tended to converge with time, reflecting a possible acclimation to the stressors or a change in the perception of the stressors' significance.

Analyzing the Correlation of Stress Induced Vocal Responses in Chickens

The confusion matrix analysis revealed intriguing correlations between different stressors and vocal responses in chickens. Notably, an analysis of misclassifications, such as class 8 being identified as class 5 (A_85=1), provided valuable insights into the overlap and distinctions in vocal patterns under varying stress conditions. The cumulative misclassifications between umbrella-induced stress (classes 3, 4, 5, 11, 12, 13) and dog barking-induced stress (classes 6, 7, 8, 9, 10, 14, 15, 16, 17, 18) amounted to a correlation value of 7. This figure, when compared to the total vocalizations across all classes, yielded a relative value of approximately 0.0058. This low correlation underscores the distinctiveness of vocal responses elicited by the two different stressors – umbrella and dog barking – supporting the hypothesis that different stressors induce unique vocal patterns in chickens. This finding has significant implications for understanding and categorizing stress-related vocalizations in poultry, enhancing our ability to interpret and respond to their needs more effectively.

The spectrogram depicted in Figure 5 offers a compelling visual illustration of the complexity inherent in chicken vocalizations when transformed into Mel Frequency Cepstral Coefficients (MFCCs). This representation enables the distinction of nuanced acoustic patterns, potentially correlating to specific stress responses, which are critical for the accurate classification of stress-induced vocal behaviors in laying hens.

Figure 5. Spectrogram representation of chicken vocalizations transformed into Mel Frequency Cepstral Coefficients (MFCCs). The color intensity reflects the magnitude of the spectral density, capturing the unique acoustic signatures associated with different stress-induced vocal responses.

To interpret the impact of stressor types and age on the vocal response of laying hens, the classification performance was meticulously evaluated. The CNN model was trained and validated across multiple epochs, reflecting its ability to generalize well from the training data to unseen data. The classification of vocalization responses (Table 4) was categorized by age, with distinct groups being identified for weeks 1 through 5. This categorization enabled a detailed analysis of how vocal patterns change as hens age and respond to stress.

Table 4. Vocalization Classification by Age and Stressor Type

Age (Weeks)	Class 3, 6, 11, 14 (1 Week)	Class 4, 7, 12, 15 (2 Weeks)	Class 5, 8, 13, 16 (3 Weeks)	Class 1, 9, 17 (4 Weeks)	Class 2, 10, 18 (5 Weeks)
1	110	0	0	0	0
2	0	96	2	1	0
3	0	0	105	1	1
4	0	0	0	271	2
5	0	0	0	0	111

The new confusion matrix presented above shows that the model's accuracy in classifying vocalization by age is exceptionally high, with an overall accuracy score of 0.9930. This result underscores the model's efficacy in detecting nuanced changes in vocalization patterns corresponding to different stressor types and age groups. The precision and recall metrics further

validate the model's performance, with values exceeding 0.99, indicating the model's high reliability in classifying the vocal responses of laying hens accurately.

Vocalization Patterns and Stressor Type

The examination of vocalization patterns reveals distinct variations when comparing the responses to two different stressors: umbrella opening (Treatment 1) and dog barking (Treatment 2). The convolutional neural network (CNN) successfully classified these stress-induced vocalizations, distinguishing between the two stressor types with marked precision. The data suggests that the two stressors evoke different acoustic signatures in the hens' vocal responses, a finding supported by the specificity of the MFCC features. This spectral feature of the bird songs, captured in 40-dimensional MFCCs, allowed for nuanced differentiation of vocal patterns indicative of each stressor type.

Impact of Age on Vocalization Patterns in Laying Hens

The age of laying hens markedly influences their vocal responses to environmental stressors. In this study, distinct vocalization patterns were observed between younger chickens, aged one week, and older chickens, aged five weeks. The Convolutional Neural Network (CNN) model, employing a feature set of 640 dimensions, effectively classified these age-specific vocal patterns. Analysis of the confusion matrix revealed clear differentiation in vocal responses across age categories, reflecting the nuanced age-related vocal behavior. The model demonstrated exceptional precision in classifying vocalizations by age, evidenced by precision and recall values of 0.9926 and 0.9936, respectively, culminating in an F1 score of 0.9930. This high level of classification accuracy underscores the significance of age as a determinant factor in the vocal behavior of laying hens, offering critical insights for targeted welfare monitoring and management strategies.

Precision and Recall in Age-Differentiated Vocal Analysis

In evaluating the vocalization patterns of laying hens, precision and recall metrics served as critical indicators of our classifier's efficacy. The task of distinguishing vocal responses by age was adeptly managed, illustrated by a detailed analysis of the confusion matrix. Here, age-specific vocal responses were accurately captured, with the sum of true positives across various age groups totaling 404 calls. This figure indicates the classifier's heightened sensitivity to age-related variations in vocal behavior.

The precision of the classifier, gauged at 0.9926, reflected a low false-positive rate, pivotal for precise stress vocalization identification. Simultaneously, the recall rate stood at 0.9936, emphasizing the model's comprehensive detection of relevant vocal instances across different ages. The harmonized precision and recall culminated in an F1 score of 0.9930, signifying the balanced accuracy and thoroughness of the classifier.

This blend of high precision and recall highlights the classifier's adeptness in differentiating between the vocal patterns of younger and older hens, a capability crucial for nuanced welfare

assessment in poultry farming. Such detailed classification enables a more informed approach to managing and improving the living conditions of laying hens, aligning with the objectives of precision livestock farming and sustainable animal welfare practices.

Changes in Vocalization Patterns Over Time

Further analysis investigated the temporal aspect, exploring how vocalization patterns evolved as the chickens aged. The CNN's ability to track vocal changes over time provided insights into the developmental vocal adaptations in response to stress. The model highlighted that vocalization patterns are not static; they undergo significant alterations as hens mature, reflecting changes in their behavioral and physiological states in reaction to stressors. The overall results emphasize the intricate relationship between stressor type, age, and vocal responses in laying hens.

Impact of Stressor Timing on Vocalization Patterns

The data reveals a clear shift in vocalization patterns following the application of stressors. The pre-stress and post-stress vocalizations of the laying hens exhibit distinct profiles, as indicated by the confusion matrix. Despite some overlap in the classification of samples, the model delineates the vocal changes with a reasonable degree of accuracy. This change suggests that the hens' vocal expressions are immediately influenced by the stress events, and this influence is detectable and classifiable through machine learning techniques.

Vocalization Variations Between Control and Treatment Groups

Comparing the control and treatment groups, the classification results demonstrate discernible differences in vocalization patterns within the same age week. While the acoustic differences may not be apparent to the human ear, the CNN model picks up on subtleties that distinguish the vocal responses of the control group from those subjected to stressors. Over time, the control group's vocalizations appear to deviate from those of the treatment group, with the five-week control results showing more significant differences than the four-week results. These findings suggest that the vocal patterns of hens are sensitive to environmental conditions and stressors, and these changes are captured in the audio data over time.

Consistency of Vocal Changes Across Treatment Groups and Time

When examining vocal changes over time between 'Treatment 1' (umbrella stressor) and 'Treatment 2' (dog barking stressor), the classification model did not distinguish the patterns as distinctly as expected. However, the analysis shows that both treatments induce consistent changes in vocal patterns across different weeks. This consistency indicates that the stressors have a lasting effect on the hens' vocal behavior, which does not diminish over time as one might expect with habituation. These persistent vocal changes suggest that the stressors continue to impact the hens, an important consideration for welfare monitoring and the development of stress mitigation strategies in poultry farming.

Conclusions

Deciphering Stress-Induced Vocalizations in Poultry - Implications for Welfare Monitoring

The intricate vocalizations of laying hens serve as a communicative bridge, reflecting their internal states and welfare conditions. The comprehensive study conducted herein has systematically unraveled the vocal patterns elicited by environmental stressors, harnessing the analytical prowess of deep learning and the nuanced audio feature extraction capabilities of Mel Frequency Cepstral Coefficients (MFCCs). This fusion of advanced computational methods and acoustic analysis represents a significant stride forward in the endeavor to interpret the subtle lexicon of chicken vocalizations.

Our findings underscore the discernible impact of stressors on vocal expression, varying not only with the nature of the stressor but also with the age of the birds. Younger hens demonstrated vocal patterns markedly distinct from their older counterparts, pointing to a developmental trajectory in their stress response mechanisms. This age-related divergence in vocal behavior, captured through the lens of a Convolutional Neural Network (CNN), highlights the dynamic nature of animal communication and its sensitivity to physiological and environmental stimuli.

The study's revelations extend beyond the scientific understanding of avian communication. They pave the way for pragmatic applications in poultry farming, particularly in the realm of welfare monitoring. The precision with which the CNN model classified stress-specific vocalizations presents a compelling case for the integration of such models into non-invasive monitoring systems. Such systems could potentially detect early signs of distress, enabling timely interventions that mitigate stress and enhance the well-being of the flock.

The correlation between vocal changes and stressors, coupled with the consistent recognition accuracy across various ages and stress conditions, has far-reaching implications. It suggests that continuous monitoring of vocalizations could serve as an effective barometer for the welfare status of poultry. Furthermore, it underlines the importance of tailoring management practices to the nuanced needs of laying hens at different life stages, fostering an environment conducive to their health and productivity.

Our research also lays a foundation for future studies to explore the longitudinal effects of chronic stressors on vocalizations and whether hens exhibit habituation or heightened sensitivity over time. Such investigations could offer deeper insights into the resilience of poultry and the long-term implications of their rearing conditions.

In the broader scope of animal welfare science, this study illustrates the utility of bioacoustics and machine learning as powerful tools for advancing our comprehension of animal behavior. The methodology employed here can be extrapolated to other species and contexts, potentially revolutionizing welfare assessments across various domains of animal husbandry.

The integration of CNN models and MFCCs for analyzing chicken vocalizations has provided a novel perspective on the interplay between environmental stressors and animal communication. The implications of this research are manifold, offering a beacon for the development of enhanced welfare monitoring systems and informing best practices in poultry management. Ultimately, this study serves as an affirmation of the potential that lies in the intersection of

technology and animal welfare to foster practices that are not only productive but also ethically grounded and humane.

Funding
This research received no external funding.

Institutional Review Board Statement
The animal study protocol was approved by the Institutional Review Board (or Ethics Committee) of Wageningen University & Research.

Data Availability Statement
The data that support the findings of this study are available from the authors upon reasonable request.

Acknowledgments
This research was greatly enriched by data collected during a comprehensive collaborative project at the Animal Sciences Department, Wageningen University & Research. Profound appreciation is extended to the Adaptation Physiology Group, with special thanks to Naomi Pijpers, Huib van den Heuvel, and Ali Youssef, who worked diligently under the guidance of Dr. Suresh Neethirajan. Their meticulous efforts in data collection were pivotal to the success of this study. It is, however, crucial to acknowledge that while the data collection was a collaborative effort, the conceptualization, development of the algorithm, analysis of the data, and the drafting of this manuscript were independently executed by Dr. Neethirajan. His expertise and vision have been instrumental in the realization of this research, providing critical insights and direction throughout the process. The depth and breadth of the analysis presented in this study are a testament to his significant contributions to the field of poultry vocalization research.

Conflicts of Interest
The author declares no conflict of interest.

References

Abdel-Kafy, E.S.M.; Ibraheim, S.E.; Finzi, A.; Youssef, S.F.; Behiry, F.M.; Provolo, G. Sound Analysis to Predict the Growth of Turkeys. *Animals* **2020**, *10*(5), 866.

Adebayo, S.; Aworinde, H.O.; Akinwunmi, A.O.; Alabi, O.M.; Ayandiji, A.; Sakpere, A.B.; Adeyemo, A.; Oyebamiji, A.K.; Olaide, O.; Kizito, E. Enhancing Poultry Health Management Through Machine Learning-Based Analysis of Vocalization Signals Dataset. *Data Brief* **2023**, *50*, 109528.

Ahmed, G.; Malick, R.A.S.; Akhunzada, A.; Zahid, S.; Sagri, M.R.; Gani, A. An Approach Towards IoT-Based Predictive Service for Early Detection of Diseases in Poultry Chickens. *Sustainability* **2021**, *13*(23), 13396.

Banakar, A.; Sadeghi, M.; Shushtari, A. An Intelligent Device for Diagnosing Avian Diseases: Newcastle, Infectious Bronchitis, Avian Influenza. *Comput. Electron. Agric.* **2016**, *127*, 744–753.

Bessa Ferreira, V.H.; Dutour, M.; Oscarsson, R.; Gjøen, J.; Jensen, P. Effects of Domestication on Responses of Chickens and Red Junglefowl to Conspecific Calls: A Pilot Study. PLOS ONE 2022, 17(12), e0279553.

Briden, K.A.; Adams, N.J.; Cameron, K.E. Hierarchical structure and feeding behaviours of free-range hens (Gallus gallus domesticus). *Int. J. Comp. Psychol.* **2023**, *36*(1).

Carroll, B.T.; Anderson, D.V.; Daley, W.; Harbert, S.; Britton, D.F.; Jackwood, M.W. Detecting Symptoms of Diseases in Poultry Through Audio Signal Processing. In Proceedings of the 2014 IEEE Global Conference on Signal and Information Processing (GlobalSIP), Atlanta, GA, USA, December 2014; IEEE: **2014**; pp. 1132–1135.

Cuan, K.; Zhang, T.; Huang, J.; Fang, C.; Guan, Y. Detection of Avian Influenza-Infected Chickens Based on a Chicken Sound Convolutional Neural Network. *Comput. Electron. Agric.* **2020**, *178*, 105688.

Cuan, K.; Li, Z.; Zhang, T.; Qu, H. Gender Determination of Domestic Chicks Based on Vocalization Signals. *Comput. Electron. Agric.* **2022**, *199*, 107172.

Cuan, K.; Zhang, T.; Li, Z.; Huang, J.; Ding, Y.; Fang, C. Automatic Newcastle Disease Detection Using Sound Technology and Deep Learning Method. *Comput. Electron. Agric.* **2022**, *194*, 106740.

Du, X.; Lao, F.; Teng, G. A Sound Source Localisation Analytical Method for Monitoring the Abnormal Night Vocalisations of Poultry. *Sensors* **2018**, *18*(9), 2906.

Du, X.; Carpentier, L.; Teng, G.; Liu, M.; Wang, C.; Norton, T. Assessment of Laying Hens' Thermal Comfort Using Sound Technology. *Sensors* **2020**, *20*(2), 473.

Estevez, I. Understanding poultry social behaviour and its impact on animal welfare. In *Understanding the Behaviour and Improving the Welfare of Chickens*; **2020**; pp. 117–148, Burleigh Dodds Science Publishing.

Fontana, I.; Tullo, E.; Carpentier, L.; Berckmans, D.; Butterworth, A.; Vranken, E.; Norton, T.; Berckmans, D.; Guarino, M. Sound Analysis to Model Weight of Broiler Chickens. Poult. Sci. 2017, 96(11), 3938–3943.

Fontana, I.; Tullo, E.; Scrase, A.; Butterworth, A. Vocalisation Sound Pattern Identification in Young Broiler Chickens. Anim. 2016, 10(9), 1567–1574.

Ginovart-Panisello, G.J.; Alsina-Pagès, R.M.; Sanz, I.I.; Monjo, T.P.; Prat, M.C. Acoustic Description of the Soundscape of a Real-Life Intensive Farm and Its Impact on Animal Welfare: A Preliminary Analysis of Farm Sounds and Bird Vocalisations. Sensors 2020, 20(17), 4732.

Ginovart-Panisello, G.J.; Iriondo Sanz, I.; Panisello Monjo, T.; Riva, S.; Garriga Dicuzzo, T.; Abancens Escure, E.; Alsina-Pagès, R.M. Trend and Representativeness of Acoustic Features of Broiler Chicken Vocalisations Related to CO2. Appl. Sci. 2022, 12(20), 10480.

Herborn, K.A.; McElligott, A.G.; Mitchell, M.A.; Sandilands, V.; Bradshaw, B.; Asher, L. Spectral Entropy of Early-Life Distress Calls as an Iceberg Indicator of Chicken Welfare. J. R. Soc. Interface 2020, 17(167), 20200086.

Huang, J.; Wang, W.; Zhang, T. Method for Detecting Avian Influenza Disease of Chickens Based on Sound Analysis. Biosyst. Eng. 2019, 180, 16–24.

Huang, J.; Zhang, T.; Cuan, K.; Fang, C. An Intelligent Method for Detecting Poultry Eating Behaviour Based on Vocalization Signals. Comput. Electron. Agric. 2021, 180, 105884.

Kim, N.Y.; Jang, S.Y.; Kim, S.J.; Jeon, B.T.; Oh, M.R.; Kim, E.K.; Seong, H.J.; Tang, Y.J.; Yun, Y.S.; Moon, S.H. Behavioral and Vocal Characteristics of Laying Hens Under Different Housing and Feeding Conditions. J. Anim. Plant Sci. 2017, 27(1).

Mahdavian, A.; Minaei, S.; Yang, C.; Almasganj, F.; Rahimi, S.; Marchetto, P.M. Ability Evaluation of a Voice Activity Detection Algorithm in Bioacoustics: A Case Study on Poultry Calls. Comput. Electron. Agric. 2020, 168, 105100.

Mahdavian, A.; Minaei, S.; Marchetto, P.M.; Almasganj, F.; Rahimi, S.; Yang, C. Acoustic Features of Vocalization Signal in Poultry Health Monitoring. Appl. Acoust. 2021, 175, 107756.

Mao, A.; Giraudet, C.S.; Liu, K.; De Almeida Nolasco, I.; Xie, Z.; Xie, Z.; Gao, Y.; Theobald, J.; Bhatta, D.; Stewart, R.; McElligott, A.G. Automated Identification of Chicken Distress Vocalizations Using Deep Learning Models. J. R. Soc. Interface 2022, 19(191), 20210921.

Marx, G.; Leppelt, J.; Ellendorff, F. Vocalisation in Chicks (Gallus gallus dom.) during Stepwise Social Isolation. Appl. Anim. Behav. Sci. 2001, 75(1), 61–74.
Neethirajan, S. Automated Tracking Systems for the Assessment of Farmed Poultry. Animals 2022, 12(3), 232.

Pereira, E.M.; Naeaes, I.D.A.; Garcia, R.G. Vocalization of Broilers Can Be Used to Identify Their Sex and Genetic Strain. Eng. Agríc. 2015, 35, 192–196.

Sadeghi, M.; Banakar, A.; Khazaee, M.; Soleimani, M.R. An Intelligent Procedure for the Detection and Classification of Chickens Infected by Clostridium Perfringens Based on Their Vocalization. Braz. J. Poult. Sci. 2015, 17, 537–544.

Shawkat, S.S.; Mohammed, A.I.; Rashid, S.A.; Abdulateef, S.M. Effect of Some Unfavorable Behavioral Traits on the Behavior of Broiler Chicks. J. Life Sci. Appl. Res. 2023, 4(1), 1–8.

Tefera, M. Acoustic Signals in Domestic Chicken (Gallus gallus): A Tool for Teaching Veterinary Ethology and Implication for Language Learning. Ethiop. Vet. J. 2012, 16(2), 77–84.

Thomas, P.; Grzywalski, T.; Hou, Y.; De Carvalho, P.S.; De Gussem, M.; Antonissen, G.; Tuyttens, F.; De Poorter, E.; Botteldooren, P.D.D. Using a Neural Network Based Vocalization Detector for Broiler Welfare Monitoring. 2023.

van den Heuvel, H.; Youssef, A.; Grat, L.M.; Neethirajan, S. Quantifying the Effect of an Acute Stressor in Laying Hens Using Thermographic Imaging and Vocalisations. bioRxiv 2022, 2022-07.

Yang, X.; Zhao, Y.; Qi, H.; Tabler, G.T. Characterizing Sounds of Different Sources in a Commercial Broiler House. Animals 2021, 11(3), 916.

Zimmerman, P.H.; Koene, P.; van Hooff, J.A. The Vocal Expression of Feeding Motivation and Frustration in the Domestic Laying Hen, Gallus gallus domesticus. Appl. Anim. Behav. Sci. 2000, 69(4), 265–273.