SECE: accurate identification of spatial domain by incorporating global spatial proximity and local expression proximity

Yuanyuan Yu[1], Yao He[1*], Zhi Xie[1,2*]

[1] State Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-sen University, Guangzhou, China

[2] Center for Precision Medicine, Sun Yat-sen University, Guangzhou, China

[*] Email: Z.X: xiezhi@gmail.com; Y.H: scheyao@hotmail.com

Abstract

Identification of spatial domains is a critical step in spatial transcriptomics (ST) analysis. Here, we introduce SECE, a method that incorporates global spatial proximity and local expression proximity to identify spatial domains. SECE can also obtain spatial features for low-dimensional visualization and trajectory inference. In addition, SECE can identify cell types when dealing with ST data with single-cell resolution. Compared with eight spatial domain identification and five cell type identification methods, SECE consistently achieved top performance on spatial and cellular characterization, accurately providing biological insights into ST data.

Keywords

Spatial transcriptome, Spatial domain, Spatial embedding, Cell type

Background

Spatial transcriptomics (ST) allows for the acquisition of expression profiles with spatial information, providing insight into the cellular composition and function of cells in different locations within a tissue microenvironment. This technology has led to novel discoveries in various fields such as neuroscience [1], developmental biology [2], and cancer biology [3].

A variety of ST platforms have been developed with varying throughput and resolution. *In situ* hybridization-based ST platforms, including STARmap [4], seqFISH [5, 6], seqFISH+ [7], MERFISH [8] and FISSEQ [9], provide highly accurate gene expression measurement at single cell resolution but only for a limited number of known genes [10]. On the other hand, barcode-based ST platforms, such as Spatial transcriptomics [11] and its commercial version 10X Genomics Visium, Slide-seqV2 [12], HDST [13], Seq-Scope [14] and Stereo-seq [15], can perform high-throughput sequencing on a genome-wide scale. With recent advancements, barcode-based ST platforms have greatly improved their resolution. The early ST method, Spatial transcriptomics, has an initial resolution of 200 μm, while Visium improves that to 55 μm. Slide-seqV2 has a resolution of 10 μm, which is close to a single-cell level. Stereo-seq, Seq-Scope, and HDST have subcellular resolutions ranging from 0.5 to 2 μm, with Seq-Scope and Stereo-seq capable of merging spots based on cellular location to achieve single-cell resolution.

In ST data, spatial domains refer to spatially consistent regions in terms of gene expression and physical location, each with specific anatomical structures [1, 16]. To address the challenge of spatial region identification, several algorithms have been developed. For instance, BayesSpace [17] employs a Bayesian model with a Markov random field and encourages neighbor spots to belong to the same cluster. SpatialPCA [18] introduces a spatially aware dimension reduction method to extract a low-dimensional representation of the ST data while preserving the spatial correlation structure. Another algorithm, BASS [19], takes a coherent approach by jointly inferring labels for cell types and spatial regions based on a Bayesian hierarchical model. Moreover, graph neural networks have been widely used since the spatial positional relationship can be naturally transformed into a graph structure. SpaGCN [20] and STAGATE [21] employ graph convolutional neural networks and graph attention neural networks to aggregate neighbor

information respectively, while GraphST [22] , SpaceFlow [23] , and conST [24] use self-

supervised graph embedding learning strategies.

The way these approaches use similarity information between spots can be classified into

global or local similarity. Global similarity information measures pairwise spot similarity, yielding

high similarity for close spots and low similarity for distant ones, including SpaGCN and

SpatialPCA. Local similarity information refers to selecting neighboring spots based on

algorithms like K-nearest neighbors (KNN) and ensuring high consistency among them, including

STAGATE, BayesSpace, BASS, GraphST, SpaceFlow, and conST. However, the simultaneous

capture of local and global information which takes both advantages is lacking. In addition, some

existing methods have limited effectiveness in extracting gene expression features. They use

scaled values of highly variable genes (HVG) (e.g., GraphST, SpatialPCA, STAGATE,

SpaceFlow) or employ dimensionality reduction techniques like principal component analysis

(PCA) (e.g., BASS, SpaGCN, BayesSpace, conST) for expression features. However, these

features cannot handle expression with high noise, which is very common in barcode-based

sequencing ST methods, particularly in high-resolution techniques such as Stereo-seq [20].

To this end, we developed SECE, an accurate spatial domain identification method for ST

data. In contrast to the existing approaches, SECE incorporates global spatial proximity and local

expression proximity of data to derive spatial domains. The spatial embedding (SE) obtained by

SECE enables downstream analysis including low-dimensional visualization and trajectory

inference. To infer trajectories, we utilizeed Partition-based Graph Abstraction (PAGA) [25] at the

domain level and Monocle3 [26] at the single-cell level. Moreover, when applied to ST data with

single-cell resolution, SECE can accurately assign cell type labels by clustering cell type-related

embedding (CE). SECE was versatile on datasets from different ST platforms, for both high-

resolution platforms, such as STARmap, Slide-seqV2 and Stereo-seq, and relatively low-

resolution ones such as Visium (Additional file 1: Table S1). For ease of use, SECE was

implemented and integrated into the workflow of ST analysis using the Anndata data structure,

which can be used in the SCANPY package [27] for visualization and downstream analyses. The

capability of highly accurate identification of spatial domains and cell types enables SECE to

facilitate the discovery of biological insights from ST data.

Results

Overview of SECE

SECE can model ST data at various levels of resolutions, including subcellular, single-cell, near-single-cell, and multicellular (Fig. 1A). SECE obtains the result of spatial region division by clustering spatial domain-related embedding (SE). Specifically, SECE performs a two-stage procedure, consisting of an autoencoder (AE) module and a graph attention network (GAT) module. (Fig. 1B). The AE module is employed to accurately capture the CE of each spot by modeling the sequencing noise of expression profile (Methods). Then SE is achieved in GAT module by incorporating local expression proximity of neighbor spots and global spatial proximity of all spots. Expression proximity, which is based on CE, is measured using an attention mechanism, while spatial proximity, which is based on physical distance, is measured using a spatial similarity matrix (SSM). Finally, we obtain spatial domain, low-dimensional visualization, and spatial trajectory inference by running clustering, dimensionality reduction and trajectory inference algorithms on SE respectively (Fig. 1C). In addition, clustering of CE results in the identification of cell types.

To assess the contributions of CE learning and global spatial proximity in learning SE, we tested three alternatives to SECE. "SECE-PCA" eliminated the influence of CE by substituting it with PCA of equal dimension. "SECE-NG" removed the impact of SSM by setting the weight of the global spatial similarity constraint λ_{global} to 0. "SECE-PCA-NG" used PCA and set λ_{global} to 0. We tested these methods on a mouse hippocampus dataset generated by Slide-seqV2 [12], a technology that provided transcriptome-wide RNA detection with a near-cellular resolution of 10 μm. This dataset included 53,208 cells and 23,264 genes from the hippocampus, cortex, and thalamus, with a high sparsity of 98.19%.

Compared to SECE, three SECE-derived methods had reduced capability in spatial domain identification or trajectory inference. SECE-PCA-NG could not detect differences in the cortical region, while SECE-PCA detected differences but did not correctly identify near-parallel structures (Fig. 1D, left). This hinted at the importance of accurately modeling gene expression using CE. SECE-NG achieved the most similar results to SECE but it could not divide the Cornu Ammonis (CA) correctly, ignoring the region CA2. In addition, it divided Layer 5 into two sub-

layers with less smoothness. Although SECE-NG recognized more of the yellow area Dentate

gyrus (DG)-molecular layer (mo) around DG compared to SECE at n=14, we observed that as the

number of clusters increased, SECE was also able to identify the subtle CA1-(stratum oriens)

so/(stratum radiatum) sr in the CA1 region. For instance, at n=15, SECE detected a purple region,

and at n=16, it detected a brown region (Additional file 1: Fig. S1, 2). In contrast, increasing the

number of categories did not lead to improved recognition of SECE-NG in the CA area or Layer

5. Spatial trajectory inference of cortex using SE generated from SECE-derived methods also

showed that a spuriously high pseudo-time of deep spots in SECE-PCA-NG (Fig. 1D, right).

These results showed that both CE and SSM are important to learn accurate SE.

Application to STARmap data

We tested SECE on the mouse visual cortex data generated by STARmap [4]. STARmap is a *in*

situ sequencing-based ST method with single-molecule resolution. Despite its low gene

throughput (160 to 1020 genes), it offers high sensitivity at single cell resolution with high

efficiency and reproducibility [4]. This dataset has a total of 1,207 cells, 1,020 genes, and a

sparsity of 76.88%. The cells were annotated into distinct layers, including L1, L2/3, L4, L5, L6,

as well as the corpus callosum (CC) and hippocampus (HPC) of the visual cortex (Fig. 2A).

First, we compared the spatial region identification results of SECE with eight existing

methods, including BASS, SpaceFlow, GraphST, STAGATE, SpatialPCA, SpaGCN, BayesSpace,

and conST (Fig. 2B left, C). We assessed the consistency of each algorithm's spatial regions with

the ground truth layers using Adjusted Rand Index (ARI), Normalized Mutual Information (NMI),

and Kuhn-Munkres [28] based Accuracy (ACC). We found that SECE achieved the highest

agreement with the ground truth (Fig. 2D). It was closely followed by BASS. However, BASS

incorrectly combined HPC and L5. The accuracy of other algorithms was relatively low, and they

exhibited weak spatial aggregation, as indicated by the Local inverse Simpson's index (LISI)

(Additional file 1: Fig. S3A). Additionally, these methods mistakenly classified aggregated

endothelial cells (Endo) in L1 and L2/3 into the same region, such as region 2 in SpaceFlow,

region 3 in STAGATE and BayesSpace, region 5 in GraphST and SpatialPCA, and region 6 in

SpaGCN.

Next, we compared SE of SECE with SpaceFlow, GraphST, STAGATE, and SpatialPCA.

Due to the difficulty of conST in generating clear spatial regions, its SE learning comparison was

excluded. Our results demonstrated that SE of SECE explained the ground truth layers most

effectively (ASW=0.16), followed by SpaceFlow (ASW=0.12), GraphST (ASW=0.07),

STAGATE (ASW=0.07), and SpatialPCA (ASW=0.05) (Fig. 2E). SECE had a clearer and

continuous pattern in UMAP visualization compared to the other methods (Fig. 2F). When

comparing trajectory inference results, we selected the cortex part in ground truth, namely L1,

L2/3, L4, L5 and L6. The PAGA analysis based on SECE revealed a linear and continuous

relationship between these layers (Fig. 2G). Conversely, SpaceFlow mistakenly made the

connection between L1 versus L4 and L5, while the patterns in the other three methods were more

unclear. For individual cells, SECE exhibited a sequential increase of pseudo-time from L6 to L1

(Additional file 1: Fig. S3B, C), and the similar trend was also observed in SpaceFlow, while

GraphST, STAGATE, and SpatialPCA ordered L1 and L2/3 incorrectly (Additional file 1: Fig.

S3D, E).

Finally, we examined the results of cell type identification. We compared SECE with Seurat

[29], SC3 [30], CIDR [31], BASS, and SpiceMix [32]. First, we compared the identified cell types

with the annotations provided in the original research [4]. BASS (ARI=0.51), SECE (ARI=0.49)

and SC3 (ARI=0.49) had the top three performances (Additional file 1: Fig. S4A), followed by

SpiceMix (ARI=0.36) and Seurat (ARI=0.41), while CIDR was lowest (ARI=0.09). Specifically,

SECE identified seven excitatory neuronal (EX-Neuron) subtypes, two inhibitory neuronal (IN-

Neuron) subtypes, and seven non-neuronal cell types (Fig. 2B, right; Additional file 1: Fig. S4B,

S5). Among them, the seven EX-Neuron subtypes identified were all distributed in the

corresponding cortex (Additional file 1: Fig. S6A). Interestingly, both astrocytes (Astro) and

oligodendrocytes (Oligo) were divided into two subtypes, one concentrated in the CC (Astro-1,

Oligo-2) and one diffused in the HPC and cortical layers (Astro-2, Oligo-1). Additionally, SECE

was the only method that could identify subtypes of Endo with one subtype (Endo-1) concentrated

between L1 and L2/3 and the other (Endo-2) scattered elsewhere. Endo-1 and Endo-2 had

different highly expressed genes, suggesting that they might also have different functions

(Additional file 1: Fig. S6B). In contrast, SpiceMix failed to recognize EX L5, Seurat recognized a

mixed neuronal cell class, while SpiceMix, Seurat, and SC3 all failed to recognize subtypes of IN-

Neuron (Additional file 1: Fig. S4B, C; Fig. S5). BASS could identify the IN-Neuron subtype VIP

not detected by other methods. However, it assigned Astro and EX L4 into the same cluster, and it

detected a class (NA) with no marker genes. CIDR could only clearly identify microglia (Micro),

SST, Smc and two Oligo subtypes.

To sum, using a dataset generated by STARmap, we showed that SECE accurately identified

spatial domains and cell types. SECE also improved the performance in SE visualization and

spatial trajectory inference compared to the previous methods.

Application to Slide-seqV2 data

Next, we tested SECE on the hippocampal dataset generated by Slide-seqV2 platform. SECE

identified 14 distinct regions, and we annotated them according to the known structure of Allen

Brain Atlas (ABA) (Fig. 3A). These regions included the hippocampus (CA1, CA2, CA3, DG, and

CA slm/so/sr), cortex (Layers 4, 5a, 5b, 6), third ventricle, CC, and three subregions of the

thalamus (Fig. 3B, left). The identified regions were supported by their marker genes. For

example, the four important components of the hippocampal region, CA1, CA2, CA3, and DG,

were clearly demarcated, as evidenced by the high expression of their known markers *Wsf1*,

Rgs14, *Nptxr* and *C1ql2* [33] (Fig. 3C, Additional file 1: Fig. S7A). Layer 5 in cortical regions

was identified as two sublayers, Layer 5a and 5b, with different gene expression levels (Additional

file 1: Fig. S7). For comparison, we evaluated the performance of eight existing methods for

spatial region identification (Fig. 3D). Except for GraphST, none of these methods could identify

the CA2 region. Furthermore, for cortical areas, BASS failed to distinguish between Layer 4 and

Layer 5, SpaceFlow mixed Layer 5 and Layer 6, SpatialPCA exhibited suboptimal division

smoothness, and the remaining algorithms were unable to generate clear cortical regions.

Additionally, SECE exhibited the highest spatial aggregation performance as it had the smallest

LISI values (Additional file 1: Fig. S8A).

Then, we compared the performance of low-dimensional embeddings. UMAP generated from

SECE clearly displayed clustering patterns for the hippocampus, cortex, thalamus, third ventricle,

and their subregions (Fig. 3B, right). In addition, the sublayers of them like Layer4 and Layer6b

were arranged in a sequence, while UMAP of STAGATE and GraphST missed them (Additional

file 1: Fig. S8B). Moreover, we conducted spatial trajectory inference for the cortex due to its

continuous relationship between subregions. We selected the regions corresponding to the cortex

in each method, starting with the deepest clusters, such as Layer 6 for SECE, domain 1 for

SpaceFlow, domain 2 for GraphST, and domain 3 for SpatialPCA and STAGATE (Fig. 3E, F).

For SECE, the pseudo-time consistently increased with decreasing cortical depth, clearly

capturing the pseudo-time relationship between different layers. In contrast, SpaceFlow exhibited

less distinguishable differences between the learned layers. SpatialPCA reversed the relationship

between Layer 5b and Layer 6, while GraphST and STAGATE failed to identify the relationship

within the cortical region.

Next, we tested cell type identification and CIDR was excluded from the test due to its huge

memory consumption. SECE identified eleven subtypes of EX-Neuron, one IN-Neuron, and eight

subtypes of non-neuronal cells (Additional file 1: Fig. S9 A to C). Each cell type exhibited a

unique set of differentially expressed genes (Additional file 1: Fig. S10). Specifically, the EX-

Neurons in CA comprised of EX CA1, EX CA2, and EX CA3, while the non-neuronal subtypes

included Astro, Micro, two types of Oligos, polydendrocyte (Poly), mural cells (Mural),

ependymal cells (Epen), and choroid plexus cells (Chor). In contrast, none of the existing methods

were able to identify EX CA2, Poly, or Micro. Furthermore, BASS assigned Astro and the EX-

Neuron in the same class and recognized a cluster labeled NA without specific marker genes.

Seurat could not identify IN Neuron and misclassified EX-Neuron in CA into two subtypes with

less distinct, namely cluster 12 and 15. SC3 and SpiceMix failed to detect differences among EX-

Neuron in the hippocampal region, assigning both EX CA and EX DG to a single broad class,

namely cluster16.

Finally, our study revealed a compelling concordance between the spatial regions and cell

types identified by SECE (Additional file 1: Fig. S9D, E; Fig. S11). Each subregion of the

hippocampus, cortex, and thalamus corresponded to specific EX-Neuron subtypes. For instance,

EX CA1, EX CA2, EX CA3, and DG were predominantly localized within their respective

regions of CA1, CA2, CA3, and DG. Moreover, we observed that Layer 5a and Layer 5b had

differential distributions of EX-Neuron subtypes, and Layer 5b exhibited a higher proportion of IN

neuron, which further confirmed the existence of Layer 5 sublayer. Additionally, distinct spatial

distributions were evident among subtypes of cell types. For instance, Oligo-1 exhibited a

predominant presence in the CC, whereas Oligo-2 was primarily concentrated in the thalamus. In

contrast, the subtypes of Oligo identified by BASS did not have this spatial pattern (Additional file

1: Fig. S9F).

In conclusion, using a sparse dataset generated by Slide-seqV2, SECE outperformed the other

methods in various tasks. We identified functional subregions of the hippocampus, cerebral

cortex, and thalamus, particularly the CA2 region, and correctly reproduced the developmental

trajectory of the cerebral cortex. The cohesive correlation between spatial regions and cell types

further supported the discovery of cell subtypes and spatial subregions.

Application to Stereo-seq data

Stereo-seq is an emerging technique for ST with genome-wide throughput and subcellular

resolution [15]. It obtains the expression profile and spatial coordination of each DNA nanoball

(DNB), followed by image-based cell segmentation to identify DNBs belong to the same cell.

Then, an expression matrix with single-cell resolution is obtained by summing the expression

values within the same cell. However, the high noise levels associate with high resolution prevent

accurate identification of cell types and spatial domains at the single-cell level. Here, we analyzed

the mouse olfactory bulb data generated by Stereo-seq [34], which contained 19,527 cells and

27,106 genes, with 98.69% of the values being zero.

First, we assessed the effectiveness of cell type recognition. SECE identified 6 types of

neurons and 4 types of non-neurons (Fig. 4A; Additional file 1: Fig. S12; Fig. S13). Neurons were

three types of Granule cells (GC), two types of Mitral and tufted cells (M/TC), and Periglomerular

cells (PGC), while non-neurons included Astro, Oligo, and two types of Olfactory ensheathing

cells (OEC). In contrast, BASS produced an NA category without any marker genes. SpiceMix

and Seurat did not identify Astro and Oligo, respectively, while SC3 failed to obtain Astro, Oligo

and PGC.

Next, we segmented this data into spatial regions using SECE and annotated the results based

on the known olfactory bulb layers (Fig. 4B, left). They were Rostral migratory stream (RMS),

Granule cell layer (GCL), Inner plexiform layer (IPL), Mitral cell layer (MCL), External

plexiform layer (EPL), Glomerular layer (GL) and Olfactory nerve layer (ONL) from inside to

outside (Fig. 4B, right; Additional file 1: Fig. S14A). Our findings were validated using known

markers [33] for each layer (Additional file 1: Fig. S14B). Notably, besides the known seven

layers, we made a finer division of GCL and ONL, and the sublayers were named GCL-Inner,

GCL-Outer, ONL-Inner and ONL-Outer respectively. Several evidences confirmed these

sublayers. First, we found that marker gene expression differed between them, where the GCL-

inner highly expressed *Nrgn* and the GCL-outer highly expressed *Pcp4*. The expression level of

markers was also higher in the ONL-outer than in the ONL-inner. Then, we calculated the

relationship between layers and the corresponding cell types (Additional file 1: Fig. S14C). We

found that RMS contained Oligo and a small amount of GC; EPL mainly contained M/TC-2 and

Astro; MCL and GL mainly contained M/TC-1 and PGC, respectively, which were consistent with

known biological knowledge [35, 36]. Most of all, GCL-Outer was almost composed of GC, while

GCL-Inner contained a certain amount of Oligo, and the GC subtypes in GCL-Inner and GCL-

Outer also differed. Moreover, the ONL-Outer contained almost exclusively OEC cells, while the

ONL-Inner contained a fraction of OEC and more Astro. This supported previous findings that the

ONL, as a part of the olfactory bulb blood-brain barrier, had fine internal and external subregions

with different cell types [37] . Our study provided further support for this finding at the spatially

resolved single-cell level.

We further compared the performance of SECE to the existing methods (Fig. 4C). BASS

failed to separate the RMS from the GCL layer and could not distinguish GL from EPL. It also

identified some irrelevant regions with few cells, such as domain5, 8, and 9. STAGATE and

SpaGCN mixed GL and EPL. SpaceFlow, GraphST, and SpatialPCA encountered challenges in

accurately dividing the GCL layer. BayesSpace and conST did not yield a clear domain

identification. Additionally, BASS had the most spatial aggregation performance with lowest LISI

value, while SECE ranked second (Additional file 1: Fig. S14D). Moreover, SECE showed

robustness when tested with different number of clusters (7, 8, and 10), consistently providing

well-bounded ring stratification (Additional file 1: Fig. S15).

Finally, we assessed the SE learning capabilities of SECE, SpaceFlow, GraphST, STAGATE,

and SpatialPCA. The low-dimensional UMAP visualization based on SECE exhibited a

continuous low-dimensional pattern with nine layers arranged in order of spatial position from

inside to outside (Fig. 4D). Moreover, the trajectory inference for these layers demonstrated

approximately linear relationship (Fig. 4E). SpatialPCA also displayed linear patterns, except for

regions 6 and 7. However, the trajectory inference results of SpaceFlow, GraphST, and

STAGATE were challenging to interpret.

Collectively, SECE accurately identified cell types and spatial domains in the mouse

olfactory bulb. Furthermore, SECE provides a higher resolution in spatial stratification compared

to the existing methods.

Application to complex hemibrain Stereo-seq data

We further tested SECE to a dataset with a more complex anatomic structure generated by Stereo-

seq [15]. This dataset was a coronal slice of the mouse hemibrain, containing 50,140 cells and

25,879 genes, of which 96.94% had a value of 0. The original study annotated cell types and

anatomic regions using single cell and bin50 data, respectively (Fig. 2 in [15]). Bin50 data binned

the neighboring 50×50 DNBs with fixed area, but it did not match the cell type label. Here, single-

cell data was utilized for both cell type and spatial region identification, although its higher

sparsity increased the difficulty.

Clustering of SE resulted in spatial domain annotations that were in highly consistent with

ABA anatomy (Fig. 5A, B, left), with clear separations between regions including cortex,

hippocampus, midbrain, thalamus, and fiber tracts (FT). The cortex included five layers: L1, L2/3,

L4, L5, L6, LVC and CAA. In contrast, other methods failed to accurately identify these cortical

layers (Fig. 5C). Specifically, all of them encountered difficulties in identifying Cortex L5 with

high smoothness. In addition, LISI showed that spatial domains identified by SECE had the

strongest spatial pattern (Fig. 5D). Furthermore, we performed a low-dimensional visualization of

SE using t-SNE [38], which showed the effect of aggregation in the same region and separation in

different regions (Fig. 5B, right; Additional file 1: Fig. S16).

Then, we tested the ability of cell type recognition. Using the SECE, we identified 15 types

of EX-Neuron, two types of IN-Neuron, dopaminergic neuron (DA-Neuron), and granule cell of

dentate gyrus (GN-DG). Among the non-neuronal cell types, we distinguished four subtypes of

Astro, three subtypes of Oligo, Micro, erythrocytes (Ery), and meninges (Fig. 5E). However, other

algorithms yielded mixed results (Additional file 1: Fig. S17, 18). BASS, for instance, identified

two hybrid classes that grouped Astro and Ery together instead of identifying them separately. It

also failed to delineate meninges and EX-Thalamus, while Seurat and SpiceMix mixed meninges

with Ery and Astro, respectively. Seurat and SpiceMix also identified a mixture of two EX CA1

and EX CA3 classes, rather than detecting them separately. Additionally, Seurat encountered

challenges in accurately separating neurons in specific regions of the cortex, leading to the

identification of three EX L5/6 classes and two EX L2/3/4 classes. SpiceMix did not identify IN-

Neurons in the telencephalon but mixed them with three EX-Neuron subtypes. Moreover, it did

not recognize the DA-Neuron. The performance of the SC3 algorithm was relatively weaker,

possibly due to the large volume and high sparsity of the data. Classes 16 and 27 contained more

than one-third of the cells, but no distinct pattern of high expression was observed. Furthermore,

none of these algorithms were able to identify Micro accurately. These findings were further

supported by analyzing differentially expressed genes and assessing spatial distribution

(Additional file 1: Fig. S19 to S24).

Finally, we verified that the spatial domains and cell types identified by SECE were

accurately matched (Additional file 1: Fig. S25, 26). The GN-DG, EX CA1, EX CA3, and EX

Subiculum cell types primarily constituted the DG, CA1, CA3, and Subiculum regions,

respectively. Cortex L2 to L6 mainly consisted of their corresponding EX-Neuron subtypes, while

the Cortex L1 contained a significant number of Astro compared to other cortical layers.

Additionally, the substantia nigra/ventral tegmental area (SN/VTA) region was predominantly

populated by DA-Neuron. Furthermore, we observed distinct spatial distribution differences

among the four Astro subtypes and three Oligo subtypes. Astro-1 was predominantly distributed in

the midbrain, while Astro-2 and Astro-4 were primarily found in the telencephalon region. Astro-3

exhibited a concentrated presence in Cortex L1, adjacent to meningeal cells. As for Oligo, Oligo-1

and Oligo-2 were distributed in FT and midbrain, respectively, while Oligo-3 was found in both

regions.

In conclusion, the Stereo-seq dataset with complex spatial structure and high sparsity was

effectively annotated by SECE, which accurately matched spatial domains and cell types for each

cell and provided robust support for cell subtypes from spatial distribution disparities.

Application to Visium data

Finally, we tested the applicability of SECE on Visium, which had a relatively low resolution. Visium sequences a fixed number of 55 μm spots, with 1-10 cells per spot, depending on the tissue type [39]. We first applied SECE to a breast cancer dataset, containing 3,798 spots and 24,923 genes, of which only 77.44% are zero values. Based on pathological images, the dataset covered four phenotypic regions, including ductal carcinoma in situ/lobular carcinoma in situ (DCIS/LCIS), healthy tissue (Healthy), invasive ductal carcinoma (IDC), and tumor surrounding regions with low features of malignancy (Tumor edge), as well as 20 annotated subdivisions [34] (Fig. 6A; Additional file 1: Fig. S27A).

First, we segmented 20 regions using each algorithm and performed phenotype annotations, which were then compared with image-based manual annotations (Fig. 6B; Additional file 1: Fig. S27B, C). Notably, in the context of a dataset characterized by a low missing rate, all algorithms demonstrated commendable performance. Most of the algorithms had ARI values ranging from 0.55 to 0.62, NMI values ranging from 0.5 to 0.6, and ACC values around 0.8 (Additional file 1: Fig. S27D). Among these nine algorithms, SECE's ARI ranked fifth, NMI and ACC ranked third (ARI=0.60, NMI=0.55, ACC=0.82). BayesSpace and BASS, which shared the top performance, attained ARI values of 0.62, NMI values of 0.56, and ACC values of 0.83.

We found that SECE divided individual tumor regions into wrapped layers more clearly. Specifically, cluster 20 and cluster 15 divided the IDC-5 into internal and external layers, while cluster 11 and cluster 10 split the DCIS/LCIS-1. (Fig. 6B; Additional file 1: Fig. S27A). To investigate the biological significance of the refined stratification. We named clusters 20,15,11 and 10 as IDC-inner, IDC-outer, DCIS/LCIS-inner and DCIS/LCIS-outer, respectively. We further focused on the parts of clusters 4 and 5 that were located near the tumor edge, referred to as IDC-edge and DCIS/LCIS-edge. (Fig. 6C). We analyzed the cell type composition of each spot of these regions by integrating the annotated breast cancer scRNA-seq data [40] and deconvoluting each spot using cell2location [41] (Additional file 1: Fig. S28, S29). Our findings showed that the proportion of tumor cells gradually decreased as the tumor moves towards the edge, while the that of immune cells and stromal cells gradually increased (Fig. 6D), confirming the correctness of subregions.

We further explored the characteristics of these subregions (-edge, -outer and -inner) of

DCIS/LCIS and IDC (Fig. 6E). Differential expression analysis showed that DCIS/LCIS-edge had

a high level of humoral immunity, which could be confirmed by the enrichment of B cell receptor

signaling pathway related genes (*IGLC2*, *IGLC3*, *IGHG1*, *IGHG3*, *IGHA1*). DCIS/LCIS-outer

overexpressed *KRT14* of the keratin family, which was a key regulator of metastasis, suggesting

invasive potential [42, 43]. In contrast, DCIS/LCIS-inner showed non-metastatic traits but also

had tumor-promoting abilities, indicated by a high expression of *LDHA*. *LDHA*, linked to

metastasis-free survival, catalyzes the conversion of pyruvate to lactate, creating a hypoxic

environment in the tumor core, and inducing metabolic changes in both tumor cells and

fibroblasts, promoting tumor progression [44].

For the IDC subregions, IDC-edge showed elevated expression of biomarkers linked to tumor

proliferation, invasion, and migration including tumor-associated macrophage (*APOE*),

complement components (*C1qA*, *C1qB*), cathepsin (*CTSD*), and apolipoprotein (*APOC1*) [45-49].

IDC-outer had a high level of immunity and some transferability, which was derived from the high

expression of MHC class I-related genes (*HLA-A*, *HLA-B*, *B2M*) [50, 51]. Besides, there were

increased expression of genes such as *MGST1* and *MRPS30-DT*, which were known to promote

breast carcinoma cell growth and metastasis [52, 53]. In IDC-inner, there were higher levels of

tumor activity and lower levels of immune response. *LINC00052*, known to promote breast cancer

cell proliferation by increasing signals of epidermal growth factor receptor (EGFR) such as *HER3*

[54-56], was overexpressed. Upregulation of *COX6C* and *FAM234B* implied higher levels of

cellular respiration [57] and lower immune response function [58], respectively.

Next, we inferred developmental trajectories for the IDC and DCIS/LCIS regions. The

starting points for calculating pseudo-time in IDC and DCIS/LCIS were the interior of the tumor

regions (Fig. 6F). There were increased pseudo-time values as the region of the tumor moves

outward, effectively mimicking the gradual progression of tumor development.

Moreover, to further evaluate the power of SECE in spatial domain identification, we tested

12 dorsolateral prefrontal cortex (DLPFC) data generated from the Visium platform. The original

study [1] had manually annotated the spatial regions of these datasets, encompassing white matter

and six cortical layers (Additional file 1: Fig S30, S31). Among the slices, eight contained all

seven regions, while the remaining four slices contained five of these regions. In our evaluation,

we ensured that the number of regions per slice remained consistent with the annotated regions.

We observed that SECE exhibited the highest agreement with the original annotations. The

median ARI for SECE was 0.58, surpassing the second and third ranked algorithms, STAGATE

and SpatialPCA, which achieved median ARI values of 0.55 and 0.54, respectively. These

findings highlighted the superior performance of SECE in accurately delineating the spatial

regions within the low-resolution data.

In summary, SECE divided tumors into finer regions compared to manual annotation based

on pathological images in the Visium breast cancer dataset, providing a deeper understanding of

the heterogeneity within a single tumor. Meanwhile, SE was able to reproduce the dynamic

process of tumor development.

Discussion

SECE was designed to accurately identify spatial domains of ST data by effectively learning low-

dimensional spatial features. It only uses ST data as input without additional single cell or

histology data and achieved superior performances on ST data generated from different platforms.

It effectively balanced expression and spatial approximation to learn a highly representative SE,

and models noise explicitly to obtain an accurate CE. SECE had excellent performance for spatial

domain identification, low-dimensional visualization, and trajectory inference. Its CE component

could accurately label cell types, even in sparse Stereo-seq datasets. To further illustrate the ability

of SECE to learn cell type features, we compared the feature learning capabilities of the AE

module in SECE with scVI on real and simulated single-cell data (details in the Additional file 1:

Supplementary note1, Fig S32, S33), showing the effectiveness of SECE for single-cell datasets.

SECE was primarily designed for high-throughput and high-resolution ST data. When

dealing with higher resolution datasets, such as Slide-seqV2 and Stereo-seq, which often

contained tens of thousands of cells with higher sparsity and noise, the identification of spatial

regions became more challenging (Additional file 1: Fig S34A). In such cases, SECE provided

more additional improvements that significantly enhance the analysis of these datasets. SECE also

achieved excellent performance on low-throughput data, such as STARmap, and coarse-resolution

data, such as Visium.

With the increasing captured area of ST data and advancements in resolution, there was a

growing demand for computational methods to exhibit higher efficiency and scalability. We

recorded the runtime and GPU memory consumption for each dataset (Additional file 1: Fig

S34B, C). For Slide-seqV2 hippocampus data and the Stereo-seq hemibrain data, which contained

over 50,000 cells, SECE achieved a running time of less than 4.2 minutes while utilizing less than

5GB of GPU memory. These results demonstrated the superior computational efficiency and

scalability of SECE when dealing with large-scale datasets.

For these emerging high-resolution sequencing methods such as Stereo-seq, our exploration

of spatial identification and cell type is just beginning. More algorithms are needed to uncover

expression patterns as well as cell-cell communication. Additionally, methods for the integration

of ST and the other types of datasets are also necessary. Ultimately, continuous improvements in

ST resolution and analysis tools will provide deep biological insights and facilitate both basic and

medical research.

Conclusion

447   In conclusion, our results demonstrated that SECE achieved great spatial domain identification,

448   particularly for ST datasets with high resolution. It outputs spatial embedding which can be used

449   for low dimensional visualization and other downstream analysis such as trajectory inference.

450   Additionally, for data with single-cell resolution, accurate cell types can also be identified.

Methods

Data preprocessing

Assuming that the spatial transcriptome data expression profile contains N spots (It means that

there are N cells for data with single-cell resolution, here we collectively refer to spots for the

sake of convenience.) and G genes, the dimensions of spatial gene expression X and spatial

coordinates Y are $N \times G$ and $N \times 2$ respectively. The raw gene expression counts X are

normalized based on library size and then transformed to a natural logarithmic scale to obtain the

normalized expression matrix $\tilde{X}$, which was used as input of SECE together with the coordinate

matrix Y.

Learning CE with AE module

463 We learn cell type-related embedding (CE) based on gene expression data. We assume that each

464 element in expression count matrix follows a zero-inflated negative binomial (ZINB) distribution

465 [59, 60]. For x_{ng}, that is the expression count of gene g at spot n, we have

$$ZINB(x_{ng}; r_{ng}, p_g) = \pi_{ng}\delta_0(x_{ng}) + (1 - \pi_{ng})\frac{\Gamma(x_{ng} + r_{ng})}{x_{ng}!\,\Gamma(r_{ng})}p_g^{\,r_{ng}}(1 - p_g)^{x_{ng}},$$

where δ_0 is Dirac delta function, π_{ng} is the probability that the true gene expression value

is observed to be 0, (r_{ng}, p_g) are the standard parametrization for the negative binomial

distribution, x_{ng} is the number of successes of independently repeated Bernoulli trials before

reaching r_{ng} failures, and p_g is the probability of failure for each Bernoulli trial.

We use the auto-encoder in deep learning to learn CE and distribution parameters at the same

time. The ZINB parameters are the $N \times G$-dimensional parameter matrix Π, R and the G-

dimensional parameter vector P, which consist of the elements π_{ng}, r_{ng} and p_g separately.

The relationships between auto-encoder and ZINB parameters are $r_{ng} = s'_n r'_{ng}$ and $p_g =$

$\frac{\exp(l_g)}{1+\exp(l_g)}$, where s'_n is the scaling factor for spot n, l_g is the logits of p_g. Here, we draw the

assumption in SCT [61], that is, there is a linear relationship between the logarithm of gene

expression level and the logarithm of the base 10 of sequencing depth, $s'_n = \exp(log_{10}s_n)$, where

s_n is the sum of UMI in cell n, so the reconstructed expression is $\hat{x}_{ng} = r_{ng} * \exp(l_g)$.

The formulation of the autoencoder is given below:

$$Z = f_1(\tilde{X})$$

$$Z' = f_2(Z)$$

$$R' = f_3(Z')$$

$$\Pi = f_4(Z'),$$

f_1 is a two-layer neural network with nonlinear activation function ReLU, it takes $\tilde{X}$ as

input, and outputs Z, which has dimension $N \times m_1$ and is used as CE. f_2 is a one-layer neural

network with nonlinear activation function ReLU. f_3 is a one-layer neural network with

exponential functions, it outputs R' consisting of elements r'_{ng}. f_4 is a one-layer neural network

with activation function sigmoid. It outputs Π, a $N \times G$- dimensional matrix consists of π_{ng},

which is the zero-inflated probability of gene g at spot n. In addition, the parameter vector L

consisting of elements l_g is obtained from a learnable vector with exponential transformation.

The loss function of this module is the negative likelihood of the ZINB distribution:

492
$$Loss_1 = NLL_{ZINB}(\mathrm{X}; \Pi, \mathrm{R}, \mathrm{P}).$$

We note that zero-inflation term in the ZINB model allows for better modeling of dropout

event. For datasets with lower sparsity, such as STARmap and Visium, negative binomial (NB) is

sufficient for modeling. Both models are implemented in SECE. We used ZINB for Stereo-seq

and Slide-seq which have high degree of sparseness and NB for STARmap and Visium whose

proportion of 0 values are relatively low.

Learning SE with GAT module

After obtaining CE of individual spots using the expression information, we can fuse the spatial

coordinate information to learn the SE. We construct the ADM and SSM from spatial location,

followed by learning SE with two Graph Attention (GAT) layers.

Construction of ADM

To learn a smooth and reliable SE, we need to construct an adjacency matrix A based on the

coordinate position Y of spots, so that each spot can aggregate the features of neighbor spots. A

is an $n \times n$-dimensional symmetric matrix with elements taking the value 0 or 1. Let the set of

spots written by $V = \{v_1, v_2, \dots, v_N\}$, then for each element a_{ij} of the adjacency matrix A,

509
$$a_{ij} = \begin{cases} 0, & v_i \in \mathcal{N}(v_j) \\ 1, & v_i \notin \mathcal{N}(v_j) \end{cases}.$$

$\mathcal{N}(v_j)$ is the set of v_j-neighbor spots, which can be calculated by KNN or distance cutoff. We

use KNN to calculate neighbors by default, and the default number of neighbors is 6 for Visium

and 10 for other data.

Construction of SSM

We hypothesize that the correlation of SE between cells decreases with increasing distance and

tends to decay exponentially. So ideally Σ is the $n \times n$-dimensional spatial correlation matrix

based on a Gaussian kernel,

$$\Sigma_{ij} = \exp\left(-\frac{|y_i - y_j|^2}{\gamma}\right),$$

where the bandwidth parameter γ controls the degree of influence of spatial location on features,

with larger γ having a greater influence. We set γ as the quantile value of the distance matrix,

the default is 0.05 quantile.

GAT Layer

Let the input feature matrix of one GAT layer is $H = (h_1, h_2, \ldots, h_N)$, which has $N \times m$

dimensions with N samples and m features. The output of GAT layer is $H' = (h'_1, h'_2, \ldots, h'_N)$

with $N \times m'$ dimensions. h'_j can be formulated as

$$h'_j = \sigma\left(\sum_{v_i \in \mathcal{N}(v_j)} \alpha_{ij} W h_i\right),$$

where W is the weight matrix with $m' \times m$ dimensions, $\mathcal{N}(v_j)$ is the set of v_j-neighbor spots,

α_{ij} is the normalized attention coefficient matrix using the SoftMax function:

$$\alpha_{ij} = softmax_j(e_{ij}) = \frac{\exp(e_{ij})}{\sum_{k \in \mathcal{N}_i} \exp(e_{ik})},$$

$e_{ij} = a^T(Wh_i \parallel Wh_j)$ where a is learnable vector and $\parallel$ is the concatenation operation. We

used Exponential Linear Unit (ELU) as activation function σ in GAT layers.

GAT module

GAT module consists of two GAT layers. It takes CE features Z and spatial coordinates Y as

input, and outputs SE features U, which is a $n \times m_2$-dimensional matrix.

$$U = GAT_2(GAT_1(Z)).$$

The goal of GAT module is to make SE at different locations satisfy the constraints of spatial

approximation while preserving as much information as possible in CE, that is, the correlation of

each SE at n positions UU^T is close to Σ and U is close to Z. Thus, the objective function can

be written as:

$$Loss_2 = \lambda_{global} * L_{global} + \lambda_{local} * L_{local},$$

where $L_{global} = MSE(UU^T, \Sigma)$, and $L_{local} = MSE(U, Z)$.

Architecture of SECE

For the AE module, the dimension of hidden layer is 128, and that of bottle layer is 32. We use

Adam optimizer to minimize $Loss_1$ with learning rate of 1e-3 and dropout rate of 0.1. For the

GAT module, the dimension of two GAT layers is consistent with bottle layer. We use Adam

optimizer to minimize $Loss_2$ with learning rate of 1e-2 and dropout rate of 0.2. The numbers of

iterations for the two modules are set as 40 and 50 by default respectively.

λ_{local} and λ_{global} are set as 1.0 and 0.08 by default. To be more specific, for data with

single-cell resolution or close to single-cell resolution (e.g., Stereo-seq and Slide-seqV2), we

employed hyperparameters 1 and 0.08. For data without single-cell resolution, such as

Visium, we employed hyperparameters 1 and 0.3. For STARmap data, which provided single-

molecule resolution, we used hyperparameters 1 and 2. When working with spatial

transcriptome data from other platforms, we recommended adjusting the value of λ_{global}

based on the number of spots. This was because λ_{global} governed the role of spatial global

information. Generally, as the number of spots increased, a smaller value of λ_{global} tended to

be more appropriate.

Preprocessing of datasets

For Stereo-seq datasets, cells with an expression level below 200 were removed according to

Stereo-seq studies [15], then we further screened genes expressed in less than 20 cells. For other

datasets, we screened spots with an expression level below 20 and genes that expressed less than

20 spots. The filtered expression matrix and its corresponding coordinates were input into SECE

for analysis.

Clustering

We use Leiden [62] and mClust [63] to cluster CE and SE, respectively. We cluster the CE based

on Leiden and then assigned the cell type labels. We use the Leiden algorithm implemented in

SCANPY package by calling scanpy.tl.leiden. We cluster the SE based on mClust and then

annotate the spatial domains. Clustering was performed based on SE by rmclust in mclust

package.

Spatial trajectory inference

We perform the trajectory inference using SE generated from SECE and STAGATE to verify their

reliability. We use PAGA [25] implemented in the SCANPY [27] package to perform the

trajectory inference between spatial domains. The PAGA graphs are calculated by scanpy.tl.paga

and visualized by scanpy.pl.paga. We use Monocle3 [26] to perform the trajectory inference for

individual spots. UMAP embeddings generated from SE are used to calculate the pseudo times by

learn_graph and order_cells in Monocle3 package.

Deconvolution

584　For Visium dataset without single-cell resolution, we perform deconvolution to analyze the cell

585　type composition using software Cell2location and annotated scRNA-seq dataset [40].

586　Cell2location is run according to the tutorial and default parameters.

Differential expression analysis

589　Differential expression analysis was used to identify cluster-specific marker genes where all the

590　clusters are pairwise compared using the Wilcoxon method. Each identified marker gene was

591　expressed in a minimum of 25% of cells and at a minimum log fold change threshold of 0.25.

Comparison of spatial domain identification approaches

We compared the existing spatial domain identification methods.

BayesSpace. BayesSpace was downloaded from https://github.com/edward130603/BayesSpace.

It was run using default parameters specified as top 2,000 HVGs performing PCA and top 15 PCs

to infer spatial domains.

SpaGCN. SpaGCN was downloaded from https://github.com/jianhuupenn/SpaGCN. SpaGCN was

run using default training parameters specified as 0.05 for learning rate and 200 for max epochs.

For the Visium data we set histology=True and for the other data we set histology=False. In

addition, we set s=1 for STARmap data as recommended in the article, and the rest of the data

follow the default parameters.

STAGATE. STAGATE was downloaded from https://github.com/QIFEIDKN/STAGATE. We run

the default parameters except for rad_cutoff. We followed the tutorial of STAGATE and set

rad_cutoff=50 for Stereo-seq and Slide-seqV2 datasets, rad_cutoff=150 for Visium datasets, and

rad_cutoff=400 for STARmap datasets.

BASS. We downloaded BASS from https://github.com/zhengli09/BASS, and ran BASS following

the vignettes https://zhengli09.github.io/BASS-Analysis.

SpaceFlow. We downloaded SpaceFlow from https://github.com/hongleir/SpaceFlow, and ran

SpaceFlow following https://github.com/hongleir/SpaceFlow/blob/master/tutorials.

GraphST. We downloaded GraphST from https://github.com/JinmiaoChenLab/GraphST, and ran

GraphST following https://deepst-tutorials.readthedocs.io/en/latest/index.html.

SpatialPCA. We downloaded the R package from https://github.com/shangll123/SpatialPCA, and

ran SpatialPCA following http://lulushang.org/SpatialPCA_Tutorial/.

conST. We referred to https://github.com/ys-zong/conST to ran conST.

Compared methods for cell type identification

We compared the existing cell type identification methods.

SpiceMix. We referred to https://github.com/ma-compbio/SpiceMix to ran SpiceMix.

Seurat. We used the R package of Seurat v 4.2.1.

SC3. We downloaded SC3 from https://github.com/hemberg-lab/SC3, and ran SC3 following

http://www.bioconductor.org/packages/release/bioc/vignettes/SC3/inst/doc/SC3.html.

CIDR. We referred to https://github.com/VCCRI/CIDR to ran SpiceMix..

Evaluation

Local inverse Simpson's index (LISI). LISI [64] is used to measure the degree of local mixing,

and we use it to evaluate the level of spatial pattern. For every sample i, *LISI* can be formulated

as

641
$$LISI(i) = \frac{1}{\sum_{l \in L} p_i(l)},$$

where $p(l)$ is the probability that the spatial domain cluster label l exists in the local

neighborhood of sample i, and L is the set of spatial domains.

In this paper, local neighborhoods are generated by spatial location, labels are spatial domain

labels predicted by each algorithm. The value of LISI is $[1, \infty)$, and smaller LISI indicates better

spatial patterns.

Adjusted Rand index (ARI). ARI evaluates the degree of overlap between the two divisions. In

this paper, we use ARI to evaluate the consistency of the spatial domains divided by various

methods with the ground truth domains. ARI can be written as

651
$$ARI = \frac{\sum_{ij} \binom{n_{ij}}{2} - [\sum_i \binom{a_i}{2} \sum_j \binom{b_j}{2}] / \binom{n}{2}}{\frac{1}{2}[\sum_i \binom{a_i}{2} + \sum_j \binom{b_j}{2}] - [\sum_i \binom{a_i}{2} \sum_j \binom{b_j}{2}] / \binom{n}{2}},$$

where n_{ij}, a_i and b_i are values from the contingency table. Here, the ground truth labels are the

manual annotation labels of domains, and the labels to be evaluated are the results of various

spatial domain identification methods. The value of ARI is $[-1,1]$, and greater ARI indicates

better agreements with true label.

Normalized mutual information (NMI). NMI evaluates the similarity between the predicted and

true labels. In this paper, we use NMI to evaluate the consistency of the spatial domains divided

by various methods with the ground truth domains. NMI can be written as

660
$$NMI(P,T) = \frac{2MI(P,T)}{H(P)+H(T)},$$

661 where MI is Mutual Information, and $H(P)$ is Shannon entropy. The value of NMI is [0,1], and

662 greater NMI indicates higher similarity with true label.

664 *Accuracy (ACC)*. ACC is calculated as

665
$$ACC = \frac{\sum_{i=1}^{N} \delta(r_i, o(s_i))}{N},$$

666 where N is the number of spots, r_i and s_i are the real and predicted spatial domain label of spot

667 i, respectively. δ is a function that can be defined as

668
$$\delta(x,y) = \begin{cases} 1, & x = y \\ 0, & x \neq y \end{cases}$$

669 o is a mapping function that takes the real label r_i as the reference label and then rearranges s_i

670 in the same arrangement, which is implemented using the classical Kuhn-Munkres algorithm [28].

672 *Average Silhouette width (ASW)*. ASW is used to describe the sharpness for each category after

673 clustering, and we use ASW to evaluate how well the SE obtained by various methods explain the

674 known spatial layers. For every sample i, Silhouette width $S(i)$ is calculated as

675
$$S(i) = \frac{b(i)-a(i)}{\max\{a(i),b(i)\}},$$

where $a(i)$ is the average distance from i to all data points in the same cluster C_i, and

$b(i)$ is calculated similarly to $a(i)$, just need to get multiple average distance from all clusters

except C_i, then choose the smallest value from it as the final $b(i)$.

In this paper, distance is calculated by SE of various methods, and cluster is the layer

annotated by original author. The value of ASW is $[-1,1]$, and greater ASW indicates better

embeddings.

Adjusted Rand index of cell types (ARI-Cell Type). For data with original cell type annotations,

we assessed the consistency of various cell type identification algorithms with the original

annotations using ARI. The calculation formula of ARI is in the Adjusted Rand index (ARI)

section. In order to distinguish it from the evaluation index of the spatial domain identification, we

named this ARI as ARI-Cell Type. Here, the ground truth labels are the annotation labels of the

original articles, and the labels to be evaluated are the results of various cell type identification

algorithms.

691   Declarations

Ethics approval and consent to participate

No ethnical approval was required for this study.

Consent for publication

Not applicable.

Availability of data and materials

Slide-seq datasets are available at https://portals.broadinstitute.org/single_cell/study/slide-seq-s

tudy. Stereo-seq mouse hemibrain dataset is available at https://db.cngb.org/stomics/mosta/.

Stereo-seq olfactory bulb dataset is available at https://github.com/JinmiaoChenLab/SEDR_a

nalyses. Slide-seqV2 hippocampus datasets are available at https://singlecell.broadinstitute.or

g/. STARmap mouse visual cortex dataset is available at http://clarityresourcecenter.org/. Vi

sium human breast cancer dataset is available at https://www.10xgenomics.com/resources/dat

asets/human-breast-cancer-block-a-section-1-1-standard-1-1-0. Visium DLPFC dataset is avail

able within the spatialLIBD package (http:://spatial.libd.org/spatialLIBD). We also organized

these raw data, including expression counts and coordinates, as well as H&E images of

Visium, into a format that is easy to read by SCANPY, which were available at https://dr

ive.google.com/drive/my-drive. The SECE algorithm is implemented and provided as a pip

installable Python package which is available on Github https://github.com/yuyuanyuana/SE

CE. The source code and datasets are available at Zenodo https://zenodo.org/record/813068

2.

Competing interests

The authors declare that they have no competing interests.

Funding

This project was supported by National Key R&D Program of China (2019YFA0904400, Z.X.), Guangzhou Science and Technology Project (202201020336, Z.X.).

Authors' contributions

Z.X. conceived and supervised the study. YY.Y. and Z.X. designed the study. YY.Y. analyzed the data. All the authors interpreted the data. YY.Y. and Z.X. wrote the manuscript. All authors read and approved the manuscript.

Acknowledgements

We would like to thank all the data and software contributors who make this research possible. We also thank the Center for Precision Medicine at Sun Yat-sen University for the long-term support.

Authors' information

State Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-sen University, Guangzhou, China

Yuanyuan Yu, Yao He & Zhi Xie

References

1. Maynard KR, Collado-Torres L, Weber LM, Uytingco C, Barry BK, Williams SR, Catallini JL, 2nd, Tran MN, Besich Z, Tippani M, et al: **Transcriptome-scale spatial gene expression in the human dorsolateral prefrontal cortex.** *Nat Neurosci* 2021, **24:**425-436.

2.   Wang M, Hu Q, Lv T, Wang Y, Lan Q, Xiang R, Tu Z, Wei Y, Han K, Shi C, et al: **High-**

resolution 3D spatiotemporal transcriptomic maps of developing Drosophila embryos and

larvae. *Dev Cell* 2022, **57:**1271-1283 e1274.

3.   Hunter MV, Moncada R, Weiss JM, Yanai I, White RM: **Spatially resolved transcriptomics**

reveals the architecture of the tumor-microenvironment interface. *Nat Commun* 2021,

12:6278.

4.   Wang X, Allen WE, Wright MA, Sylwestrak EL, Samusik N, Vesuna S, Evans K, Liu C,

Ramakrishnan C, Liu J, et al: **Three-dimensional intact-tissue sequencing of single-cell**

transcriptional states. *Science* 2018, **361**.

5.   Lubeck E, Coskun AF, Zhiyentayev T, Ahmad M, Cai L: **Single-cell in situ RNA profiling by**

sequential hybridization. *Nat Methods* 2014, **11:**360-361.

6.   Shah S, Lubeck E, Zhou W, Cai L: **In Situ Transcription Profiling of Single Cells Reveals**

Spatial Organization of Cells in the Mouse Hippocampus. *Neuron* 2016, **92:**342-357.

7.   Eng CL, Lawson M, Zhu Q, Dries R, Koulena N, Takei Y, Yun J, Cronin C, Karp C, Yuan GC,

Cai L: **Transcriptome-scale super-resolved imaging in tissues by RNA seqFISH.** *Nature* 2019,

568:235-239.

8.   Moffitt JR, Bambah-Mukku D, Eichhorn SW, Vaughn E, Shekhar K, Perez JD, Rubinstein ND,

Hao J, Regev A, Dulac C, Zhuang X: **Molecular, spatial, and functional single-cell profiling of**

the hypothalamic preoptic region. *Science* 2018, **362**.

9.   Lee JH, Daugharthy ER, Scheiman J, Kalhor R, Yang JL, Ferrante TC, Terry R, Jeanty SS, Li C,

Amamoto R, et al: **Highly multiplexed subcellular RNA sequencing in situ.** *Science* 2014,

343:1360-1363.

10.   Moses L, Pachter L: **Museum of spatial transcriptomics.** *Nat Methods* 2022, **19:**534-546.

11.   Stahl PL, Salmen F, Vickovic S, Lundmark A, Navarro JF, Magnusson J, Giacomello S, Asp M,

Westholm JO, Huss M, et al: **Visualization and analysis of gene expression in tissue sections**

by spatial transcriptomics. *Science* 2016, **353:**78-82.

12.   Stickels RR, Murray E, Kumar P, Li J, Marshall JL, Di Bella DJ, Arlotta P, Macosko EZ, Chen F:

Highly sensitive spatial transcriptomics at near-cellular resolution with Slide-seqV2. *Nat*

Biotechnol 2021, **39:**313-319.

13.   Vickovic S, Eraslan G, Salmen F, Klughammer J, Stenbeck L, Schapiro D, Aijo T, Bonneau R,

771   Bergenstrahle L, Navarro JF, et al: **High-definition spatial transcriptomics for in situ tissue**

772   **profiling.** *Nat Methods* 2019, **16:**987-990.

773   14.   Cho CS, Xi J, Si Y, Park SR, Hsu JE, Kim M, Jun G, Kang HM, Lee JH: **Microscopic**

774   **examination of spatial transcriptome using Seq-Scope.** *Cell* 2021, **184:**3559-3572 e3522.

775   15.   Chen A, Liao S, Cheng M, Ma K, Wu L, Lai Y, Qiu X, Yang J, Xu J, Hao S, et al:

776   **Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA nanoball-patterned**

777   **arrays.** *Cell* 2022, **185:**1777-1792 e1721.

778   16.   Ortiz C, Navarro JF, Jurek A, Martin A, Lundeberg J, Meletis K: **Molecular atlas of the adult**

779   **mouse brain.** *Sci Adv* 2020, **6:**eabb3446.

780   17.   Zhao E, Stone MR, Ren X, Guenthoer J, Smythe KS, Pulliam T, Williams SR, Uytingco CR,

781   Taylor SEB, Nghiem P, et al: **Spatial transcriptomics at subspot resolution with BayesSpace.**

782   *Nat Biotechnol* 2021, **39:**1375-1384.

783   18.   Shang L, Zhou X: **Spatially aware dimension reduction for spatial transcriptomics.** *Nat*

784   *Commun* 2022, **13:**7203.

785   19.   Li Z, Zhou X: **BASS: multi-scale and multi-sample analysis enables accurate cell type**

786   **clustering and spatial domain detection in spatial transcriptomic studies.** *Genome Biol* 2022,

787   **23:**168.

788   20.   Hu J, Li X, Coleman K, Schroeder A, Ma N, Irwin DJ, Lee EB, Shinohara RT, Li M: **SpaGCN:**

789   **Integrating gene expression, spatial location and histology to identify spatial domains and**

790   **spatially variable genes by graph convolutional network.** *Nat Methods* 2021, **18:**1342-1351.

791   21.   Dong K, Zhang S: **Deciphering spatial domains from spatially resolved transcriptomics with**

792   **an adaptive graph attention auto-encoder.** *Nat Commun* 2022, **13:**1739.

793   22.   Long Y, Ang KS, Li M, Chong KLK, Sethi R, Zhong C, Xu H, Ong Z, Sachaphibulkij K, Chen A,

794   et al: **Spatially informed clustering, integration, and deconvolution of spatial transcriptomics**

795   **with GraphST.** *Nat Commun* 2023, **14:**1155.

796   23.   Ren H, Walker BL, Cang Z, Nie Q: **Identifying multicellular spatiotemporal organization of**

797   **cells with SpaceFlow.** *Nat Commun* 2022, **13:**4076.

24. Zong Y, Yu T, Wang X, Wang Y, Hu Z, Li Y: **conST: an interpretable multi-modal**

contrastive learning framework for spatial transcriptomics. *bioRxiv*

2022**:**2022.2001.2014.476408.

25. Wolf FA, Hamey FK, Plass M, Solana J, Dahlin JS, Gottgens B, Rajewsky N, Simon L, Theis FJ:

PAGA: graph abstraction reconciles clustering with trajectory inference through a topology

preserving map of single cells. *Genome Biol* 2019, **20:**59.

26. Cao J, Spielmann M, Qiu X, Huang X, Ibrahim DM, Hill AJ, Zhang F, Mundlos S, Christiansen

L, Steemers FJ, et al: **The single-cell transcriptional landscape of mammalian organogenesis.**

Nature 2019, **566:**496-502.

27. Wolf FA, Angerer P, Theis FJ: **SCANPY: large-scale single-cell gene expression data analysis.**

Genome Biol 2018, **19:**15.

28. Munkres J: **Algorithms for the Assignment and Transportation Problems.** *Journal of the*

Society for Industrial and Applied Mathematics 1957, **5:**32-38.

29. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, Lee MJ, Wilk AJ, Darby

C, Zager M, et al: **Integrated analysis of multimodal single-cell data.** *Cell* 2021, **184:**3573-3587

e3529.

30. Kiselev VY, Kirschner K, Schaub MT, Andrews T, Yiu A, Chandra T, Natarajan KN, Reik W,

Barahona M, Green AR, Hemberg M: **SC3: consensus clustering of single-cell RNA-seq data.**

Nat Methods 2017, **14:**483-486.

31. Lin P, Troup M, Ho JW: **CIDR: Ultrafast and accurate clustering through imputation for**

single-cell RNA-seq data. *Genome Biol* 2017, **18:**59.

32. Chidester B, Zhou T, Alam S, Ma J: **SPICEMIX enables integrative single-cell spatial**

modeling of cell identity. *Nat Genet* 2023, **55:**78-88.

33. Saunders A, Macosko EZ, Wysoker A, Goldman M, Krienen FM, de Rivera H, Bien E, Baum M,

Bortolin L, Wang S, et al: **Molecular Diversity and Specializations among the Cells of the**

Adult Mouse Brain. *Cell* 2018, **174:**1015-1030.e1016.

34. Fu H, Xu H, Chong K, Li M, Ang KS, Lee HK, Ling J, Chen A, Shao L, Liu L, Chen J:

Unsupervised Spatially Embedded Deep Representation of Spatial Transcriptomics. *bioRxiv*

2021.

35.   Su X, Vasilkovska T, Frohlich N, Garaschuk O: **Characterization of cell type-specific S100B**

expression in the mouse olfactory bulb. *Cell Calcium* 2021, **94:**102334.

36.   Nagayama S, Homma R, Imamura F: **Neuronal organization of olfactory bulb circuits.** *Front*

Neural Circuits 2014, **8:**98.

37.   Beiersdorfer A, Wolburg H, Grawe J, Scheller A, Kirchhoff F, Lohr C: **Sublamina-specific**

organization of the blood brain barrier in the mouse olfactory nerve layer. *Glia* 2020,

68:631-645.

38.   Van der Maaten L, Hinton G: **Visualizing data using t-SNE.** *Journal of machine learning*

research 2008, **9**.

39.   Larsson L, Frisen J, Lundeberg J: **Spatially resolved transcriptomics adds a new dimension to**

genomics. *Nat Methods* 2021, **18:**15-18.

40.   Wu SZ, Al-Eryani G, Roden DL, Junankar S, Harvey K, Andersson A, Thennavan A, Wang C,

Torpy JR, Bartonicek N, et al: **A single-cell and spatially resolved atlas of human breast**

cancers. *Nat Genet* 2021, **53:**1334-1347.

41.   Kleshchevnikov V, Shmatko A, Dann E, Aivazidis A, King HW, Li T, Elmentaite R, Lomakin A,

Kedlian V, Gayoso A, et al: **Cell2location maps fine-grained cell types in spatial**

transcriptomics. *Nat Biotechnol* 2022, **40:**661-671.

42.   Bilandzic M, Rainczuk A, Green E, Fairweather N, Jobling TW, Plebanski M, Stephens AN:

Keratin-14 (KRT14) Positive Leader Cells Mediate Mesothelial Clearance and Invasion by

Ovarian Cancer Cells. *Cancers (Basel)* 2019, **11**.

43.   Cheung KJ, Padmanaban V, Silvestri V, Schipper K, Cohen JD, Fairchild AN, Gorin MA,

Verdone JE, Pienta KJ, Bader JS, Ewald AJ: **Polyclonal breast cancer metastases arise from**

collective dissemination of keratin 14-expressing tumor cell clusters. *Proc Natl Acad Sci U S A*

2016, **113:**E854-863.

44.   Martinez-Ordonez A, Seoane S, Avila L, Eiro N, Macia M, Arias E, Pereira F, Garcia-Caballero

T, Gomez-Lado N, Aguiar P, et al: **POU1F1 transcription factor induces metabolic**

reprogramming and breast cancer progression via LDHA regulation. *Oncogene* 2021,

40:2725-2740.

45.   Nalio Ramos R, Missolo-Koussou Y, Gerber-Ferder Y, Bromley CP, Bugatti M, Nunez NG,

Tosello Boari J, Richer W, Menger L, Denizeau J, et al: **Tissue-resident FOLR2(+)**

macrophages associate with CD8(+) T cell infiltration in human breast cancer. *Cell* 2022,

185:1189-1207 e1125.

46. Zhang H, Wang Y, Liu C, Li W, Zhou F, Wang X, Zheng J: **The Apolipoprotein C1 is involved**

in breast cancer progression via EMT and MAPK/JNK pathway. *Pathol Res Pract* 2022,

229:153746.

47. Seo SU, Woo SM, Im SS, Jang Y, Han E, Kim SH, Lee H, Lee HS, Nam JO, Gabrielson E, et al:

Cathepsin D as a potential therapeutic target to enhance anticancer drug-induced apoptosis

via RNF183-mediated destabilization of Bcl-xL in cancer cells. *Cell Death Dis* 2022, **13:**115.

48. Zhang C, Zhang M, Song S: **Cathepsin D enhances breast cancer invasion and metastasis**

through promoting hepsin ubiquitin-proteasome degradation. *Cancer Lett* 2018, **438:**105-

115.

49. Revel M, Sautes-Fridman C, Fridman WH, Roumenina LT: **C1q+ macrophages: passengers or**

drivers of cancer progression. *Trends Cancer* 2022, **8:**517-526.

50. Noblejas-Lopez MDM, Nieto-Jimenez C, Morcillo Garcia S, Perez-Pena J, Nuncia-Cantarero M,

Andres-Pretel F, Galan-Moya EM, Amir E, Pandiella A, Gyorffy B, Ocana A: **Expression of**

MHC class I, HLA-A and HLA-B identifies immune-activated breast tumors with favorable

outcome. *Oncoimmunology* 2019, **8:**e1629780.

51. Nomura T, Huang W-C, E Zhau H, Josson S, Mimata H, WK Chung L: **β2-Microglobulin-**

mediated signaling as a target for cancer therapy. *Anti-Cancer Agents in Medicinal Chemistry*

(Formerly Current Medicinal Chemistry-Anti-Cancer Agents) 2014, **14:**343-352.

52. Wu B, Pan Y, Liu G, Yang T, Jin Y, Zhou F, Wei Y: **MRPS30-DT Knockdown Inhibits Breast**

Cancer Progression by Targeting Jab1/Cops5. *Front Oncol* 2019, **9:**1170.

53. Zeng B, Ge C, Li R, Zhang Z, Fu Q, Li Z, Lin Z, Liu L, Xue Y, Xu Y, et al: **Knockdown of**

microsomal glutathione S-transferase 1 inhibits lung adenocarcinoma cell proliferation and

induces apoptosis. *Biomed Pharmacother* 2020, **121:**109562.

54. Huang X, Yu J, Lai S, Li Z, Qu F, Fu X, Li Q, Zhong X, Zhang D, Li H: **Long Non-Coding RNA**

LINC00052 Targets miR-548p/Notch2/Pyk2 to Modulate Tumor Budding and Metastasis of

Human Breast Cancer. *Biochem Genet* 2022.

55. Xiong D, Wang D, Chen Y: **Role of the long non-coding RNA LINC00052 in tumors.** *Oncol*

Lett 2021, **21:**316.

56. Salameh A, Fan X, Choi BK, Zhang S, Zhang N, An Z: **HER3 and LINC00052 interplay**

promotes tumor growth in breast cancer. *Oncotarget* 2017, **8:**6526-6539.

57. Grzybowska-Szatkowska L, Slaska B: **Mitochondrial NADH dehydrogenase polymorphisms**

are associated with breast cancer in Poland. *J Appl Genet* 2014, **55:**173-181.

58. Lyu L, Wang M, Zheng Y, Tian T, Deng Y, Xu P, Lin S, Yang S, Zhou L, Hao Q, et al:

Overexpression of FAM234B Predicts Poor Prognosis in Patients with Luminal Breast

Cancer. *Cancer Manag Res* 2020, **12:**12457-12471.

59. Eraslan G, Simon LM, Mircea M, Mueller NS, Theis FJ: **Single-cell RNA-seq denoising using a**

deep count autoencoder. *Nat Commun* 2019, **10:**390.

60. Lopez R, Regier J, Cole MB, Jordan MI, Yosef N: **Deep generative modeling for single-cell**

transcriptomics. *Nat Methods* 2018, **15:**1053-1058.

61. Hafemeister C, Satija R: **Normalization and variance stabilization of single-cell RNA-seq data**

using regularized negative binomial regression. *Genome Biol* 2019, **20:**296.

62. Traag VA, Waltman L, van Eck NJ: **From Louvain to Leiden: guaranteeing well-connected**

communities. *Sci Rep* 2019, **9:**5233.

63. Scrucca L, Fop M, Murphy TB, Raftery AE: **mclust 5: Clustering, Classification and Density**

Estimation Using Gaussian Finite Mixture Models. *R J* 2016, **8:**289-317.

64. Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K, Baglaenko Y, Brenner M, Loh

PR, Raychaudhuri S: **Fast, sensitive and accurate integration of single-cell data with**

Harmony. *Nat Methods* 2019, **16:**1289-1296.

Figures

Fig.1. Overview of SECE. (A) SECE is applicable to a wide range of spatial transcriptome data

with different resolution, including subcellular, single-cell, near-single-cell, and multicellular

resolutions. **(B)** SECE takes gene expression profile and spatial coordinates as input. SECE begins

with AE module, which learns CE under the count distribution. Then, GAT module learns the SE

by balancing the expressive proximity and the spatial proximity, which are measured by the

distance of the CE and the distance of the coordinate, respectively. **(C)** The main output includes

spatial region and cell type labels, which are obtained by clustering CE and SE separately. SE can

also perform downstream tasks such as low dimensional visualization and trajectory inference. **(D)**

Spatial domain identification and trajectory inference of SEAL_PCA (top), SEAL_NG (middle)

and SEAL_PCA_NG (bottom).

Fig. 2. Application of SECE to mouse cortex STARmap data. (A) Layer structure of the tissue

section from the original study. **(B)** Left: Spatial domains identified by SECE. Right: Cell types

identified by SECE. Astro, astrocytes; Oligo, oligodendrocytes; EX, excitatory neurons; IN

Neuron, inhibitory neurons; Endo, endothelial cells; SMC, smooth muscle cells. **(C)** Spatial

domains identified by BASS, SpaceFlow, GraphST, STAGATE, SpatialPCA, SpaGCN,

BayesSpace, and conST. **(D)** Accuracy of spatial domain identification for different methods

928 measured by ARI, NMI, and ACC. (**E**) Barplot of ASW shows the discrimination capability in

929 ground truth layers of SEs learned by SECE, SpaceFlow, GraphST, STAGATE, and SpatialPCA.

930 (**F**) UMAP visualization generated by SECE, SpaceFlow, GraphST, STAGATE, and SpatialPCA,

931 colored by annotated layers. (**G**) PAGA graph generated by SECE, SpaceFlow, GraphST,

932 STAGATE, and SpatialPCA.

Fig. 3. Application of SECE to mouse hippocampus Slide-seqV2 data. (A) The annotation of

hippocampus structures from the Allen Reference Atlas of an adult mouse brain. **(B)** Left: Spatial

visualization of domains identified by SECE. Right: UMAP visualization of domains identified by

SECE. Spatial domains were annotated by the structure annotation showed in the Allen Reference

Atlas. CA, Ammon's horn; DG, Dentate gyrus. **(C)** Spatial visualization of CA1, CA2, and CA3

domains identified by SECE (Left) and the corresponding marker genes *Wsf1*, *Rgs14*, and *Nptxr*

(right). **(D)** 14 spatial regions identified by BASS, SpaceFlow, GraphST, STAGATE, SpatialPCA,

SpaGCN, BayesSpace, and conST. **(E)** Pseudo-time of each cell calculated by Monocle3 based on

embedding of SECE, SpaceFlow, GraphST, STAGATE, and SpatialPCA. **(F)** Pseudo-time of the

944 cells in each layer of the isocortex based on SECE, SpaceFlow, GraphST, STAGATE, and

945 SpatialPCA.

Fig. 4. Application of SECE to mouse olfactory bulb Stereo-seq data. (A) Cell types identified

by SECE. GC, Granule cells; M/TC, Mitral and tufted cells; PGC, Periglomerular cells; OEC,

Olfactory ensheathing cells. **(B)** Left: The annotation of mouse olfactory bulb structures from the

Allen Reference Atlas. Right: Spatial visualization of domains identified by SECE. RMS, Rostral

migratory stream; GCL, Granule cell layer; IPL, Inner plexiform layer; MCL, Mitral cell layer;

EPL, External plexiform layer; GL, Glomerular layer; ONL, Olfactory nerve layer. **(C)** 9 spatial

regions identified by BASS, SpaceFlow, GraphST, STAGATE, SpatialPCA, SpaGCN,

BayesSpace, and conST. **(D)** UMAP visualization generated by SECE, SpaceFlow, GraphST,

956 STAGATE, and SpatialPCA, colored by annotated layers. (**E**) PAGA graph generated by SECE,

957 SpaceFlow, GraphST, STAGATE, and SpatialPCA.

Fig. 5. Application of SECE to mouse brain Stereo-seq data. (A) The annotation of mouse

olfactory bulb structures from the Allen Reference Atlas. **(B)** Left: Spatial visualization of

domains identified by SECE. Right: UMAP visualization of domains identified by SECE. Spatial

domains were annotated by the structure annotation showed in the Allen Reference Atlas. sl/r,

stratum lacunosum / raditum cornu ammonis; DG, dentate gyrus; FT, fiber tract; MLDG,

molecular layer of dentate gyrus; MRN, midbrain reticular nucleus; SN, substantia nigra; VTA,

ventral tegmental area. **(C)** 20 spatial regions identified by BASS, SpaceFlow, GraphST,

STAGATE, SpatialPCA, SpaGCN, BayesSpace, and conST. **(D)** Boxplot of LISI measuring the

968   degree of spatial aggregation of domains identified by different methods. (**E**) Cell types identified

969   by SECE. Astro, astrocytes; DA Neuron: dopaminergic neuron; EX, excitatory neurons; CA,

970   Cornu Ammonis; CAA, cortical amygdalar area; LVC, lateral-ventral cortex; MB, midbrain; Ery:

971   erythrocyte; GN DG: dentate gyrus of granule neurons; IN, inhibitory neurons; Tele:

972   telencephalon; Oligo, oligodendrocytes; Micro, microglia.

Fig. 6 Application of SECE to human breast cancer Visium data. (A) Pathology annotation of the tissue section from the original study. **(B)** Spatial regions identified by SECE. **(C)** Fine regions identified by SECE, that is, DCIS/LCIS-inner, DCIS/LCIS-outer, DCIS/LCIS-edge, IDC-inner, IDC-outer, and IDC-edge. **(D)** Cell type compositions of 6 fine regions. **(E)** Heatmaps of normalized expression of signature genes identified in the DE analysis based on 6 fine regions. **(F)** Pseudo-times of spots calculated by Monocle3 in DCIS/LCIS (top) and IDC (bottom).