Optical manipulation reveals macrophage mediated mesoscale brain mechanical homeostasis *in vivo*

Woong Young So[1], Bailey Johnson[1], Patricia B. Gordon[1], Kevin S. Bishop[1], Hyeyeon Gong[1,3],

Hannah A Burr[1], Jack Rory Staunton[1], Chenchen Handler[3], Raman Sood[2], Giuliano Scarcelli[3]

and Kandice Tanner[1]

[1]National Cancer Institute, National Institutes of Health (NIH), MD, USA

[2]National Human Genome Research Institute, NIH, MD, USA

[3]University of Maryland - College Park, MD, USA

* Corresponding author information:

Dr. Kandice Tanner; Center for Cancer Research, National Cancer Institute, Building 37, Room

2132, Bethesda, MD 20892; Ph: 260-760-6882; Email: kandice.tanner@nih.gov

Abstract

Tissues are active materials where epithelial turnover, immune surveillance and remodeling of stromal cells such as macrophages all regulate form and function. There is a long-standing interest in harnessing the mechanical properties as a readout of tissue function. Optical techniques can provide the ability to assess the interplay between function, form and the measured mechanical properties in living tissues. However, *in vivo* microscale longitudinal mapping of entire tissues remains challenging. Here, we employed optical tweezer active microrheology and Brillouin microscopy to address two questions: a) how do we reconcile methods to probe mechanical properties of mm tissues with sub cellular resolution? and b) how do we link the measured mechanical properties to stromal mediated-tissue homeostasis? We determined that the corrected Longitudinal modulus measured using Brillouin microscopy correlated well with shear modulus obtained from the optical tweezer measurements at lower frequencies. We determined that the brain tissue mechanical properties are dependent on both the structural presence of the macrophages and the CSF1R receptor that facilitates stromal remodeling linking mechanical measurements of Brillouin microscopy to biological determinants. Ablation of macrophages altered mechanical properties concomitantly with neural structures. Moreover, tissue mechanics and architecture do not return to those of unperturbed age-matched control animals after transient ablation has been resolved. Thus, consideration of the contribution of macrophage remodeling in the maintenance of mechanical homeostasis may be needed to harness macrophage-based therapeutics in disease management.

Main

Organs have distinct mechanical properties[1, 2]. For example, the Young's modulus of the brain is ~200Pa whereas comparable values are ~ 10kPa for muscle tissue[1-3]. Within a given organ, there are regional differences in mechanical properties that are linked to functional properties[3, 4]. A mechanical readout in one functional unit may correspond to normal physiology whereas a comparable measurement in a different unit may indicate the presence of disease. Methods to measure tissue mechanics largely rely on techniques that quantitate the elastic response usually a Young's modulus averaged over large regions[3]. Moreover, mechanical properties are probed at different temporal and force scales[5]. One area of active exploration focuses on what do measurements of "compliance" or "stiffness", "solid-like" or "liquid-like" mean when considering a cell or tissues[5, 6]. Simply, can we link a quantitative mechanical measurement to an underlying biological process? A stand-alone method to assess what aspects of tissue mechanics are relevant for diagnostics or biological mechanisms remains an ongoing process.

Organ homeostasis requires an intricate balance between diverse and dynamic tissue constituents[7, 8]. Within the brain, homeostasis is possible due to specialized networks of diverse neuronal architectures, protected by specialized vasculature systems such as the blood brain barrier and meningeal lymphatics and shaped by tissue resident immune cells[9-12]. During development, tissue resident immune cells such as microglia play a significant role in sculpting the developing brain and neuronal connectivity to establish and maintain normal brain function[13]. Not restricted to developmental processes, these stromal cells also impact the onset of different pathologies such as cancers and neurodegenerative disease[14-17]. As first suggested by D'Arcy Thompson more than a century ago, biophysical properties of tissue are also key

determinants of development[18]. However, probing these in large intact tissues with sub cellular

resolution non-invasively remains challenging. There are several techniques that allows

measurement of shear, young's or longitudinal moduli as a function of different length,

frequency and force scales. But tissue anisotropy complicates our ability to reliably move

interchangeably between these different methods. Thus, we also sought to compare two

independent mechanical mapping methods to compare/contrast the measured physical property

and the underlying physical structure. Here, we focused our studies on optically probing brain

morphogenesis to decipher the interplay between a mechanical signature and stromal dynamics

as a first step in our goal of defining tissue mechanics as a biomarker.

The zebrafish has long been utilized as a model organism as many organs are conserved

in mammalian counterparts. Moreover, the transparency of larval zebrafish allows for non-

invasive optical techniques[19-26]. Here, we applied two optical based techniques: optical

tweezer based active microrheology and Brillouin microscopy in an animal model that

recapitulates spatial anisotropy and temporal evolution of mechanical properties in native intact

tissues. We aimed to determine the biological mechanisms that drive the observed mechanical

signatures. Using a suite of tools where we either chemo- genetically or chemically ablated

stromal constituents, we probed the role of mechanical mediated tissue remodeling in larval

zebrafish. Tissue viscoelasticity was driven by both macrophages and csf1r receptor mediated

crosstalk within the developing brain. Our data provided evidence to link viscoelastic properties

of tissue to macrophage mediated tensional homeostasis in the brain. These data also suggest that

stromal dynamics that induce acute changes in mechanical properties may play an augmented

role in changes in both normal tissue architecture and for therapies targeted against them.

Optical trap active microrheology and Brillouin microscopy show strong correlation in mapping microscale mechanics

The developmental program that drives the formation of the architecturally distinct forebrain, midbrain and hindbrain and blood brain barrier are conserved between zebrafish and mammals and it is heavily reliant on the balance between mechanical forces and tissue stiffness[26, 27] (Extended Data 1a). To measure the mechanical properties of the developing brain from 3 - 4dpf, we employed optical trap based active microrheology (OT-AMR) (Figure 1a, c). First, we introduced 1 micron-diameter polystyrene spheres directly into the brain parenchyma at 2 days post fertilization (2dpf). Using these spheres as mechanical probes, we applied a multiplexed-sinusoidal oscillation based on ~nm amplitude to each bead over a range of frequencies from 7Hz to 15kHz. The displacements of the bead are simultaneously tracked via back focal plane interferometry. Quantitation of the local viscoelasticity, $G^*(\omega)$ at that bead position is achieved by correlating the trap and bead positions. The complex modulus, $G^*(\omega) = |G^*|\exp(i\delta) = G'(\omega) + iG''(\omega)$ which then gives storage ($G'(\omega)$) and loss moduli ($G''(\omega)$). Then,

comparing G' (elastic modulus) and G'' (viscous modulus) by G''/G' would result into loss tangent or hystersivity to determine the relative state between solid-like (more G' contribution) or liquid-like characteristics (more G'' contribution) based on crossover frequency. We show that brain tissue is mechanically heterogeneous as a function of region of the brain and age (Figure 1c, Extended Data 1b). At 3 dpf, the mid-brain is more liquid-like than the hind and forebrain respectively. In contrast, the forebrain is more rigid than the mid and hindbrain. As the brain develops, all regions become more solid-like with the greatest transition occurring for the mid brain. An increase in the complex modulus is observed for all regions with the hind brain showing the greatest increase. G' and G" monotonically increase as a function of frequency and

show power law dependence at frequencies >400Hz (Figure 1c, Extended Data 1b and c). The

normalized complex modulus data from whole frequency range by 3 dpf hindbrain shows that

each brain regions has distinct mechanical property with statistical significance based on paired

t-test (Figure 1e and g, Extended Data 1d and e).

OT-AMR offers the advantage of quantifying absolute values of viscoelasticity[19, 28].

However, the technique necessitates the introduction of probes, that if introduced in sufficient

numbers can perturb normal physiology. Mechanical phenotypes of large tissues can only be

obtained by averaging across many samples assuming random bead distribution. Thus, assessing

the viscoelasticity of an individual fish is difficult. Brillouin microscopy (BM) can overcome this

challenge as it is a purely non-invasive technique, based on the intrinsic inelastic scattering due

to incident photon-phonon interactions in materials, whose spectral characterization provides

information correlated with the underlying mechanical properties (Figure 1b)[29, 30].

Formulaically, the frequency shift (v_B) of Brillouin scattered photons (on the order of GHz)

relate to the local refractive index (n), mass density (ρ), and the longitudinal elastic modulus (M')

of samples (Figure 1b)[29, 30]. Thus, stiffer samples would result in higher frequency shifts. We

determined that Brillouin shifts were comparable in detecting mechanical heterogeneity of brain

regions at 3 and 4dpf (Figure 1d and f). Furthermore, the trends in stiffness at 3 and 4 dpf based

on Brillouin microscopy are the same as those measured using OT-AMR (Figure 1e and g). At 3

dpf, hindbrain is the softest but becomes the stiffest at 4dpf with a significant increase in

frequency shift (Figure 1e and g, Extended Data 1f).

Brillouin longitudinal modulus using calculated density factor shows greater correlation

for OT measurements obtained at lower frequencies in the zebrafish brain

Mechanical heterogeneity is comparable at different temporal scales for BM (~GHz) and

OT-AMR (7 – 15kHz) *in vivo*. We then aimed to link tissue architecture that provide the contrast

for the scattering signals measured by BM (Figure 2a). One of the most dramatic changes is that

the neuronal networks increase in size and complexity as the brain develops. Recently, it has

been proposed that the acetylation of lysin 40 of α-tubulin, a key component of neurons, is

important for stiffness of developing tissue as well as cell mechanics [31, 32]. Whole-mount

immunofluorescence of acetylated tubulin (acTub) of the brain shows differential structure and

clustering of neuronal networks between 3 and 4 dpf, especially in the midbrain and hindbrain.

An increase in normalized intensity of acTub was observed for all brain regions during

development (Figure 2b-d). BM has been recently used to non-invasively infer the mechanical

properties of engineered thick tissues such as organoids and in living animals[29, 30, 33]. The

BM technique requires a direct calibration of microscale mechanical properties for quantitative

readouts of tissue stiffness. Thus, we quantified the relative areas of the different regions of the

brain as well as the number of cells to determine a correction of the density factor for all of the

cells present (Figure 2e-g). During development, not only are resident cells increasing in

numbers but there are infiltrating cells such as macrophages (Figure 2g). We next estimated the

differences in these infiltrating cells by quantifying macrophages at the mid-brain as a function

of development (Figure 2g). Moreover, there are also differences in the timescale between the

OT measurements and that due to the Brillouin phenomenon. Therefore, we compared the raw

and corrected correlation between Brillouin (longitudinal) modulus, M' and OT at 7Hz, 907 Hz

and 15kHz in terms of complex modulus, elastic modulus, and viscous modulus for development

(3-4dpf) (Figure 2h-i, Extended data 2). Comparison of linear fits between log($|G^*|$, G', and G''

for forebrain, midbrain, and hindbrain) and log(Brillouin calculated) revealed poor correlation

between techniques before corrections based on estimated brain refractive index and density

(Figure 2h and i, Extended data 2). While the overall correlations improved post correction,

better fits were obtained for values measured at 7 and 907Hz compared to 15kHz (Figure 2h and

i, Extended data 2).

OT and Brillouin microscopy can detect changes in tissue mechanical properties in CSF1r

zebrafish mutant with delayed macrophage invasion.

157   As we observed difference in macrophage numbers as a function of development, we

then asked if the presence of these cells also contributed to the overall mechanical properties of

the brain. Moreover, if OT and Brillouin would be sensitive to microscale perturbations due to

these infiltrating cells. To assess if both the presence and function of microglia regulated tissue

mechanics, we employed the zebrafish mutant, Panther, a mutant which lacks a functional fms

(M-CSF receptor) gene [34]. Colony stimulating factor 1 receptor (csf1ra) is a key determinant

in microglia development[35-37]. In these fish, early macrophages differentiate and behave

normally in the yolk sac[37], however, there is a delayed invasion of macrophages into the brain.

Colonization of the larval brain is eventually achieved with ~ 4-day delay[37] (Figure 3a and b,

Extended data 3a and b). We determined that the mutant showed a difference in brain

morphology and gene expression at 4 dpf compared to WT (Figure 3a and b, Extended data 3a

and b). Furthermore, there are decreased numbers of macrophages in the mutant brain compared

to the WT (Figure 3a). Additionally, the blood brain barrier (BBB) is impaired in the mutant fish

where increased leakage of 150kDa dextran is observed (Extended Data 3b) compared to WT

(Extended Data 1a). OT measurements revealed that the regional heterogeneity observed in WT

is lost where the magnitude of the complex modulus is comparable for all regions (Figure 3c and

d). Direct comparison to WT revealed that the mutant brain is softer in all brain regions at 4 dpf

(Figure 3e). Additionally, the fore and hindbrain in the mutant become more liquid-like during

development as the crossover frequency where viscous modulus (G'') crosses over elastic

modulus (G') shifts to lower frequency (Figure 3c). G' and G'' monotonically increase as a

function of frequency and show power law dependence at frequencies >400Hz (Figure 3c,

Extended Data 3e and f). However, the measured mechanical property of the mid-brain showed

modest increase during development concomitant with minimal changes in the exponent of the

power law (Extended Data 3e, f, h, and i). The normalized complex modulus data from whole

frequency range by 3 dpf hindbrain revealed a more mechanically homogeneous brain (Figure

3d and e). In addition to reduced macrophages, reduced staining of acetylated tubulin was

observed in the mid brain of the panther mutant (Figure 3f, Extended Data 3g). Similar

examination using Brillouin microscopy revealed that all brain regions at 4 dpf show reduced

Brillouin shifts in the mutant compared to WT(Figure 3g and h).

Acetylated tubulin is a major driver of brain mechanical properties.

As immunofluorescence revealed that there is massive remodeling and increase in

acetylated tubulin. After demonstrating that BM is sensitive to mechanical changes due to

development and changes due to genetic mutants, we then investigated the role of acetylated

tubulin in driving mechanical properties. Hence, we modulated expression of tubulin using two

pharmacological inhibitors, Nocodozale and Blebbistatin. Nocodozale inhibits self-assembly of

tubulin and/or associated proteins. Moreover, it also depolymerizes preformed microtubules in

vitro (Figure 4a-d). Brillouin frequency shifts decreased ~ 150 – 400 MHz for all regions of the

brain parenchyma compared to control (Figure 4a-b). We observed that the mid brain was more

affected where we observed the greatest decrease in BM shift upon treatment with nocodozale

compared to the fore and hind brain (Figure 4 c-d). Myosin II plays a role in cell contractility

which in turn can regulate tissue architecture. Furthermore, during development, Myosin II

activity has been shown to contribute to spindle orientation for cell-division orientation in

epiboly, a key developmental step for zebrafish[38]. Thus, we also probed the role of Myosin II

in regulating the mechanical properties of the brain parenchyma using the pharmacological agent,

Blebbistatin (Figure 4e-f). We confirmed that brain tissue was altered upon Blebbistatin

treatment as acTub staining showed a decreased normalized fluorescence intensity (Figure 4g

and h). Inhibition of Myosin II activity revealed reduced Brillouin frequency shifts decreased

about 15 – 50 MHz for all regions of the brain parenchyma compared to control (Figure 4e-f).

Moreover, the regional trend in stiffness persisted as the hindbrain remains the stiffest region.

However, the forebrain showed the largest decrease as we measured a Brillouin shift of 50 MHz

reflecting the greatest observed reduction in stiffness. acTub intensity at the forebrain was most

affected compared to the other regions. As we saw differences in macrophage infiltration as a

function of development and mutant status, we then investigated if there would be difference

upon inhibition of mechanical properties via modulation of tubulin. We determined that for each

of these treatments there were differences in the number of macrophages that were present in the

mid- brain (Figure 4i and j) .

CSF1r macrophages density regulates brain mechanics

To confirm that the presence of macrophages or their remodeling capabilities can directly

perturb mechanical properties of the brain, we ablated macrophages using two methods and

blocked the receptor using pharmacological methods. First, we injected nanoparticles (liposomes

encapsulating clodronate) directly into the brain (Figure 5a-c). Second, we employed a

chemogenetic model, Tg(mpeg1.1:NTR-IRES-eGFP-CAAX) where macrophages express an

enhanced variant of Escherichia coli nitroreductase (NfsBT41Q/N71S/F124T) (NTR) resulting

in macrophage specific ablation (Figure 5d-f) [39]. Upon treatment with Metronidazole,

mpeg1.1$^+$ cells are ablated. In each condition, we observed a reduction of macrophages. We then

used a small molecule drug, PLX5622, against the csf1r receptor to assess if macrophage

mediated tissue remodeling is important for mechanical homeostasis (Figure 5g-i). After 1 day

post injection (dpi) of clodronate, there was ~55% decrease in macrophage population and ~ 65%

reduction using the tissue specific ablation based on neutral red staining assay (Figure 5b and e)

compared to control fish. Macrophage depletion due to clodronate liposomes or due to

macrophage specific ablation caused softening of whole brain parenchyma (Figure 5c and f) as

all brain regions had a decrease in frequency shifts in the range of 70 – 90 MHz compared to

control counterparts. The hindbrain showed the greatest reduction in stiffness concomitant with

alterations in acTub expression (Extended Data 5). We observed a reduction of macrophages in

the treated fish concomitant with a softening of all brain regions as measured using BM (Figure

5g-i). However, the greatest decrease in BM shift was observed for fish with the specific genetic

ablation of the macrophages.

Discussion

Mechanical properties have been used as metrics to identify the presence of pathologies

since antiquity where touch can sometimes serve as a sensile guide to assist in surgical resection

of cancerous tissue[40]. Not restricted to cancers, a differential mechanical phenotype is also a

hallmark of diseases such as fibrosis and diabetes[41]. In spite of historical and modern

approaches, tissue mechanics are not yet used as either a diagnostic technique or as a biomarker

in the clinic. As a first step, we must quantify these metrics in native tissue. Elastography

methods, such as Brillouin scattering microscopy, form a class of non-invasive techniques that

are used in preclinical models and in the clinic[42-44]. However, a key goal in using these

techniques as stand alone diagnostic tools. Specifically, one outstanding question is what

structural elements regulate the scattering changes as measured using BM. Recent studies in the

zebrafish show the broad utility of optical based techniques in probing biophysical properties

such as viscoelasticity and hemodynamic forces *in vivo*[19-21, 23, 25, 28, 45, 46]. Here, we

provide evidence for the biological origin of the BM scattering signal in the larval zebrafish

brain. Our data suggest that these signals are largely driven by microscale neuronal architecture,

macrophage dynamics, and sub-cellular receptor modulation. Beyond linking BM to its

biological determinants, we validated the extracted micro-mechanical signatures by performing a

direct comparison against a gold-standard microscale mechanical test, OT-AMR. We first show

that there is good agreement between the techniques despite the fact that they probe mechanical

properties at different timescales. BM probes the longitudinal modulus on the GHz timescale

which differs from both the Young's and shear moduli measured by OT-AMR. Our results

confirm congruence between traditional rheological methods in cells and tissues. There are some

differences, such as the fact that a significant shift was not observed as the fish ages for BM

measurements as observed for the OT-AMR (Figure 1d-f). This could be due to the contributions

of the "viscous" component of the material that changes as a function of aging as we observed

regional differences in viscous modulus as a function of age, where the forebrain became more

"liquid-like", and the hindbrain became more "solid-like" from 3 to 4 dpf. In addition, the power

law analysis revealed age dependent transitions between semi-flexible and flexible polymers at

high frequencies (Extended Data 1c). These results can possibly drive differences in the

refractive indices and densities that are needed to interpret the longitudinal modulus as measured

by BM (Figure 2e-i, Extended Data 2).

Recent advancements in contact-based atomic force microscopy (AFM) have generated

valuable insights on tissue mechanics during embryogenesis[31, 47]. Our results confirm that

morphogenetic programs such as invading immune cells and proliferation of neurons [18, 48-51].

Stromal cells and the associated signaling pathways associated with the sculpting embryonic

tissue are also important in microenvironmental regulation of tumor etiology and metastatic

colonization[17]. One such cell is the macrophage, a member of the innate immune system[17].

Here, we identified csfr1 macrophages as a key factor in brain tensional homeostasis. We

determined that the mechanical properties are dependent on both the structural presence of the

macrophages and the receptor that facilitates stromal remodeling. The mechanical homeostasis

was modulated by both acute and permanent alterations of macrophage dynamics. Csf1r has

pleiotropic functions in humans, rodents and zebrafish where loss of the receptor resulted in

central nervous system and skeletal deformities[52]. In humans, variants contribute to a spectrum

of development disorders that are classified collectively as BANDDOS: brain abnormalities,

neurodegeneration and dysosteosclerosis[53]. More recently, variants of CSF1R have been

implicated in adult-onset leukodystrophy[14]. The appearance of white matter lesions, axonal

spheroids, and cerebral calcifications have deleterious effects on motor and cognitive functional

resulting in death[14]. Although our results are obtained using larval fish, these observations

may have consequences for adult tissues. Our data suggest that mapping longitudinal mechanical

properties may aid in diagnosing the onset of disease and monitoring its progression. An

important area for future research exists in bridging the knowledge gap between our microscale

techniques and the macroscale techniques such as magnetic resonance elastography which are

already employed in the clinic.

Moreover, in cancers, tissue resident and recruited macrophages can help promote or

suppress tumor growth, tumor cell escape, and distal organ colonization[16, 17]. Consequently,

macrophage specific receptors have been pursued as druggable targets in the treatment of cancer

[54]. These approaches have yielded mixed results in clinical studies. One missing factor could

be that targeting the biochemical signaling of macrophages may alter mechanical properties of

the microenvironment. These changes in turn could either act synergistically or blunt desired

results of tumor clearance. Thus, it suggests that understanding the contribution of macrophage

remodeling in the maintenance of mechanical homeostasis may be needed to harness

macrophage-based therapeutics in disease management. Moreover, it underscores our need to

understand normal tissue homeostasis if we are to design therapeutics that target the

microenvironment. With this information in hand, we can begin to understand the functional role

of tissue mechanics in the establishment and maintenance of normal tissue and what may go

awry at the onset of and progression of disease.

Methods

302   All animal experiments were done under protocols approved by the National Cancer

303 Institute (NCI) and the National Institutes of Health (NIH) Animal Care and Use Committee.

Zebrafish husbandry

Zebrafish were maintained at 28.5 °C on a 14-hour light/10-hour dark cycle according to

standard procedures. Larvae were obtained from natural spawning, raised at 28.5 °C, and

maintained in fish water, 60 mg sea salt (Instant Ocean, #SS15-10) per liter of DI water. For all

experiments, larvae were transferred to fish water supplemented with N-phenylthiourea (PTU;

Millipore Sigma, # P7629-25G) between 18-22 hours post-fertilization to inhibit melanin

formation for enhanced optical transparency. PTU fish embryo water was prepared by dissolving

400 µl of PTU stock (7.5% w/v in DMSO) per 1 L of fish water. Water was replaced twice per

day.

Zebrafish lines

For wild type, AB strain zebrafish were crossed for brain mechanics measurement at 3

and 4 dpf. Additionally, csf1ra mutant zebrafish (Panther[j4as]) were crossed to validate the role of

csf1ra on brain mechanics and development at 3 and 4 dpf. To investigate the role of

macrophage on brain mechanics, Tg(mpeg1.1:NTR-IRES-eGFP-CAAX)[co57] was crossed where

macrophage, mpeg1.1 promoter, is expressed under eGFP. Tg(foxd3:eGFP)[zdf10] zebrafish line

was crossed with Tg(kdlr:mCherry) zebrafish to visualize foxd3 and kdrl. eGFP is expressed

under the foxd3 promoter to label neural crest while mCherry is expressed under kdrl promoter

to visualize vasculature. Tg(foxd3:eGFP) zebrafish were kindly gifted from Ajay Chitnis lab

(NICHD, NIH). For analysis of the zebrafish neural crest, 3 dpf and 4dpf were used for

quantification. Tg(fli1a:eGFP)[y1] zebrafish was crossed to visualize the integrity of blood-brain

barrier upon 150kDa dextran circulation injection at 3 and 4 dpf compared to panther. For all

experiments, at around 22 – 24 hpf, larvae were transferred to fish water supplemented with PTU

to inhibit melanin formation for increased optical transparency. Larvae were then returned to the

incubator at 28.5 ˚C and checked for normal development.

Optical Tweezer (OT) based active microrheology measurement

Zebrafish larvae at 2 dpf were anesthetized using 0.4% buffered tricaine. 2nl of 1 µm

polystyrene beads (ThermoFisher Scientific, #F8816) resuspended in PBS at a final

concentration of 10^7 beads/ml was injected directly into brain parenchyma (midbrain). Injected

larvae were transferred into fresh PTU water and maintained at 28.5 ˚C. For OT measurement,

injected larvae at 3 and 4dpf were anesthetized using 0.4% buffered tricaine and embedded in

1.25% low melting point agarose gel and allowed to polymerize in with cover glass (no. 1.5

thickness). Then, fish water supplemented with tricaine was added to the agarose hydrogel for

the entire time of data acquisition as previously described[23, 24]. The details of our home-built

OT setup are described[19, 29]. Mainly, the equipment is composed of two near infrared lasers, a

1064 nm trapping laser (IPG Photonics, #YLR 20 1064 Y11) to trap and oscillate a bead and a

975 nm detection laser (Lumics, #LU0975M00 1002F10D) to detect the movement of bead.

Along with the two near infrared lasers, there are two quadrant photodiodes (QPD; First Sensor,

#QP154-QHVSD) to detect trapping laser displacement (trap QPD) and the motion of bead

(detection QPD). The trap laser is oscillated by a dual axis acousto-optic deflector (AOD;

IntraAction, #DTD274HD6), which is operated by radio frequency generating cards (Analog

Devices, #AD9854/PCBZ) with on-board temperature-controlled crystal oscillators[19, 29].

Then, the cards are digitally controlled by a data acquisition (DAQ) card (National Instruments,

PCIe 5871R FPGA). Right after AOD, a small fraction of trap laser power (~1%) is directed to

trap QPD. Both trapping and detection lasers go through the backport of an inverted microscope

(Nikon, Eclipse Ti-U) with a long working distance water immersive objective (Nikon,

MRDO7602 CFIPLAN APO VC60XA WI 1.2 NA). Then, a long working distance (WD) and

high numerical aperture (NA) condenser (Nikon, WI 0.9NA) collects the light from objective.

After the condenser, a dichroic mirror (Chroma, ZT1064RDC-2P) reflects both trap and

detection lasers. A bandpass filter (Chroma, #ET980/20x) excludes the trap laser so that the

detection laser reaches detection QPD to record the motion of bead. Then, time-correlated trap

and detection QPD signals from control and samples are collected and conducted by the DAQ

card and custom program LabVIEW (National Instruments). During experiments, condenser is

placed in Kohler illumination to locate beads via a piezo XYZ nanopositioning stage (Prior,

#77011201) and a charge-coupled device (CCD) camera (Andor, Ixon DU-987E-C50-#BV). A

selected bead is precisely positioned at the center of trap after scanning it by detection laser in

three dimensions using a piezo nanopositioning while recording the voltages (V) from the

detection QPD. The relationship between V and displacement (nm) relationship, β, from the

detection QPD is calibrated *in situ* by fitting the central linear line of the detector responses to

scanning the bead through the detection laser in the direction of the trap laser oscillations,

yielding β in V/nm. A trap QPD records the position of the trap laser to find the relative phase

lag between the trap oscillations and bead. The trap stiffness, k, is determined *in situ* for selected

beads based on the active-passive calibration method [29]. The stiffness, k, is determined based

on the active power spectrum, $\tilde{R}_L(\omega)$, and passive power spectrum, $P_U(\omega)$. The stiffness (Eq. 1)

is

$$k = \frac{\Re\{\tilde{R}_L(\omega)\}}{P_U(\omega)}$$

(

1)

373   Active power spectrum, $\tilde{R}_L(\omega)$, is recorded while trap laser is oscillating. The active

374   power spectrum (Eq. 2) is

$$\tilde{R}_L(\omega) \equiv \frac{\tilde{x}_{dr}(\omega)}{-i\omega \tilde{x}_L(\omega)}$$

(

2)

377   Where $\tilde{x}_L(\omega)$ and $\tilde{x}_{dr}(\omega)$ are the Fourier transforms of the time series of the positions of

378   the trap laser and the driven bead respectively.

380   Passive power spectrum, $P_U(\omega)$, is measured while trap laser is held stationary. The

381   passive power spectrum (Eq. 3) is

$$P_U(\omega) \equiv \langle |\tilde{x}_U(\omega)|^2 \rangle$$

(

3)

384   where $\tilde{x}_U(\omega)$ is the Fourier transform of the time series of the undriven bead's thermally

385   fluctuating position while trap laser is stationary.

387   With all the information from the trajectories of bead positions along with β, k, mass of

388   bead (m), and radius (a), the generalized Stokes-Einstein relationship yield the complex modulus,

389    $G*(\omega)$, shown in (Eq. 4).

$$G^*(\omega) = \frac{i\omega \tilde{\gamma}_D(\omega)}{6\pi a}$$

(

4)

391   where $\tilde{\gamma}_D(\omega)$ corresponds to the friction relaxation spectrum (Eq. 5) based on the active

392   power spectrum, $\tilde{R}_L(\omega)$.

$$\tilde{\gamma}_D(\omega) + i\omega m = -\frac{k}{i\omega}\left(\frac{1}{i\omega\tilde{R}_L(\omega)} + 1\right)$$

(

5)

395   Then, the complex modulus can be broken into elastic modulus, $G'(\omega)$, and viscous

396   modulus, $G''(\omega)$ (Eq. 6).

$$G^*(\omega) = G'(\omega) + iG''(\omega)$$

(

6)

For all the measurements, laser power was set to 100 mW at the microscope back port

while the amplitude of trap laser oscillations was set to 100nm. Experiments were controlled

using custom LabVIEW programs.

Brillouin microscopy

Larvae were anesthetized using 0.4% buffered tricaine and embedded in a dorsal

orientation in 1.25% low melting point agarose gel and allowed to polymerize in with cover glass

(no. 1.5 thickness) at 3 and 4 dpf. Dorsal orientation has been chosen to scan all brain regions

(fore-, mid-, and hind- brain) but also to avoid eye interference during data acquisition. Then,

fish water supplemented with tricaine was added to the agarose hydrogel for the entire time of

measurement. Brillouin microscopy setup has been described elsewhere[58, 59]. Briefly, the

setup is composed of 660 nm laser (Laser Quantum, #Torus-660), which illuminates the larvae

brain at 20-30mW *via* 40x air objective (Olympus, LUCPLFLN40X 0.6 NA) after passing

through the backport of microscopy body (Olympus, #IX81+DSU). Then, the backscattering

from the focused voxel is collected through a single mode fiber (Thorlabs, #P1-460Y-FC-2),

which serves as confocal pinhole. Brillouin light is then directed and analyzed by a two-stage

VIPA etalon (FSR 15 GHz, LightMachinery, #OP-6721-6743-3) spectrometer in cross-axis

configuration[60]. The VIPA spectrometer separates the frequencies of light, which is imaged on

a high-sensitivity electron-multiplying charge-coupled device (EMCCD) camera (Andor, #iXon

897). Adjustable slits in the spectrometer blocks the stray light from elastic scattering to yield

two Brillouin peaks, which are the anti-Stokes Brillouin scattering peak and the Stokes peak of

the next diffraction order. At the end, the graph of intensity versus two Brillouin peaks

(frequency) is obtained. The raw data is fitted with a Lorentzian function in a custom MATLAB

program based on nonlinear least squares fitting to localize peak centers between the two

Brillouin peaks. Brillouin microscopy measures spontaneous Brillouin scattering from the

interactions between the incident light and inherent acoustic phonons (thermal fluctuations)

inside the sample. Then, a frequency shift or Brillouin shift can be measured from the scattering.

The Brillouin shift, v_B, is calculated based on (Eq. 7).

$$v_B = \frac{2n}{\lambda} \sqrt{\frac{M'}{\rho}} \sin\left(\frac{\theta}{2}\right)$$

(

7)

where n is refractive index of the material, λ is the laser wavelength, M' is the

longitudinal elastic modulus of measured sample, ρ is the density, and θ is the collection angle

of the scattered light. For our setup, backward scattered light was collected, yielding $\theta = 180$ °.

Before actual sample measurements, Brillouin scattering calibration has been done by collecting

500 Brillouin spectra of methanol and water at 10ms exposure time. With the known literature

values of Brillouin shift for methanol and water, the effective free spectral range and the spectral

dispersion parameter (GHz per pixel) could be calculated. Calibration was done at least once an

hour throughout whole measurements. The experimental parameters for brain scanning were

done at 50-100ms exposure time and a pixel size at 2 µm x 2 µm.

Corrected Brillouin modulus was calculated based on the conversion of experimental

Brillouin shifts from each brain regions to longitudinal modulus (M'). The longitudinal modulus

is correlated to OT's complex modulus (lG*l), elastic modulus (G'), and viscous modulus (G'')

at 7Hz, 907Hz, and 15kHz based on log-log linear relationship, $\log(M') = $ a * log(lG*l, G', or

G'') + b, where the relative change in Brillouin modulus is related to the relative change in OT's

lG*l, G', or G''. Thus, the slope (a) is multiplied to log(lG*l, G', or G'') in order to get corrected

Brillouin modulus, where log(Brillouin modulus) = a * log(lG*l, G', or G'') + log(M'). During

the conversion process, density ($\rho = 1081$ g/m³) and refractive index ($n = 1.395$) values are

kept constant to understand the correlation between OT and Brillouin.

Drug treatments

- *Mysoin II inhibition*: (±)-Blebbistatin (Millipore Sigma, #203390-5MG) was diluted to

10 µM in 0.1% DMSO in PTU fish embryo water right before use to inhibit non muscle Myosin

II. Larvae were treated overnight from 3 dpf. Thus, the control was 0.1% DMSO in PTU fish

embryo water.

- *Macrophage ablation*: Metronidazole (Millipore Sigma, #M1547) was diluted to 10

mM in 0.1% DMSO in embryo medium immediately before use for nitroreductase-mediated cell

ablation. The control was 0.1% DMSO in PTU fish embryo water. As Metronidazole is light

sensitive, all treatment groups were kept in the incubator in the dark including control groups.

The treatment started at 3dpf for overnight.

-*Macrophage depletion injection*: Zebrafish larvae at 3 dpf were anesthetized using 0.4%

buffered tricaine. 1.2nl of clodronate liposomes (FormuMax, #F7010C-AH) was injected directly

into brain parenchyma (midbrain) to deplete macrophages at the brain. As a control, 1.2nl of

empty liposomes (FormuMax, #F70101-AH) was injected. Injected larvae were immediately

transferred into fresh PTU fish embryo water.

-CSF1R inhibitor: PLX5622 (Selleckchem, #S8874-5mg) was diluted to 10 µM in 0.1%

DMSO in PTU fish embryo water right before use to inhibit CSF1R. Larvae were treated

overnight from 3 dpf. Thus, the control was 0.1% DMSO in PTU fish embryo water.

Macrophage labelling

Neutral red (Millipore Sigma, # N7005) was diluted to 2.5 µg/ml in embryo medium

and administered to 3-4 dpf larvae for 2.5 h at 28.5 ˚C in the dark. Following staining,

larvae were washed two or three times in fresh PTU fish embryo water and monitored until

nonspecific tissue redness had washed out (30 - 60 min), before mounting dorsal-down in low-

melt agarose for brightfield imaging of optic tectum (midbrain) microglia.

Zebrafish immunofluorescence and brain area quantification

Larvae from 3 - 4dpf were fixed for 4 hours at 4˚C in 4% paraformaldehyde in PBS. Fixed larvae were washed three times in PBDT (PBS supplemented with 1% DMSO and 0.5% Triton X-100). Washed larvae were gone sequential dehydration to methanol and then were stored at -20˚C for at least 12˚C hours. Dehydrated larvae in methanol were sequentially hydrated in PBDT. Then, larvae were permeabilized with 10 µg/ml Proteinase K (Millipore Sigma, #3115879001) in PBDT for 15 min at room temperature to remove epidermis, followed by 4% paraformaldehyde in PBS for 30 min. Larvae were washed three times with PBDT and blocked for 1 h at room temperature in PBDT containing 5% goat serum. Larvae were then incubated in a 1:200 dilution of mouse monoclonal anti-tubulin (acetal Lys40) antibody (GeneTex, #16292) in PBDT with 5% goat serum for at least three days at 4˚C to stain acetylated tubulin. Stained larvae were washed quickly in PBDT three times and then washed an additional two times in PBDT for 15 min each. Antibody-stained larvae were then transferred to a 1:250 secondary antibody cocktail containing an AlexaFluor Plus 488 goat anti-mouse secondary antibody (Thermo Fisher Scientific, #A-11001) in PBDT with 5% goat serum. The larvae were stained for overnight at 4˚C. Then, the fish were washed three times in PBDT for 15 min each and imaged with a Zeiss 780 LSM confocal microscopy.

Confocal z-stacks were acquired at 0.5 µm steps with the pinhole diameter set at 90.1 µm. 12-bit images were acquired with a Zeiss 20x EC Plan-Apochromat, 0.8 NA objective. Samples were excited with 488 nm light from an argon laser at 2% total power of 25 mW for normalized fluorescence intensity comparison. Transmittance spectrum was also recorded. The master gain was set at or below 650. Pixel dwell times of 1.58 ms were used. Then, images were max

projected in terms of z-axis and y-axis for further image analysis by using ImageJ. Each brain

region was quantified based on acetylated tubulin area.

High-Throughput Zebrafish Imaging via Vertebrate Automated Screening

Technology (VAST)

At 2dpf, embryos were removed from the incubator Fish were put into flat-bottom 96-

well plate in 150uL fish water containing PTU. Using LP Sampler (Union Biometrica), fish were

loaded into capillary and imaged by a Vertebrate Automated Screening Technology (VAST)

BioImager (Union Biometrica). The camera of the VAST BioImager was used to quantify brain

development and morphology. The fish were scored based on development of the hindbrain.

After acquisition, the embryos were dispensed into a 96-well plate, rinsed from tricaine, and

placed back into the incubator.

Bulk RNA sequencing

RNA from embryos was extracted using TriZol (Thermo Fisher Scientific Cat No

15596026) according to manufacturer protocol. 1.25ug of Total RNA was submitted for RNA-

Sequencing to the NCI CCR Genomics Core. The libraries were made using Poly(A) Selection

kit (illumina). 150bp fragments were sequenced by paired-end 75bp utilizing Illumina NextSeq.

Reads were aligned to Ensembl Zebrafish Genome Build 11, GRCz11, using Histat21. SAMtools

was used for sorting and indexing2. PartekFlow was used for ANOVA and pathway analysis.

Statistical analysis

All data are displayed as representative or the results from multiple independent

experiments. Data comparisons were made by paired two-tailed t-test to distinguish the

heterogeneity of brain mechanics within brain regions and unpaired two-tailed t-test to determine

the drug efficacy on brain mechanics in GraphPad Prism. Error bars are expressed as mean ±

standard of error for optical tweezer data and as mean□±□standard deviation for everything else,

with * p□<□0.05, ** p□<□0.01, *** p□<□0.001 and **** p□<□0.0001.

Acknowledgements

This effort was supported by the Intramural Research Program of the National Institutes

of Health, the National Cancer Institute (KT). This work was also supported by grants from the

National Science Foundation (DBI-1942003) and the National Institutes of Health

(R21CA258008, R01HD095520) (GS). This work utilized the computational resources of the

NIH HPC Biowulf cluster. (http://hpc.nih.gov) Services were provided by the CCR Genomics

Core at National Cancer Institute. The panther zebrafish line and advice were kindly provided by

Dr. David Parichy (University of Virginia). The Tg(mpeg1.1:NTR-IRES-eGFP-CAAX) was also

kindly provided by Dr. Bruce Appel (University of Colorado, Denver.) We also thank Dr.

Mayssa Mokalled (Washington University) St. Louis for helpful discussion. All schemes are

created with BioRender.com.

References

1.   Barnes, J.M., L. Przybyla, and V.M. Weaver, *Tissue mechanics regulate brain development,*
homeostasis and disease. J Cell Sci, 2017. **130**(1): p. 71-82.
2.   Keung, A.J., K.E. Healy, S. Kumar, and D.V. Schaffer, *Biophysics and dynamics of natural and*
engineered stem cell microenvironments. Wiley Interdiscip Rev Syst Biol Med, 2010. **2**(1): p. 49-
64.
3.   Klatt, D., et al., *Noninvasive assessment of the rheological behavior of human organs using*
multifrequency MR elastography: a study of brain and liver viscoelasticity. Phys Med Biol, 2007.
52(24): p. 7281-94.
4.   Budday, S., et al., *Mechanical properties of gray and white matter brain tissue by indentation.* J
Mech Behav Biomed Mater, 2015. **46**: p. 318-30.
5.   Chaudhuri, O., et al., *Effects of extracellular matrix viscoelasticity on cellular behaviour.* Nature,
2020. **584**(7822): p. 535-546.
6.   Kasza, K.E., et al., *The cell as a material.* Current Opinion in Cell Biology, 2007. **19**(1): p. 101-107.
7.   Nobs, S.P. and M. Kopf, *Tissue-resident macrophages: guardians of organ homeostasis.* Trends in
Immunology, 2021. **42**(6): p. 495-507.
8.   Yu, F.-X., B. Zhao, and K.-L. Guan, *Hippo Pathway in Organ Size Control, Tissue Homeostasis, and*
Cancer. Cell, 2015. **163**(4): p. 811-828.
9.   Kiecker, C. and A. Lumsden, *Compartments and their boundaries in vertebrate brain*
development. Nature Reviews Neuroscience, 2005. **6**(7): p. 553-564.
10.   Daneman, R. and A. Prat, *The blood-brain barrier.* Cold Spring Harb Perspect Biol, 2015. **7**(1): p.
a020412.
11.   Schafer, D.P. and B. Stevens, *Microglia Function in Central Nervous System Development and*
Plasticity. Cold Spring Harb Perspect Biol, 2015. **7**(10): p. a020545.
12.   Louveau, A., et al., *Understanding the functions and relationships of the glymphatic system and*
meningeal lymphatics. J Clin Invest, 2017. **127**(9): p. 3210-3219.
13.   Cserép, C., et al., *Microglia monitor and protect neuronal function through specialized somatic*
purinergic junctions. Science, 2020. **367**(6477): p. 528-537.
14.   Berdowski, W.M., et al., *Dominant-acting CSF1R variants cause microglial depletion and altered*
astrocytic phenotype in zebrafish and adult-onset leukodystrophy. Acta Neuropathol, 2022.
144(2): p. 211-239.
15.   Mattiace, L.A., P. Davies, S.H. Yen, and D.W. Dickson, *Microglia in cerebellar plaques in*
Alzheimer's disease. Acta Neuropathol, 1990. **80**(5): p. 493-8.
16.   Cassetta, L. and J.W. Pollard, *Targeting macrophages: therapeutic approaches in cancer.* Nat Rev
Drug Discov, 2018. **17**(12): p. 887-904.
17.   Ruffell, B. and L.M. Coussens, *Macrophages and therapeutic resistance in cancer.* Cancer Cell,
2015. **27**(4): p. 462-72.
18.   Thompson, D.W., *On growth and form.* 1942: Cambridge Univ. Press. 116 pp.
19.   Blehm, B.H., A. Devine, J.R. Staunton, and K. Tanner, *In vivo tissue has non-linear rheological*
behavior distinct from 3D biomimetic hydrogels, as determined by AMOTIV microscopy.
Biomaterials, 2016. **83**: p. 66-78.
20.   Harlepp, S., F. Thalmann, G. Follain, and J.G. Goetz, *Hemodynamic forces can be accurately*
measured in vivo with optical tweezers. Mol Biol Cell, 2017. **28**(23): p. 3252-3260.
21.   Johansen, P.L., et al., *Optical micromanipulation of nanoparticles and cells inside living zebrafish.*
Nat Commun, 2016. **7**: p. 10974.

22.   Lieschke, G.J. and P.D. Currie, *Animal models of human disease: zebrafish swim into view.* Nat
Rev Genet, 2007. **8**(5): p. 353-67.
23.   Paul, C.D., et al., *Tissue Architectural Cues Drive Organ Targeting of Tumor Cells in Zebrafish.* Cell
Syst, 2019. **9**(2): p. 187-206 e16.
24.   Paul, C.D., et al., *Human macrophages survive and adopt activated genotypes in living zebrafish.*
Sci Rep, 2019. **9**(1): p. 1759.
25.   Schlussler, R., et al., *Mechanical Mapping of Spinal Cord Growth and Repair in Living Zebrafish*
Larvae by Brillouin Imaging. Biophys J, 2018. **115**(5): p. 911-923.
26.   White, R., K. Rose, and L. Zon, *Zebrafish cancer: the state of the art and the path forward.* Nat
Rev Cancer, 2013. **13**(9): p. 624-36.
27.   Renshaw, S.A. and N.S. Trede, *A model 450 million years in the making: zebrafish and vertebrate*
immunity. Dis Model Mech, 2012. **5**(1): p. 38-47.
28.   Staunton, J.R., B. Blehm, A. Devine, and K. Tanner, *In situ calibration of position detection in an*
optical trap for active microrheology in viscous materials. Opt Express, 2017. **25**(3): p. 1746-1761.
29.   Zhang, J., M. Nikolic, K. Tanner, and G. Scarcelli, *Publisher Correction: Rapid biomechanical*
imaging at low irradiation level via dual line-scanning Brillouin microscopy. Nat Methods, 2023.
30.   Nikolic, M., G. Scarcelli, and K. Tanner, *Multimodal microscale mechanical mapping of cancer*
cells in complex microenvironments. Biophys J, 2022. **121**(19): p. 3586-3599.
31.   Marchant, C.L., A.N. Malmi-Kakkada, J.A. Espina, and E.H. Barriga, *Cell clusters softening triggers*
collective cell migration in vivo. Nature Materials, 2022. **21**(11): p. 1314-1323.
32.   Chitnis, A. and J. Kuwada, *Axonogenesis in the brain of zebrafish embryos.* The Journal of
Neuroscience, 1990. **10**(6): p. 1892-1905.
33.   Bevilacqua, C., et al., *High-resolution line-scan Brillouin microscopy for live imaging of*
mechanical properties during embryo development. Nat Methods, 2023.
34.   Parichy, D.M., et al., *An orthologue of the kit-related gene fms is required for development of*
neural crest-derived xanthophores and a subpopulation of adult melanocytes in the zebrafish,
Danio rerio. Development, 2000. **127**(14): p. 3031-3044.
35.   Li, Q. and B.A. Barres, *Microglia and macrophages in brain homeostasis and disease.* Nat Rev
Immunol, 2018. **18**(4): p. 225-242.
36.   Ferrero, G., M. Miserocchi, E. Di Ruggiero, and V. Wittamer, *A csf1rb mutation uncouples two*
waves of microglia development in zebrafish. Development, 2021. **148**(1).
37.   Herbomel, P., B. Thisse, and C. Thisse, *Zebrafish Early Macrophages Colonize Cephalic*
Mesenchyme and Developing Brain, Retina, and Epidermis through a M-CSF Receptor-Dependent
Invasive Process. Developmental Biology, 2001. **238**(2): p. 274-288.
38.   Campinho, P., et al., *Tension-oriented cell divisions limit anisotropic tissue tension in epithelial*
spreading during zebrafish epiboly. Nat Cell Biol, 2013. **15**(12): p. 1405-14.
39.   Hughes, A.N. and B. Appel, *Microglia phagocytose myelin sheaths to modify developmental*
myelination. Nature Neuroscience, 2020. **23**(9): p. 1055-1066.
40.   Michor, F., J. Liphardt, M. Ferrari, and J. Widom, *What does physics have to do with cancer?* Nat
Rev Cancer, 2011. **11**(9): p. 657-70.
41.   So, W.Y. and K. Tanner, *Emerging principles of cancer biophysics.* Fac Rev, 2021. **10**: p. 61.
42.   Jordan, J.E.L., et al., *Microscopic multifrequency MR elastography for mapping viscoelasticity in*
zebrafish. Magn Reson Med, 2022. **87**(3): p. 1435-1445.
43.   Scarcelli, G., R. Pineda, and S.H. Yun, *Brillouin optical microscopy for corneal biomechanics.*
Invest Ophthalmol Vis Sci, 2012. **53**(1): p. 185-90.
44.   Zhang, H., et al., *Motion Tracking Brillouin Microscopy Evaluation of Normal, Keratoconic, and*
Post-Laser Vision Correction Corneas: Motion Tracking Brillouin Microscopy in Keratoconus and
Laser Vision Correction. American Journal of Ophthalmology, 2023.

45.　Bambardekar, K., et al., *Direct laser manipulation reveals the mechanics of cell contacts in vivo.*
Proc Natl Acad Sci U S A, 2015. **112**(5): p. 1416-21.
46.　Serwane, F., et al., *In vivo quantification of spatially varying mechanical properties in developing*
tissues. Nat Methods, 2017. **14**(2): p. 181-186.
47.　Thompson, A.J., et al., *Rapid changes in tissue mechanics regulate cell behaviour in the*
developing embryonic brain. eLife, 2019. **8**: p. e39356.
48.　Lu, P., K. Takai, V.M. Weaver, and Z. Werb, *Extracellular matrix degradation and remodeling in*
development and disease. Cold Spring Harb Perspect Biol, 2011. **3**(12).
49.　Petridou, N.I., Z. Spiro, and C.P. Heisenberg, *Multiscale force sensing in development.* Nat Cell
Biol, 2017. **19**(6): p. 581-588.
50.　Shook, D. and R. Keller, *Mechanisms, mechanics and function of epithelial-mesenchymal*
transitions in early development. Mech Dev, 2003. **120**(11): p. 1351-83.
51.　Vu, T.H. and Z. Werb, *Matrix metalloproteinases: effectors of development and normal*
physiology. Genes Dev, 2000. **14**(17): p. 2123-33.
52.　Hume, D.A., et al., *Phenotypic impacts of CSF1R deficiencies in humans and model organisms.*
Journal of Leukocyte Biology, 2019. **107**(2): p. 205-219.
53.　Guo, L. and S. Ikegawa, *From HDLS to BANDDOS: fast-expanding phenotypic spectrum of*
disorders caused by mutations in CSF1R. Journal of Human Genetics, 2021. **66**(12): p. 1139-1144.
54.　Cannarile, M.A., et al., *Colony-stimulating factor 1 receptor (CSF1R) inhibitors in cancer therapy.*
Journal for ImmunoTherapy of Cancer, 2017. **5**(1): p. 53.

Figure 1: *in vivo* brain mechanics show region heterogeneity revealed by optical trap active microrheology and Brillouin microscopy.

(a) Scheme of brain mechanics measurement for optical trap where 1 μm beads are directly injected into midbrain parenchyma for mechanical mapping at 3 days-post-fertilization (3dpf) / 1 day post injection (dpi) and 4dpf / 2dpi. High power trap laser oscillates the bead while the stationary low power detection laser records the bead motion, which is translated into complex modulus (G^*), elastic modulus (G'), and viscous modulus (G'') (b) Scheme of Brillouin microscopy measurement on brain parenchyma where the scattering is collected upon 660nm laser radiation (c) log-log plot of WT brain mechanics (elastic modulus, G', and viscous modulus, G'') and frequencies (7Hz to 15kHz) in terms of development (3dpf and 4dpf) and brain regions (forebrain, midbrain, and hindbrain) with crossover frequency labelled where viscous modulus becomes greater than elastic modulus; 3dpf in black and 4dpf in magenta. Error bars in standard of error (3dpf fish = 4 and 4dpf fish = 6 - 3dpf forebrain: number of beads = 78, 3dpf midbrain: n = 90, 3dpf hindbrain: n = 77, 4dpf forebrain: n = 202, 4dpf midbrain: n = 224, 4dpf hindbrain: n = 259) (d) 3dpf WT Brillouin microscopy image with Brillouin shift (GHz) in terms of 'Fore-', 'Mid-', and 'Hind-' (n = 11). ** $p<0.01$, paired two-tailed t-tests. (e) Normalized bar graph of Brillouin microscopy on WT 3dpf in respect to average Brillouin shift (GHz) of WT 3dpf hindbrain and normalized bar graph of WT 3dpf complex modulus (G^*) from optical tweezer in respect to WT 3dpf hindbrain based on 19 different frequencies. * $p<0.05$, and **** $p<0.0001$, paired two-tailed t-tests. (f) 4dpf WT Brillouin microscopy image with Brillouin shift in terms of 'Fore-', 'Mid-', and 'Hind-' (n = 12). * $p<0.05$, paired two-tailed t-tests. (g) Normalized bar graph of Brillouin microscopy on WT 4dpf in respect to average Brillouin shift (GHz) of WT 3dpf hindbrain and normalized bar graph of WT 4dpf complex modulus (G^*) from optical

tweezer in respect to WT 3dpf hindbrain based on 19 different frequencies. * p<0.05, and ****

p<0.0001, paired two-tailed t-tests.

Figure 2: Brillouin longitudinal modulus using calculated density factor shows greater

correlation for OT measurements obtained at lower frequencies in the zebrafish brain

(a) Immunofluorescence scheme of acetylated tubulin in different axis projections, z-axis and y-

axis focusing on brain parenchyma excluding eye (b) y-axis maximum projection of WT

acetylated tubulin over the development (c) z-axis maximum projection of WT acetylated tubulin

intensity excluding eye over the development (d) Quantification of normalized acetylated tubulin

intensity in terms of development and brain regions (WT 3dpf: n = 14, WT 4dpf: n = 13). *

p<0.05, *** p<0.001, **** p<0.0001, unpaired two-tailed t-tests. (e) Quantification of brain

region area based on acetylated tubulin in terms of development, xz (y-axis maximum

projection), and xy (z-axis maximum projection) (WT 3dpf: n = 14, WT 4dpf: n = 13). * p<0.05,

**** p<0.0001, unpaired two-tailed t-tests. (f) Dissociated head cells per fish based on the

cluster of 15 fish in terms of development (WT 3dpf: n = 45, WT 4dpf: n = 45). Each dot

represents a group of 15 fish head. ** p<0.01, unpaired two-tailed t-tests. (g) Macrophages at

midbrain labelled by neutral red assay for WT in terms of development (WT 3dpf: n = 15 and

WT 4dpf: n = 18). **** p<0.0001, unpaired two-tailed t-tests. (h) Raw correlation between

Brillouin (longitudinal) modulus and OT at 15kHz in terms of complex modulus, elastic modulus,

and viscous modulus for development (3-4dpf). Linear fit (top) between log(|G*|, G', and G'' for

forebrain, midbrain, and hindbrain) and log(Brillouin calculated based on estimated brain

refractive index and density) to get a slope to get corrected Brillouin modulus (bottom) in respect

to |G*|, G', and G'' (i) Density factor corrected correlation between Brillouin (longitudinal)

modulus and OT at 15kHz in terms of complex modulus, elastic modulus, and viscous modulus

for development (3-4dpf). Linear fit (top) between log(|G*|, G', and G'' for forebrain, midbrain,

and hindbrain) and log(Brillouin calculated based on estimated brain refractive index and density)

to get a slope to get corrected Brillouin modulus (bottom) in respect to |G*|, G', and G''

Figure 3: OT and Brillouin microscopy can detect changes in tissue mechanical properties

in CSF1r zebrafish mutant with delayed macrophage invasion.

(a) Macrophages at midbrain labelled by neutral red assay comparing between WT and

csf1ranull at 4dpf (WT 4dpf : n = 18 and csf1ra^{null} 4dpf : n = 18). **** p<0.0001, unpaired two-

tailed t-tests (b) Bulk RNA sequence at 2dpf and 4dpf in terms of csf1ra mutation. (c) log-log

plot of csf1ranull brain mechanics (elastic modulus, G', and viscous modulus, G'') and

frequencies (7Hz to 15kHz) in terms of development (3dpf and 4dpf) and brain regions

(forebrain, midbrain, and hindbrain) with crossover frequency labelled where viscous modulus

becomes greater than elastic modulus; 3dpf in black and 4dpf in magenta. Error bars in standard

of error (3dpf fish = 4 and 4dpf fish = 5 - 3dpf forebrain: n = 65, 3dpf midbrain: n = 88, 3dpf

hindbrain: n = 73, 4dpf forebrain: n = 141, 4dpf midbrain: n = 140, 4dpf hindbrain: n = 168). (d)

normalized complex modulus, |G*|, of csf1ra^{null} at 19 different frequencies (7Hz to 15kHz) over

the development in terms of brain regions normalized by 3dpf csf1ra hindbrain |G*|. (n=19), *

p<0.05, **** p<0.0001, paired two-tailed t-test (e) normalized complex modulus, |G*|, of WT

4dpf at 19 different frequencies in respect to 3dpf WT hindbrain |G*| to show the degree of

increase compared to csf1ra^{null}. (n=19), **** p<0.0001, paired two-tailed t-test (f) z-axis

maximum projection of csf1ra^{null} acetylated tubulin over the development and its normalized

intensity in terms of development and brain regions (3dpf n = 9 and 4dpf n = 7) (g) 4dpf

csf1ra[null] Brillouin microscopy image with Brillouin shift (GHz) in terms of brain regions

comparing with WT (WT = 12 and csf1ranull = 12). ** p<0.01, ***p<0.001, unpaired two-tailed

t-tests. (h) Normalized bar graph of Brillouin microscopy on 4dpf csf1ra[null] in respect to average

Brillouin shift (GHz) of WT 3dpf hindbrain. * p<0.05, unpaired two-tailed t-test.

Figure 4: Acetylated tubulin is a major driver of brain mechanical properties.

(a) Brillouin microscopy images for WT 4dpf control/DMSO and WT 4dpf Nocodazole

treatment. (b) Brillouin shift comparison between WT 4dpf control/DMSO and WT 4dpf

Nocodazole treatment in terms of brain regions. (control: n = 7 and Nocodazole: n = 11). ****

p<0.0001, unpaired two tailed t-tests. (c) z-axis maximum projection (xy) and y-axis maximum

projection (xz) for WT 4dpf acetylated tubulin in terms of Nocodazole treatment (d)

Quantification of normalized acetylated tubulin intensity in terms of brain regions upon

Nocodazole treatment (control: n = 12 and Nocodazole: n = 13). **** p<0.0001, unpaired two-

tailed t-tests. (e) Brillouin microscopy images for WT 4dpf control/DMSO and WT 4dpf

Blebbistatin treatment (f) Brillouin shift comparison between WT 4dpf control/DMSO and WT

4dpf Blebbistatin treatment in terms of brain regions. (control: n = 5 and Blebbistatin: n = 7). *

p<0.05, ** p<0.01, unpaired two tailed t-tests. (g) y-axis maximum projection (xz) of WT 4dpf

acetylated tubulin in terms of Blebbistatin treatment (h) Quantification of normalized acetylated

tubulin intensity in terms of brain regions upon Blebbistatin treatment (control: n = 13 and

Blebbistatin: n = 14). * p<0.05, ** p<0.01, unpaired two-tailed t-tests. (i) Macrophages at

midbrain labelled by neutral red assay for WT upon Nocodazole treatment (WT 4dpf control: n =

15 and WT 4dpf Nocodazole: n = 16). **** p<0.0001, unpaired two-tailed t-tests. (j)

Macrophages at midbrain labelled by neutral red assay for WT upon Blebbistatin treatment (WT

4dpf control: n = 17 and WT 4dpf Blebbistatin: n = 20). **** p<0.0001, unpaired two-tailed t-

tests.

Figure 5: CSF1r macrophages density regulates brain mechanics

(a) Macrophage depletion scheme by injecting nanoparticles (liposomes encapsulating

clodronate) into WT 3dpf midbrain. (b) Macrophages at midbrain labelled by neutral red assay

for WT 4dpf upon nanoparticle injection (WT 4dpf control NP: n = 20 and WT 4dpf clodronate

NP: n = 19). **** p<0.0001, unpaired two-tailed t-tests (c) Brillouin shift comparison between

WT 4dpf control NP and WT 4dpf clodronate NP in terms of brain regions. (control NP: n = 5

and clodronate NP: n = 8). * p<0.05, unpaired two tailed t-tests. (d) Macrophage depletion

scheme by using mpeg:NTR fish upon Metronidazole treatment at 3dpf (e) Macrophages at

midbrain labelled by neutral red assay at for mpeg:NTR 4dpf upon Metronidazole (control NP: n

= 9 and WT 4dpf clodronate NP: n = 10). **** p<0.0001, unpaired two-tailed t-tests. (f)

Brillouin shift comparison between 4dpf mpeg:NTR control/DMSO and 4dpf mpeg:NTR

Metronidazole in terms of brain regions. (control: n = 5 and Metronidazole: n = 5). * p<0.05, **

p<0.01, unpaired two tailed t-tests. (g) Scheme of CSF1R inhibition using small molecule drug,

PLX5622, on WT at 3dpf (h) Macrophages at midbrain labelled by neutral red assay at for WT

4dpf upon PLX5622 (control/DMSO: n = 12 and PLX5622: n = 12). **** p<0.0001, unpaired

two-tailed t-tests. (i) Brillouin shift comparison between WT 4dpf control/DMSO and WT 4dpf

PLX5622 in terms of brain regions. (control: n = 6 and PLX5622: n = 9). * p<0.05, ** p<0.01,

unpaired two tailed t-tests.

Extended Data1:

(a) circulation injection of 150kDa Dextran to look at the integrity of blood-brain barrier for WT in terms of development (b) log-log plot of complex modulus (G*) and frequencies from 7Hz to 15kHz with frequency-dependent power law fit (from 400Hz to 15kHz) (c) the exponent β of power law for WT in terms of development and brain regions (d) Normalized bar graph of WT G' (elastic modulus) based on 19 different frequencies in respect to WT 3dpf hindbrain G' (n = 19). ** p<0.01, **** p<0.0001, paired two-tailed t-tests. (e) Normalized bar graph of WT G'' (viscous modulus) based on 19 different frequencies in respect to WT 3dpf hindbrain G'' (n = 19). ** p<0.01, **** p<0.0001, paired two-tailed t-tests (f) log2 of normalized Brillouin shifts in respect to WT 3dpf hindbrain Brillouin shift (GHz) in terms of development and brain regions. * p<0.05, unpaired two-tailed t-tests.

Extended Data2:

(a) Raw correlation between Brillouin (longitudinal) modulus and OT at 7Hz in terms of complex modulus, elastic modulus, and viscous modulus for development (3-4dpf). Linear fit (top) between log(|G*|, G', and G'' for forebrain, midbrain, and hindbrain) and log(Brillouin calculated based on estimated brain refractive index and density) to get a slope to get corrected Brillouin modulus (bottom) in respect to |G*|, G', and G'' (b) Density factor corrected correlation between Brillouin (longitudinal) modulus and OT at 7Hz in terms of complex modulus, elastic modulus, and viscous modulus for development (3-4dpf). Linear fit (top) between log(|G*|, G', and G'' for forebrain, midbrain, and hindbrain) and log(Brillouin calculated based on estimated brain refractive index and density) to get a slope to get corrected

Brillouin modulus (bottom) in respect to |G*|, G', and G''. (a) Raw correlation between Brillouin

(longitudinal) modulus and OT at 907Hz in terms of complex modulus, elastic modulus, and

viscous modulus for development (3-4dpf). Linear fit (top) between log(|G*|, G', and G'' for

forebrain, midbrain, and hindbrain) and log(Brillouin calculated based on estimated brain

refractive index and density) to get a slope to get corrected Brillouin modulus (bottom) in respect

to |G*|, G', and G'' (b) Density factor corrected correlation between Brillouin (longitudinal)

modulus and OT at 907Hz in terms of complex modulus, elastic modulus, and viscous modulus

for development (3-4dpf). Linear fit (top) between log(|G*|, G', and G'' for forebrain, midbrain,

and hindbrain) and log(Brillouin calculated based on estimated brain refractive index and density)

to get a slope to get corrected Brillouin modulus (bottom) in respect to |G*|, G', and G''.

Extended Data3:

(a) The role of csf1ra on brain development at 3dpf and 4dpf (b) circulation injection of 150kDa

Dextran to look at the integrity of blood-brain barrier for csf1ra^{null} in terms of development (e)

log-log plot of csf1ra^{null} complex modulus (|G*|) and frequencies from 7Hz to 15kHz with

frequency-dependent power law fit (from 400Hz to 15kHz) (f) the exponent β of power law for

WT in terms of development and brain regions (g) y-axis maximum projection of csf1ra^{null}

acetylated tubulin intensity excluding eye over the development and quantification of normalized

acetylated tubulin intensity in terms of development and brain regions (csf1ranull 3dpf: n = 8,

csf1ra^{null} 4dpf: n = 8). **** $p<0.0001$, unpaired two-tailed t-tests. (h) normalized bar graph of

csf1ra^{null} G' (elastic modulus) based on 19 different frequencies in respect to csf1ra^{null} 3dpf

hindbrain G' (n = 19). **** $p<0.0001$, paired two-tailed t-tests. (e) Normalized bar graph of

csf1ra^{null} G'' (viscous modulus) based on 19 different frequencies in respect to csf1ra^{null} 3dpf

hindbrain G'' (n = 19). * $p<0.05$, **** $p<0.0001$, paired two-tailed t-tests

Extended Data4:

(a) y-axis maximum projection (xz) of WT 4dpf acetylated tubulin in terms of clodronate NP

injection (b) Quantification of normalized acetylated tubulin intensity in terms of brain regions

upon clodronate NP injection (control NP: n = 12 and clodronate NP: n = 10). ** $p<0.01$,

unpaired two-tailed t-tests.

(a)

(b) Acetylated tubulin / z-axis

WT 3dpf

Fore- Mid- Hind-

100 µm

x-axis

WT 4dpf

Fore- Mid- Hind-

(c) Acetylated tubulin / y-axis

WT 3dpf

Fore- Mid- Hind-

x-axis

WT 4dpf

Fore- Mid- Hind-

100 µm

(d) WT (csf1ra^WT)

Normalized intensity

WT 3dpf_forebrain
WT 3dpf_midbrain
WT 3dpf_hindbrain
WT 4dpf_forebrain
WT 4dpf_midbrain
WT 4dpf_hindbrain

(e) WT development

xz area [zprojectmax]
Area [µm²]

WT 3dpf forebrain
WT 3dpf midbrain
WT 3dpf hindbrain
WT 4dpf forebrain
WT 4dpf midbrain
WT 4dpf hindbrain

WT development

xy area [zprojectmax]
Area [µm²]

WT 3dpf forebrain
WT 3dpf midbrain
WT 3dpf hindbrain
WT 4dpf forebrain
WT 4dpf midbrain
WT 4dpf hindbrain

(f) Total cells per 1 head

csf1ra^WT 3dpf
csf1ra^WT 4dpf

(g) Macrophage (Neutral Red)

3dpf

4dpf

100 µm

Neutral Red per midbrain (n)

3dpf
4dpf

(h) Raw

3-4dpf 15kHz |G*|
Y = 0.001125*X + 9.403
R²=0.09388
log(Brillouin) vs log(G*-15kHz)

3-4dpf 15kHz G'
Y = 0.0009499*X + 9.403
R²=0.05972
log(Brillouin) vs log(G'-15kHz)

3-4dpf 15kHz G''
Y = 0.001160*X + 9.403
R²=0.1086
log(Brillouin) vs log(G''-15kHz)

3-4dpf Brillouin vs |G*| 15kHz
R²=0.2930
Brillouin modulus vs |G*| - 15kHz (Pa)

3-4dpf Brillouin vs G' 15kHz
R²=0.2028
Brillouin modulus vs G' - 15kHz (Pa)

3-4dpf Brillouin vs G'' 15kHz
R²=0.8028
Brillouin modulus vs G'' - 15kHz (Pa)

(i) Density factor considered

3-4dpf 15kHz |G*|
Y = 0.4495*X + 7.945
R²=0.7094
log(Brillouin) vs log(G*-15kHz)

3-4dpf 15kHz G'
Y = 0.4441*X + 8.072
R²=0.6182
log(Brillouin) vs log(G'-15kHz)

3-4dpf 15kHz G''
Y = 0.4384*X + 8.019
R²=0.7343
log(Brillouin) vs log(G''-15kHz)

3-4dpf Brillouin vs |G*| 15kHz
R²=0.9664
Brillouin modulus vs |G*| - 15kHz (Pa)

3-4dpf Brillouin vs G' 15kHz
R²=0.9380
Brillouin modulus vs G' - 15kHz (Pa)

3-4dpf Brillouin vs G'' 15kHz
R²=0.9722
Brillouin modulus vs G'' - 15kHz (Pa)

*:p<0.05, **:p<0.01, ***:p<0.001, ****:p<0.0001

(a) WT 4dpf   csf1ra^null 4dpf

Macrophage (Neutral Red)

100 μm

4dpf

Neutral Red per midbrain (n)

WT   csf1ra^null

****

(b) Heatmap Selected Genes log₁₀(Normalized Gene Counts)

csf1ra
csf1rb
flt1
flt4
il34
kdr
kdrl
vegfaa
vegfab

2dpf csf1ra^wt 2dpf csf1ra^null 4dpf csf1ra^wt 4dpf csf1ra^null

(c) csfr1a^null

G' G'' (Pa)

4dpf ω   3dpf ω

3dpf Forebrain G'
3dpf Forebrain G''
4dpf Forebrain G'
4dpf Forebrain G''

Frequency (Hz)

3dpf Midbrain G'
3dpf Midbrain G''
4dpf Midbrain G'
4dpf Midbrain G''

Frequency (Hz)

3dpf Hindbrain G'
3dpf Hindbrain G''
4dpf Hindbrain G'
4dpf Hindbrain G''

Frequency (Hz)

(d)

IG*| / |G*| panther 3dpf hindbrain

3dpf
4dpf

forebrain midbrain hindbrain

*   ****   ****

(e) OT csf1ra^null 4dpf

|G*| normalized by WT 4dpf hindbrain

Forebrain Midbrain Hindbrain

****   ****

(f) Acetylated tubulin

csfr1a^null 3dpf   csfr1a^null 4dpf

Fore- Mid- Hind-

Panther (csf1ra^null)

Normalized intensity

3dpf_forebrain 3dpf_midbrain 3dpf_hindbrain 4dpf_forebrain 4dpf_midbrain 4dpf_hindbrain

***   ****   ***   ***

(g) csf1ra^null 4dpf

μm

Fore- Mid- Hind-

μm

GHz

4 dpf

Brillouin shift (GHz)

WT - Forebrain WT - Midbrain WT - Hindbrain csf1ra^null Forebrain csf1ra^null Midbrain csf1ra^null Hindbrain

***   ***   **

(h) Brillouin csf1ra^null 4dpf

Shift normalized by WT 3dpf hindbrain

Forebrain Midbrain Hindbrain

*   *

*:p<0.05, **:p<0.01, ***:p<0.001, ****:p<0.0001

*:p<0.05, **:p<0.01, ****:p<0.0001

*:p<0.05, **:p<0.01

(a) Inject - 2 dpf

Inject — Mechanical mapping

2dp

3dpf
1dpi

4dpf
2dpi

beads

Trap Laser
Detection Laser

$$G^*(\omega) = G'(\omega) + iG''(\omega)$$

(b)

Scattering

660nm laser

v_B v_B (Brillouin Shift)

Intensity

v_0

v (GHz)

$$v_B = \frac{2n}{\lambda} \sqrt{\frac{M'}{\rho}} \sin\left(\frac{\theta}{2}\right)$$

(c) 3dpf - 4dpf brain mechanics: G' and G'' change

3dpf Forebrain G'
3dpf Forebrain G''
4dpf Forebrain G'
4dpf Forebrain G''

3dpf ω   4dpf ω

3dpf Midbrain G'
3dpf Midbrain G''
4dpf Midbrain G'
4dpf Midbrain G''

3dpf ω   4dpf ω

3dpf Hindbrain G'
3dpf Hindbrain G''
4dpf Hindbrain G'
4dpf Hindbrain G''

3dpf ω   4dpf ω

G' G'' (Pa)

Frequency (Hz)

(d) 3dpf

Hind-
Mid-
Fore-

μm

μm

GHz

Brillouin
WT 3dpf

Brillouin shift (GHz)

Forebrain Midbrain Hindbrain

(e) Brillouin
WT 3dpf

Shift normalized by
WT 3dpf hindbrain

Forebrain Midbrain Hindbrain

OT
WT 3dpf

IG'I normalized by
WT 3dpf hindbrain

Forebrain Midbrain Hindbrain

(f) 4dpf

Hind-
Mid-
Fore-

μm

μm

GHz

Brillouin
WT 4dpf

Brillouin shift (GHz)

Forebrain Midbrain Hindbrain

(g) Brillouin
WT 4dpf

Shift normalized by
WT 3dpf hindbrain

Forebrain Midbrain Hindbrain

OT
WT 4dpf

IG'I normalized by
WT 3dpf hindbrain

Forebrain Midbrain Hindbrain

*:p<0.05, **:p<0.01, ****:p<0.0001

*:p<0.05, **:p<0.01, ****:p<0.0001

WT 3dpf - 4dpf development: Brillouin vs OT

(a) Raw (7Hz)

(b) Density factor considered (7Hz)

(c) Raw (907Hz)

(d) Density factor considered (907Hz)

(a)

3dpf   4dpf

csfr1a wt

csfr1a null

3dpf

4dpf

■ Normal Brain Development
■ Abnormal Brain Development

(b)

csfr1a null   3dpf: 1hpi

csfr1a null   4dpf: 1hpi

DEXTRAN

100 µm

100 µm

(e)

csfr1a null
panther - 3dpf

■ forebrain
● midbrain
▲ hindbrain

$|G^*(\omega)| \propto \omega^\beta$

csfr1a null
panther - 4dpf

■ forebrain
● midbrain
▲ hindbrain

$|G^*(\omega)| \propto \omega^\beta$

(f)

● 3dpf
○ 4dpf

Power law (exponent b)

Forebrain   Midbrain   Hindbrain

(g)

csfr1a null

3dpf

csfr1a null

4dpf

100 µm

Acetylated tubulin

Panther (csf1ra null)

Normalized intensity

****

*:p<0.05, ***:p<0.001, ****:p<0.0001

(h)

■ 3dpf
■ 4dpf

G' / G' Panther$_{3dpf}$ hindbrain

****   ****

forebrain   midbrain   hindbrain

(i)

■ 3dpf
■ 4dpf

G'' / G'' Panther$_{3dpf}$ hindbrain

****   ****   *

forebrain   midbrain   hindbrain

(a)

Acetylated tubulin

WT 4dpf - Blank NP

WT 4dpf - Clodronate NP

100 µm

(b)

4dpf upon
clodronate NP injection

Normalized intensity

Forebrain$_{Blank\ NP}$
Midbrain$_{Blank\ NP}$
Hindbrain$_{Blank\ NP}$
Forebrain$_{Clodronate\ NP}$
Midbrain$_{Clodronate\ NP}$
Hindbrain$_{Clodronate\ NP}$

**

*:p<0.05, **:p<0.01