JNK signaling-mediated exocytosis coordinates epithelial cell-turnover in *Drosophila* ribosomal protein mutants

Nanami Akai[1], Tatsushi Igaki[2], and Shizue Ohsawa[1, *]

[1]Group of Genetics, Division of Biological Science, Graduate School of Science, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, Aichi, 464-8602, Japan

[2]Laboratory of Genetics, Graduate School of Biostudies, Kyoto University, Yoshida-Konoe-cho, Sakyo-ku, Kyoto 606-8501, Japan.

[*]To whom correspondence should be addressed:
Shizue Ohsawa, Ph.D.
E-mail: ohsawa.shizue.x5@f.mail.nagoya-u.ac.jp
TEL: +81-52-789-2994
FAX: +81-89-789-2995

Running Title: JNK-mediated exocytosis drives epithelial cell-turnover

Abstract

Robust tissue growth is orchestrated by the precise coordination of cell death
and cell proliferation. Our previous study found that in the developing wing
pouches of *Drosophila Minute*/+ animals, both cell death and compensatory cell
proliferation are dramatically increased, which contributes to robust growth of
mutant tissue. The induction of this cell-turnover depends on activation of JNK
signaling, although the mechanism by which JNK activation induces cell-
turnover remained unclear. Here, we show that JNK-mediated elevation of
exocytosis in dying cells is crucial for triggering cell-turnover in *M*/+ wing
morphogenesis. Mechanistically, elevated JNK signaling in dying cells
upregulates exocytosis-related genes and Wingless (Wg), leading to enhanced Wg
secretion. Furthermore, this exocytosis-mediated Wg secretion generally occurs
downstream of JNK signaling, regardless of the genetic background. Overall,
our findings provide mechanistic insights into robust tissue growth through the
orchestration of cell-turnover, which is primarily governed by JNK-mediated
exocytosis during *Drosophila Minute*/+ wing morphogenesis.

Introduction

Ribosomes are essential molecular machines responsible for protein synthesis in living cells, and thus fundamental to life. In humans, heterozygosity of genes related to ribosomal proteins or ribosomal biogenesis factors leads to a group of genetic disorders collectively termed ribosomopathies. For instance, heterozygous mutations in various ribosomal protein genes, including *RPS17* [1]*, RPS26* [2]*, RPS19* [3]*, RPS24* [4], and *RPL27* [5], are associated with Diamond-Blackfan anaemia (DBA), with some patients exhibiting tissue-specific developmental anomalies such as limb defects, cleft palate, and abnormal heart development [6,7]. In addition, heterozygous mutations in the *RpSA* gene are linked to isolated congenital asplenia (ICA), a disorder characterized by the absence of a spleen at birth [8]. However, the precise mechanisms by which a reduction in ribosomal protein gene dosage leads to these genetic disorders remain elusive.

In *Drosophila*, a series of heterozygous mutants for ribosomal protein genes, called *Minute/+* (*M/+*), exhibit a pronounced developmental delay during larval stages [9]. Despite this significant delay, *M/+* animals are essentially normal flies without noticeable morphological defects, except for the thinner bristles, suggesting that *M/+* animals exert certain mechanisms to overcome developmental perturbations caused by reduced ribosomal protein levels. It has been reported that extensive cell death [10], as well as cellular stress [10-13], is a distinctive characteristic of the *M/+* wing pouch. Our previous study [14] has shown that apoptotic cell death and the subsequent cell proliferation are dramatically increased in the *M/+* wing pouch. Blocking this cell-turnover by inhibiting cell death resulted in morphological abnormalities, indicating the essential role of cell-turnover in *M/+* wing morphogenesis. Genetic analyses have revealed that the induction of this cell-turnover depends on activation of JNK signaling. However, downstream events of JNK activation remain to be elucidated.

The coordination of cell death and proliferation through cell-cell communications is crucial for proper development and homeostasis of multicellular organisms. Apoptotic cells, for example, can secrete mitogens such as Wingless (Wg; a Wnt homolog), dpp (a BMP homolog), and Hh, which could promote the proliferation of nearby cells in the *Drosophila* epithelium [15-19]. Cell competition, a phenomenon in which cells with higher fitness ("winners") eliminate neighboring less fit cells ("losers") by inducing

cell death [20], is another aspect of cell-turnover. It has been reported that secretory

factors from prospective loser cells contribute to cell competition. For instance, the

cytokine-like ligand unpaired 3 (upd3) produced by losers promotes cell competition

between wild-type (winners) and *M/+* cells (losers) in the *Drosophila* epithelium [12].

Furthermore, Madin-Darby canine kidney cells depleted of the polarity regulator gene

scribble (*scrib*[KD] MDCK cells) secrete Fibroblast Growth Factor 21 (FGF21), which

in turn promotes cell competition with neighboring wild-type MDCK cells [21].

However, the mechanism by which these secreted factors are released from dying

cells is still not fully understood.

Here, we found that exocytosis, acting as a downstream event of JNK signaling in

dying cells, contributes to cell-turnover in the *M/+* wing pouch, which is essential for

robust wing development in *M*/+ animals. Our genetic analyses suggest that elevated

exocytosis likely facilitates the release of Wg from dying cells, which in turn

stimulates their own cell death and the proliferation of neighboring living cells.

Furthermore, we found that this exocytosis-mediated Wg release is not limited to *M*/+

context but represents a general event downstream of JNK signaling. Our

observations could provide new mechanistic insights into how dying cells coordinate

robust development through cell-cell interactions.

Results

M/+ wing pouch elevates exocytosis downstream of JNK signaling

To elucidate the downstream event of JNK signaling in the *M/+* wing pouch, we

conducted RNA sequence (RNA-seq) analyses on GFP-labeled fluorescence-activated

cell sorting (FACS)-sorted wing pouch cells from one of the *M/+* mutants *RpS3/+*,

compared to GFP-labeled wild-type control or *RpS3/+* wing pouch cells

overexpressing JNK inhibitor Puckered (Puc) (Figure S1A; the wing pouch is green-

marked oval domain that becomes the adult wing blade). Among the 1,097 genes that

were upregulated or downregulated in the *RpS3/+* wing pouch relative to the wild-

type control, we identified positive regulators of JNK signaling such as *reaper* and

pvf1, JAK/STAT targets including *Socs36E* and c*hinmo*, Gr64 gustatory receptors and

numerous genes associated with oxidative stress and DNA repair (Figure S1B, Tables

S1 and S2), which aligns with previous reports on genes upregulated in the *RpS3/+* or

RpS17/+ wing disc [12,13]. These data validate our experimental conditions for

identifying genes required for JNK-mediated events in the *M/+* wing pouch.

Mining the list of genes differentially expressed in the *RpS3/+* wing pouch cells

dependent on JNK signaling (Figure S1C and Table S3), we noticed that among the

genes associated with the "secretion by cell" GO term (Figure S1B), the

evolutionarily conserved exocytosis-related genes *unc-13*, *SNAP25,* and *cadps*

(*Calcium-dependent secretion activator*) were upregulated in a JNK-dependent

manner (2.29-fold increase, *RpS3*/+ compared to wild-type; 3.44-fold increase,

RpS3/+ compared to *RpS3*/+ + Puc) (Figure S1D). Among these exocytosis-related

genes, the upregulation of *SNAP25* and *unc-13* was consistent with the previous

reports concerning genes upregulated in *RpS3/+* or *RpS17/+* cells [12,13].

Calcium-dependent exocytosis is the process whereby cells release intracellular

contents into the extracellular space. It has been shown that *unc-13, SNAP25*, and

cadps collectively regulate the docking process of secretory vesicles to the plasma

membrane during Ca^{2+}-mediated exocytosis [22]. Specifically, UNC-13, a conserved

presynaptic protein with calcium-binding domains, interacts with syntaxin to prime

vesicles for fusion, crucial for calcium-regulated exocytosis [23]. SNAP-25, in

conjunction with syntaxin-1 and synaptobrevin, forms the pivotal SNARE complex

for neuronal exocytosis, assembling into a four-helix bundle that is essential for

drawing vesicle and plasma membranes close together to enable membrane fusion

[24,25]. Like UNC-13, CAPS/Cadps possesses conserved C-terminal domains that are

instrumental in the assembly of SNARE complexes, thus priming vesicles for Ca^{2+}-

induced exocytosis [26]. First, to investigate whether exocytosis is upregulated in the

M/+ wing pouch, we utilized the exosome marker CD63 tagged with EGFP (EGFP-

CD63) [27-30]. Additionally, we utilized Syt1 tagged with EGFP (Syt1-EGFP), a widely

used vesicle marker prevalent in both neuronal and non-neuronal cells [31-33]. We

observed a significant increase in EGFP-CD63-positive and Syt1-EGFP-positive

vesicles in the *RpS3/+* wing pouch compared to the wild-type control (Figures 1A, 1B,

1G and 1H, corresponding to the boxed region in Figure 1E, quantified in Figures 1F

and 1K). EGFP-CD63-positive vesicles were also found to be increased in the

RpL19/+ wing pouch (Figure S1E, quantified in Figure S1F). Notably, these vesicles

were predominantly observed in regions of the *RpS3/+* wing pouch undergoing

massive cell death (as indicated by arrowheads in Figures 1B, 1H and S1E), a

phenomenon we previously reported [14] and also confirmed in this study (shown in

Figure 2H, compared to Figure 2G, quantified in Figure 2T). Furthermore, we noted

an increase in vesicles positive for the ESCRT protein Hrs, an additional exosome

marker crucial for exosome secretion [34,35], in the *RpS3/+* wing pouch relative to the

wild-type control (Figures S1G and S1H, quantified in Figure S1I). To further

investigate the enhanced exocytosis in the *RpS3/+* wing pouch, we utilized the

GCaMP6m fluorescent calcium reporter [36] to monitor Ca^{2+} activity, a known inducer

of exocytosis. We observed persistent Ca^{2+} activity in morphologically dying cells,

along with spontaneous Ca^{2+} flashes similar to those in the wild-type control (Figures

1L and 1M, quantified in Figure 1P). This persistent activity suggested an enhanced

exocytic process in dying cells in the *M/+* wing pouch, potentially driven by elevated

calcium levels.

We then examined whether exocytosis is the downstream event of JNK

activation in the *M/+* wing pouch. We found that blocking JNK signaling by

overexpressing Puc significantly suppressed the increased number of EGFP-

CD63/Syt1-EGFP-positive vesicles, as well as the emergence of cells exhibiting

persistent Ca^{2+} signaling (Figure 1C, quantified in Figure 1F; Figure 1I, quantified in

Figure 1K; Figure 1N, quantified in Figure 1P). Together, these observations suggest

that exocytosis is elevated in dying cells within the *M/+* wing pouch as a downstream

event of JNK signaling.

160 Exocytosis is required for massive cell-turnover in the *M/+* wing pouch

We have previously demonstrated that JNK signaling is required for cell-turnover in

the *M/+* wing pouch [14]. Blocking JNK signaling by overexpressing Puc significantly

reduced cell death and mitoses in the *RpS3/+* wing pouch, as assayed by the CD8-

PARP-Venus probe for caspase activity in the imaginal disc [37-44], and the M phase

marker phospho-Histone H3 (Figures 2A-2C, quantified in Figure 2F; Figures 2G, 2H

and S2A, quantified in Figure S2N). In addition, we intriguingly found that

downregulating *unc-13*, a docking factor involved in exocytosis, similarly diminished

cell death and mitoses (Figures 2I and S2B, quantified in Figures 2T and S2N,

respectively; Figure 2D, quantified in Figure 2F), and also reduced EGFP-CD63-

positive and Syt1-EGFP-positive vesicles in the *RpS3/+* wing pouch (Figures 1D and

1J, quantified in Figures 1F and 1K, respectively). Knocking down *rab3*, a "Secretory

GTPase Rabs" required for vesicle exocytosis, also significantly reduced cell death

and mitoses in the *RpS3/+* wing pouch (Figures 2J and S2C, quantified in Figures 2T

and S2N, respectively; Figure 2E, quantified in Figure 2F). Furthermore,
overexpressing Puc or downregulating either *unc-13* or *rab3* led to morphological
defects in the *RpS3/+* adult wing, while their individual overexpression or
downregulation did not affect wing development (Figures 2O, 2P and S2O). These
results suggest that exocytosis, as a downstream component of JNK signaling, is
pivotal for cell turnover and subsequent normal morphogenesis in the *M/+* wing
pouch. Indeed, downregulating other docking factors such as *SNAP25* or *Syt1* [45],
another Secretory GTPase *Rab27*, or *ALIX*, an ESCRT (endosomal sorting complexes
required for transport)-related protein involved in the biogenesis of extracellular
vesicles [46,47], also significantly suppressed cell death in the *RpS3/+* wing pouch
(Figure 2K, quantified in Figure 2T; Figures S2D-S2J, quantified in Figure S2N).
Additionally, downregulating factors such as $G\alpha q$, *Plc21C*, and *norpA,* which are
required for intracellular Ca^{2+} release that triggers exocytosis process [32], also
significantly suppressed cell death in the *RpS3*/+ wing pouch (Figures 2L-2N,
quantified in Figure 2T; Figures S2K-S2M, quantified in Figure S2N). Together,
these data reinforce the vital role of exocytosis in cell turnover within the M/+ wing
pouch, functioning as a downstream process of JNK signaling. Intriguingly, blocking
Dronc activity by overexpressing Dronc[DN] suppressed the elevation of exocytosis in
the *RpS3*/+ wing pouch (Figure 2S, compared to Figure 2R, quantified in Figure 2U).
This suggests the presence of a signal amplification loop between exocytosis and
Dronc within the JNK signaling pathway, consistent with previous studies showing
Dronc-JNK amplification loop that induce cell death [48,49].

Dying cells secrete Wg via exocytosis in the *M/+* wing pouch
We next sought to identify the protein secreted through exocytosis from dying
cells in the *M/+* wing pouch. Our previous genetic analyses [14] suggested that cell-
turnover in the *M/+* wing pouch is initiated by a mechanism akin to Wg-dependent
cell competition, a phenomenon by which cells with higher Wg signaling activity
eliminate neighboring cells with lower Wg signaling activity in the wing disc [50].
Indeed, as demonstrated through previous genetic manipulations [14], and confirmed
here, genetically reducing the Wg signaling gradient in the entire *M/+* pouch
significantly suppressed cell death (Figure S3A, quantified in Figure S3B).
Additionally, it has been shown that apoptotic cells produce Wg through JNK

signaling [18,51,52]. These observations led us to hypothesize that dying cells in the *M/+*

wing pouch might release Wg via JNK-mediated exocytosis, potentially conferring a

survival advantage to their neighboring living cells. Supporting this hypothesis, it has

been reported that Wnt/Wg can be released through extracellular vesicles including

exosome [28,53-56]. We found that GFP-Wg-positive puncta, derived from a knock-in

allele [34], were more abundant in the *RpS3/+* wing pouch compared to the wild-type

control (Figures 3A and 3B). This increase in GFP-Wg-positive puncta, particularly in

the area with massive cell death within the *RpS3*/+ wing pouch (Figures S3D-S3E''',

compared to Figure S3C), was more evident when using a membrane-tethered anti-

GFP nanobody (Vhh4-CD8), which immobilizes GFP-Wg on the cell surface [34]

(Figures 3F and 3G, quantified in Figure 3J). Intriguingly, these GFP-Wg-positive

puncta, observed both without and with the Vhh4-CD8 morphotrap, sometimes

colocalized with CD63-mCherry-positive vesicles (Figures 3E and S3G, indicated by

arrowheads), suggesting that GFP-Wg might be secreted via extracellular vesicles

through JNK-mediated exocytosis. Supporting this possibility, we found that blocking

JNK signaling by overexpressing Puc or knocking down *unc-13* significantly reduced

the number of GFP-Wg-positive puncta in the *RpS3/+* wing pouch (Figures 3H and 3I,

quantified in Figure 3J). In addition, we observed a significant increase in

extracellular GFP-positive puncta and extracellular CD63-mCherry-positive vesicles

in the *RpS3/+* wing pouch when GFP-Wg was captured on the cell surface using the

Vhh4-CD8 morphotrap, compared to the wild-type control (Figures S3H and S3I,

quantified in Figures S3J and S3K). Furthermore, these extracellular GFP-Wg-

positive puncta occasionally colocalized with extracellular CD63-mCherry-positive

vesicles (6.88% in the wild-type, and 16.45% in the *RpS3/+* wing pouch region,

characterized by massive cell death, as indicated by the arrowheads in Figures S3H

and S3I). Together, these data suggest that dying cells secrete Wg via exocytosis in

the *M/+* wing pouch.

JNK signaling generally triggers exocytosis-mediated Wg secretion

Finally, we investigated whether JNK activation generally upregulates

exocytosis and subsequent Wg secretion in contexts beyond the *M/+* wing pouch. We

generated JNK-activating clones by overexpressing Eiger, the *Drosophila* homolog of

tumor necrosis factor (TNF), in the eye-antennal discs. Although dying cells were

observed in Eiger-overexpressing clones (Figure 4G, compared to Figure 4F), the

tissue size remained unchanged (data not shown), suggesting that neighboring cells

proliferate to maintain tissue homeostasis. We found that the elimination of Eiger-

expressing clones and the number of dying cells were significantly suppressed by

knocking down *unc-13* in these clones (Figures 4D and 4H, quantified in Figure 4E

and 4I), whereas unc-13-RNAi expression alone did not affect tissue growth (Figures

4A and 4C, quantified in Figure 4E). In addition, we observed a significant increase in

Hrs-positive vesicles and GFP-Wg-positive puncta, derived from a knock-in allele [34],

in Eiger-overexpressing clones compared to the control (Figure 4G' and 4L''),

quantified in Figure 4J and 4M). Furthermore, the elimination of Eiger-expressing

clones was significantly suppressed when the Wg signaling gradient was reduced

either by deleting one copy of the *wg* gene (Figure 4N-4, quantified in Figure 4Q),

suggesting the intriguing possibility that exocytosis-mediated Wg secretion from

dying cells may be essential for both their cell death and proliferation of neighboring

cells. Together, these data suggest that JNK signaling generally triggers exocytosis

and the subsequent Wg secretion, which is crucial for maintaining tissue homeostasis.

Discussion

Our current study reveals that exocytosis is crucial for massive cell-turnover in

M/+ wing morphogenesis, acting as a downstream event of JNK signaling. In dying

cells, JNK signaling upregulates the expression of exocytosis-related genes and Wg,

leading to increased Wg secretion. In addition, elevated exocytosis induces cell death

via an exocytosis-Dronc amplification loop (Figure 5). Notably, impairment of

exocytosis in dying cells in *RpS3/+* animals leads to phenotypic variations in adult

wings (Figures 2O and 2P in this study), suggesting that increased exocytosis in dying

cells, followed by subsequent cell-turnover, may provide the flexibility needed to

adjust developmental programs.

We identified Wg as an important factor secreted through exocytosis from dying

cells, potentially essential for the proliferation of neighboring living cells to maintain

tissue homeostasis. Previous studies have shown that 'undead' cells, stimulated

toward apoptosis but kept alive by the caspase inhibitor p35, are able to promote the

proliferation of neighboring cells in the *Drosophila* epithelium [57-59]. Several mitogens

including Wg, Dpp, and Spitz (a EGF homolog) have been identified as being

produced by apoptotic cells in a JNK-dependent manner [18,51,60]. In addition, Hedgehog

(Hh) [15] have also been identified as products of apoptotic cells. Furthermore, during

cell competition between wild-type (winners) and *M/+* cells (losers) in the *Drosophila* epithelium*,* prospective loser cells secrete upd3, which promotes competitive elimination by winner cells [12]. Hence, in addition to Wg, other mitogens could also be secreted thorough exocytosis from dying cells, potentially contributing to massive cell-turnover in the *M/+* wing pouch.

Previous studies have identified exosomes as carriers of Wnt/Wg in the extracellular space of both mammalian and *Drosophila* cells, including wing disc cells [28,53-56,61]. Concurrently, alternative mechanisms for Wg transport, such as those involving lipoprotein particles or a lipocalin Swim in the *Drosophila* wing disc, have been reported [29,62]. Additionally, the cell-surface proteoglycan Dally-like-protein (Dlp) has been reported to enable long-range signaling of the palmitoylated Wg [34]. Intriguingly, Wnt/Wg transport has also been observed through filopodia-like cellular extensions known as cytonemes [63-66]. In our study of the *M/+* wing disc, a model characterized by massive cell-turnover, we observed partial colocalization of Wg-GFP puncta with exosomal markers such as CD63-mCherry and Hrs. Our data also suggest that Wg secretion through exocytosis may not uniformly occur among all dying cells within this context. This disparity suggests that while exosomes from dying cells significantly contribute to Wg transport within the *M/+* wing pouch, other pathways may also be operative.

Rp genes are crucial in a broad range of organisms, from yeast to humans, frequently exhibiting haploinsufficient mutant phenotypes. Notably, heterozygous mutations in genes encoding ribosomal proteins or ribosomal biogenesis factors are linked to tissue-specific human diseases known as ribosomopathies [67]. In patients affected by these conditions, some tissues may preserve normal patterning and functions, potentially by buffering stresses caused by ribosomal mutations. Given the evolutionary conservation of the molecules we identified in *Drosophila*, regulating cell turnover through exocytosis in dying cells may represent evolutionarily conserved strategy to prevent developmental abnormalities in ribosomal protein mutants.

Author Contributions

Conceptualization: Nanami Akai, Tatsushi Igaki, Shizue Ohsawa.
Data curation: Nanami Akai, Tatsushi Igaki, Shizue Ohsawa.

Formal analysis: Nanami Akai.

Funding acquisition: Shizue Ohsawa, Tatsushi Igaki.

Investigation: Nanami Akai, Shizue Ohsawa.

Methodology: Nanami Akai.

Project administration: Shizue Ohsawa.

Resources: Shizue Ohsawa.

Supervision: Shizue Ohsawa.

Visualization: Nanami Akai, Tatsushi Igaki, Shizue Ohsawa.

Writing – original draft: Shizue Ohsawa, Nanami Akai.

Writing – review & editing: Nanami Akai, Tatsushi Igaki, Shizue Ohsawa.

Acknowledgments

We thank Takefumi Kondo, Keisuke Ikawa, Kiichiro Taniguchi, Rina Nagata, and
Yukari Sando for discussions, Susumu Tsutsumi, Nana Watanabe, Mina Hoshino,
Tomoko Furukawa, and Sayako Suzuki for technical support, and NGS core facility
of the Graduate Schools of Biostudies, Kyoto University for supporting the RNA-seq
analysis. We also thank Yash Hiromi, Gary Struhl, Maoto Sato, Jean-Paul Vincent,
Sharad Kumar, Konrad Basler, the Bloomington *Drosophila* Stock Center (Indiana),
the Vienna *Drosophila* Resource Center (Vienna), the National Institute of Genetics
Stock Center (Mishima), and the *Drosophila* Genomics and Genetic Resources
(Kyoto) for fly stocks. Cell sorting using BD FACS Aria II were performed at the
Medical Research Support Center, Graduate School of Medicine, Kyoto University,
supported by Basis for Supporting Innovative Drug Discovery and Life Science
Research (BINDS) from AMED (Grant No. JP22ama121034). This work was
supported in part by MEXT Grant-in-Aid (KAKENHI) for Transformative Research
Area (A) (Grant Nos. 20H05945 to SO, 21H05284 to TI), the Scientific Research (B)
(Grant No. 22H02616 to SO), and Challenging Exploratory Research (Grant No.
21K19257 to SO), Japan Science and Technology Agency (Moonshot Research &
Development: Grant Number JPMJPS2022 to SO), AMED-CREST, Japan Agency
for Medical Research and Development (22gm1710002h0001 to TI), and Japan
Agency for Medical Research and Development (21gm5010001 to TI) .

Declaration of interests

343   The authors declare no competing interests.

Figure legends

Figure 1. *M/+ wing pouch elevates exocytosis downstream of JNK signaling*

(A-D'') The exosome marker EGFP-CD63 was expressed in the wing pouch of wild-type (A), *RpS3/+* (B), *RpS3/+, nub-Gal4, UAS-Puc* (C), or *RpS3/+, nub-Gal4, UAS-unc-13-RNAi* (D) flies (white). The images correspond to the boxed area in the schematic diagram of the wing disc (E). Dying cells in the wing disc were visualized by anti-cleaved Dcp-1 staining (white). EGFP-CD63-positive vesicles were frequently observed in the area of morphological dying cells in the *RpS3/+* wing pouch (Indicated by Orange arrowheads). Scale bar, 50 μm.

(F) Boxplot with dots representing the number of EGFP-CD63-positive puncta in the pouch of genotypes shown in (A) (n=13, number of wing pouches), (B) (n=14), (C) (n=16), and (D) (n=13). Error bars, SEM; ***, p<0.001; Wilcoxon rank-sum test.

(G-J'') The vesicle marker Syt1-EGFP was expressed in the wing pouch of wild-type (G), *RpS3/+* (H), *RpS3/+, nub-Gal4, UAS-Puc* (I), or *RpS3/+, nub-Gal4, UAS-unc-13-RNAi* (J) flies (white). The images correspond to the boxed area in the schematic diagram of the wing disc (E). Dying cells in the wing disc were visualized by anti-cleaved Dcp-1 staining (white). Syt1-EGFP-positive vesicles were frequently observed in the area of morphological dying cells in the *RpS3/+* wing pouch (Indicated by Orange arrowheads). Scale bar, 50 μm.

(K) Boxplot with dots representing the number of Syt1-EGFP-positive puncta in the wing pouch of genotypes shown in (G) (n=15), (H) (n=12), (I) (n=14), and (J) (n=16). Error bars, SEM; ***, p<0.001; Wilcoxon rank-sum test.

(L-N) Time-lapse imaging of Ca^{2+} signaling in cultured wing discs. The calcium reporter GCaMP6m (shown in pseudo color) was expressed in the wing pouch of wild-type (L), *RpS3/+* (M), or *RpS3/+, nub-Gal4, UAS-Puc* (N) flies. Images were acquired at 10-second intervals. White arrowheads indicate the persistent activation of Ca^{2+} signaling. White asterisks indicate spontaneous Ca^{2+} flashes. The images correspond to the boxed area in the schematic diagram of the wing disc (O). Scale bar, 5 μm.

(P) Quantification of cells exhibiting continuous Ca^{2+} activity for 20 min in the wing pouch. The respective genotypes are shown in (L) (n=12), (M) (n=10), and (N) (n=9). These images are Z-stacked images from 3 confocal images (4μm). Error bars, SEM; **, p<0.01; Wilcoxon rank-sum test.

Figure 2. Exocytosis is required for massive cell-turnover in the *M/+* wing pouch

(A-E) Wing discs of wild-type (A), *RpS3/+* (B), *RpS3/+, nub-Gal4, UAS-Puc* (C),

RpS3/+, nub-Gal4, UAS-unc-13-RNAi (D), or *RpS3/+, nub-Gal4, UAS-Rab3-RNAi*

(E) flies were stained with anti-phospho-histone H3 (pH3) (Ser10) antibody (white).

Wing pouches were marked in pale green. Scale bar, 100 μm.

(F) Boxplot with individual dots representing pH3-positive areas in the pouch of

genotypes shown in (A) (n=18), (B) (n=22), (C) (n=23), (D) (n=12), and (E) (n=21).

Error bars, SEM; ***, $p<0.001$; Wilcoxon rank-sum test.

(G-N) The activated-caspase-3 indicator CD8-PARP-Venus was expressed in the

wing pouch of wild-type (G), *RpS3/+, nub-Gal4, UAS-lexA-RNAi* (H), *RpS3/+, nub-*

Gal4, UAS-unc-13-RNAi (I), *RpS3/+, nub-Gal4, UAS-Rab3-RNAi* (J), *RpS3/+, nub-*

Gal4, UAS-SNAP25-RNAi (K), *RpS3/+, Gαq/+* (L), *RpS3/+, nub-Gal4, UAS-Plc21C-*

RNAi (M), or *RpS3/+, norpA/+* (N) flies, and dying cells in the wing pouch were

visualized by anti-cleaved PARP staining (white). Orange arrowheads indicate

massive cell death in the *RpS3/+* wing pouch. Scale bar, 100μm.

(O) Adult wing phenotypes were classified as following five types: (a) normal, (b)

vein (weak) (weakly bearing a brunched vein in the point of "landmark 14" (arrow) [68]

(c) vein (strong) (bearing other additional vein phenotypes (arrowheads) in addition to

"landmark 14"), (d) notch, and (e) severe.

(P) The rate of defective wings in the genotypes of wild-type (n=55), *RpS3/+* (n=44),

RpS3/+, nub-Gal4, UAS-Puc (n=63), *RpS3/+, nub-Gal4, UAS-unc-13-RNAi* (n=101),

and *RpS3/+, nub-Gal4, UAS-Rab3-RNAi* (n=99).

(Q-S) The exosome marker CD63-mCherry was expressed in the wing pouch of wild-

type (Q), *RpS3/+* (R), or *RpS3/+, nub-Gal4, UAS-Dronc^{DN}* (S) flies (white). The

images correspond to the area enclosed by rectangle 1 in the schematic diagram of the

wing disc (Figure 3C). Scale bar, 20 μm.

(T) Boxplot with individual dots representing the number of cleaved-PARP-positive

dying cells per pouch in genotypes shown in (G) (n=18), (H) (n=24), (I) (n=18), (J)

(n=13), (K) (n=13), (L) (n=15), (M) (n=16), and (N) (n=22). Error bars, SEM; ***,

$p<0.001$, **, $p<0.01$, *, $p<0.05$; Wilcoxon rank-sum test.

(U) Boxplot with dots representing the number of CD63-mCherry-positive puncta in the pouch of genotypes shown in (Q) (n=11, number of wing pouches), (R) (n=11), and (S) (n=14). Error bars, SEM; ***, p<0.001, **, p<0.01; Wilcoxon rank-sum test.

Figure 3. Dying cells secrete Wg via exocytosis in the *M/+* wing pouch

(A-B'') Wing discs of *GFP-Wg/+* (A), or *RpS3/+, GFP-Wg/+* (B) flies, and dying cells in the wing pouch were visualized by anti-cleaved Dcp-1 staining (white). GFP-Wingless was visualized by anti-GFP staining (white). Arrowheads indicate massive cell death in the *RpS3/+* wing pouch. Scale bar, 100 μm.

(D-E''') Immunofluorescent localization of the exosome marker CD63-mCherry, as visualized by anti-dsRed staining (magenta), and GFP-Wingless (white) in the wing pouch of *GFP-Wg/+* (D), or *RpS3/+, GFP-Wg/+* (E) flies expressing CD63-mCherry. The nuclei were visualized by DAPI staining (blue). The images correspond to the area enclosed by rectangle 1 in the schematic diagram of the wing disc (C). Orange arrowheads indicate the colocalization of GFP-Wingless-positive puncta with CD63-mCherry-positive puncta. Scale bar, 20μm.

(F-I') The membrane-tethered anti-GFP nanobody (Vhh4-CD8-HA) was expressed in the wing pouch of *GFP-Wg/+* (F), *RpS3/+, GFP-Wg/+* (G), *RpS3/+, GFP-Wg/+, nub-Gal4, UAS-Puc* (H), or *RpS3/+, GFP-Wg/+, nub-Gal4, UAS-unc-13-RNAi* (I) flies. GFP-Wingless was visualized by anti-GFP staining (white). Dying cells were visualized by anti-cleaved Dcp-1 staining in the wing discs (white). The images correspond to the area enclosed by the rectangle 2 in the schematic diagram of the wing disc (C). Immobilized Wg-GFP-positive puncta were frequently observed in the area of morphological dying cells in the *RpS3/+* wing pouch (Indicated by orange arrowheads). Scale bar, 50μm.

(J) Boxplot with individual dots representing the number of GFP-Wingless-positive puncta per rectangle-2 in the wing pouch (C). The respective genotypes are shown in (F) (n=9), (G) (n=14), (H) (n=11), and (I) (n=11). Error bars, SEM; ***, p<0.001, **, p<0.01; Wilcoxon rank-sum test.

Figure 4. JNK signaling generally triggers exocytosis-mediated Wg secretion

(A-D) Eye disc bearing eyFLP-induced MARCM clones of *UAS-yellow-RNAi* (A),

UAS-Eiger^W +*UAS-yellow-RNAi* (B), *UAS-unc-13-RNAi* (C), or *UAS-Eiger^W* +*UAS-*

unc-13-RNAi (D) cells. Scale bar, 100μm.

(E) Boxplot with individual dots representing the clone size (% of total clone area per

eye disc area) in genotypes shown in (A) (n=23), (B) (n=32), (C) (n=26), and (D)

(n=17). Error bars, SEM; ***, p<0.001; Wilcoxon rank-sum test.

(F-H') Eye disc bearing eyFLP-induced MARCM clones of *wild-type* (F), *UAS-*

Eiger^W (G), or *UAS-Eiger^W* +*UAS-unc-13-RNAi* (H) cells stained with anti-cleaved

Dcp-1 and anti-Hrs antibody (magenta). Scale bar, 20μm.

(I) Boxplot with individual dots representing the number of cleaved-Dcp-1-positive

dying cells per clone area in genotypes shown in (F) (n=14, number of clones), (G)

(n=15), and (H) (n=15). Error bars, SEM; ***, p<0.001; Wilcoxon rank-sum test.

(J) Boxplot with individual dots representing the relative Hrs-positive area in the eye

disc for genotypes shown in (F') (n=12, number of clones), (G') (n=14), and (H')

(n=13). Error bars, SEM; ***, p<0.001; Wilcoxon rank-sum test.

(K-L''') Eye disc bearing eyFLP-induced MARCM clones of *GFP-Wg/+* (K), or

UAS-Eiger^W, *GFP-Wg/+* (L) cells stained with anti-GFP antibody (white). Scale bar,

20μm.

(M) Boxplot with individual dots representing the relative GFP-Wg area per clone

area of the eye disc for genotypes shown in (K) (n=11), and (L) (n=9). Error bars,

SEM; ***, p<0.001; Wilcoxon rank-sum test.

(N-P) Eye disc bearing eyFLP-induced MARCM clones of wild-type (N), *UAS-*

Eiger^{KB} (O), or *UAS-Eiger^{KB}*, *wg/+* (P) cells. Scale bar, 100μm.

(Q) Boxplot with individual dots representing the clone size (% of total clone area

per eye disc area) in genotypes shown in (N) (n=12), (O) (n=11), and (P) (n=7). Error

bars, SEM; ***, p<0.001, **, p<0.01; Wilcoxon rank-sum test.

Figure 5. A model for ensuring robust coordination of tissue growth in *M/+*

animals

During the larval period of *M/+* animals, JNK signaling is activated in the *M/+* wing

pouch, which induces the expression of exocytosis-related genes and Wg, leading to

the secretion of Wg through exocytosis in dying cells. In addition, elevated exocytosis

induces cell death via an exocytosis-Dronc amplification loop. This process

subsequently promotes the proliferation of neighboring cells, establishing their

apoptotic/proliferative status and resulting in significant cell turnover in the $M/+$ wing

pouch.

STAR Methods

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to and

will be fulfilled by the lead contact, Shizue Ohsawa

(ohsawa.shizue.x5@f.mail.nagoya-u.ac.jp).

Materials availability

Drosophila lines generated in this study are available from the lead contact without

restriction.

Experimental model and subject detail

Fly strains

Fly stocks were cultured at 25°C on standard fly food. To detect dying cells in the

wing pouch, the following strains were used: *nub-Gal4; UAS-CD8-PARP-Venus*

(control); *nub-Gal4; UAS-CD8-PARP-Venus, RpS3^{Plac92}* (BL5627)*/TM6B* (*RpS3*/+

tester)*. Additional strains were used as follows: *UAS-EGFP.CD63* (BL91390), *UAS-*

CD63.mCherry (BL91389), *UAS-Syt1-eGFP* (BL6925), *RpL19^{K03704}* (DGRC102285),

20XUAS-IVS-GCaMP6m (BL42750), *UAS-unc-13-RNAi* (VDRC33609, NIG2999R-

2), *UAS-LexA-RNAi* (BL67947), *UAS-Rab3-RNAi* (NIG7576R-3, BL31691), *UAS-*

SNAP25-RNAi (HMS01367, BL27306), *UAS-Syt1-RNAi* (NIG3139R-1, BL31668),

UAS-Rab27-RNAi (NIG14791R-2, BL31887), *UAS-ALiX-RNAi* (HMS00298,

VDRC32049), *Gαq^{221C}* (BL30744), *UAS-Gαq-RNAi* (NIG17759R-2), *UAS-Plc21C-*

RNAi (NIG4574R-2, 4574R-3), *norpA^{7}* (DGRC108362)*, UAS-norpA-RNAi*

(VDRC105676), *UAS-y-RNAi* (NIG3757R-1), *wg^{l-8}* (BL5351), *UAS-Puc* [68], *UAS-*

Dronc^{DN}(Sharad Kumar) [69]*, GFP-Wingless^{S239A}* (JP. Vincent) [34], *UAS-VHH4-CD8-HA*

(JP. Vincent) [34], *UAS-Eiger^{+W}* [70], and *UAS-Eiger^{KB}* (Konrad Basler) [71].

Method details

Histology

Wandering third instar larvae were dissected in phosphate buffered saline (PBS) and

fixed with 4%-Paraformaldehyde (PFA) for 5 min on ice, followed by 20 min at room

temperature. The fixed larvae were washed 3 times with PBT (PBS + 0.1% TritonX100) for 20 min each, and were then blocked with PBTn (5% donkey serum + PBT) solution for 30 min. For immunostaining, they were incubated overnight at 4°C with primary antibodies in PBTn. After four 30-min washes with PBT, the samples were re-blocked in PBTn for 30 minutes, then incubated with secondary antibodies for 2 hours at room temperature. Following another set of four 30-minute washes with PBT, the samples were mounted using either SlowFade Diamond Antifade Mountant (Invitrogen, #S36972) or an alternative anti-fade mounting medium. The latter was composed of 70% Glycerol (Sigma, #G5516), 0.2% n-Propyl Gallate (KANTO CHEMICAL, #32465-31), and supplemented with DAPI (Sigma, #D9542). For extracellular immunostaining, larvae were dissected in ice-cold Schneider's *Drosophila* Medium (Gibco, #21720024) and incubated with primary antibodies for 2 hours on ice. The larvae were then rinsed with ice-cold Schneider's *Drosophila* Medium followed by PBS and fixed with 4% PFA for 40 min at room temperature. After fixation, samples were washed with PBS. Subsequent processing was the same as in the preceding protocol. Images were taken with Leica SP5, Leica SP8, Leica SPE, Leica STELLARIS and ZEISS LSM900 confocal microscopes. Primary antibodies used are as follows; rat anti-GFP antibody (Nacalai Tesque, #04404-26, 1:1000) (Extracellular GFP, 1:300), rabbit anti-cleaved PARP (Asp214) antibody (Cell Signaling Technology (CST), #9541, 1:200), rabbit anti-phospho-histone H3 (Ser10) antibody (CST, #9706, 1:100), rabbit anti-DsRed antibody (Takara Bio Clontech (CLN), #632496, 1:250) (Extracellular DsRed, 1:75), mouse anti-Hrs antibody (DSHB, #27-4-s, 1:250), and rabbit anti-Cleaved *Drosophila* Dcp-1 (Asp216) antibody (CST, #9678, 1:100) for 3 days. Secondary antibodies used are as follows; Goat anti-rabbit Alexa 488 (1:250, Invitrogen #A11034), Goat anti-rabbit Alexa 546 (1:250, Invitrogen #A11035), Goat anti-rat Alexa 488 (1:250, Invitrogen #A11006), Goat anti-chicken Alexa 488 (1:250, Invitrogen, #A11039), and Goat anti-mouse Alexa 647 (1:250, Invitrogen #A21237) antibodies.

Quantification of puncta number

The vesicle marker EGFP-CD63, CD63-mCherry, and Syt1-eGFP were expressed in the wing pouch under the control of *nub-gal4 driver*. In the region where massive cell death is observed in the *M/+* wing disc, these markers, in addition to GFP-Wingless signals, were acquired at the confocal plane exhibiting the highest signal intensity.

CD63-mCherry were also expressed within the mosaic clones of the eye-antenna disc.

Puncta were manually counted using Fiji software and the data were analyzed with

KaleidaGraph (Synergy Software).

Quantitative Analysis of GFP-Wingless and CD63-mCherry Colocalization

To analyze the overlap between GFP-Wingless-positive puncta and CD63-mcherry-

positive vesicles, their colocalization was manually quantified in the regions

exhibiting massive cell death. This quantification was conducted using the confocal

plane that displayed the highest signal intensity. Images were manually thresholded to

eliminate background signals, and the areas of overlap were calculated using Fiji

software, followed by further analyses with KaleidaGraph.

Calcium Time-lapse imaging and quantification

To monitor Ca^{2+} activity, wandering third instar larvae were dissected and cultured in

Schneider's *Drosophila* medium supplemented with 2% FBS, 1% Insulin solution

(Sigma-Aldrich, #I0516), 3% fly extract (VDRC, #VFE1.25), and 2% low-melting

agarose (Sigma-Aldrich, #A4018)/PBS. Imaging was conducted at 10 sec intervals for

20-min with a Leica STELLARIS confocal microscope [42,72]. For quantitative analysis,

dying cells exhibiting continuous calcium activity were manually counted using Fiji

software and the data were analyzed with KaleidaGraph.

Detection of dying cells and quantification using CD8-PARP-Venus

To detect dying cells, the activated-caspase-3 indicator CD8-PARP-Venus was

expressed in the pouch under the control of *nub-gal4* driver. Cleaved PARP signals

were acquired at the confocal plane where signal intensity was highest. The number of

these signals in the wing pouch was manually measured using Fiji software and the

data were analyzed with KaleidaGraph.

Quantification of pH3-positive area

For the analysis of pH3-positive cells, total area of pH3-positive cells and the size of

the wing pouch area were automatically measured from confocal slices of Z-stack

images compressed using maximum projection functions in Fiji software, as

previously described [73]. The acquired data were then analyzed with KaleidaGraph.

Measurement of adult wings

Left and right wings of female flies were rinsed with xylene and mounted in a Canada Balsam (Nacalai Tesque). Images of the wings were acquired using Leica M205C stereo microscope.

Quantification of clone size

For the analysis of clone size in the eye disc, total clone area as a percentage of the disc area (%) was measured from XY confocal images using Fiji software, as previously described [73]. The acquired data were then analyzed with KaleidaGraph.

FACS and mRNA-seq analysis

GFP+ wing pouch cells were isolated using a FACS Aria II cell sorter (BD Bioscience). Total RNA was then extracted for RNA sequencing analysis, following the method previously described [74]. Briefly, third instar larvae were dissected in ice-cold PBS. The wing disc cells were dissociated using Trypsin (10x TrypLE Select) (Gibco, #A1217701) at 37°C for 20 min. The enzymatic reaction was stopped with Schneider's medium containing 5% FBS, and GFP-positive cells from wing pouches were sorted using a FACS Aria II cell sorter (BD Bioscience). RNA samples were extracted using NucleoSpin RNA XS Kit (TaKaRa, #740902). The quality of the extracted total RNA was assessed using an Agilent 2100 Bioanalyzer, and then Strand-specific Libraries for mRNA-seq were prepared using the NEBNext Ultra II Directional RNA Library Prep Kit for Illumina (E7760S). For RNA-seq, NextSeq 500/550 High output Kit v2.5 was utilized on a NextSeq 500 (Illumine), generating single-end reads at a length of 85 bases.

RNA-seq data analysis

Each genotype was analyzed across three independent replicates. Each sequencing experiment generated more than 16 million raw reads. These reads were then quality-filtered using Trim_Galore! (v0.6.5) (https://www.bioinformatics.babraham.ac.uk/projects/ trim_galore/) with the default setting to remove low-quality reads and adaptor sequences. The filtered reads were mapped to the *Drosophila* melanogaster genome (Ensembl BDGP6.32), obtained from Illumina iGenomes using STAR [75]. In the *RpS3/+, nub-Gal4, UAS-Puc* experiment, one sample exhibited only 65.69% of reads being uniquely mapped,

which was lower compared to other experiments where more than 80% of reads were
uniquely mapped in each sample. The number of reads mapped to each gene was
quantified using RSEM (v1.3.3) [76]. Normalization was performed with
calcNormFactors function in edgeR (v3.30.3) [77]. Significantly differentially expressed
genes were identified with an FDR<0.05 using the glmQLFit and glmQLFTest
function in edgeR (R version 4.0.1, limma 3.44.3).
Furthermore, GO enrichment analysis was performed on the 362 genes regulated by
JNK signaling, utilizing the online tool available at "Gene Ontology"
(https://geneontology.org).
623
Statistical analysis
Statistical analysis was performed, and boxplot graphs were generated using
KaleidaGraph ver4.5 (Synergy Software). All experiments represented in the same
graph were conducted at the same time. In bar graphs, error bars indicate the standard
deviation (SD). Significance levels are indicated as follows: * $P<0.05$, ** $P<0.01$, and
*** $P<0.001$. All statistical data were summarized in Table S4.
630
631
632

Supplemental information

Document S1. Figures S1-S3 and Table S5.

Video S1. Calcium Time-lapse imaging of wing discs. Related to Figure 1.

The calcium reporter GCaMP6m was expressed in wing pouches of wild-type larvae. Wing discs were dissected and subjected to time-lapse imaging at 10-second intervals with a Leica STELLARIS confocal microscope. See *Methods* for details.

Video S2. Calcium Time-lapse imaging of wing discs. Related to Figure 1.

The calcium reporter GCaMP6m was expressed in wing pouches of *RpS3/+* larvae. Wing discs were dissected and subjected to time-lapse imaging at 10-second intervals with a Leica STELLARIS confocal microscope. See *Methods* for details.

Video S3. Calcium Time-lapse imaging of wing discs. Related to Figure 1.

The calcium reporter GCaMP6m was expressed in wing pouches of *RpS3/+*, *nub-Gal4, UAS-Puc* larvae. Wing discs were dissected and subjected to time-lapse imaging at 10-second intervals with a Leica STELLARIS confocal microscope. See *Methods* for details.

Table S1

List of genes identified as differentially expressed by RNA-seq between RpS3/+ and WT, and between *RpS3*/+, +Puc and WT pouch cells.

Table S2

List of genes differentially expressed in *RpS3*/+ pouch cells, associated with JNK or JAK/STAT targets, Gr64 gustatory receptors, oxidative stress, and DNA repair.

Table S3

List of genes identified as differentially expressed by RNA-seq between *RpS3*/+ and *RpS3*/+, +Puc pouch cells.

668   **Table S4**

669   **Summary of statistical analysis.**

672
673 **References**
674

1.   Cmejla, R., Cmejlova, J., Handrkova, H., Petrak, J., and Pospisilova, D.
(2007). Ribosomal protein S17 gene (RPS17) is mutated in Diamond-Blackfan
anemia. Hum Mutat *28*, 1178-1182. 10.1002/humu.20608.

2.   Doherty, L., Sheen, M.R., Vlachos, A., Choesmel, V., O'Donohue, M.F.,
Clinton, C., Schneider, H.E., Sieff, C.A., Newburger, P.E., Ball, S.E., et al.
(2010). Ribosomal protein genes RPS10 and RPS26 are commonly mutated in
Diamond-Blackfan anemia. Am J Hum Genet *86*, 222-228.
10.1016/j.ajhg.2009.12.015.

3.   Campagnoli, M.F., Ramenghi, U., Armiraglio, M., Quarello, P., Garelli, E.,
Carando, A., Avondo, F., Pavesi, E., Fribourg, S., Gleizes, P.E., et al. (2008).
RPS19 mutations in patients with Diamond-Blackfan anemia. Hum Mutat *29*,
911-920. 10.1002/humu.20752.

4.   Choesmel, V., Fribourg, S., Aguissa-Touré, A.H., Pinaud, N., Legrand, P.,
Gazda, H.T., and Gleizes, P.E. (2008). Mutation of ribosomal protein RPS24
in Diamond-Blackfan anemia results in a ribosome biogenesis disorder. Hum
Mol Genet *17*, 1253-1263. 10.1093/hmg/ddn015.

5.   Wang, R., Yoshida, K., Toki, T., Sawada, T., Uechi, T., Okuno, Y., Sato-
Otsubo, A., Kudo, K., Kamimaki, I., Kanezaki, R., et al. (2015). Loss of
function mutations in RPL27 and RPS27 identified by whole-exome
sequencing in Diamond-Blackfan anaemia. Br J Haematol *168*, 854-864.
10.1111/bjh.13229.

6.   Danilova, N., and Gazda, H.T. (2015). Ribosomopathies: how a common root
can cause a tree of pathologies. Dis Model Mech *8*, 1013-1026.
10.1242/dmm.020529.

7.   Lipton, J.M., and Ellis, S.R. (2009). Diamond-Blackfan anemia: diagnosis,
treatment, and molecular pathogenesis. Hematol Oncol Clin North Am *23*,
261-282. 10.1016/j.hoc.2009.01.004.

8.   Bolze, A., Mahlaoui, N., Byun, M., Turner, B., Trede, N., Ellis, S.R.,
Abhyankar, A., Itan, Y., Patin, E., Brebner, S., et al. (2013). Ribosomal
protein SA haploinsufficiency in humans with isolated congenital asplenia.
Science *340*, 976-978. 10.1126/science.1234864.

9.   Brehme, K.S. (1939). A Study of the Effect on Development of "Minute"
Mutations in Drosophila Melanogaster. Genetics *24*, 131-161.
10.1093/genetics/24.2.131.

10.   Recasens-Alvarez, C., Alexandre, C., Kirkpatrick, J., Nojima, H., Huels, D.J.,
Snijders, A.P., and Vincent, J.P. (2021). Ribosomopathy-associated mutations
cause proteotoxic stress that is alleviated by TOR inhibition. Nat Cell Biol *23*,
127-135. 10.1038/s41556-020-00626-1.

11.   Baumgartner, M.E., Dinan, M.P., Langton, P.F., Kucinski, I., and Piddini, E.
714   (2021). Proteotoxic stress is a driver of the loser status and cell competition.
715   Nat Cell Biol *23*, 136-146. 10.1038/s41556-020-00627-0.

716   12.   Kucinski, I., Dinan, M., Kolahgar, G., and Piddini, E. (2017). Chronic
717   activation of JNK JAK/STAT and oxidative stress signalling causes the loser
718   cell status. Nat Commun *8*, 136. 10.1038/s41467-017-00145-y.

719   13.   Lee, C.H., Kiparaki, M., Blanco, J., Folgado, V., Ji, Z., Kumar, A., Rimesso,
720   G., and Baker, N.E. (2018). A Regulatory Response to Ribosomal Protein
721   Mutations Controls Translation, Growth, and Cell Competition. Dev Cell *46*,
722   456-469.e454. 10.1016/j.devcel.2018.07.003.

723   14.   Akai, N., Ohsawa, S., Sando, Y., and Igaki, T. (2021). Epithelial cell-turnover
724   ensures robust coordination of tissue growth in Drosophila ribosomal protein
725   mutants. PLoS Genet *17*, e1009300. 10.1371/journal.pgen.1009300.

726   15.   Fan, Y., and Bergmann, A. (2008). Distinct mechanisms of apoptosis-induced
727   compensatory proliferation in proliferating and differentiating tissues in the
728   Drosophila eye. Dev Cell *14*, 399-410. 10.1016/j.devcel.2008.01.003.

729   16.   Huh, J.R., Guo, M., and Hay, B.A. (2004). Compensatory proliferation
730   induced by cell death in the Drosophila wing disc requires activity of the
731   apical cell death caspase Dronc in a nonapoptotic role. Curr Biol *14*, 1262-
732   1266. 10.1016/j.cub.2004.06.015.

733   17.   Pérez-Garijo, A., Martín, F.A., and Morata, G. (2004). Caspase inhibition
734   during apoptosis causes abnormal signalling and developmental aberrations in
735   Drosophila. Development *131*, 5591-5598. 10.1242/dev.01432.

736   18.   Ryoo, H.D., Gorenc, T., and Steller, H. (2004). Apoptotic cells can induce
737   compensatory cell proliferation through the JNK and the Wingless signaling
738   pathways. Dev Cell *7*, 491-501. 10.1016/j.devcel.2004.08.019.

739   19.   Smith-Bolton, R.K., Worley, M.I., Kanda, H., and Hariharan, I.K. (2009).
740   Regenerative growth in Drosophila imaginal discs is regulated by Wingless
741   and Myc. Dev Cell *16*, 797-809. 10.1016/j.devcel.2009.04.015.

742   20.   Morata, G., and Ripoll, P. (1975). Minutes: mutants of drosophila
743   autonomously affecting cell division rate. Dev Biol *42*, 211-221.
744   10.1016/0012-1606(75)90330-9.

745   21.   Ogawa, M., Kawarazaki, Y., Fujita, Y., Naguro, I., and Ichijo, H. (2021).
746   FGF21 Induced by the ASK1-p38 Pathway Promotes Mechanical Cell
747   Competition by Attracting Cells. Curr Biol *31*, 1048-1057.e1045.
748   10.1016/j.cub.2020.11.052.

749   22.   Omar-Hmeadi, M., and Idevall-Hagren, O. (2021). Insulin granule biogenesis
750   and exocytosis. Cell Mol Life Sci *78*, 1957-1970. 10.1007/s00018-020-03688-
751   4.

23. Richmond, J.E., Weimer, R.M., and Jorgensen, E.M. (2001). An open form of
syntaxin bypasses the requirement for UNC-13 in vesicle priming. Nature *412*,
338-341. 10.1038/35085583.

24. Megighian, A., Scorzeto, M., Zanini, D., Pantano, S., Rigoni, M., Benna, C.,
Rossetto, O., Montecucco, C., and Zordan, M. (2010). Arg206 of SNAP-25 is
essential for neuroexocytosis at the Drosophila melanogaster neuromuscular
junction. J Cell Sci *123*, 3276-3283. 10.1242/jcs.071316.

25. Rizo, J., and Rosenmund, C. (2008). Synaptic vesicle fusion. Nat Struct Mol
Biol *15*, 665-674. 10.1038/nsmb.1450.

26. James, D.J., and Martin, T.F. (2013). CAPS and Munc13: CATCHRs that
SNARE   Vesicles.   Front   Endocrinol   (Lausanne)   *4*,   187.
10.3389/fendo.2013.00187.

27. Eugster, C., Panáková, D., Mahmoud, A., and Eaton, S. (2007). Lipoprotein-
heparan sulfate interactions in the Hh pathway. Dev Cell *13*, 57-71.
10.1016/j.devcel.2007.04.019.

28. Gross, J.C., Chaudhary, V., Bartscherer, K., and Boutros, M. (2012). Active
Wnt proteins are secreted on exosomes. Nat Cell Biol *14*, 1036-1045.
10.1038/ncb2574.

29. Panáková, D., Sprong, H., Marois, E., Thiele, C., and Eaton, S. (2005).
Lipoprotein particles are required for Hedgehog and Wingless signalling.
Nature *435*, 58-65. 10.1038/nature03504.

30. Redhai, S., Hellberg, J.E., Wainwright, M., Perera, S.W., Castellanos, F.,
Kroeger, B., Gandy, C., Leiblich, A., Corrigan, L., Hilton, T., et al. (2016).
Regulation of Dense-Core Granule Replenishment by Autocrine BMP
Signalling in Drosophila Secondary Cells. PLoS Genet *12*, e1006366.
10.1371/journal.pgen.1006366.

31. Sugita, S., Han, W., Butz, S., Liu, X., Fernández-Chacón, R., Lao, Y., and
Südhof, T.C. (2001). Synaptotagmin VII as a plasma membrane Ca(2+) sensor
in exocytosis. Neuron *30*, 459-473. 10.1016/s0896-6273(01)00290-2.

32. Yamanaka, N., Marqués, G., and O'Connor, M.B. (2015). Vesicle-Mediated
Steroid Hormone Secretion in Drosophila melanogaster. Cell *163*, 907-919.
10.1016/j.cell.2015.10.022.

33. Zhang, Y.Q., Rodesch, C.K., and Broadie, K. (2002). Living synaptic vesicle
marker: synaptotagmin-GFP. Genesis *34*, 142-145. 10.1002/gene.10144.

34. McGough, I.J., Vecchia, L., Bishop, B., Malinauskas, T., Beckett, K., Joshi,
D., O'Reilly, N., Siebold, C., Jones, E.Y., and Vincent, J.P. (2020). Glypicans
shield the Wnt lipid moiety to enable signalling at a distance. Nature *585*, 85-
90. 10.1038/s41586-020-2498-z.

35. Vietri, M., Radulovic, M., and Stenmark, H. (2020). The many functions of
ESCRTs. Nat Rev Mol Cell Biol *21*, 25-42. 10.1038/s41580-019-0177-4.

36. Chen, T.W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A.,
Schreiter, E.R., Kerr, R.A., Orger, M.B., Jayaraman, V., et al. (2013).
Ultrasensitive fluorescent proteins for imaging neuronal activity. Nature *499*,
295-300. 10.1038/nature12354.

37. Enomoto, M., Kizawa, D., Ohsawa, S., and Igaki, T. (2015). JNK signaling is
converted from anti- to pro-tumor pathway by Ras-mediated switch of Warts
activity. Dev Biol *403*, 162-171. 10.1016/j.ydbio.2015.05.001.

38. Florentin, A., and Arama, E. (2012). Caspase levels and execution efficiencies
determine the apoptotic potential of the cell. J Cell Biol *196*, 513-527.
10.1083/jcb.201107133.

39. Gorelick-Ashkenazi, A., Weiss, R., Sapozhnikov, L., Florentin, A., Tarayrah-
Ibraheim, L., Dweik, D., Yacobi-Sharon, K., and Arama, E. (2018). Caspases
maintain tissue integrity by an apoptosis-independent inhibition of cell
migration and invasion. Nat Commun *9*, 2806. 10.1038/s41467-018-05204-6.

40. Liang, C.J., Chang, Y.C., Chang, H.C., Wang, C.K., Hung, Y.C., Lin, Y.E.,
Chan, C.C., Chen, C.H., Chang, H.Y., and Sang, T.K. (2014). Derlin-1
regulates mutant VCP-linked pathogenesis and endoplasmic reticulum stress-
induced apoptosis. PLoS Genet *10*, e1004675. 10.1371/journal.pgen.1004675.

41. Ohsawa, S., Sato, Y., Enomoto, M., Nakamura, M., Betsumiya, A., and Igaki,
T. (2012). Mitochondrial defect drives non-autonomous tumour progression
through Hippo signalling in Drosophila. Nature *490*, 547-551.
10.1038/nature11452.

42. Ohsawa, S., Sugimura, K., Takino, K., Xu, T., Miyawaki, A., and Igaki, T.
(2011). Elimination of oncogenic neighbors by JNK-mediated engulfment in
Drosophila. Dev Cell *20*, 315-328. 10.1016/j.devcel.2011.02.007.

43. Shinoda, N., Obata, F., Zhang, L., and Miura, M. (2016). Drosophila SETDB1
and caspase cooperatively fine-tune cell fate determination of sensory organ
precursor. Genes Cells *21*, 378-386. 10.1111/gtc.12348.

44. Williams, D.W., Kondo, S., Krzyzanowska, A., Hiromi, Y., and Truman, J.W.
(2006). Local caspase activity directs engulfment of dendrites during pruning.
Nat Neurosci *9*, 1234-1236. 10.1038/nn1774.

45. de Wit, H., Walter, A.M., Milosevic, I., Gulyás-Kovács, A., Riedel, D.,
Sørensen, J.B., and Verhage, M. (2009). Synaptotagmin-1 docks secretory
vesicles to syntaxin-1/SNAP-25 acceptor complexes. Cell *138*, 935-946.
10.1016/j.cell.2009.07.027.

46. Baietti, M.F., Zhang, Z., Mortier, E., Melchior, A., Degeest, G., Geeraerts, A.,
Ivarsson, Y., Depoortere, F., Coomans, C., Vermeiren, E., et al. (2012).
Syndecan-syntenin-ALIX regulates the biogenesis of exosomes. Nat Cell Biol
14, 677-685. 10.1038/ncb2502.

47. Ghossoub, R., Lembo, F., Rubio, A., Gaillard, C.B., Bouchet, J., Vitale, N.,
Slavík, J., Machala, M., and Zimmermann, P. (2014). Syntenin-ALIX

exosome biogenesis and budding into multivesicular bodies are controlled by
ARF6 and PLD2. Nat Commun *5*, 3477. 10.1038/ncomms4477.

48. Shlevkov, E., and Morata, G. (2012). A dp53/JNK-dependant feedback
amplification loop is essential for the apoptotic response to stress in
Drosophila. Cell Death Differ *19*, 451-460. 10.1038/cdd.2011.113.

49. Wada, Y., Ohsawa, S., and Igaki, T. (2021). Yorkie ensures robust tissue
growth in Drosophila ribosomal protein mutants. Development *148*.
10.1242/dev.198705.

50. Vincent, J.P., Kolahgar, G., Gagliardi, M., and Piddini, E. (2011). Steep
differences in wingless signaling trigger Myc-independent competitive cell
interactions. Dev Cell *21*, 366-374. 10.1016/j.devcel.2011.06.021.

51. Pérez-Garijo, A., Shlevkov, E., and Morata, G. (2009). The role of Dpp and
Wg in compensatory proliferation and in the formation of hyperplastic
overgrowths caused by apoptotic cells in the Drosophila wing disc.
Development *136*, 1169-1177. 10.1242/dev.034017.

52. Wells, B.S., Yoshida, E., and Johnston, L.A. (2006). Compensatory
proliferation in Drosophila imaginal discs requires Dronc-dependent p53
activity. Curr Biol *16*, 1606-1615. 10.1016/j.cub.2006.07.046.

53. Beckett, K., Monier, S., Palmer, L., Alexandre, C., Green, H., Bonneil, E.,
Raposo, G., Thibault, P., Le Borgne, R., and Vincent, J.P. (2013). Drosophila
S2 cells secrete wingless on exosome-like vesicles but the wingless gradient
forms independently of exosomes. Traffic *14*, 82-96. 10.1111/tra.12016.

54. Koles, K., Nunnari, J., Korkut, C., Barria, R., Brewer, C., Li, Y., Leszyk, J.,
Zhang, B., and Budnik, V. (2012). Mechanism of evenness interrupted (Evi)-
exosome release at synaptic boutons. J Biol Chem *287*, 16820-16834.
10.1074/jbc.M112.342667.

55. Korkut, C., Ataman, B., Ramachandran, P., Ashley, J., Barria, R., Gherbesi,
N., and Budnik, V. (2009). Trans-synaptic transmission of vesicular Wnt
signals through Evi/Wntless. Cell *139*, 393-404. 10.1016/j.cell.2009.07.051.

56. Won, J.H., and Cho, K.O. (2021). Wg secreted by conventional Golgi
transport diffuses and forms Wg gradient whereas Wg tethered to extracellular
vesicles do not diffuse. Cell Death Differ *28*, 1013-1025. 10.1038/s41418-
020-00632-8.

57. Fogarty, C.E., and Bergmann, A. (2017). Killers creating new life: caspases
drive apoptosis-induced proliferation in tissue repair and disease. Cell Death
Differ *24*, 1390-1400. 10.1038/cdd.2017.47.

58. Morata, G., Shlevkov, E., and Pérez-Garijo, A. (2011). Mitogenic signaling
from apoptotic cells in Drosophila. Dev Growth Differ *53*, 168-176.
10.1111/j.1440-169X.2010.01225.x.

59.   Su, T.T. (2015). Non-autonomous consequences of cell death and other perks
of being metazoan. AIMS Genet *2*, 54-69.
10.3934/genet.2015.1.54#sthash.dNy9tFhS.dpuf.

60.   Fan, Y., Wang, S., Hernandez, J., Yenigun, V.B., Hertlein, G., Fogarty, C.E.,
Lindblad, J.L., and Bergmann, A. (2014). Genetic models of apoptosis-
induced proliferation decipher activation of JNK and identify a requirement of
EGFR signaling for tissue regenerative responses in Drosophila. PLoS Genet
10, e1004131. 10.1371/journal.pgen.1004131.

61.   Chen, Q., Takada, R., Noda, C., Kobayashi, S., and Takada, S. (2016).
Different populations of Wnt-containing vesicles are individually released
from polarized epithelial cells. Sci Rep *6*, 35562. 10.1038/srep35562.

62.   Mulligan, K.A., Fuerer, C., Ching, W., Fish, M., Willert, K., and Nusse, R.
(2012). Secreted Wingless-interacting molecule (Swim) promotes long-range
signaling by maintaining Wingless solubility. Proc Natl Acad Sci U S A *109*,
370-377. 10.1073/pnas.1119197109.

63.   Huang, H., and Kornberg, T.B. (2015). Myoblast cytonemes mediate Wg
signaling from the wing imaginal disc and Delta-Notch signaling to the air sac
primordium. Elife *4*, e06114. 10.7554/eLife.06114.

64.   Mattes, B., Dang, Y., Greicius, G., Kaufmann, L.T., Prunsche, B., Rosenbauer,
J., Stegmaier, J., Mikut, R., Özbek, S., Nienhaus, G.U., et al. (2018). Wnt/PCP
controls spreading of Wnt/β-catenin signals by cytonemes in vertebrates. Elife
7. 10.7554/eLife.36953.

65.   Stanganello, E., Hagemann, A.I., Mattes, B., Sinner, C., Meyen, D., Weber, S.,
Schug, A., Raz, E., and Scholpp, S. (2015). Filopodia-based Wnt transport
during vertebrate tissue patterning. Nat Commun *6*, 5846.
10.1038/ncomms6846.

66.   Zhang, C., Brunt, L., Ono, Y., Rogers, S., and Scholpp, S. (2024). Cytoneme-
mediated transport of active Wnt5b-Ror2 complexes in zebrafish. Nature *625*,
126-133. 10.1038/s41586-023-06850-7.

67.   Xue, S., and Barna, M. (2012). Specialized ribosomes: a new frontier in gene
regulation and organismal biology. Nat Rev Mol Cell Biol *13*, 355-369.
10.1038/nrm3359.

68.   Martín-Blanco, E., Gampel, A., Ring, J., Virdee, K., Kirov, N., Tolkovsky,
A.M., and Martinez-Arias, A. (1998). puckered encodes a phosphatase that
mediates a feedback loop regulating JNK activity during dorsal closure in
Drosophila. Genes Dev *12*, 557-570. 10.1101/gad.12.4.557.

69.   Quinn, L.M., Dorstyn, L., Mills, K., Colussi, P.A., Chen, P., Coombe, M.,
Abrams, J., Kumar, S., and Richardson, H. (2000). An essential role for the
caspase dronc in developmentally programmed cell death in Drosophila. J Biol
Chem *275*, 40416-40424. 10.1074/jbc.M002935200.

70. Igaki, T., Pagliarini, R.A., and Xu, T. (2006). Loss of cell polarity drives
tumor growth and invasion through JNK activation in Drosophila. Curr Biol
16, 1139-1146. 10.1016/j.cub.2006.04.042.

71. Moreno, E., Yan, M., and Basler, K. (2002). Evolution of TNF signaling
mechanisms: JNK-dependent apoptosis triggered by Eiger, the Drosophila
homolog of the TNF superfamily. Curr Biol *12*, 1263-1268. 10.1016/s0960-
9822(02)00954-5.

72. Aldaz, S., Escudero, L.M., and Freeman, M. (2010). Live imaging of
Drosophila imaginal disc development. Proceedings of the National Academy
of Sciences of the United States of America *107*, 14217-14222.
10.1073/pnas.1008623107.

73. Iida, C., Ohsawa, S., Taniguchi, K., Yamamoto, M., Morata, G., and Igaki, T.
(2019). JNK-mediated Slit-Robo signaling facilitates epithelial wound repair
by extruding dying cells. Scientific reports *9*, 19549. 10.1038/s41598-019-
56137-z.

74. Cong, B., Nakamura, M., Sando, Y., Kondo, T., Ohsawa, S., and Igaki, T.
(2021). JNK and Yorkie drive tumor malignancy by inducing L-amino acid
transporter 1 in Drosophila. PLoS Genet *17*, e1009893.
10.1371/journal.pgen.1009893.

75. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S.,
Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal
RNA-seq aligner. Bioinformatics *29*, 15-21. 10.1093/bioinformatics/bts635.

76. Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification
from RNA-Seq data with or without a reference genome. BMC Bioinformatics
12, 323. 10.1186/1471-2105-12-323.

77. Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a
Bioconductor package for differential expression analysis of digital gene
expression data. Bioinformatics *26*, 139-140. 10.1093/bioinformatics/btp616.
940

Akai *et al.*, Figure 1

Akai *et al.*, Figure 2

A *wt*

B–E *RpS3/+, nub-gal4*

B | **C** Puc | **D** *unc-13* RNAi VDRC33609 | **E** *Rab3* RNAi 7576R-3

pH3

F

% area pH3(+)/ pouch

*** *** *** *** ***

wt control +Puc +*unc-13*RNAi

RpS3/+ +*Rab3*RNAi

G *wt, nub-gal4*

RpS3/+, nub-gal4

H *LexA*RNAi HMS05772 | **I** *unc-13*RNAi VDRC33609 | **J** *Rab3*RNAi 7576R-3

Cleaved PARP

K *SNAP25*RNAi HMS01367 | **L** *Gαq*²²¹C/+ BL30744 | **M** *Plc21C*RNAi 4574R-3 | **N** *norpA*⁷/+ DGRC108362

O

(a) normal

(b) vein (weak)

(c) vein (strong)

(d) notch

(e) severe

P

% adult wing

a

b
c
d
e

wt control +Puc +*unc-13*RNAi +*Rab3*RNAi

RpS3/+

T

No. of c-PARP(+) cells/ pouch

900
600
300
0

*** *** * ***

wt +*LexA*RNAi +*unc-13*RNAi +*Rab3*RNAi +*Snap25*RNAi +*Gαq*/+ +*Plc21C*RNAi +*norpA*/+

RpS3/+

U

Relative CD63-mCherry area /pouch

1.5

0.5

*** **

wt control +*Dronc*^{DN}

RpS3/+

Q *nub-gal4*

RpS3/+, nub-gal4

R | **S** *Dronc*^{DN}

CD63-mCherry

Minute/+ wing pouch

Living cell   **Dying cell**

compensatory proliferation

cell death