The genetic consequences of population marginality: a case study in

maritime pine

Adélaïde Theraroz[1], Carlos Guadaño-Peyrot[2], Juliette Archambeau[1,3], Sara Pinosio[4],

Francesca Bagnoli[4], Andrea Piotti[4], Camilla Avanzi[4], Giovanni G. Vendramin[4], Ricardo

Alía[2], Delphine Grivet[2], Marjana Westergren[5], Santiago C. González-Martínez[1,*]

[1]INRAE, Univ. Bordeaux, BIOGECO, Cestas, France

[2]Department of Forest Ecology and Genetics, Institute of Forest Science (ICIFOR-INIA),

CSIC, Carretera de la Coruña km 7,5, Madrid, Spain

[3]UK Centre for Ecology & Hydrology, Bush Estate, Penicuik, United Kingdom

[4]National Research Council, Institute of Biosciences and BioResources, Sesto Fiorentino,

Italy

[5]Slovenian Forestry Institute, Večna Pot 2, Ljubljana, Slovenia

*Corresponding author:

Santiago C. González-Martínez

UMR 1202 INRAE – Univ. Bordeaux

69 route d'Arcachon, F-33610 Cestas, France

Tel: +33 (0) 5 35 38 53 20

santiago.gonzalez-martinez@inrae.fr

Abstract (no more than 250 words)

Marginal tree populations, either those located at the edges of the species' range or in

suboptimal environments, are often a valuable genetic resource for biological conservation.

However, there is a lack of knowledge about the genetic consequences of population's

marginality, estimated across entire species' ranges. Using 10,185 SNPs across 82

populations of *Pinus pinaster* Ait., a widespread and economically important conifer

characterised by a fragmented range, we modelled the relationship of five genetic indicators

potentially related to ecological resilience, population vulnerability and adaptability (genetic

diversity, inbreeding, genetic differentiation, recessive genetic load and genomic offset) with

population geographical, demo-historical and ecological marginality (as estimated by nine

quantitative indices). Models were constructed for both regional (gene pool) and range-wide

spatial scales. We showed an overall trend towards decreasing genetic diversity and

increasing differentiation with geographic marginality, supporting the centre-periphery

hypothesis. Moreover, we found no correlation between population inbreeding and

marginality, while geographically marginal populations had a lower recessive genetic load

(only models without the gene pool effect) and ecologically marginal populations had a

higher genomic offset. Overall, our results suggest that marginal populations of maritime pine

could be more vulnerable to climate change than core populations, despite reduced levels of

genetic load, a risk that is exacerbated by typically small effective population sizes and

increasing human impact.

Keywords (no more than 6): marginal populations, centre-periphery hypothesis, ecological

modelling, genetic indicators, forest genetic resources, *Pinus pinaster*

Introduction

All species thrive and reproduce within an environmentally limited geographic area, which sets the boundaries of their range. Current global warming may restrict the climatic suitability of some parts of the species' ranges and modify their geographic and ecological margins. The 'centre-periphery hypothesis' (hereafter CPH), a major paradigm in biogeography that aims to disentangle the genetic, demographic, and ecological causes of species' range limits (Gaston, 2009; Sexton et al., 2009), defines marginality as the level of geographic isolation from the species' centre of distribution, which in turn is related to the species' suitability for its environment (Brown, 1984; Hengeveld & Haeck, 1982). According to the CPH, marginal populations are expected to be less abundant and more prone to extinction than those in the centre, due to harsher environmental conditions at the periphery (Birch, 1957; Gaston, 2003; Nicholson, 1958; Richards, 1961; Whittaker, 1971). However, as specific environmental conditions (topography, soils) in the core of the species distribution may also induce harsher environmental conditions than in the periphery, Soulé (1973) distinguished between geographical and ecological marginality when describing centre-periphery gradients, defining ecological marginality as a population's exposure to extreme environmental variables irrespectively of their geographical location.

While several studies have identified marginal populations based on different criteria (e.g., for maritime pine, Alía et al., 1996; Burban & Petit, 2003), we often lack evidence on the extent to which population marginality is associated with particular genetic features. Classic studies at the range-wide scale suggested a general trend for lower genetic diversity and higher genetic differentiation in marginal populations, but they typically did not distinguish between geographical and ecological marginality (Eckert et al., 2008; Johannesson & André, 2006; Pironon et al., 2015). More recent studies often called for decoupling the roles of geography and ecology in shaping range-wide patterns of genetic

variation (Lira-Noriega & Manthey, 2014; Sexton et al., 2013). To that end, a clear

distinction needs to be made between different kinds of population marginality. In a recent

multispecies study, Picard et al. (2022) evaluated the ability of quantitative measures to

distinguish between geographical, demo-historical (i.e., related to the demographic history of

the target species) and ecological marginality.

Past global climate dynamics rather than demographic stochasticity seem to have

played a determining role in the establishment of current range limits (e.g., Hampe & Petit,

2005, for forest trees). Postglacial migrations after the Last Glacial Maximum (LGM) would

have resulted in a mosaic of relatively small and isolated populations in the rear and, at a

lesser extent, front edges of the expansion. In this context, the level of connectivity among

marginal populations and between a given marginal population and the species distribution

core is especially important (Sachdeva et al., 2022), as gene flow may increase genetic

diversity and reduce differentiation (e.g., Lynch et al., 1995; Young et al., 1996). Gene flow

can also bring adaptive alleles and contribute to the evolutionary rescue of small, isolated

populations by buffering the effect of genetic drift, and reducing the fixation and

accumulation of deleterious alleles within populations (Sachdeva et al., 2022). However,

marginal populations may also carry specific alleles derived from local adaptation to atypical

environments, constituting valuable genetic resources, and making the contribution of

external gene flow harmful (i.e., the so-called 'migration load'; Kimura et al., 1963).

Overall, marginal populations are considered to be more vulnerable to climate change

than core populations (Kolzenburg, 2022; Soulé, 1973). Marginal populations are expected to

accumulate deleterious variants (i.e., genetic load), a process governed by effective

population size, N_e (Kimura & Ohta, 1969). However, genetic purging due to inbreeding

tends to reduce genetic load over time, even in relatively small populations (Hedrick &

Garcia-Dorado, 2016). Current developments in population genomics have provided metrics

to estimate components of population vulnerability to future climates (e.g., by estimating genomic 'offsets' or 'gaps'; Capblancq et al., 2020; Fitzpatrick & Keller, 2015; Gougherty et al., 2021; Rellstab et al., 2021), allowing for much needed systematic studies on the connection between population marginality and vulnerability.

Maritime pine (*Pinus pinaster* Ait., Pinaceae) is an outcrossing, wind-pollinated conifer, with a widespread but fragmented natural distribution in southwestern Europe and North Africa, covering a wide range of contrasted environments from coastal dunes next to the Atlantic Ocean in France to the High Atlas Mountains in Morocco. Maritime pine population genetic structure is a consequence of historical and current dynamics of range expansion-contraction, resulting in distinct genetic clusters (Jaramillo-Correa et al., 2015). This species is also characterised by a fragmented distribution due to human-induced habitat loss and ecological disturbances such as forest fires (De-Lucas et al., 2009). Nowadays, some maritime pine populations are found in ecologically marginal environments (e.g., under very dry conditions in southern Spain and northern Morocco). In addition, the species' range margins are characterised by small, geographically isolated populations, in particular in the southern and eastern parts of the distribution (Alía et al., 1996; Wahid et al., 2004). Reduced dispersal with distant core populations, coupled with demographic and environmental stochasticity, may push such populations into an 'extinction vortex' (Lande, 1988). Few studies have focused on describing the particular genetic characteristics of marginal populations of maritime pine, whose potentially valuable genetic resources could be lost in the near future.

In this study, we used 10,185 Single Nucleotide Polymorphisms (SNPs) genotyped in 82 maritime pine populations (1,510 individuals), including ecologically and geographically marginal ones, to compute genetic indicators potentially related to ecological resilience, population vulnerability and adaptability. The main objective of this study is to assess the

relationship between marginality and population genetic features at the scale of the whole species range, by testing predictions of the CPH and adding new elements to its general framework. More specifically, we i) assessed losses of genetic diversity and increased genetic differentiation in marginal populations; ii) evaluated levels of maladaptation brought about by the accumulation of recessive genetic load; and iii) tested whether marginal populations are more vulnerable to climate change, applying genomic approaches that consider the contribution of pre-adapted variants to future climates (i.e., genomic offset models). To this end, we estimated a wide range of genetic indicators and correlated them with quantitative measures of marginality, providing valuable insights for understanding processes governing genetic resources in maritime pine.

Material and Methods

Plant material and molecular markers

Needles were collected from 1,510 individuals in 82 maritime pine populations covering all previously identified gene pools throughout the species range (Jaramillo-Correa et al., 2015; Milesi et al., 2023). This sampling is, to date, the most complete in the species (see Figure 1 and Table S1 in Supplementary Information), and includes several populations from the distribution margins, as well as isolated populations that have not been considered in genetic studies before. Population Cómpeta (COM), with only three samples, was removed from all data analyses but the gene-environment association (GEA) methods used to estimate the genomic offset. The accuracy of landscape genomic approaches, such as GEA, is highly improved by increasing the number of populations and environments, while being less sensitive to unbalanced sampling designs (Santos & Gaiotto, 2020). In addition, the two stands of Maures population (MAU) (see Table S1) were kept separate for these analyses as they were sampled at different altitudes.

DNA was extracted using the DNeasy 96 plant kit (QIAGEN, Germany), following

the manufacturer's instructions. All the samples were genotyped for SNPs using the

multispecies 4TREE Axiom array (Thermo Fisher Scientific, USA). Only SNPs with high-

quality scoring following the Best Practices Workflow implemented in the Axiom™ Analysis

Suite v5.2 were selected and filtered by missing data (<30%), yielding a total of 10,185

SNPs. SNP annotation based on SnpEff v5.1 was retrieved from Piñeiro et al. (forthcoming)

for a set of 1,325 common SNPs between the two studies.

Data analysis

Population marginality indices

To assess population marginality, we first produced a new distribution map for maritime pine

that includes only natural populations (Figure S1), building on that of Caudullo et al. (2017)

but adding information from National Forest Inventories and specific publications for less-

known parts of the distribution range (Alía et al., 1996; Marques et al., 2012; Abad-Viñas et

al., 2016; Fkiri et al., 2019; Wahid et al., 2004, 2006). Second, we computed eight

quantitative marginality indices that consider both geographical distribution and demographic

history, following Picard et al. (2022) (see Table 1 and Table S2 in Supplementary

Information). The demo-historical indices, which are related to the postglacial colonization

history of the species (see, e.g., Bucci et al., 2007; Jaramillo-Correa et al., 2015), and the

centroid index were computed from the distribution map. The centroid of the species'

distribution corresponded to the location whose geographic coordinates were the average of

all populations included in the study. Then, the centroid index was defined as the cost

distance between any population and the centroid location in terms of gene flow circulation,

as a function of terrestrial and marine connectivity with respect to the species distribution

(see Table 1). This index reflects the level of long-distance gene flow. The other geographical

indices relied on a morphological spatial pattern analysis (MSPA), which considers a binary

image (1-presence/0-absence and NA for water) with emphasis on the connectivity within the

image (Soille & Vogt, 2009). Presence/absence maps were structured in three categories

(cores, edges, and other classes including loops, islets, bridges and branches) using

BioManager package in R version 4.2.2. Third, we calculated an index of ecological

marginality (henceforth 'ecological index') based on climatic data for the period 1901-1970,

as follows. For each population, we extracted the climatic information provided by the

Climate Downscaling Tool (ClimateDT, https://www.ibbr.cnr.it/climate-dt/) for the Summer

Heat Moisture (SHM) aridity index, an indicator of exposure to drought (see e.g., De La

Torre et al., 2014; Marchi et al., 2020), and temperature (bio4) and precipitation seasonality

(bio15). This set of climatic variables were selected because they best explained the climatic

variation across the species range in previous studies (e.g., Archambeau et al., forthcoming).

The ecological index was constructed by computing the standardized Euclidean distance for

SHM, bio4 and bio15 between each population and the overall average. Thus, this index

represents the climatic distance of the population from the average climate (Table S2).

Finally, we reduced the set of marginality indices by removing those that were highly

correlated based on Pearson's correlation coefficients (Pearson's correlation ≥ 0.6) and a

principal component analysis (PCA) using *FactoMineR* package in R v4.2.2 (Figure S2).

Population genetic structure and genetic indicators

Population genetic structure was evaluated using the Bayesian clustering algorithm

implemented in STRUCTURE v2.3.4 (Pritchard et al., 2000). To assess the optimal

individual's assignment probabilities ($Q_{ancestry}$) in K genetic clusters (or gene pools), we ran

models from $K=1$ (no structure) to $K=10$ with a burn-in length of 100,000 and run lengths of

200,000 MCMC steps. The number of K that best describes the genetic structure was

determined based on the delta *K* method (Evanno et al., 2005) and the visual observation of

bar plots.

All genetic indicators were computed at the population level (Table S3). In a few

cases where more than one stand was sampled for a population (see Table S1), genetic

indicator values were averaged. Genetic diversity was estimated as $1-Q_{inter}$, with Q_{inter} being

the observed frequencies of identical pairs of alleles among individuals within populations,

using GenePop v4.7.5 (Rousset, 2008), after standard correction for sample size (N), using

(N/N-1). Notice that this estimate averages over monomorphic and polymorphic loci.

Population inbreeding (F_{IS}) was estimated following Weir & Cockerham (1984). Population-

specific divergence was estimated as the genetic differentiation (population-specific F_{ST}) of

each population from a common ancestral gene pool using BayeScan v2.1 (Foll & Gaggiotti,

2008), as well as Jost's *D* (Jost, 2008) estimated with MMOD package in R v4.2.2.

Finally, we computed two genetic indicators related to population vulnerability based

on genomic data, the recessive genetic load and, more specifically related to population

vulnerability to climate change, the genomic offset. The recessive genetic load represents the

accumulation of predicted deleterious mutations in the population standardized by the

population genetic diversity. This statistic was estimated by counting different kind of

mutations (annotated by SnpEff v5.1) averaged over individuals, as the number of derived

moderate- (i.e., non-synonymous) and high-impact (i.e., loss of function) mutations in

homozygosity divided by the number of derived low-impact (i.e., synonymous) mutations in

homozygosity, following González-Martínez et al. (2017). Population vulnerability to climate

change is defined by three major components (Foden et al., 2013): exposure (which refers to

the variation of climatic factors within an organism's physical space), sensitivity (which

characterizes an organism's response to changes in its environment), and adaptive capacity

(which refers to a species' ability to tolerate, adapt or recover from changes in its

environment). The genomic offset is estimated as the change in genetic composition required

to maintain the current gene-climate relationships under future climates (see Fitzpatrick &

Keller, 2015) and thus captures the exposure/sensitivity component of population

vulnerability (Razgour et al., 2017). Briefly, we first identified candidate SNPs for climate

adaptation with two univariate GEA methods, BAYPASS (Gautier, 2015) and Latent Factor

Mixed Model (LFMM; Frichot et al., 2013), and three multivariate ones, Redundancy

Analysis (RDA; Capblancq et al., 2020), partial RDA (Van Den Wollenberg, 1977) and

Gradient Forest (GF; Ellis et al., 2012). Then, the genomic offset was estimated using the

candidate SNPs and the GF approach, which showed the best empirical validation in a

previous study based on a smaller sample of populations and SNPs (Archambeau et al.,

forthcoming), and six climatic variables related to maritime pine expected exposure to

climate change (Table S4; see also Archambeau et al., forthcoming). Future climates for 2070

were described using the predictions from the moderately alarming shared socio-economic

pathway (SSP) SSP3-7.0 and five global circulation models (GCMs; IPCC, 2021). As

genomic offset predictions across GCMs were highly correlated (Pearson's correlation

coefficient >0.75; Figure S3), we used population averages for the five GCMs. Additional

details and scripts are available at

https://github.com/JulietteArchambeau/ReadyToGO_Pinpin. All analyses were undertaken

under R v4.2.2 (R Core Team, 2022).

Effects of population marginality on genetic indicators

We estimated the association between genetic indicators and population marginality by fitting

two series of five linear mixed-models (one model for each genetic indicator), using the R

packages LME4 and LMER, respectively. Models M1 to M5 included population marginality

indices as fixed effects irrespectively of the gene pool of origin while models M6 to M10 also

included the gene pool of origin as random effects. Note that the two gene pools with a single

population (Fuencaliente, FUE; and Point Cires, PCI) were assigned to the geographically

closest gene pool (Southeastern Spain and Morocco, respectively). Model goodness-of-fit

was evaluated with R^2 and both the Akaike Information Criterion (AIC) and the Bayesian

Information Criterion (BIC). Then, the best models were selected by considering goodness-

of-fit (higher R^2 and lower AIC/BIC) and parsimony criteria (i.e., including only significant

effects at $\alpha = 0.01$).

Results

Marginality indices

Pairwise Pearson's correlations and PCA identified some strongly correlated marginality

indices (Figure S2). For example, the second nearest-core was positively correlated with the

isolation index and negatively correlated with the edge index, and the ecological index was

negatively correlated with the East-West index. Thus, only five indices with low correlation

(<0.6) were retained for further analysis, namely three geographical indices (centroid, second

nearest-core and gravity; see definitions in Table 1), one demo-historical (North-South), and

the ecological index based on climate distances (Figure S4). Interestingly, the centroid index

was positively correlated with the longitude (Pearson's correlation of 0.70) and the North-

South index was negatively correlated with the elevation (Pearson's correlation of -0.56),

with southern populations being, generally, at higher elevation (Figure S2).

Population genetic structure

Population genetic structure analyses identified ten distinct gene pools, among which two

included only a single population (FUE in southern Spain and PCI in northern Morocco;

Figure 1). Interestingly, these two single-population gene pools were not identified as

marginal populations by the geographical or demo-historical marginality indices whereas one

of them, PCI, was characterized by high values of the ecological index (standardized value of

5.013; Table S2), indicating persistence in a marginal climate. FUE and PCI had also low

levels of admixture with nearby gene pools. In contrast, the eight main gene pools (with the

exception of Tunisia - Pantelleria, see below) were not genetically isolated from each other,

with populations often showing admixture with nearby gene pools. This suggests either

historical or recent gene flow across neighbouring gene pools along a latitudinal cline in the

western range of the species, and substantial shared ancestry among French (including

Corsica) and Italian Mediterranean populations in the eastern one.

Effects of population marginality on genetic indicators

Our models revealed a decline in genetic diversity (corrected 1-Q_{inter}, Figure 2a) and an

increase in genetic differentiation (population-specific F_{ST}, Figure 2b; and Jost's D statistic,

Table S5) with population marginality based on the centroid, second-nearest core and North-

South indices (Tables 2 and S4). Indeed, models M1 and M3 predicted lower genetic

diversity and higher genetic differentiation in marginal populations based on the centroid

index (Tables 2 and S4), in particular in isolated populations from the southern maritime pine

range (as shown by significant coefficients for the second nearest-core and North-South

indices and their interaction; see also Figures 2 and S4d). Models including the gene pool of

origin as a random effect (M6 and M8; Tables 2 and S4) significantly improved the fit by

10% for genetic diversity and 20% for population-specific F_{ST} (29% for Jost's D statistic),

but retained only the centroid and North-South indices and their interaction as explanatory

factors. Interestingly, we found no relationship between inbreeding and the indices of

population marginality (see Figure S5a) for any model (M2, M7; Table 2).

The model for recessive genetic load (M4) showed reduced genetic load for marginal

populations based on the centroid index, this association probably stemming mainly from the

high recessive genetic load found in some Iberian core populations (e.g., Carbonero el

Mayor, CAR, and Boniches, BON; see Figure S5b). However, this model had a relatively low

goodness of fit ($R^2 = 12\%$) and we found no relationship between recessive genetic load and

any of the population marginality indices when we added the gene pool of origin as a random

effect (M9, Table 2). Temperature seasonality (bio4) was the most important predictor

contributing to the genetic turnover in genomic offset estimates (Figure 3a). Noticeably, this

variable showed a steep slope between -2 and 0°C, which may indicate a rapid turnover in

allele frequency in this range (see Figure S6a). Both models including genomic offset,

without (M5) and with (M10) the gene pool of origin as a random effect, predicted an

increased genomic offset for populations in marginal climatic conditions (Table 2 and Figure

3b). Accordingly, we observed a trend for higher genomic offset in the western gene pools

along the Atlantic coast (average genomic offset of 0.047 ± 0.026 and 0.047 ± 0.016 for the

Atlantic Iberian Peninsula and the French Atlantic gene pools, respectively), with remarkably

high population values for Segurde (SEG), Alto de la Llama (ALT), Armayán (ARM) and

Olonne sur Mer (OLO) as well as for some southern Mediterranean populations (e.g., Point

Cires, PCI; Fuencaliente, FUE; Estepona, EST; and Pantelleria, PAN; Figure 3a).

Discussion

In this study, we revealed a trend towards declining genetic diversity and increased genetic

differentiation in geographically and demo-historically marginal populations of maritime

pine. We also found lower recessive genetic load in geographically marginal populations and

higher genomic offset in ecologically marginal ones. Models including the gene pool of

origin as a random effect were similar to those without (with the notable exception of the

models for recessive genetic load), suggesting that the underlying processes operate at both

the regional and range wide geographical scales in this species. These results, taken together,

provide support for the CPH and suggest that climate change may endanger valuable and

untapped genetic resources in maritime pine.

Patterns of genetic diversity and differentiation support the CPH

The lower genetic diversity and higher genetic differentiation of geographical and demo-

historical marginal populations support the main predictions of the CPH. Our results are

consistent with those of Eckert et al. (2008) and Pironon et al. (2016), who found a decline in

genetic diversity and an increase in differentiation towards the limits of the species ranges in

47% and 45% of studies in various taxa, respectively. As in our study, lower genetic variation

in peripheral compared to central populations was found for some other conifers, i.e., in Sitka

spruce (*Picea sitchensis*; Gapare et al., 2005), Swiss stone pine (*Pinus cembra*; Gugerli et al.,

2009) and common yew (*Taxus baccata*; Hilfiker et al., 2004), but not in Norway spruce

(*Picea abies*), for which only genetic differentiation conformed to the CPH (Westergren et

al., 2018).

In addition to geographical marginality, demo-historical marginality showed a

significant association with genetic diversity and differentiation in maritime pine (see M1 and

M3 in Table 2a, and M3bis in Table S5). Southern populations from the Moroccan and

Tunisia-Pantelleria gene pools and northern populations from the North-East and Corsican

gene pools exhibited the lowest genetic diversity and the highest genetic differentiation in

maritime pine (see Figure 2). Maritime pine gene pools are probably the result of population

expansion from multiple glacial refugia, both in Mediterranean and Atlantic regions of the

species (Bucci et al., 2007; Jaramillo-Correa et al., 2015), with Naydenov et al. (2014) also

suggesting that, for this species, the high level of overall genetic differentiation may have

resulted from long historical isolation predating the Last Glacial Maximum (~18,000 years

ago). This was confirmed by estimates of genetic divergence between North-African

(Moroccan) and Iberian populations of maritime pine, which dated back to 1.90 Ma (95%

credibility interval: 1.41-2.76), probably due to the Strait of Gibraltar's effect as a major

biogeographic barrier to pollen and seed gene flow (Jaramillo-Correa et al., 2010).

Consequently, southernmost Moroccan populations are likely to have been pre-glacial relict

populations that survived in North-African refugia (Baradat & Marpeau, 1988; Vendramin et

al., 1998). Lower genetic diversity and high differentiation of Moroccan populations may

have been exacerbated by a reduction of effective population size (N_E) due to human impact

(Wahid et al., 2004), as well as the incidence of recurrent forest fires resulting in population

bottlenecks and genetic drift (Vendramin et al., 1998). The demographic history of the north-

eastern part of the continental range of maritime pine and Corsica has not been clearly

assessed (Naydenov et al., 2014), however, the presence of an endemic mitotype in this

region also suggests long-term isolation of North-East and Corsican gene pools (Burban &

Petit, 2003).

Lower recessive genetic load in geographically marginal populations

Our models revealed a significant, albeit weak, reduction in recessive genetic load with

increased geographical population marginality. However, these models were only significant

when the gene pool effect was not accounted for, suggesting the existence of gene pools with

reduced/increased recessive genetic load and an important role of demographic history (see

below). This was a surprising finding, as we were expecting marginal populations to be

characterised by an accumulation of recessive genetic load due to the reduced effectiveness

of purifying selection in small and isolated populations with high demographic stochasticity

(Caballero et al., 2017; Sachdeva et al., 2022). The level of accumulation of deleterious

mutations and the extent to which it represents a risk to a given population depends primarily on its effective size (N_E). Previous empirical studies of animal and plant populations that underwent historical range expansions or declines have often shown an increase in genetic load (e.g., Günther & Schmid, 2010, in *Arabidopsis thaliana*; Feng et al., 2019, in *Nipponia nippon*; González-Martínez et al., 2017, in *Mercurialis annua*; or Peischl et al., 2013, in humans). However, as in maritime pine, recent empirical studies based on genomic data suggested that recessive genetic load can also be purged in long-term isolated and inbred populations (see review in Dussex et al., 2023). As an example, Dussex et al. (2021) found that current island populations of *Strigops habroptilus*, a New-Zealand flightless parrot, had lower deleterious mutation load compared to mainland populations. A similar pattern was found for the Alpine ibex (*Capra ibex*), which suffered severe population bottlenecks and nearly became extinct (Grossen et al., 2020), and for the Channel Island fox (*Urocyon littoralis*) that has remained at small population sizes with low diversity for many generations (Robinson et al., 2018).

Present-day levels of inbreeding in maritime pine are low and not significantly higher in marginal than core populations. Therefore, we hypothesise maritime pine marginal populations to have effectively purged recessive genetic load during past inbreeding events, operating at the regional geographical scale. These events may have occurred during the range contractions and/or expansions associated to Quaternary glacial and interglacial forest tree migration episodes (Bucci et al., 2007; Naydenov et al. 2014; Jaramillo-Correa et al., 2015). Polyembryony, which is ubiquitous in gymnosperms such as maritime pine (Willson & Burley, 1983), could have also played a role in purging recessive genetic load, as it tends to dampen self-fertilisation's deleterious effects by more effectively removing the mutational load through selection between viable embryos (see Latta, 1995).

Higher genomic offset in ecologically marginal populations

Our models also found a higher genomic offset for ecologically marginal populations, but not for geographically or demo-historically marginal ones suggesting that the gap between the current and required genetic composition in future climates will be greater for populations in marginal climatic conditions. This is consistent with the findings of Fréjaville et al. (2019), who observed that adaptation lags in several forest trees, including maritime pine, are consistently higher in climatically marginal populations (cold/warm, dry/wet) than in populations growing under climatically optimal conditions. Although studies estimating genomic offset in marginal populations are still scarce, two recent studies in widespread Asian forest trees provided some support to our findings, as they showed a relatively high genomic offset in the northern and southern margins of the Chinese thuja tree (*Platycladus orientalis*; Jia et al., 2020) and the sawtooth oak (*Quercus acutissima*; Yuan et al., 2023); but whether these range margins represented also ecologically extreme environments was not assessed and more detailed studies are thus needed.

Estimation of the genomic offset is becoming a popular approach to assess population vulnerability in the face of climate change, but it is not free of pitfalls (see Ahrens et al., 2023; Rellstab et al., 2021; Lind et al., 2023; Archambeau et al., forthcoming). Moreover, the genomic offset can gauge for some components of population vulnerability (i.e., exposure and sensitivity; Estoque et al., 2023; IPCC, 2007) but not others (notably the adaptive capacity of populations; Fitzpatrick et al., 2021; Archambeau et al., forthcoming). Marginal populations of forest trees can retain notable adaptive capacity, as shown in common garden experiments for the handful of species with available data (e.g., for the Sierra Nevada Scots pine population in southern Spain, an isolated marginal population at the species southern distribution limit; Alía et al., 2001; Castro et al., 2004). Thus, the potential for both genetic

adaptation and plastic responses needs to be integrated in predictive models of the expected

responses of marginal populations to climate change.

Conclusion

In maritime pine, a trend of decreasing genetic diversity and increasing differentiation with

geographic marginality supported the well-established centre-periphery hypothesis, at both

range-wide and regional (gene pool) spatial scales. However, geographically marginal

populations displayed also lower recessive genetic load compared to core populations, which,

together with expected novel adaptations in the species range margins, highlight their

importance in the context of future adaptation to climate change. Higher genomic offset in

ecologically marginal populations suggests higher vulnerability of these populations to future

climates and the urgency to develop specific management actions to preserve populations in

marginal climatic conditions. Overall, our study shows the importance of combining

quantitative marginality indices and diverse genetic indicators, gauging for multiple

evolutionary processes, to have a sound basis for conservation decisions.

Acknowledgements

We thank Carmen García-Barriga, Francisco Auñón, Eduardo Ballestero, Diana Barba, Maria

Regina Chambel, Fernando Del-Caño and Rodrigo Pulido-Sanz for their contribution to

genetic data production. We are also very grateful to Dr Bruno Fady and Dr Nicolas Picard

for their valuable recommendations and help at different stages of this study.

Funding

This study was funded by the European Union's Horizon 2020 research and innovation

program under grant agreement No 862221 (FORGENIUS). The study also received financial

support from the French government in the framework of the IdEx Bordeaux University

'Investments for the Future' program / GPR Bordeaux Plant Sciences (France) and the

MITECO contract for Conservation of Genetic Resources (Spain).

Author Contribution

Data accessibility

Data are openly available in a public INRAE repository: https://doi.org/10.57745/SPUDQQ.

Conflict of interest

The authors declare no conflict of interest.

References

Abad-Viñas, R., Caudullo, G., Oliveira, S., & de Rigo, D. (2016). *Pinus pinaster* in Europe: distribution, habitat, usage and threats. In: San-Miguel-Ayanz, J., de Rigo, D., Caudullo, G., Houston Durrant, T., Mauri, A. (Eds.), *European Atlas of Forest Tree Species*. Publ. Off. EU, Luxembourg, pp. e012d59+

Ahrens, C. W., Rymer, P. D., & Miller, A. D. (2023). Genetic offset and vulnerability modelling: misinterpretations of results and violations of evolutionary principles.

bioRxiv (Cold Spring Harbor Laboratory). [Preprint].

https://doi.org/10.22541/au.168727971.18670759/v1

Alía, R., Martín, S., de Miguel, J., Galera, R., Agúndez, D., Gordo, J., Salvador, L., Catalán, G., & Gil, L. (1996). *Las regiones de procedencia de Pinus pinaster Aiton en España* (Organismo Autónomo Parques Nacionales-ETSI de Montes).

Alía, R., Moro-Serrano, J., & Notivol, E. (2001). Genetic variability of Scots pine (*Pinus sylvestris*) provenances in Spain: growth traits and survival. *Silva Fennica*, *35*(1). https://doi.org/10.14214/sf.601

Baradat, P., Marpeau-Bezard, A. (1988). Le pin maritime *Pinus pinaster* Ait. Biologie et génétique des terpènes pour la connaissance et l'amélioration de l'espèce. Thèse doctorale d'État, Université Bordeaux I

Birch, L. C. (1957). The role of weather in determining the distribution and abundance of animals. *Cold Spring Harbor Symposia on Quantitative Biology*, *22*, 203-218. https://doi.org/doi:10.1101/SQB.1957.022.01.021

Brown, J. H. (1984). On the relationship between abundance and distribution of species. *The American Naturalist*, *124*(2), 255-279. https://doi.org/10.1086/284267

Bucci, G., González-Martínez, S. C., Le Provost, G., Plomion, C., Ribeiro, M. M., Sebastiani, F., Alía, R., & Vendramin, G. G. (2007). Range-wide phylogeography and gene zones in *Pinus pinaster* Ait. revealed by chloroplast microsatellite markers. *Molecular Ecology*, *16*(10), 2137-2153. https://doi.org/10.1111/j.1365-294X.2007.03275.x

Burban, C., & Petit, R. J. (2003). Phylogeography of maritime pine inferred with organelle markers having contrasted inheritance. *Molecular Ecology*, *12*(6), 1487-1495. https://doi.org/10.1046/j.1365-294X.2003.01817.x

Caballero, A., Bravo, I., & Wang, J. (2017). Inbreeding load and purging: implications for the

short-term survival and the conservation management of small populations. *Heredity*,

118(2), 177-185. https://doi.org/10.1038/hdy.2016.80

Capblancq, T., Fitzpatrick, M. C., Bay, R. A., Expósito-Alonso, M., & Keller, S. R. (2020).

Genomic prediction of (mal)adaptation across current and future climatic landscapes.

Annual Review of Ecology, Evolution, and Systematics, *51*(1), 245-269.

https://doi.org/10.1146/annurev-ecolsys-020720-042553

Castro, J., Zamora, R., Hódar, J. A., & Gómez, J. M. (2004). Seedling establishment of a

boreal tree species (*Pinus sylvestris*) at its southernmost distribution limit:

consequences of being in a marginal Mediterranean habitat. *Journal of Ecology*,

92(2), 266-277. https://doi.org/10.1111/j.0022-0477.2004.00870.x

Caudullo, G., Welk, E., & San-Miguel-Ayanz, J. (2017). Chorological maps for the main

European woody species. *Data in Brief*, *12*, 662-666.

https://doi.org/10.1016/j.dib.2017.05.007

De La Torre, A. R., Wang, T., Jaquish, B., & Aitken, S. N. (2014). Adaptation and exogenous

selection in a *Picea glauca* × *Picea engelmannii* hybrid zone: implications for

forest management under climate change. *New Phytologist*, *201*(2), 687-699.

https://doi.org/10.1111/nph.12540

De-Lucas, A. I., González-Martínez, S. C., Vendramin, G. G., Hidalgo, E., & Heuertz, M.

(2009). Spatial genetic structure in continuous and fragmented populations of *Pinus*

pinaster Aiton. *Molecular Ecology*, *18*(22), 4564-4576.

https://doi.org/10.1111/j.1365-294X.2009.04372.x

Dussex, N., Morales, H. E., Grossen, C., Dalén, L., & van Oosterhout, C. (2023). Purging and

accumulation of genetic load in conservation. *Trends in Ecology & Evolution*, *38*(10),

961-969. https://doi.org/10.1016/j.tree.2023.05.008

Dussex, N., van der Valk, T., Morales, H. E., Wheat, C. W., Díez-del-Molino, D., von Seth, J., Foster, Y., Kutschera, V. E., Guschanski, K., Rhie, A., Phillippy, A. M., Korlach, J., Howe, K., Chow, W., Pelan, S., Mendes Damas, J. D., Lewin, H. A., Hastie, A. R., Formenti, G., … Dalén, L. (2021). Population genomics of the critically endangered kākāpō. *Cell Genomics*, *1*(1), 100002. https://doi.org/10.1016/j.xgen.2021.100002

Eckert, C. G., Samis, K. E., & Lougheed, S. C. (2008). Genetic variation across species' geographical ranges: the central–marginal hypothesis and beyond. *Molecular Ecology*, *17*(5), 1170-1188. https://doi.org/10.1111/j.1365-294X.2007.03659.x

Ellis, N., Smith, S. J., & Pitcher, C. R. (2012). Gradient forests: calculating importance gradients on physical predictors. *Ecology*, *93*(1), 156-168. https://doi.org/10.1890/11-0252.1

Estoque, R. C., Ishtiaque, A., Parajuli, J., Athukorala, D., Rabby, Y. W., & Ooba, M. (2023). Has the IPCC's revised vulnerability concept been well adopted? *Ambio*, *52*(2), 376-389. https://doi.org/10.1007/s13280-022-01806-z

Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology*, *14*(8), 2611-2620. https://doi.org/10.1111/j.1365-294X.2005.02553.x

Feng, S., Fang, Q., Barnett, R., Li, C., Han, S., Kuhlwilm, M., Zhou, L., Pan, H., Deng, Y., Chen, G., Gamauf, A., Woog, F., Prys-Jones, R., Marques-Bonet, T., Gilbert, M. T. P., & Zhang, G. (2019). The genomic footprints of the fall and recovery of the crested Ibis. *Current Biology*, *29*(2), 340-349.e7. https://doi.org/10.1016/j.cub.2018.12.008

Fitzpatrick, M. C., Chhatre, V. E., Soolanayakanahally, R. Y., & Keller, S. R. (2021). Experimental support for genomic prediction of climate maladaptation using the machine learning approach Gradient Forests. *Molecular Ecology Resources*, *21*(8), 2749-2765. https://doi.org/10.1111/1755-0998.13374

Fitzpatrick, M. C., & Keller, S. R. (2015). Ecological genomics meets community-level

modelling of biodiversity: mapping the genomic landscape of current and future

environmental adaptation. *Ecology Letters*, *18*(1), 1-16.

https://doi.org/10.1111/ele.12376

Fkiri, S., Guibal, F., Elkhorchani, A., Khouja, M. L., Khaldi, A., & Nasr, Z. (2019).

Relationship between climate and growth of two North African varieties of *Pinus*

pinaster Arn. *African Journal of Ecology*, *57*(3), 327-334.

https://doi.org/10.1111/aje.12610

Foden, W., Butchart, S. H. M., Stuart, S. N., Vié, J., Akçakaya, H. R., Angulo, A., DeVantier,

L., Gutsche, A., Turak, E., Cao, L., Donner, S. D., Katariya, V., Bernard, R., Holland,

R. A., Hughes, A., O'Hanlon, S. E., Garnett, S. T., Şekercioğlu, Ç. H., & Mace, G. M.

(2013). Identifying the world's most climate change vulnerable species: a systematic

trait-based assessment of all birds, amphibians and corals. *PLOS ONE*, *8*(6), e65427.

https://doi.org/10.1371/journal.pone.0065427

Foll, M., & Gaggiotti, O. (2008). A genome-scan method to identify selected loci appropriate

for both dominant and codominant markers: a Bayesian perspective. *Genetics*, *180*(2),

977-993. https://doi.org/10.1534/genetics.108.092221

Fréjaville, T., Vizcaíno-Palomar, N., Fady, B., Kremer, A., & Benito Garzón, M. (2019).

Range margin populations show high climate adaptation lags in European trees.

Global Change Biology, 26*(2)*, 484-495. https://doi.org/10.1111/gcb.14881

Frichot, E., Schoville, S. D., Bouchard, G., & François, O. (2013). Testing for associations

between loci and environmental gradients using Latent Factor Mixed Models.

Molecular Biology and Evolution, *30*(7), 1687-1699.

https://doi.org/10.1093/molbev/mst063

Gapare, W. J., Aitken, S. N., & Ritland, C. E. (2005). Genetic diversity of core and peripheral

Sitka spruce (*Picea sitchensis* (Bong.) Carr) populations: implications for

conservation of widespread species. *Biological Conservation*, *123*(1), 113-123.

https://doi.org/10.1016/j.biocon.2004.11.002

Gaston, K. J. (2003). *The structure and dynamics of geographic ranges.* Oxford University

Press, Oxford.

Gaston, K. J. (2009). Geographic range limits: achieving synthesis. *Proceedings of the Royal*

Society B: Biological Sciences, *276*(1661), 1395-1406.

https://doi.org/10.1098/rspb.2008.1480

Gautier, M. (2015). Genome-wide scan for adaptive divergence and association with

population-specific covariates. *Genetics*, *201*(4), 1555-1579.

https://doi.org/10.1534/genetics.115.181453

González-Martínez, S. C., Ridout, K., & Pannell, J. R. (2017). Range expansion compromises

adaptive evolution in an outcrossing plant. *Current Biology*, *27*(16), 2544-2551.e4.

https://doi.org/10.1016/j.cub.2017.07.007

Gougherty, A. V., Keller, S. R., & Fitzpatrick, M. C. (2021). Maladaptation, migration and

extirpation fuel climate change risk in a forest tree species. *Nature Climate Change*,

11(2), 166-171. https://doi.org/10.1038/s41558-020-00968-6

Grossen, C., Guillaume, F., Keller, L. F., & Croll, D. (2020). Purging of highly deleterious

mutations through severe bottlenecks in Alpine ibex. *Nature Communications*, *11*(1),

1001. https://doi.org/10.1038/s41467-020-14803-1

Gugerli, F., Rüegg, M., & Vendramin, G. G. (2009). Gradual decline in genetic diversity in

Swiss stone pine populations (*Pinus cembra*) across Switzerland suggests postglacial

re-colonization into the Alps from a common eastern glacial refugium. *Botanica*

Helvetica, *119*(1), 13-22. https://doi.org/10.1007/s00035-009-0052-6

Günther, T., & Schmid, K. J. (2010). Deleterious amino acid polymorphisms in *Arabidopsis*

thaliana and rice. *Theoretical and Applied Genetics*, *121*(1), 157-168.

https://doi.org/10.1007/s00122-010-1299-4

Hampe, A., & Petit, R. J. (2005). Conserving biodiversity under climate change: the rear edge

matters. *Ecology Letters*, *8*(5), 461-467. https://doi.org/10.1111/j.1461-

0248.2005.00739.x

Hedrick, P. W., & García-Dorado, A. (2016). Understanding inbreeding depression, purging,

and genetic rescue. *Trends in Ecology & Evolution*, *31*(12), 940-952.

https://doi.org/10.1016/j.tree.2016.09.005

Hengeveld, R., & Haeck, J. (1982). The distribution of abundance. I. Measurements. *Journal*

of Biogeography, *9*(4), 303. https://doi.org/10.2307/2844717

Hilfiker, K., Gugerli, F., Schütz, J.-P., Rotach, P., & Holderegger, R. (2004). Low RAPD

variation and female-biased sex ratio indicate genetic drift in small populations of the

dioecious conifer *Taxus baccata* in Switzerland. *Conservation Genetics*, *5*(3),

357-365. https://doi.org/10.1023/B:COGE.0000031144.95293.1b

IPCC (2007). *Contribution of Working Group II to the Fourth Assessment Report of the*

the Intergovernmental Panel on Climate Change. Cambridge University Press,

Cambridge, United Kingdom and New York, NY, USA.

IPCC (2014). *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Summaries,*

frequently asked questions, and cross-chapter boxes. A contribution of Working

Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate

Change. World Meteorological Organization, Geneva, Switzerland.

IPCC (2021). *Climate Change 2021: The Physical Science Basis: Contribution of Working*

Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate

Change. Cambridge University Press, Cambridge, United Kingdom and New York,

NY, USA.

Jaramillo-Correa, J. P., Grivet, D., Terrab, A., Kurt, Y., De-Lucas, A. I., Wahid, N.,

Vendramin, G. G., & González-Martínez, S. C. (2010). The Strait of Gibraltar as a

major biogeographic barrier in Mediterranean conifers: a comparative

phylogeographic survey. *Molecular Ecology*, *19*(24), 5452-5468.

https://doi.org/10.1111/j.1365-294x.2010.04912.x

Jaramillo-Correa, J.-P., Rodríguez-Quilón, I., Grivet, D., Lepoittevin, C., Sebastiani, F.,

Heuertz, M., Garnier-Géré, P. H., Alía, R., Plomion, C., Vendramin, G. G., &

González-Martínez, S. C. (2015). Molecular proxies for climate maladaptation in a

long-lived tree (*Pinus pinaster* Aiton, Pinaceae). *Genetics*, *199*(3), 793-807.

https://doi.org/10.1534/genetics.114.173252

Jaramillo-Correa, J.-P., Grivet, D., Terrab, A., Kurt, Y., De-Lucas, AI., Wahid, N.,

Vendramin, G.G., González-Martínez, S.C. (2010). The Strait of Gibraltar as a major

biogeographic barrier in Mediterranean conifers: a comparative phylogeographic

survey. *Molecular Ecology*, *19*(24), 5452-68. https://doi:10.1111/j.1365-

294X.2010.04912.x

Jia, K., Zhao, W., Maier, P. A., Hu, X., Jin, Y., Zhou, S., Jiao, S., El-Kassaby, Y. A., Wang,

T., Wang, X., & Mao, J. (2020). Landscape genomics predicts climate change-related

genetic offset for the widespread *Platycladus orientalis* (Cupressaceae). *Evolutionary*

Applications, *13*(4), 665-676. https://doi.org/10.1111/eva.12891

Johannesson, K., & André, C. (2006). Life on the margin: genetic isolation and diversity loss

in a peripheral marine ecosystem, the Baltic Sea. *Molecular Ecology*, *15*(8),

2013-2029. https://doi.org/10.1111/j.1365-294X.2006.02919.x

Jost, L. (2008). G_{ST} and its relatives do not measure differentiation. *Molecular Ecology*,

17(18), 4015-4026. https://doi.org/10.1111/j.1365-294X.2008.03887.x

Kimura, M., Maruyama, T., & Crow, J. F. (1963). The mutation load in small populations.

Genetics, *48*(10), 1303–1312. https://doi.org/10.1093/genetics/48.10.1303

Kimura, M., & Ohta, T. (1969). The average number of generations until fixation of a mutant

gene in a finite population. *Genetics*, *61*(3), 763-771.

https://doi.org/10.1093/genetics/61.3.763

Kolzenburg, R. (2022). The direct influence of climate change on marginal populations: a

review. *Aquatic Sciences*, *84*(2), 24. https://doi.org/10.1007/s00027-022-00856-5

Lande, R. (1988). Genetics and demography in biological conservation. *Science*, *241*(4872),

1455-1460. https://doi.org/10.1126/science.3420403

Latta, R. G. (1995). The effects of embryo competition with mixed mating on the genetic

load in plants. *Heredity*, *75*(6), 637-643. https://doi.org/10.1038/hdy.1995.183

Lind, B. M., Candido-Ribeiro, R., Singh, P., Lu, M., Vidaković, D. O., Booker, T. R.,

Whitlock, M. C., Yeaman, S., Isabel, N., & Aitken, S. N. (2023). How useful is

genomic data for predicting maladaptation to future climate? *bioRxiv (Cold Spring*

Harbor Laboratory). [Preprint]. https://doi.org/10.1101/2023.02.10.528022

Lira-Noriega, A., & Manthey, J. D. (2014). Relationship of genetic diversity and niche

centrality: a survey and analysis. *Evolution*, *68*(4), 1082-1093.

https://doi.org/10.1111/evo.12343

Lynch, M., Conery, J., & Bürger, R. (1995). Mutation accumulation and the extinction of

small populations. *The American Naturalist*, *146*(4), 489-518.

https://doi.org/10.1086/285812

662 Marchi, M., Castellanos-Acuña, D., Hamann, A., Wang, T., Ray, D., & Menzel, A. (2020).

663  ClimateEU, scale-free climate normals, historical time series, and future projections

664  for Europe. *Scientific Data*, *7*(1), 428. https://doi.org/10.1038/s41597-020-00763-0

665 Marques, H., Pinto, G., Pinto, P., Teixeira, C. (2012). Regiões de Proveniência. Portugal.

666  ICNF, 47-51.

667 Milesi, P., Kastally, C., Dauphin, B., Cervantes, S., Bagnoli, F., Budde, K. B., Cavers, S.,

668  Fady, B., Faivre-Rampant, P., González-Martínez, S. C., Grivet, D., Gugerli, F.,

669  Jorge, V., Kupin, I. L., Ojeda, D. I., Olsson, S., Opgenoorth, L., Pinosio, S., Plomion,

670  C.,. . . Pyhäjärvi, T. (2023). Synchronous effective population size changes and

671  genetic stability of forest trees through glacial cycles. *bioRxiv (Cold Spring Harbor*

672  *Laboratory)*. [Preprint]. https://doi.org/10.1101/2023.01.05.522822

673 Naydenov, K. D., Alexandrov, A., Matevski, V., Vasilevski, K., Naydenov, M. K., Gyuleva,

674  V., Carcaillet, C., Wahid, N. and Kamary, S. (2014). Range-wide genetic structure of

675  maritime pine predates the last glacial maximum: evidence from nuclear DNA.

676  *Hereditas*, *151*(1), 1–13. https://doi.org/10.1111/j.1601-5223.2013.00027.x

677 Nicholson, A. J. (1958). Dynamics of insect populations. *Annual Review of Entomology*, *3*(1),

678  107-136. https://doi.org/10.1146/annurev.en.03.010158.000543

679 Peischl, S., Dupanloup, I., Kirkpatrick, M., & Excoffier, L. (2013). On the accumulation of

680  deleterious mutations during range expansions. *Molecular Ecology*, *22*(24),

681  5972-5982. https://doi.org/10.1111/mec.12524

682 Picard, N., Marchi, M., Serra-Varela, M. J., Westergren, M., Cavers, S., Notivol, E., Piotti,

683  A., Alizoti, P., Bozzano, M., González-Martínez, S. C., Grivet, D., Aravanopoulos, F.

684  A., Vendramin, G. G., Ducci, F., Fady, B., & Alía, R. (2022). Marginality indices for

685  biodiversity conservation in forest trees. *Ecological Indicators*, *143*, 109367.

686  https://doi.org/10.1016/j.ecolind.2022.109367

Pironon, S., Villellas, J., Morris, W. F., Doak, D. F., & García, M. B. V. (2015). Do

geographic, climatic or historical ranges differentiate the performance of central

versus peripheral populations? *Global Ecology and Biogeography*, *24*(6), 611-620.

https://doi.org/10.1111/geb.12263

Pironon, S., Papuga, G., Villellas, J., Angert, A. L., García, M. B. V., & Thompson, J. D.

(2016). Geographic variation in genetic and demographic performance: new insights

from an old biogeographical paradigm. *Biological Reviews*, *92*(4), 1877-1909.

https://doi.org/10.1111/brv.12313

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using

multilocus genotype data. *Genetics*, *155*(2), 945-959.

https://doi.org/10.1093/genetics/155.2.945

R Core Team. (2022). *R: A Language and Environment for Statistical Computing*. R

Foundation for Statistical Computing. https://www.R-project.org/

Razgour, O., Taggart, J. B., Manel, S., Juste, J., Ibáñez, C., Rebelo, H., Alberdi, A., Jones,

G., & Park, K. (2017). An integrated framework to identify wildlife populations under

threat from climate change. *Molecular Ecology Resources*, *18*(1), 18-31.

https://doi.org/10.1111/1755-0998.12694

Rellstab, C., Dauphin, B., & Expósito-Alonso, M. (2021). Prospects and limitations of

genomic offset in conservation management. *Evolutionary Applications*, *14*(5),

1202-1212. https://doi.org/10.1111/eva.13205

Richards, O. W. (1961). The theoretical and practical study of natural insect populations.

Annual Review of Entomology, *6*(1), 147-162.

https://doi.org/10.1146/annurev.en.06.010161.001051

Robinson, J. A., Brown, C., Kim, B. Y., Lohmueller, K. E., & Wayne, R. K. (2018). Purging

of strongly deleterious mutations explains long-term persistence and absence of

inbreeding depression in island foxes. *Current Biology*, *28*(21), 3487-3494.e4.

https://doi.org/10.1016/j.cub.2018.08.066

Rousset, F. (2008). Genepop'007: a complete re-implementation of the genepop software for

Windows and Linux. *Molecular Ecology Resources*, *8*(1), 103-106.

https://doi.org/10.1111/j.1471-8286.2007.01931.x

Sachdeva, H., Olusanya, O., & Barton, N. (2022). Genetic load and extinction in peripheral

populations: the roles of migration, drift and demographic stochasticity. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *377*(1846).

https://doi.org/10.1098/rstb.2021.0010

Santos, A. S., & Gaiotto, F. A. (2020). Knowledge status and sampling strategies to

maximize cost-benefit ratio of studies in landscape genomics of wild plants. *Scientific*

Reports, *10*(1), 3706. https://doi.org/10.1038/s41598-020-60788-8

Sexton, J. P., McIntyre, P. J., Angert, A. L., & Rice, K. J. (2009). Evolution and ecology of

species range limits. *Annual Review of Ecology, Evolution, and Systematics*, *40*(1),

415-436. https://doi.org/10.1146/annurev.ecolsys.110308.120317

Sexton, J. P., Hangartner, S. B., & Hoffmann, A. A. (2013). Genetic isolation by environment

or distance: which pattern of gene flow is most common? *Evolution*, *68*(1), 1-15.

https://doi.org/10.1111/evo.12258

Soille, P., & Vogt, P. (2009). Morphological segmentation of binary patterns. *Pattern*

Recognition Letters, *30*(4), 456-459. https://doi.org/10.1016/j.patrec.2008.10.015

Soulé, M.E. (1973). The epistasis cycle: a theory of marginal populations. *Annual Review of*

Ecology and Systematics, *4*(1), 165-187.

https://doi.org/10.1146/annurev.es.04.110173.001121

van den Wollenberg, A. L. (1977). Redundancy analysis an alternative for canonical

correlation analysis. *Psychometrika*, *42*(2), 207-219.

https://doi.org/10.1007/BF02294050

Vendramin, G., Anzidei, M., Madaghiele, A., Bucci, G. (1998). Distribution of genetic

diversity in *Pinus pinaster* Ait. as revealed by chloroplast microsatellites. *Theoretical*

and Applied Genetics, 97, 456–463. https://doi.org/10.1007/s001220050917

Wahid, N., González-Martínez, S. C., Hadrami, I. E., & Boulli, A. (2004). Genetic Structure

and variability of natural populations of maritime pine (*Pinus pinaster* Aiton) in

Morocco. *Silvae Genetica*, *53*(1-6), 93-99. https://doi.org/10.1515/sg-2004-0017

Wahid, N., González-Martínez, S. C., El Hadrami, I., & Boulli, A. (2006). Variation of

morphological traits in natural populations of maritime pine (*Pinus pinaster* Ait.) in

Morocco. *Annals of Forest Science*, *63*(1), 83-92.

https://doi.org/10.1051/forest:20050100

Weir, B. S., & Cockerham, C. C. (1984). Estimating F-statistics for the analysis of population

structure. *Evolution*, *38*(6), 1358. https://doi.org/10.2307/2408641

Westergren, M., Božič, G., Kraigher, H. (2018). Genetic diversity of core *vs.* peripheral

Norway spruce native populations at a local scale in Slovenia. *iForest -*

Biogeosciences and Forestry, *11*(1), 104-110. https://doi.org/10.3832/ifor2444-011

Whittaker, J. B. (1971). Population changes in *Neophilaenus lineatus* (L.) (Homoptera :

Cercopidae) in different parts of its range. *The Journal of Animal Ecology*, *40*(2), 425.

https://doi.org/10.2307/3253

Willson, M. F., & Burley, N. (1983). *Mate Choice in Plants (MPB-19), Volume 19: Tactics,*

Mechanisms, and Consequences. (MPB-19). Princeton University Press.

https://doi.org/10.2307/j.ctvx5wbss

Young, A.G., Boyle, T., & Brown, T.L. (1996). The population genetic consequences of

habitat fragmentation for plants. *Trends in Ecology & Evolution*, *11*(10), 413-418.

https://doi.org/10.1016/0169-5347(96)10045-8

Yuan, S., Shi, Y., Zhou, B., Liang, Y., Chen, X., An, Q., Fan, Y., Shen, Z., Ingvarsson, P. K.,

& Wang, B. (2023). Genomic vulnerability to climate change in *Quercus acutissima*,

a dominant tree species in East Asian deciduous forests. *Molecular Ecology*, *32*(7),

1639-1655. https://doi.org/10.1111/mec.16843

Figure legends

Figure 1. Genetic pools in *P. pinaster*. Pie charts depict membership proportions of each

genetic cluster (K=10) for each studied population, calculated by STRUCTURE v2.3.4. The

natural distribution of the species is shadowed in light green (see details in Figure S1).

Population codes are only provided for those populations specifically mentioned in the main

text (see full population information, including population names and geographical

coordinates, in Table S1).

Figure 2. Geographical distribution of a) genetic diversity (1-Q_{inter}) and b) genetic

differentiation (population-specific F_{ST}), and two marginality indices involved in significant

correlations with these genetic indicators, c) centroid index and d) second nearest-core index,

for *P. pinaster* populations. See Table S1 for population information, including population

names and geographical coordinates.

Figure 3. a) Geographical distribution of genomic offset for *P. pinaster* populations and b)

correlation between the genomic offset and the ecological index based on climate distances

(standardized values), as shown by linear regression in *ggplot2* R package. Population codes

are only provided for populations specifically mentioned in the main text (see full population

information, including population names and geographical coordinates, in Table S1).

Table 1. Geographical (*geo*) and demo-historical (*histo*) marginality indices computed for maritime pine populations; '(-)' and '(+)' indicate that

higher values of the index correspond to lower and higher population marginality, respectively; '(-)(+)' indicates that both higher or lower values

of the index correspond to higher population marginality. Adapted after Picard et al. (2022).

Marginality index	Category	Link with marginality	Description
Area	*geo*	(-)	Size in ha of the core (i.e., continuous patches of the species' presence) that is the nearest to the location of the studied population.
Gravity	*geo*	(-)	Spatially continuous alternative to the area index, computed as the weighted mean of the core areas with weights inversely proportional to the distance squared from the location to the cores.
Centroid	*geo*	(+)	Eccentricity computed as the cost distance from the centroid of the species distribution taking into account the terrestrial and marine connectivity. Cost distances are based on a conductance matrix (the inverse of a resistance matrix) reflecting the permeability of land and water cells to gene flow. This index is expected to capture the level of long-distance gene flow.
Edge	*geo*	(-)	Distance to the nearest border of the species distribution.
Isolation	*geo*	(+)	Isolation with respect to the core species distribution, computed as the distance from the focal population to the nearest core greater than 100 ha and further than 50 km away.
Second nearest-core	*geo*	(+)	Spatially continuous alternative to the isolation index, computed as the distance from a location to the second nearest-core greater than 100 ha.

North-South	*histo*	(-) (+)	Proximity to the species rear-edge (southernmost limit of the species distribution) or leading-edge (northernmost limit of the species distribution) along a latitudinal gradient.
East-West	*histo*	(-) (+)	Proximity to the species leading-edge or rear-edge along a longitudinal gradient.

794 **Table 2**. Models evaluating the effect of population marginality on the estimated genetic indicators; a) Models without gene pool effect (M1-

795 M5); b) Models including gene pool effect as a random factor (M6-M10). Significant fixed effect(s) are listed with their associated point

796 estimates and 95% confidence intervals [in brackets]. R^2: variance explained by fixed factors (M1-M5) or both fixed (marginal R^2) and random

797 (conditional R^2) factors (M6-M10); ns: not significant.

799 a) Models without gene pool effect.

Model [R^2]	Genetic indicator	Fixed-effect #1	Estimate [95% CIs]	Fixed-effect #2	Estimate [95% CIs]	Fixed-effect #3	Estimate [95% CIs]	Interaction	Estimate [95% CIs]
M1 [0.59]	*Genetic diversity*	Centroid	-1.01 [-1.34, -0.67]	Second nearest-core	-1.47 [-2.43, -0.51]	North-South	1.61 [0.66, 2.55]	Second nearest-core × North-South	1.44 [0.50, 2.38]
M2 [<0.01]	*Inbreeding*	ns		ns		ns		ns	
M3 [0.66]	*Population-specific F_{ST}*	Centroid	1.05 [0.74, 1.35]	Second nearest-core	1.43 [0.55, 2.31]	North-South	-1.90 [-2.76, -1.04]	Second nearest-core × North-South	-1.66 [-2.51, -0.80]
M4 [0.12]	*Recessive genetic load*	Centroid	-0.69 [-1.08, -0.30]	ns		ns		ns	
M5 [0.27]	*Genetic offset*	Ecological index	0.48 [0.28, 0.68]	ns		ns		ns	

b) Models with gene pool effect as random factor.

803
804

Model [R^2]	Genetic indicator	Fixed-effect #1	Estimate [95% CIs]	Fixed-effect #2	Estimate [95% CIs]	Interaction	Estimate [95% CIs]
M6 [0.41, 0.69]	*Genetic diversity*	Centroid	-1.04 [-1.58, -0.50]	North-South	1.04 [0.34, 1.75]	Centroid × North-South	0.75 [0.25, 1.24]
M7 [<0.01]	*Inbreeding*	ns		ns		ns	
M8 [0.31, 0.86]	*Population-specific F_{ST}*	Centroid	0.74 [0.21, 1.28]	North-South	-1.40 [-2.15, -0.65]	Centroid × North-South	-0.96 [-1.45, -0.46]
M9 [<0.01]	*Recessive genetic load*	ns		ns		ns	
M10 [0.22, 0.40]	*Genetic offset*	Ecological index	0.44 [0.24, 0.65]	ns		ns	

Figure 1

816   **Figure 2**

Figure 3