1   **The genetic consequences of population marginality: a case study in**

2   **maritime pine**

3
Adélaïde Theraroz[1], Carlos Guadaño-Peyrot[2], Juliette Archambeau[1,3], Sara Pinosio[4],

Francesca Bagnoli[4], Andrea Piotti[4], Camilla Avanzi[4], Giovanni G. Vendramin[4], Ricardo

Alía[2], Delphine Grivet[2], Marjana Westergren[5], Santiago C. González-Martínez[1,*]

[1]INRAE, Univ. Bordeaux, BIOGECO, Cestas, France

[2]Department of Forest Ecology and Genetics, Institute of Forest Science (ICIFOR-INIA),

CSIC, Carretera de la Coruña km 7,5, Madrid, Spain

[3]UK Centre for Ecology & Hydrology, Bush Estate, Penicuik, United Kingdom

[4]National Research Council, Institute of Biosciences and BioResources, Sesto Fiorentino,

Italy

[5]Slovenian Forestry Institute, Večna Pot 2, Ljubljana, Slovenia

*Corresponding author:

Santiago C. González-Martínez

UMR 1202 INRAE – Univ. Bordeaux

69 route d'Arcachon, F-33610 Cestas, France

Tel: +33 (0) 5 35 38 53 20

santiago.gonzalez-martinez@inrae.fr

Abstract

Aim: Marginal tree populations, either those located at the edges of the species' range or in suboptimal environments, are often a valuable genetic resource for biological conservation. However, there is a lack of knowledge about the genetic consequences of population's marginality, estimated across entire species' ranges. Our study addresses this gap by providing information about several genetic indicators and their variability in marginal and core populations identified using quantitative marginality indices.

Location: Southwestern Europe and North Africa.

Methods: Using 10,185 SNPs across 82 populations of maritime pine (*Pinus pinaster* Ait.), a widespread and economically important conifer characterised by a fragmented range, we modelled the relationship of seven genetic indicators potentially related to population evolutionary resilience, namely genetic diversity (based on all SNPs and two kind of outlier loci), inbreeding, genetic differentiation, recessive genetic load and genomic offset, with population geographical, demo-historical and ecological marginality (as estimated by nine quantitative indices). Models were constructed for both regional (introducing gene pool as random factor) and range-wide spatial scales.

Results: We showed a trend towards decreasing overall genetic diversity (albeit not necessarily for genetic diversity estimates based on outlier loci) and increasing differentiation with geographic marginality, supporting the centre-periphery hypothesis (CPH). Moreover, we found no correlation between population inbreeding and marginality, while geographically marginal populations had a lower recessive genetic load (only models without

the gene pool effect) and ecologically marginal populations had a higher genomic offset,

suggesting higher maladaptation to future climate.

Main conclusions: Overall, our results indicate that marginal populations could be at higher

risk of maladaptation to climate change than core populations, despite reduced levels of

genetic load, a risk that is exacerbated by typically small effective population sizes and

increasing human impact.

Keywords: centre-periphery hypothesis, ecological modelling, forest genetic resources,

genetic indicators, marginal populations, Mediterranean and Atlantic regions

Introduction

All species thrive and reproduce within an environmentally limited geographic area, which sets the boundaries of their range. Current global warming may restrict the climatic suitability of some parts of the species' ranges and, together with other factors (e.g., typically competitive biotic interactions, Loehle, 1998; but also positive, often unaccounted-for, interactions with other species, Stephan et al., 2021), modify their geographic and ecological margins. The 'centre-periphery hypothesis' (hereafter CPH), a major paradigm in biogeography that aims to disentangle the genetic, demographic, and ecological causes of species' range limits (Gaston, 2009; Sexton et al., 2009), defines marginality as the level of geographic isolation from the species' centre of distribution, which in turn is related to the species' suitability for its environment (Brown, 1984; Hengeveld & Haeck, 1982). According to the CPH, marginal populations are expected to be less abundant and more prone to extinction than those in the centre, due to harsher environmental conditions at the periphery (Birch, 1957; Gaston, 2003; Nicholson, 1958; Richards, 1961; Whittaker, 1971). However, as specific environmental conditions in the core of the species distribution may also induce harsher environmental conditions than in the periphery (e.g., temperature extremes, rugged topography, peculiar edaphic features), Soulé (1973) distinguished between geographical and ecological marginality when describing centre-periphery gradients, defining ecological marginality as a population's exposure to extreme environmental variables irrespectively of their geographical location.

While several studies have identified marginal populations based on different criteria (e.g., for maritime pine, Alía et al., 1996; Burban & Petit, 2003), we often lack evidence on the extent to which population marginality is associated with particular genetic features. Classic studies at the range-wide scale suggested a general trend for lower genetic diversity and higher genetic differentiation in marginal populations, but they typically did not

distinguish between geographical and ecological marginality (Eckert et al., 2008;

Johannesson & André, 2006; Pironon et al., 2015). More recent studies often called for

decoupling the roles of geography and ecology in shaping range-wide patterns of genetic

variation (Lira-Noriega & Manthey, 2014; Sexton et al., 2013). To that end, a clear

distinction needs to be made between different kinds of population marginality. In a recent

multispecies study, Picard et al. (2022) evaluated the ability of quantitative measures to

distinguish between geographical, demo-historical (i.e., related to the demographic history of

the target species) and ecological marginality.

Past global climate dynamics rather than demographic stochasticity seem to have

played a crucial role in the establishment of current range limits (e.g., Hampe & Petit, 2005,

for forest trees). Postglacial migrations after the Last Glacial Maximum (LGM) would have

resulted in a mosaic of relatively small and isolated populations at the rear and, in a lesser

extent, at the front edges of the expansion. In this context, the level of connectivity among

marginal populations and between a given marginal population and the species distribution

core is especially important (Sachdeva et al., 2022), as gene flow may increase genetic

diversity and reduce differentiation (e.g., Lynch et al., 1995; Young et al., 1996). Gene flow

can also bring adaptive alleles and contribute to the evolutionary rescue of small, isolated

populations by buffering the effect of genetic drift, and reducing the fixation and

accumulation of deleterious alleles within populations (Sachdeva et al., 2022). However,

marginal populations may also carry specific alleles derived from local adaptation to atypical

environments, constituting valuable genetic resources, and making the contribution of

external gene flow harmful (i.e., the so-called 'migration load'; Kimura et al., 1963).

Overall, marginal populations are considered to be more vulnerable to climate change

than core populations (Kolzenburg, 2022; Soulé, 1973). Marginal populations are expected to

accumulate deleterious variants (i.e., genetic load), a process governed by effective

population size, N_e (Kimura & Ohta, 1969). However, genetic purging due to inbreeding

tends to reduce genetic load over time, even in relatively small populations (Hedrick &

Garcia-Dorado, 2016), and the overall outcome is context-dependant (Sachdeva et al., 2021).

Current developments in population genomics have provided metrics to estimate

maladaptation to future climates e.g., by estimating genomic 'offsets' or 'gaps' (a measure of

the mismatch in genotype-climate association between current and potential future climates,

Capblancq et al., 2020; Fitzpatrick & Keller, 2015; Gougherty et al., 2021; Rellstab et al.,

2021). Studies aimed at validating genomic offset predictions with data from common garden

experiments and natural populations have shown that populations with higher genomic offset

exhibit a reduction in population growth performance (e.g., Fitzpatrick et al., 2021; for

balsam poplar) and survival (Archambeau et al., 2024; for maritime pine), and concluded on

the potential of this indicator to provide an estimate of the degree of expected maladaptation

to future climate. Thus, despite several limitations (Rellstab et al., 2021; Láruson et al. 2022;

Ahrens et al., 2023; Lotterhos, 2023), the calculation of genomic offsets may still enable for

much needed systematic studies on the connection between population marginality and

maladaptation in the face of climate change (Archambeau et al., 2024).

Maritime pine (*Pinus pinaster* Ait., Pinaceae) is an outcrossing, wind-pollinated

conifer, with a widespread but fragmented natural distribution in southwestern Europe and

North Africa, covering a wide range of contrasted environments from coastal dunes next to

the Atlantic Ocean in France to the High Atlas Mountains in Morocco. Maritime pine

population genetic structure is a consequence of historical and current dynamics of range

expansion-contraction, resulting in distinct genetic clusters (Jaramillo-Correa et al., 2015).

This species is also characterised by a fragmented distribution due to human-induced habitat

loss and ecological disturbances such as forest fires (De-Lucas et al., 2009). Nowadays, some

maritime pine populations are found in ecologically marginal environments (e.g., under very

dry conditions in southern Spain and northern Morocco). In addition, the species' range margins are characterised by small, geographically isolated populations, in particular in the southern and eastern parts of the distribution (Alía et al., 1996; Wahid et al., 2004). Reduced dispersal with distant core populations, coupled with demographic and environmental stochasticity, may push such populations into an 'extinction vortex' (Lande, 1988). Few studies have focused on describing the particular genetic characteristics of marginal populations of maritime pine (Salvador et al., 2000; Wahid et al., 2004; González-Martínez et al., 2007), whose potentially valuable genetic resources could be lost in the near future.

In this study, we used 10,185 Single Nucleotide Polymorphisms (SNPs) genotyped in 82 maritime pine populations (1,510 individuals), including ecologically and geographically marginal ones, to compute seven genetic indicators potentially related to population evolutionary resilience (defined as the property of an ecosystem to undergo adaptive evolution in response to biotic or abiotic disturbances; Sgrò et al., 2010) and correlated them with quantitative measures of marginality. The main objective of this study is to assess the relationship between marginality and population genetic features at the scale of the whole species range, by testing predictions of the CPH and adding new elements to its general framework. More specifically, we i) assessed the losses of genetic diversity and increases of genetic differentiation in marginal populations, distinguishing overall genetic diversity from that estimated using different kind of outlier loci; ii) evaluated the levels of accumulation of recessive genetic load based on counts of deleterious alleles; and iii) tested whether marginal populations are maladapted to future climate conditions, applying genomic approaches that consider the contribution of pre-adapted variants to future climates (i.e., genomic offset models).

Materials and Methods

Plant materials and molecular markers

Needles were collected from 1,510 individuals in 82 maritime pine populations covering all previously identified gene pools throughout the species range (Jaramillo-Correa et al., 2015; Milesi et al., 2023). Populations were selected based on gene pool size, density of maritime pine in the area, and representativeness in terms of climatic conditions specific to their geographical position (e.g., orography), as to achieve a regular sampling across the full distribution range of the species. This sampling is, to date, the most complete in the species (see Figure 1 and Table S1 in Supplementary Information), and includes several populations from the distribution margins, as well as isolated populations that have not been considered in genetic studies before. Population Cómpeta (COM), with only three samples, was removed from all data analyses but the gene-environment association (GEA) methods used to estimate the genomic offset. The accuracy of landscape genomic approaches, such as GEA, is highly improved by increasing the number of populations and environments, while being less sensitive to unbalanced sampling designs (Santos & Gaiotto, 2020). In addition, the two stands of Maures population (MAU) (see Table S1) were kept separate for these analyses as they were sampled at different altitudes.

DNA was extracted using the DNeasy 96 plant kit (QIAGEN, Germany), following the manufacturer's instructions. All the samples were genotyped for SNPs using the multispecies 4TREE Axiom array (Thermo Fisher Scientific, USA). For maritime pine, the 4TREE Axiom array combines SNPs identified in two previous studies: the 9K Illumina Infinium array generated by Plomion et al. (2016) and the exome capture experiment used in Milesi et al. (2023). The new array has a conversion rate of 79%, as well as 99% genotype reproducibility (based on genotyping of duplicated samples). Apart from potentially neutral genetic polymorphisms, this array comprises also SNPs from candidate genes that showed

signatures of natural selection or significant environmental associations with climate at the

range-wide spatial scale, orthologs for gene families with important adaptive functions in

model species, and coding regions with differential expression under biotic and abiotic stress

in maritime pine (see details in Plomion et al., 2016; Milesi et al., 2023). Only SNPs with

high-quality scoring following the Best Practices Workflow implemented in the Axiom™

Analysis Suite v5.2 were selected and filtered by missing data (<30%), yielding a total of

10,185 SNPs. To assess the impact of missing data in the calculation of genetic indicators,

SNPs were also filtered for missing data using a 5% threshold, resulting in a subset of 6,390

SNPs. SNP annotation based on SnpEff v5.1 (Cingolani et al., 2012) was retrieved from Cahn

(2023) for a set of 1,325 SNPs in common with our study.

Data analysis

Population marginality indices

To assess population marginality, we first produced a new distribution map for maritime pine

that includes only natural populations (Figure S1), building on that of Caudullo et al. (2017)

but adding information from National Forest Inventories and specific publications for less-

known parts of the distribution range (Alía et al., 1996; Marques et al., 2012; Abad-Viñas et

al., 2016; Fkiri et al., 2019; Wahid et al., 2004, 2006). Second, we computed eight

quantitative marginality indices that consider both geographical distribution and demographic

history, following Picard et al. (2022) (see Table 1 and Table S2 in Supplementary

Information). The demo-historical indices, which are related to the postglacial colonization

history of the species (see, e.g., Bucci et al., 2007; Jaramillo-Correa et al., 2015), and the

centroid index were computed from the distribution map. The centroid of the species'

distribution corresponded to the location whose geographic coordinates were the average of

the geographic coordinates of all locations where the species is present (in our case, eastern

Spain, a known glacial refugia of maritime pine; Salvador et al., 2000; see Figure S2). Then,

the centroid index was defined as the 'cost distance' between any population and the centroid

location. Cost distances were computed using a conductance matrix (the inverse of a

resistance matrix), reflecting the conductance of gene flow. In the conductance matrix, sea

cells were assigned low conductance, land cells where the species was absent were assigned

intermediate conductance, and land cells where the species was present were assigned high

conductance (see Table 1). This index reflects the level of long-distance gene flow between a

given population and the geographical core of the species distribution. The other

geographical indices relied on a morphological spatial pattern analysis (MSPA), which

considers a binary image (1-presence/0-absence and NA for water) with emphasis on the

connectivity within the image (Soille & Vogt, 2009). Presence/absence maps were structured

in three categories (cores, edges, and other classes including loops, islets, bridges and

branches) using BioManager package in R version 4.2.2. Third, we calculated an index of

ecological marginality (henceforth 'ecological index') based on climatic data for the period

1901-1970, as follows. For each population, we extracted the climatic information provided

by the Climate Downscaling Tool (ClimateDT, https://www.ibbr.cnr.it/climate-dt/) for the

Summer Heat Moisture (SHM) aridity index, an indicator of exposure to drought (see, e.g.,

De La Torre et al., 2014; Marchi et al., 2020), and temperature (bio4) and precipitation

seasonality (bio15). This set of climatic variables was selected because it best explained the

climatic variation across the species range in previous studies (e.g., Archambeau et al., 2024).

The ecological index was constructed by computing the standardized Euclidean distance for

SHM, bio4 and bio15 between each population and the overall average. Thus, this index

represents the climatic distance of the population from the average climate (Table S2).

Finally, we reduced the set of marginality indices by removing those that were highly

correlated based on Pearson's correlation coefficients (Pearson's correlation ≥ 0.6) and a

principal component analysis (PCA) using *FactoMineR* package in R v4.2.2 (Figure S3).

Population genetic structure and genetic indicators

Population genetic structure was evaluated using the Bayesian clustering algorithm

implemented in STRUCTURE v2.3.4 (Pritchard et al., 2000). To assess the optimal

individual's assignment probabilities ($Q_{ancestry}$) in K genetic clusters (or gene pools), we ran

models from K=1 (no structure) to K=10 with a burn-in length of 100,000 and run lengths of

200,000 MCMC steps. The number of K that best describes the genetic structure was

determined based on the delta K method (Evanno et al., 2005) and the visual observation of

bar plots.

All genetic indicators were computed at the population level using the full dataset of

10,185 SNPs (Table S3). In a few cases where more than one stand was sampled for a

population (see Table S1), genetic indicator values were averaged. The computation of

genetic indicators was robust to the inclusion of missing data, as shown by the high

correlations (Pearson's correlation > 0.7) with genetic indicator estimates computed using the

dataset (6,390 SNPs) with missing data lower than 5% (Figure S4a,b).

Genetic diversity was estimated as 1-Q_{inter}, with Q_{inter} being the observed frequencies

of identical pairs of alleles among individuals within populations, using GenePop v4.7.5

(Rousset, 2008), after standard correction for sample size (N), using (N/N-1). Notice that this

estimate averages over monomorphic and polymorphic loci. In addition to overall genetic

diversity, we also calculated two indicators of genetic diversity based on outlier loci. The first

was calculated on the basis of climate-associated (GEA) outliers (i.e., 73 outlier SNPs

selected for the genomic offset computation, described below). The second indicator was

calculated on the basis of general outliers due to unknown factors (i.e., 151 outlier SNPs

common to two environment-independent F_{ST}-outlier-detection methods, as implemented in

the R package *pcadapt* and BayeScan v2.1; Foll & Gaggiotti, 2008). Population inbreeding

(F_{IS}) was estimated following Weir & Cockerham (1984). Population-specific divergence

was estimated as the genetic differentiation (population-specific F_{ST}) of each population from

a common ancestral gene pool using BayeScan v2.1 (Foll & Gaggiotti, 2008), as well as

Jost's *D* (Jost, 2008) estimated with MMOD package in R v4.2.2.

    Finally, we computed two additional genetic indicators more specifically related to

potential maladaptation to climate change, i.e., the recessive genetic load and the genomic

offset. The recessive genetic load represents the accumulation of predicted deleterious

mutations in the population standardized by the population genetic diversity. This statistic

was estimated by counting different kind of mutations (annotated by SnpEff v5.1; Cingolani

et al., 2012) averaged over individuals, as the number of derived moderate- (i.e., non-

synonymous) and high-impact (i.e., loss of function) mutations in homozygosity divided by

the number of derived low-impact (i.e., synonymous) mutations in homozygosity, following

González-Martínez et al. (2017). The genomic offset is estimated as the change in genetic

composition required to maintain the current gene-climate relationships under future climates

(see Fitzpatrick & Keller, 2015) and thus captures the degree of maladaptation a population

will undergo when the environment to which it is currently adapted will change, either from a

spatial or temporal perspective (Rellstab et al., 2021; Lotterhos, 2024). Briefly, we first

identified outlier SNPs for climate adaptation with two univariate GEA methods, BAYPASS

(Gautier, 2015) and Latent Factor Mixed Model (LFMM; Frichot et al., 2013), and three

multivariate ones, Redundancy Analysis (RDA; Capblancq et al., 2020), partial RDA (Van

Den Wollenberg, 1977) and Gradient Forest (GF; Ellis et al., 2012). Then, the genomic offset

was estimated using the set of outlier SNPs identified by at least two methods (see the

GitHub detailing this analysis and referenced below) and the GF approach, which showed the

best empirical validation in a previous study based on a smaller sample of populations and

SNPs (Archambeau et al., 2024), and six climatic variables related to maritime pine expected

exposure to climate change (Table S4; see also Archambeau et al., 2024). Future climates for

2070 were described using the predictions from the moderately alarming shared socio-

economic pathway (SSP) SSP3-7.0 and five global circulation models (GCMs; IPCC, 2021).

As genomic offset predictions across GCMs were highly correlated (Pearson's correlation

coefficient >0.75; Figure S5), we used population averages for the five GCMs. Additional

details and scripts are available at https://anonymous.4open.science/r/ReadyToGO_Pinpin-

FA56/README.md. All analyses were undertaken under R v4.2.2 (R Core Team, 2022).

Effects of population marginality on genetic indicators

We estimated the relationship between genetic indicators and population marginality by

fitting two series of seven linear mixed-models with pairwise interaction terms (one model

for each genetic indicator), using the R packages LME4 and LMER, respectively. Models M1 to

M7 included population marginality indices as fixed effects irrespectively of the gene pool of

origin while models M8 to M14 also included the gene pool of origin as random effects.

Random effects in linear mixed-models allow the inclusion of non-independent data from a

nested structure (populations sampled within gene pools), allowing each level of the grouping

factor (gene pool) to have its own random intercept. The gene pools with a single population

(Fuencaliente, FUE; and Point Cires, PCI) were assigned to the geographically closest gene

pool (Southeastern Spain and Morocco, respectively). Model goodness-of-fit was evaluated

with R^2 and both the Akaike Information Criterion (AIC) and the Bayesian Information

Criterion (BIC). A visual evaluation of model fit to data was also performed using diagnostic

plots (QQ and residual plots; Figures S12 to S17). Then, the best models were selected by

considering goodness-of-fit (higher R^2 and lower AIC/BIC) and parsimony criteria (i.e.,

including only significant effects at $\alpha = 0.01$).

Results

Marginality indices

Pairwise Pearson's correlations and PCA identified some strongly correlated marginality

indices (Figure S3). For example, the second nearest-core was positively correlated with the

isolation index (Pearson's correlation of 0.64) and negatively correlated with the edge index

(Pearson's correlation of -0.67), and the ecological index was negatively correlated with the

East-West index (Pearson's correlation of -0.51). Thus, only five indices with low correlation

(<0.6) were retained for further analysis, namely three geographical indices (centroid, second

nearest-core and gravity; see definitions in Table 1), one demo-historical (North-South), and

the ecological index based on climate distances (Figure S6). Interestingly, the centroid index

was positively correlated with the longitude (Pearson's correlation of 0.70) and the North-

South index was negatively correlated with the elevation (Pearson's correlation of -0.56),

with southern populations being, generally, at higher elevation (Figure S3).

Population genetic structure

Population genetic structure analyses identified ten distinct gene pools, among which two

included only a single population (FUE in southern Spain and PCI in northern Morocco;

Figure 1). Remarkably, these two single-population gene pools were not identified as

marginal populations by the geographical or demo-historical marginality indices whereas one

of them, PCI, was characterized by high values of the ecological index (standardized value of

5.013; Table S2), indicating persistence in a marginal climate. FUE and PCI had also low

levels of admixture with nearby gene pools. In contrast, the eight main gene pools (with the

exception of the highly isolated Tunisia-Pantelleria one, see below) were not genetically

isolated from each other, with populations often showing admixture with nearby gene pools.

This suggests either historical or recent gene flow across neighbouring gene pools along a

latitudinal cline in the western range of the species, and substantial shared ancestry among

French (including Corsica) and Italian populations in the eastern one.

Effects of population marginality on genetic indicators

Our models revealed a decline in overall genetic diversity (corrected $1\text{-}Q_{inter}$, Figure 2a) and

an increase in genetic differentiation (population-specific F_{ST}, Figure 2b; and Jost's D

statistic, Table S5) with population marginality based on the centroid, second-nearest core

and North-South indices (Tables 2a and S4). Indeed, models M1 and M5 predicted lower

genetic diversity and higher genetic differentiation in marginal populations based on the

centroid index (Tables 2a and S4), in particular in isolated populations from the southern

maritime pine range (as shown by significant coefficients for the second nearest-core and

North-South indices and their interaction; see also Figures 2 and S7). Model M2 revealed a

decrease in genetic diversity based on GEA outliers for southern range populations (Table 2a;

Figure S8a). Interestingly, model M3 revealed an increase in genetic diversity based on

general outliers in marginal northern range populations, especially for those with high values

of the centroid index (as shown by significant coefficients for North-South and centroid

indices and their interaction; Table 2a, Figure S9). However, this unexpected pattern may just

result from confounded effects due to high genetic diversity for general outlier loci in the two

northernmost gene pools (Atlantic France and North-East; compare Figure 1 with Figure S8b

and see M10 below). Models including the gene pool of origin as a random effect (M8 and

M12; Tables 2b) significantly improved the fitting by 10% for overall genetic diversity and

20% for population-specific F_{ST} (29% for Jost's D statistic; Table S5), but retained only the

centroid and North-South indices and their interaction as explanatory factors (with the

interaction having a similar interpretation as that for models without gene pool as random

effect; Figure S10). For genetic diversity based on GEA outliers, the model including the

gene pool as a random effect (M9) also improved the fitting (by over 30%) and revealed the

same relationship with the North-South index as M2 while no significant relationship was

found between genetic diversity based on general outliers and population marginality at the

gene pool level (M10). Interestingly, we found no relationship between inbreeding and the

indices of population marginality (see Figure S8c) for any model (M4, M11; Table 2a,b).

The model for recessive genetic load (M6) showed reduced genetic load for marginal

populations based on the centroid index, this association probably stemming mainly from the

high recessive genetic load found in some Iberian core populations (e.g., Carbonero el

Mayor, CAR, and Boniches, BON; see Figure S8d). However, this model had a relatively low

goodness-of-fit (R^2 = 12%) and we found no relationship between recessive genetic load and

any of the population marginality indices when we added the gene pool of origin as a random

effect (M13, Table 2b). Temperature seasonality (bio4) was the most important predictor

contributing to the genetic turnover in genomic offset estimates (see Figure S11b and the

GitHub detailing this analysis). Noticeably, this variable showed a steep slope between -2 and

0°C, which may indicate a rapid turnover in allele frequency in this range (see Figure S11a).

Despite relatively poor goodness-of-fit (Table 2a,b and Figures S16 and S17), both models

including the genomic offset, without (M7) and with (M14) the gene pool of origin as a

random effect, predicted an increased genomic offset for populations in marginal climatic

conditions (Table 2a,b and Figure 3b). Accordingly, we observed a trend for higher genomic

offset in the western gene pools along the Atlantic coast (average genomic offset of 0.047 ±

0.026 and 0.047 ± 0.016 for the Atlantic Iberian Peninsula and the French Atlantic gene

pools, respectively), with remarkably high values for Segurde (SEG), Alto de la Llama

(ALT), Armayán (ARM) and Olonne sur Mer (OLO) as well as for some southern

Mediterranean populations (e.g., Point Cires, PCI; Fuencaliente, FUE; Estepona, EST; and

Pantelleria, PAN; Figure 3a).

Discussion

In this study, we revealed a trend towards declining genetic diversity and increased genetic

differentiation in geographically and demo-historically marginal populations of maritime

pine; however, this genetic diversity trend would not necessarily apply to genes potentially

involved in local adaptation, as shown by genetic diversity analyses based on different sets of

outlier loci. We also found lower recessive genetic load in geographically marginal

populations and higher genomic offset in ecologically marginal ones. Models including the

gene pool of origin as a random effect were similar to those without (with the notable

exception of the models for genetic diversity for general outliers and recessive genetic load),

suggesting that the underlying processes operate at both the regional and range wide

geographical scales in this species. These results, taken together, provide support for the CPH

and suggest that climate change may endanger valuable and untapped genetic resources in

maritime pine.

Patterns of genetic diversity and differentiation support the CPH

The lower overall genetic diversity and higher genetic differentiation of geographical and

demo-historical marginal populations support the main predictions of the CPH. Our results

are consistent with those of Eckert et al. (2008) and Pironon et al. (2016), who found a

decline in genetic diversity and an increase in differentiation towards the limits of the species

ranges in 47% and 45% of studies in various taxa, respectively. As in our study, lower

genetic variation in peripheral compared to central populations was found for some other

conifers, i.e., in Sitka spruce (*Picea sitchensis*; Gapare et al., 2005), Swiss stone pine (*Pinus*

cembra; Gugerli et al., 2009) and common yew (*Taxus baccata*; Hilfiker et al., 2004), but not

in Norway spruce (*Picea abies*), for which only genetic differentiation conformed to the CPH

(Westergren et al., 2018). Interestingly, unlike the distribution of overall and climate-

associated (GEA) outlier genetic diversity with population marginality, genetic diversity

based on general outliers (identified by the R package *pcadapt* and BayeScan v2.1; see

Methods) increased with geographic marginality in terms of distance from the centroid,

especially for populations in the northern margins. There are several ways to interpret this

result. Firstly, geographically marginal populations may show more variation in outlier SNPs

linked to unknown biotic or abiotic factors (other than climate) than central populations.

Secondly, this pattern could result from a confounding effect linked to the distinct level of

genetic variation in outlier SNPs in the different gene pools, which could be corroborated by

the fact that no significant relationship with population marginality remains when models

included the gene pool of origin as a random factor. Finally, we could hypothesize that no

robust biological explanation can be drawn from this result, due to the uncertainty of these

outliers to be related or not to adaptation. Indeed, Lotterhos et al. (2014, 2015) and Hoban et

al. (2016) showed that most outliers detected with F_{ST}-outlier tests are likely to be false

positives when calculated on species with peculiar demographic histories such as range

expansion (e.g., Travis et al. 2007; Excoffier et al. 2009), as is the case in maritime pine.

In addition to geographical marginality, demo-historical marginality showed a

significant association with overall genetic diversity (M1 in Table 2a), as well as with genetic

diversity based on outlier loci (M2 and M3). This significant association is consistent with

the pattern observed in plants in the Mediterranean basin (Fady & Conord, 2010) but differs

from a more global pattern in which the neutral genetic diversity of plants (including pines)

does not decrease significantly with latitude (De Kort et al., 2021). Genetic differentiation,

which decreases with latitude in plants across their ranges (Gamba & Muchhala, 2020),

showed a significant association with demo-historical marginality in the same direction as

well (see M5 in Table 2a and M5bis in Table S5). However, despite the general trend of

decreasing genetic diversity with latitude, southern populations from the Moroccan and

Tunisia-Pantelleria gene pools and northern populations from the North-East and Corsican

gene pools exhibited the lowest overall and GEA outlier genetic diversity, and the highest

genetic differentiation in maritime pine (see Figures 2 and S8a). Maritime pine gene pools are

probably the result of population expansion from multiple glacial refugia, both in

Mediterranean and Atlantic regions of the species (Bucci et al., 2007; Jaramillo-Correa et al.,

2015), with Naydenov et al. (2014) also suggesting that, for this species, the high level of

overall genetic differentiation may have resulted from long historical isolation predating the

Last Glacial Maximum (~18,000 years ago). This was confirmed by estimates of genetic

divergence between North-African (Moroccan) and Iberian populations of maritime pine,

which dated back to 1.90 Ma (95% credibility interval: 1.41-2.76), probably due to the Strait

of Gibraltar's effect as a major biogeographic barrier to pollen and seed gene flow (Jaramillo-

Correa et al., 2010). Consequently, southernmost Moroccan populations are likely to have

been pre-glacial relict populations that survived in North-African refugia (Baradat &

Marpeau, 1988; Vendramin et al., 1998). After range expansion from glacial refugia, the

persistence of maritime pine in several isolated groups characterized by contrasting climatic

conditions may have resulted in populations that are locally adapted to the climate in some

way. However, our models revealed that the southernmost populations of this species

(Moroccan and Tunisia-Pantelleria gene pools) may be more at risk of not displaying enough

diversity, neutral or adaptive, than the central populations. This pattern may have been

exacerbated by a reduction of effective population size (N_E) due to human impact (Wahid et

al., 2004), as well as the incidence of recurrent forest fires resulting in population bottlenecks

and genetic drift in this region (Vendramin et al., 1998). The demographic history of

maritime pine in the north-eastern part of the continental range and the Corsica island has not

been clearly assessed (Naydenov et al., 2014), however, the presence of an endemic mitotype

in this region suggests long-term isolation of North-East and Corsican gene pools (Burban &

Petit, 2003). Nevertheless, both overall and GEA outlier genetic diversity for these

populations seem to be larger than for those from the southernmost gene pools (see Figures 2

and S8a).

Lower recessive genetic load in geographically marginal populations

Our models revealed a significant, albeit weak, reduction in recessive genetic load with

increased geographical population marginality. However, these models were only significant

when the gene pool effect was not accounted for, suggesting the existence of gene pools with

reduced/increased recessive genetic load and an important role of demographic history (see

below). This was a surprising finding, as we were expecting marginal populations to be

characterised by an accumulation of recessive genetic load due to the reduced effectiveness

of purifying selection in small and isolated populations with high demographic stochasticity

(Caballero et al., 2017; Sachdeva et al., 2022). The level of accumulation of deleterious

mutations and the extent to which it represents a risk to a given population depends primarily

on its effective size (N_E). Previous empirical studies of animal and plant populations that

underwent historical range expansions or declines have often shown an increase in genetic

load (e.g., Günther & Schmid, 2010, in *Arabidopsis thaliana*; González-Martínez et al., 2017,

in *Mercurialis annua*; Feng et al., 2019, in *Nipponia nippon*; or Peischl et al., 2013, in

humans). However, as in maritime pine, recent empirical studies based on genomic data

suggested that recessive genetic load can also be purged in long-term isolated and inbred

populations (see review in Dussex et al., 2023). As an example, Dussex et al. (2021) found

that current island populations of *Strigops habroptilus*, a New-Zealand flightless parrot, had

lower deleterious mutation load compared to mainland populations. A similar pattern was

found for the Alpine ibex (*Capra ibex*), which suffered severe population bottlenecks and

nearly became extinct (Grossen et al., 2020), and for the Channel Island fox (*Urocyon*

littoralis) that has remained at small population sizes with low diversity for many generations

(Robinson et al., 2018).

Present-day levels of inbreeding in maritime pine are low and not significantly higher

in marginal than core populations. Therefore, we hypothesise maritime pine marginal

populations to have effectively purged recessive genetic load during past inbreeding events,

operating at the regional scale. These events may have occurred during the range contractions

and/or expansions associated to Quaternary glacial and interglacial forest tree migrations

(Bucci et al., 2007; Naydenov et al. 2014; Jaramillo-Correa et al., 2015). Polyembryony,

which is ubiquitous in gymnosperms such as maritime pine (Willson & Burley, 1983), could

have also played a role in purging recessive genetic load, as it tends to dampen self-

fertilisation's deleterious effects by more effectively removing the mutational load through

selection between viable embryos (see Latta, 1995).

Higher genomic offset in ecologically marginal populations

Our models also revealed a higher genomic offset for ecologically marginal populations, but

not for geographically or demo-historically marginal ones suggesting that the gap between

the current and required genetic composition in future climates will be greater for populations

in marginal climatic conditions. This is consistent with the findings of Fréjaville et al. (2019),

who observed that adaptation lags in several forest trees, including maritime pine, are

consistently higher in climatically (cold/warm, dry/wet) marginal populations than in

populations growing under climatically optimal conditions. Although studies estimating

genomic offset in marginal populations are still scarce, two recent studies in widespread

Asian forest trees provided support to our findings, as they showed a relatively high genomic

offset in the northern and southern distribution margins of the Chinese thuja tree (*Platycladus*

orientalis; Jia et al., 2020) and the sawtooth oak (*Quercus acutissima*; Yuan et al., 2023); but

whether these range margins represented also ecologically extreme environments was not

assessed and more detailed studies are thus needed.

Estimation of the genomic offset is becoming a popular approach to assess population

vulnerability in the face of climate change. Genomic offset predictions have been validated

using data from common garden experiments and natural populations (e.g., Fitzpatrick et al.,

2021), including for maritime pine (Archambeau et al., 2024), but it is not free of pitfalls (see

Ahrens et al., 2023; Rellstab et al., 2021; Lind et al., 2024; Archambeau et al., 2024).

Moreover, genomic offsets can gauge (at some extent) for maladaptation to future climates

but not for the adaptive capacity of populations (Fitzpatrick et al., 2021; Archambeau et al.,

2024). Despite some of our model predictions based on general outlier loci, which showed a

reduction in potentially adaptive diversity towards the distribution margins, marginal

populations of forest trees can retain notable adaptive capacity, as shown in common garden

experiments for the handful of species with available data (e.g., for the Sierra Nevada Scots

pine, *Pinus sylvestris*, population in southern Spain, an isolated marginal population at the

species southern distribution limit; Alía et al., 2001; Castro et al., 2004). Thus, the potential

for both genetic adaptation and plastic responses needs to be integrated in models predicting

the responses of marginal populations to climate change. Furthermore, species' range limits

are not determined solely by climate and demographic processes. Loehle (1998) showed that

the range limits of many low-latitude tree species are set by competitive interactions with

other tree species. Other biotic factors, such as the positive interactions between species

(Stephan et al., 2021), are known to have a strong influence on the definition of tree range

boundaries. To better disentangle the relationship between genetic indicators and population

marginality within a species' range, future research should address the development of

predictive models that include species-specific indicators related to biotic interactions.

Conclusion

In maritime pine, a trend of decreasing in both overall and climate-associated outlier genetic

diversity, and increasing differentiation with geographic marginality supported the well-

established centre-periphery hypothesis, at both range-wide and regional (gene pool) spatial

scales. However, geographically marginal populations displayed also lower recessive genetic

load compared to core populations, which, together with expected novel adaptations in the

species range margins, highlight their importance in the context of future adaptation to

climate change. Higher genomic offset in ecologically marginal populations suggests higher

potential maladaptation of these populations to future climates and the urgency to develop

specific management actions to preserve populations in marginal climatic conditions. Overall,

our study shows the importance of combining quantitative marginality indices and diverse

genetic indicators, gauging for multiple evolutionary processes, to have a sound basis for

conservation decisions.

Author Contribution

SCG-M, MW, GGV, DG, AP and RA conceptualized the study. SP, FB, GGV, CA and SCG-

M produced, cleaned and formatted the genetic data. SCG-M, MW, DG and RA conceived

the paper methodology. AT, CG-P, JA and SCG-M analysed the data. AT, SCG-M and MW

drafted the article. All authors contributed to revision and editing of the final version, and

gave their final approval for publication.

Acknowledgements

We thank Carmen García-Barriga, Francisco Auñón, Eduardo Ballestero, Diana Barba, Maria

Regina Chambel, Fernando Del-Caño and Rodrigo Pulido-Sanz for their contribution to

genetic data production. We are also very grateful to Dr Bruno Fady and Dr Nicolas Picard

for their valuable recommendations and help at different stages of this study.

Funding statement

This study was funded by the European Union's Horizon 2020 research and innovation

program under grant agreement No 862221 (FORGENIUS). Views and opinions expressed

are however those of the author(s) only and do not necessarily reflect those of the European

Union. Neither the European Union nor the granting authority can be held responsible for

them. The study also received financial support from the French government in the

framework of the IdEx Bordeaux University 'Investments for the Future' program / GPR

Bordeaux Plant Sciences (France) and the MITECO contract for Conservation of Genetic

Resources (Spain).

Data archiving statement

Data are openly available in a public repository (Data INRAE) and was provided for the

anonymized review process as a file. Scripts for preparation of data, selection of climate

variables and computation of genomic offset statistics can be found in a GitHub public

repository, and were cloned and anonymised for the review process at

https://anonymous.4open.science/r/ReadyToGO_Pinpin-FA56/README.md.

Conflict of interest statement

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

Biosketch

This study is the result of a long-term collaboration among four public research institutions in Spain, Italy, Slovenia and France, which are working together on elucidating the distribution of genetic diversity and patterns of local adaptation in European forest trees, in particular those from highly diverse Mediterranean and Alpine forests. Our research interests are broad, including population and quantitative genetics, biogeography, conservation genetics, ecological modelling and genomics. The paper constitutes the first PhD manuscript of the first author, Adélaïde Theraroz, developed at the University of Bordeaux (France), and supervised by Marjana Westergren (Slovenian Forestry Institute) and Santiago C. González-Martínez (INRAE).

Short title

Genetic effects of population marginality

References

Abad-Viñas, R., Caudullo, G., Oliveira, S., & de Rigo, D. (2016). *Pinus pinaster* in Europe: distribution, habitat, usage and threats. In: San-Miguel-Ayanz, J., de Rigo, D., Caudullo, G., Houston Durrant, T., Mauri, A. (Eds.), *European Atlas of Forest Tree Species*. Publ. Off. EU, Luxembourg, pp. e012d59+.

Ahrens, C. W., Rymer, P. D., & Miller, A. D. (2023). Genetic offset and vulnerability modelling: misinterpretations of results and violations of evolutionary principles. *Authorea*. https://doi.org/10.22541/au.168727971.18670759/v1.

Alía, R., Martín, S., de Miguel, J., Galera, R., Agúndez, D., Gordo, J., Salvador, L., Catalán, G., & Gil, L. (1996). *Las regiones de procedencia de Pinus pinaster Aiton en España* (Organismo Autónomo Parques Nacionales-ETSI de Montes).

Alía, R., Moro-Serrano, J., & Notivol, E. (2001). Genetic variability of Scots pine (*Pinus*

sylvestris) provenances in Spain: growth traits and survival. *Silva Fennica*, *35*(1), 601.

https://doi.org/10.14214/sf.601.

Archambeau, J., Benito-Garzón, M., de-Miguel, M., Changenet, A., Bagnoli, F., Barraquand,

F., Marchi, M., Vendramin, G.G., Cavers, S., Perry, A., & González-Martínez, S.G.

(2024). Evaluating genomic offset predictions in a forest tree with high population

genetic structure. *bioRxiv*. https://doi.org/10.1101/2024.05.17.594631.

Baradat, P., Marpeau-Bezard, A. (1988). Le pin maritime *Pinus pinaster* Ait. Biologie et

génétique des terpènes pour la connaissance et l'amélioration de l'espèce. Thèse

doctorale d'État, Université Bordeaux I.

Birch, L. C. (1957). The role of weather in determining the distribution and abundance of

animals. *Cold Spring Harbor Symposia on Quantitative Biology*, *22*, 203-218.

https://doi.org/doi:10.1101/SQB.1957.022.01.021.

Brown, J. H. (1984). On the relationship between abundance and distribution of species. *The*

American Naturalist, *124*(2), 255-279. https://doi.org/10.1086/284267.

Bucci, G., González-Martínez, S. C., Le Provost, G., Plomion, C., Ribeiro, M. M., Sebastiani,

F., Alía, R., & Vendramin, G. G. (2007). Range-wide phylogeography and gene zones

in *Pinus pinaster* Ait. revealed by chloroplast microsatellite markers. *Molecular*

Ecology, *16*(10), 2137-2153. https://doi.org/10.1111/j.1365-294X.2007.03275.x.

Burban, C., & Petit, R. J. (2003). Phylogeography of maritime pine inferred with organelle

markers having contrasted inheritance. *Molecular Ecology*, *12*(6), 1487-1495.

https://doi.org/10.1046/j.1365-294X.2003.01817.x.

Caballero, A., Bravo, I., & Wang, J. (2017). Inbreeding load and purging: implications for the

short-term survival and the conservation management of small populations. *Heredity*,

118(2), 177-185. https://doi.org/10.1038/hdy.2016.80.

Cahn, L. (2023). *Genetic basis of local adaptation of maritime pine in Corsica* [Unpublished

master's thesis]. Université de Rennes 1.

Capblancq, T., Fitzpatrick, M. C., Bay, R. A., Expósito-Alonso, M., & Keller, S. R. (2020).

Genomic prediction of (mal)adaptation across current and future climatic landscapes.

Annual Review of Ecology, Evolution, and Systematics, *51*(1), 245-269.

https://doi.org/10.1146/annurev-ecolsys-020720-042553.

Castro, J., Zamora, R., Hódar, J. A., & Gómez, J. M. (2004). Seedling establishment of a

boreal tree species (*Pinus sylvestris*) at its southernmost distribution limit:

consequences of being in a marginal Mediterranean habitat. *Journal of Ecology*,

92(2), 266-277. https://doi.org/10.1111/j.0022-0477.2004.00870.x.

Caudullo, G., Welk, E., & San-Miguel-Ayanz, J. (2017). Chorological maps for the main

European woody species. *Data in Brief*, *12*, 662-666.

https://doi.org/10.1016/j.dib.2017.05.007.

Cingolani, P., Platts, A., Wang, L. L., Coon, M., Nguyen, T., Wang, L., Land, S. J., Lu, X., &

Ruden, D. M. (2012). A program for annotating and predicting the effects of single

nucleotide polymorphisms, SnpEff. *Fly, 6*(2), 80–92.

https://doi.org/10.4161/fly.19695.

De La Torre, A. R., Wang, T., Jaquish, B., & Aitken, S. N. (2014). Adaptation and exogenous

selection in a *Picea glauca* × *Picea engelmannii* hybrid zone: implications for forest

management under climate change. *New Phytologist*, *201*(2), 687-699.

https://doi.org/10.1111/nph.12540.

De-Lucas, A. I., González-Martínez, S. C., Vendramin, G. G., Hidalgo, E., & Heuertz, M.

(2009). Spatial genetic structure in continuous and fragmented populations of *Pinus*

pinaster Aiton. *Molecular Ecology*, *18*(22), 4564-4576.

https://doi.org/10.1111/j.1365-294X.2009.04372.x.

De Kort, H., Prunier, J. G., Ducatez, S., Honnay, O., Baguette, M., Stevens, V. M., & Blanchet, S. (2021). Life history, climate and biogeography interactively affect worldwide genetic diversity of plant and animal populations. *Nature Communications*, *12*(1). https://doi.org/10.1038/s41467-021-20958-2.

Dussex, N., Morales, H. E., Grossen, C., Dalén, L., & van Oosterhout, C. (2023). Purging and accumulation of genetic load in conservation. *Trends in Ecology & Evolution*, *38*(10), 961-969. https://doi.org/10.1016/j.tree.2023.05.008.

Dussex, N., van der Valk, T., Morales, H. E., Wheat, C. W., Díez-del-Molino, D., von Seth, J., Foster, Y., Kutschera, V. E., Guschanski, K., Rhie, A., Phillippy, A. M., Korlach, J., Howe, K., Chow, W., Pelan, S., Mendes Damas, J. D., Lewin, H. A., Hastie, A. R., Formenti, G., … Dalén, L. (2021). Population genomics of the critically endangered kākāpō. *Cell Genomics*, *1*(1), 100002. https://doi.org/10.1016/j.xgen.2021.100002.

Eckert, C. G., Samis, K. E., & Lougheed, S. C. (2008). Genetic variation across species' geographical ranges: the central–marginal hypothesis and beyond. *Molecular Ecology*, *17*(5), 1170-1188. https://doi.org/10.1111/j.1365-294X.2007.03659.x.

Ellis, N., Smith, S. J., & Pitcher, C. R. (2012). Gradient forests: calculating importance gradients on physical predictors. *Ecology*, *93*(1), 156-168. https://doi.org/10.1890/11-0252.1.

Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology*, *14*(8), 2611-2620. https://doi.org/10.1111/j.1365-294X.2005.02553.x.

Excoffier, L., Foll, M., & Petit, R. J. (2009). Genetic consequences of range expansions. *Annual Review of Ecology, Evolution, and Systematics*, *40*(1), 481–501. https://doi.org/10.1146/annurev.ecolsys.39.110707.173414.

Fady, B., & Conord, C. (2010). Macroecological patterns of species and genetic diversity in

vascular plants of the Mediterranean basin. *Diversity and Distributions*, *16*(1), 53–64.

https://doi.org/10.1111/j.1472-4642.2009.00621.x.

Feng, S., Fang, Q., Barnett, R., Li, C., Han, S., Kuhlwilm, M., Zhou, L., Pan, H., Deng, Y.,

Chen, G., Gamauf, A., Woog, F., Prys-Jones, R., Marques-Bonet, T., Gilbert, M. T.

P., & Zhang, G. (2019). The genomic footprints of the fall and recovery of the crested

Ibis. *Current Biology*, *29*(2), 340-349.e7. https://doi.org/10.1016/j.cub.2018.12.008.

Fitzpatrick, M. C., Chhatre, V. E., Soolanayakanahally, R. Y., & Keller, S. R. (2021).

Experimental support for genomic prediction of climate maladaptation using the

machine learning approach Gradient Forests. *Molecular Ecology Resources*, *21*(8),

2749-2765. https://doi.org/10.1111/1755-0998.13374.

Fitzpatrick, M. C., & Keller, S. R. (2015). Ecological genomics meets community-level

modelling of biodiversity: mapping the genomic landscape of current and future

environmental adaptation. *Ecology Letters*, *18*(1), 1-16.

https://doi.org/10.1111/ele.12376.

Fkiri, S., Guibal, F., Elkhorchani, A., Khouja, M. L., Khaldi, A., & Nasr, Z. (2019).

Relationship between climate and growth of two North African varieties of *Pinus*

pinaster Arn. *African Journal of Ecology*, *57*(3), 327-334.

https://doi.org/10.1111/aje.12610.

Foll, M., & Gaggiotti, O. (2008). A genome-scan method to identify selected loci appropriate

for both dominant and codominant markers: a Bayesian perspective. *Genetics*, *180*(2),

977-993. https://doi.org/10.1534/genetics.108.092221.

Fréjaville, T., Vizcaíno-Palomar, N., Fady, B., Kremer, A., & Benito Garzón, M. (2019).

Range margin populations show high climate adaptation lags in European trees.

Global Change Biology, *26*(2), 484-495. https://doi.org/10.1111/gcb.14881.

Frichot, E., Schoville, S. D., Bouchard, G., & François, O. (2013). Testing for associations

between loci and environmental gradients using Latent Factor Mixed Models.

Molecular Biology and Evolution, *30*(7), 1687-1699.

https://doi.org/10.1093/molbev/mst063.

Gamba, D., & Muchhala, N. (2020). Global patterns of population genetic differentiation in

seed plants. *Molecular Ecology, 29*(18), 3413–3428.

https://doi.org/10.1111/mec.15575.

Gapare, W. J., Aitken, S. N., & Ritland, C. E. (2005). Genetic diversity of core and peripheral

Sitka spruce (*Picea sitchensis* (Bong.) Carr) populations: implications for

conservation of widespread species. *Biological Conservation*, *123*(1), 113-123.

https://doi.org/10.1016/j.biocon.2004.11.002.

Gaston, K. J. (2003). *The structure and dynamics of geographic ranges.* Oxford University

Press, Oxford.

Gaston, K. J. (2009). Geographic range limits: achieving synthesis. *Proceedings of the Royal*

Society B: Biological Sciences, *276*(1661), 1395-1406.

https://doi.org/10.1098/rspb.2008.1480.

Gautier, M. (2015). Genome-wide scan for adaptive divergence and association with

population-specific covariates. *Genetics*, *201*(4), 1555-1579.

https://doi.org/10.1534/genetics.115.181453.

González-Martínez, S.C., Gómez, A., Carrión, J.S., Agúndez, D., Alía, R., Gil, L. (2007).

Spatial genetic structure of an explicit glacial refugium of maritime pine (*Pinus*

pinaster Aiton) in southeastern Spain. In: Weiss, S., Ferrand, N. (eds)

Phylogeography of Southern European Refugia. Springer, Dordrecht.

González-Martínez, S. C., Ridout, K., & Pannell, J. R. (2017). Range expansion compromises

adaptive evolution in an outcrossing plant. *Current Biology*, *27*(16), 2544-2551.e4.

https://doi.org/10.1016/j.cub.2017.07.007.

Gougherty, A. V., Keller, S. R., & Fitzpatrick, M. C. (2021). Maladaptation, migration and

extirpation fuel climate change risk in a forest tree species. *Nature Climate Change*,

11(2), 166-171. https://doi.org/10.1038/s41558-020-00968-6.

Grossen, C., Guillaume, F., Keller, L. F., & Croll, D. (2020). Purging of highly deleterious

mutations through severe bottlenecks in Alpine ibex. *Nature Communications*, *11*(1),

1001. https://doi.org/10.1038/s41467-020-14803-1.

Gugerli, F., Rüegg, M., & Vendramin, G. G. (2009). Gradual decline in genetic diversity in

Swiss stone pine populations (*Pinus cembra*) across Switzerland suggests postglacial

re-colonization into the Alps from a common eastern glacial refugium. *Botanica*

Helvetica, *119*(1), 13-22. https://doi.org/10.1007/s00035-009-0052-6.

Günther, T., & Schmid, K. J. (2010). Deleterious amino acid polymorphisms in *Arabidopsis*

thaliana and rice. *Theoretical and Applied Genetics*, *121*(1), 157-168.

https://doi.org/10.1007/s00122-010-1299-4.

Hampe, A., & Petit, R. J. (2005). Conserving biodiversity under climate change: the rear edge

matters. *Ecology Letters*, *8*(5), 461-467. https://doi.org/10.1111/j.1461-

0248.2005.00739.x.

Hedrick, P. W., & García-Dorado, A. (2016). Understanding inbreeding depression, purging,

and genetic rescue. *Trends in Ecology & Evolution*, *31*(12), 940-952.

https://doi.org/10.1016/j.tree.2016.09.005.

Hengeveld, R., & Haeck, J. (1982). The distribution of abundance. I. Measurements. *Journal*

of Biogeography, *9*(4), 303. https://doi.org/10.2307/2844717.

Hilfiker, K., Gugerli, F., Schütz, J.-P., Rotach, P., & Holderegger, R. (2004). Low RAPD

variation and female-biased sex ratio indicate genetic drift in small populations of the

dioecious conifer *Taxus baccata* in Switzerland. *Conservation Genetics*, *5*(3),

357-365. https://doi.org/10.1023/B:COGE.0000031144.95293.1b.

Hoban, S., Kelley, J. L., Lotterhos, K. E., Antolin, M. F., Bradburd, G., Lowry, D. B., Poss,

M. L., Reed, L. K., Storfer, A., & Whitlock, M. C. (2016). Finding the genomic basis

of local adaptation: pitfalls, practical solutions, and future directions. *The American*

Naturalist, 188(4), 379–397. https://doi.org/10.1086/688018.

IPCC (2021). *Climate Change 2021: The Physical Science Basis: Contribution of Working*

Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate

Change. Cambridge University Press, Cambridge, United Kingdom and New York,

NY, USA.

Jaramillo-Correa, J. P., Grivet, D., Terrab, A., Kurt, Y., De-Lucas, A. I., Wahid, N.,

Vendramin, G. G., & González-Martínez, S. C. (2010). The Strait of Gibraltar as a

major biogeographic barrier in Mediterranean conifers: a comparative

phylogeographic survey. *Molecular Ecology*, *19*(24), 5452-5468.

https://doi.org/10.1111/j.1365-294x.2010.04912.x.

Jaramillo-Correa, J.-P., Rodríguez-Quilón, I., Grivet, D., Lepoittevin, C., Sebastiani, F.,

Heuertz, M., Garnier-Géré, P. H., Alía, R., Plomion, C., Vendramin, G. G., &

González-Martínez, S. C. (2015). Molecular proxies for climate maladaptation in a

long-lived tree (*Pinus pinaster* Aiton, Pinaceae). *Genetics*, *199*(3), 793-807.

https://doi.org/10.1534/genetics.114.173252.

Jia, K., Zhao, W., Maier, P. A., Hu, X., Jin, Y., Zhou, S., Jiao, S., El-Kassaby, Y. A., Wang,

T., Wang, X., & Mao, J. (2020). Landscape genomics predicts climate change-related

genetic offset for the widespread *Platycladus orientalis* (Cupressaceae). *Evolutionary Applications*, *13*(4), 665-676. https://doi.org/10.1111/eva.12891.

Johannesson, K., & André, C. (2006). Life on the margin: genetic isolation and diversity loss in a peripheral marine ecosystem, the Baltic Sea. *Molecular Ecology*, *15*(8), 2013-2029. https://doi.org/10.1111/j.1365-294X.2006.02919.x.

Jost, L. (2008). G_{ST} and its relatives do not measure differentiation. *Molecular Ecology*, *17*(18), 4015-4026. https://doi.org/10.1111/j.1365-294X.2008.03887.x.

Kimura, M., Maruyama, T., & Crow, J. F. (1963). The mutation load in small populations. *Genetics*, *48*(10), 1303–1312. https://doi.org/10.1093/genetics/48.10.1303.

Kimura, M., & Ohta, T. (1969). The average number of generations until fixation of a mutant gene in a finite population. *Genetics*, *61*(3), 763-771. https://doi.org/10.1093/genetics/61.3.763.

Kolzenburg, R. (2022). The direct influence of climate change on marginal populations: a review. *Aquatic Sciences*, *84*(2), 24. https://doi.org/10.1007/s00027-022-00856-5.

Lande, R. (1988). Genetics and demography in biological conservation. *Science*, *241*(4872), 1455-1460. https://doi.org/10.1126/science.3420403.

Láruson, Á. J., Fitzpatrick, M. C., Keller, S. R., Haller, B. C., & Lotterhos, K. E. (2022). Seeing the forest for the trees: Assessing genetic offset predictions from gradient forest. *Evolutionary Applications, 15*, 403-416. https://doi.org/10.1111/eva.13354.

Latta, R. G. (1995). The effects of embryo competition with mixed mating on the genetic load in plants. *Heredity*, *75*(6), 637-643. https://doi.org/10.1038/hdy.1995.183.

Lind, B. M., Candido-Ribeiro, R., Singh, P., Lu, M., Vidaković, D. O., Booker, T. R., Whitlock, M. C., Yeaman, S., Isabel, N., & Aitken, S. N. (2024). How useful is genomic data for predicting maladaptation to future climate? *Global Change Biology, 30*(4). https://doi.org/10.1111/gcb.17227.

Lira-Noriega, A., & Manthey, J. D. (2014). Relationship of genetic diversity and niche

centrality: a survey and analysis. *Evolution*, *68*(4), 1082-1093.

https://doi.org/10.1111/evo.12343.

Loehle, C. (1998). Height growth rate trade-offs determine northern and southern range limits

for trees. *Journal of Biogeography, 25*(4), 735–742.

http://www.jstor.org/stable/2846146.

Lotterhos, K. E., & Whitlock, M. C. (2014). Evaluation of demographic history and neutral

parameterization on the performance of F_{ST} outlier tests. *Molecular Ecology, 23*(9),

2178–2192. https://doi.org/10.1111/mec.12725.

Lotterhos, K. E., & Whitlock, M. C. (2015). The relative power of genome scans to detect

local adaptation depends on sampling design and statistical method. *Molecular*

Ecology, 24(5), 1031–1046. https://doi.org/10.1111/mec.13100.

Lotterhos, K. E. and Whitlock, M. C. (2014). Evaluation of demographic history and neutral

parameterization on the performance of FST outlier tests. *Molecular Ecology, 23*,

2178– 2192. https://doi.org/10.1111/mec.12725.

Lotterhos, K. E. (2024). Interpretation issues with "genomic vulnerability" arise from

conceptual issues in local adaptation and maladaptation. *Evolution Letters, 8*(3), 331-

339. https://doi.org/10.1093/evlett/qsrae004.

Lynch, M., Conery, J., & Bürger, R. (1995). Mutation accumulation and the extinction of

small populations. *The American Naturalist*, *146*(4), 489-518.

https://doi.org/10.1086/285812.

Marchi, M., Castellanos-Acuña, D., Hamann, A., Wang, T., Ray, D., & Menzel, A. (2020).

ClimateEU, scale-free climate normals, historical time series, and future projections

for Europe. *Scientific Data*, *7*(1), 428. https://doi.org/10.1038/s41597-020-00763-0.

Marques, H., Pinto, G., Pinto, P., Teixeira, C. (2012). Regiões de Proveniência. Portugal.

ICNF, 47-51.

Milesi, P., Kastally, C., Dauphin, B., Cervantes, S., Bagnoli, F., Budde, K. B., Cavers, S.,

Fady, B., Faivre-Rampant, P., González-Martínez, S. C., Grivet, D., Gugerli, F.,

Jorge, V., Kupin, I. L., Ojeda, D. I., Olsson, S., Opgenoorth, L., Pinosio, S., Plomion,

C., ... Pyhäjärvi, T. (2023). Synchronous effective population size changes and

genetic stability of forest trees through glacial cycles. *bioRxiv*.

https://doi.org/10.1101/2023.01.05.522822.

Naydenov, K. D., Alexandrov, A., Matevski, V., Vasilevski, K., Naydenov, M. K., Gyuleva,

V., Carcaillet, C., Wahid, N. and Kamary, S. (2014). Range-wide genetic structure of

maritime pine predates the last glacial maximum: evidence from nuclear DNA.

Hereditas, *151*(1), 1–13. https://doi.org/10.1111/j.1601-5223.2013.00027.x.

Nicholson, A. J. (1958). Dynamics of insect populations. *Annual Review of Entomology*, *3*(1),

107-136. https://doi.org/10.1146/annurev.en.03.010158.000543.

Peischl, S., Dupanloup, I., Kirkpatrick, M., & Excoffier, L. (2013). On the accumulation of

deleterious mutations during range expansions. *Molecular Ecology*, *22*(24),

5972-5982. https://doi.org/10.1111/mec.12524.

Picard, N., Marchi, M., Serra-Varela, M. J., Westergren, M., Cavers, S., Notivol, E., Piotti,

A., Alizoti, P., Bozzano, M., González-Martínez, S. C., Grivet, D., Aravanopoulos, F.

A., Vendramin, G. G., Ducci, F., Fady, B., & Alía, R. (2022). Marginality indices for

biodiversity conservation in forest trees. *Ecological Indicators*, *143*, 109367.

https://doi.org/10.1016/j.ecolind.2022.109367.

Pironon, S., Villellas, J., Morris, W. F., Doak, D. F., & García, M. B. V. (2015). Do

geographic, climatic or historical ranges differentiate the performance of central

versus peripheral populations? *Global Ecology and Biogeography*, *24*(6), 611-620.

https://doi.org/10.1111/geb.12263.

Pironon, S., Papuga, G., Villellas, J., Angert, A. L., García, M. B. V., & Thompson, J. D.

(2016). Geographic variation in genetic and demographic performance: new insights

from an old biogeographical paradigm. *Biological Reviews*, *92*(4), 1877-1909.

https://doi.org/10.1111/brv.12313.

Plomion, C., Bartholomé, J., Lesur, I., Boury, C., Rodríguez-Quilón, I., Lagraulet, H., &

González-Martínez, S. C. (2016). High-density SNP assay development for genetic

analysis in maritime pine (*Pinus pinaster*). *Molecular Ecology Resources, 16*(2), 574-

587. https://doi.org/10.1111/1755-0998.12464.

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using

multilocus genotype data. *Genetics*, *155*(2), 945-959.

https://doi.org/10.1093/genetics/155.2.945.

R Core Team. (2022). *R: A Language and Environment for Statistical Computing*. R

Foundation for Statistical Computing. https://www.R-project.org/.

Rellstab, C., Dauphin, B., & Expósito-Alonso, M. (2021). Prospects and limitations of

genomic offset in conservation management. *Evolutionary Applications*, *14*(5),

1202-1212. https://doi.org/10.1111/eva.13205.

Richards, O. W. (1961). The theoretical and practical study of natural insect populations.

Annual Review of Entomology, *6*(1), 147-162.

https://doi.org/10.1146/annurev.en.06.010161.001051.

Robinson, J. A., Brown, C., Kim, B. Y., Lohmueller, K. E., & Wayne, R. K. (2018). Purging

of strongly deleterious mutations explains long-term persistence and absence of

inbreeding depression in island foxes. *Current Biology*, *28*(21), 3487-3494.e4.

https://doi.org/10.1016/j.cub.2018.08.066.

Rousset, F. (2008). Genepop'007: a complete re-implementation of the genepop software for

Windows and Linux. *Molecular Ecology Resources*, *8*(1), 103-106.

https://doi.org/10.1111/j.1471-8286.2007.01931.x.

Sachdeva, H., Olusanya, O., & Barton, N. (2022). Genetic load and extinction in peripheral

populations: the roles of migration, drift and demographic stochasticity. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *377*(1846).

https://doi.org/10.1098/rstb.2021.0010.

Salvador, L., Alía, R., Agúndez, D., & Gil, L. (2000). Genetic variation and migration

pathways of maritime pine (*Pinus pinaster* Ait) in the Iberian Peninsula. *Theoretical*

and Applied Genetics, 100(1), 89–95. https://doi.org/10.1007/s001220050013.

Santos, A. S., & Gaiotto, F. A. (2020). Knowledge status and sampling strategies to

maximize cost-benefit ratio of studies in landscape genomics of wild plants. *Scientific*

Reports, *10*(1), 3706. https://doi.org/10.1038/s41598-020-60788-8.

Sexton, J. P., McIntyre, P. J., Angert, A. L., & Rice, K. J. (2009). Evolution and ecology of

species range limits. *Annual Review of Ecology, Evolution, and Systematics*, *40*(1),

415-436. https://doi.org/10.1146/annurev.ecolsys.110308.120317.

Sexton, J. P., Hangartner, S. B., & Hoffmann, A. A. (2013). Genetic isolation by environment

or distance: which pattern of gene flow is most common? *Evolution*, *68*(1), 1-15.

https://doi.org/10.1111/evo.12258.

Sgrò, C. M., Lowe, A. J., & Hoffmann, A. A. (2010). Building evolutionary resilience for

conserving biodiversity under climate change. *Evolutionary Applications*, *4(2)*, 326–

337. https://doi.org/10.1111/j.1752-4571.2010.00157.x.

Soille, P., & Vogt, P. (2009). Morphological segmentation of binary patterns. *Pattern*

Recognition Letters, *30*(4), 456-459. https://doi.org/10.1016/j.patrec.2008.10.015.

Soulé, M.E. (1973). The epistasis cycle: a theory of marginal populations. *Annual Review of*

Ecology and Systematics, *4*(1), 165-187.

https://doi.org/10.1146/annurev.es.04.110173.001121.

Stephan, P., Mora, B. B., & Alexander, J. M. (2021). Positive species interactions shape

species' range limits. *Oikos*, *130*(10), 1611–1625. https://doi.org/10.1111/oik.08146.

Travis, J. M. J., Munkemuller, T., Burton, O. J., Best, A., Dytham, C., & Johst, K. (2007).

Deleterious mutations can surf to high densities on the wave front of an expanding

population. *Molecular Biology and Evolution*, *24*(10), 2334–2343.

https://doi.org/10.1093/molbev/msm167.

van den Wollenberg, A. L. (1977). Redundancy analysis an alternative for canonical

correlation analysis. *Psychometrika*, *42*(2), 207-219.

https://doi.org/10.1007/BF02294050.

Vendramin, G., Anzidei, M., Madaghiele, A., Bucci, G. (1998). Distribution of genetic

diversity in *Pinus pinaster* Ait. as revealed by chloroplast microsatellites. *Theoretical*

and Applied Genetics, 97, 456–463. https://doi.org/10.1007/s001220050917.

Wahid, N., González-Martínez, S. C., Hadrami, I. E., & Boulli, A. (2004). Genetic Structure

and variability of natural populations of maritime pine (*Pinus pinaster* Aiton) in

Morocco. *Silvae Genetica*, *53*(1-6), 93-99. https://doi.org/10.1515/sg-2004-0017.

Wahid, N., González-Martínez, S. C., El Hadrami, I., & Boulli, A. (2006). Variation of

morphological traits in natural populations of maritime pine (*Pinus pinaster* Ait.) in

Morocco. *Annals of Forest Science*, *63*(1), 83-92.

https://doi.org/10.1051/forest:20050100.

Weir, B. S., & Cockerham, C. C. (1984). Estimating F-statistics for the analysis of population

structure. *Evolution*, *38*(6), 1358. https://doi.org/10.2307/2408641.

Westergren, M., Božič, G., Kraigher, H. (2018). Genetic diversity of core *vs.* peripheral

Norway spruce native populations at a local scale in Slovenia. *iForest -*

Biogeosciences and Forestry, *11*(1), 104-110. https://doi.org/10.3832/ifor2444-011.

Whittaker, J. B. (1971). Population changes in *Neophilaenus lineatus* (L.) (Homoptera :

Cercopidae) in different parts of its range. *The Journal of Animal Ecology*, *40*(2), 425.

https://doi.org/10.2307/3253.

Willson, M. F., & Burley, N. (1983). *Mate Choice in Plants (MPB-19), Volume 19: Tactics,*

Mechanisms, and Consequences. (MPB-19). Princeton University Press.

https://doi.org/10.2307/j.ctvx5wbss.

Young, A.G., Boyle, T., & Brown, T.L. (1996). The population genetic consequences of

habitat fragmentation for plants. *Trends in Ecology & Evolution*, *11*(10), 413-418.

https://doi.org/10.1016/0169-5347(96)10045-8.

Yuan, S., Shi, Y., Zhou, B., Liang, Y., Chen, X., An, Q., Fan, Y., Shen, Z., Ingvarsson, P. K.,

& Wang, B. (2023). Genomic vulnerability to climate change in *Quercus acutissima*,

a dominant tree species in East Asian deciduous forests. *Molecular Ecology*, *32*(7),

1639-1655. https://doi.org/10.1111/mec.16843.

 Figure legends

 Figure 1. Genetic pools in *P. pinaster*. Pie charts depict membership proportions of each

 genetic cluster (K=10) for each studied population, calculated by STRUCTURE v2.3.4. The

 natural distribution of the species is shadowed in light green (see details in Figure S1).

 Population codes are only provided for those populations specifically mentioned in the main

 text (see full population information, including population names and geographical

 coordinates, in Table S1).

 Figure 2. Geographical distribution of a) overall genetic diversity (1-Q_{inter}) and b) genetic

 differentiation (population-specific F_{ST}), and two marginality indices involved in significant

 correlations with these genetic indicators, c) centroid index and d) second nearest-core index,

 for *P. pinaster* populations. See Table S1 for population information, including population

 names and geographical coordinates.

 Figure 3. a) Geographical distribution of genomic offset for *P. pinaster* populations and b)

 correlation between the genomic offset and the ecological index based on climate distances

 (standardized values), as shown by linear regression in *ggplot2* R package. Population codes

 are only provided for populations specifically mentioned in the main text (see full population

 information, including population names and geographical coordinates, in Table S1).

962 **Table 1**. Geographical (*geo*) and demo-historical (*histo*) marginality indices computed for maritime pine populations; '(-)' and '(+)' indicate that

963 higher values of the index correspond to lower and higher population marginality, respectively; '(-)(+)' indicates that both higher or lower values

964 of the index correspond to higher population marginality. Adapted after Picard et al. (2022).

Marginality index	Category	Link with marginality	Description
Area	*geo*	(-)	Size in ha of the core (i.e., continuous patches of the species' presence) that is the nearest to the location of the studied population.
Gravity	*geo*	(-)	Spatially continuous alternative to the area index, computed as the weighted mean of the core areas with weights inversely proportional to the distance squared from the location to the cores.
Centroid	*geo*	(+)	Eccentricity computed as the cost distance from the centroid of the species distribution to any location, taking into account the terrestrial and marine connectivity linked to pollen circulation capacity. Cost distances are based on a conductance matrix (the inverse of a resistance matrix) reflecting the permeability of land and water cells to gene flow. This index is expected to capture the level of long-distance gene flow.
Edge	*geo*	(-)	Distance to the nearest border of the species distribution.
Isolation	*geo*	(+)	Isolation with respect to the core species distribution, computed as the distance from the focal population to the nearest core greater than 100 ha and further than 50 km away.
Second nearest-core	*geo*	(+)	Spatially continuous alternative to the isolation index, computed as the distance from a location to the second nearest-core greater than 100 ha.

North-South	*histo*	(-) (+)	Proximity to the species rear-edge (southernmost limit of the species distribution) or leading-edge (northernmost limit of the species distribution) along a latitudinal gradient.
East-West	*histo*	(-) (+)	Proximity to the species leading-edge or rear-edge along a longitudinal gradient.

968   **Table 2**. Models evaluating the effect of population marginality on the estimated genetic indicators; a) Models without gene pool effect (M1-

969   M7); b) Models including gene pool effect as a random factor (M8-M14). Significant fixed effect(s) are listed with their associated point

970   estimates and 95% confidence intervals [in brackets]. R^2: variance explained by fixed factors (M1-M7) or both fixed (marginal R^2) and random

971   (conditional R^2) factors (M8-M14); ns: not significant.

973   a) Models without gene pool effect.

Model [R^2]	Genetic indicator	Fixed-effect #1	Estimate [95% CIs]	Fixed-effect #2	Estimate [95% CIs]	Fixed-effect #3	Estimate [95% CIs]	Interaction	Estimate [95% CIs]
M1 [0.59]	*Overall genetic diversity*	Centroid	-1.01 [-1.34, -0.67]	Second nearest-core	-1.47 [-2.43, -0.51]	North-South	1.61 [0.66, 2.55]	Second nearest-core × North-South	1.44 [0.50, 2.38]
M2 [0.26]	*Genetic diversity for GEA outliers*	North-South	0.62 [0.40, 0.85]	ns		ns		ns	
M3 [0.79]	*Genetic diversity for general outliers*	Centroid	1.26 [1.07, 1.45]	North-South	1.60 [1.32, 1.88]	ns		Centroid × North-South	1.16 [0.93, 1.40]
M4 [<0.01]	*Inbreeding*	ns		ns		ns		ns	
M5 [0.66]	*Population-specific F_{ST}*	Centroid	1.05 [0.74, 1.35]	Second nearest-core	1.43 [0.55, 2.31]	North-South	-1.90 [-2.76, -1.04]	Second nearest-core × North-South	-1.66 [-2.51, -0.80]
M6 [0.12]	*Recessive genetic load*	Centroid	-0.69 [-1.08, -0.30]	ns		ns		ns	

Model [R^2]	Genetic indicator	Fixed-effect #1	Estimate [95% CIs]	Fixed-effect #2	Estimate [95% CIs]	Interaction	Estimate [95% CIs]
M7 [0.27]	*Genetic offset*	Ecological index	0.48 [0.28, 0.68]	ns		ns		ns

975   b) Models with gene pool effect as random factor.

Model [R^2]	Genetic indicator	Fixed-effect #1	Estimate [95% CIs]	Fixed-effect #2	Estimate [95% CIs]	Interaction	Estimate [95% CIs]
M8 [0.41, 0.69]	*Overall genetic diversity*	Centroid	-1.04 [-1.58, -0.50]	North-South	1.04 [0.34, 1.75]	Centroid × North-South	0.75 [0.25, 1.24]
M9 [0.13, 0.5]	*Genetic diversity for GEA outliers*	North-South	0.44 [0.10, 0.77]	ns		ns	
M10 [<0.01]	*Genetic diversity for general outliers*	ns		ns		ns	
M11 [<0.01]	*Inbreeding*	ns		ns		ns	
M12 [0.31, 0.86]	*Population-specific F_{ST}*	Centroid	0.74 [0.21, 1.28]	North-South	-1.40 [-2.15, -0.65]	Centroid × North-South	-0.96 [-1.45, -0.46]
M13 [<0.01]	*Recessive genetic load*	ns		ns		ns	
M14 [0.22, 0.40]	*Genetic offset*	Ecological index	0.44 [0.24, 0.65]	ns		ns	

Figure 1

980   **Figure 2**

983 **Figure 3**