Castration Model Illuminates Sex Differences in Healthy Aging: Insights from Metabolome and Transcriptome Analyses

Jianjun Jiang[1,2#], Na Ge[2], Yuzhi Wang[2], Juntao Qi[3], Guibiao Wen[3], Xiufen Gu[2], Xuewen Yu[4], Muming Shao[4], Yueming Luo[4], Kangshuyun Gu[2], Feng Lin[3], Shudong Yang[2#], Wei Wei [1#],

[1] Institute for Systems Biology, Seattle, USA;
[2] Department of Nephrology, [3] Department of Urology, [4] Department of Pathology, [5] Department of Geriatrics, Shenzhen Traditional Chinese Medicine Hospital, The Fourth Clinical Medical College of Guangzhou University of Chinese Medicine, Shenzhen, China;

Authors' e-mails:
Jianjun Jiang : jjiang@systemsbiology.org; Na Ge: gena.sz@gzucm.edu.cn;
Yuzhi Wang : wangyuzhi997@gmail.com; Juntao Qi: juntaoqi@outlook.com;
Guibiao Wen: guibiaowen@163.com; Xiufen Gu: nodo1007@163.com;
Xuewen Yu:yuxuewen365@163.com; Muming Shao: smm1320@gzucm.edu.cn;
Yueming Luo: yuemingluo@163.com; Kangshuyun Gu: gukangshuyun@foxmail.com;
Feng Lin: lf0237@gzucm.edu.cn;Shudong Yang: ysd0870@gzucm.edu.cn;
Wei Wei:wwei@systemsbiology.org;

[*] These authors contribute equally to this work
[#] Corresponding authors
Jianjun Jiang, Ph.D., E-mail: jjiang@systemsbiology.org
Shudong Yang: ysd0870@gzucm.edu.cn
Wei Wei, Ph.D., E-mail: wwei@systemsbiology.org;

Abstract

Females typically outlive males, a disparity mitigated by castration, yet the molecular underpinnings remain elusive. Our study leverages untargeted metabolomics and RNA sequencing to uncover the pivotal compounds and genes influencing healthy aging post-castration, examining serum, kidney, and liver biospecimens from 12-week and 18-month old castrated male mice and their unaltered counterparts. Behavioral tests and LC-MS/MS metabolomics reveal that castrated males exhibit altered steroid hormones, superior cognitive performance, and higher levels of anti-oxidative compounds like taurine, despite identical diets. Integrated metabolome-transcriptome analysis confirms reduced lipid peroxidation and oxidative stress in female and castrated male mice, suggesting a protective mechanism against aging. Histological examinations post-cisplatin treatment highlight the model's applicability in studying drug toxicity and reveal varying susceptibility in organ-specific toxicities, underlining the crucial role of sex hormones in physiological defenses. In essence, our castration model unveils a feminized metabolic and transcriptomic intermediary, serving as a robust tool for studying gender-specific aspects of healthy aging and exploring sex hormone-induced differences in diverse biomedical domains.

INTRODUCTION

Sex-related differences in human are easy to find, where women consistently live longer than men [1]. In the longevous population - centenarians, women always outnumber men [2]. While past research have attributed longer female lifespans to healthier lifestyles, such as lower alcohol and tobacco consumption [2-4], a compiled demographic study of 134 mammal populations across 101 species revealed that females have a median lifespan that is on average 18.6% longer than their male counterparts. This suggests the intrinsic biological mechanisms underlie the sex differences in lifespan [1]. Sex hormones were considered as key factors in explaining the sex difference patterns of longevity [5]. Estrogen was proposed to play protective roles against many diseases, while testosterone may increase the risk of developing diseases and loss of testosterone perhaps promote healthy aging and longevity [6, 7]. Nevertheless, more evidence from clinical trials is needed to elucidate it.

Aging stands as a paramount risk factor for a myriad of chronic diseases, with differential impacts on males and females. The absence of a stable and broadly applicable model continues to pose a significant challenge in the quest to unravel the intricacies of sex differences in healthy aging [8]. In this study, we present castration as an effective model for understanding the sex-based differences in healthy aging. This is supported by a growing body of evidence demonstrating castration can extend the lifespan across multiple mammal species and decelerate intrinsic aging processes by slowing the DNA methylation clock [9]. For example, castration at prepubertal period can significantly prolong male mice lifespan, which eliminates the sex differences in lifespan and growth trajectories [10]. Castration in human, with statistical evidence shown that Korean eunuchs lived 14.4–19.1 years longer than normal men of similar socio-economic status [11]. However, the extent to which castration promotes healthy aging, and the underlying mechanisms, remain largely unexplored. Considered that genome sequences retain the same pre- and post-castration, we hypothesized that alterations in hormones spectrum and associated changes in metabolome, transcriptome and epigenome by castration possibly act as healthy aging regulators and affects lifespan extension. In this study, we utilized untargeted liquid chromatography with tandem mass spectrometry (LC-MS/MS) metabolomics and deep RNA sequencing (RNA-seq) to interrogate serum, kidney, and liver samples collected from 12 weeks and 18 months old male, female and castrated male mice to explore the versatile

84   perspectives of sex-related healthy aging mechanisms.

RESULTS

Castration is an effective model of healthy aging

To elucidate the effects of castration on lifespan extension and healthy aging, we employed multi-omics approaches to resolve the potential roles at both molecular and phenotypic levels (**Figure 1A)**. Young adult C57/BL6 mice were castrated at 8 weeks old (casM) and exhibited weight gain post-castration, but attained similar body weights as the sham males (shamM) by 60 weeks. This pattern align with findings in prepubertal castration models using UM-HET3 mice (**Figure 1B)** [10]**.** Hair graying, a common aging marker in both mice and humans [12], was notably delayed in casM compared to shamM at 13 months, indicating a more youthful phenotype in the castrated group (**Figure 1C**). Given the known correlation between ages 10 – 14 months in mice and 38 – 47 years in humans, where senescence-related changes become prominent [13], we conducted aging-related behavioral analyses. These included Barnes maze (BM), open field (OF) and novel object recognition (NOR) tasks to evaluate the spatial learning and memory, locomotor activity, and recognition memory in 12 – 13 month-old casM, shamM and sham female (shamF) mice. Previous studies have shown pronounced cognitive decline in middle-aged mice (8 – 12 months) as assessed by the BM test [14]. However, our results showed that middle-aged casM demonstrated superior performance, spending less time and making fewer errors with less immobile time in locating the escape hole than shamM and/or shamF (**Figure 1D-F**), hinting at a mitigated cognitive decline in casM (**Figure 1E-F**). Non-significant difference was found in OF and NOR tasks across the three groups (**Supplementary Figure 1**). More than possessing longer lifespans, our results revealed that castration confers a protective effect against cognitive decline. This selective enhancement in spatial learning and memory may reflect the differential influence of castration on brain regions, particularly affecting areas like the hippocampus that are crucial for navigation tasks, while exerting less impact on regions involved in locomotor and recognition memory functions.

Castration affected the secretion of steroid hormone spectrum

Hormones, serving as vital chemical messengers, travel through from bloodstream to tissues for coordinating a range of biological functions, with their dysregulation impacting development [15], aging [16] and disease progression [17]. The effect of surgical castration, specifically the removal of the testes, leads to a marked reduction

in testosterone (TS) levels. However, the broader hormone changes resulting from castration remain underexplored. Here, we employed ultra-high-performance liquid chromatography coupled with multiple reaction monitoring tandem mass spectrometry (UHPLC-MRM-MS/MS) to analyze 17 targeted steroid hormones (SHs) in the serum of casM, shamM and shamF mice, four weeks post-operation.

Given the limited volume of mouse serum available for SH measurement, we could only detect and compare 10 SH types across three groups. Notably, castration significantly altered SH secretion in casM (**Figure 2A**). As expected, testosterone and its more potent form, dihydrotestosterone, were remarkably reduced in CasM, to the point of being barely detectable (**Figure 2B-C**). Intriguingly, CasM mice exhibited a significant higher level of 17α-Hydroxyprogesterone (17α-OHP) (**Figure 2D**) and lower androstenedione (A4) compared to shamM (p < 0.05) (**Figure 2E**). These hormones, precursors to TS, might be part of a compensatory mechanism in response to TS deprivation [18]. In addition, we observed slight increases in aldosterone (ALD) and corticosterone (CORT) in casM, with p-values of 0.11 and 0.083, respectively (**Supplementary Figure 2**). ALD, a mineralocorticoid, and CORT, a glucocorticoid, are primarily produced by the adrenal cortex. Their upregulation may suggest an activated stress response and changes in mineral and glucose homeostasis consequent to TS deprivation.

To elucidate metabolites associated with TS, we conducted an untargeted metabolomics analysis by LC-MS/MS on serum samples from three groups of mice, four weeks post-operation. A total of 1495 compounds with MS^2 scores ≥ 0.3 were identified. Notably, coexpression analysis revealed that 17 of these compounds, which include 11 bioactive lipids, showed a significant correlation with TS levels ($|r| \geq 0.8$, p ≤ 0.001, **Figure 2F**), These 11 lipids encompass a range of molecules such as prostaglandin B2 (PGD2), a specific phosphatidylcholine (PC) variant, lysophosphatidylserine (LPS), lysophosphatidic acid (LPA), cyclic PA (CPA), glucuronosyldiacylglycerol (GlcADG), sulfurHexosylceramide (SHexCer), and four types of ceramide fatty acids (Cer) (**Supplementary Table 1**). While further investigation is required to solidify these findings, our coexpression analysis suggests that TS deprivation may significantly influence peripheral lipid metabolism, particularly impacting ceramide and lysophosphatide pathways.

Characterization of sex-related serum metabolomic features

To assess the serum metabolomic profile differences among shamM, shamF, and CasM mice at 12 weeks old (12w, 4 weeks post sham/castration procedures), principal component analysis (PCA) was performed using data from 1495 compounds identified by LC-MS/MS (**Figure 3A**). Lipid and lipid-like molecules constituted the largest category (~33%) of these compounds (Figure 3B). A comparison of shamM and shamF groups to explore sex-specific differences in serum metabolic activity revealed 255 significantly altered metabolites (SAMs) were identified (**Supplementary Table 2**), with the top SAMs (ranked by fold changes) displayed in **Supplementary Figure 3A.** Despite all mice receiving the same diet, we observed notably higher levels of antioxidants and anti-inflammatory compounds in shamF compared to shamM. These compounds, likely derive from food intake, included carnosic acid [19], arachidonic acid [20], choline [21], nicotinamide [22], D-pantothenic acid [23], gamma-crocetin [24], ruscogenin [25], 4-hydroxycinnamic acid [26], oleamide [27, 28] (**Figure 3C).** Conversely, shamM mice exhibited elevated levels of metabolites associated with resistance to high-fat diet-related diseases, such as 5-HEPE [29] and 12-HEPE [30], and a higher level of protective L-Anserine [31] than shamF (**Figure 3C**). Consistently, obesity-related compounds such as LysoPE(0:0/20:3(11Z,14Z,17Z)/0:0) (LysoPE) [32] and 7−Ketocholesterol (7-KC) [33] were found to be lower in male (**Figure 3C**), offering insights into why obesity is more prevalent in women than men [34].

Differential abundance (DS) score of the 255 SAMs was calculated to inform the overall changes of each pathway (Figure 3D). These SAMs were found to be highly enriched in lipid metabolism related pathways such as "sphingolipid signaling pathway", "regulation of lipolysis in adipocytes" and "arachidonic acid metabolism", suggesting a sex dimorphism in lipid metabolism. Additionally, a serum untargeted metabolomic analysis in the same groups at 18 months old identified 868 compounds, with "lipid and lipid-like molecules" again being the predominant category" (**Supplementary Figure 3B**). PCA analysis suggested that the metabolic profile of 18-month-old casM mice closely resembled that of shamF (**Supplementary Figure 3C**). Similar to the younger mice, older shamF demonstrated higher levels of protective compounds derived from the same diet compared to shamM (**Figure S3D, Supplementary Table 3**), suggesting a potentially stronger ability in females to generate healthy compounds from the same chow.

Metabolomic feminization and taurine enrichment in castrated male mice

In our analysis of 12-week-old mice serum, a comparative study between casM and shamF mice identified 170 SAMs. In contrast, comparisons of "casM vs shamM" and "shamF vs shamM" revealed 280 and 255 SAMs, respectively (**Figure 4A, Supplementary Table 4**). Hierarchical Clustering of the 1495 identified compound demonstrated a closer metabolic profile between casM and shamF than between casM and shamM (**Figure 4B**). These findings, achieved without altering genetic background, suggest that castration feminizes the metabolomic traits of male mice. When ranking these SAMs by fold change (FC), notable differences were observed between shamM and casM. The most significantly upregulated metabolite in casM was the unsaturated fatty acid, PI (16:1/18:2), with $\log_2 FC$ of 13.14, while the most downregulated was guanosine, with a $\log_2 FC$ of -6.79. CasM mice exhibited elevated levels of various metabolites associated with healthy aging, anti-skin aging, and anti-inflammation, such as ectoine [35], resolvin D2 [36], oleamide [27, 28], taurine [37], taurocholic acid [38], taurodeoxycholic acid [39], nicotinamide [40], ascorbic acid (vitamin C) [41], D-pantothenic acid (vitamin B5) [42, 43] and choline [44] (**Figure 4C**). Similar to shamF, casM mice also showed an enrichment of anti-oxidative metabolites likely derived from plant-based foods, including ganoderenic acid C [45], oleuropeic acid [46], fasciculol C [47], graveoline [48] (**Figure 4C**).

Enrichment analysis of the 280 SAMs from the "casM *vs* shamM" comparison categorized them into pathways such as "bile secretion", "vitamin digestion and absorption", "HIF-1 signaling pathway", "pyrimidine metabolism", various fatty acid metabolism pathways including "arachidonic acid metabolism" and "regulation of lipolysis in adipocytes". (**Figure 4D**). Interestingly, lipid metabolism-related pathways were consistently enriched in both the "casM vs shamM" and "shamF vs shamM" comparisons, indicating a potential sensitivity of lipid metabolism to the levels of SHs.

To investigate the metabolomic changes particularly derived from castration, union SAMs among 12w-old casM, shamM and shamF groups were further classified by c-means fuzzy clustering (**Figure 4E**, **Supplementary Table 5**). This analysis revealed that cluster1 metabolites were notably elevated in casM compared to both shamM and shamF. Topology-based pathway enrichment analysis indicated that cluster1 metabolites were significantly ($p < 0.05$) enriched in pathways like "primary bile acid biosynthesis", "D-glutamine and D-glutamate metabolism", "taurine and hypotaurine metabolism" among others (**Figure 4F**). Notably, taurine levels were higher in shamF than in shamM, and castration further augmented taurine levels along

with two key components of the taurine and hypotaurine metabolism pathway: taurodeoxycholic acid (TDCA) and taurocholic acid (TCA) (**Figure 4G**). However, in 18m-old mice, a higher level of taurine with no significant changes of TDAC and TCA were observed in shamM compared to shamF and casM, (**Figure 4H**), suggesting an age-related sex dimorphism of taurine concentrations in mice serum. Bile acids (BA) are crucial for maintaining lipid homeostasis and have been recognized as significant signaling molecules during aging process [48]. Given that the SAMs from both "casM vs shamM" and "shamF vs shamM" comparisons were significantly enriched in lipid-related metabolic pathways, and the primary function of taurine is the conjugation of cholesterol into bile acids [49], our findings suggest a potential link between taurine and the sex and/or SH-related differences in lipid metabolism. Taurine, along with TDCA and TCA, serve as antioxidants and protective factors against many degenerative diseases [50]. A recent study has demonstrated that blood taurine levels decline with age, but intake of taurine supplement at middle age can extend the healthy lifespan of mice and monkeys [37]. Considering the established longevity of females and castrated males over unaltered males, our results imply that mice with elevated taurine levels at younger ages may reap more significant benefits than those with higher levels at older ages.

Co-expression analysis across serum, livers and kidneys identifies potential taurine-related regulators

Kidney and liver are two major organs of organisms, the sex differences of renal and hepatic function have been aware over years [51]. To elucidate the sex and age-related regulatory networks in kidney and liver functions, we analyzed metabolome and transcriptome of renal cortex and left liver lobe from three groups of mice at 12 week (12w) and 18 months (18m) old. We detected an average of 899 compounds across both tissues, with 121 compounds common among kidneys, livers and serum samples in both age groups of sham, shamF, and casM mice (**Figure 5A** and **Supplementary Table 6**). Coexpression analysis of these 121 commonly detected metabolites (CDMs) revealed 4 major clusters (**Figure 5B**). Metabolite Set Enrichment Analysis (MSEA) indicated that Clusters 1, 2 and 3 are primarily involved in "Bile Acid Biosynthesis" and "Glycine and Serine Metabolism" pathways (**Supplementary Figure 5A-C**). Notably, two compounds known for their health-protective properties, taurine and alpha-linolenic acid (ALA), were grouped in Cluster 4. They exhibited a strong correlation ($r > 0.75$, p

< 0.01) with adenosine, xanthosine, 1-palmitoylglycerophosphoinositol, L-tyrosine,

erythrulose, glycolic acid, PC-M5', pyruvic acid and 5 other lipids include LPC(15:0),

LPE(18:2), LPE(20:3), LPS(20:4), FAHFA(18:1/22:3) (**Figure 5C**), suggesting a

potential link between Cluster 4 compounds and healthy aging. A recent study has

demonstrated that taurine intake can attenuate lipid peroxidative stress (LPOS) and lipid

metabolism disorder in mice, evidenced by decreased levels of malondialdehyde

(MDA), a marker of LPOS [52]. Similarly, ALA, an omega-3 polyunsaturated fatty acid

derived from plants, has also been shown to markedly reduce oxidative stress and

increase the lifespan of *Caenorhabditis elegans* in a dose-dependent manner [53].

Considering the fact that a linoleic acid-rich diet increases hepatic taurine [54] and the

observed strong association (r = 0.934) between taurine and ALA across three sample

types from three mouse groups and at two age stages, these results collectively suggest

potential role for ALA in taurine-mediated antioxidant defense and lipid metabolism

processes.

Due to the limited range of lipids identifiable by KEGG compounds database,

Cluster 4 compounds were showing enrichment in a few terms include "glyoxylate and

dicarboxylate metabolism", "tyrosine metabolism" and others (**Figure 5D**).

Noteworthy, xanthosine and adenosine, two key components of "purine metabolism"

pathway, were strikingly co-expressed with taurine (r = 0.973 and 0.874, respectively,

Figure 5C). Dysregulated purine metabolism has been implicated in many aging-

related disorders such as Alzheimer's diseases [55] and age-associated lower urinary

tract dysfunctions [56]. Considering taurine's role in aging delay [9], we hypothesized

that its influence on lifespan extension might also be linked to xanthosine and

adenosine-mediated purine metabolism, though further evidence is required to

substantiate this.

Considering the significance of taurine in aging and healthy aging, co-expression

analysis and c-means clustering were applied to identify genes that positively associated

with taurine. A total of 51 genes, whose expression levels showed similar dynamic trend

with taurine across casM, shamM and shamF mice in kidney and liver tissues at both

12w and 18m ages, were identified (**Figure 5E-G**). Among these, 15 genes exhibited a

generally decrease in expression in 18m tissues than 12w, such as *Hdac5, Preb, Plk2,*

Atxn2 and *Trit* (**Figure 5E**). Two key lipid metabolism genes, *Srebf1* and *Ppard* were

among the 51 taurine-associated genes. *Srebf1* encodes sterol regulatory element-

binding protein 1 (SREBP1), a critical transcription factor regulating lipogenes

involved in fatty acid and cholesterol biosynthesis [57]. *Ppard* encodes Peroxisome proliferator-activated receptor-delta (PPAR-δ), a member of the PPAR group in the nuclear receptor superfamily, known for its ligand-binding domains (LBD) that interacts with a broad range of lipid and lipid-like molecules [58]. This includes multiple types of polyunsaturated fatty acids, such as arachidonic and linoleic acid [58]. Both PPAR-δ and SREBP1 are also involved in retinoid X receptors/ Vitamin D Receptor (RXR/VDR) pathway (**Supplementary Table 7**). Given previous research showing that intestinal-specific deletion of VDR increased taurine levels [59], it is posited that taurine may also be associated with PPAR-δ and SREBP1 involved RXR/VDR pathway. Taken together, our results from both metabolomics and RNA-seq analysis strongly suggest that taurine is a key regulator in lipid peroxidation and metabolism, potentially playing a significant role in the process of healthy aging.

Joint metabolomic-transcriptomic analysis reveals a stronger capacity of defensing lipid peroxidation in female and castrated male

From deep RNA-seq analysis, we identified differentially expressed genes (DEGs) among casM, shamM and shamF groups in kidney and liver tissues (**Figure 6A-B, Supplementary Table 6**). In the kidneys of 12w old mice, over 4500 DEGs were noted in the comparisons of "shamF *vs* shamM" and "casM *vs* shamM," while only 987 genes showed differential expression between shamF and casM. Interestingly, livers seemed to be less affected by sex and/or the changes of SH levels than kidneys as only 2053 and 2533 DEGs were detected in "shamF *vs* shamM" and "casM *vs* shamM" comparisons, respectively, in 12w-old mice, and 546 DEGs between shamF and casM. Hierarchical clustering of RNA-seq data also revealed that the transcriptomic profiles of casM samples more closely resembled those of shamF than shamM samples, with the exception of 18m-old livers, suggesting a feminizing effect of castration that creates a crossover zone between male and female transcriptomic profiles in both kidney and liver (**Figure 6C**). However, the impact of sex and/or SH levels on transcriptomes, pronounced at 4 weeks post-castration, appeared to diminish in 18-month-old mice. In this older age group, the numbers of DEGs among the 3 groups were reduced, especially the comparisons of shamM with casM. In 18-month-old kidneys and livers of "casM *vs* shamM", it showed only 435 and 353 DEGs, respectively.

Mapping the DEGs onto mouse chromosomes, we found that DEGs in "shamF vs shamM" and "casM vs shamM" from both 12w and 18m-old liver tissues were

significantly clustered to chr4B3 (FDR < 0.05). Of the 145 annotated genes in chr4B3, 46 belong to the Major Urinary Proteins (*Mup*) family, with 25 being pseudogenes. Except for *Mup4* and *Mup5*, which were barely detectable, the other *Mup* family genes in chr4B3 were almost completely silenced in casM and shamF livers, showing an average log2FC = -5.57 (**Figure 6SA-B**). *Mup14*, *7*, *11*, and *1* were the top 4 differentially expressed *Mup* genes. It is known that *Mup* genes, predominantly expressed in the liver, exhibit sexually dimorphic traits. Male mice typically excrete 2–8 times more Mup protein in their urine compared to females [60]. Our results indicate that the expression of *Mup* genes is influenced not only by sex genotypes but also by testosterone levels.

To identify potential long-term salutary regulatory elements brought by sex and/or castration at old age, we conducted a joint metabolomic-transcriptomic analysis. Here, shamM mice, typically having a shorter liferspan, served as controls, while shamF and casM mice, associated with healthier and longer lifespans, were the focus [61]. We utilized the SAMs and DEGs from "shamF *vs* shamM" for the joint analysis to investigate female-specific signatures; meanwhile, the joint metabolomic-transcriptomic analysis was also performed using the SAMs and DEGs from "casM *vs* shamM" for interpreting castration-specific features. In 18m-old kidneys, both shamF and casM groups showed enrichment in KEGG programs related to "lysosome" and "cholesterol metabolism" (**Supplementary Table 8-9**). In the liver of the same age group, the significantly enriched KEGG pathways in the "casM vs shamM" comparison are illustrated in **Figure 6D**. Notably, 20 terms, such as "linoleic acid metabolism", "arachidonic acid metabolism", "retinol metabolism", "Jak-stat3 signaling" and "glutathione metabolism" were consistent with those in the "shamF vs shamM" comparison (terms marked by orange in **Figure 6D and 6E, , Supplementary Table 10-11**). These KEGG terms are known to play critical roles in lipid peroxidation.

To further validate these findings, we measured levels of malondialdehyde (MDA) - a final product of polyunsaturated fatty acids peroxidation, in the three groups of mice. Results indicated that shamF mice exhibited the lowest, while shamM mice had the highest MDA levels in 18m-old livers, suggesting that females and castrated males experience less lipid oxidation stress than their male counterparts (**Figure 6F**). Additional oxidative stress markers, SOD1 and SOD2, were also tested and showed similar trends to MDA among three groups (**Supplementary Figure 6C**). Taken together, our analysis suggests a close association between lipid metabolism and sex

and SH levels. CasM and shamF mice are likely to absorb more health-beneficial compounds from their diet and experience reduced lipid peroxidative stress, potentially contributing to their longer lifespans.

Evaluating sex-dependent cisplatin toxicity through castration models

Drug-induced tissue toxicity presents a significant challenge in the clinical application of anti-tumor drugs. Platinum-based drugs (PBDs), such as cisplatin, is the mainstay of cancer therapy, with nearly half of all patients receiving chemotherapy treated with PBDs [62]. However, the therapeutic application of these drugs is tempered by adverse side effects, including nephrotoxicity and hepatotoxicity, including nephrotoxicity and hepatotoxicity, which exhibit differential impacts between male and female patients [63]. To evaluate the utility of the castration model in studying sex-dependent toxicity studies, we treated 12w-old mice (4 weeks post-castration/sham procedure) with a low dose of cisplatin twice, every 72 hours. It is well-established that the kidney, and to a lesser extent the liver, are susceptible to cisplatin-induced damage. Consequently, we harvested renal cortex and left liver lobe tissues post-treatment for histological analysis.

Hematoxylin and eosin (H&E) staining of liver tissues showed the most severe liver injury in shamF mice, characterized by marked steatosis, sinusoidal congestion, and hemorrhage. In contrast, sham-operated male (shamM) and castrated male (casM) mice displayed comparatively less hepatic damage (**Figure 6G**). Conversely, Periodic Acid Schiff (PAS) staining of kidney tissues showed that casM mice had more severe kidney damages than both shamF and shamM, with a notable increase in acute renal tubule injury (**Figure 6H**). Previous studies on cisplatin-induced hepatotoxicity have yielded limited insights regarding sex differences. However, substantial evidence indicates that males are more susceptible to cisplatin-induced nephrotoxicity [63]. Studies administering cisplatin to rats at both 2 weeks (1 mg/kg/day) and a single dose (7.5 mg/kg) have shown that males experience more severe nephrotoxic effects than females [64]. Our pathological observations, combined with previous findings and the fact that castrated male (casM) mice have the lowest testosterone (TS) levels, suggest that intermediate TS levels, similar to those in shamF mice, may offer greater protection against renal toxicity induced by cisplatin.

DISCUSSION

The burgeoning aging population has become a global concern, poised to increase the

prevalence of age-related diseases [65]. Hence, it is imperative to gain a better

understanding of the regulatory factors contributing to healthy longevity. Historically,

sexual dimorphism in aging has been largely overlooked, despite the fact that females

consistently outlive males. While castration is a common practice in animal husbandry,

known to extend the lifespan of domesticated dogs and cats in veterinary medicine, its

implications for aging research have only recently been explored [66]. For instance,

castrated domestic sheep exhibit both extended lifespan and decelerated epigenetic

aging compared to their intact male counterparts [9]. Similarly, survival assays in

genetically diverse mice reveal that castration extends longevity, implicating post-

pubertal testicular actions as key determinants of sex differences in lifespan and growth

trajectories [10]. Moreover, castration has been shown to induce metabolic remodeling

in male hearts [67] and mitigate the formation of aging-induced abdominal aortic

aneurysms [68].

Nevertheless, the utility of castration as a model for studying sex differences in

healthy aging and the underlying molecular mechanisms remain underexplored. In this

study, we employed multi-omics approaches to systematically investigate sex

differences in lifespan at both young (12w) and old (18m) stages of mice. Intriguingly,

despite being fed the same diet, shamF and casM exhibited elevated levels of multiple

food-derived health-promoting compounds than shamM. Given the influential role of

gut microbiome in aging and its known sex-dependent modulation [69, 70], we

hypothesize that sex hormones may involve in the regulation the microbial community,

thereby affecting the bioavailability of these protective compounds from food.

Previous research has offered mixed insights into the impact of castration on

cognitive function. One such study employed a dual-task approach – utilizing an 8-arm

radial maze and a step-down avoidance task – to assess spatial and short-term memory

in rats. Conducted approximate 82 days post-surgical and chemical castration, the study

concluded that castration deteriorates memory function [71]. In contrast, our findings,

based on BM test in castrated mice, suggest an enhancement in cognitive performance.

Several factors could account for these divergent outcomes. First, the timing of

cognitive assessments post-castration differed between the studies. Our behavioral tests

were performed on 12-13month old mice, nearly 10-11 months after castration or sham

operations. It is plausible that the benefits of castration on spatial memory may not be

readily apparent in younger rodents but become more pronounced with age. Second,

the choice of behavioral tasks could also contribute to the observed discrepancies. The

step-down inhibitory avoidance task, employed in the previous study, uses immediate electric shocks as stimuli to evaluate aversive memory, while the 8-arm radial maze task requires deprivation of food and/or water for animals. Electric shock is a type of psychological stress for rodents [72]. Mounting evidence suggests that testosterone significantly modulates physiological responses to stressors [73]; thus, the markedly reduced testosterone levels in castrated mice could potentially confound the evaluation of cognitive performance. To mitigate potential confounders, we opted for less stressful tasks – namely BM, OF and NOR tests. Our results from the BM test indicate that castration not only extends lifespan but also attenuates cognitive decline in older mice. This suggests that the castration model offers a nuanced understanding of aging, one that incorporates both longevity and cognitive preservation.

Sex as a key determinant biological variable has been historically overlooked despite sexual dimorphism can be observed everywhere. On one hand, the bias toward using male subjects in biomedical research has often been justified by the desire to minimize variability, particularly traits influenced by the female estrous cycle [8]. This male-centric focus has inadvertently limited our understanding of tissue function and disease mechanisms from a sex-specific perspective, thereby highlighting the pressing need for more inclusive animal models. On the other hand, straightforwardly comparing male with female is generally applied in the increasingly emphatic but still limited sex-difference studies, which made the leading mechanisms hard to recognize due to complicated variables between male and female in both genetic and non-genetic aspects. To broaden the feasibility and practicability of applying castration model in sex-dependent biological queries, we tested the differences of tumor drugs-induced tissue injuries among the three groups of mice. More sever renal damages were observed in male than female after cisplatin treatment at both young and old ages [74]; however, the roles of genetic and non-genetic co-determinants involved in this process are still elusive. Rostami et al. found that 1 week post-castration rats with low dosage treatment of TS rather than high dosage, were more tolerant with cisplatin nephrotoxicity [75], while our pathological assays showed that castration deteriorated the renal toxicity. Taken together, we believe that the balanced levels of TS are essential to defense the toxicity. In summary, our study, in conjunction with methylation data in sheep [9], advocates for the castration model as a promising approach. It effectively "feminizes" males at multiple molecular levels through alterations in SHs, while preserving the somatic male genotype. This unique duality positions the castration model as a useful

tool for distinguishing between hormone-induced sex differences and genotypic sex,
broadening its utility beyond the scope of aging research.

Our comprehensive metabolomic and transcriptomic profiling across multiple
tissues at both young and old life stages offers compelling insights. We observed
multiple lipid related pathways were identified recurrently in comparisons of both
"shamF *vs* shamM" and "casM *vs* shamM", and reduced lipid peroxidation in both
casM and shamF, indicative of a shared molecular pathway that may contribute to
healthier aging. Moreover, our data indicates that castration significantly attenuates the
molecular differences mediated by sex and/or sex hormones across multiple omic layers.
Driven by the need for a stable and broadly applicable model for the study of sex-
dependent responses to cancer therapy, we further extended our exploration to include
the feasibility of the castration model in assessing cancer drug toxicities, in addition to
examining the healthy aging phenotype. These explorations and findings collectively
underscore the castration model's versatility and its untapped potential for advancing
our understanding of sex differences across various physiological and pathological
landscapes.

METHODS

Animals, surgical castration procedures, and sample collections

C57/BL6 mice were purchased from Zhuhai BesTest Bio-Tech Co,.Ltd. and housed in the animal facility of Shenzhen TopBiotech Co., Ltd (Shenzhen, China) under pathogen-free condition with free access to chow and water. Male and female mice at 8 weeks old were anesthetized by isoflurane for castration (testicles removed) or sham procedures. Mice were placed on a warm pad after suture, then transferred to cages while awake. 4-weeks post-castration/sham operations, serum was collected for steroid hormone detection and untargeted metabolites analysis while liver and kidney were dissected for untargeted metabolites analysis and RNA-seq (n=3-6/group). Castrated/sham mice at 12-13m old were subjected to behavioral tests and were sacrificed for sample collection at 18m. At 18m, serum was used for untargeted metabolomics while liver and kidneys were dissected for both untargeted metabolomics and RNA-seq (n=3-6/group). The animal study protocol was approved by the Animal Ethics Review Board of Shenzhen TopBiotech Animal Facilities (No. TOP-IACUC-2021-0100).

Sample treatment for targeted steroid hormone spectrum detection

50 μL serum was thawed and vortexed for 30s following by mixture with purified H_2O, internal standard working solution and methanol. The well mixed samples were then subjected to repeated 13000 rpm centrifuge for 10 minutes (mins) under 4°C. The supernatant was retained further purified with OASIS PRiME HLB uElution (SPE). 200 μL methanol and 200 μL H_2O were separately used to wash and equilibrate the SPE cartridges. The supernatant was loaded to the equilibrated cartridges and then washed by 200 μL 10% ACN/H_2O (v/v) and 200 μL Hexane. The flow-through fraction was discarded. Rinse the cartridge was with 40 μL 90% ACN/H_2O (v/v) and collect the cluent. Add 60 μL H_2O to the cluent and shake the mixture for 3 mins. The final mixture and the same volume of isotopically-labelled internal standards were subjected to UHPLC-MS/MS analysis.

Serum sample preparation for untargeted MS analysis

150 μL methanol which was premixed with isotopically-labelled internal standard compounds was added to 50 μL mouse serum and vortexed for 30 seconds to mix well. The mixture was then sonicated for 10 mins in ice water bath and incubated for 1 h at -

40 °C to precipitate proteins. After 15-min high-speed centrifugation, the supernatant was used for MS analysis. An equal aliquot of the supernatants obtained from each serum sample was mixed as the quality control (QC).

Kidney/liver sample preparation for untargeted MS analysis

10-25 mg tissues were submerged with extraction solution (methanol:water = 3: 1, with isotopically-labelled internal standard mixture) and homogenized in ice-water bath. Incubate homogenized samples for 1 h at -40°C and centrifuge at 12000 rpm for 15 min at 4 °C. The supernatant was retained for MS analysis. A volume matched quality control (QC) sample was prepared for MS.

UHPLC-MS/MS analysis for targeted steroid hormone spectrum

EXIONLC System (Sciex) equipped with a Waters Acquity Uplc Bch C8 column was utilized for UHPLC analysis at Biotree Technology Co., Ltd. Assay development was applied by SCIEX 6500+ triple quadrupole mass spectrometer (Sciex) with an IonDrive Turbo V electrospray ionization (ESI) interface, under multiple reaction monitoring (MRM) module. MassLynx Work Station Software (Version 4.1) was employed for MRM data acquisition and preprocessing.

UHPLC-MS/MS untargeted metabolomics analysis

T3 chromatographic column with affinities to both lipid and hydrophile at Biotree Technology Co., Ltd was used to acquire the metabolome of castrated male (casM), sham male (shamM) and sham female (shamF) mice by Orbitrap mass spectrometer. Joint negative (NEG) and positive (POS) ionization modes were employed for detecting metabolites based on their mass to charge ratios. Raw data was converted to the mzXML format using ProteoWizard platform (https://proteowizard.sourceforge.io/) and processed with an in-house R script and XCMS tool [76] for peak detection, extraction, alignment, and integration. Data was subjected to fragment extracted ion chromatogram (MS^2) annotations as previously described [77], MS^2 metabolites with scores > 0.3 were retained for next-step analysis. These compounds were mapped to Human Metabolome Database (HMDB) and classified into multiple categories. The biggest category under both NEG and POS modes is lipid and lipid-like molecules, which counts for ≥ 30% identified metabolites (**Figure 3a-b, Supplementary Table 2-3**). Compounds from NEG and POS modes were merged for down-stream analysis.

Behavioral tests

Three groups of mice include casM, shamM and shamF at 12-13 months old (n= 6-13) were subjected to BM, OF and NOR tasks. A 60 cm high, 120 cm-diameter and 60 mm thick circular wheeled PVC slab with 25 cm-diameter holes was used for BM tests. A modified protocol was employed following the standard methods in previous study [78]. Briefly, mice were firstly trained in continuous 7 days to find the escape hole within 4 mins; if failed, mice were finally guided to the target hole gently. Mice were subjected to the probe trial on day 8. OF test was applied to test the exploratory behavior and general activity of mice, mice were placed in an enclosure, rectangular tank with surrounding walls for 5 mins, distance moved, time spent moving, rearing, and change in activity over time for each mouse were recorded. NOR test was performed following previous study [79]. Mice were placed in the empty tank for 10 mins on day1; day 2, two objects were placed in the tank, and mice were allowed for 10 mins to explore; one of the objects was replaced by a new one on day 3, and time spent for exploration (TE) within 5 mins for each object was recorded. NOR index was calculated via this formulation: NOR index = new object TE- old object TE/ Total TE. Mice behaviors during these tests were recorded and analyzed by ANY-MAZE software (https://www.any-maze.com/).

Western-blot

Total protein was isolated on ice by cell & tissue lysis buffer (Cat#P0013, Beyontime, China) and quantitated using BCA assay kit (Cat#P0012, Beyontime, China). Approximate 15ug protein was used for western-blot assays following previous protocols [65]. Antibodies for MDA, SOD1 and SOD2were purchased from abcam (Cat# ab243066, ab13498 and ab13533), GAPDH was purchased from Beyontime (Cat# AF0006).

RNA extraction for RNA-seq analysis

Total RNA was extracted using Trizol reagents (Invitrogen Cat#15596026) as we have described before [80]. PolyA-tailed RNA was enriched for constructing 150bp pair-end RNA-sequencing libraries using MGISEQ-2000RS kit (MGI Cat#1000012555), and sequenced by DNBseq MGI2000 platform at BGI genomics Co,. Ltd.

Cisplatin administration

Mice were administrated for intraperitoneal injection of cisplatin (Cat# 232120, sigma) at 4 weeks post-castration/sham operations, with a dosage of 5 mg/kg every72 hours for 2 times. Mice were sacrificed 48 hours after cisplatin treatment for the collection of liver and kidney tissues.

Histopathological analysis

For liver and kidney tissue examination, samples were first fixed in 10% formalin solution for 48 hours, followed by paraffin embedding. Sections of 3 μm thickness were then prepared for histological analysis. Hematoxylin and Eosin (H & E) and Periodic Acid-Schiff (PAS) staining were performed according to established protocols. Liver injury was evaluated by visually inspecting the patterns of steatosis, sinusoidal congestion, and hemorrhage. Renal injury was quantitatively assessed using the scoring criteria established in a previous study [81].

Identification of significant altered metabolites (SAMs) and differentially expressed genes (DEGs)

A strict pipeline was applied to identify SAMs. Firstly, the normalized peak values of metabolites were subjected to unsupervised principal component analysis (PCA) and then administrated to supervised orthogonal projections to latent structures-discriminate analysis (OPLS-DA) for data modeling by commercial SIMCA software (V16.0.2). A 7-fold cross validation was employed for model selection. To evaluate the predictive power and robustness of OPLS-DA model, a 200-times-permutations was performed and the value of variable importance in the projection (VIP) of the first principal component in OPLS-DA analysis was calculated which represents the contribution of each variable to the model. Metabolites with VIP > 1 and students' t-test p-value < 0.05 were recognized as SAMs. For DEG identification, quality control was performed for raw RNA-seq reads by SolexaQA (-h 30, -l 30) [82]. Cleaned reads were subjected to Hisat2 [83] for mapping. Cufflinks package [84] was utilized to calculate the Fragments Per Kilobase of transcript per Million mapped reads (FPKM) values and to identify DEGs use FDR < 0.05 as the cutoff value.

Expression trend clustering analysis

To identify expression trend patterns of metabolites among the serum samples of 12w

casM, shamM and shamF, the union of SAMs across three pairwise comparisons in Figure 4A were analyzed by Mfuzz package (c=9, m =1.25) (Figure 4E) [85]. To illustrate genes that potentially related with taurine, the expression levels (FPKM values) of all the annotated genes in kidneys and livers at 12w and 18m casM, shamM and shamF mice, were also subjected to Mfuzz package [85] for clustering (c=9, m =1.25). The semi-quantitative concentrations of taurine were plotted as stack lines. Genes that show similar expression trend with taurine in all the tissues were retained (Figure 5F-G).

Hierarchical clustering analysis

Hierarchical clustering analysis for SH levels after castration/sham treatments (Figure 2A), the expression of 51 taurine related genes (Figure 5E), serum metabolomic profiles among casM, shamM and shamF using (Figure 4B) and tissue RNA-seq gene expression profiles (Figure 6C), were performed by Hclust function in R platform using Euclidean distance metric and complete lineage. The z-score standardized SH levels and expression levels of taurine related genes after clustering were visualized via heatmap using pheatmap 1.0.12 package.

Enrichment analysis of SAMs

SAMs were subjected for topology-based KEGG pathway enrichment analysis using MetaboAnalyst with default parameters [86]. Differential abundance (DA) scores [87] were calculated for the significant enrichment terms using this formula: DA scores =

$$\frac{\text{Up regulated metabolite No.in the pathway}-\text{Downregulated metabolite No. in the pathway}}{\text{Total metabolite No.of the pathway}}$$

Joint metabolome & transcriptome analysis

To examine the interplay between metabolomics and other transcriptomics layers, SAMs and DEGs were used for joint metabolome & transcriptome enrichment analysis against KEGG pathways via MetaboAnalystR under joint-analysis module by hypergeometric test with default parameters. FDR <.0.05 was applied as the cut-off value [86, 88].

REFERENCE

1. Lemaître, J.-F., et al., *Sex differences in adult lifespan and aging rates of mortality across wild mammals.* Proceedings of the National Academy of Sciences, 2020. **117**(15): p. 8546-8553.

2. Martínez-Sellés, M., et al., *Sex-Related Differences in Centenarians and Their Hearts.* J Am Geriatr Soc, 2016. **64**(2): p. 444-6.

3. Evert, J., et al., *Morbidity profiles of centenarians: survivors, delayers, and escapers.* J Gerontol A Biol Sci Med Sci, 2003. **58**(3): p. 232-7.

4. Miszko, T.A., et al., *Effect of strength and power training on physical function in community-dwelling older adults.* J Gerontol A Biol Sci Med Sci, 2003. **58**(2): p. 171-5.

5. Marais, G.A.B., et al., *Sex gap in aging and longevity: can sex chromosomes play a role?* Biol Sex Differ, 2018. **9**(1): p. 33.

6. Horstman, A.M., et al., *The role of androgens and estrogens on healthy aging and longevity.* J Gerontol A Biol Sci Med Sci, 2012. **67**(11): p. 1140-52.

7. Iorga, A., et al., *The protective role of estrogen and estrogen receptors in cardiovascular disease and the controversial use of estrogen therapy.* Biol Sex Differ, 2017. **8**(1): p. 33.

8. Mauvais-Jarvis, F., A.P. Arnold, and K. Reue, *A Guide for the Design of Pre-clinical Studies on Sex Differences in Metabolism.* Cell Metab, 2017. **25**(6): p. 1216-1230.

9. Sugrue, V.J., et al., *Castration delays epigenetic aging and feminizes DNA methylation at androgen-regulated loci.* Elife, 2021. **10**.

10. Jiang, N., et al., *Prepubertal castration eliminates sex differences in lifespan and growth trajectories in genetically heterogeneous mice.* Aging Cell, 2023. **22**(8): p. e13891.

11. Min, K.J., C.K. Lee, and H.N. Park, *The lifespan of Korean eunuchs.* Curr Biol, 2012. **22**(18): p. R792-3.

12. Zhang, Z., et al., *Wnt/β-catenin signaling promotes aging-associated hair graying in mice.* Oncotarget, 2017. **8**(41): p. 69316-69327.

13. Bresilla, D., et al., *The sex-specific metabolic signature of C57BL/6NRj mice during aging.* Sci Rep, 2022. **12**(1): p. 21050.

14. Raldiris, T.L., et al., *Cognitive control beliefs and cognitive functioning in mid-to late-life.* Br J Dev Psychol, 2021. **39**(2): p. 269-281.

15. Kraemer, W.J., et al., *Growth Hormone(s), Testosterone, Insulin-Like Growth Factors, and Cortisol: Roles and Integration for Cellular Development and Growth With Exercise.* Front Endocrinol (Lausanne), 2020. **11**: p. 33.

16. van den Beld, A.W., et al., *The physiology of endocrine systems with ageing.* Lancet Diabetes Endocrinol, 2018. **6**(8): p. 647-658.

17. Crafa, A., et al., *The Burden of Hormonal Disorders: A Worldwide Overview With a Particular Look in Italy.* Front Endocrinol (Lausanne), 2021. **12**: p. 694325.

18. Turcu, A., et al., *Adrenal androgens and androgen precursors-definition, synthesis, regulation and physiologic actions.* Compr Physiol, 2014. **4**(4): p. 1369-81.

19. Lin, C., et al., *Effects on longevity extension and mechanism of action of carnosic acid in Caenorhabditis elegans.* Food Funct, 2019. **10**(3): p. 1398-1410.

20. Das, U.N., *Ageing: Is there a role for arachidonic acid and other bioactive lipids? A review.* J Adv Res, 2018. **11**: p. 67-79.

21. Wortmann, S.B. and J.A. Mayr, *Choline-related-inherited metabolic diseases-A mini review.* J Inherit Metab Dis, 2019. **42**(2): p. 237-242.

22. Igarashi, M., et al., *Chronic nicotinamide mononucleotide supplementation elevates blood nicotinamide adenine dinucleotide levels and alters muscle function in healthy older men.* npj Aging, 2022. **8**(1): p. 5.

23. Hong, Y., et al., *Association between plasma Vitamin B5 levels and all-cause mortality: A nested case-control study.* J Clin Hypertens (Greenwich), 2022. **24**(7): p. 945-954.

24. Mohammadi, E., et al., *Protective effect of crocin against d-galactose-induced aging in mice.* Avicenna J Phytomed, 2018. **8**(1): p. 14-23.

25. Lu, H.J., et al., *Ruscogenin ameliorates diabetic nephropathy by its anti-inflammatory and anti-fibrotic effects in streptozotocin-induced diabetic rat.* BMC Complement Altern Med, 2014. **14**: p. 110.

26. Sova, M. and L. Saso, *Natural Sources, Pharmacokinetics, Biological Activities and Health Benefits of Hydroxycinnamic Acids and Their Metabolites.* Nutrients, 2020. **12**(8).

27. Heo, H.J., et al., *Effects of oleamide on choline acetyltransferase and cognitive activities.* Biosci Biotechnol Biochem, 2003. **67**(6): p. 1284-91.

28. Ano, Y., et al., *Preventive effects of a fermented dairy product against Alzheimer's disease and identification of a novel oleamide with enhanced microglial phagocytosis and anti-inflammatory activity.* PLoS One, 2015. **10**(3): p. e0118512.

29. Wang, C., et al., *Hydroxyeicosapentaenoic acids and epoxyeicosatetraenoic acids attenuate early occurrence of nonalcoholic fatty liver disease.* Br J Pharmacol, 2017. **174**(14): p. 2358-2372.

30. Nagatake, T., et al., *12-Hydroxyeicosapentaenoic acid inhibits foam cell formation and ameliorates high-fat diet-induced pathology of atherosclerosis in mice.* Sci Rep, 2021. **11**(1): p. 10426.

31. Rokicki, J., et al., *Daily Carnosine and Anserine Supplementation Alters Verbal Episodic Memory and Resting State Network Connectivity in Healthy Elderly Adults.* Front Aging Neurosci, 2015. **7**: p. 219.

32. Cai, H., et al., *Metabolomic signatures for liver tissue and cecum contents in high-fat diet-induced obese mice based on UHPLC-Q-TOF/MS.* Nutr Metab (Lond), 2021. **18**(1): p. 69.

33. Anderson, A., et al., *7-Ketocholesterol in disease and aging.* Redox Biol, 2020. **29**: p. 101380.

34. Kanter, R. and B. Caballero, *Global gender disparities in obesity: a review.* Advances in nutrition, 2012. **3**(4): p. 491-498.

35. Bilstein, A., et al., *Ectoine in the Treatment of Irritations and Inflammations of the Eye Surface.* Biomed Res Int, 2021. **2021**: p. 8885032.

36. Zhao, M., et al., *Resolvin D2 and its receptor GPR18 in cardiovascular and metabolic diseases: A promising biomarker and therapeutic target.* Pharmacol Res, 2023. **195**: p. 106832.

37. Singh, P., et al., *Taurine deficiency as a driver of aging.* Science, 2023. **380**(6649): p. eabn9257.

38. Warden, C., J.M. Barnett, and M.A. Brantley, Jr., *Taurocholic acid inhibits features of age-related macular degeneration in vitro.* Exp Eye Res, 2020. **193**: p. 107974.

39. Wolf, P.G., et al., *Effects of taurocholic acid metabolism by gut bacteria: A controlled feeding trial in adult African American subjects at elevated risk for*

colorectal cancer. Contemp Clin Trials Commun, 2020. **19**: p. 100611.

40.   Ur Rahman, S., A. Qadeer, and Z. Wu, *Role and Potential Mechanisms of*
Nicotinamide Mononucleotide in Aging. Aging Dis, 2023.

41.   Harrison, F.E., *A critical review of vitamin C for the prevention of age-related*
cognitive decline and Alzheimer's disease. J Alzheimers Dis, 2012. **29**(4): p.
711-26.

42.   Pelton, R.B. and R.J. Williams, *Effect of pantothenic acid on the longevity of*
mice. Proc Soc Exp Biol Med, 1958. **99**(3): p. 632-3.

43.   Ishiguro, K., *Aging effect of blood pantothenic acid content in female.* Tohoku
J Exp Med, 1972. **107**(4): p. 367-72.

44.   Dave, N., et al., *Dietary choline intake is necessary to prevent systems-wide*
organ pathology and reduce Alzheimer's disease hallmarks. Aging Cell, 2023.
22(2): p. e13775.

45.   Wang, J., et al., *Emerging Roles of Ganoderma Lucidum in Anti-Aging.* Aging
Dis, 2017. **8**(6): p. 691-707.

46.   Kim, Y.J., et al., *Oleanolic Acid Protects the Skin from Particulate Matter-*
Induced Aging. Biomol Ther (Seoul), 2021. **29**(2): p. 220-226.

47.   Ványolós, A., et al., *Extracts and Steroids from the Edible Mushroom*
Hypholoma lateritium Exhibit Anti-Inflammatory Properties by Inhibition of
COX-2 and Activation of Nrf2. Chem Biodivers, 2020. **17**(9): p. e2000391.

48.   Li, Z., et al., *Graveoline Analogs Exhibiting Selective Acetylcholinesterase*
Inhibitory Activity as Potential Lead Compounds for the Treatment of
Alzheimer's Disease. Molecules, 2016. **21**(2): p. 132.

49.   Murakami, S., et al., *Taurine ameliorates cholesterol metabolism by stimulating*
bile acid production in high-cholesterol-fed rats. Clin Exp Pharmacol Physiol,
2016. **43**(3): p. 372-8.

50.   Jakaria, M., et al., *Taurine and its analogs in neurological disorders: Focus on*
therapeutic potential and molecular mechanisms. Redox Biol, 2019. **24**: p.
101223.

51.   Guy, J. and M.G. Peters, *Liver disease in women: the influence of gender on*
epidemiology, natural history, and patient outcomes. Gastroenterol Hepatol (N
Y), 2013. **9**(10): p. 633-9.

52.   Guo, Q., et al., *Taurine Attenuates Oxidized Fish Oil-Induced Oxidative Stress*
and Lipid Metabolism Disorder in Mice. Antioxidants (Basel), 2022. **11**(7).

53.   Qi, W., et al., *The ω-3 fatty acid α-linolenic acid extends Caenorhabditis*
elegans lifespan via NHR-49/PPARα and oxidation to oxylipins. Aging Cell,
2017. **16**(5): p. 1125-1135.

54.   Hajri, T., A. Pronczuk, and K. Hayes, *Linoleic acid-rich diet increases hepatic*
taurine and cholesterol 7α-hydroxylase activity in conjunction with altered bile
acid composition and conjugation in gerbils. The Journal of nutritional
biochemistry, 1998. **9**(5): p. 249-257.

55.   Wang, H., et al., *Mutations in SREBF1, Encoding Sterol Regulatory Element*
Binding Transcription Factor 1, Cause Autosomal-Dominant IFAP Syndrome.
Am J Hum Genet, 2020. **107**(1): p. 34-45.

56.   Birder, L.A. and E.K. Jackson, *Dysregulated Purine Metabolism Contributes to*
Age-Associated Lower Urinary Tract Dysfunctions. Adv Geriatr Med Res, 2021.
3(4).

57.   Sekiya, M., et al., *SREBP-1-independent regulation of lipogenic gene*
expression in adipocytes. J Lipid Res, 2007. **48**(7): p. 1581-91.

58.   Liu, Y., et al., *The Role of PPAR-δ in Metabolism, Inflammation, and Cancer:*

793   *Many Characters of a Critical Transcription Factor.* Int J Mol Sci, 2018. **19**(11).

794   59.   Chatterjee, I., et al., *Vitamin D receptor promotes healthy microbial metabolites*
795   *and microbiome.* Sci Rep, 2020. **10**(1): p. 7340.

796   60.   Penn, D.J., S.M. Zala, and K.C. Luzynski, *Regulation of Sexually Dimorphic*
797   *Expression of Major Urinary Proteins.* Front Physiol, 2022. **13**: p. 822073.

798   61.   Lu, Y., Z. Pang, and J. Xia, *Comprehensive investigation of pathway enrichment*
799   *methods for functional interpretation of LC-MS global metabolomics data.*
800   Brief Bioinform, 2023. **24**(1).

801   62.   Johnstone, T.C., G.Y. Park, and S.J. Lippard, *Understanding and improving*
802   *platinum anticancer drugs--phenanthriplatin.* Anticancer Res, 2014. **34**(1): p.
803   471-6.

804   63.   Marcu, L.G., *Gender and Sex-Related Differences in Normal Tissue Effects*
805   *Induced by Platinum Compounds.* Pharmaceuticals (Basel), 2022. **15**(2).

806   64.   Nematbakhsh, M., et al., *Gender difference in Cisplatin-induced nephrotoxicity*
807   *in a rat model: greater intensity of damage in male than female.* Nephrourol
808   Mon, 2013. **5**(3): p. 818-21.

809   65.   Jiang, J., et al., *Decoding the role of long noncoding RNAs in the healthy aging*
810   *of centenarians.* Brief Bioinform, 2021. **22**(5).

811   66.   Kent, M.S., et al., *Longevity and mortality in cats: A single institution necropsy*
812   *study of 3108 cases (1989-2019).* PLoS One, 2022. **17**(12): p. e0278199.

813   67.   Svedlund Eriksson, E., et al., *Castration of Male Mice Induces Metabolic*
814   *Remodeling of the Heart.* J Endocr Soc, 2022. **6**(11): p. bvac132.

815   68.   Zhang, X., et al., *Castration of male mice prevents the progression of*
816   *established angiotensin II-induced abdominal aortic aneurysms.* J Vasc Surg,
817   2015. **61**(3): p. 767-76.

818   69.   Ghosh, T.S., F. Shanahan, and P.W. O'Toole, *The gut microbiome as a modulator*
819   *of healthy ageing.* Nat Rev Gastroenterol Hepatol, 2022. **19**(9): p. 565-584.

820   70.   McGee, J.S. and C. Huttenhower, *Of mice and men and women: Sexual*
821   *dimorphism of the gut microbiome.* Int J Womens Dermatol, 2021. **7**(5Part A):
822   p. 533-538.

823   71.   Shin, M.S., et al., *Long-term Surgical and Chemical Castration Deteriorates*
824   *Memory Function Through Downregulation of PKA/CREB/BDNF and c-*
825   *Raf/MEK/ERK Pathways in Hippocampus.* Int Neurourol J, 2019. **23**(2): p. 116-
826   124.

827   72.   Kim, J.Y., et al., *Mice subjected to uncontrollable electric shocks show*
828   *depression-like behaviors irrespective of their state of helplessness.* Behav
829   Brain Res, 2017. **322**(Pt A): p. 138-144.

830   73.   Kutlikova, H.H., et al., *The effects of testosterone on the physiological response*
831   *to social and somatic stressors.* Psychoneuroendocrinology, 2020. **117**: p.
832   104693.

833   74.   Eshraghi-Jazi, F. and M. Nematbakhsh, *Sex Difference in Cisplatin-Induced*
834   *Nephrotoxicity: Laboratory and Clinical Findings.* J Toxicol, 2022. **2022**: p.
835   3507721.

836   75.   Rostami, B., et al., *Effect of testosterone on Cisplatin-induced nephrotoxicity in*
837   *surgically castrated rats.* Nephrourol Mon, 2014. **6**(5): p. e21546.

838   76.   Xia, J. and D.S. Wishart, *MSEA: a web-based tool to identify biologically*
839   *meaningful patterns in quantitative metabolomic data.* Nucleic Acids Res, 2010.
840   **38**(Web Server issue): p. W71-7.

841   77.   Alseekh, S., et al., *Mass spectrometry-based metabolomics: a guide for*
842   *annotation, quantification and best reporting practices.* Nat Methods, 2021.

18(7): p. 747-756.

78.   Ederer, M.L., et al., *Voluntary Wheel Running in Old C57BL/6 Mice Reduces*
Age-Related Inflammation in the Colon but Not in the Brain. Cells, 2022. **11**(3).

79.   Lueptow, L.M., *Novel Object Recognition Test for the Investigation of Learning*
and Memory in Mice. J Vis Exp, 2017(126).

80.   Luo, Y.M., et al., *Insights into the mechanisms of triptolide nephrotoxicity*
through network pharmacology-based analysis and RNA-seq. Front Plant Sci,
2023. **14**: p. 1144583.

81.   Rong, S., et al., *The TIM-1:TIM-4 pathway enhances renal ischemia-*
reperfusion injury. J Am Soc Nephrol, 2011. **22**(3): p. 484-95.

82.   Cox, M.P., D.A. Peterson, and P.J. Biggs, *SolexaQA: At-a-glance quality*
assessment of Illumina second-generation sequencing data. BMC
Bioinformatics, 2010. **11**: p. 485.

83.   Kim, D., B. Langmead, and S.L. Salzberg, *HISAT: a fast spliced aligner with*
low memory requirements. Nat Methods, 2015. **12**(4): p. 357-60.

84.   Trapnell, C., et al., *Differential gene and transcript expression analysis of RNA-*
seq experiments with TopHat and Cufflinks. Nat Protoc, 2012. **7**(3): p. 562-78.

85.   Kumar, L. and E.F. M, *Mfuzz: a software package for soft clustering of*
microarray data. Bioinformation, 2007. **2**(1): p. 5-7.

86.   Lu, Y., Z. Pang, and J. Xia, *Comprehensive investigation of pathway enrichment*
methods for functional interpretation of LC–MS global metabolomics data.
Briefings in Bioinformatics, 2022. **24**(1).

87.   Su, G.-H., et al., *Radiogenomic-based multiomic analysis reveals imaging*
intratumor heterogeneity phenotypes and therapeutic targets. Science Advances,
2023. **9**(40): p. eadf0837.

88.   Lu, Y., Z. Pang, and J. Xia, *Comprehensive investigation of pathway enrichment*
methods for functional interpretation of LC–MS global metabolomics data.
Briefings in Bioinformatics, 2023. **24**(1): p. bbac553.

Author Contributions

J.J. conceived and designed this study; J.J. and W.W. wrote and revised this manuscript; S.Y. and F.L. provided necessary resources to this study; J.J. performed the metabolomics, RNA-seq analysis, and the western blot assay; N.G. and X.G performed the metabolomics analysis; Y.W. and G.W. performed the animal behavior tests; J. Q., and K. G. performed the sham/castration procedure; J.J., Y.W., G.W., J. Q., K. G., X.G and Y.L. did the mouse colony and tissue samples collection.

Acknowledgement

We acknowledge Ms. Jiahui Liu for her generous help for sample collection and Dr. Xinhui Liu for his kind help for assessing renal injury scores. This work was supported by Shenzhen Science and Technology Program (Grant No.RCBS20210706092341003) to J.J.

Declaration of interests

The authors declare that none of them has competing interests.

FIGURES

Figure 1. Overall study design as well as phenotypic and behavior tests of castration-mediated longevity. (**A**) Graphic illustration of the overall experimental design. (**B**) Body weight of casM, shamM and shamF mice from 8 to 60 weeks (n = 13, 12, 13 for casM, shamM, and shamF, respectively; unpaired students' t-test,***$p <$ 0.001, ns: not significant) (**C**) More gray hairs (see red arrows) were observed in shamM than casM at 12-13 months old; (**D-F**) Analysis of BM tests results: duration time that mice have spent to find the correct target escape hole (**D**), exploration of

898   wrong holes numbers (**E**), and immobile time during the test (**F**) among casM (n=6),

899   shamM (n= 13) and shamF (n = 10) by One-Way ANOVA, $^{*}p < 0.05$, $^{**}p < 0.01$, ns:

900   not significant.

Figure 2. Alterations of the steroid hormones secretion induced by castration. (**A**) Hierarchical cluster heatmap of z-score standardized expression levels of 10 detectable steroid hormones among shamM, casM and shamF; (**B-E**) Comparisons of the quantitative levels of testosterone (**B**), dihydroestosterone (**C**), 17-hydroxyprogesterone (**D**) and androstenedione (**E**) between shamM and casM (unpaired students' t-test, *p*-values were shown on top); (**F**) List of compounds highly correlated with TS (Spearman's |r| ≥ 0.8, *p* ≤ 0.001).

Figure 3. Characterizations of the 12w serum metabolomics. (**A**) PCA clustering of the serum untargeted metabolomics among casM, shamM and shamF; (**B**) Categories of the 1495 12w serum metabolites with high quality MS^2 annotations; (**C**) Representative SAMs between shamM and shamF that related with aging and healthy aging (unpaired students' t-test, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$); (**D**) Differential abundance (DA) scores were calculated for the significant enrichment terms between shamM and shamF ($*p < 0.05$, $**p < 0.01$, $***p < 0.001$).

Figure 4. Identification of castration-specific metabolomic features in 12w mice.
(**A**) Venny diagram of the serum SAMs identified from "shamF *vs* ShamM", "shamF *vs* casM" and "casM *vs* ShamM"; (**B**) Hierarchical clustering tree of different mouse groups using levels of compounds as input; (**C**) Representative food derived SAMs between casM and shamM (unpaired students' t-test, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$); (D) Barplot of the KEGG enrichment terms using SAMs of "casM *vs* ShamM", percentage of hit compounds to total SAMs was calculated for each term; (**E**)

Expression patterns of merged SAMs among casM, shamM and shamF analyzed using
mfuzzing clustering. Clusters 1 and 8 distinctively represent metabolites that are,
respectively, highly and lowly detected specifically in casM. (**F**) Significantly enriched
KEGG terms of metabolites highly expressed metabolites in casM (cluster 1 in **E**)
calculated by topology-based pathway enrichment with topology impact shown in x-
axis; (**G-H**) Comparisons of taurine, taurocholic acid and taurodeoxycholic acid among
12w (**G**) and 18m (**H**) shamF, sham and casM (One-Way ANOVA, *$p < 0.05$, **$p <$
0.01, ns: not significant).

Figure 5. Co-expression analysis across serum, livers and kidneys identifies potential taurine-related regulators. (**A**) Venn diagram of detected compounds among serum, livers and kidneys across all the mouse groups at 12w and 18m, 121 commonly detected compounds (CDBs) were identified; (**B**) Co-expression analysis of the 121 CDBs revealed four major consensus clusters; (**C**) Correlation circos plot of the cluster 4 compounds in **B**; (**D**) Topology-based pathway enrichment analysis of the cluster 4 compounds in **B**, x-axis is the topology enrichment impact of hinted

compounds; (**E**) Hierarchical cluster heatmap of z-score standardized expression levels

of 51 taurine related genes in 12w and 18m kidneys and livers among shamM, shamF

and casM mice. Genes selectively reduced in 18m tissues are highlighted in red; (**F**)

Trends of taurine levels among shamM, shamF and casM across liver and kidney tissues

in 12w and 18m mice; (**F**) Mfuzzing clustering of the 51 taurine co-expressed genes

across liver and kidney tissues in 12w and 18m mice.

Figure 6. Joint metabolomic-transcriptomic analysis reveals a stronger capacity of

defensing lipid peroxidation in female and castrated male. (**A-B**) The numbers of

DEGs among livers and kidneys tissues of shamM, shamF and casM mice collected at

12w (**A**) and 18m (**B**); (**C**) Hierarchical clustering of 12w and 18m liver and kidney

samples demonstrating casM's closer alignment with shamF than sham; (**D, E**)

Representative significantly enriched KEGG pathways in 18m liver samples from

'casM vs ShamM,' (**D**) and 'shamF vs ShamM' (**E**), identified by joint transcriptomic-

metabolomic analysis using DEGs and SAMs. Pathways marked in orange indicate common terms between these two comparisons. (**F**) Immunoblotting of MDA in 18m livers; (**G**) H & E staining of livers (upper panel) and PAS staining of kidneys (lower panel) in 12w mice (4 week post castration/sham procedure) with cisplatin treatment, blue arrows indicate representative injury spots in livers; (**H**) Bar plots of renal injury scores in three groups of mice by One-Way ANOVA test (**$p < 0.01$, ns: not significant).