Effects of local brain temperature on somatosensory evoked potentials in rats

Mizuho Gotoh[a,b,c], Shinnosuke Dezawa[a,b,d], Ichiro Takashima[a,b,e], Shinya Yamamoto[a,b,*]

[a] Human Informatics and Interaction Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba, Japan

[b] Graduate School of Comprehensive Human Sciences, University of Tsukuba, Tsukuba Japan

[c] Department of Rehabilitation for Brain Functions, Research Institute of National Rehabilitation Center for Persons with Disabilities, Tokorozawa, Japan

[d] Faculty of Medical and Health Sciences, Tsukuba International University, Tsuchiura, Japan

[e] Department of Informatics and Electronics, Daiichi Institute of Technology, Tokyo, Japan

* Corresponding author: Shinya Yamamoto, Human Informatics and Interaction Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), 1-1-1, Umezono, Tsukuba 305-8568, Japan

Email address: yamamoto-s@aist.go.jp

Abstract

Although the focal brain cooling technique is widely used to examine brain function, the effects of variations in cortical temperature on sensory information processing and neural mechanisms remain underexplored. To elucidate the mechanisms of temperature modulation in somatosensory processing, this study aimed to examine how P1 and N1 deflections of somatosensory evoked potentials (SEPs) depend on cortical temperature and how excitatory and inhibitory inputs contribute to this temperature dependency. SEPs were generated through electrical stimulation of the contralateral forepaw in anesthetized rats. The SEPs were recorded while cortical temperatures were altered between 18–37°C: without any antagonists; with gamma-aminobutyric acid type A ($GABA_A$) receptor antagonist, gabazine; with aminomethylphosphonic acid (AMPA) receptor antagonist (NBQX), or N-Methyl-D-aspartic acid (NMDA) receptor antagonist ([R]-CPP). The effects of different gabazine concentrations (0, 1, and 10 µM) were examined in the 35–38°C range. The P1/N1 amplitudes and their peak-to-peak differences plotted against cortical temperature showed an inverted U relationship with their maximum at approximately 27.5°C when no antagonists were administered. The negative correlation between these amplitudes and temperatures ≥27.5°C plateaued after gabazine administration, which occurred progressively as the gabazine concentration increased. In contrast, the correlation remained negative after the administration of NBQX and (R)-CPP. GABAergic inhibitory inputs contribute to the negative correlation between SEP amplitude and cortical temperature around the physiological cortical temperature by suppressing SEPs to a greater extent at higher temperatures.

Keywords

1. Introduction[1]

The focal brain cooling (FBC) technique has been widely used in neuroscience to inactivate various brain areas (Bauer and Fuster, 1976; Fuster and Bauer, 1974; Fuster et al., 1981; Hupé et al., 1998; Lomber and Malhotra, 2008; Moseley et al., 1972; Payne et al., 1996; Takei et al., 2021; Trendelenburg, 1911). The FBC technique was first developed more than 100 years ago, whereby motor function was inactivated by cooling the motor cortex (Trendelenburg, 1911). Since then, this technique has contributed significantly to the exploration of various neural mechanisms. This technique has also been applied to treat epilepsy in humans (Karkar et al., 2002; Nomura et al., 2017; Sartorius and Berger, 1998). Although previous neuroscience research has developed and utilized a variety of other inactivation techniques including the physical removal (Deuel and Mishkin, 1977; Johnson et al., 2015; Minamimoto et al., 2010), pharmacological manipulation (Hikosaka et al., 1985; Hikosaka and Wurtz, 1985a; Hikosaka and Wurtz, 1985b; Newsome et al., 1985; Schwarcz et al., 1979), light-induced lesion (Nakata et al., 2018; Nakata et al., 2021; Suzuki et al., 2012), and optogenetic techniques (Chow et al., 2010; Han and Boyden, 2007; Han et al., 2011), the FBC technique has an advantage in that it can inactivate wide brain areas simultaneously in a reversible manner.

Somatosensory evoked potentials (SEPs) are field potentials generated in the somatosensory cortex in response to stimulation of the skin or peripheral nerves (e.g., the median and radial nerves). SEPs are useful in examining neural circuits and their plasticity (All et al., 2021; Dezawa et al., 2021; Franceschini et al., 2008). In addition, SEPs are used clinically to evaluate the somatosensory pathway in patients with neurological disorders and monitor the condition of patients during surgery (Carter and Butt, 2005; Cruccu et al., 2008; Florence et al., 2004; MacDonald et al., 2019). The SEPs are composed of alternating positive and negative deflections. The SEPs recorded from the primary somatosensory cortex in rats include two components: the first positive deflection after the stimulation is termed as P1, and the subsequent negative deflection is termed as N1 (Bruyns-Haylett et al., 2017; Di and Barth, 1991; Franceschini et al., 2008; Freeman and Sohmer, 1996; Jellema et al., 2004). Each deflection corresponds to a different somatosensory process in each phase.

---

[1] FBC, focal brain cooling; SEPs, somatosensory evoked potentials; P1, positive deflection; N1, negative deflection; AMPA, aminomethylphosphonic acid; NMDA, *N*-Methyl-D-aspartic acid; GABA, gamma-aminobutyric acid; GABA$_A$, gamma-aminobutyric acid type A

The FBC technique has been used to examine sensory information processing. For example, unilateral cooling of the superior colliculus or the temporo-occipito-parietal junction induces visual hemineglect in cats (Payne et al., 1996). Bilateral cooling of the posterior part of the non-primary auditory cortex impairs sound localization, whereas cooling its anterior part impairs sound identification in cats (Lomber and Malhotra, 2008). In these studies, the temperature of the sensory cortices was sufficiently decreased (≤20°C). Regarding somatosensory processing, previous studies have measured SEPs while cooling the somatosensory cortex. For example, Schwerdtfeger et al. (Schwerdtfeger et al., 1999) showed that the peak-to-peak amplitudes between P1 and N1 increased while cooling the somatosensory cortex in rats. Bindman et al. (Bindman et al., 1963) also reported that cooling the cortical surface over the somatosensory cortex increased N1 amplitudes in rats. Although these findings seemingly contradict the assumption that cooling the cortex attenuates SEPs, the estimated decrease in cortical temperature in these SEP studies was smaller than that in FBC studies (approximately 10°C from the physiological temperature at a depth of approximately 1 mm). Hence, the extent to which broader fluctuations in cortical temperature influence SEP amplitudes remains unknown.

A recent study proposed that the net effects of brain temperature on neural activity involve a trade-off between excitatory and inhibitory inputs. Gotoh et al. (Gotoh et al., 2020) recorded evoked potentials in the frontal cortex that were triggered by stimulation of the mesocortical pathway. Their results showed that the increase in amplitude by focal cortical cooling was caused by a decrease in the contribution of inhibitory inputs compared to that of excitatory inputs. However, the contributions of excitatory and inhibitory inputs to the temperature dependence of sensory information processing are underexplored. In particular, elucidating their effects on each component of the SEPs (i.e., P1 and N1) would contribute significantly to a better understanding of the role of temperature modulation in sensory information processing. To address these issues, the present study examined: (1) how the amplitudes of P1 and N1 deflections of SEPs (as well as their peak-to-peak voltages) depend on the temperature of the somatosensory cortex, and (2) how their temperature dependencies are affected by glutamate (aminomethylphosphonic acid [AMPA] and *N*-Methyl-D-aspartic acid [NMDA]) and gamma-aminobutyric acid (GABA) receptor antagonists.

2. Results

2.1 Experiment 1: Effects of cortical temperature on SEPs

We recorded SEPs in the primary somatosensory cortex that were elicited by electrical stimulation of the contralateral forepaw while altering the local cortical temperature (Fig. 1A; n = 24 rats). A custom-made thermal control chamber was placed above the somatosensory cortex, which contained 100 µL of liquid (saline or an antagonist solution). Cortical temperature was controlled by circulating temperature-controlled water through a stainless-steel tube embedded in the chamber. The cortical temperature at a depth of 1 mm was linearly correlated with the temperature of the water circulating in the tube ($R^2 = 0.99$; Fig. 1B).

The average waveforms of the SEPs from 24 rats at various temperatures are shown in Fig. 2A. The SEP waveforms changed depending on cortical temperature. To evaluate the effects of cortical temperature on the SEPs, we focused on the initial positive deflection (P1) and subsequent negative deflection (N1). We defined the individual peak amplitudes for positive (V_p) and negative peaks (V_n), as well as the peak-to-peak amplitudes (V_{p-n}) between these two peaks and the individual peak latencies (L_p and L_n), as shown in Fig. 2B (see also *4.8 Data analysis*).

The relationship between temperature and V_{p-n} exhibited a non-monotonic inverted-U shape (Fig. 2C). The V_{p-n} was negatively correlated with the cortical temperature at $\geq$27.5°C ($R = -0.70$, $P = 1.9 \times 10^{-15}$; Supplementary Table 1), while it was positively correlated at $\leq$27.5°C ($R = 0.31$, $P = 0.0019$; Supplementary Table 1). A two-way ANOVA (temperature by animal) revealed significant main effects for temperature (F (6,138) = 84.2, $P = 1.2 \times 10^{-43}$; Supplementary Table 2).

The V_p was negatively correlated with the cortical temperatures at $\geq$27.5°C ($R = -0.65$, $P = 9.6 \times 10^{-13}$; Supplementary Table 1), but was not significantly correlated with the cortical temperatures at $\leq$27.5°C ($R = 0.15$, $P = 0.15$; Supplementary Table 1, Fig. 2D). The V_n was negatively correlated with the cortical temperatures at $\geq$27.5°C ($R = -0.67$, $P = 5.9 \times 10^{-14}$; Supplementary Table 1) and was positively correlated with the cortical temperatures at $\leq$27.5°C ($R = 0.36$, $P = 3.4 \times 10^{-4}$; Supplementary Table 1, Fig. 2E). Two-way ANOVAs (temperature by animal) for positive and negative peaks revealed significant main effects for temperature (V_p: F (6,138) = 40.9, $P = 2.4 \times 10^{-28}$, V_n: F (6,138) = 83.0, $P = 2.5 \times 10^{-43}$; Supplementary Table 2).

The latencies of the SEPs also changed depending on cortical temperature. Peak latency for P1 (L_p) and N1 (L_n) was defined as the time between the onset of stimulation and the positive and negative peaks, respectively (Fig. 2B, see *4.8 Data analysis*). Both L_p and L_n decreased monotonically as cortical temperature increased (Fig. 2F–G). A two-way ANOVA (temperature by animal) for negative and positive peaks revealed significant main effects for temperature (L_p: $F(6,138) = 229.6$, $P = 3.2 \times 10^{-69}$, L_n: $F(6,138) = 343.2$, $P = 2.5 \times 10^{-80}$; Supplementary Table 3).

2.2 Experiment 2: Effects of gamma-aminobutyric acid type A (GABAₐ) receptor antagonist

To examine how the contribution of GABAergic inputs to the temperature dependency of SEPs, we administered the GABAₐ receptor antagonist gabazine (10 µM, 100 µL) to the chamber in eight of 24 rats after Experiment 1 (Fig. 3A). Gabazine administration increased $V_{p\text{-}n}$. In particular, an increase in $V_{p\text{-}n}$ at higher temperatures resulted in the disappearance of the negative correlation at ≥27.5°C observed in Experiment 1 (Fig. 3B; $R = –0.11$, $P = 0.54$; Supplementary Table 4). A three-way ANOVA for $V_{p\text{-}n}$ (antagonist by temperature by animal) revealed significant main effects for antagonist ($F(1,42) = 3502.1$, $P = 4.4 \times 10^{-42}$) and temperature ($F(6,42) = 37.2$, $P = 2.8 \times 10^{-15}$), and significant interaction between antagonist and temperature ($F(6,42) = 35.9$, $P = 5.2 \times 10^{-15}$; Supplementary Table 5). These results suggest that GABAergic inhibitory inputs contribute to the negative correlation between SEP amplitude and cortical temperature around the physiological cortical temperature.

Gabazine administration increased individual peak amplitudes (V_p and V_n). The correlations between peak amplitudes (V_p and V_n) and temperature at ≥27.5°C were not significantly different from 0 in Experiment 2 (Fig. 3C–D, V_p: $R = –0.23$, $P = 0.21$, V_n: $R = 0.010$, $P = 0.96$; Supplementary Table 4). A three-way ANOVA for V_p and V_n (antagonist by temperature by animal) revealed significant main effects for the antagonist (V_p: $F(1,42) = 166.8$, $P = 3.2 \times 10^{-16}$, V_n: $F(1,42) = 4155.4$, $P = 1.2 \times 10^{-43}$) and temperature (V_p: $F(6,42) = 25.0$, $P = 2.2 \times 10^{-12}$, V_n: $F(6,42) = 35.5$, $P = 6.5 \times 10^{-15}$) and significant interaction between antagonist and temperature (V_p: $F(6,42) = 6.9$, $P = 3.8 \times 10^{-5}$, V_n: $F(6,42) = 36.8$, $P = 3.5 \times 10^{-15}$) (Supplementary Table 5).

To evaluate the contribution of GABAergic inputs, we calculated the difference and ratio between the amplitudes before and after gabazine administration at each cortical temperature level. Both the difference and the ratio of the amplitudes increased at higher temperatures for $V_{p\text{-}n}$, V_p, and V_n (Fig. S1). The correlation between *difference*($V_{p\text{-}n}$) and temperature, that between

difference(V_p) and temperature, and that between *difference*(V_n) and temperature were significantly positive (V_{p-n}: $R = 0.73$, $P = 1.7 \times 10^{-10}$, V_p: $R = 0.41$, $P = 1.7 \times 10^{-3}$, V_n: $R = 0.70$, $P = 1.4 \times 10^{-9}$) (Supplementary Table 6). The correlation between *ratio*(V_{p-n}) and temperature, that between *ratio*(V_p) and temperature, and that between *ratio*(V_n) and temperature were significantly positive (V_{p-n}: $R = 0.57$, $P = 5.2 \times 10^{-6}$, V_p: $R = 0.51$, $P = 5.0 \times 10^{-5}$, V_n: $R = 0.55$, $P = 1.4 \times 10^{-5}$) (Supplementary Table 7). These results suggest that the effect of gabazine (i.e., an increase in V_{p-n}, V_p, and V_n) was greater at higher temperatures.

Gabazine administration influenced the latency. In Experiment 2, both L_p and L_n exhibited values higher than those recorded in Experiment 1; nonetheless, they decreased monotonically as the temperature rose (Fig. 3E–F). A three-way ANOVA (antagonist by temperature by animal) revealed significant main effects for the antagonist (L_p: $F_{(1,42)} = 466.2$, $P = 2.3 \times 10^{-24}$, L_n: $F_{(1,42)} = 1159.4$, $P = 3.2 \times 10^{-32}$) and temperature (L_p: $F_{(6,42)} = 199.3$, $P = 3.3 \times 10^{-29}$, L_n: $F_{(6,42)} = 211.2$, $P = 1.0 \times 10^{-29}$) and significant interaction between antagonist and temperature (L_p: $F_{(6,42)} = 13.3$, $P = 2.2 \times 10^{-8}$, L_n: $F_{(6,42)} = 34.1$, $P = 1.3 \times 10^{-14}$; Supplementary Table 8). These results suggest that GABAergic inputs increase the peak latencies of SEPs.

2.3 Experiment 3: Effects of different concentrations of GABA$_A$ receptor antagonist around the physiological temperature

To evaluate the effects of concentration of gabazine around the physiological temperature, we recorded the SEPs at 35, 36.5, and 38°C with the chamber filled with saline, 1 µM gabazine, and 10 µM gabazine (Fig. 4). As the gabazine concentration increased, V_{p-n}, V_p, and V_n increased, and the negative correlation between the amplitude and temperature with saline plateaued (Fig. 4A–C). Three-way ANOVAs for V_{p-n}, V_p, and V_n (antagonist by temperature by animal) revealed significant main effects for the antagonist (V_{p-n}: $F_{(2,28)} = 3286.2$, $P = 6.1 \times 10^{-3}$, V_p: $F_{(2, 28)} = 534.9$, $P = 4.9 \times 10^{-23}$, V_n: $F_{(2,28)} = 3430.9$, $P = 3.4 \times 10^{-34}$) and temperature (V_{p-n}: $F_{(2, 28)} = 32.3$, $P = 5.3 \times 10^{-8}$, V_p: $F_{(2, 28)} = 20.4$, $P = 3.5 \times 10^{-6}$, V_n: $F_{(2, 28)} = 27.3$, $P = 2.7 \times 10^{-7}$), and significant interaction between antagonist and temperature (V_{p-n}: $F_{(4, 28)} = 8.4$, $P = 1.4 \times 10^{-4}$, V_p: $F_{(4, 28)} = 4.0$, $P = 1.0 \times 10^{-2}$, V_n: $F_{(4, 28)} = 8.0$, $P = 2.0 \times 10^{-4}$; Supplementary Table 9).

To evaluate the effects of cortical temperature around the physiologically normal temperature on SEPs, we fitted V_{p-n}, V_p, and V_n at each concentration using a linear model of the temperatures (Supplementary Table 10; see also *4.8 Data Analysis*). The constant terms (the mean amplitudes

at 36.5°C) of the fitted linear models increased at higher concentrations (Fig. 4F, top). The two-way ANOVA for the constant terms of V_{p-n}, V_p, and V_n (antagonist by animal) revealed significant main effects of antagonist (V_{p-n}: $F_{(2,14)} = 99.3$, $P = 5.4 \times 10^{-9}$, V_p: $F_{(2,14)} = 43.8$, $P = 9.5 \times 10^{-7}$, V_n: $F_{(2,14)} = 95.9$, $P = 6.8 \times 10^{-9}$; Supplementary Table 11). The slopes also significantly increased from -0.28 mV/°C to -0.0065 mV/°C in V_{p-n}, -0.041 mV/°C to -0.0084 mV/°C in V_p, -0.24 mV/°C to 0.0019 mV/°C in V_n (Fig. 4F, bottom). Two-way ANOVA of the slopes (antagonist by animal) revealed significant main effects of the antagonist (V_{p-n}: $F_{(2,14)} = 10.4$, $P = 1.7 \times 10^{-3}$, V_p: $F_{(2,14)} = 4.9$, $P = 0.024$, V_n: $F_{(2,14)} = 10.3$, $P = 1.8 \times 10^{-3}$; Supplementary Table 12). The slopes in the saline and 1-μM gabazine conditions were significantly smaller than 0 (V_{p-n}: $P = 6.6 \times 10^{-4}$ and $P = 0.015$; V_p: $P = 2.0 \times 10^{-3}$ and $P = 7.7 \times 10^{-3}$; V_n: $P = 6.9 \times 10^{-4}$ and $P = 0.025$, t-test) but not in the 10-μM gabazine condition (V_{p-n}: $P = 0.83$; V_p: $P = 0.39$; V_n: $P = 0.94$, t-test; Supplementary Table 10). The results showed that a higher concentration of gabazine had a greater effect on the peak amplitudes.

The effects of gabazine on the latencies (L_p and L_n) were also greater at higher concentrations (Fig. 4D–E). A three-way ANOVA (antagonist by temperature by animal) revealed significant main effects for antagonist (L_p: $F_{(2,28)} = 168.6$, $P = 2.4 \times 10^{-16}$, L_n: $F_{(2,28)} = 1564.3$, $P = 1.9 \times 10^{-29}$) and temperature (L_p: $F_{(2, 28)} = 12.1$, $P = 1.6 \times 10^{-4}$, L_n: $F_{(2, 28)} = 30.8$, $P = 8.5 \times 10^{-8}$). The interaction between antagonist and temperature was significant for L_n ($F_{(4, 28)} = 5.1$, $P = 3.3 \times 10^{-3}$), but not for L_p ($F_{(4, 28)} = 1.2$, $P = 0.35$; Supplementary Table 13).

2.4 Experiment 4: Effects of AMPA receptor antagonist

To examine the effects of glutamatergic inputs to AMPA receptors on SEPs, we administered the AMPA receptor antagonist NBQX (10 μM, 100 μL) to the chamber in eight of 24 rats after Experiment 1 (Fig. 5A). The administration of NBQX consistently reduced V_{p-n} across all tested temperatures (Fig. 5B). The correlation between V_{p-n} and temperature at ≥27.5°C remained negative ($R = -0.69$, $P = 1.3 \times 10^{-5}$; Supplementary Table 14). A three-way ANOVA (antagonist by temperature by animal) revealed significant main effects for the antagonist ($F_{(1,42)} = 224.0$, $P = 2.0 \times 10^{-18}$) and temperature ($F_{(6,42)} = 49.8$, $P = 1.6 \times 10^{-17}$). The interaction between the antagonist and temperature was not significant ($F_{(6,42)} = 1.0$, $P = 0.42$, Supplementary Table 15). Both the difference and ratio of V_{p-n} before and after the administration of NBQX were almost constant across temperatures (Fig. S2A). The correlation between the *difference*(V_{p-n}) and temperature, as well as the *ratio*(V_{p-n}) and temperature, did not significantly deviate from 0 (*difference*[V_{p-n}]: $R = 0.21$, $P = 0.12$, *ratio*[V_{p-n}]: $R = 0.057$, $P = 0.68$; Supplementary Tables 16–

17). This implies that the impact of NBQX (i.e., the decrease in $V_{p\text{-}n}$) remained constant across temperatures.

Administration of NBQX decreased V_p with the correlation at $\geq 27.5°C$ remained negative (Fig. 5C; $R = -0.63$, $P = 9.5 \times 10^{-5}$; Supplementary Table 14). A three-way ANOVA (antagonist by temperature by animal) revealed significant main effects for the antagonist ($F(1,42) = 275.3$, $P = 4.7 \times 10^{-20}$) and temperature ($F(6,42) = 67.6$, $P = 5.5 \times 10^{-20}$) and the interaction between antagonist and temperature ($F(6,42) = 12.3$, $P = 5.9 \times 10^{-8}$, Supplementary Table 15). The difference and ratio of V_p before and after NBQX administration—referred to as $difference(V_p)$ and $ratio(V_p)$—increased at elevated temperatures (Fig. S2B). The correlation between $difference(V_p)$ and temperature and that between $ratio(V_p)$ and temperature were significantly positive ($difference(V_p)$: $R = 0.55$, $P = 1.1 \times 10^{-5}$, $ratio(V_p)$: $R = 0.52$, $P = 3.8 \times 10^{-5}$; Supplementary Tables 16–17), suggesting that the effect of NBQX (i.e., the decrease of V_p) was larger at the lower temperatures.

Administration of NBQX also decreased V_n with the correlation at $\geq 27.5°C$ remaining negative (Fig. 5D; $R = -0.63$, $P = 9.7 \times 10^{-5}$; Supplementary Table 14). A three-way ANOVA (antagonist by temperature by animal) revealed significant main effects for the antagonist ($F(1,42) = 158.6$, $P = 7.5 \times 10^{-16}$) and temperature ($F(6,42) = 34.6$, $P = 9.8 \times 10^{-15}$). The interaction between the antagonist and temperature was not significant ($F(6,42) = 0.4$, $P = 0.87$, Supplementary Table 15). Both the difference and ratio of V_n before and after the administration of NBQX were almost constant across temperatures (Fig. S2C). The correlation between $difference(V_n)$ and temperature and that between $ratio(V_n)$ and temperature were not significantly different from 0 ($difference(V_n)$: $R = 0.045$, $P = 0.74$, $ratio(V_n)$: $R = -0.14$, $P = 0.31$; Supplementary Tables 16–17), supporting that the effect of NBQX (i.e., the decrease of V_n) was constant across temperatures.

Administering NBQX caused a slight but significant increase in the peak latencies (Fig. 5E–F). L_p remained negatively correlated with the cortical temperature after administering NBQX but was 0.48 ms longer on average. A three-way ANOVA (antagonist by temperature by animal) revealed significant main effects for the antagonist ($F(1,42) = 35.5$, $P = 4.5 \times 10^{-7}$), for temperature ($F(6,42) = 280.1$, $P = 3.3 \times 10^{-32}$), and significant interaction between antagonist and temperature ($F(6,42) = 3.0$, $P = 0.015$). Administering NBQX also caused a slight but significant increase in L_n by 0.85 ms on average while remaining negatively correlated with the cortical temperature. A three-way ANOVA (antagonist by temperature by animal) revealed significant main effects for the antagonist ($F(1,42) = 21.7$, $P = 3.2 \times 10^{-5}$) and temperature (F

$(6,42) = 406.0$, $P = 1.6 \times 10^{-35}$). The interaction between the antagonist and temperature was not significant ($F (6,42) = 0.1$, $P = 0.999$; Supplementary Table 18).

2.5 Experiment 5: Effects of NMDA receptor antagonist

To examine the effects of glutamatergic inputs to NMDA receptors on SEPs, we administered the NMDA receptor antagonist (R)-CPP (10 µM, 100 µL) to the chamber in eight of 24 rats after Experiment 1 (Fig. 6A). V_{p-n} decreased after administration of (R)-CPP, particularly at higher temperatures (Fig. 6B). The correlation between V_{p-n} and temperature at $\geq$27.5°C remained negative ($R = -0.86$, $P = 4.6 \times 10^{-10}$; Supplementary Table 19). A three-way ANOVA (antagonist by temperature by animal) revealed significant main effects for the antagonist (F $(1,42) = 33.3$, $P = 8.5 \times 10^{-7}$) and temperature ($F (6,42) = 135.2$, $P = 7.9 \times 10^{-26}$) and the interaction between antagonist and temperature ($F (6,42) = 5.6$, $P = 2.4 \times 10^{-4}$, Supplementary Table 20). Both the difference and ratio of V_{p-n} before and after the administration of (R)-CPP (that is, the *difference*(V_{p-n}) and *ratio*(V_{p-n})) were smaller at higher temperatures (Fig. S3A). The correlation between *difference*(V_{p-n}) and temperature and that between *ratio*(V_{p-n}) and temperature were significantly negative (*difference*(V_{p-n}): $R = -0.52$, $P = 3.8 \times 10^{-5}$, *ratio*(V_{p-n}): $R = -0.64$, $P = 1.0 \times 10^{-7}$; Supplementary Tables 21–22), suggesting that the effect of (R)-CPP (i.e., the decrease of V_{p-n}) was larger at the higher temperatures.

Unlike V_{p-n}, the administration of (R)-CPP increased V_p (Fig. 6C). The correlation between V_p and temperature at $\geq$27.5°C remained negative ($R = -0.74$, $P = 1.2 \times 10^{-6}$; Supplementary Table 19). A three-way ANOVA (antagonist by temperature by animal) revealed significant main effects for the antagonist ($F (1,42) = 120.2$, $P = 6.7 \times 10^{-14}$) and temperature ($F (6,42) = 111.4$, $P = 3.6 \times 10^{-24}$). The interaction between the antagonist and temperature was not significant (F $(6,42) = 2.0$, $P = 0.094$, Supplementary Table 20). Both the difference and ratio of V_p before and after administering (R)-CPP (i.e., the *difference*(V_p) and *ratio*(V_p)) were almost constant across temperatures (Fig. S3B). The correlation between *difference*(V_p) and temperature, as well as *ratio*(V_p) and temperature, did not significantly deviate from 0 (*difference*(V_p): $R = -0.21$, $P = 0.12$, *ratio*(V_{p-n}): $R = -5.0 \times 10^{-3}$, $P = 0.97$; Supplementary Tables 21–22), implying that the effect of (R)-CPP (i.e., the decrease of V_p) was constant across temperatures.

The administration of (R)-CPP decreased V_n (Fig. 6D). The correlation between V_n and temperature at $\geq$27.5°C remained negative ($R = -0.86$, $P = 1.9 \times 10^{-10}$; Supplementary Table 19). A three-way ANOVA (antagonist by temperature by animal) revealed significant main effects for the antagonist ($F (1,42) = 114.6$, $P = 1.4 \times 10^{-13}$) and temperature ($F (6,42) = 107.1$, $P = 7.9 \times 10^{-24}$) and the interaction between antagonist and temperature ($F (6,42) = 6.5$, $P = 6.5 \times 10^{-5}$,

Supplementary Table 20). Both the difference and ratio of V_n before and after the administration of (R)-CPP (that is, the *difference*(V_n) and *ratio*(V_n)) were smaller at higher temperatures (Fig. S3C). The correlation between *difference*(V_n) and temperature and that between *ratio*(V_n) and temperature were significantly negative (*difference*(V_n): $R = -0.55$, $P = 1.1 \times 10^{-5}$, *ratio*(V_n): $R = -0.69$, $P = 3.5 \times 10^{-9}$; Supplementary Tables 21–22), suggesting that the effect of (R)-CPP (i.e., the decrease of V_n) was larger at higher temperatures.

The administration of (R)-CPP caused a slight but significant decrease in peak latencies (Fig. 6E–F). After the administration of (R)-CPP, L_p maintained a negative correlation with cortical temperature but was, on average, 0.41 ms shorter. A three-way ANOVA (antagonist by temperature by animal) revealed significant main effects for the antagonist ($F(1,42) = 31.8$, $P = 1.3 \times 10^{-6}$), temperature ($F(6,42) = 342.8$, $P = 5.2 \times 10^{-34}$). The interaction between the antagonist and temperature was not significant ($F(6,42) = 1.1$, $P = 0.39$). After the administration of (R)-CPP, L_n maintained a negative correlation with the cortical temperature but was, on average, 1.3 ms shorter. A three-way ANOVA (antagonist by temperature by animal) revealed significant main effects for the antagonist ($F(1,42) = 81.1$, $P = 2.4 \times 10^{-11}$) and temperature ($F(6,42) = 545.1$, $P = 3.6 \times 10^{-38}$), and significant interaction between antagonist and temperature ($F(6,42) = 3.2$, $P = 0.012$) (Supplementary Table 23).

3. Discussion

In the present study, we showed that the amplitudes of SEPs triggered by forepaw stimulation changed non-monotonically with the local temperature in the somatosensory cortex. For either V_{p-n}, V_n, or V_p, the relationship between the amplitude and the temperature showed an inverted U-curve, where the SEP amplitudes increased by cooling up to approximately 27.5°C and then decreased by further cooling. This temperature-dependent pattern was modulated by the administration of the GABA$_A$ receptor antagonist, gabazine. The negative correlation between these amplitudes and cortical temperature at ≥27.5°C disappeared and plateaued after gabazine administration. In addition, the negative correlation progressively disappeared as gabazine concentration increased. Notably, despite the increased SEP amplitude being associated with higher gabazine concentration, the negative correlation vanished. The effects of GABAergic inputs on SEP amplitudes, evaluated using the difference and the ratio before and after gabazine administration, were larger at higher temperatures. Note that the negative correlation at ≥27.5°C remained after the administration of either the AMPA receptor antagonist, NBQX, or the NMDA receptor antagonist, (R)-CPP. Taken together, these results suggest that GABAergic inhibitory inputs contribute to the negative correlation between SEP amplitude and cortical temperature around the physiological cortical temperature by inducing stronger suppression of SEPs at higher temperatures.

The event-related potentials recorded from the cortical surface reflect the postsynaptic activity of pyramidal neurons, whose apical dendrites are located vertically to the cortical surface (Jackson and Bolger, 2014). Excitatory inputs to the pyramidal neurons in the deep layers cause a positive potential at the surface electrode, whereas inhibitory inputs to the pyramidal neurons in the deep layers work in opposition. In contrast, excitatory inputs at the superficial layers cause negative potentials, whereas inhibitory inputs at the superficial layers work oppositely (Jackson and Bolger, 2014). Our data showed that the administration of NBQX decreased V_p and gabazine increased V_p, suggesting that excitatory inputs via AMPA receptors and inhibitory inputs via GABA$_A$ receptors in the deep layer contribute to the positive deflection of the SEP. In fact, the granular layer (and the infragranular layer) receive afferent inputs from the thalamus (Jackson and Bolger, 2014; Lund, 1973), and early sinks were observed in these layers in the current source density analysis of the SEPs (Jellema et al., 2004; Mitzdorf, 1985). Thus, thalamocortical inputs to the deep cortical layers may have caused the first positive deflection (P1) in our experiments.

In contrast to the AMPA receptor antagonist, the NMDA receptor antagonist, (R)-CPP, increased V_p. This suggests that pyramidal neurons receive glutamatergic inputs via NMDA receptors in superficial layers during the early phase of SEPs. In fact, it has been demonstrated that the apical dendritic tufts in layer I receive input from the thalamus (Rubio-Garrido et al., 2009). Glutamatergic inputs via NMDA receptor channels on thin dendrites generate spike-like responses called NMDA spikes (Larkum et al., 2009; Palmer, 2014), which support the initiation of calcium spikes (Larkum et al., 2009). Thus, the present data support the possibility that glutamatergic inputs to apical dendritic tufts via NMDA receptors in layer I may contribute to the generation of the initial positive deflection (P1).

V_n increased upon gabazine administration and decreased following the administration of NBQX or (R)-CPP administration. These results suggest that excitatory and inhibitory inputs to the pyramidal neurons in the superficial layers cause negative deflection (N1). The superficial layer (i.e., the supragranular layer) receives input from the granular layer within the same cortical area (Bannister, 2005; Rockland and Pandya, 1979). In addition, the supragranular layer receives feedback input from the infragranular layer of higher cortical areas (Bannister, 2005; Rockland and Pandya, 1979). These feed-forward and feedback inputs to the superficial cortical layers can generate subsequent negative deflection (N1), either directly or indirectly via excitatory and inhibitory interneurons.

The influence of the antagonists on P1/N1 deflections varied with temperature. To evaluate the contribution of excitatory and inhibitory inputs, we calculated the difference and ratio of the amplitudes before and after antagonist administration. The effects of gabazine on V_p, V_n, and V_{p-n} (i.e., the increase in amplitude) were greater at higher temperatures. The impact of NBQX on V_p (i.e., the decrease in amplitude) were smaller at higher temperatures, whereas the effects of R)-CPP on V_p (i.e., the decrease in amplitude) did not show significant differences between different temperatures. The impact of NBQX on V_n and V_{p-n} did not show significant differences among the temperatures, whereas those of (R)-CPP on V_n and V_{p-n} were greater at higher temperatures than at lower temperatures. In the present study, either V_p, V_n, or V_{p-n} showed a negative correlation with the cortical temperature at $\geq 27.5°C$ without antagonists. Thus, the present data suggest that the larger/smaller contribution of GABAergic/glutamatergic inputs at higher temperatures leads to a negative correlation for V_p. Additionally, a heightened contribution of GABAergic inputs at higher temperatures results in negative correlations for both V_n and V_{p-n}.

The temperature dependency observed in the present study was quantitatively different from that observed in the frontal cortex in a previous study (Gotoh et al., 2020). The temperature at which the amplitude was maximum was approximately 17°C in the frontal cortex. In contrast, the temperature of the maximum amplitude was as high as 27.5°C in the somatosensory cortex. One possible reason for this difference is the impact of inhibitory inputs that contribute to the evoked potentials. Although the proportion of rat GABAergic neurons is reported to be approximately 15% in both the frontal and somatosensory cortices and the total number of neurons is even smaller in the frontal cortex than in the somatosensory cortex (Beaulieu, 1993), data from autoradiographic studies indicate that the densities of binding sites of [^{3}H]GABA or [^{3}H]muscimol to $GABA_A$ receptors are higher in the frontal cortex than in the somatosensory cortex (Bowery et al., 1987; Que et al., 1999). Therefore, the contribution of GABAergic inhibitory inputs may be greater in the frontal cortex than in the somatosensory cortex. In addition, the temperature of maximum amplitude shifted toward a higher temperature than 17°C when the concentration of gabazine increased in the frontal cortex (Gotoh et al., 2020). Therefore, we propose that the extent of the contribution of GABAergic inhibitory inputs determines the temperature-dependent pattern in the cortex and that the increased influence of inhibitory inputs lowers the temperature of the maximum amplitude in the inverted U-curve.

In the present study, focal cortical cooling also changed the SEP peak latency. Cooling the somatosensory cortex from 38°C to 17°C monotonically increased the P1 and N1 peak latencies. One possible reason for this delay is that cooling slows molecular processes. The slowing down of these processes can result in delays in signal conduction and synaptic transmission (Asztely et al., 1997; Cais et al., 2008; De Koninck and Mody, 1994; Esmann and Skou, 1988; Hartmann and Nothwang, 2011; Iversen and Neal, 1968; Jenkins et al., 1999; Koike and Nagata, 1979; Korogod and Demianenko, 2017; Owen et al., 2019; Postlethwaite et al., 2007; Pyott and Rosenmund, 2002; Sabatini and Regehr, 1996). In addition to delays in signal transduction, another possible reason may be the inactivation of GABAergic inhibition. A decrease in inhibition caused by focal cortical cooling may lead to a sustained excitation (Bruyns-Haylett et al., 2017; Kunori et al., 2014), resulting in a delay in the peak latency. In the present study, the application of a $GABA_A$ receptor antagonist increased SEP latency, whereas the application of AMPA and NMDA receptor antagonists induced only small changes. Thus, the sustained excitation caused by disinhibition (i.e., a decrease in inhibition) at lower temperatures might have contributed to the elongated latencies in the no-antagonist conditions.

In this study, we recorded SEPs by stimulating the contralateral forepaw while altering the local brain temperature. The results demonstrated that the relationship between SEP amplitude and

cortical temperature was reflected by an inverted U-curve and that inhibitory inputs played a key role in determining this relationship. Recently, autism spectrum disorder (ASD), whose typical symptoms include tactile hypersensitivity, has been associated with an increase in the balance of excitation and inhibition in the cortex (Chen et al., 2020; Kondo and Lin, 2020; Lee et al., 2017; Nelson and Valakh, 2015; Rubenstein and Merzenich, 2003). Interestingly, autistic behaviors have been reported to be relieved during fever in ASD patients (Curran et al., 2007; Good, 2017), although the autistic traits evaluated using questionnaires were shown to have no significant relationship with normal body (axillary) temperatures in the general population (Hidaka et al., 2023).

In conclusion, the negative correlation between SEP amplitudes and cortical temperature around the physiological cortical temperature may be caused by the high sensitivity of GABAergic inhibitory signaling to the temperature. Upon administration of the $GABA_A$ receptor antagonist, gabazine, the negative correlation plateaued, and this change occurred progressively as the concentration of gabazine increased. In contrast, a negative correlation remained after administration of the AMPA receptor antagonist, NBQX, or NMDA receptor antagonist, (R)-CPP. These results suggest that GABAergic inhibitory inputs contribute to this negative correlation by suppressing SEPs to a greater extent at higher temperatures. Our findings contribute not only to the elucidation of how cortical temperature influences sensory processing under physiological conditions but also to our understanding of psychiatric disorders and establishing new treatments via future research.

4. Experimental Procedure

4.1 Animals

Thirty-two male Wistar rats (253–331 g; Japan SLC Inc., Shizuoka, Japan) were used in this study. All experimental procedures were performed in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals and approved by the Animal Care and Use Committee of the National Institute of Advanced Industrial Science and Technology (AIST).

4.2 Experimental setup

Before surgery, anesthesia was induced using 3% isoflurane. For electrophysiological recordings, the skull above the left somatosensory cortex was removed to create a square hole (anteroposterior, ± 2 mm; mediolateral, 2–6 mm from the bregma). The dura mater covering the exposed area was left intact. A custom-made thermal control chamber (inner diameter of approximately 5 mm) developed previously (Gotoh et al., 2020) was positioned over the skull window (Fig. 1A).

The chamber was filled with 100 µL saline or saline containing an antagonist. The chamber was covered with a plastic sheet to prevent evaporation. Cortical temperature was regulated by circulating temperature-controlled water through a stainless-steel tube (Fig. 1A). Cortical temperature was monitored by inserting a thermocoupled electrode (MT-29/2; Physitemp Instruments, Inc., Clifton, NJ, USA) at a depth of 1 mm into the cortex. The cortical temperature and the temperature of the water circulating in the tube were linearly correlated ($R^2 = 0.99$; Fig. 1B). The body (rectal) temperature was maintained at 38°C using a heating pad (BWT-100A; BioResearch Center, Nagoya, Japan).

A silver ball electrode (UL-3010; Unique Medical Co., Ltd., Tokyo, Japan; diameter = 1 mm) was placed on the cortical surface to record cortical field potentials. The field potentials were band-pass filtered at 0.1–1,000 Hz. The location of the ball electrode was 4–5 mm lateral from the midline and ± 0 mm anteroposterior to the bregma, and the thermocoupled electrode was located approximately 1.5 mm away from the ball electrode. To induce SEP, a single electrical pulse was delivered to the contralateral (right) forepaw (amplitude: 1 mA; duration: 1 ms). During electrophysiological recordings, anesthesia was maintained using 1.5% isoflurane.

4.3 Experiment 1: Effects of cortical temperature on SEPs

Twenty-four rats were used in Experiment 1. The chamber was filled with a saline solution. The cortical temperature was changed within the range of 17–38°C in 3.5°C steps. The SEPs were recorded in 30 trials at each temperature. Electrical pulses were delivered every 9.9 s.

Thirty min after the surgery, the chamber was filled with 100 µL saline. The cortical temperature was maintained at 38°C for 1 h. The order of temperature change was counterbalanced (12 rats in ascending order and the other 12 rats in descending order). In rats assigned to the descending order, the initial SEP recording took place at 38°C. On the other hand, for those in the ascending order, the cortical temperature decreased before SEP recording, with the first recording conducted at 17°C (approximately 30 min after the cortical temperature reached 18°C). After each 30-trial recording, the cortical temperature was changed to the next temperature. The recording started ≥10 min after the temperature reached the range of ± 1°C.

4.4 Experiment 2: Effects of GABA$_A$ receptor antagonist

Eight of the 24 rats used in Experiment 1 were used in Experiment 2. After Experiment 1, the saline in the chamber was replaced with 10 µM (in 100 µL saline) of the GABA$_A$ receptor antagonist SR-95531 (gabazine; Toronto Research Chemicals; Toronto, ON, Canada). After incubation for 1 h at 38°C, the SEPs were recorded using the same procedure as in Experiment 1. The order of assignment in Experiments 1 and 2 was consistent within each rat.

4.5 Experiment 3: Effects of different concentrations of GABA$_A$ receptor antagonist around the physiological temperature

Eight rats were used in Experiment 3. The cortical temperature was changed within the range of 35–38°C in 1.5°C steps (i.e., 35, 36.5, and 38°C). First, SEPs were recorded using a chamber filled with saline. The order of the temperature changes was counterbalanced (four for the descending order and four for the ascending order). Second, the saline in the chamber was replaced with 1 µM gabazine (in 100 µL saline). After an incubation of 1 h at 38°C, SEPs were recorded at three temperatures. Third, the liquid in the chamber was replaced with 10 µM gabazine (in 100 µL saline). After incubation at 38°C for 1 h, SEPs were recorded. The order of temperature change was consistent within each rat. Time intervals and other procedures followed those used in Experiments 1 and 2.

4.6 Experiment 4: Effects of AMPA receptor antagonist

Eight of the 24 rats used in Experiment 1 were included in Experiment 4. Following Experiment 1, the saline in the chamber was substituted with a 10 µM solution (in 100 µL saline) of the AMPA receptor antagonist NBQX disodium salt (NBQX; Tocris Bioscience; Bristol, UK). Time intervals and other procedures followed those used in Experiment 2.

4.7 Experiment 5: Effects of NMDA receptor antagonist

Eight of the 24 rats used in Experiment 1 were included in Experiment 5. Following Experiment 1, the saline in the chamber was substituted with a 10 µM solution (in 100 µL saline) of the NMDA receptor antagonist (R)-CPP (Tocris Bioscience; Bristol, UK). Time intervals and other procedures followed those used in Experiment 2.

4.8 Data analysis

The SEP data between −1,000 and 200 ms from the onset of the electrical pulse (0 ms) were used for the data analysis. The SEP data for each trial were recalibrated for an average of −50 and 0 ms to obtain the baseline (0 mV). According to a preliminary analysis, SEPs were sometimes disturbed by spontaneous activity close to or during the evoked response. To precisely evaluate the effect of temperature on SEPs, data from trials in which spontaneous activity was not observed close to or during stimulation were analyzed. First, the data of trials in which the difference between maximum and minimum values in any 25-ms time bin between −1,000 and 0 ms exceeded 1.0 mV were excluded from the analyses (1,022 of 12,240 trials). Second, the data of trials in which the absolute values between 2 and 7 ms (i.e., for 5 ms after the electrical noise disappeared) exceeded 0.5 mV were excluded from the analyses (36 of the remaining 11,218 trials). In total, data from 1,058 of 12,240 (8.6%) trials were excluded from the analyses. Following exclusion, the SEP data for each temperature in each animal were averaged.

Positive peak, negative peak, and peak-to-peak amplitudes were calculated for the averaged SEP data at each temperature for each animal. The averaged data were smoothed using a moving-average filter with five-time bins (approximately 0.4 ms). To automatically find the peaks of P1 and N1 deflections, we used the following algorithm: 1) Find the local maximums during 2–20 ms (2–30 ms in Experiment 2) after stimulation. 2) The largest local maximum was defined as the P1 peak. 3) Find the first local minimum ≥2 ms after the P1 peak. 4) Find the local minimums during 0–5 ms after the first local minimum. 5) The smallest local minimum was defined as the N1 peak. The P1 peak amplitude (V_p) was defined as the voltage at P1 and the N1 peak amplitude (V_n) was defined as the sign-inverted voltage at N1. Peak-to-peak amplitude (V_{p-}

$_n$) was defined as the difference between P1 and N1. The P1 (L_p) and N1 peak latencies (L_n) were defined as the time between the onset of stimulation (0 ms) and the respective peaks.

To examine temperature dependency on V_p, V_n, and V_{p-n}, we calculated correlation coefficients between these amplitudes and cortical temperatures in the range of $\leq 27.5°C$ and $\geq 27.5°C$, separately, in Experiments 1, 2, 4, and 5. To examine the effects of cortical temperature on the amplitudes and latencies with no antagonist, we performed a two-way ANOVA (temperature by animal) for the amplitudes and latencies in Experiment 1. To examine the effects of cortical temperature and antagonists, we performed three-way ANOVAs (antagonist by temperature by animal) for the amplitudes and latencies in Experiment 2–5. To examine the effects of antagonists on the amplitudes at each temperature, we calculated the difference and ratio between the amplitudes before (Experiment 1) and after antagonist application (Experiments 2, 4, and 5) as follows:

$$difference(V) = V_{after} - V_{before}$$

$$ratio(V) = V_{after}/V_{before}$$

where V denotes the amplitude (V_p, V_n, or V_{p-n}), and V_{before} and V_{after} denote the amplitudes before and after the antagonist application, respectively. For these differences and ratios, we calculated the correlation coefficients with the temperature. In Experiment 3, the amplitudes were fitted using the following linear function:

$$V(T) = C_1(T - 36.5) + C_0$$

where T denotes the cortical temperature, V denotes the amplitude, C_1 denotes the slope of the amplitude, and C_0 denotes the constant term (that is, the mean amplitudes at 36.5°C). Tests for the null hypothesis that the slope is equal to zero were performed using a t-test. Additionally, we performed two-way ANOVAs (antagonist by animal) for C_0 and C_1. MATLAB (R2020b, The MathWorks, Inc., Natick, MA, USA) was used for all data analyses.

Declaration of Interests: none.

Acknowledgments

We thank Shigeru Yamane, Yoshihisa Tachibana, and Makoto Wada for their valuable discussions.

Funding

This work was supported by the Japan Society for the Promotion of Science (JSPS) [KAKENHI Grant Numbers 19K22990 and 21H03532 to I.T. and 19K22585 and 21H03788 to S.Y.].

Figure Captions

Figure 1. Experimental procedures. **(A)** Schematic illustration of the experimental setup. The right forepaw was stimulated, and somatosensory evoked potentials (SEPs) were recorded from the contralateral somatosensory cortex in anesthetized rats. The thermal control chamber was placed above the somatosensory cortex. **(B)** Regulation of the cortical temperature by the thermal control chamber. Cortical temperatures (T_{cortex}), measured at 1 mm from the surface during experiments without the antagonist, are plotted against the temperature of the circulating water (T_{water}) through a stainless-steel tube inside the chamber and fitted by a linear function ($T_{cortex} = 0.068\text{x} \times T_{water} + 10.56$, $R^2 = 0.99$, $n = 32$ animals).

Figure 2. Effects of local cortical temperature on somatosensory evoked potentials (SEPs) (Experiment 1). **(A)** Averaged SEP waveforms at various temperatures from 24 animals. The time 0 indicates the onset of the electrical stimulation. **(B)** Indices of SEPs. The peak amplitudes of the positive peak (V_p) and negative peak (V_n) were defined as the absolute voltage of the respective peaks. The peak-to-peak amplitude (V_{p-n}) was defined as the difference between positive and negative peaks. The peak latencies of the positive peak (L_p) and negative peak (L_n) were defined as the time between the onset of stimulation (0 ms) and the respective peaks. **(C–E)** Effects of cortical temperature on V_{p-n} (C), V_p (D)**,** and V_n (E). Amplitudes are plotted against the cortical temperatures (average data: filled circles and thick lines, individual data: semitransparent circles and thin lines; $n = 24$ animals). Error bars indicate mean $\pm$ standard error. **(F–G)** Effects of cortical temperature on L_p (F) and L_n (G). Peak latencies are plotted against cortical temperatures (average data: filled circles and thick lines, individual data: semitransparent circles and lines; $n = 24$ animals). Error bars indicate mean $\pm$ SE.

Figure 3. Effects of the gamma-aminobutyric acid type A (GABA$_A$) receptor antagonist, gabazine, on the temperature dependency of somatosensory evoked potentials (SEPs) (Experiment 2). **(A)** Averaged SEP waveforms before (left panel) and after 10 μM gabazine administration (right panel) at various temperatures from eight animals. **(B–F)** Effects of cortical temperature on V_{p-n} (B), V_p (C), V_n (D), L_p (E), and L_n (F) before (black circles and lines) and after (red squares and lines) gabazine administration ($n = 8$ animals). Error bars indicate mean $\pm$ standard error.

Figure 4. Effects of the concentration of the gamma-aminobutyric acid type A (GABA$_A$) receptor antagonist, gabazine, on the temperature dependency of somatosensory evoked potentials (SEPs) (Experiment 3). **(A–E)** Effects of cortical temperature on V_{p-n} (A), V_p (B), V_n

(C), L_p (D), and L_n (E) in the saline (black circles and lines), 1-μM gabazine (red squares and lines), and 10-μM gabazine (blue triangles and lines) condition ($n = 8$ animals). Error bars indicate mean ± standard error. **(F)** The constant terms (top panel; the mean amplitudes at 36.5°C) and slopes (bottom panel) of the fitted linear models for $V_{p\text{-}n}$, V_p, V_n plotted against the gabazine concentration (black circles and lines; $n = 8$ animals).

Figure 5. Effects of the aminomethylphosphonic acid (AMPA) receptor antagonist, NBQX, on the temperature dependency of somatosensory evoked potentials (SEPs) (Experiment 4). **(A)** Averaged SEP waveforms before (left panel) and after (right panel) NBQX administration at various temperatures from eight animals. **(B–F)** Effects of cortical temperature on $V_{p\text{-}n}$ (B), V_p (C), V_n (D), L_p (E), and L_n (F) before (black circles and lines) and after (red squares and lines) NBQX administration ($n = 8$ animals). Error bars indicate mean ± standard error.

Figure 6. Effects of the NMDA receptor antagonist (R)-CPP on the temperature dependency of somatosensory evoked potentials (SEPs) (Experiment 5). **(A)** Averaged SEP waveforms before (left panel) and after (right panel) (R)-CPP administration at various temperatures from eight animals. **(B–F)** Effects of cortical temperature on $V_{p\text{-}n}$ (B), V_p (C), V_n (D), L_p (E), and L_n (F) before (black circles and lines) and after (red squares and lines) (R)-CPP administration ($n = 8$ animals). Error bars indicate mean ± standard error.

References

All, A.H., Luo, S., Liu, X., Al-Nashash, H., 2021. Effect of thoracic spinal cord injury on forelimb somatosensory evoked potential. Brain Res. Bull. 173**,** 22–27. https://doi.org/10.1016/j.brainresbull.2021.05.005

Asztely, F., Erdemli, G., Kullmann, D.M., 1997. Extrasynaptic glutamate spillover in the hippocampus: dependence on temperature and the role of active glutamate uptake. Neuron. 18**,** 281–293. https://doi.org/10.1016/s0896-6273(00)80268-8

Bannister, A.P., 2005. Inter- and intra-laminar connections of pyramidal cells in the neocortex. Neurosci. Res. 53**,** 95–103. https://doi.org/10.1016/j.neures.2005.06.019

Bauer, R.H., Fuster, J.M., 1976. Delayed-matching and delayed-response deficit from cooling dorsolateral prefrontal cortex in monkeys. J. Comp. Physiol. Psychol. 90**,** 293–302. https://doi.org/10.1037/h0087996

Beaulieu, C., 1993. Numerical data on neocortical neurons in adult rat, with special reference to the GABA population. Brain Res. Bull. 609**,** 284–292. https://doi.org/10.1016/0006-8993(93)90884-p

Bindman, L.J., Lippold, O.C., Redfearn, J.W., 1963. Comparison of the effects on electrocortical activity of general body cooling of the surface of the brain. Electroencephalogr Clin Neurophysiol. 15**,** 238-245. https://doi.org/10.1016/0013-4694(63)90093-2

Bowery, N.G., Hudson, A.L., Price, G.W., 1987. GABAA and GABAB receptor site distribution in the rat central nervous system. Neuroscience. 20**,** 365–383. https://doi.org/10.1016/0306-4522(87)90098-4

Bruyns-Haylett, M., Luo, J., Kennerley, A.J., Harris, S., Boorman, L., Milne, E., Vautrelle, N., Hayashi, Y., Whalley, B.J., Jones, M., Berwick, J., Riera, J., Zheng, Y., 2017. The neurogenesis of P1 and N1: A concurrent EEG/LFP study. Neuroimage. 146**,** 575–588. https://doi.org/10.1016/j.neuroimage.2016.09.034

Cais, O., Sedlacek, M., Horak, M., Dittert, I., Vyklicky, L., Jr., 2008. Temperature dependence of NR1/NR2B NMDA receptor channels. Neuroscience. 151**,** 428–438. https://doi.org/10.1016/j.neuroscience.2007.11.002

Carter, B.G., Butt, W., 2005. Are somatosensory evoked potentials the best predictor of outcome after severe brain injury? A systematic review. Intensive Care Med. 31**,** 765–775. https://doi.org/10.1007/s00134-005-2633-1

Chen, Q., Deister, C.A., Gao, X., Guo, B., Lynn-Jones, T., Chen, N., Wells, M.F., Liu, R., Goard, M.J., Dimidschstein, J., Feng, S., Shi, Y., Liao, W., Lu, Z., Fishell, G., Moore,

C.I., Feng, G., 2020. Dysfunction of cortical GABAergic neurons leads to sensory hyper-reactivity in a Shank3 mouse model of ASD. Nat. Neurosci. 23, 520–532. https://doi.org/10.1038/s41593-020-0598-6

Chow, B.Y., Han, X., Dobry, A.S., Qian, X., Chuong, A.S., Li, M., Henninger, M.A., Belfort, G.M., Lin, Y., Monahan, P.E., Boyden, E.S., 2010. High-performance genetically targetable optical neural silencing by light-driven proton pumps. Nature. 463, 98–102. https://doi.org/10.1038/nature08652

Cruccu, G., Aminoff, M.J., Curio, G., Guerit, J.M., Kakigi, R., Mauguiere, F., Rossini, P.M., Treede, R.D., Garcia-Larrea, L., 2008. Recommendations for the clinical use of somatosensory-evoked potentials. Clin. Neurophysiol. 119, 1705–1719. https://doi.org/10.1016/j.clinph.2008.03.016

Curran, L.K., Newschaffer, C.J., Lee, L.C., Crawford, S.O., Johnston, M.V., Zimmerman, A.W., 2007. Behaviors associated with fever in children with autism spectrum disorders. Pediatrics. 120, e1386-e1392. https://doi.org/10.1542/peds.2007-0360

De Koninck, Y., Mody, I., 1994. Noise analysis of miniature IPSCs in adult rat brain slices: properties and modulation of synaptic GABAA receptor channels. J. Neurophysiol. 71, 1318–1335. https://doi.org/10.1152/jn.1994.71.4.1318

Deuel, R., Mishkin, M., 1977. Limbic and prefrontal contributions to somesthetic learning in monkeys. Brain Res. Bull. 132, 521–535. https://doi.org/10.1016/0006-8993(77)90199-8

Dezawa, S., Nagasaka, K., Watanabe, Y., Takashima, I., 2021. Lesions of the nucleus basalis magnocellularis (Meynert) induce enhanced somatosensory responses and tactile hypersensitivity in rats. Exp. Neurol. 335, 113493. https://doi.org/10.1016/j.expneurol.2020.113493

Di, S., Barth, D.S., 1991. Topographic analysis of field potentials in rat vibrissa/barrel cortex. Brain Res. 546, 106–112. https://doi.org/10.1016/0006-8993(91)91164-v

Esmann, M., Skou, J.C., 1988. Temperature-dependencies of various catalytic activities of membrane-bound Na+/K+-ATPase from ox brain, ox kidney and shark rectal gland and of C12E8-solubilized shark Na+/K+-ATPase. Biochim. Biophys. Acta. 944, 344–50. https://doi.org/10.1016/0005-2736(88)90504-4

Florence, G., Guerit, J.M., Gueguen, B., 2004. Electroencephalography (EEG) and somatosensory evoked potentials (SEP) to prevent cerebral ischaemia in the operating room. Neurophysiol. Clin. 34, 17–32. https://doi.org/10.1016/j.neucli.2004.01.001

Franceschini, M.A., Nissilä, I., Wu, W., Diamond, S.G., Bonmassar, G., Boas, D.A., 2008. Coupling between somatosensory evoked potentials and hemodynamic response in the

rat. NeuroImage. 41, 189–203. https://doi.org/10.1016/j.neuroimage.2008.02.061

Freeman, S., Sohmer, H., 1996. A comparison of forepaw and vibrissae somatosensory cortical evoked potentials in the rat. Electroencephalogr. Clin. Neurophysiol. 100, 362–369. https://doi.org/10.1016/0168-5597(96)95690-7

Fuster, J.M., Bauer, R.H., 1974. Visual short-term memory deficit from hypothermia of frontal cortex. Brain Res. 81, 393–400. https://doi.org/10.1016/0006-8993(74)90838-5

Fuster, J.M., Bauer, R.H., Jervey, J.P., 1981. Effects of cooling inferotemporal cortex on performance of visual memory tasks. Exp. Neurol. 71, 398–409. https://doi.org/10.1016/0014-4886(81)90098-4

Good, P., 2017. Simplifying study of fever's dramatic relief of autistic behavior. Clin. Nutr. ESPEN. 17, 1–7. https://doi.org/10.1016/j.clnesp.2016.09.002

Gotoh, M., Nagasaka, K., Nakata, M., Takashima, I., Yamamoto, S., 2020. Brain temperature alters contributions of excitatory and inhibitory inputs to evoked field potentials in the rat frontal cortex. Front. Cell. Neurosci. 14, 593027. https://doi.org/10.3389/fncel.2020.593027

Han, X., Boyden, E.S., 2007. Multiple-color optical activation, silencing, and desynchronization of neural activity, with single-spike temporal resolution. PLOS ONE. 2, e299. https://doi.org/10.1371/journal.pone.0000299

Han, X., Chow, B.Y., Zhou, H., Klapoetke, N.C., Chuong, A., Rajimehr, R., Yang, A., Baratta, M.V., Winkle, J., Desimone, R., Boyden, E.S., 2011. A high-light sensitivity optical neural silencer: development and application to optogenetic control of non-human primate cortex. Front. Syst. Neurosci. 5, 18. https://doi.org/10.3389/fnsys.2011.00018

Hartmann, A.M., Nothwang, H.G., 2011. Opposite temperature effect on transport activity of KCC2/KCC4 and N(K)CCs in HEK-293 cells. BMC Res. Notes. 4, 526. https://doi.org/10.1186/1756-0500-4-526

Hidaka, S., Gotoh, M., Yamamoto, S., Wada, M., 2023. Exploring relationships between autistic traits and body temperature, circadian rhythms, and age. Sci. Rep. 13, 5888. https://doi.org/10.1038/s41598-023-32449-z

Hikosaka, O., Tanaka, M., Sakamoto, M., Iwamura, Y., 1985. Deficits in manipulative behaviors induced by local injections of muscimol in the first somatosensory cortex of the conscious monkey. Brain Res. 325, 375–380.

Hikosaka, O., Wurtz, R.H., 1985a. Modification of saccadic eye movements by GABA-related substances. I. Effect of muscimol and bicuculline in monkey superior colliculus. J. Neurophysiol. 53, 266–291.

Hikosaka, O., Wurtz, R.H., 1985b. Modification of saccadic eye movements by GABA-related

substances. II. Effects of muscimol in monkey substantia nigra pars reticulata. J. Neurophysiol. 53, 292–308.

Hupé, J.M., James, A.C., Payne, B.R., Lomber, S.G., Girard, P., Bullier, J., 1998. Cortical feedback improves discrimination between figure and background by V1, V2 and V3 neurons. Nature. 394, 784–787. https://doi.org/10.1038/29537

Iversen, L.L., Neal, M.J., 1968. The uptake of [3H]GABA by slices of rat cerebral cortex. J. Neurochem. 15, 1141–1149. https://doi.org/10.1111/j.1471-4159.1968.tb06831.x

Jackson, A.F., Bolger, D.J., 2014. The neurophysiological bases of EEG and EEG measurement: a review for the rest of us. Psychophysiology. 51, 1061–1071. https://doi.org/10.1111/psyp.12283

Jellema, T., Brunia, C.H., Wadman, W.J., 2004. Sequential activation of microcircuits underlying somatosensory-evoked potentials in rat neocortex. Neuroscience. 129, 283–295. https://doi.org/10.1016/j.neuroscience.2004.07.046

Jenkins, A., Franks, N.P., Lieb, W.R., 1999. Effects of temperature and volatile anesthetics on GABA(A) receptors. Anesthesiology. 90, 484–491. https://doi.org/10.1097/00000542-199902000-00024

Johnson, V.E., Meaney, D.F., Cullen, D.K., Smith, D.H., 2015. Animal models of traumatic brain injury. Handb. Clin. Neurol. 127, 115–128. https://doi.org/10.1016/B978-0-444-52892-6.00008-8

Karkar, K.M., Garcia, P.A., Bateman, L.M., Smyth, M.D., Barbaro, N.M., Berger, M., 2002. Focal cooling suppresses spontaneous epileptiform activity without changing the cortical motor threshold. Epilepsia. 43, 932–935. https://doi.org/10.1046/j.1528-1157.2002.03902.x

Koike, H., Nagata, Y., 1979. Intra-axonal diffusion of [3H]acetylcholine and [3H]gamma-aminobutyric acid in a neurone of Aplysia. J. Physiol. 295, 397–417. https://doi.org/10.1113/jphysiol.1979.sp012976

Kondo, H.M., Lin, I.F., 2020. Excitation-inhibition balance and auditory multistable perception are correlated with autistic traits and schizotypy in a non-clinical population. Sci. Rep. 10, 8171. https://doi.org/10.1038/s41598-020-65126-6

Korogod, S.M., Demianenko, L.E., 2017. Temperature effects on non-TRP ion channels and neuronal excitability. Opera Med. Physiol. 3, 84–92.

Kunori, N., Kajiwara, R., Takashima, I., 2014. Voltage-sensitive dye imaging of primary motor cortex activity produced by ventral tegmental area stimulation. J. Neurosci. 34, 8894–8903. https://doi.org/10.1523/JNEUROSCI.5286-13.2014

Larkum, M.E., Nevian, T., Sandler, M., Polsky, A., Schiller, J., 2009. Synaptic integration in tuft

dendrites of layer 5 pyramidal neurons: a new unifying principle. Science. 325, 756–60. https://doi.org/10.1126/science.1171958

Lee, E., Lee, J., Kim, E., 2017. Excitation/inhibition imbalance in animal models of autism spectrum disorders. Biol. Psychiatry. 81, 838–847. https://doi.org/10.1016/j.biopsych.2016.05.011

Lomber, S.G., Malhotra, S., 2008. Double dissociation of 'what' and 'where' processing in auditory cortex. Nat. Neurosci. 11, 609–616. https://doi.org/10.1038/nn.2108

Lund, J.S., 1973. Organization of neurons in the visual cortex, area 17, of the monkey (Macaca mulatta). J. Comp. Neurol. 147, 455–496. https://doi.org/10.1002/cne.901470404

MacDonald, D.B., Dong, C., Quatrale, R., Sala, F., Skinner, S., Soto, F., Szelényi, A., 2019. Recommendations of the International Society of Intraoperative Neurophysiology for intraoperative somatosensory evoked potentials. Clin. Neurophysiol. 130, 161–179. https://doi.org/10.1016/j.clinph.2018.10.008

Minamimoto, T., Saunders, R.C., Richmond, B.J., 2010. Monkeys quickly learn and generalize visual categories without lateral prefrontal cortex. Neuron. 66, 501–507. https://doi.org/10.1016/j.neuron.2010.04.010

Mitzdorf, U., 1985. Current source-density method and application in cat cerebral cortex: investigation of evoked potentials and EEG phenomena. Physiol. Rev. 65, 37–100. https://doi.org/10.1152/physrev.1985.65.1.37

Moseley, J.I., Ojemann, G.A., Ward, A.A., Jr., 1972. Unit activity during focal cortical hypothermia in the normal cortex. Exp. Neurol. 37, 152–63. https://doi.org/10.1016/0014-4886(72)90232-4

Nakata, M., Nagasaka, K., Shimoda, M., Takashima, I., Yamamoto, S., 2018. Focal brain lesions induced with ultraviolet irradiation. Sci. Rep. 8, 7968. https://doi.org/10.1038/s41598-018-26117-w

Nakata, M., Shimoda, M., Yamamoto, S., 2021. UV-induced neuronal degeneration in the rat cerebral cortex. Cereb. Cortex Commun. 2, tgab006.

Nelson, S.B., Valakh, V., 2015. Excitatory/inhibitory balance and circuit homeostasis in autism spectrum disorders. Neuron. 87, 684–698. https://doi.org/10.1016/j.neuron.2015.07.033

Newsome, W.T., Wurtz, R.H., Dürsteler, M.R., Mikami, A., 1985. Punctate chemical lesions of striate cortex in the macaque monkey: effect on visually guided saccades. Exp. Brain Res. 58, 392–399. https://doi.org/10.1007/BF00235320

Nomura, S., Inoue, T., Imoto, H., Suehiro, E., Maruta, Y., Hirayama, Y., Suzuki, M., 2017. Effects of focal brain cooling on extracellular concentrations of neurotransmitters in patients with epilepsy. Epilepsia. 58, 627–634. https://doi.org/10.1111/epi.13704

Owen, S.F., Liu, M.H., Kreitzer, A.C., 2019. Thermal constraints on in vivo optogenetic manipulations. Nat. Neurosci. 22, 1061–1065. https://doi.org/10.1038/s41593-019-0422-3

Palmer, L.M., 2014. Dendritic integration in pyramidal neurons during network activity and disease. Brain Res. Bull. 103, 2–10. https://doi.org/10.1016/j.brainresbull.2013.09.010

Payne, B.R., Lomber, S.G., Geeraerts, S., van der Gucht, E., Vandenbussche, E., 1996. Reversible visual hemineglect. Proc. Natl. Acad. Sci. U. S. A. 93, 290–294. https://doi.org/10.1073/pnas.93.1.290

Postlethwaite, M., Hennig, M.H., Steinert, J.R., Graham, B.P., Forsythe, I.D., 2007. Acceleration of AMPA receptor kinetics underlies temperature-dependent changes in synaptic strength at the rat calyx of Held. J. Physiol. 579, 69–84. https://doi.org/10.1113/jphysiol.2006.123612

Pyott, S.J., Rosenmund, C., 2002. The effects of temperature on vesicular supply and release in autaptic cultures of rat and mouse hippocampal neurons. J. Physiol. 539, 523–535. https://doi.org/10.1113/jphysiol.2001.013277

Que, M., Witte, O.W., Neumann-Haefelin, T., Schiene, K., Schroeter, M., Zilles, K., 1999. Changes in GABA(A) and GABA(B) receptor binding following cortical photothrombosis: a quantitative receptor autoradiographic study. Neuroscience. 93, 1233–1240. https://doi.org/10.1016/s0306-4522(99)00197-9

Rockland, K.S., Pandya, D.N., 1979. Laminar origins and terminations of cortical connections of the occipital lobe in the rhesus monkey. Brain Res. 179, 3–20. https://doi.org/10.1016/0006-8993(79)90485-2

Rubenstein, J.L., Merzenich, M.M., 2003. Model of autism: increased ratio of excitation/inhibition in key neural systems. Genes Brain Behav. 2, 255–267. https://doi.org/10.1034/j.1601-183x.2003.00037.x

Rubio-Garrido, P., Pérez-de-Manzo, F., Porrero, C., Galazo, M.J., Clascá, F., 2009. Thalamic input to distal apical dendrites in neocortical layer 1 is massive and highly convergent. Cereb. Cortex. 19, 2380–2395. https://doi.org/10.1093/cercor/bhn259

Sabatini, B.L., Regehr, W.G., 1996. Timing of neurotransmission at fast synapses in the mammalian brain. Nature. 384, 170–172. https://doi.org/10.1038/384170a0

Sartorius, C.J., Berger, M.S., 1998. Rapid termination of intraoperative stimulation-evoked seizures with application of cold Ringer's lactate to the cortex. Technical note. J. Neurosurg. 88, 349–351. https://doi.org/10.3171/jns.1998.88.2.0349

Schwarcz, R., Hökfelt, T., Fuxe, K., Jonsson, G., Goldstein, M., Terenius, L., 1979. Ibotenic acid-induced neuronal degeneration: a morphological and neurochemical study. Exp.

Brain Res. 37**,** 199–216. https://doi.org/10.1007/BF00237708

Schwerdtfeger, K., von Tiling, S., Kiefer, M., Strowitzki, M., Mestres, P., Booz, K.H., Steudel, W.I., 1999. Identification of somatosensory pathways by focal-cooling-induced changes of somatosensory evoked potentials and EEG-activity--an experimental study. Acta Neurochir. (Wien). 141**,** 647–654. https://doi.org/10.1007/s007010050355

Suzuki, T., Sakata, H., Kato, C., Connor, J.A., Morita, M., 2012. Astrocyte activation and wound healing in intact-skull mouse after focal brain injury. Eur. J. Neurosci. 36**,** 3653–3664. https://doi.org/10.1111/j.1460-9568.2012.08280.x

Takei, T., Lomber, S.G., Cook, D.J., Scott, S.H., 2021. Transient deactivation of dorsal premotor cortex or parietal area 5 impairs feedback control of the limb in macaques. Curr. Biol. 31**,** 1476–1487. https://doi.org/10.1016/j.cub.2021.01.049

Trendelenburg, W., 1911. Untersuchungen über reizlose Vorübergehende Ausschaltung am Zentralnervensystem: III. Mitteilung. Die Extremitätenregion der Grosshirnrinde. Pflüger's Arch. 137**,** 515–544.

A

Thermocoupled electrode

Ball electrode

Chamber

Stainless-steel tube

Temperature-controlled water

B