Polarised Moonlight Guides Nocturnal Bull Ants Home

Cody A Freas*, Ajay Narendra; Trevor Murray; Ken Cheng

School of Natural Sciences, Macquarie University, Sydney, NSW Australia

* Corresponding Author

Address for correspondence:
Cody A Freas
School of Natural Sciences
Macquarie University
Sydney, NSW 2113 Australia
Email: cody.freas@mq.edu.au

Abstract

For the first time in any animal, we show that nocturnal bull ants use the exceedingly dim polarisation pattern produced by the moon for overnight navigation. The sun or moon can provide directional information via their position; however, they can often be obstructed by clouds, canopy or the horizon. Despite being hidden, these bodies can still provide compass information through the polarised light pattern they produce/reflect. Sunlight produces polarised light patterns across the overhead sky as it enters the atmosphere, and solar polarised light is a well-known compass cue for navigating animals. Moonlight produces an analogous pattern, albeit a million times dimmer than sunlight. Here we show the first evidence that polarised moonlight forms part of the celestial compass of navigating nocturnal ants. Nocturnal bull ants leave their nest at twilight and rely heavily on the overhead solar polarisation pattern to navigate. Yet many foragers return home overnight when the sun cannot guide them. We demonstrate that these bull ants use polarised moonlight to navigate home during the night, by rotating the ambient polarisation pattern above homing ants, who alter their headings in response. Furthermore, these ants can detect this cue throughout the lunar month, even under crescent moons, when polarised light levels are at their lowest. Finally, we show long-term incorporation of this moonlight pattern into the ants' path integration system throughout the night for homing, as polarised sunlight is incorporated throughout the day.

Keywords: ant navigation, lunar cues, route maintenance, cue detection, polarised light, vector

Introduction

Many navigating animals attend to the position of the sun or moon to guide their movement (Jander 1957; Koltz and Reid 1993; Perez et al. 1997; Dacke et al. 2014; Warrant and Dacke 2016; Freas and Cheng 2022). Yet these celestial bodies are not always directly visible, often obscured by clouds, the canopy or after passing below the horizon, resulting in gaps for navigation relying solely upon direct visual detection. Animals hence rely on the pattern of polarised skylight which is accessible even when the celestial bodies are occluded to some extent (Horváth et al., 2014). Polarised light comprises light waves which occur along a single plane and are produced as a by-product of light passing through the upper atmosphere (Horváth & Varjú 2004; Horváth et al., 2014). E-vectors are arranged in concentric circles around the sun/moon with the maximum degree of polarisation located 90° from the source. Hence when the sun/moon is near the horizon, the pattern of polarised skylight is particularly simple with uniform direction of polarisation approximately parallel to the north-south axes (Dacke et al., 1999, 2003; Reid et al. 2011; Zeil et al., 2014). The pattern's stability makes the sky's polarisation a useful directional cue for orientation (Wehner and Müller 2006; Reid et al. 2011; Lebhardt and Ronacher 2013; Warrant and Dacke 2016; Freas et al. 2017a, 2019; Freas and Spetch 2023), which insects detect through specialised photoreceptors located in the dorsal rim area of their eyes (Labhart and Meyer 1999; Homberg and Paech 2002; el Jundi et al. 2015). Like solar polarisation, though a million times weaker, the moon reflects sunlight, producing a polarised moonlight pattern emanating from the moon's position in the night sky (Gál et al. 2001). Given that the moon creates a much dimmer version of the polarisation pattern formed around the sun, only night-navigating insects with eyes highly specialised for low light detection may be able to rely on this pattern to orient and navigate to goals.

Currently, only nocturnal dung beetles (*Scarabaeus satyrus* and *S. zambesianus*) are known to attend to moonlight polarisation patterns during their movement (Dacke et al. 2003, 2004, 2011; Foster et al. 2019). Yet interestingly, these beetles do not use moonlight to navigate, instead relying on this pattern to keep moving straight in order to roll their dung balls expeditiously away from a central dung pile. As such, this cue has only been documented for heading maintenance over short periods. While it has been theorised that this ability to detect the much dimmer polarisation pattern produced by the moon may be present across nocturnal insects more broadly, including nocturnal bees and crickets (Herzmann and Labhart 1989; Greiner et al. 2007; Rost and Honegger 1987), there is currently no behavioural evidence for its use in goal-directed navigation.

The large-eyed *Myrmecia* ants have several species that restrict majority of their navigation to evening twilight (outbound) and morning twilight (inbound) respectively (Narendra et al., 2017). We know that two well-studied nocturnal ant species, *Myrmecia pyriformis* and *Myrmecia midas*, use the overhead solar polarised light pattern, which is still visible during the twilight period to derive compass information (Reid et al. 2011; Freas et al. 2017ab; Freas and Cheng 2018). Because the information required for visual navigation degrades beyond twilight, it has been suggested that animals tend to be less active at night.

However, a small proportion of *M. pyriformis* foragers leave the nest (10.7% of daily foraging
force) or return home during night (13.3% of daily foraging force) (Reid et al., 2013). In *M.*
midas, this nocturnal activity is even more pronounced with a majority of the foraging force
returning during night (62.8% of daily foraging force) along with a minority of foragers
leaving the nest during night (26.2% of daily foraging force) (Freas et al. 2017a).

Nocturnal bull ants navigate using a combination of learned visual cues and homing
vectors obtained by integrating pedometric and celestial compass information. This 'true
nocturnal navigation' is likely aided by the increase in the light intensity provided by the
moon's presence. In *M. pyriformis*, on 'full-moon' nights there was a significantly greater
proportion of foragers leaving the nest at night compared to a 'new-moon' night (Reid et al.,
2013). This additional light at night may enhance terrestrial visual features foragers have
learned and therefore assist in visual guidance. In addition, the moon and the lunar
polarisation pattern also likely provides compass information, allowing foragers acquire or
follow a homing vector. Several arthropods, including ants and bees directly track the
moon's position to obtain compass information (Jander 1957; Koltz and Reid 1993; Dacke et
al. 2004; Ugolini et al. 2003). But given the moon may be occluded by clouds or overhanging
canopy we aim to identify here whether the lunar polarised skylight can also be used by ants
for homing.

In the current study, we provide the first evidence of polarised moonlight detection
and guidance in ants for goal-directed navigation. We studied foraging ants' ability to orient
during lunar twilight, by placing and rotating a linear polarising filter over them as they
returned to the nest (Figure 1A). We also explored whether these navigators weigh their
attendance to polarised moonlight across the lunar calendar, since during quarter-moon and
crescent-moon nights, smaller portions of the moon's surface reflect sunlight (as well as
moonless nights when no ambient e-vector is present). Finally, we characterised changes in
weighting that these ant navigators give polarised moonlight, as a function of the moon's
overnight consistency in the sky (waxing vs. waning) and the length of their accumulated
path integrator, which should increase the weight given to celestial compass portion of the
path integrator when in conflict with terrestrial visual cues (Burkhalter 1972; Narendra 2007;
Wystrach et al. 2015; Freas and Cheng 2019).

Methods
Study site
Experiments were conducted from April through October 2023 on two *Myrmecia midas*
nests on the Macquarie University Wallumattagal campus in Sydney, Australia (33°46 11S,
151°06 40E). *M. midas* nests are typically located within stands of *Eucalyptus* trees with the
nest entrance located near (<30cm) a tree. *M. midas* is nocturnal, with foraging onset
occurring ~20min after sunset when foragers leave the nest to travel to and up one of
several surrounding foraging trees overnight (Freas et al. 2018). Inbound navigation is more
variable with foragers returning to the nest entrance overnight and into morning twilight
(Freas et al. 2017a). For this study, we chose two nests under open canopies to ensure

foragers had unobstructed visual access to the overhead sky (Figure 1B). The understory of
these areas is naturally barren of vegetation, and we cleared the foraging column of debris
to aid visual tracking.
168
Apparatus: Polarisation filter
For each condition, we altered the ambient pattern of polarised moonlight by
rotating a linear polarisation filter (30cm diameter; same apparatus used in Freas et al. 2019)
above each ant along their inbound journey. This rotation alters the dominant ambient e-
vector direction of the sky above the navigator. This filter was held by a circular 1cm thick
ring and lifted 10cm off the ground by four equally spaced thin metal legs (Figure 1). All
testing was conducted overnight before morning solar twilight, during the lunar twilight for
each tested lunar phase. For each night we obtained the moon's position as it reached the
horizon based on the Astronomical Almanac (http://asa.usno.navy.mil) and set the ambient
lunar e-vector perpendicular from the moon's position at moonset. We relied on a digital
compass mobile application (Apple™) confirmed by an analogue compass to locate the
ambient e-vector and rotated the linear polariser by ±45° from this direction for each ant.
Across all conditions, we recorded four positions: the ant's release point, position
when the filter was placed overhead, the filter exit point, and reorientation after ~1m, taking
care to slowly follow the ant and mark their positions so as to not disturb their travel. These
positions determine each forager's initial orientation, exit orientation and reorientation
directions (Figure 1). After testing, each forager was marked with acrylic paint (Tamiya™) to
prevent retesting. Testing was conducted at distinct lunar phases which predicably occur
throughout the lunar month cycle (29.5 days).
188
Full moon
For full moon testing we chose nights during which the waxing moon's lunar phase was near
full but with clear separation between solar and lunar twilights. Testing on true full moon
nights is problematic as solar and lunar twilights fully overlap and the solar polarised light
pattern would overpower the lunar counterpart. Testing during the nights preceding the full
moon (waxing phase) ensured the moon's presence in the night sky overnight and testing
occurred on nights in which the lunar twilight (1am – 4am) was clearly separated from the
start of morning solar twilight (5:22am) with illumination above 80% of the lunar surface.
Outbound *M. midas* foragers from two nests were followed as they left the nest
during evening twilight and collected as they climbed onto their foraging tree, (Nest 1: 6.0m;
Nest 2: 3.1m from the nest entrance). Each forager was provided a small amount of honey
and held within a clear plastic phial on the ground 5m from the foraging tree with an
unobstructed view of the sky (Figure 1B). Foragers were held in these phials overnight until
the moon's position was within ±10º of the horizon (large stands of trees and buildings near
the western horizon occluded the moon's position during all testing; Figure 1B).
204
Waxing lunar phase

We observed clear shifts in forager headings at both nests under *Full Moon* conditions, yet
nights with over 80% lunar illumination only account for nine nights per lunar cycle. To
assess if polarised moonlight can be used throughout the lunar month, we tested foragers in
three further conditions representing distinct lunar phases: a *Waxing Quarter Moon, a*
Waxing Crescent Moon, and a *No Moon* control*.* For the *Quarter Moon* and *Crescent Moon*
conditions, we tested ants identically to full moon conditions; for overnight testing, however,
the moon has a different temporal period when it is positioned near the horizon (12am and
9pm, respectively).
The procedure was slightly modified for *No Moon* testing as we did not test these
foragers on the new moon night (the new moon is only present during the day). We
hypothesised that foragers with no available ambient polarisation and suddenly presented
with an e-vector pattern might fall back on a memory of the morning solar e-vector as many
foragers return during morning twilight and this direction remains consistent across nights.
In order to test on a night when there were distinct directional differences between the
lunar and morning solar e-vectors (Lunar e-vector: 329º; Morning solar e-vector: 7º; Evening
Solar e-vector: 353º), we chose to test on a quarter moon night when the moon was well
below the horizon. Testing commenced at 9:00pm, when the moon was 30º below the
horizon and we rotated the filter around to the future lunar e-vector direction (moonrise at
12:39am). If foragers were relying on a solar e-vector memory, we expected to see unequal
shifts between ±45° (smaller shifts in the +45º condition and larger shifts in the −45º
condition).
227
Waning lunar phases
While there is a consistent presence of polarised light during the waxing phase, during
waning lunar phases there is a gap which may impede this pattern's use as a compass cue.
Testing during the waxing lunar phase, as the moon's illuminated surface increases,
corresponds with periods in which the moon rises prior to sunset and sets overnight. This
creates a consistent presence of polarised light (solar then lunar) that foragers could use to
continuously update their celestial compass and path integrator. In contrast, the waning
lunar phase corresponds with the moon rising overnight, leading to a gap in this cue as solar
twilight ends. We hypothesised that the absence of the moon's presence as a cue overnight
might decrease its weighting or degrade its integration into the celestial compass. We added
two conditions to test this hypothesis: a *Waning Full Moon* and *Waning Quarter Moon*
conditions. Foragers were tested identical to previous conditions; only, they were tested
during moonrise (10pm and 1am) overnight rather than moonset.
241
Vector testing
After noticing differences in heading shift magnitude between nests which correlated with PI
vector lengths, we tested the hypothesis that, similar to solar polarisation (Freas et al.
2017b), the navigator's accumulated vector length impacts orientation to rotated lunar
polarised light. While foragers at Nest 1 (6.0m vector) reoriented almost fully, those at Nest

2 (3.1m vector) reoriented only halfway on near-full-moon nights. To test this hypothesis, we
collected foragers of Nest 1 on near (waxing) full moon nights and tested them at a site close
(2m) to their nest tree with either a 2.0m remaining vector, or with a larger (~6m) full
corridor remaining vector (*Halfway Collect & Release* or *Halfway Release* conditions). In the
Halfway Collect & Release condition, outbound foragers were collected at the halfway point,
held overnight and released at this site to be tested (±45° rotations) at ~2m from the nest. In
the *Halfway Release* condition, we followed outbound foragers to their foraging tree,
collected and held them until lunar twilight (2-4am) identically to previous full moon
conditions. These foragers were released along the route at the halfway point (3.0m) from
the nest/foraging tree and we tested each forager with their vector state near full but their
true location close (~2m) to the nest tree. In both conditions, we again recorded the initial
orientation, filter exit orientation and post-filter reorientation of each forager.
259
Statistical analysis
Data were analysed with circular statistics with the statistics package Oriana Version
4 (Kovach Computing Services). Each ant had a slightly different inbound heading due to
their stereotypical route along the foraging corridor and we corrected this variance by
designating the initial headings (pre-filter) as 0° for calculating shift magnitudes under the
filter. To assess shift magnitude between −45° and +45° foragers within conditions, we
calculated the mirror of shift in each −45° condition, allowing shift magnitude comparisons
within each condition. As −45° and +45° shifts were not significantly different in any
condition (Watson–Williams F-tests), they were combined for between condition
comparisons. Within-individual comparisons (Initial Orientation vs. Filter Exit and Filter Exit
vs. Reorientation) were analysed using Moore's Paired Tests. Across-condition shift
magnitudes were analysed using Watson–Williams F-tests. In the lunar phase comparisons
where full, quarter and crescent shift magnitudes, as well as Waning and Waxing phases
were compared, Holm-Bonferroni corrections were applied to the p-value to account for
multiple comparisons.
275
Results
Full moon testing
Under a (waxing) *Full Moon* with lunar illumination above 80%, when the linear polariser
was rotated clockwise (+45°), exit orientations were shifted to the right of initial headings
(mean ± s.e.m. Nest 1: 38.4±4.8°; Nest 2: 23.4±4.2°; Figure 2AB), and these changes were
significant (Moore's Paired Test, Nest 1: R=1.639, p<0.001; Nest 2: R=1.592, p<0.001).
Forager headings also changed predictably when the filter was rotated counter-clockwise (−
45°) with exit orientations to the left of initial headings (Nest 1: −41.1±5.5°; Nest 2: −
27.1±7.4°; Figure 2AB), and these changes were significant (Moore's Paired Test, Nest 1:
R=1.794, p<0.001; Nest 2: R=1.310, p<0.01). After exiting the +45° or −45º rotated filter,
foragers reoriented significantly to the left (Moore's Paired Test, Nest 1: R=1.598, p < 0.001;
Nest 2: R=1.383, p<0.005) or right respectively (Moore's Paired Test, Nest 1: R=1.604,

$p<0.001$; Nest 2: $R=1.328$, $p<0.005$). Shift magnitudes did not differ between +45° and −45°
conditions (Watson–Williams F-test, Nest 1: $F_{(1,23)}=0.155$, $p=0.697$; Nest 2: $F_{(1,20)}=0.234$,
$p=0.634$). Shift magnitudes were significantly larger at Nest 1 compared to Nest 2 (Watson–
Williams F-test, $F_{(1,45)}=8.672$, $p=0.005$), exhibiting shifts near the full e-vector change
(39.7±3.2°) while foragers at Nest 2 only exhibited shift magnitudes at about half the 45º
rotation (25.2±3.7°).

Waxing lunar phases
Under a *Waxing Quarter Moon* (lunar illumination ~50%), when the e-vector was
rotated clockwise (+45°), exit orientations were again significantly shifted (Moore's Paired
Test, $R=1.787$, $p<0.001$) to the right of their initial heading (mean ± s.e.m.; 37.6±4.1°; Figure
3A). Forager headings were similarly significantly altered (Moore's Paired Test, $R=1.734$,
$p<0.001$) when the overhead e-vector was rotated counter-clockwise (−45°) with exit
orientations to the left of initial headings (mean ± s.e.m. −38.5±6.4°; Figure 3A). After exiting
the filter in both conditions, foragers reoriented significantly back to the ambient lunar e-
vector either to the left (+45°: Moore's Paired Test, Nest 1: R=1.616, $p<0.001$) or right (−45°:
Moore's Paired Test, Nest 1: R=1.664, $p<0.001$) of their filter exit heading direction.
Foragers continued to show evidence of attending to the lunar polarisation pattern
even under a *Waxing Crescent Moon* (lunar illumination ~20%). Here, when the ambient e-
vector was rotated clockwise (+45°), exit orientations were again significantly shifted
(Moore's Paired Test, $R=1.175$, $p<0.025$) to the right of their initial heading (mean ± s.e.m.;
29.3±9.9°; Figure 3B). Forager headings were similarly significantly altered (Moore's Paired
Test, $R=1.39$, $p<0.005$) when the overhead e-vector was rotated −45º counter-clockwise,
with exit orientations to the left of initial headings (mean ± s.e.m.; −27.6±4.7°; Figure 3B).
After exiting the filter in both conditions, foragers reoriented significantly back to the
ambient lunar e-vector either to the left (+45°: Moore's Paired Test, Nest 1: R=1.324,
$p<0.005$) or right (−45°: Moore's Paired Test, Nest 1: R=1.223, $p<0.025$) of their filter exit
heading direction. Shift magnitudes were not significantly different between ±45° conditions
(Watson–Williams F-test, $F_{(1,20)}=0.03$, $p=0.863$).
When no ambient lunar e-vector was present (*No Moon*) and the polariser was
rotated either clockwise (+45°) or counter-clockwise (−45°), foragers did not significantly
alter their paths under the filter (+45°: Moore's Paired Test, $R=0.226$, $p>0.50$; mean ± s.e.m.:
−5.4±7.4°; −45°: Moore's Paired Test, $R = 0.650$, $p > 0.10$; mean ± s.e.m.: 0.7±7.6°; Figure 3C).
Foragers also did not significantly reorient after exiting the filter (+45°: Moore's Paired Test,
$R=0.294$, $p>0.50$; mean ± s.e.m.: −7.1±5.6°; −45°: Moore's Paired Test, $R=0.611$, $p>0.10$;
mean ± s.e.m.: 1.8±5.1°). Shift magnitudes were not significantly different between ±45°
conditions (Watson–Williams F-test, $F_{(1,24)}=0.016$, $p=0.899$).
When comparing shift magnitudes across lunar phases, *Full Moon f*oragers were not
significantly different from either the *Waxing Quarter Moon* or *Waxing Crescent Moon*
(Watson–Williams F-test, $p>0.05$; Figure 5).

Waning lunar phases

Under a *Waning Full Moon*, when the ambient e-vector was rotated clockwise (+45°), exit

orientations were shifted significantly to the right of initial headings (mean ± s.e.m. =

16.5±5.3°; Moore's Paired Test, R=1.468, p<0.005; Figure 4A). Headings also shifted

significantly to the left of initial headings (mean ± s.e.m.: −15.4±5.1°; Moore's Paired Test,

R=1.513, p<0.001; Figure 4A) when the overhead e-vector was rotated −45º counter-

clockwise. After exiting the filter in both conditions, foragers reoriented significantly back to

the ambient lunar e-vector either to the left (+45°: Moore's Paired Test, Nest 1: R=1.548,

p<0.001) or right (−45°: Moore's Paired Test, Nest 1: R=1.247, p<0.025) of their filter exit

heading direction. Shift magnitudes were not significantly different between the ±45°

conditions (Watson–Williams F-test, $F_{(1,30)}$=0.022, p=0.884).

Under a *Waning Quarter Moon* (lunar illumination ~50%), when the ambient e-

vector was rotated clockwise (+45°), exit orientations were shifted to the right of initial

headings (mean ± s.e.m. = 20.5±8.4°; Figure 4B), and these changes were significant

(Moore's Paired Test, R=1.33, p<0.005). Headings also changed significantly (Moore's Paired

Test, R=1.31, p<0.01) when the overhead e-vector was rotated counter-clockwise (−45°) with

exit orientations to the left of initial headings (mean ± s.e.m.: −22.5±11.7°; Figure 4B). After

exiting the filter in both conditions, foragers reoriented significantly back to the ambient

lunar e-vector either to the left (+45°: Moore's Paired Test, Nest 1: R=1.504, p<0.001) or

right (−45°: Moore's Paired Test, Nest 1: R=1.246, p<0.025) of their filter exit heading

direction. Shift magnitudes were not significantly different between the ±45° conditions

(Watson–Williams F-test, $F_{(1,18)}$=0.154, p=0.700).

When comparing shift magnitudes between Waxing and Waning phases, shift

magnitude was significantly higher in both *Waxing Full Moon* and *Waxing Quarter Moon*

foragers when compared to their *Waning* counterparts (39.8º vs. 16.0º and 38.1º vs. 22.7º

respectively; Full Moon: Watson–Williams F-test, $F_{(1,55)}$=21.62, p<0.001; Quarter Moon:

Watson–Williams F-test, $F_{(1,44)}$=5.889, p=0.038; Figure 5).

Vector testing

Both *Halfway Release* (~6m vector) and *Halfway Collection & Release* (~2m vector) foragers

exhibited significant shifts (*Halfway Release*: Moore's Paired Test, R=1.728, p<0.001; *Halfway*

Collection & Release: Moore's Paired Test, R=1.380, p<0.005) to the right of initial headings

(mean ± s.e.m.: 41.9±4.9° and 6.8±4.3°; Figure 6AB) when the ambient e-vector was rotated

clockwise (+45°). Forager headings also significantly shifted (*Halfway Release*: Moore's

Paired Test, R=1.664, p<0.001; *Halfway Collection & Release:* Moore's Paired Test, R=1.07,

p<0.05) when the overhead e-vector was rotated counter-clockwise (−45°) with exit

orientations to the left of initial headings (mean ± s.e.m.: −46.4±6.3° and 12.7±8.5°; Figure

6AB). After exiting the filter in both conditions, foragers reoriented significantly back to the

ambient lunar e-vector either to the left (+45º: *Halfway Release*: Moore's Paired Test,

R=1.692, p<0.001; *Halfway Collection & Release:* Moore's Paired Test, R=1.600, p<0.001) or

right (−45°: *Halfway Release*: Moore's Paired Test, R=1.604, p<0.001; *Halfway Collection &*

Release: Moore's Paired Test, *R*=1.274, p<0.01) of their exit heading direction. Shift

magnitude was significantly higher in *Halfway Release* foragers compared to *Halfway*

Collection & Release foragers tested at the same location 2m from the nest entrance (44.1º

and 14.8º respectively; Watson–Williams F-test, $F_{(1,40)}$ = 29.105, p < 0.001; Figure 6AB).

Discussion

These results constitute the first instance of polarised moonlight use for homing and only

the second reported instance of its use for orientation in any animal (Figure 7). *Myrmecia*

midas foragers predictably altered their heading directions in response to experimental

rotations in the ambient overhead lunar polarisation pattern. This ability to detect and

attend to polarised moonlight persisted throughout the lunar cycle, with foragers attending

to the pattern even under a crescent moon with ~20% lunar illumination. This indicates that

polarised moonlight is detectable across the lunar month, making it a stable cue these ants

can use when moving or updating their path integrator overnight.

384   While they can utilise the lunar polarisation pattern through the lunar cycle, foragers

exhibited reduced heading shifts during the waning lunar phases, during which the moon's

absence for a portion of the night leads to a cue gap overnight. These reductions in observed

shifts are likely due to either decreased weighting or a degradation in the celestial compass

due to periods when no celestial cues are available. The shift magnitude differences

between conditions also point to moonlight polarisation being continuously tracked

throughout the overnight period, in line with this celestial cue being integrated into the

navigator's path integrator. Further evidence of this cue's integration into the path integrator

is illustrated in our *Halfway* testing conditions. Here, foragers with longer home vectors

under a full moon responded almost fully to e-vector changes regardless of their distance to

the nest, while short vector distances led to less than half the e-vector shift, corresponding

with decreases in vector cue strength at small distances (Wystrach et al. 2015). These

changes align exactly with how *M. midas*, uses polarised sunlight as part of its vector-based

homing during evening and morning twilight (Freas et al. 2017b; Figure 7), suggesting that

polarised moonlight is detected and integrated into its path integrator along the same visual

pathways as sunlight.

Moonlight vs. sunlight

The nocturnal bull ants *Myrmecia pyriformis* and *Myrmecia midas* are known to use the

solar polarised light pattern during the twilight periods (Reid et al. 2011; Freas et al.

2017ab), yet both species are active after twilight, when solar polarisation cues are absent

(Reid et al. 2011, 2013; Freas et al. 2017ab). The observed true nocturnal navigation in these

animals could be driven by the moon's presence with *M. pyriformis*, showing more foraging

activity on full moon nights (Reid et al., 2013). *M. midas* exhibits a high level of overnight

activity, with almost half of foragers returning before the morning twilight. *M midas* also

navigates through heavily canopied forest habitat where the moon may be occluded but its

polarisation pattern across the sky remains unobstructed. Thus, *M. midas* makes for an
interesting species to characterise lunar and solar polarised light detection.
While we cannot compare solar and moonlight polarisation navigation in outbound
ants (outbound foraging is highly correlated with evening twilight when the solar light would
overpower any moonlight polarisation pattern), striking similarities occur when comparing
solar and moonlight polarisation navigation in ants homing to the nest (Figure 7). In the solar
polarisation study (Freas et al. 2017b), inbound foragers tested during morning twilight at 4–
6 m from their nest altered their paths under the filter for almost the full 45º solar e-vector
manipulation (−45° rotation: −41.16°/ +45° rotation: 34.13°; Figure 7) but only compensated
for around half of the rotation when tested 1–2 m from the nest (−45° rotation: −24.86°/
+45° rotation: 19.73º). We see the same pattern with polarised moonlight with foragers
exhibiting near full shifts 5m from Nest 1 (−45° rotation: −41.4°/ +45° rotation: 38.4°) and
half shifts 2m from Nest 2 (−45° rotation: −27.1°/ +45° rotation: 23.4°; Figure 7).
We also observed consistent slight under-estimation in the shifts even when foragers
had a long vector. Observations of ants after the filter was placed overhead suggest that
heading updates are not immediate, occurring only after the ant travels along its original
heading a few centimetres (~5cm). This means that even if the ant fully shifts its heading,
the delay will cause our measurements at filter exit to slightly underestimate each
individual's position since we measure the angle from where the lunar cue changes (as filter
was placed overhead) and not the position at which the ant altered its heading.
430
Moonlight and the path integrator
We also see the same full heading changes to both solar- and moonlight-polarisation-
pattern rotations when foragers were released halfway to the nest and tested 2m with a
larger vector (Solar: −45° rotation: −35.77°/ +45° rotation: 39.42°; Moonlight: −45° rotation:
−46.4°/ +45° rotation: 41.9; Figure 7). In the *Halfway collect & Release* condition, this shift
significantly decreased at the same testing location, indicating that the vector length and not
the testing site dictated this adjustment in cue weighting. These findings suggest that under
the filter, foragers use any available celestial and terrestrial cues that are still visible, yet the
weighting of the polarisation pattern appears to change in accordance with the vector state
(Burkhalter 1972; Narendra 2007) and not its test locaiton close to the nest tree, which
could be a potentially highly salient landmark. This leads to several interesting implications.
First, these ants weight the polarisation cue more highly and perform larger heading shifts
when their current path integrator distance is longer. These increased shift magnitudes align
with the hypothesis that longer accumulated vectors, ants increase the weighting given this
cue (Burkhalter 1972; Narendra 2007; Wystrach et al. 2015; Freas et al. 2017b). Secondly,
these ants are using polarised moonlight precisely the same way they use solar polarisation,
meaning that just as polarised moonlight is likely integrated into the forager's path
integrator throughout the night. Moonlight polarised light is likely processed through the
same visual pathways as polarised sunlight, meaning that these ants can use the same
underlying neural architecture for polarised solar light and polarised moonlight cues. The

only limiting factor would be the navigator's detection threshold (at least 20% lunar
luminance), which can be accomplished through the specialised eye morphology of these
ants (Zeil et al. 2014).
454
Polarised moonlight and lunar phase
Foragers showed clear evidence of detecting and employing polarised moonlight when
homing to the nest across the lunar cycle, even on waxing crescent moon nights. This aligns
with polarised moonlight's use in dung beetles, with individuals able to maintain their
straight-line paths under quarter and crescent moon e-vectors (Dacke et al. 2004).
Furthermore, the lack of a shift-magnitude reduction between full and crescent nights
suggests no reduction in detection. While we could have continued to test with smaller
portions of the moon surface illuminated, a reduction in shift magnitude could result from
either physiological limits in detection or 'decisional' processes in how much weight to
accord the cue. Behavioural responses and physiological limits cannot be untangled
behaviourally, and detection thresholds would require intracellular recordings under dim
polarised light. Finally, the lack of shifts with *No Moon* foragers indicates that these
navigators do not fall back on memories of the evening or morning solar e-vector when
presented one overnight.
One unexpected finding was the reduction in shift magnitude under waning moons
relative to waxing moons. When we first tested foragers under *Waning Full Moons* and
Waning *Quarter Moons* they showed clear evidence of attending to the polarisation cue, but
shift magnitudes were significantly smaller compared to the *Waxing Moon* conditions. This
reduction in shift magnitude suggests that polarised moonlight was being detected but it
was weighted weakly, perhaps due to the celestial compass incorporation into the path
integrator being interrupted. This observed reduction in shift magnitude likely results from
the period overnight in which the waning moon was absent from the night sky, meaning
foragers could not attend to a consistent polarisation pattern throughout the night. The
moonlight pattern only becomes visible in the sky once the moon reaches −18º below the
horizon; thus, waning-moon nights present periods when there are neither solar nor lunar
cues available to maintain the compass, potentially degrading the compass portion of the
vector estimate. This suggests that when the moon is waxing and present throughout the
overnight period (at least until moonset), *M. midas* foragers are continuously tracking it and
integrating this compass cue into their path integrator. When there are periods overnight
when the moon and its polarisation pattern are absent, it is either weighted weakly or it
becomes misaligned with the true homing direction since the celestial compass
incorporation into the path integrator has been interrupted with overnight periods with no
detectable celestial cues.
488
Conclusions
In the first evidence of polarised moonlight being used to navigate to a goal location,
inbound *M. midas* foragers detect and respond predictably to rotations of the moonlight e-

vector orientation under a filter and reorient back to the ambient e-vector after filter exit.
This ability occurs across lunar phase, suggesting that polarised moonlight is a detectable
cue throughout the lunar month. Heading changes due to polarised moonlight align with
responses to polarised sunlight as part of the path integrator during solar twilight. This
indicates that polarised moonlight is likely detected and integrated into the ant's path
integrator for inbound homing along the same visual pathways as polarised sunlight.
Reductions in heading shifts due to differences in PI vector lengths, and periods without
access to polarised light patterns suggest that these animals can weight the information
provided by celestial polarised light. In so doing these foragers can cater their navigational
decisions proportionate to closely match the reliability of available navigational information.
502

Acknowledgments
Funding
This project was funded by a Macquarie University Research Fellowship (MQRF0001094), by
Macquarie University, and by an ARC Discovery Grant (DP200102337).
Land Acknowledgment
This work was conducted upon the Wallumattagal campus of Macquarie University. We
acknowledge the traditional custodians of the land on which Macquarie University sits, the
Wallumattagal clan of the Dharug Nation. Their culture and customs have nurtured and
sustained this land since the Dreamtime and continue to do so. We pay our respects to their
Elders, past, present, and future.
Conflicts of interest
The authors declare no conflicts of interest associated with this work.
Ethics
There are no state or federal governmental regulations guiding the research of
invertebrates in Australia.
Data Availability
All data, documentation and code will be made available online at osf.io

References
Burkhalter, A. (1972). Distance measuring as influenced by terrestrial cues in *Cataglyphis*
bicolor. In Information Processing in the Visual System of Arthropods (ed. R. Wehner),
pp. 303-308. Berlin: Springer.
Dacke, M., Byrne, M. J., Baird, E., Scholtz, C. H., & Warrant, E. J. (2011). How dim is dim?
Precision of the celestial compass in moonlight and sunlight. Philosophical
Transactions of the Royal Society B: Biological Sciences, 366(1565), 697-702.

Dacke, M., Byrne, M. J., Scholtz, C. H., & Warrant, E. J. (2004). Lunar orientation in a beetle.
Proceedings of the Royal Society of London. Series B: Biological Sciences, 271(1537),
361-365.
Dacke, M., Nilsson, DE., Warrant, E. et al. Built-in polarizers form part of a compass organ in
spiders. Nature, 401, 470–473 (1999). https://doi.org/10.1038/46773
Dacke, M., Nilsson, D. E., Scholtz, C. H., Byrne, M., & Warrant, E. J. (2003). Insect orientation
to polarized moonlight. Nature, 424(6944), 33-33.
Dacke, M., el Jundi, B., Smolka, J., Byrne, M., & Baird, E. (2014). The role of the sun in the
celestial compass of dung beetles. Philosophical Transactions of the Royal Society B:
Biological Sciences, 369(1636), 20130036.
el Jundi, B., Warrant, E. J., Byrne, M. J., Khaldy, L., Baird, E., Smolka, J., & Dacke, M. (2015).
Neural coding underlying the cue preference for celestial orientation. Proceedings of
the National Academy of Sciences, 112, (36), 11395–11400.
Foster, J. J., Kirwan, J. D., El Jundi, B., Smolka, J., Khaldy, L., Baird, E., & Dacke, M. (2019).
Orienting to polarized light at night–matching lunar skylight to performance in a
nocturnal beetle. Journal of Experimental Biology, 222(2), jeb188532.
Freas, C. A., & Cheng, K. (2019). Panorama similarity and navigational knowledge in the
nocturnal bull ant *Myrmecia midas*. Journal of Experimental Biology, 222(11),
jeb193201.
Freas, C. A., & Cheng, K. (2022). The basis of navigation across species. Annual Review of
Psychology, 73, 217-241.
Freas, C. A., & Spetch, M. L. (2023). Varieties of visual navigation in insects. Animal cognition,
26(1), 319-342.
Freas, C. A., Narendra, A., & Cheng, K. (2017a). Compass cues used by a nocturnal bull ant,
Myrmecia midas. Journal of Experimental Biology, 220(9), 1578-1585.
Freas, C. A., Narendra, A., Lemesle, C., & Cheng, K. (2017b). Polarized light use in the
nocturnal bull ant, *Myrmecia midas*. Royal Society Open Science, 4(8), 170598.
Freas, C. A., Plowes, N. J., & Spetch, M. L. (2019). Not just going with the flow: foraging ants
attend to polarised light even while on the pheromone trail. Journal of Comparative
Physiology A, 205(5), 755-767.
Freas, C. A., Wystrach, A., Narendra, A., & Cheng, K. (2018). The view from the trees:
nocturnal bull ants, Myrmecia midas, use the surrounding panorama while descending
from trees. Frontiers in Psychology, 9, 16.
Gál, J., Horváth, G., Meyer-Rochow, V. B., & Wehner, R. (2001). Polarization patterns of the
summer sky and its neutral points measured by full–sky imaging polarimetry in Finnish
Lapland north of the Arctic Circle. Proceedings of the Royal Society of London A,
457(2010), 1385-1399.
Greiner, B., Cronin, T. W., Ribi, W. A., Wcislo, W. T., & Warrant, E. J. (2007). Anatomical and
physiological evidence for polarisation vision in the nocturnal bee *Megalopta genalis*.
Journal of Comparative Physiology A, 193, 591-600.
Herzmann, D., & Labhart, T. (1989). Spectral sensitivity and absolute threshold of
polarization vision in crickets: a behavioral study. Journal of Comparative Physiology A,
165, 315-319.
Homberg, U., & Paech, A. (2002). Ultrastructure and orientation of ommatidia in the dorsal
rim area of the locust compound eye. Arthropod Structure & Development, 30(4), 271-
280.

Horváth, G., & Varjú, D. (2004). Polarized light in animal vision: polarization patterns in
nature. Springer Science & Business Media.
Horváth, G. Lerner, A., & Shashar, N. (2014). Polarized light and polarization vision in animal
sciences (Vol. 2,). G. Horváth (Ed.). Berlin: Springer.
Jander, R. (1957) Die optische Richtungsorientierung der Roten Waldameise (Formica rufa L.)
Zeitschrift tiir vergMehende Physiologie Bd, 40, 162-238.
Klotz, J. H., & Reid, B. L. (1993). Nocturnal orientation in the black carpenter ant
Camponotus pennsylvanicus (DeGeer)(Hymenoptera: Formicidae). Insectes Sociaux,
40, 95-106.
Labhart, T., & Meyer, E. P. (1999). Detectors for polarized skylight in insects: a survey of
ommatidial specializations in the dorsal rim area of the compound eye. Microscopy
research and technique, 47(6), 368-379.
Lebhardt, F., & Ronacher, B. (2014). Interactions of the polarization and the sun compass in
path integration of desert ants. Journal of Comparative Physiology A, 200, 711-720.
Narendra, A. (2007). Homing strategies of the Australian desert ant *Melophorus bagoti II*.
Interaction of the path integrator with visual cue information. Journal of Experimental
Biology, 210(10), 1804-1812.
Narendra A, Kamhi JF & Ogawa Y. 2017. Moving in dim light: behavioural and visual
adaptations in nocturnal ants. Integrative and Comparative Biology, 57, 1104-1116.
Perez, S., Taylor, O. & Jander, R. 1997. A sun compass in monarch butterflies. Nature, 387, 29.
https://doi.org/10.1038/387029a0
Reid, S. F., Narendra, A., Hemmi, J. M., & Zeil, J. (2011). Polarised skylight and the landmark
panorama provide night-active bull ants with compass information during route
following. Journal of Experimental Biology, 214(3), 363-370.
Reid SF, Narendra A, Taylor RW & Zeil J. 2013. Foraging ecology of the night-active bull
ant, Myrmecia pyriformis. Australian Journal of Zoology, 61, 170-177.
Rost, R., & Honegger, H. W. (1987). The timing of premating and mating behavior in a field
population of the cricket *Gryllus campestris L*. Behavioral Ecology and Sociobiology, 21,
279-289.
Ugolini, A., Galanti, G., & Mercatelli, L. (2013). Do sandhoppers use the skylight polarization
as a compass cue?. Animal Behaviour, 86(2), 427-434.
Warrant, E., & Dacke, M. (2016). Visual navigation in nocturnal insects. Physiology, 31(3),
182-192.
Wehner, R., & Müller, M. (2006). The significance of direct sunlight and polarized skylight in
the ant's celestial system of navigation. Proceedings of the National Academy of
Sciences, 103(33), 12575-12579.
Wehner, R. and Srinivasan, M.V. (2003). Path integration in insects. In: Jeffery KJ (ed) The
neurobiology of spatial behaviour. Oxford University Press, Oxford, pp 9–30.
Wystrach, A., Mangan, M., & Webb, B. (2015). Optimal cue integration in ants. Proceedings
of the Royal Society B, 282(1816), 20151484.
Zeil J, Ribi WA, Narendra A. 2014 Polarization vision in ants, bees and wasps. In Polarized
light and polarization visionin animal sciences, 2nd edn (ed.Gábor Horváth), pp. 41–60.
Berlin, Germany:Springer Series in Vision Research.
617
618
619
620

  Figure Captions

Figure 1. Diagram of the polarisation filter and changes to the ambient lunar polarisation
pattern. (A) During the inbound journey, a linear polarisation filter was placed over the
forager, rotating the overhead e-vector by ±45°. Panel depicts the positional measurements
recorded during testing. Initial orientation routes were measured from the foraging tree
release point (a) to when the polarisation filter was placed over each forager (b). Exit
orientations were measured from the filter centre (b) to the forager's exit location at the
filter edge (c). Route directions under the filter (b°) were calculated from the forager's initial
route direction zeroed. Reorientations were measured from the forager's exit location from
the polarisation filter (c) to the forager's path 1m after exit (d). Reorientation directions (c°)
were calculated from the under-filter route direction zeroed. (B) Images of the sky and
canopy cover at both nests. Photos were taken at the on-route midpoint between the
foraging and nest trees. (C) Lunar phases denote the sunlit part of the moon's surface and
where this area is increasing (Waxing) or decreasing (Waning). The lunar phase cycle repeats
every 29.5 days. moon images are public domain art accessed through wiki commons
(https://commons.wikimedia.org/).

639
Figure 2. Circular distributions of headings during the original full moon conditions. In both
conditions, testing occurs in the nights preceding the full moon (illumination > 80%) with the
moon waxing. Circular plot shifts show the exit orientations of individual foragers from their
initial headings while the reorientation represents the change in headings 1m after exiting
the filter. Triangles denote the ±45° e-vector rotation. The arrow denotes the length and
direction of the mean vector. (*A*) Nest 1 foragers, 5m from the nest (6.0m foraging route). (*B*)
Nest 2 foragers, 2m from the nest (3.1m foraging route). *n*, number of individuals; Ø, mean
vector; *r*, length of the mean vector.

648
649
650
651

652
653
Figure 3. Circular distributions of headings during (A) Waxing *Quarter Moon,* (B*) Waxing*
Crescent Moon and (C) *No Moon* conditions. Circular plot shifts show the exit orientations of
individual foragers from their initial headings while the reorientation represents the change
in headings 1m after exiting the filter. Triangles denote ±45° e-vector rotation. The arrow
denotes the length/direction of the mean vector. *n*, number of individuals; Ø, mean
vector; *r*, length of the mean vector.

Figure 4. Circular distributions of headings during (A) Waning Full *Moon,* (B*) Waning Quarter Moon* conditions. Circular plot shifts show the exit orientations of individual foragers from their initial headings while the reorientation represents the change in headings 1m after exiting the filter. Triangles denote ±45° e-vector rotation. The arrow denotes the length/direction of the mean vector. *n*, number of individuals; *Ø*, mean vector; *r*, length of the mean vector.

Figure 5. Shift magnitudes for lunar phase conditions at Nest 1. Each circular plot shows the ±45° combined shifts for each condition. Triangles denote +45° e-vector rotation; data from −45º were mirrored and combined). The arrow denotes the length and direction of the mean vector. *n*, number of individuals; *Ø*, mean vector; *r*, length of the mean vector.

685
Figure 6. Circular distributions of *M. midas* headings during (A) *Released Halfway, (*B*)*
Collected & Released Halfway conditions. Foragers in both conditions were tested at 2m
from the nest with *Released Halfway* foragers having a long 6.0m vector and *Collected &*
Released Halfway foragers having a 3.1m vector. Circular plot shifts show the exit
orientations of individual foragers from their initial headings while the reorientation
represents the change in headings 1m after exiting the filter. Triangles denote ±45° e-vector
rotation. The arrow denotes the length and direction of the mean vector. *n*, number of
individuals; Ø, mean vector; *r*, length of the mean vector.

699
700
701 **Figure 7.** Mean shift magnitudes, +45º and −45º combined, reported as percentages of the
702 45º e-vector rotation (100% = 45º). Vector data for solar polarised light is reported from
703 Freas et al. 2017, Royal Society Open Science. Sun and moon images are public domain art
704 accessed through wiki commons (https://commons.wikimedia.org/).

A

Ambient Angle of Lunar Polarization

Initial Heading

Filter Polarization

45°

1m

b

b°

15cm

a

Initial Heading

a°

Reorientation

1m

d

c°

c

Exit Heading

B

Foraging Tree

N

Nest 1

Foraging Tree

N

Nest 2

C

Lunar Phases Southern Hemisphere (Sydney, AU)

New Moon   Waxing Crescent   Waxing Quarter   Full Moon   Waning Quarter   Waning Crescent   New Moon