Identification of nonsense-mediated decay inhibitors that alter the tumor immune landscape

Ashley L Cook[1,2], Surojit Sur[1,3,4], Laura Dobbyn[3], Evangeline Watson[1], Joshua D Cohen[1,3,4], Blair Ptak[3], Bum Seok Lee[3], Suman Paul[1,3,4], Emily Hsiue[1], Maria Popoli[1], Bert Vogelstein[1,2,3,4,5,6], Nickolas Papadopoulos[1,3,4], Chetan Bettegowda[1,2,3,4,7], Kathy Gabrielson[3,4,8], Shibin Zhou[1,3,4], Kenneth W Kinzler[1,2,3,4,6,]*, Nicolas Wyhs[1,3,4,9,]*

Affiliations

[1] Ludwig Center for Cancer Genetics and Therapeutics, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA.

[2] Cellular and Molecular Medicine Graduate Program, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA.

[3] Department of Oncology, Johns Hopkins Medical Institutions, Baltimore, MD 21287, USA.

[4] Sidney Kimmel Cancer Center, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA.

[5] Howard Hughes Medical Institute, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA.

[6] Sol Goldman Pancreatic Cancer Research Center, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA.

[7] Department of Neurosurgery, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA.

[8] Department of Molecular and Comparative Pathobiology, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA.

[9] Lead contact

* Correspondence: wyhs@jhmi.edu, kinzlke@jhmi.edu

Summary

Despite exciting developments in cancer immunotherapy, its broad application is limited by the paucity of targetable antigens on the tumor cell surface. As an intrinsic cellular pathway, nonsense-mediated decay (NMD) conceals neoantigens through the destruction of the RNA products from genes harboring truncating mutations. We developed and conducted a high throughput screen, based on the ratiometric analysis of transcripts, to identify critical mediators of NMD. This screen revealed disruption of kinase SMG1's phosphorylation of UPF1 as a potent disruptor of NMD. This led us to design a novel SMG1 inhibitor, KVS0001, that elevates the expression of transcripts and proteins resulting from truncating mutations *in vivo* and *in vitro*. Most importantly, KVS0001 concomitantly increased the presentation of immune-targetable HLA class I-associated peptides from NMD-downregulated proteins on the surface of cancer cells. KVS0001 provides new opportunities for studying NMD and the diseases in which NMD plays a role, including cancer and inherited diseases.

One Sentence Summary

Disruption of the nonsense-mediated decay pathway with a newly developed SMG1 inhibitor with *in-vivo* activity increases the expression of T-cell targetable cancer neoantigens resulting from truncating mutations.

Key Words

SMG1; nonsense-mediated decay; small molecule; high throughput screen; cancer; immunotherapy; drug repurposing

Introduction

Despite success with cancer immunotherapies, approved immunotherapies are not available for the majority of cancer patients and only a minority of treated patients realize a durable response[1-4]. Current studies largely focus on discovering new agents and identifying patients most likely to benefit from existing immunotherapies[5]. While many studies correlate tumor insertion and deletion (indel) mutation load with immunotherapeutic response, not all tumors with high indel mutational loads respond to checkpoint inhibitors[6-13].

The typical adult solid tumor contains a median of 54 coding somatic nucleotide variants, many

of which have the potential to create novel neoantigens or Mutation Associated NeoAntigens (MANA)[14-

16]. Approximately 5% of mutations are insertions/deletions (indels) or splice site changes that alter the

open reading frame (ORF) of the transcript[14-16]. This subset of mutations is of particular interest because

they can result in proteins and derived peptides that are foreign to a host's healthy cells, giving rise to

neoantigens[17-19]. In normal cells, nonsense-mediated decay (NMD) plays an important role in messenger

RNA (mRNA) quality control, as well as normal gene expression[20,21]. In cancer cells, however, NMD

may aid immuno-evasion by eliminating RNA transcripts coming from genes that carry truncating

mutants[19]. This prevents translation and presentation of peptides from these proteins on MHC class I

complexes, rendering them invisible to immune cells[22]. Previous work has shown knockdown of the

NMD pathway with siRNA enhances the anti-tumor immune response[18,23]. Analysis of cell lines with loss

of the UPF1 RNA helicase and ATPase gene (*UPF1*), a key mediator of NMD degradation, also showed

increased levels of aberrant transcripts and mutant proteins in the alleles targeted by NMD[24]. This is

reminiscent of the pharmacological modulation of splicing, which has also been shown to increase the

number of neoantigens present on the cancer cell surface due to similar underlying mechanisms[25]. Despite

early studies showing little toxicity with NMD inhibition, there is as of yet no reports of a specific

chemical inhibitor of the pathway with good bioavailability[26-30].

In this work we create a cell-based high throughput assay to query the effects of a curated library

of small molecules on NMD function. From this screen, we identify a lead compound capable of

inhibiting NMD and ascertain its protein target as nonsense mediated mRNA decay associated PI3K

related kinase (SMG1), a kinase that is a critical mediator of NMD. As the lead compound produced

unacceptable toxicity in animal models, we then designed a specific and bioavailable small molecule

inhibitor of SMG1 (KVS0001) that is well tolerated *in vivo*. We then demonstrate that targeted inhibition

of SMG1 by KVS0001 leads to the presentation of novel neoantigens identifiable by T cells leading to

tumor growth inhibition *in vitro* and *in vivo*.

Results

Development of a high throughput assay to identify NMD inhibitors

To develop an assay to find NMD inhibitors, we identified isogenic cell lines with out-of-frame

indel mutations, hereinafter referred to as truncating mutations, targeted by NMD activity. We previously

reported a panel of non-cancerous cell lines in which 19 common tumor suppressor genes were

inactivated using the Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR)-Cas9

system[31]. Through evaluation of this panel, we discovered two genes, Stromal Antigen 2 (*STAG2*) and

Tumor Protein p53 (*TP53*), which did not express their expected proteins when assessed by western blots

(STAG2) or immunohistochemistry (p53) (Figure S1). Because the inactivation of these two genes was

the result of frame-shift mutations, we suspected that the absence of the proteins was due to NMD. This

suspicion was supported by whole transcriptome RNA-sequencing (Figure S2)[31]. Notably we saw an

average decrease of STAG2 RNA transcripts by 20-fold and TP53 by 6-fold relative to their respective

wild-type transcripts in the parental cell lines.

We selected two *STAG2* knockout clones (clone 2 and 8) and one *TP53* knockout clone (clone

221) derived from the Retinal Pigmented Epithelial (RPE1) cell line to design a high-throughput screen

(HTS). Treatment with the canonical NMD inhibitor emetine demonstrated up to a 60-fold recovery of

mutant RNA transcript expression, establishing this panel as appropriate for HTS (Figure S3)[32].

We then designed a Next Generation Sequencing (NGS) assay for NMD (Figure 1A). We mixed

the three cell lines in equal proportions and plated the mixture in 96-wells plates followed by treatment

with one compound (drug) per well. We determined NMD inhibition efficacy by comparing the wild-type

and truncating mutant transcript expression levels in a ratiometric manner. Specifically, the wild-type

sequences of the reciprocally knocked out clone (i.e., wild-type *STAG2* sequence from the *TP53* knockout

clones, and wild-type *TP53* sequence from the *STAG2* knockout clones) served as internal references,

providing a ratiometric assay of mutant to wild-type transcript abundance. This ratiometric assay

minimized confounders introduced by nonspecific transcriptional activators or generally toxic agents.

The use of three cell lines with different truncating mutations from two different target genes minimized

the possibility that drugs identified in the HTS were cell-line clone or mutation-specific. Note that the use

of these cell lines, carefully mixed, banked, and preserved, did not substantially increase the amount of

time or work required to screen a single cell line. A combination of well and plate barcodes allowed the

pooling and scoring of over 1920 assays in a single NGS lane (see methods).

Execution of a high throughput screen to identify NMD inhibitors

Previous human clinical trials suggest that off-target toxicity at doses required for NMD

inhibition makes emetine and other well-known NMD inhibitors unsuitable for human use[33-37]. We

performed a high throughput screen (HTS) to identify more specific NMD inhibitors by treating the

isogenic cell line panel described above with a commercially available library consisting of 2,658 FDA-

approved or in late-phase clinical trial small molecules and natural products (Table S1). After purifying

RNA from cells 16 hours post-treatment, we scored NMD inhibition using the strategy described in

Figure 1A and Figure S4. Predictably, emetine increased the relative expression of mutant to wild-type

transcripts by 3 to 4 fold on average (Figure S5). Eight compounds (0.3% of the library) increased the

ratiometric mutant transcript fraction more than five standard deviations above the DMSO controls in all

three cell lines (Figure 1B). This hit threshold was chosen as it was the minimum required to ensure no

false positives were observed in the DMSO controls. One of these eight compounds, anisomycin, is a

known inhibitor of protein synthesis and commonly used NMD inhibitor for *in vitro* studies, providing

independent validation of the screen[32]. The other seven compounds increased mutant RNA transcript

levels 5 to 10-fold relative to untreated cells.

LY3023414 inhibits NMD and causes re-expression of mutant RNA and protein

To validate the eight hit compounds described above, we tested their effects in additional lines

with mutations targeted by NMD. First, we assessed them on isogenic *STAG2* and *TP53* knockouts in

RPtec cells, another non-cancerous cell line (Figures S6A, S6B, and S7)[31]. Four of the original eight hit

compounds increased mutant RNA expression in a dose-dependent manner (Figures 1C and S8). Next, we

examined the effects of these four compounds on additional RPE1 TP53 knockout cell lines with different

mutations predicted to generate truncated TP53 proteins (Figures S6C and S7). While treatment with all

four potential NMD inhibitors restored expression of the truncated mutant TP53 proteins, two of them

(LY3023414 and halofuginone) did so most robustly (Figures 1D and 1E). Additionally, *TP53* has an

isoform, TP53β, whose expression is known to be controlled by the NMD pathway[38]. Using cell lines

with intact TP53β isoform transcripts, we observed an increase in the TP53β isoform in cell lines treated

with both LY3023414 and halofuginone (Figure 1E middle and left column). Quantitative real time PCR

(qPCR) of both the full-length TP53 (TP53α) and NMD sensitive (TP53β) transcripts in parental RPE1

and RPTec cells treated with LY3023414 or halofuginone confirmed the selective upregulation of TP53β,

but not TP53α, transcripts (Figure 1F). Based on a consistently stronger effect of LY3023414 over

halofuginone across multiple isogenic cell lines and assays, LY3023414 was chosen to be the initial lead

compound for further studies involving NMD inhibition.

Figure 1: LY3023414 is a small molecule capable of increasing transcription of NMD targets.
A) Schematic of HTS used to identify inhibitors of NMD. Mutant transcripts are represented by a smaller length in the cartoon for illustrative purposes only. **B)** Mutant RNA reads relative to wild-type reads for the top 8 hits from the HTS. The dotted line represents the minimum fraction required to be considered a hit (> 5 standard deviations above DMSO control). Full screen results are presented in Figure S4. **C)** Targeted RNA sequencing results of isogenic RPTec knockout clones treated with the 8 best hits from the HTS at 10 µM. The dotted line represents a relative RNA expression level of 1, equal to that of DMSO treated wells. Data for ceritinib, which did not validate on any line, is presented only in Figure S8. **D)** TP53 western blot on RPE TP53 224, containing a homozygous *TP53* mutation, using the four hit compounds that validated in RPTec isogenic lines. **E)** Western blot analysis of full-length TP53α and isoform TP53β after treatment with two NMD inhibitor lead candidates. TP53β (expression known to be controlled by NMD) as well as mutant TP53 are prominently induced by LY3023414 whereas full-length is not. Note that RPE TP53 223 is a heterozygous knockout clone with one near wild-type allele whereas RPTec TP53 588 contains a homozygous *TP53* indel mutation. **F)** qPCR showing LY3023414 treatment causes increased expression of the NMD controlled alternative transcript for TP53, TP53β, in parent cell lines for RPE1 and RPTec. Significance determined by Student's t-test. Unless indicated otherwise cells were exposed to 5 µM of test compound for 16 hours.

LY3023414 increases expression of NMD repressed mutant RNA transcripts and proteins *in vitro*

and *in vivo*.

To evaluate whether LY3023414 could relieve NMD repression of naturally occurring heterozygous mutant transcripts, we chose the NCI-H358 and LS180 cancer cell lines. Treatment with LY3023414 followed by whole transcriptome RNA sequencing revealed increased expression of the mutant allele in 42% and 67% of heterozygous, out-of-frame, indel mutations in these two lines (Figure 2A). A third of the nonsense mutations in NCI-H358 were also "recovered" (i.e., mutant transcripts increased relative to wildtype transcripts) after treatment with LY3023414 (Figure 2A middle). In LS180 there was only one nonsense mutation meeting the required coverage, so it could not be evaluated in depth. Single base pair substitutions not resulting in stop codons were not affected by LY3023414 in either line (Figure 2A right).

The whole transcriptome sequencing data was confirmed by targeted deep RNA sequencing of two naturally occurring heterozygous truncating mutations from each cell line. Increased expression of the truncated mutant alleles relative to the wild-type alleles of the *EXOC1* and *SPTAN1* genes was observed in NCI-H358 cells after treatment with LY3023414 (Figure 2B left). Likewise, increased expression of *HNRNPA3* and *LRPPRC* mutant transcripts was observed in LS180 after LY3023414 treatment (Figure 2B right). We noted no significant effects on two coding region heterozygous single nucleotide polymorphisms (SNP's) after LY3023414 treatment in the genes *RNF43* or *DROSHA* in either cell line (Figure 2B). To substantiate these transcriptomic effects at the protein level, we treated NCI-H358 cells for 24 hours with LY3023414. Western blotting revealed truncated mutant protein for both EXOC1 and SPTAN1 after treatment with LY3023414, whereas the full-length protein levels remained present and unchanged in both the treated and control (DMSO) samples (Figure 2C and 2D left)[39]. An antibody that recognizes the C-terminus of SPTAN1, encoded downstream of the frameshift mutation, did not detect mutant protein, as expected (Figure 2D right).

LY3023414 was originally developed as a phosphatidylinositol 3-kinase (*PI3K*) inhibitor with activity against AKT Serine/Threonine Kinase 1 (*AKT1*) and Mammalian Target of Rapamycin (*mTOR*)[40]. It was tested in a number of clinical trials and has a well-known pharmacokinetic profile *in vivo*[41-45]. To determine whether LY3023414 affects NMD *in vivo*, we established xenograft tumors of both NCI-H358 and LS180 in nude mice. Treatment of these mice with a single oral dose of 60 mg/kg LY3023414 led to a significant increase in the expression of mutant RNA transcripts relative to wild-type transcripts 16 hours later (Figure 2E). The *RNF43* gene, which harbors a coding region heterozygous SNP, served as a control (Figure 2E). Severe drug-associated toxicity, including weight loss bordering on cachexia and

 near total inactivity, precluded the longer-term dosing required for anti-tumorigenic effects of

 LY3023414 in both BALB/c and C57BL/6N immunocompetent mouse strains.

Figure 2: Inhibiting NMD in cancer cells increases broad expression of truncated gene mRNA and protein. A) Mutant transcript recovery rates for genes containing heterozygous indel mutations based on RNA-sequencing results in cell lines treated with 5µM LY3023414 for 16 hours. Strict inclusion criteria were used, such that only mutations with sufficient sequencing coverage are shown (see methods). Recovery is defined as at least 2-fold increase over DMSO treatment. **B)** Targeted high coverage RNA-sequencing confirms recovery of mutant transcript levels in NCI-H358 and LS180 cancer cell lines treated with LY3023414. *RNF43* and *DROSHA* contain common heterozygous SNPs and the mutant allele refers to the non-reference genome allele. Error bars indicate 95% confidence limits. **C)** Western blot analyses of NCI-H358 cells showing mutant and wild-type protein levels in EXOC1 and **D)** SPTAN1 with and without 5 µM LY3023414 treatment. The black arrow indicates the expected size of the mutant protein. The C-terminal SPTAN1 antibody is downstream of the out-of-frame indel mutation and is not expected to identify the mutant allele. **E)** Fold change in the number of mutant RNA transcripts from deep-targeted RNA sequencing of heterozygous mutated genes in NCI-H358 and LS180 xenografts treated by oral gavage with 60 mg/kg LY3023414 assayed 16 hours post-treatment. Student's t-test for target genes are all p < 0.05, while the null hypothesis holds for *RNF43* (common SNP).

The kinase SMG1 is the target for NMD inhibition by LY3023414

To investigate the mechanism of NMD inhibition by LY3023414, we evaluated the six kinases with the highest reported inhibition by LY3023414[40]. siRNA-mediated knockdown of each of these kinases in NCI-H358 and LS180 cancer cell lines was performed for this purpose (Figure S9). Only knockdown of *SMG1* resulted in significant changes in the amount of truncating mutant transcript relative to wild-type transcript in all four genes evaluated (Figure 3A top). *RNF43* and *DROSHA* contain heterozygous coding region SNP's and served as controls in these experiments and as expected showed no changes despite siRNA treatment (Figure 3A bottom).

SMG1 is known to regulate the NMD pathway by activating *UPF1*, an enzyme with RNA helicase and ATPase activity[46,47]. We therefore knocked down *UPF1* with siRNA and found that it restored expression of the NMD-downregulated transcripts, at levels similar to those observed after the knock-down of *SMG1* (Figure 3B). Additionally, treatment of NCI-H358 and LS180 cancer cell lines with a previously reported *SMG1* specific small molecule inhibitor, SMG1i-11 resulted in specific increases in the transcripts and proteins from genes with truncating mutations, just as did LY3023414 (Figures S10 and S11)[30]. Although SMG1i-11 displayed considerable *SMG1* specificity and marked inhibition of NMD, it was highly insoluble. We were unable to find a vehicle to administer it *in vivo*, despite numerous attempts at various formulations and administration routes. This may explain why SMG1i-11 has been demonstrated to be an effective *SMG1* inhibitor *in vitro*, but no peer-reviewed reports of its *in vivo* activity have been reported to date[30,48,49].

Development of an improved NMD inhibitor targeting SMG1

In view of the problems with LY3023414 and SMG1i-11, we sought to develop a new SMG1 inhibitor based on the cryo-electron microscopy structure of the binding pocket of SMG1[50,51]. We attempted the synthesis of eleven compounds (KVS0001 to KVS0011) and tested for bioavailability and preservation of target specificity. Among these, KVS0001 stood out due to its solubility while preserving SMG1 inhibitory activity (Table S2, Figures 3C and S12). Mass spectrometry-based assays showed that KVS0001 inhibits SMG1 protein more than any of the other 246 protein or lipid kinases tested at concentrations from 10nM to 1µM (Table S3 and Figure S13)[52]. Noteworthy off-target kinase inhibition was not observed until doses of 1µM and above. Experiments using NCI-H358 and LS180 cells showed that KVS0001 is bioactive in the nanomolar range and subverts the NMD-mediated down-regulation of truncating mutant transcripts and proteins (Figures 3D, 3E, S14, S15). Inhibition at concentrations as low as 600nM led to equal expression of wild-type and mutant transcripts, suggesting near total blockade of the NMD pathway (Figures 3D, S14). Western blotting showed that KVS0001 substantially decreases the

213 amount of phosphorylated UPF1, the downstream target of SMG1 kinase activity, in three different cell

214 lines (Figure 3F). Finally, KVS0001 treatment of NCI-H358 and LS180-derived xenograft tumors in nude

215 mice resulted in significant increases in transcript levels in each of six tested endogenous genes with

216 truncating mutations, while having had no measurable effects on genes containing heterozygous coding

217 region SNP's (Figures 3G and 3H).

Figure 3: Novel NMD inhibitor KVS0001 is SMG1 specific and induces expression of NMD

targeted genes *in vitro* and *in vivo*. A) Fraction of mutant allele transcripts in genes with heterozygous indels previously established in this study as sensitive to NMD inhibition. Results show mutant levels after siRNA treatment targeting kinases inhibited by LY3023414. *RNF43* and *DROSHA* are common heterozygous SNPs (shaded gray) and serve as negative controls. **B)** Fraction of mutant allele transcripts in genes with truncating mutations known to be sensitive to NMD inhibition after siRNA treatment with siUPF1 or non-targeting siRNA. Data from deep-targeted RNA-sequencing. **C)** Structure of novel NMD inhibitor KVS0001. **D)** Targeted RNA-sequencing on three genes with heterozygous, out-of-frame, indel mutations in LS180 cancer cells treated in a dose-response with KVS0001 or SMG1i-11. *RNF43* serves as a control (common heterozygous SNP) and the mutant allele refers to the non-reference genome allele. **E)** Western blot of EXOC1 protein in NCI-H358 cells treated with 5µM novel inhibitor KVS0001, LY3023414 or SMG1i-11 for 24 hours. **F)** Western blot of phosphorylated UPF1 on three cell lines treated with KVS0001, SMG1i-11, or DMSO. Note that total UPF1 and p-UPF1 were run on different gels, loading controls correspond to indicated gel. **G)** Fold change in the number of mutant allele transcripts measured by targeted RNA-Seq in genes containing heterozygous out-of-frame indel mutations in NCI-H358 or **H)** LS180 subcutaneous xenografts in bilateral flanks of nude mice. Mice were treated once with IP injection of vehicle or 30mg/kg KVS0001 and tumors harvested 16hrs post IP treatment. All genes shown contain heterozygous out-of-frame truncating mutations except *RNF43* and *DROSHA* which serve as controls (contain heterozygous SNP's).

NMD inhibition with KVS0001 causes MHC class I display of hidden neoantigens

Cancer cells may evade immune surveillance by downregulating genes with truncating mutations

as a result of NMD[19,53]. Indeed, previous studies with non-specific or toxic NMD inhibitors have shown

an increase in selected antigens from tumor-specific mutations when NMD is inhibited[54]. We therefore

investigated whether KVS0001 could similarly alter the cell surface presentation of proteins from genes

harboring truncating mutations in NCI-H358 and LS180 cells. Using quantitative HPLC-mass

spectrometry (MS) we observed a striking (45 to 90 fold) increase in the *EXOC1* and *RAB14* derived

neoantigens, and a significant (2 fold) increase in the *ZDHHC16* derived neoantigen (Figures 4A, 4B,

S16, S17)[55]. This is consistent with the re-expression of mutant transcript and protein shown previously in

this study (Figures 3D and 3E). These three peptides were chosen based on an *in-silico* review of

potentially presented peptides and a preliminary experiment that looked at qualitative (present or not

present) presentation of the predicted peptides[56,57].

Targetable peptide presentation occurs with NMD inhibition by KVS0001

To test whether cancer cell neoantigens presented as a result of NMD inhibition could be

targeted, we evaluated two cancer cell lines, NCI-H716 and NCI-H2228. Both lines contain homozygous

truncating mutations in *TP53* which produce transcripts down-regulated by NMD, with each exhibiting

RNA levels less than 20% of the median level of expression among 675 cancer cell lines (Figures 4C and

S18)[58,59]. Treatment with KVS0001 increased the expression of TP53 in both lines, while the commonly

used therapeutic agents 5-fluorouracil and etoposide did not affect TP53 protein abundance (Figures 4D
and S19).

To determine whether this disruption of NMD repression is targetable by T-cells, we developed a
bispecific antibody (KVS-BI043) that recognizes a peptide-HLA complex on one end and CD3 on the
other end. CD3 is expressed only on T-cells, and this bispecific antibody functions as a T-cell engager,
linking target cells to cytotoxic T-cells, which then kill the targets[60,61]. KVS-BI043 recognizes a ten
amino acid peptide (residues 125 to 134 of TP53) bound to HLA-A24. NCI-H2228 naturally express A24
whereas NCI-H716 cells were engineered to express A24 using a retrovirus (NCI-H716_A24). Treatment
of NCI-H716_A24 or NCI-H2228 cells with KVS0001 and the KVS-BI043 bispecific antibody in the
presence of normal T-cells caused a significant increase in interferon (IFN-γ) release (Figure 4E). We
observed no changes in IFN-γ levels in the absence of KVS-BI043 or normal T-cells (Figure 4E grey
boxed lanes). Most importantly, treatment of NCI-H716_A24 with KVS0001 in the presence of KVS-
BI043 and T-cells led to significant killing of the target cancer cells, which was also not observed in the
absence of T-cells or the absence of HLA-24 in the target cells (Figure 4F). NCI-H2228 was assessed for
killing but expressed too much TP53 at baseline (Figure 4D left most lane) and thus displayed substantial
killing even in the DMSO controls.

Figure 4: KVS0001 treatment induces targetable cell surface presentation of peptides known to be downregulated by NMD. A) MHC class I HLA presentation of mutant specific peptide sequences from NCI-H358 and **B)** LS180 cells by quantitative HPLC-Mass Spectrometry. The gene name, type of mutation (in parenthesis) and presented peptide are shown on the y-axis for each gene. Colors indicate different ions. **C)** *TP53* gene structure and mutant DNA sequence for NCI-H716 and NCI-H2228 cancer cell lines, both contain a homozygous splice site mutation in *TP53*. Capital letters represent exonic sequence; lowercase letters represent intronic sequence. DNA mutation reflected by gold bases. **D)** Western blot against TP53 in the presence or absence of NMD inhibitor in NCI-H716_A24 and NCI-H2228 cell lines. NCI-H2228 has an expected size of 46.6 kDa and NCI-H716 of 34.7 kDa. **E)** IFN-γ levels over baseline based on ELISA in a co-culture assay with NCI-H716_A24 and NCI-H2228 cells, NMD inhibitor, human CD3+ T-cells, and bispecific antibody for TP53 and CD3. Chemotherapy (5-Fluorouracil) is shown as a control. **F)** Cell killing based on luciferase levels in a co-culture assay in NCI-H716 cells with and without A24 expression, treated with TP53-CD3 bispecific antibody, NMD inhibitor and human CD3+ T-cells.

Tumor growth is slowed in mice treated with KVS0001

Finally, we investigated whether KVS0001 could impact tumor growth in syngeneic models in

which the native immune system might play a role. For this purpose, we first used murine RENCA (renal

cancer) and LLC (lung cancer) as they are known to have a relatively large number of out-of-frame indel

mutations (Table S4). Although murine and human SMG1 are highly related (98% at the amino acid

level), it was important to show that KVS0001 could actually inhibit NMD in murine cells. For this

purpose, we tested eight genes in LLC, and four in RENCA, which contained out-of-frame indel

mutations potentially targeted by NMD. We also assessed the expression of three genes without any

mutations known to have their normal expression controlled by NMD[62]. We found that six of the twelve

truncating mutation-containing genes and five of the six expression controlled by NMD genes had

significantly increased RNA following *in vitro* treatment with KVS0001 (Figure 5A and 5B).

We then implanted LLC and RENCA cancer cells in the mammary fat pad of C57BL/6N and

BALB/c mice respectively and treated with KVS0001 or vehicle control (Figure 5C). The dose of

KVS0001 was based on experiments showing that the maximum solubility limit was reached around 2 to

3mg/mL, leading to a maximum single dose of 30mg/kg per mouse per treatment. At this dose, the only

toxicity noted was transient weight loss (Figure S20), and no other pathology was observed. Both tumor

types experienced significant slowing of tumor growth (Figure 5D). However, when the same tumors

were implanted in immunocompromised mice, there was no statistically significant difference in tumor

growth between mice treated with KVS0001 or vehicle control (Figure 5E).

We evaluated a total of seven syngeneic mouse cancer cell lines. Four contained a relatively high

number of truncating indel mutations, while the remaining three contained a moderate or low number.

None of the three models with moderate or low numbers of truncating mutations responded to KVS0001

(EMT6, B16-F10, M3 Melanoma) (Figure S21). Of the four models with a higher number of truncating

mutations, two (LLC and RENCA) responded in a statistically significant fashion while the other two

failed to reach significance (CT26, MC38) (Figures 5D and S21).

Figure 5: *In vivo* **treatment of murine tumors with KVS0001 yield differential tumor growth compared with vehicle treatment. A)** Fold change in RNA transcript levels in LLC or **B)** RENCA cells treated in-vitro with 5µM of NMD inhibitor KVS0001 or DMSO. Orange bars indicate genes with homozygous indel mutations potentially targeted by NMD. Purple bars show genes with no mutations but that are known to have their normal transcription levels controlled by NMD. Green bar is a control gene that should not change with treatment. The dotted line shows relative expression of DMSO treatment (equal to 1). **C)** Treatment schedule for syngeneic tumor mouse experiments. **D)** Average tumor size of LLC (left) and RENCA (right) syngeneic tumors in immune-competent mice (n=8) treated with 30 mg/kg KVS0001 or vehicle control. Difference is statistically significant after day 10 based on one-way ANOVA with Dunnett´s test p<0.001 (p<0.05 for day 23 RENCA data point) for both tumors tested. **E)** Average tumor size of LLC (left) and RENCA (right) in immunodeficient mice (n=8) treated with 30 mg/kg KVS0001 or vehicle control. Error bars show 95% confidence intervals in all plots.

Discussion

We report the design and execution of a high throughput screen to identify small molecule

inhibitors of the nonsense-mediated decay pathway. The method employs a ratiometric output, allowing

quantitative and controlled determination of RNA expression changes associated with NMD. Unlike

previous NMD screens, this assay is directed at NMD as a process rather than as a way to identify

compounds for treating specific mutations related to a disease state, such as cystic fibrosis or β-

thalassemia[26,28,63]. The genetically modified isogenic cell lines employed here are uniquely suited for this

approach, producing a high signal-to-noise ratio. Follow- up experiments confirmed a low false positive

rate in the screen: four of the eight hits re-expressed transcripts from genes with truncating mutations in a

dose-responsive manner when re-tested. Note that our assay allows for the detection of NMD inhibitors

regardless of effects on protein translation by virtue of the comparison of wild-type to mutant transcripts

for each queried gene. This is an important feature, as many previously reported inhibitors of NMD rely

on inhibition of protein translation and thus are unlikely to be optimal for specific restoration of NMD-

targeted RNA and protein expression[33,36,37,64].

The optimal inhibitor of NMD found in our screen was LY3023414, and we were able to show

that this compound inhibited the SMG1 protein kinase. *SMG1* acts as the gatekeeper of *UPF1* activity

through its phosphorylation at multiple sites[46]. Unfortunately, issues likely related to off-target (pan-

kinase) toxicity at the doses required for NMD inhibition prevented LY3023414 from being utilized in

further *in vivo* work. While there are previously described SMG1 inhibitors, all have undesirable off-

target or biophysical characteristics complicating their use as therapeutic agents[26-29,37,54,65]. For example,

SMG1i-11 was previously identified as a potent and specific SMG1 inhibitor[30]. However, in our hands, it

was insoluble and neither we nor others have been able to formulate it for administration *in vivo*. Another

unrelated compound has shown successful *in vivo* re-expression of mutant RNA transcripts, but the effect

is reported as less than a two-fold change, compared to the three to four-fold demonstrated here by

KVS0001, and little work has been done to develop the compound further in subsequent studies[66].

In light of these issues, we attempted to improve the biophysical properties of SMG1i-11 through

the development of KVS0001. As reported in this study, KVS0001 was a specific inhibitor of SMG1 that

was soluble and could easily be administered to mice. Treatment of KVS0001 not only increased levels of

transcripts and proteins from genes with truncating mutations, but also increased the levels of

corresponding peptide bound-HLA complexes on the cell surfaces. We developed a new bispecific

antibody, KVS-B043, which could recognize these pHLA complexes when the truncating mutations were

in the *TP53* gene and destroy cancer cells harboring such mutations. In syngeneic mouse models,

KVS0001 slowed the growth of some tumors containing multiple truncating mutations, but only when the

mice were immunocompetent. This dependence on an intact immune system supports the idea that the

slowed growth of the treated tumors was due to recognition of neoantigens.

Despite these encouraging data, the potential utility of KVS0001 for therapeutic purposes remains

speculative, for several reasons. First, we do not understand why KVS0001 (and other NMD inhibitors)

increase the expression of many, but not all, genes with truncating mutations. More research on the

biochemistry of NMD will undoubtedly be required for such understanding, and perhaps the tools

described here may facilitate that effort. Second, treatment with KVS0001 reduced the growth of some

but not all syngeneic tumor types tested in mice. There appeared to be a correlation between the ability of

KVS0001 to inhibit growth and the number of truncating mutations in the tumor. This situation is

reminiscent of that encountered with immune checkpoint inhibitors, in which there is a positive but

imperfect correlation with tumor mutation burden[8,12]. Undoubtedly, the ability of the immune system to

react to neoantigens is related to their quality as well as their quantity, with quality defined as the ability

of the neoantigen to be bound to the host's particular MHC constitution and initiate an immune

response[67,68]. Third, treatment with KVS0001 led to decreased tumor growth in immunocompetent mice,

but did not lead to tumor regressions of the type mandated by RECIST criteria in human clinical trials[69,70].

Whether this failure to cure tumors in mice results from the contrived nature of the models used –

injection of large numbers of rapidly growing cancer cells into animals with relatively little time to react

prior to their demise – complicates the interpretation of many pre-clinical immunotherapeutic approaches.

Finally, KVS0001, though it was soluble and could be administered to animals for at least a

month, was not entirely non-toxic, causing transient weight loss in some of the mice. Previous work has

raised concerns that the importance of NMD in development and normal gene expression make it a

difficult pathway to safely disrupt[21,71-74]. The limited impact on body weight or observed physical activity

of mice undergoing KVS0001 therapy at therapeutically relevant doses suggests that NMD inhibition may

have acceptable toxicity and be tolerated in developed animals. Another example of this lies in the natural

genetic variants of NMD components, which convey variability in the efficiency of NMD between

humans, and lends itself to supporting that knockdown of this pathway may be tolerable[75-77]. Studies in

mice also support this observation, as post-development knockdown of UPF1 displayed minimal

phenotype change[62]. The broad use of NMD in normal development and growth, coupled with the

observations here, suggests future work with inhibitors of this pathway should continue with a close eye

for on-target off-tissue toxicity.

Though there is much work to be done, we hope that the tools, approaches, and compounds

described here will facilitate that work. Should the administration of KVS0001 or a related compound

prove non-toxic and well-tolerated in humans, we end with a speculation about the possibility of

KVS0001 being used as a preventative rather than as a therapeutic agent. Its potential use to prevent the

onset of symptoms in pediatric syndromes caused by germline truncating mutations is obvious[17,78]. Less

obvious is the potential for it to prophylactically reduce cancer incidence in patients with hereditary non-

polyposis colorectal cancer. These patients inherit heterozygous mutations of a mismatch repair gene, and

they do not develop tumors until biallelic mutations of that mismatch repair gene are acquired in a rare

stem cell during the second or third decade of life. Similar speculations can be made about the potential of

KVS0001 to be used to prevent cancer initiation or progression in patients with other inherited mutations

in repair genes, or in individuals exposed to high levels of exogenous mutagens.

References

1.   Haslam, A., and Prasad, V. (2019). Estimation of the Percentage of US Patients With Cancer Who
Are Eligible for and Respond to Checkpoint Inhibitor Immunotherapy Drugs. JAMA Netw Open *2*,
e192535. 10.1001/jamanetworkopen.2019.2535.
2.   Ben-Aharon, O., Magnezi, R., Leshno, M., and Goldstein, D.A. (2018). Association of
Immunotherapy With Durable Survival as Defined by Value Frameworks for Cancer Care. JAMA
Oncol *4*, 326-332. 10.1001/jamaoncol.2017.4445.
3.   Darvin, P., Toor, S.M., Sasidharan Nair, V., and Elkord, E. (2018). Immune checkpoint inhibitors:
recent progress and potential biomarkers. Exp Mol Med *50*, 1-11. 10.1038/s12276-018-0191-1.
4.   Yang, K., Halima, A., and Chan, T.A. (2023). Antigen presentation in cancer - mechanisms and
clinical implications for immunotherapy. Nat Rev Clin Oncol *20*, 604-623. 10.1038/s41571-023-
00789-4.
5.   Kraehenbuehl, L., Weng, C.H., Eghbali, S., Wolchok, J.D., and Merghoub, T. (2022). Enhancing
immunotherapy in cancer by targeting emerging immunomodulatory pathways. Nat Rev Clin
Oncol *19*, 37-50. 10.1038/s41571-021-00552-7.
6.   Mandal, R., Samstein, R.M., Lee, K.W., Havel, J.J., Wang, H., Krishna, C., Sabio, E.Y., Makarov, V.,
Kuo, F., Blecua, P., et al. (2019). Genetic diversity of tumors with mismatch repair deficiency
influences anti-PD-1 immunotherapy response. Science *364*, 485-491. 10.1126/science.aau0447.
7.   Turajlic, S., Litchfield, K., Xu, H., Rosenthal, R., McGranahan, N., Reading, J.L., Wong, Y.N.S.,
Rowan, A., Kanu, N., Al Bakir, M., et al. (2017). Insertion-and-deletion-derived tumour-specific
neoantigens and the immunogenic phenotype: a pan-cancer analysis. Lancet Oncol *18*, 1009-
1021. 10.1016/S1470-2045(17)30516-8.
8.   Yarchoan, M., Hopkins, A., and Jaffee, E.M. (2017). Tumor Mutational Burden and Response
Rate to PD-1 Inhibition. N Engl J Med *377*, 2500-2501. 10.1056/NEJMc1713444.
9.   Chan, T.A., Yarchoan, M., Jaffee, E., Swanton, C., Quezada, S.A., Stenzinger, A., and Peters, S.
(2019). Development of tumor mutation burden as an immunotherapy biomarker: utility for the
oncology clinic. Ann Oncol *30*, 44-56. 10.1093/annonc/mdy495.
10.   Van Allen, E.M., Miao, D., Schilling, B., Shukla, S.A., Blank, C., Zimmer, L., Sucker, A., Hillen, U.,
Foppen, M.H.G., Goldinger, S.M., et al. (2015). Genomic correlates of response to CTLA-4
blockade in metastatic melanoma. Science *350*, 207-211. 10.1126/science.aad0095.

11. Samstein, R.M., Lee, C.H., Shoushtari, A.N., Hellmann, M.D., Shen, R., Janjigian, Y.Y., Barron,
D.A., Zehir, A., Jordan, E.J., Omuro, A., et al. (2019). Tumor mutational load predicts survival
after immunotherapy across multiple cancer types. Nat Genet *51*, 202-206. 10.1038/s41588-
018-0312-8.
12. Ma, X., Riaz, N., Samstein, R.M., Lee, M., Makarov, V., Valero, C., Chowell, D., Kuo, F., Hoen, D.,
Fitzgerald, C.W.R., et al. (2022). Functional landscapes of POLE and POLD1 mutations in
checkpoint blockade-dependent antitumor immunity. Nat Genet *54*, 996-1012. 10.1038/s41588-
022-01108-w.
13. Rizvi, N.A., Hellmann, M.D., Snyder, A., Kvistborg, P., Makarov, V., Havel, J.J., Lee, W., Yuan, J.,
Wong, P., Ho, T.S., et al. (2015). Cancer immunology. Mutational landscape determines
sensitivity to PD-1 blockade in non-small cell lung cancer. Science *348*, 124-128.
10.1126/science.aaa1348.
14. Consortium, I.T.P.-C.A.o.W.G. (2020). Pan-cancer analysis of whole genomes. Nature *578*, 82-93.
10.1038/s41586-020-1969-6.
15. Vogelstein, B., Papadopoulos, N., Velculescu, V.E., Zhou, S., Diaz, L.A., Jr., and Kinzler, K.W.
(2013). Cancer genome landscapes. Science *339*, 1546-1558. 10.1126/science.1235122.
16. Segal, N.H., Parsons, D.W., Peggs, K.S., Velculescu, V., Kinzler, K.W., Vogelstein, B., and Allison,
J.P. (2008). Epitope landscape in breast and colorectal cancer. Cancer Res *68*, 889-892.
10.1158/0008-5472.CAN-07-3095.
17. Lindeboom, R.G.H., Vermeulen, M., Lehner, B., and Supek, F. (2019). The impact of nonsense-
mediated mRNA decay on genetic disease, gene editing and cancer immunotherapy. Nat Genet
51, 1645-1651. 10.1038/s41588-019-0517-5.
18. Pastor, F., Kolonias, D., Giangrande, P.H., and Gilboa, E. (2010). Induction of tumour immunity
by targeted inhibition of nonsense-mediated mRNA decay. Nature *465*, 227-230.
10.1038/nature08999.
19. Nogueira, G., Fernandes, R., Garcia-Moreno, J.F., and Romao, L. (2021). Nonsense-mediated RNA
decay and its bipolar function in cancer. Mol Cancer *20*, 72. 10.1186/s12943-021-01364-0.
20. Frischmeyer, P.A., and Dietz, H.C. (1999). Nonsense-mediated mRNA decay in health and
disease. Hum Mol Genet *8*, 1893-1900. 10.1093/hmg/8.10.1893.
21. Mendell, J.T., Sharifi, N.A., Meyers, J.L., Martinez-Murillo, F., and Dietz, H.C. (2004). Nonsense
surveillance regulates expression of diverse classes of mammalian transcripts and mutes
genomic noise. Nat Genet *36*, 1073-1078. 10.1038/ng1429.
22. Dalton, W.B., Helmenstine, E., Walsh, N., Gondek, L.P., Kelkar, D.S., Read, A., Natrajan, R.,
Christenson, E.S., Roman, B., Das, S., et al. (2019). Hotspot SF3B1 mutations induce metabolic
reprogramming and vulnerability to serine deprivation. J Clin Invest *129*, 4708-4723.
10.1172/JCI125022.
23. Castle, J.C., Loewer, M., Boegel, S., de Graaf, J., Bender, C., Tadmor, A.D., Boisguerin, V., Bukur,
T., Sorn, P., Paret, C., et al. (2014). Immunomic, genomic and transcriptomic characterization of
CT26 colorectal carcinoma. BMC Genomics *15*, 190. 10.1186/1471-2164-15-190.
24. Oka, M., Xu, L., Suzuki, T., Yoshikawa, T., Sakamoto, H., Uemura, H., Yoshizawa, A.C., Suzuki, Y.,
Nakatsura, T., Ishihama, Y., et al. (2021). Aberrant splicing isoforms detected by full-length
transcriptome sequencing as transcripts of potential neoantigens in non-small cell lung cancer.
Genome Biol *22*, 9. 10.1186/s13059-020-02240-8.
25. Lu, S.X., De Neef, E., Thomas, J.D., Sabio, E., Rousseau, B., Gigoux, M., Knorr, D.A., Greenbaum,
B., Elhanati, Y., Hogg, S.J., et al. (2021). Pharmacologic modulation of RNA splicing enhances
anti-tumor immunity. Cell *184*, 4032-4047 e4031. 10.1016/j.cell.2021.05.038.

26. Martin, L., Grigoryan, A., Wang, D., Wang, J., Breda, L., Rivella, S., Cardozo, T., and Gardner, L.B.
(2014). Identification and characterization of small molecules that inhibit nonsense-mediated
RNA decay and suppress nonsense p53 mutations. Cancer Res *74*, 3104-3113. 10.1158/0008-
5472.CAN-13-2235.
27. Durand, S., Cougot, N., Mahuteau-Betzer, F., Nguyen, C.H., Grierson, D.S., Bertrand, E., Tazi, J.,
and Lejeune, F. (2007). Inhibition of nonsense-mediated mRNA decay (NMD) by a new chemical
molecule reveals the dynamic of NMD factors in P-bodies. J Cell Biol *178*, 1145-1160.
10.1083/jcb.200611086.
28. Zhao, J., Li, Z., Puri, R., Liu, K., Nunez, I., Chen, L., and Zheng, S. (2022). Molecular profiling of
individual FDA-approved clinical drugs identifies modulators of nonsense-mediated mRNA
decay. Mol Ther Nucleic Acids *27*, 304-318. 10.1016/j.omtn.2021.12.003.
29. Gotham, V.J., Hobbs, M.C., Burgin, R., Turton, D., Smythe, C., and Coldham, I. (2016). Synthesis
and activity of a novel inhibitor of nonsense-mediated mRNA decay. Org Biomol Chem *14*, 1559-
1563. 10.1039/c5ob02482j.
30. Gopalsamy, A., Bennett, E.M., Shi, M., Zhang, W.G., Bard, J., and Yu, K. (2012). Identification of
pyrimidine derivatives as hSMG-1 inhibitors. Bioorg Med Chem Lett *22*, 6636-6641.
10.1016/j.bmcl.2012.08.107.
31. Cook, A.L., Wyhs, N., Sur, S., Ptak, B., Popoli, M., Dobbyn, L., Papadopoulos, T., Bettegowda, C.,
Papadopoulos, N., Vogelstein, B., et al. (2022). An isogenic cell line panel for sequence-based
screening of targeted anticancer drugs. iScience *25*, 104437. 10.1016/j.isci.2022.104437.
32. Carter, M.S., Doskow, J., Morris, P., Li, S., Nhim, R.P., Sandstedt, S., and Wilkinson, M.F. (1995). A
regulatory mechanism that detects premature nonsense codons in T-cell receptor transcripts in
vivo is reversed by protein synthesis inhibitors in vitro. J Biol Chem *270*, 28995-29003.
10.1074/jbc.270.48.28995.
33. Siddiqui, S., Firat, D., and Olshin, S. (1973). Phase II study of emetine (NSC-33669) in the
treatment of solid tumors. Cancer Chemother Rep *57*, 423-428.
34. Moertel, C.G., Schutt, A.J., Hahn, R.G., and Reitemeier, R.J. (1974). Treatment of advanced
gastrointestinal cancer with ememtine (NSC-33669). Cancer Chemother Rep *58*, 229-232.
35. Mastrangelo, M.J., Grage, T.B., Bellet, R.E., and Weiss, A.J. (1973). A phase I study of emetine
hydrochloride (NSC 33669) in solid tumors. Cancer *31*, 1170-1175. 10.1002/1097-
0142(197305)31:5<1170::aid-cncr2820310520>3.0.co;2-4.
36. Tang, Z., Xing, F., Chen, D., Yu, Y., Yu, C., Di, J., and Liu, J. (2012). In vivo toxicological evaluation
of Anisomycin. Toxicol Lett *208*, 1-11. 10.1016/j.toxlet.2011.10.001.
37. Bongiorno, R., Colombo, M.P., and Lecis, D. (2021). Deciphering the nonsense-mediated mRNA
decay pathway to identify cancer cell vulnerabilities for effective cancer therapy. J Exp Clin
Cancer Res *40*, 376. 10.1186/s13046-021-02192-2.
38. Cowen, L.E., and Tang, Y. (2017). Identification of nonsense-mediated mRNA decay pathway as a
critical regulator of p53 isoform beta. Sci Rep *7*, 17535. 10.1038/s41598-017-17283-4.
39. Williams, D.S., Bird, M.J., Jorissen, R.N., Yu, Y.L., Walker, F., Zhang, H.H., Nice, E.C., and Burgess,
A.W. (2010). Nonsense mediated decay resistant mutations are a source of expressed mutant
proteins in colon cancer cell lines with microsatellite instability. PLoS One *5*, e16012.
10.1371/journal.pone.0016012.
40. Smith, M.C., Mader, M.M., Cook, J.A., Iversen, P., Ajamie, R., Perkins, E., Bloem, L., Yip, Y.Y.,
Barda, D.A., Waid, P.P., et al. (2016). Characterization of LY3023414, a Novel PI3K/mTOR Dual
Inhibitor Eliciting Transient Target Modulation to Impede Tumor Growth. Mol Cancer Ther *15*,
2344-2356. 10.1158/1535-7163.MCT-15-0996.

41. Bendell, J.C., Varghese, A.M., Hyman, D.M., Bauer, T.M., Pant, S., Callies, S., Lin, J., Martinez, R.,
Wickremsinhe, E., Fink, A., et al. (2018). A First-in-Human Phase 1 Study of LY3023414, an Oral
PI3K/mTOR Dual Inhibitor, in Patients with Advanced Cancer. Clin Cancer Res *24*, 3253-3262.
10.1158/1078-0432.CCR-17-3421.
42. Rubinstein, M.M., Hyman, D.M., Caird, I., Won, H., Soldan, K., Seier, K., Iasonos, A., Tew, W.P.,
O'Cearbhaill, R.E., Grisham, R.N., et al. (2020). Phase 2 study of LY3023414 in patients with
advanced endometrial cancer harboring activating mutations in the PI3K pathway. Cancer *126*,
1274-1282. 10.1002/cncr.32677.
43. Wei, L., Chintala, S., Ciamporcero, E., Ramakrishnan, S., Elbanna, M., Wang, J., Hu, Q., Glenn,
S.T., Murakami, M., Liu, L., et al. (2016). Genomic profiling is predictive of response to cisplatin
treatment but not to PI3K inhibition in bladder cancer patient-derived xenografts. Oncotarget *7*,
76374-76389. 10.18632/oncotarget.13062.
44. Zou, Y., Ge, M., and Wang, X. (2017). Targeting PI3K-AKT-mTOR by LY3023414 inhibits human
skin squamous cell carcinoma cell growth in vitro and in vivo. Biochem Biophys Res Commun
490, 385-392. 10.1016/j.bbrc.2017.06.052.
45. Sweeney, C.J., Percent, I.J., Babu, S., Cultrera, J.L., Mehlhaff, B.A., Goodman, O.B., Morris, D.S.,
Schnadig, I.D., Albany, C., Shore, N.D., et al. (2022). Phase Ib/II Study of Enzalutamide with
Samotolisib (LY3023414) or Placebo in Patients with Metastatic Castration-Resistant Prostate
Cancer. Clin Cancer Res *28*, 2237-2247. 10.1158/1078-0432.CCR-21-2326.
46. Yamashita, A., Ohnishi, T., Kashima, I., Taya, Y., and Ohno, S. (2001). Human SMG-1, a novel
phosphatidylinositol 3-kinase-related protein kinase, associates with components of the mRNA
surveillance complex and is involved in the regulation of nonsense-mediated mRNA decay.
Genes Dev *15*, 2215-2228. 10.1101/gad.913001.
47. Yamashita, A. (2013). Role of SMG-1-mediated Upf1 phosphorylation in mammalian nonsense-
mediated mRNA decay. Genes Cells *18*, 161-175. 10.1111/gtc.12033.
48. Keenan, M.M., Huang, L., Jordan, N.J., Wong, E., Cheng, Y., Valley, H.C., Mahiou, J., Liang, F.,
Bihler, H., Mense, M., et al. (2019). Nonsense-mediated RNA Decay Pathway Inhibition Restores
Expression and Function of W1282X CFTR. Am J Respir Cell Mol Biol *61*, 290-300.
10.1165/rcmb.2018-0316OC.
49. Valley, H.C., Bukis, K.M., Bell, A., Cheng, Y., Wong, E., Jordan, N.J., Allaire, N.E., Sivachenko, A.,
Liang, F., Bihler, H., et al. (2019). Isogenic cell models of cystic fibrosis-causing variants in
natively expressing pulmonary epithelial cells. J Cyst Fibros *18*, 476-483.
10.1016/j.jcf.2018.12.001.
50. Zhu, L., Li, L., Qi, Y., Yu, Z., and Xu, Y. (2019). Cryo-EM structure of SMG1-SMG8-SMG9 complex.
Cell Res *29*, 1027-1034. 10.1038/s41422-019-0255-3.
51. Langer, L.M., Bonneau, F., Gat, Y., and Conti, E. (2021). Cryo-EM reconstructions of inhibitor-
bound SMG1 kinase reveal an autoinhibitory state dependent on SMG8. Elife *10*.
10.7554/eLife.72353.
52. Patricelli, M.P., Nomanbhoy, T.K., Wu, J., Brown, H., Zhou, D., Zhang, J., Jagannathan, S., Aban,
A., Okerberg, E., Herring, C., et al. (2011). In situ kinase profiling reveals functionally relevant
properties of native kinases. Chem Biol *18*, 699-710. 10.1016/j.chembiol.2011.04.011.
53. Hu, Z., Yau, C., and Ahmed, A.A. (2017). A pan-cancer genome-wide analysis reveals tumour
dependencies by induction of nonsense-mediated decay. Nat Commun *8*, 15943.
10.1038/ncomms15943.
54. Becker, J.P., Helm, D., Rettel, M., Stein, F., Hernandez-Sanchez, A., Urban, K., Gebert, J., Kloor,
M., Neu-Yilik, G., von Knebel Doeberitz, M., et al. (2021). NMD inhibition by 5-azacytidine

525   augments presentation of immunogenic frameshift-derived neoepitopes. iScience *24*, 102389.
526   10.1016/j.isci.2021.102389.

527 55. Wang, Q., Douglass, J., Hwang, M.S., Hsiue, E.H., Mog, B.J., Zhang, M., Papadopoulos, N., Kinzler,
528   K.W., Zhou, S., and Vogelstein, B. (2019). Direct Detection and Quantification of Neoantigens.
529   Cancer Immunol Res *7*, 1748-1754. 10.1158/2326-6066.CIR-19-0107.

530 56. Reynisson, B., Alvarez, B., Paul, S., Peters, B., and Nielsen, M. (2020). NetMHCpan-4.1 and
531   NetMHCIIpan-4.0: improved predictions of MHC antigen presentation by concurrent motif
532   deconvolution and integration of MS MHC eluted ligand data. Nucleic Acids Res *48*, W449-
533   W454. 10.1093/nar/gkaa379.

534 57. Schmidt, J., Smith, A.R., Magnin, M., Racle, J., Devlin, J.R., Bobisse, S., Cesbron, J., Bonnet, V.,
535   Carmona, S.J., Huber, F., et al. (2021). Prediction of neo-epitope immunogenicity reveals TCR
536   recognition determinants and provides insight into immunoediting. Cell Rep Med *2*, 100194.
537   10.1016/j.xcrm.2021.100194.

538 58. Lindeboom, R.G., Supek, F., and Lehner, B. (2016). The rules and impact of nonsense-mediated
539   mRNA decay in human cancers. Nat Genet *48*, 1112-1118. 10.1038/ng.3664.

540 59. Klijn, C., Durinck, S., Stawiski, E.W., Haverty, P.M., Jiang, Z., Liu, H., Degenhardt, J., Mayba, O.,
541   Gnad, F., Liu, J., et al. (2015). A comprehensive transcriptional portrait of human cancer cell
542   lines. Nat Biotechnol *33*, 306-312. 10.1038/nbt.3080.

543 60. Mack, M., Riethmuller, G., and Kufer, P. (1995). A small bispecific antibody construct expressed
544   as a functional single-chain molecule with high tumor cell cytotoxicity. Proc Natl Acad Sci U S A
545   *92*, 7021-7025. 10.1073/pnas.92.15.7021.

546 61. Middelburg, J., Kemper, K., Engelberts, P., Labrijn, A.F., Schuurman, J., and van Hall, T. (2021).
547   Overcoming Challenges for CD3-Bispecific Antibody Therapy in Solid Tumors. Cancers (Basel) *13*.
548   10.3390/cancers13020287.

549 62. Echols, J., Siddiqui, A., Dai, Y., Havasi, V., Sun, R., Kaczmarczyk, A., and Keeling, K.M. (2020). A
550   regulated NMD mouse model supports NMD inhibition as a viable therapeutic option to treat
551   genetic diseases. Dis Model Mech *13*. 10.1242/dmm.044891.

552 63. Salvatori, F., Cantale, V., Breveglieri, G., Zuccato, C., Finotti, A., Bianchi, N., Borgatti, M., Feriotto,
553   G., Destro, F., Canella, A., et al. (2009). Development of K562 cell clones expressing beta-globin
554   mRNA carrying the beta039 thalassaemia mutation for the screening of correctors of stop-codon
555   mutations. Biotechnol Appl Biochem *54*, 41-52. 10.1042/BA20080266.

556 64. Bhuvanagiri, M., Lewis, J., Putzker, K., Becker, J.P., Leicht, S., Krijgsveld, J., Batra, R., Turnwald,
557   B., Jovanovic, B., Hauer, C., et al. (2014). 5-azacytidine inhibits nonsense-mediated decay in a
558   MYC-dependent fashion. EMBO Mol Med *6*, 1593-1609. 10.15252/emmm.201404461.

559 65. Cheruiyot, A., Li, S., Nickless, A., Roth, R., Fitzpatrick, J.A.J., and You, Z. (2018). Compound C
560   inhibits nonsense-mediated RNA decay independently of AMPK. PLoS One *13*, e0204978.
561   10.1371/journal.pone.0204978.

562 66. Keeling, K.M., Wang, D., Dai, Y., Murugesan, S., Chenna, B., Clark, J., Belakhov, V., Kandasamy, J.,
563   Velu, S.E., Baasov, T., and Bedwell, D.M. (2013). Attenuation of nonsense-mediated mRNA decay
564   enhances in vivo nonsense suppression. PLoS One *8*, e60478. 10.1371/journal.pone.0060478.

565 67. Halima, A., Vuong, W., and Chan, T.A. (2022). Next-generation sequencing: unraveling genetic
566   mechanisms that shape cancer immunotherapy efficacy. J Clin Invest *132*. 10.1172/JCI154945.

567 68. Ribas, A., and Wolchok, J.D. (2018). Cancer immunotherapy using checkpoint blockade. Science
568   *359*, 1350-1355. 10.1126/science.aar4060.

569 69. Eisenhauer, E.A., Therasse, P., Bogaerts, J., Schwartz, L.H., Sargent, D., Ford, R., Dancey, J.,
570   Arbuck, S., Gwyther, S., Mooney, M., et al. (2009). New response evaluation criteria in solid

tumours: revised RECIST guideline (version 1.1). Eur J Cancer *45*, 228-247.
10.1016/j.ejca.2008.10.026.
70.   Seymour, L., Bogaerts, J., Perrone, A., Ford, R., Schwartz, L.H., Mandrekar, S., Lin, N.U., Litiere, S.,
Dancey, J., Chen, A., et al. (2017). iRECIST: guidelines for response criteria for use in trials testing
immunotherapeutics. Lancet Oncol *18*, e143-e152. 10.1016/S1470-2045(17)30074-8.
71.   Kurosaki, T., Popp, M.W., and Maquat, L.E. (2019). Quality and quantity control of gene
expression by nonsense-mediated mRNA decay. Nat Rev Mol Cell Biol *20*, 406-420.
10.1038/s41580-019-0126-2.
72.   Tarpey, P.S., Raymond, F.L., Nguyen, L.S., Rodriguez, J., Hackett, A., Vandeleur, L., Smith, R.,
Shoubridge, C., Edkins, S., Stevens, C., et al. (2007). Mutations in UPF3B, a member of the
nonsense-mediated mRNA decay complex, cause syndromic and nonsyndromic mental
retardation. Nat Genet *39*, 1127-1133. 10.1038/ng2100.
73.   Nguyen, L.S., Kim, H.G., Rosenfeld, J.A., Shen, Y., Gusella, J.F., Lacassie, Y., Layman, L.C., Shaffer,
L.G., and Gecz, J. (2013). Contribution of copy number variants involving nonsense-mediated
mRNA decay pathway genes to neuro-developmental disorders. Hum Mol Genet *22*, 1816-1825.
10.1093/hmg/ddt035.
74.   McIlwain, D.R., Pan, Q., Reilly, P.T., Elia, A.J., McCracken, S., Wakeham, A.C., Itie-Youten, A.,
Blencowe, B.J., and Mak, T.W. (2010). Smg1 is required for embryogenesis and regulates diverse
genes via alternative splicing coupled to nonsense-mediated mRNA decay. Proc Natl Acad Sci U S
A *107*, 12186-12191. 10.1073/pnas.1007336107.
75.   Linde, L., Boelz, S., Neu-Yilik, G., Kulozik, A.E., and Kerem, B. (2007). The efficiency of nonsense-
mediated mRNA decay is an inherent character and varies among different cells. Eur J Hum
Genet *15*, 1156-1162. 10.1038/sj.ejhg.5201889.
76.   Nguyen, L.S., Wilkinson, M.F., and Gecz, J. (2014). Nonsense-mediated mRNA decay: inter-
individual variability and human disease. Neurosci Biobehav Rev *46 Pt 2*, 175-186.
10.1016/j.neubiorev.2013.10.016.
77.   MacArthur, D.G., Balasubramanian, S., Frankish, A., Huang, N., Morris, J., Walter, K., Jostins, L.,
Habegger, L., Pickrell, J.K., Montgomery, S.B., et al. (2012). A systematic survey of loss-of-
function variants in human protein-coding genes. Science *335*, 823-828.
10.1126/science.1215040.
78.   Miller, J.N., and Pearce, D.A. (2014). Nonsense-mediated decay in genetic disease: friend or foe?
Mutat Res Rev Mutat Res *762*, 52-64. 10.1016/j.mrrev.2014.05.001.
79.   Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol *37*, 907-915.
10.1038/s41587-019-0201-4.
80.   Pertea, M., Kim, D., Pertea, G.M., Leek, J.T., and Salzberg, S.L. (2016). Transcript-level expression
analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. Nat Protoc *11*, 1650-1667.
10.1038/nprot.2016.095.
81.   Robinson, J.T., Thorvaldsdottir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and
Mesirov, J.P. (2011). Integrative genomics viewer. Nat Biotechnol *29*, 24-26. 10.1038/nbt.1754.
82.   Koboldt, D.C., Zhang, Q., Larson, D.E., Shen, D., McLellan, M.D., Lin, L., Miller, C.A., Mardis, E.R.,
Ding, L., and Wilson, R.K. (2012). VarScan 2: somatic mutation and copy number alteration
discovery in cancer by exome sequencing. Genome Res *22*, 568-576. 10.1101/gr.129684.111.
83.   Hsiue, E.H., Wright, K.M., Douglass, J., Hwang, M.S., Mog, B.J., Pearlman, A.H., Paul, S., DiNapoli,
S.R., Konig, M.F., Wang, Q., et al. (2021). Targeting a neoantigen derived from a common TP53
mutation. Science *371*. 10.1126/science.abc8697.

84.   Skora, A.D., Douglass, J., Hwang, M.S., Tam, A.J., Blosser, R.L., Gabelli, S.B., Cao, J., Diaz, L.A., Jr.,
Papadopoulos, N., Kinzler, K.W., et al. (2015). Generation of MANAbodies specific to HLA-
restricted epitopes encoded by somatically mutated genes. Proc Natl Acad Sci U S A *112*, 9967-
9972. 10.1073/pnas.1511996112.
85.   Hwang, M.S., Mog, B.J., Douglass, J., Pearlman, A.H., Hsiue, E.H., Paul, S., DiNapoli, S.R., Konig,
M.F., Pardoll, D.M., Gabelli, S.B., et al. (2021). Targeting loss of heterozygosity for cancer-specific
immunotherapy. Proc Natl Acad Sci U S A *118*. 10.1073/pnas.2022410118.

Acknowledgements

The authors thank Dr. Qing Wang from Complete Omics Inc. for his help and advice with the neoantigen presentation by mass spectrometry studies. We would also like to thank Dr. Xiao Chen from Ascendex LLC. for helpful suggestions during medicinal chemistry experiments. Biophysical predictions for small molecules was done using Molinspiration Cheminformatics free web services, https://www.molinspiration.com, Slovensky Grob, Slovakia, accessed 9/2023. This work was supported by Oncology Core CA 06973 (B.V., K.W.K., N.P.); The Virginia and D.K. Ludwig Fund for Cancer Research (B.V., K.W.K., N.P., C. B.); The Sol Goldman Sequencing Facility at Johns Hopkins (B.V.); NIH Cancer Center Support Grant P30 CA006973. SP was supported by NCI Grant K08CA270403, the Leukemia Lymphoma Society Translation Research Program award, the American Society of Hematology Scholar award, and the Swim Across America Translational Cancer Research Award.

Author contributions

A.L.C, K.W.K., and N.W. conceptualized the study. A.L.C, S.S., B.V., N.P., C.B., K.G., S.Z., K.W.K., and N.W. developed the methodology. A.L.C., S.S., L.D., E.W., B.P., B.S.L., S.P., E.H., M.P., and N.W. conducted the experiments. A.L.C., S.S., J.D.C., B.S.L., and N.W. performed the formal data analysis. A.L.C, N.W., and K.W.K. wrote the paper.

Declaration of interests

BV, KWK, & NP are founders of Thrive Earlier Detection, an Exact Sciences Company. KWK, NP are consultants to Thrive Earlier Detection. BV, KWK, NP, SZ hold equity in Exact Sciences. BV, KWK, NP, SS and SZ, are founders of or consultants to and own equity in ManaT Bio., Haystack Oncology, Neophore, CAGE Pharma and Personal Genome Diagnostics. NP is consultant to Vidium. BV is a consultant to and holds equity in Catalio Capital Management. SZ has a research agreement with BioMed Valley Discoveries, Inc. CB is a consultant to Depuy-Synthes, Bionaut Labs, Haystack Oncology, Privo Technologies and Galectin Therapeutics. CB is a co-founder of OrisDx and Belay Diagnostics. The

companies named above, as well as other companies, have licensed previously described technologies
related to the work from this lab but not this paper from Johns Hopkins University. BV, KWK, NP, JDC,
and SS are inventors on some of these technologies. Licenses to these technologies are or will be
associated with equity or royalty payments to the inventors as well as to Johns Hopkins University.
Provisional patent applications on the work described in this paper have been filed by Johns Hopkins
University. The terms of all these arrangements are being managed by Johns Hopkins University in
accordance with its conflict of interest policies.

659 Supplemental Materials

660 Figures S1-S21

661 Tables S1-S5

663 STAR methods

664 KEY RESOURCE TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
anti goat 680RD used for Western blot at 1:10K	Licor	926-68074
anti mouse 680RD used for Western blot at 1:15K	Licor	926-68072
anti mouse HRP used for Western blot at 1:2500	Jackson Immuno Research	115-035-006
anti rabbit 800CW used for Western blot at 1:10K	Licor	926-32213
anti rabbit HRP used for Western blot at 1:2500	Jackson Immuno Research	111-035-006
Beta-actin used for Western blot at 1:5000	Cell Signaling Technology	3700S
EXOC1 used for Western blot at 0.4 µg/mL	Abcam	ab251853
LMAN1 used for Western blot at 1:200	ThermoFisher Scientific	CF502137
OKT-3 used for T-Cell Culturing at 15 ng/mL	Biolegend	317347
p21/WAF1 used for Western blot at 1:1000	Cell Signaling Technology	2947S
p53 used for Western blot at 1:1000	Cell Signaling Technology	18032S
phospho-UPF1 used for Western blot at 1:1000	Cell Signaling Technology	2851S
Sodium Potassium Pump (Na-K) used for Western blot at 1:500	Cell Signaling Technology	3010S
SPTAN1 (polyclonal) used for Western blot at 0.04 µg/mL	Bethyl	A301-249A
SPTAN1 (N-Terminal) used for Western blot at 1:1000	Abcam	ab11755
SPTAN1 (C-Terminal) used for Western blot at 1:1000	Abcam	ab75755
STAG2 used for Western blot at 1:1000	Cell Signaling Technology	5882S
UPF1 used for Western blot at 1:2500	Abcam	ab10510
Flow cytometry antibody: α-HLA-A24 used for Flow cytometry at 10 µg/mL	MBL Life Science	Cat #K0208-A64
TP53-CD3 bispecific antibody used for Co-culture at 12.5pg/mL	This Study	N/A
Bacterial and virus strains		

HLA-A24 Retrovirus	Hsiue et al. Science, 2021.	N/A
Luciferase lentivirus	OriGene	Cat #PS100071
Biological samples		
N/A		
Chemicals, peptides, and recombinant proteins		
1% Penicillin-Streptomycin	Thermo Fisher Scientific	Cat #15140122
4–15% Mini-PROTEAN TGX Precast Protein Gels	Bio-Rad	Cat #456-1086
5-Fluroruracil	Sigma Aldrich	Cat #F6627
AMPure beads	Beckman Coulter	Cat #A63880
Cremaphor	Sigma Aldrich	Cat #C5135
DharmaFECT1 transfection reagent	Horizon	Cat #T-2001-02
Dimethyl Sulfoxide (DMSO)	Sigma Aldrich	Cat #C6295
DMEM medium	Gibco	Cat #11995065
EMEM medium	ATCC	Cat #30-2003
Emetine	Sigma Aldrich	Cat #7083-71-8
EPITHELIAL CELL MEDIUM-Complete Kit	Science Cell Research	Cat #4101
Fetal Bovine Serum (FBS)	HyClone	Cat #16777-006
Glycerol	Sigma Aldrich	Cat #G5516
KVS0001	This study	N/A
LY3023414	Selleckchem	Cat #S8322
Matrigel Phenol Red Free Standard Formulation	Corning	Cat #356237
MEGM Mammary Epithelial Cell Growth Medium BulletKit	Lonza	Cat #CC3150
Methylcellulose	Sigma Aldrich	Cat #M6385
Phosphate Buffered Saline (PBS)	Thermo Fisher	Cat # J60465.K2
Phusion Flash High-Fidelity PCR Master Mix	Thermo Fisher	Cat #F548S
Pierce ECL Western Blotting Substrate	Thermo Fisher	Cat #32106
PRIMETIME Gene Expression master mix	Integrated DNA Technologies	Cat #1055770
Protease inhibitor	Millipore Sigma	Cat #4693159001
QIAshredder	Qiagen	Cat #79656
Recombinant IL-2 protein	Prometheus Therapeutics and Diagnostics	Cat #aldesleukin
Recombinant IL-7 protein	BioLegend	Cat #581908
RediJect D-Luciferin Ultra Bioluminescent Substrate	PerkinElmer	Cat #770505
RIPA Lysis and Extraction buffer	Thermo Fisher	Cat #89901
RNA Later	Invitrogen	Cat #AM7020
RPMI 1640 medium	Gibco	Cat #11875-119
Selleckchem Bioactive Compound library (Table S1)	Selleckchem	Cat # L1700
SMG1specific inhibitor: 11j	Ascendex LLC	N/A
SsoAdvanced Universal SYBR Green Supermix	Bio-Rad	Cat #1725270
Trypsin	Gibco	Cat #25300054
Critical commercial assays		
AllPrep DNA/RNA Mini Kit	Qiagen	Cat #80204
Agilent RNA ScreenTape	Agilent	Cat #5067-5576
Agilent RNA ScreenTape Sample Buffer	Agilent	Cat #5067-5577
Agilent RNA Ladder	Agilent	Cat #5067-5578
BCA Protein Assay Kit	Thermo Fisher	Cat #23227

Bio-Rad SingleShot Cell Lysis kit	Bio-Rad	Cat #1725080
ELISA	R&D Systems	Cat #SIF50C
High-Capacity cDNA Reverse Transcription Kit	Advanced Biosystems	Cat #4368814
Illumina RNA library prep kit	Illumina	Cat #RS-122-2001
Kinase kinativ assay	ActivX Biosciences	N/A
Luciferase Assay System	Promega	Cat #E1501
RNeasy	Qiagen	Cat #74104
Deposited data		
Whole transcriptome RNA-seq: NCI-H358	This study	EGAS00001007597
Whole transcriptome RNA-seq: LS180	This study	EGAS00001007597
Whole transcriptome RNA-seq: Isogenic knockout cell lines	Cook, et. al. iScience, 2022.	EGAD00001008559
Experimental models: cell lines		
B16-F10	ATCC	Cat #CRL-6475
HEK293T	ATCC	Cat #CRL-3216
Lewis lung carcinoma (LLC)	ATCC	Cat #CRL-1642
LS180	ATCC	Cat #CL-187
MCF10a	ATCC	Cat #CRL-10317
NCI-H716	ATCC	Cat #CCL-251
NCI-H2228	ATCC	Cat #CRL-5935
NCI-H358	ATCC	Cat #CRL-5807
Renca	ATCC	Cat #CRL-2947
RPE1	ATCC	Cat #CRL-4000
RPE1 STAG2 2	Cook, et. al. iScience, 2022.	N/A
RPE1 STAG2 5	Cook, et. al. iScience, 2022.	N/A
RPE1 STAG2 6	Cook, et. al. iScience, 2022.	N/A
RPE1 STAG2 8	Cook, et. al. iScience, 2022.	N/A
RPE1 TP53 221	Cook, et. al. iScience, 2022.	N/A
RPE1 TP53 223	Cook, et. al. iScience, 2022.	N/A
RPE1 TP53 224	Cook, et. al. iScience, 2022.	N/A
RPTec	ATCC	Cat #CRL-4031
RPTec STAG2 3	Cook, et. al. iScience, 2022.	N/A
RPTec STAG2 943	Cook, et. al. iScience, 2022.	N/A
RPTec STAG2 943A	Cook, et. al. iScience, 2022.	N/A
RPTec STAG2 944	Cook, et. al. iScience, 2022.	N/A
RPTec STAG2 945	Cook, et. al. iScience, 2022.	N/A

RPTec STAG2 946	Cook, et. al. iScience, 2022.	N/A
RPTec STAG2 947	Cook, et. al. iScience, 2022.	N/A
RPTec STAG2 951	Cook, et. al. iScience, 2022.	N/A
RPTec STAG2 952	Cook, et. al. iScience, 2022.	N/A
RPTec STAG2 953	Cook, et. al. iScience, 2022.	N/A
RPTec TP53 544	Cook, et. al. iScience, 2022.	N/A
RPTec TP53 588	Cook, et. al. iScience, 2022.	N/A
Experimental models: Organisms/strains		
Mus musculus BALB/cAnAHsd	Harlan Laboratories	
Mus musculus C57BL/6NCrl	Charles River GmbH	
Mus musculus BALB/cAnNCrl	Charles River GmbH	
Mus musculus Hsd:Athymic Nude-Foxn1[nu]	Harlan Laboratories	
Mus musculus NOD.Cg-Prkdcscid Il2rgtm1Wjl/SzJ (NSG)	Jackson Laboratories	
Oligonucleotides		
for Primers see Table S5	This study	N/A
HTS Screen sequencing primer: STAG2 Seq Forward: AATGATACGGCGACCACCGAGATCTACACTCT TTCCCTACACGACGCTCTTCCGATCTNNNNNNN NGAATTTCTCTACAAAAAGCTCTTCA	This study	N/A
HTS Screen sequencing primer: STAG2 Seq Reverse: CAAGCAGAAGACGGCATACGAGATNNNNNNN NNNTTCATCATTCCATCCTCCTC	This study	N/A
HTS Screen sequencing primer: TP53 Seq Forward: AATGATACGGCGACCACCGAGATCTACACTCT TTCCCTACACGACGCTCTTCCGATCTNNNNNNN NGAAACTACTTCCTGAAAACAACGT	This study	N/A
HTS Screen sequencing primer: TP53 Seq Reverse: CAAGCAGAAGACGGCATACGAGATNNNNNNN NNNGCTTCATCTGGACCTGGGTC	This study	N/A
ON-TARGETplus Human UPF1 siRNA Smartpool 10nmol	Horizon Discovery	Cat #L-011763-00-0010
ON-TARGETplus Human mTOR siRNA Smartpool 10nmol	Horizon Discovery	Cat #L-003008-00-0010
ON-TARGETplus Human ATM siRNA Smartpool 10nmol	Horizon Discovery	Cat #L-003201-00-0010
ON-TARGETplus Human ATR siRNA Smartpool 10nmol	Horizon Discovery	Cat # L-003202-00-0010
SMARTpool: ON-TARGETplus SMG1 siRNA	Horizon Discovery	Cat # L-005033-00-0020
SMARTpool: ON-TARGETplus PRKDC siRNA	Horizon Discovery	Cat # L-005030-00-0010

SMARTpool: ON-TARGETplus PIK3CA siRNA	Horizon Discovery	Cat # L-003018-00-0010
ON-TARGETplus Non-targeting Pool	Horizon Discovery	Cat # D-001810-10
Phage display library	GeneArt	N/A
Recombinant DNA		
N/A		
Software and algorithms		
HISAT2 (version 2.0.5)	Kim et. al.[79]	N/A
StringTie (version 1.3.3)	Pertea et. al.[80]	N/A
Ballgown (version 2.6.0)	Pertea et. al.[80]	N/A
R (version 4.0.3)	R Core Team 2022	N/A
ggplot (version 3.4.1)	https://github.com/tidyverse/ggplot2/blob/HEAD/R/ggplot2-package.R	N/A
Pipeline for analyzing mutant and normal transcript abundance	MSSQL was used for initial data processing	N/A
Other		

RESOURCE AVAILABILITY

Contact for Reagent and Resource Sharing

Further information and requests for resources and reagents should be directed to and will be fulfilled by
the lead contact, Nicolas Wyhs (wyhs@jhmi.edu). Isogenic knockout cell lines are available through The
Genetic Resources Core Facility at Johns Hopkins School of Medicine (jhbiobank@jhmi.edu, Maryland,
USA). KVS0001 generated in this study will be made available on request if available, but availability
may be limited and we may require a payment and/or a completed materials transfer agreement if there is
potential for commercial application.

Data availability

• Whole transcriptome RNA-seq raw and processed data have been deposited at GEO and are
publicly available as of the date of publication. Accession numbers are listed in the Key Resource
Table.
• Full western blot images have been deposited at Mendeley (Mendeley Data, V1, doi:
10.17632/n5s2z393z6.1) and are publicly available as of the date of publication.
• Any additional information required to reanalyze the data reported in this paper is available from
the lead contact upon request.

EXPERIMENTAL MODELS AND SUBJECT DETAILS

Cell lines
NCI-H716, NCI-H2228, HEK293, NCI-H358, RPE1, MCF10a, RPTec, LLC, RENCA, B16-F10 and
LS180 cells were purchased from The American Type Culture Collection (Virginia, USA). RPE1 STAG2
2, RPE1 STAG2 5, RPE1 STAG2 6, RPE1 STAG2 8, RPTec STAG2 3, RPTec STAG2 943, RPTec
STAG2 943A, RPTec STAG2 944, RPTec STAG2 945, RPTec STAG2 946, RPTec STAG2 947, RPTec

STAG2 951, RPTec STAG2 952, RPTec STAG2 953, RPE1 TP53 221, RPE1 TP53 223, RPE1 TP53
224, RPTec TP53 54, RPTec TP53 588, RPTec TP53 544 isogenic knockout cell lines were generated
and grown as previously described (Cook et al, iScience 2022)[31]. NCI-H358, NCI-H2228, NCI-H716,
RENCA and RPE1 cells were grown in RPMI 1640 Medium (Gibco, California, USA, Cat #11875-119)
supplemented with 10% Fetal Bovine Serum (FBS) (HyClone, Utah, USA, Cat #16777-006). LS180 cells
were grown in EMEM (ATCC, Virginia, USA, Cat #30-2003) supplemented with 10% FBS. LLC, B16-
F10 and HEK293 was grown in DMEM (Gibco, USA, Cat #11995065) supplemented with 10% FBS.
RPTec cells were grown in EPITHELIAL CELL MEDIUM-Complete Kit (Science Cell Research,
California, USA, Cat #4101). MCF10a cells were grown in MEGM Mammary Epithelial Cell Growth
Medium BulletKit (Lonza, USA, Cat #CC3150). *In vitro,* all cells were grown at 37°C with 5% CO_2.
Mycoplasma testing was performed by The Genetic Resources Core Facility at Johns Hopkins School of
Medicine (Maryland, USA).

Quantitative real-time PCR (qPCR)
RNA was obtained from cells using a Qiagen RNeasy Kit (Qiagen, Maryland, USA, Cat #74104) per the
manufacturer's instruction. Reverse transcription of RNA to cDNA was performed using a High-Capacity
cDNA Reverse Transcription Kit (Applied Biosystems, USA, Cat #4368814) per manufacturer's
instructions. Unless otherwise specified, qPCR reactions were set up using SsoAdvanced Universal
SYBR Green Supermix (Bio-Rad, USA, Cat #1725270) following manufacturer's instructions and
performed on a QuantStudio 3 (Applied Biosystems, USA) with manufacturer's recommended plates and
plate covers. PCR thermocycling conditions were as follows: 2 min at 50°C, 2 min at 95°C, 35 cycles of
10 sec at 95°C, 10 sec at 60°C, 30 sec at 72°C. Finally a melt curve was performed starting at 50°C and
ending at 95°C with 5-second incubations for imaging. For TP53β analysis, qPCR reactions were set up
using PRIMETIME™ Gene Expression master mix (Integrated DNA Technologies, USA, Cat #1055770)
and a probe with FAM reporter and TAMRA quencher (Integrated DNA Technologies, USA) following
the manufacturer's instructions and concentrations. PCR thermocycling conditions were as follows: 3 min
at 50°C, 10min at 95°C, 40 cycles of 10 sec at 95°C, 30 sec at 60°C. Primers used for all qPCR can be
found in Table S5.

Small molecule compounds
SMG1 specific inhibitor SMG1i11 (11j)[30] and KVS0001 were synthesized by Ascendex LLC
(Pennsylvania, USA). The synthesis scheme for KVS0001 is located below. Emetine was obtained from
Sigma Aldrich (Cat #7083-71-8). All other small molecule hits from the screen were purchased from
Selleckchem (Texas, USA).

NMD screen

We screened the Selleckchem Bioactive Compound library, a 2,658 compound library (Selleckchem, Texas, USA, Cat #L1700). The library was distributed over 33 x 96-well tissue culture plates with each plate containing library compounds, 2 NMD positive controls (emetine, 12 μg/mL) and 6 NMD negative controls (DMSO). We mixed the three cell lines, RPE1 STAG2 2, RPE1 STAG2 8, and RPE1 TP53 221, in equal ratio and dosed with the compound library at 10 μM for 14 hours. Cell pools were harvested by washing plates 2x with PBS and frozen. RNA extraction and reverse transcriptase cDNA conversion were performed as described in *Targeted RNA-seq.* Quality control was performed by qPCR of one NMD positive and one NMD negative control well from each plate to confirm that increased expression of the positive control (emetine) wells was observed. All samples were prepared for sequencing by amplifying cDNA using primers containing both a plate (forward primer) and well (reverse primer) barcode which amplifies the knockout cell line specific mutation for STAG2 and TP53 (See Table S5). Primers were obtained from Integrated DNA Technologies (Iowa, USA). PCR was performed using Phusion Flash High-Fidelity PCR Master Mix (Thermo Fisher, USA, Cat #F548S) for 1 min at 98 °C, 29 cycles of 10 sec at 98 °C, 15 sec at 64 °C, 15 sec at 72 °C, and 5 min at 72 °C. Samples were then well barcoded using a similar PCR setup for 2-4 cycles. Samples were then pooled, cleaned up with AMPure beads (Beckman Coulter, California, USA, Cat #A63880), and sequenced on an Illumina HiSeq2500 using manufacturer's instructions (150 cycle single read) for an average of 56,753 reads per well. The screen was scored by calculating the number of sequencing reads matching each mutant and reference transcript and calculating a Mutant Allele Fraction (MAF) correcting for the number of cell lines and heterozygous mutations. NMD positive and negative control data were pooled, respectively, and averaged across all plates to determine the hit threshold of MAF > 0.46. This value, 5 standard deviations above the mean for the

DMSO controls, was chosen to ensure no false positives in the DMSO controls were observed. A
compound was considered a hit if the MAF for all three cell lines was greater than this value. All results
from the screen are reported as MAF.

Targeted RNA-seq
RNA extraction of *in vitro* cells was performed using BioRad SingleShot Cell Lysis kit (Bio-Rad,
California, USA, Cat #1725080) scaled down to 50 µl reactions per well of 96-well plate per
manufacturer's instructions. Tissue culture cells were lysed directly on the tissue culture plate. For RNA
extraction of *in vivo* studies, tissues were harvested and placed in RNA Later (Invitrogen, Maryland,
USA, Cat # AM7020) and stored at -80°C until RNA extraction. RNA extraction of tissues was performed
using a Qiagen RNeasy Kit (Qiagen, Maryland, USA, Cat #74104) per manufacturer's instruction with
tissue homogenization in 600 µl of RLT buffer via dounce homogenizer followed by QIAshredder
(Qiagen, Maryland, USA, Cat #79656). All RNA quality was assessed by Agilent Tapestation 2200
(Agilent, California, USA, Cat #G2964AA) and the Agilent RNA ScreenTape (Agilent, California, USA,
Cat #5067-5576) with Agilent RNA ScreenTape Sample Buffer and Ladder (Agilent, California, USA,
Cat #5067-5577, Cat #5067-5578) per manufacturer's instruction. Reverse transcription to cDNA was
performed using High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems, USA, Cat
#4368814) per the manufacturer's instructions. cDNA was amplified using cDNA-specific primers with
at least one primer (forward or reverse) covering an exon-exon boundary (See Table S5 for primer
sequences). Primers were obtained from Integrated DNA Technologies (Iowa, USA). PCR was performed
using Phusion Flash High-Fidelity PCR Master Mix (Thermo Fisher, USA, Cat #F548S) for 1 min at 98
$^{\circ}$C, 30 cycles of 10 sec at 98 $^{\circ}$C, 15 sec at 64 $^{\circ}$C, 15 sec at 72 $^{\circ}$C, and 5 min at 72 $^{\circ}$C. Sequencing libraries
were prepped from samples by addition of well barcodes using the same method described above for an
additional 2-6 cycles. Libraries were pooled, cleaned up with AMPure beads (Beckman Coulter,
California, USA, Cat #A63880) and sequenced on an Illumina Miseq using manufacturer's instructions
(150 cycle single read).

Mutant Allele Fraction (MAF) analysis
Mutant allele fraction was determined by processing fastq files using HISAT2 (version 2.0.5) and
aligning to a pseudo reference genome consisting of only the mutant or wild-type amplicon sequences for
targeted regions. The mutant allele fraction was determined by taking a ratio of the number of mutant
transcripts to the total number of transcripts from the region in question. Initial data processing was
performed in MSSQL and Excel.

Whole transcriptome RNA-seq
LS180 or NCI-H358 cells were run in biological duplicate, treated with DMSO or 5µM LY3023414. For
RNA extraction, cells were pelleted, frozen in liquid nitrogen, and stored at -80°C until RNA extraction.
RNA extraction was performed using a Qiagen AllPrep DNA/RNA Mini Kit (Qiagen, Maryland, USA,
Cat# 80204) per manufacturer's instruction with cell homogenization and lysis in RLT buffer with a
QIAshredder (Qiagen, Maryland, USA, Cat# 79656). RNA quality control using Agilent Tapestation
2200 (Agilent, California, USA, Cat# G2964AA) and the Agilent RNA ScreenTape (Agilent, California,
USA, Cat# 5067- 5576) with Agilent RNA ScreenTape Sample Buffer and Ladder (Agilent, California,
USA, Cat# 5067- 5577, Cat# 5067- 5578) per manufacturer's instruction. Library prep using Illumina
RNA library prep kit (Illumina, California, USA, Cat #RS-122-2001) and sequenced on an Illumina
HiSeq 4000 150 cycle paired-end using manufacturer's instructions.

RNA-seq analysis
Sequencing reads aligned to Hg38 using HISAT2 (version 2.0.5), RNA alignment metrics using
CollectRnaSeqMetrics (Picard, version 2.20.2). Exon skipping was determined using IGV Viewer
Sashimi Plots[81]. The average number of bases sequenced per sample and percent aligned in LS180 was
5.14e9 bases (range 5.07e9-5.20e9) and 77.8% (range 76.1%-78.7%) and for NCI-H358 was 5.58e9 bases
(5.19e9-5.87e9) and 79.4% (range 77.2%-82.7%). Mean allele frequency was determined using VarScan
2[82] by generating the ratio of Read 1 (mutant) to read2 (wild type) transcripts. Mutations were only
considered if they were heterozygous, and contained at least 5 reads at the somatic mutation or indel site
in all 4 samples being compared (biological duplicates of treated and untreated).

Compound response curves
We performed a 6-point dose-response curve by treating cell pools for 14 hours with single compounds.
Cell pools consisted of three isogenic knockout cell lines grouped by parental cell line with readout via
targeted sequencing RNA (see *Targeted RNA-seq* for details). The effect of the compound was
determined by calculating the MAF by comparing the abundance of the expected mutation and compared
to the wild type within each isogenic pool (see *Mutant Allele Fraction (MAF) analysis* for details).

Immunohistochemistry (IHC)
IHC was performed on cell lines as previously described[83].

Western blots
Cells were lysed using RIPA buffer (Thermo Fisher, USA, Cat #89901) containing 1x protease inhibitor
(Thermo Fisher, USA, Cat #4693159001) on ice for 30 minutes. Samples were then centrifuged at max
speed for 3 minutes at 4°C in a QIA shredder (Qiagen, Maryland, USA, Cat #79654) before being
transferred to a new sample collection tube. Protein was quantified using a BCA assay (Thermo Fisher,
USA, Cat #23227) per the manufacture's instructions. Gels were run by loading 50 µg of total protein per
sample into 15 well polyacrylamide gels (Bio-Rad, California, USA, Cat #456-1086) and run for 30
minutes at 200V. Gels were then transferred using the manufacturer's instructions (based on size) to
nitrocellulose membrane using a Bio-Rad turbo transfer apparatus (Bio-Rad, USA, #170-4270).
Membranes were blocked for 1 hour with 3% milk-TBS-Tween before being incubated overnight in
primary antibody (concentration dependent on antibody). Primary antibodies and concentrations can be
found in the key resources table. Membranes were washed 4 times for 5 minutes each with TBS-Tween.
Secondary antibody was applied at 1:2500 using either α-rabbit (Jackson ImmunoResearch, Pennsylvania,
USA, Cat #111-035-006) or α-mouse (Jackson ImmunoResearch, Pennsylvania, USA, Cat #115-035-
006). Membranes were imaged using Pierce™ ECL Western Blotting Substrate (Thermo Fisher, USA,
Cat #32106) following the manufacturer's instructions on a Bio-Rad Chemidoc (Bio-Rad, California,
USA). Phospho-UPF1 westerns were performed as detailed above with the following exceptions:
secondary antibodies were used at 1:10,000 either α-rabbit (Licor Biosciences, USA, Cat #926-32213) or
α-goat (Licor Biosciences, USA, Cat #926-68074). Membranes were imaged using an Odyssey CLx
(Licor Biosciences, USA) following the manufacturer's instructions.

siRNA
We performed knockdown of kinase proteins using siRNA. LS180 and NCI-H358 cells were plated in 96-
well plates (Costar, USA, Cat #3595) at 5,000 cells per well. Cells were transfected with DharmaFECT1
transfection reagent (Horizon, USA, Cat #T-2001-02) and either 50 nM control or kinase-specific pooled
siRNA. siRNA used in this experiment can be found in the key resources table. Knockdown efficacy was
confirmed using qPCR (see *Quantitative real-time PCR (qPCR)* for details). NMD target gene mutation

transcript levels were determined using next-generation sequencing and MAF (see *Targeted RNA-seq* for
details).

Bispecific scFv construction
We utilized a bispecific antibody against CD-3 and TP53 wild-type peptide 'TYSPALNKMF' (residues
125-134) presented in a HLA-A24 MHC-I molecule scFv. This bispecific antibody was identified by
panning a phage display library. The scFv-bearing phage library was constructed similarly as described in
detail previously with some modifications[84]. Briefly, oligonucleotides were synthesized by GeneArt
(Thermo Fisher, USA) using trinucleotide mutagenesis (TRIM) technology to diversify complementarity-
determining region (CDR)-L2, CDR-L3, CDR-H1, CDR-H2, and CDR-H3. A FLAG (DYKDDDDK)
epitope tag was placed immediately downstream of the scFv, which was followed in frame by the full-
length M13 pIII coat protein sequence. The total number unique clones obtained was determined to be 3.6
x 10^{10}. Panning details can be found in the reference section[83,84].

Kinase target assay
843   Kinase kinativ experiments were performed by ActivX Biosciences (San Diego, USA)[52].

Quantitative presentation of HLA-bound neoantigens via HPLC-Mass Spectrometry
Identification and quantitation of HLA-presented neoantigens was performed as previously described by
Complete Omics Inc.[55]. Briefly, cells were treated *in vitro* for 24 hours with either DMSO or 5 µM
KVS0001. Cells were cross-linked and immunoprecipitated with pan-HLA antibodies to obtain cell
surface MHC-presented peptides. Mass spectrometry was performed in the presence of heavy labeled
peptide to serve as an internal (loading) control to quantify the presence of the presented neoantigen. The
presented peptides were identified in a preliminary MS screen of 187 candidate peptides predicted by
using the union of NetMHC and Predictor of Immunogenic Epitopes (PRIME) predictions[56,57].

CD3-TP53scFv bispecific co-culture
Co-culture of bispecific antibody was performed using volunteer human donor T-Cells. T-cell enrichment
and activation was performed as previously described [85]. Briefly, PBMCs are incubated with OKT3
antibody (Biolegend) for 3 days. T-cells were then expanded in RPMI-1640 with 10% FBS and 1%
penicillin-streptomycin, recombinant IL-2 (Proleukin, Prometheus Laboratories) and IL-7 (BioLegend)
for at least 15 days before use. NCI-H716 cells were labeled with HLA-A24 using retrovirus as
previously described[83]. Briefly, the MSCV retroviral expression system (Clontech, USA, Cat #634401)
was used to overexpress HLA-A*24-T2A-GFP in target cells. Expression was confirmed by flow
cytometry. NCI-H2228 cells were not labeled as they express this specific HLA endogenously. For NCI-
H716_A24 and NCI-H2228, 40,000 cells were plated in 96-well plates, and co-incubated with 40,000
activated T-cells, and 12.5 pg/mL of TP53-CD3 bispecific antibody. Cells were then dosed with 5 µM of
KVS0001, SMG1 specific inhibitor or 200 mg/mL of 5-fluoruracil (Sigma Aldrich, USA, Cat #F6627) for
24 hours. Readout of interferon (IFN-γ) was performed by ELISA following the manufacturer's
instructions (R&D Systems, USA, Cat #SIF50C). Cell killing was assayed by luciferase levels following
the manufacturer's instructions (Promega, USA, Cat #E1501) and read out on a Synergy H1 Microplate
reader (BioTek, USA). NCI-H716_A24 cells were labeled with luciferase via lentiviral transduction
following the manufacturer's recommendations (OriGene, USA, Cat #PS100071).

Animal protocols
Animal research was approved and overseen by Johns Hopkins University Institutional Animal Care and
Use Committee (ACUC) approved research protocol M018M79. Mice are housed in individually
ventilated caging (Allentown, New Jersey, USA) at a maximum 5 animals per cage. Cages are changed

every 14 days. Enrichment is provided through paper bedding, paper hut, and some food placed in the
bottom of the cage. Facility is maintained between 70 and 72 degrees Fahrenheit on a 12-hour light-dark
cycle. Mice standard diet is ad lib Teklad Global 18% Protein Extruded Rodent Diet, autoclaved (Envigo,
Huntingdon, UK) and acidified water via sipper tube.

In Vivo tumor models

Cells for tumor inoculation in each mouse were grown to 70-80% confluency *in vitro*. Cells were
harvested with trypsin (Gibco, California, USA, Cat #25300054) and suspended in either PBS for LS180
cells or 50% PBS, 50% Matrigel Phenol Red Free Standard Formulation (Corning, New York, USA, Cat
#356237) for NCI-H358 cells. For both cell lines 1e6 cells were placed subcutaneously and grown to
approximately 200 mm^3. Animals were randomized into treatment groups prior to treatment by tumor
size. All tumor volumes were measured via caliper twice weekly. Mice with tumors <150 mm^3 at 11 days
post inoculation were excluded from experiments.

Human xenograft experiments

6 to 8 weeks old *Mus musculus* Hsd:Athymic Nude-Foxn1^{nu} mice (referred to as nude mice) were
purchased from Harlan Laboratories (Indiana, USA). Only female mice were used as gender was not
considered to be a significant confounder in the experiment. LS180 cells were inoculated at 1.0×10^6 cells
per mouse in the left mouse flank or NCI-H358 cells were inoculated at 7.5×10^5 cells per mouse in the
right flank. For single-dose experiments, mice were orally dosed via gavage at 0, 40, or 60 mg/kg of
LY3023414 (Selleckchem, Texas, USA, Cat #S8322) in 1% Methylcellulose (Sigma Aldrich, Missouri,
USA, Cat #M6385). KVS0001 was dosed at 30 mg/mL intraperitoneal (IP) in 0.5% Dimethyl sulfoxide
(DMSO)(Sigma Aldrich, USA, Cat #C6295), 10% cremaphor (Sigma Aldrich, USA, Cat #C5135) and
2% glycerol (Sigma Aldrich, USA, Cat #G5516). Mice were given physical exams prior to euthanasia at
designated endpoints according to the approved research protocol. Tumors, spleen, blood, and lungs were
harvested post-euthanasia for further analysis. Tumors were harvested at indicated times and stored in
RNA Later (Invitrogen, Maryland, USA, Cat # AM7020) at -80 for further analysis.

Syngeneic mouse tumor experiments

6 to 8 weeks old immune compromised *Mus musculus* BALB/cAnNCrl (referred to as BALB/c) and
C57BL/6NCrl (referred to as C57BL/6N) mice were purchased from Charles River GmbH (Germany).
Only female mice were used as gender was not considered to be a significant confounder in the
experiment. Murine cancer cell lines were placed in the left mammary fat pad of female mice at the
following densities: LLC 1.0×10^6, RENCA 1.0×10^6, and B16-F10 0.2×10^6. Mice were randomized
around day 7 post-implantation when tumor sizes had reached approximately 30-40 mm^3. Mice were
dosed with 30 mg/kg KVS0001 or vehicle control via IP injection alternating the right and left side daily
for 28 days (Figure 5C). Mice were given physical exams before euthanasia at designated endpoints
according to the approved research protocol.

In Vivo TP53 bispecific experiments

For the NCI-H716 *in vivo* bispecific experiments 2.5×10^6 NCI-H716_A24 expressing cells were placed
orthotopically by IP injection into NOD.Cg-Prkdcscid Il2rgtm1Wjl/SzJ (referred to as NSG) mice
(Jackson Laboratories, USA). On day 2 mice were randomized to ensure a balanced tumor burden in all
groups. Luminescence was measured by injecting mice with 150 µL of RediJect D-Luciferin Ultra
Bioluminescent Substrate (PerkinElmer, USA, Cat #770505), and anesthetized using isoflurane in an
induction chamber for 5 minutes. Readouts and analysis were performed on an IVIS Spectrum imaging
system and Living Image software (Perkin Elmer, USA).

 Data reporting
No statistical methods were used to predetermine the sample size. Statistical analysis was performed
using R and Excel. All animal experiments were randomized being sure to alter the first cage dosed and
location of cages with a row of the rack. Randomization was performed by animal tumor size, unless
otherwise indicated. The investigators responsible for weight and tumor measurements were blinded to
the allocation, treatment, and outcome assessment of experiments. The investigators processing mouse
tissue processing were blinded to the allocation, treatment, and outcome assessment of experiments. The
investigator harvesting the tissues was aware of the allocation, treatment, and outcome assessment of
experiments.

Statistical Testing
Chi-Squared testing was performed using counts of cancer cell line mutations predicted to undergo NMD
and RNA recovered and prop.test() in R. Mann-Whitney testing performed with wilcox.test() in R.
Quantiles calculated utilizing z scores with quantile() in R. Student T-Test and one-way ANOVA with
Dunnett´s test and Student's t-test were performed in R and Excel.