Common Ancestry of the *Id* Locus: Genetic Clues and Polygenic Possibilities

Ashutosh Sharma, Nagarjun Vijay*

Computational Evolutionary Genomics Lab, Department of Biological Sciences, IISER Bhopal, Bhauri, Madhya Pradesh, India

*Corresponding author: nagarjun@iiserb.ac.in

Abstract

The diversity of dermal pigmentation and plumage color in domestic chickens is intriguing.

Black bone chickens (BBC) stand out with their intense melanin hyperpigmentation, driven by a

complex chromosomal rearrangement on chromosome 20, specifically the *Fm* locus, causing

overexpression of the *EDN3* gene. In contrast, dermal pigmentation inhibition is regulated by the

sex-linked *Id* locus. Previous studies have used crosses, GWAS and gene expression analysis to

identify the genetic basis of the *Id* locus, but the puzzle remains unsolved. Our study seeks to

precisely locate the *Id* locus, identify gene(s) inhibiting dermal pigmentation, and explore its

ancestral state in BBC breeds. We analyzed public data from 270 chickens (42 BBC and 228

non-BBC) to refine the position of the *Id* locus to a ~1.6 Mb region between Z amplicon repeats

on the distal end of q-arm and found that it shares a common ancestry among all BBC breeds,

much like the *Fm* locus. The selection signatures and high fixation levels within *MTAP,*

CCDC112 and a few other genes within this region support multiple candidates. Our results

suggest that the *Id* locus is shared among BBC breeds and may have a polygenic nature,

explaining the shank and dermal pigmentation variation.

Keywords: *Id* locus, Black-bone chickens, Dermal pigmentation, Chromosome Z, Polygenic

Introduction

Phenotypic diversity in pigmentation patterns within avian species has long been a focal point of

biological research. The chicken has garnered considerable attention due to its economic and

culinary significance and the remarkable variety of plumage colors and dermal pigmentation

exhibited across different breeds. The most striking of such pigmentation phenotypes,

fibromelanosis (*Fm*), refers to hyperpigmentation throughout the tissues and organs,

encompassing the dermal layer of skin, bones, muscles, nerves, and gonads of black-bone

chicken (BBC) (Dorshorst et al., 2011; Shinomiya et al., 2012). The underlying genetic cause of

this hyperpigmentation, often referred to as the "black bone phenotype," was first investigated by

Bateson and Punnett using the Silkie BBC breed (Bateson et al., 1911). Genetic crossing

experiments identified two independent epistatically interacting factors influencing the degree of

pigmentation observed in hybrid crosses between Silkie and White Leghorn chickens (Arora et

al., 2011; Dunn and Jull, 1927; Tian et al., 2014a). These two genetic factors responsible for

dermal pigmentation are termed "fibromelanosis (*Fm*) locus" and "inhibitor of dermal melanin

(*Id*) locus" (Mukherjee et al., 1969; Tixier-Boichard, 2002).

The genetic basis of the *Fm* locus has been identified on chromosome 20 as a complex

chromosomal rearrangement encompassing three different non-paralogous genomic regions,

Dup1, Int, and Dup2, with the Dup1 and Dup2 regions being duplicated (Dharmayanthi et al.,

2017; Dorshorst et al., 2011; Shinde et al., 2023; Shinomiya et al., 2012; Sohn et al., 2018). The

most important region Dup1 contains five protein-coding genes: *EDN3*, *ZNF831*, *SLMO2*,

ATP5E, and *TUBB1*. The prominent factor contributing to the intense melanization observed in

the internal organs of BBC is the heightened expression of the Endothelin 3 gene (*EDN3*), which

plays a pivotal role in producing melanocytes, specialized cells responsible for melanin synthesis

(Baynash et al., 1994). The increased expression of *EDN3* resulting from the rearrangement is a

key determinant in the hypermelanization phenomenon (Dorshorst et al., 2011). Notably, various

BBC breeds from different geographic regions across Asia exhibit this distinctive chromosomal

rearrangement at the *Fm* locus. The shared rearrangement and the population genetic signature of

this region suggest a common ancestral origin for all BBC breeds (Shinde et al., 2023).

In contrast to the autosomal dominant *Fm* locus, the mode of inheritance of *Id* is sex-linked

incomplete dominance. As its name suggests, the *Id* locus is known to inhibit dermal

pigmentation to different degrees based on its genetic state. Gene mapping studies have

identified the physical location of the *Id* locus at the distal end of the q-arm of chromosome Z, in

proximity to the sex-linked barring loci B (Bitgood, 1988, 1985; Dorshorst and Ashwell, 2009).

The availability of genome assemblies helped identify that the chicken Z chromosome has a

distinctive genetic structure known as the Z amplicon repeat unit at the distal end of the q-arm.

The Z amplicon repeats consist of multiple copies of four genes: *ADCY10*, *C2orf3*, *ARHGAP33*,

and *MRPL19* (Bellott et al., 2010; Yan et al., 2010). Notably, parallel research on identifying the

Id locus relying upon early draft genomes confirmed its localization at the distal end of the q-arm

of chromosome Z (Dorshorst et al., 2010). Over time, with advancements in genomics and the

availability of more comprehensive chicken genome data, the approximate position of the *Id*

locus was refined. In recent years, utilizing a genome-wide SNP-trait association analysis,

researchers pinpointed a potential *Id* locus in the galGal4 genome assembly, situated

approximately at 78.4 Mb on chromosome Z (Dorshorst et al., 2010; Li et al., 2014; Siwek et al.,

2013). Several studies have been undertaken involving crosses between BBC and non-BBC

breeds, all in pursuit of identifying the candidate *Id* gene. These investigations have reported

various potential genes which are associated with dermal and shank pigmentation variations

(Arora et al., 2011; Cha et al., 2023; Li et al., 2014; Mcgibbon, 1978; Tian et al., 2014b; Xu et

al., 2017).

Numerous studies have been conducted to unravel the mysteries of the *Id* locus. Yet, the puzzle

surrounding this locus persists, and the precise genomic region or gene responsible for its

function remains elusive. With recent advancements in the quality of the chicken genome and the

availability of population genomic data from various BBC and non-BBC breeds, a newfound

opportunity exists to conduct a comprehensive evaluation and accurate identification of the *Id*

locus. Our study aims to address the following objectives: **1)** Assess the co-localization of Z

amplicon repeats and *Id* locus along the Z chromosome. **2)** Evaluate the ancestral relationship at

the *Id* locus among different BBC breeds. **3)** Identify BBC-specific genomic signatures within

the *Id* locus using population genomic statistics. **4)** Determine genomic variants associated with

dermal pigmentation and plumage.

Results

Id locus is located between Z amplicon repeat units

Using the BLASTN and read coverage approaches, we have refined the precise positions of Z-

amplicon repeat units (located at Z:73613150-82529921) situated at the distal end of the q-arm

of chromosome Z within the Gallus_gallus.GRCg6a genome assembly (**Fig. 1**). We have

identified two regions containing single-copy protein-coding genes, R1 and R2, within this

region, which divide the Z amplicon repeat units into three distinct segments. R1, spanning from

Z:78725000 to Z:80340000, extends over 1.61 Mb and harbors ten genes (*MTAP*, *CDKN2A*,

CDKN2B, *TRIM36*, *PGGT1B*, *CCDC112*, *FEMC1*, *ALDH7A1*, *GRAMD2B*, and *ZNF608)*.

Similarly, R2, spanning from Z:81040000 to Z:82226500, encompasses a length of 1.18 Mb and

hosts more than 15 protein-coding genes. Previous research (Cha et al., 2023; Li et al., 2014;

Tian et al., 2014b; Xu et al., 2017), particularly concerning dermal pigmentation and plumage

color variation, has highlighted several candidate genes predominantly located within the R1

region. Considering these earlier studies and our findings (discussed in the subsequent section), it

becomes evident that the *Id* locus is co-localized with the Z-amplicon repeat units at the distal
end of the q-arm of chromosome Z.

BBC breeds share common ancestry at *Id* locus

Our higher-resolution identification of the *Id* locus is based on a comprehensive analysis of local
phylogeny, principal component analysis (PCA), and admixture analysis of the R1 and R2
regions. Our findings suggest that R1 represents the *Id* locus for the following reasons: The
genetic structure of the R1 region demarcates BBC from non-BBC breeds (**Fig. S1-10**). In stark
contrast, when examining the R2 region, we observe an intermingling of BBC and non-BBC
breeds (**Fig. S11-15**). This intermixing implies a lack of distinct differentiation between these
two groups in the R2 region. The local phylogeny analysis of the R1, based on single nucleotide
polymorphisms (SNP) data from 270 individuals, provides clear insights into the genetic
relationships among different chicken breeds. As illustrated in **Fig. 2A** and **Fig. S1-4**, this
analysis distinctly separates BBC from non-BBC individuals. Notably, a few individuals from
the Tibetan native chicken population exhibit genetic proximity to Tibetan BBC individuals,
suggesting potential gene flow.

Principal component analysis (PCA) of the R1 region further elucidates the genetic structure,
with PC1 and PC2 accounting for 19.8% and 9.1% of the genetic variation, respectively. Notably,
the BBC individuals form a distinct cluster, reinforcing the patterns observed in the local
phylogeny analysis (**Fig. 2B** and **Fig. S5-7**). An admixture analysis of the R1 region, aimed at
elucidating ancestry information, reveals that BBC individuals are most distinct when the best
value of K (genetic clusters) is set to 6. Non-BBC native breeds exhibit limited genetic overlap
with BBC and commercial breeds, as depicted in **Fig. 2C** and **Fig. S8-10**.

BBC-specific variants and genomic regions under selection

We employed two distinct approaches to find candidate genes or genomic regions associated
with dermal pigmentation. In the first approach, we focused on identifying selective sweeps
specific to BBC by leveraging various population genetic statistics. This analysis covered the
distal end of the q-arm (Z:72000000-82529921). We sought to isolate the top 10% F_{ST} windows,
derived from pairwise comparisons between BBC and non-BBC breeds, as potential regions

subject to selection. To designate a region as being under selection, we established specific

criteria. F_{ST} values were required to be less than 0.5 in comparisons involving BBC breeds, while

in comparisons between BBC and non-BBC breeds, F_{ST} values should exceed 0.8. Furthermore,

the selected region must fall within the top 10% of weighted F_{ST} windows. This initial approach

led us to identify numerous regions displaying substantial differentiation in BBC vs. non-BBC

F_{ST} comparisons while at the same time having lower differentiation in BBC vs. BBC F_{ST}

comparisons (**Fig S16-18** and **Table S1** for detailed results).

In contrast to the signatures of selection, preexisting or standing genetic variation and breed-

specific mutations arising from genetic drift can drive rapid evolution and influence phenotypic

diversity and adaptation (Barrett and Schluter, 2008). Consequently, our second approach

investigates breed-specific genomic variants within BBC and non-BBC breeds. Our analysis has

revealed the presence of numerous genomic variants in both genic and intergenic regions that are

nearly fixed in both BBC and non-BBC breeds (**Table S2**).

Putative BBC-specific selective sweep in the *MTAP* gene

Our genomic scan of R1 has unveiled distinct regions specific to BBC, characterized by a

weighted F_{ST} exceeding the predefined threshold. As depicted in **Fig. S16** and **Table S1**, one

particularly noteworthy region spans five Kb and is situated within the last intron of the *MTAP*

gene. What sets this 5 Kb region apart is its markedly lower F_{ST} values in BBC vs. BBC

comparisons compared to BBC vs. non-BBC comparisons, as evident in **Fig. 3A-D**. To further

corroborate our findings, we examined D_{xy} results within the same region, and they reinforced

the observed F_{ST} pattern, displaying striking differences, as depicted in **Fig. 3E-G**. Assessment

of the evolutionary conservation of this region through the phastCons conservation score

suggests a level of noteworthy conservation, as illustrated in **Fig. 3H**. Within this 5 Kb region of

the *MTAP* gene, we identified 6 BBC-specific SNPs, detailed in **Table S2**. Notably, the bam-read

count data across 270 individuals underscore that these SNPs are nearly ubiquitous among BBC

individuals, irrespective of their geographic origins. A barplot depicting the percentage

composition of each base at these six positions distinctly highlights the substantial difference in

base composition between BBC and non-BBC individuals, as showcased in **Fig. 3I**.

BBC-specific variants in the promoter region of the *CCDC112* gene

Within the promoter region of the *CCDC112* gene, we identified three highly differentiated SNPs located at positions Z:79049203, Z:79049207, and Z:79049235. Our analysis across 42 BBC individuals found that ~83% are homozygous for the alternative allele, ~9% are heterozygous, and ~7% possess the reference allele. To provide a more accessible and insightful visualization of these findings, we've generated a barplot illustrating the bam-read count results for these three positions, as showcased in **Fig. 4**. In the case of the 228 non-BBC individuals, ~72% carry reference alleles, ~12% are heterozygous, and ~ 14% possess alternative alleles, akin to those observed in BBC individuals. These large differences in allele frequency underscore the genetic divergence in this gene between BBC and non-BBC breeds.

microRNAs in BBC-specific region target melanogenesis genes

We identified several other genomic regions upstream of the Z amplicon repeat units with a selection signature in BBC breeds (**Table S1**). Among these, we found a 26 Kb continuous intergenic region spanning Z:72580000-72606000 with a remarkable pattern in pairwise F_{ST} and D_{xy} comparisons, as depicted in **Fig. 5A-G**. The nucleotide diversity (π and Watterson's θ) within this region is higher in White Leghorn chickens. However, Kadaknath and Silkie chickens have markedly reduced nucleotide diversity, as visualized in **Fig. 5H-I**. An examination of the phastCons conservation score for this region indicated a notably high level of sequence conservation, as depicted in **Fig. 5J.**

Although this specific region lacks any protein-coding genes, we uncovered five pre-microRNA precursors highly similar to gga-mir-1457, gga-mir-6702, gga-mir-6551, gga-mir-6555, and gga-mir-6552 (**Fig. S19**). Of these five miRNAs, we found distinct BBC-specific SNPs within the paralogs of gga-mir-1457 and gga-mir-6552, which are predicted to target several genes (**Table S3,4**). Gene enrichment analysis of the targets of these two miRNAs revealed multiple target genes associated with various signaling pathways, including the GnRH signaling pathway, Apelin signaling pathway, Insulin signaling pathway, adrenergic signaling pathway, Wnt signaling pathway, and Hedgehog signaling pathway (**Fig. 5K, L,** and **Fig. S20-21**). Notably, our analysis in the context of gene enrichment unveiled an intriguing finding: 30 genes integral to the melanogenesis pathway (see **Table S5**) serve as targets for these miRNAs, underscoring their potential role in regulating pigmentation-related processes.

Discussion

Our study has refined the position of the *Id* locus and identified several candidate genes responsible for inhibiting dermal melanin. We demonstrate that the *Id* locus is co-localized among Z amplicon repeats and occurs within a region of ~1.61 Mb at the distal end of the q-arm. The phylogenetic and population structure analysis of 270 individuals (42 BBC and 228 non-BBC) suggest that the *Id* locus has common ancestry among all BBC breeds. Within the refined region of the *Id* locus, we identified the BBC-specific regions putatively under selection. The two candidate genes, *MTAP* and *CCDC112*, involved in the melanogenesis pathway and plumage color variation, have BBC-specific SNPs in the intronic and promoter regions, respectively. The findings of our study suggest that the *Id* locus has a common ancestral state in BBC breeds and a polygenic nature, which is responsible for the variation in dermal pigmentation.

The hyperpigmentation phenotype in the BBC breeds results from common complex chromosomal rearrangement at the *Fm* locus and distinguishes them from non-BBC breeds (Dharmayanthi et al., 2017; Shinde et al., 2023). The shared rearrangement in diverse BBC breeds supports a model of a common origin of BBC followed by dispersal globally. Therefore, we anticipated that the *Id* locus should possess unique characteristics specific to and shared among BBC breeds. Our focus on R1 as a potential *Id* locus led us to perform local phylogenetic analysis, principal component analysis (PCA), and admixture analysis. These analyses effectively separated the BBC breeds from the non-BBC breeds, emphasizing the common ancestry of the *Id* locus within the BBC breeds similar to the *Fm* locus. We observed relatedness at the *Id* locus between some BBC and Tibetan non-BBC native breeds, suggesting ongoing gene flow between these groups reported in earlier studies (Li et al., 2020). Hence, our analysis indicates that the *Id* locus emerged in BBC before their global dispersal and may share selection signatures and causative genetic variants involved in inhibiting dermal pigmentation.

One of our major findings is the signature of selection in the 5Kb genomic region of the last intron of *MTAP,* which contains 6 SNPs with an exceptionally high percentage of breed-specific fixation. *MTAP* plays a crucial role in the melanogenesis pathway within melanocytes, the specialized cells responsible for producing the pigment melanin in the skin (Stephen A. Ainger et al., 2017). It is responsible for catalyzing the phosphorylation of methylthioadenosine (MTA), a byproduct of polyamine metabolism (Behrmann et al., 2003). Methylthioadenosine, in turn,

hinders the function of phosphodiesterase (PDE), a regulator of intracellular cAMP levels. By

blocking PDE, Methylthioadenosine blocks the cAMP-PKA-CREB pathway, leading to the

inhibition of *MITF* stimulation. It has been reported that mutations in the *MTAP* gene are

associated with nevus count and melanoma development (Kvaskoff et al., 2011). Genomic

variants identified in the *MTAP* are associated with plumage and dermal pigmentation in the F2

population crossed from White Leghorn and the Yeonsan Ogye (Cha et al., 2023; Heo et al.,

2023). Functional gene expression analysis of Silkie and White Leghorn chicken embryos during

early embryonic stages indicated distinct patterns of *MTAP* expression. Before embryonic day

10, White Leghorn embryos exhibited a higher *MTAP* expression level than Silkie embryos.

However, a shift occurred after day 10, with Silkie embryos showing greater *MTAP* expression

than White Leghorn embryos (Tian et al., 2014b). In the context of migration and differentiation

studies of melanoblasts in Silkie and White Leghorn chickens, it was observed that Silkie neural

crest cells underwent a slower melanocyte differentiation process compared to their White

Leghorn counterparts. Specifically, on day 8, Silkie neural crest cells displayed less pigmentation

than their White Leghorn counterparts. However, by day 11, they had achieved full

pigmentation, signifying a distinct temporal pattern in melanocyte development between these

two chicken breeds (Faraco et al., 2001; Reedy et al., 1998). The expression of *MTAP* and the

migration and differentiation of melanocytes, both displaying a similar pattern, strongly imply

the pivotal role of *MTAP* in regulating dermal pigmentation. Our findings of putative selective

sweep with SNPs suggest that *MTAP* is the most prominent candidate gene for the dermal

melanin inhibitor.

SNPs in gene promoters can have implications for traits such as skin color and plumage in

various species. For instance, in humans, an SNP in the promoter of the *SLC45A2* gene has been

linked to variations in skin color (Graf et al., 2007). In ducks, the black plumage phenotype is

determined by a mutation within the regulatory region of the *MC1R* gene (Liu et al., 2023).

Similarly, studies on different chicken breeds have revealed that mutations in the promoter

regions of genes like *MITF*, *TYR*, *ASIP,* and *GJA5* play a role in influencing skin color (Li et al.,

2021; Wang et al., 2018; Yu et al., 2019, 2017). Interestingly, past research has established a

connection between loss-of-function mutations in *CCDC112* and variations in feather color in

geese (Ren et al., 2021). A previous gene expression analysis study conducted between Silkie

BBC and White Leghorn chickens reported that after embryonic day 10, the *CCDC112* expression within the shank of Silkie embryos was greater than that of White Leghorn embryos (Tian et al., 2014b). This finding strongly implies a potential role of *CCDC112* in shank pigmentation in chickens. The regulatory mutations identified within *CCDC112's* promoter region hold promise for shedding light on their association with dermal pigmentation and plumage color.

We found that paralogs of two miRNAs, gga-mir-1457 and gga-mir-6552, which have BBC-specific SNPs, target 30 genes associated with the melanogenesis pathway. Previous literature has extensively documented the role of miRNAs in melanocyte biology and pigmentation, primarily by modulating the expression of specific genes or pathways (Mione and Bosserhoff, 2015). Associations between miRNAs and melanin production, skin pigmentation, and plumage coloration have been observed in diverse species such as Koi fishes (Tian et al., 2018), pigs (Yuan et al., 2023), alpacas (Zhu et al., 2019), and ducks (Apopo et al., 2015). Numerous miRNAs have been identified in the case of Muchuan BBC, impacting melanin production in the breast muscle (Yu et al., 2020). Out of the *Id* locus, these miRNAs may involved in plumage color and require a comprehensive functional analysis, which can provide valuable insights into their roles within the melanogenesis pathway.

The crosses between different BBC breeds and the gene expression analysis in early embryonic days can be used to study a polygenic basis for the *Id* locus. Hence, we propose a novel genetic crossing design for conducting a comprehensive study involving various BBC and non-BBC breeds (**Fig. 6**). The proposed experiment involves a round-robin cross between three BBC breeds (Kadaknath, Silkie, and Yeonsan Ogye) and two non-BBC breeds (White Leghorn and White Plymouth Rock) to generate the F1 crosses. The F1 individuals can be categorized based on dermal pigmentation and subsequently utilized for reciprocal crosses to generate F2 crosses. The F2 individuals can be further categorized into three groups based on their pigmentation levels, namely high, intermediate, and no pigmentation. Using GWAS, Population genomic study, and gene expression analysis during the early embryonic stages, particularly concerning genes containing morph-specific SNPs, may provide valuable insights into identifying the responsible genes. The approach we propose here involves multiple intermediate morphs

290 obtained from different genetic backgrounds (i.e., different BBC breeds) and can help identify
291 shared genetic variants involved in dermal pigmentation.

Conclusion

We explored the *Id* locus and used read coverage, blastn search for amplicon repeats and
localised population genetic structure to refine its position within the Gallus_gallus.GRCg6a
genome assembly. Our investigation concludes that the *Id* locus (Z:78725000-80340000) closely
co-localizes with the Z-amplicon repeat units situated at the distal end of the q-arm of
chromosome Z. Multiple lines of evidence, such as local phylogeny, PCA, and admixture
analysis, support this region's notion of common ancestry within BBC breeds like as *Fm* locus.
Among the ten protein-coding genes in this region, *MTAP* stands out as the most prominent
candidate for the *Id*, suggesting a pivotal role in inhibiting dermal pigmentation. In addition,
CCDC112 emerges as a potential candidate for influencing variations in pigmentation patterns.
Further scrutiny of BBC breeds-specific selective sweep region upstream to the Z amplicon led
us to discover miRNAs which can target various genes involved in the melanogenesis pathway.
Our findings collectively suggest that the *Id* locus is shared among BBC breeds, likely exhibiting
a polygenic nature. Future studies considering the proposed novel genetic crossing design may
reveal the mechanism of dermal inhibition at the molecular level.

Materials and methods

Data collection

To comprehensively investigate the genomic structure and diversity of the *Id* locus, we acquired
publicly accessible whole-genome resequencing short-read Illumina data of 270 chicken
individuals. This dataset comprises three distinct wild junglefowl species: green, Ceylon, and
grey. In addition to the wild junglefowl, the dataset includes multiple breeds of domesticated
chicken. We included various BBC breeds, including Kadaknath, Silkie, Yeonsan Ogye, and
Chinese breeds. Additionally, we incorporated non-black native chicken breeds like Aseel,
Japanese gamefowl, Chinese gamefowl, Tibetan native chicken, Ethiopian native chicken, Sri
Lankan native chicken, and Korean native chicken. The collection also encompasses commercial
chicken breeds like White leghorn, Broiler, and Rhode Island Red. Of the 270 individuals used in
this study, 42 belonged to the BBC category, while 228 were categorized as non-BBC. We

sourced these data from public repositories such as the National Center for Biotechnology

Information (NCBI), the European Nucleotide Archive (ENA), the National Genomics Data

Center (NGDC), and the Korean National Agricultural Biotechnology Information Center

(KNABIC). The genome coverage of the resequencing data for each individual was at least more

than 10x. Comprehensive details for each individual can be found in **Table S6**.

Read mapping and variant calling

We started our analysis by mapping paired-end raw Illumina reads from 270 individuals to the

Gallus_gallus.GRCg6a genome assembly. The mapping was executed using the "bwa mem" flag,

employing the default parameters within the Burrows-Wheeler Aligner (BWA) software package

version 0.7.17-r1188 (Li and Durbin, 2009). We incorporated read group information to enhance

data integrity and eliminated duplicate reads from the resulting BAM files, utilizing the Picard

tools (accessible at https://github.com/broadinstitute/picard). For further analysis and variant

calling, we subsetted the bam files for chromosome Z as our region of interest *Id* locus is known

to be located on Z (Dorshorst et al., 2010). We employed two distinct tools for variant calling:

bcftools version 1.10.2 (Danecek et al., 2021; Li and Durbin, 2009) and FreeBayes version 1.0.2

(Garrison and Marth, 2012). For bcftools, we used different quality control flags such as -

mapping quality -C 50, -min base quality -Q 20, and -min mapping quality -q 20. Similarly, for

the FreeBayes, we imposed various quality control flags, including -min-alternate-count -C 10, -

min-mapping-quality -m 20, -min-base-quality -q 20, and -min-coverage, requiring a minimum

of 10 reads at a given locus. The aim was to identify SNPs with high quality and robustness. To

refine the dataset, we applied stringent filters using VCFtools version 0.1.16 (Danecek et al.,

2011). To exclude low-quality variant calls and indels, we used --remove-indels, --max-alleles 2,

and --max-missing 0.95. Additionally, to enhance our SNP dataset's precision, we extracted

common SNPs concordantly identified by both variant callers, minimizing the possibility of false

positive SNP calls.

Validation of the Z amplicon repeat units at the distal end of the q-arm of chromosome Z

In light of previous research (Bellott et al., 2010) about Z amplicon repeat units, we employed

two distinct methodologies: (1) blastn and (2) read coverage estimation, to validate the presence

of the Z amplicon repeat units (ZARU) at the distal end of q-arm of chromosome Z in the

Gallus_gallus.GRCg6a genome assembly. In the first approach, we utilized the blastn v2.9.0

(Altschul et al., 1990) to elucidate the spatial distribution of four specific genes: *ADCY10*,

C2orf3, *ARHGAP33*, and *MRPL19*. To achieve this, we used the genomic sequences of these

genes as query sequences and performed BLAST searches against the Gallus_gallus.GRCg6a

genome assembly. Subsequently, the obtained BLAST hits were saved in bed format. To

visualize the distribution pattern of these genes, we divided chromosome Z into 1 Kb non-

overlapping windows using bedtools v2.27.1 (Quinlan and Hall, 2010). For each of these

windows, we calculated the coverage of the BLAST hits using the bedtools coverage command.

In our second approach, we took a different perspective by generating 10 Kb non-overlapping

windows spanning chromosome Z. This allowed us to analyze read counts within each window,

providing valuable confirmation regarding the presence of multiple copies of the four genes. This

combined methodology aimed to deliver robust evidence and validate the Z amplicon repeat

unit's existence.

Phylogenetic, Principal component, and Admixture analysis

After validation of ZARU at the distal end of the q-arm of chromosome Z in

Gallus_gallus.GRCg6a genome assembly, we found two single-copy protein-coding regions

between them. We considered these single-copy protein-coding regions R1 (Z:78725000-

80340000) and R2 (Z:81040000-82226500) as potential *Id* locus regions. To construct a

localized phylogeny of both regions, we used VCF2PopTree (Subramanian et al., 2019) and vk

phylo implemented in the VCF-kit tool (Cook and Andersen, 2017). We applied Neighbour

Joining (NJ) and the unweighted pair group method with arithmetic mean (UPGMA) to the SNPs

identified within the R1 and R2 regions. Subsequently, for visualizing the resulting phylogenetic

tree, we utilized Figtree v1.4.2 (Rambaut, 2014).

For the Principal Component Analysis (PCA) of both regions, we used PCAngsd v1.10 (Meisner

and Albrechtsen, 2018) based on genotype likelihood estimates derived from the Analysis of

Next-Generation Sequencing Data (ANGSD) v0.935 (Korneliussen et al., 2014). The genotype

likelihood values in beagle format, obtained from ANGSD, were employed in PCAngsd to

determine principal components. Additionally, we estimated admixture proportions for both

regions through NGSadmix (Skotte et al., 2013). In the NGSadmix analysis, we executed

multiple runs with different values of K, ranging from K1 to K15, each with ten iterations. To

discern the optimal K value, we applied the Evanno best K method (Evanno et al., 2005), an

approach implemented in the Clumpak (Kopelman et al., 2015) web server, to identify the most

appropriate K for our analysis. To visualize the results of PCA and Admixture analysis, we used

R (Ihaka and Gentleman, 1996).

Genetic diversity and selection analysis

Our population genetic analysis exclusively included male (ZZ) individuals, a choice driven by

the substantial genomic and sequence divergence observed in sex chromosomes. Using female

(ZW) individuals can lead to uncertainty in the variant calls on the Z chromosome due to the

mapping of gametologous or paralogous sequences shared between W and Z chromosomes

(Ellegren, 2011; Soh et al., 2014). Specifically, we selected three distinct breeds: two BBC

breeds, Kadaknath (KADK) and Silkie (SILK), and a non-BBC breed represented by white

leghorn chickens (WHLH). To ensure robust data quality, we carefully selected ten male

individuals from each breed, each possessing sequencing coverage exceeding 20x (**see Table**

S7). In pursuit of signatures of selection, we computed various population genetic statistics for

the entire chromosome Z, employing 1 Kb windows as our analytical units. To assess genetic

diversity within breeds, we utilized the folded site frequency spectrum (SFS) approach,

implemented through ANGSD, to derive population-specific estimates of π (nucleotide

diversity), Watterson's θ, and τ (Tajima's D). We calculated inter-population genomic

differentiation (F_{ST}) using ANGSD to quantify genetic differentiation between populations. We

calculated the weighted F_{ST} across chromosome Z, dividing the sum of the F_{ST} value of BBC vs.

BBC comparison's F_{ST} value by the non-BBC comparison.

$$\frac{FST_{\text{KADK Vs. SILK}}}{FST_{\text{KADK Vs. WHLH}} + FST_{\text{SILK Vs. WHLH}}}$$

Additionally, we calculated divergence (D_{xy}) between population pairs, employing the

popgenWindows   Python   script,   which   can   be   found   at

https://github.com/simonhmartin/genomics_general. To identify regions potentially under

selection, we defined $F_{ST,}$ weighted $F_{ST,}$ and D_{xy} outlier regions as 1 Kb windows falling within

the top 10 percent of their respective estimates, signaling genomic regions of particular interest

in population differentiation and evolutionary divergence.

Identification of micro-RNA, their target genes, and gene enrichment analysis

Based on our criteria of identifying genomic regions under selection using population genetic analysis, we found a 26 Kb long (Z:72580000-72606000) intergenic region upstream of the ZARU. We searched for micro-RNAs within this region using a nucleotide sequence-based search of micro-RNAs using the miRBase (Kozomara and Griffiths-Jones, 2014) webserver. Our search yielded the identification of five pre-micro-RNA precursors situated within the designated region. Subsequently, we sought to determine the target genes of these micro-RNAs. To accomplish this, we used the TargetScanHuman (Agarwal et al., 2015; McGeary et al., 2019) web server, specifying the chicken as the species of interest, and this enabled us to identify the target genes specific to these chicken-derived micro-RNAs. We performed gene enrichment analysis with the list of identified target genes. To explore the functional implications of these genes, we employed the KEGG pathway database (Kanehisa and Goto, 2000) with default settings. This analysis was facilitated through the ShinyGO v 0.77 (Ge et al., 2020) web server, enabling us to gain insights into the potential pathways and biological processes influenced by these micro-RNAs and their associated target genes.

Identification of BBC-specific SNPs

To understand the impact of genetic variants within both coding and non-coding regions, we used the snpEff v4.3t along with GRCg6a.96 databases for annotation (Cingolani et al., 2012). This approach allowed us to comprehensively assess the genetic variants within regions putatively under selection in BBC and promoter regions of various genes. Our analysis was precisely honed to focus on alternative SNPs found exclusively in BBC individuals while their non-BBC counterparts retained reference alleles. We employed the bam-read count method to calculate the proportions of SNPs within all 270 individuals (42 BBC and 228 non-BBC) (Khanna et al., 2022).

Availability of data

Scripts and data are available at https://github.com/Ashu2195/ID_locus.

CRediT authorship contribution statement

Ashutosh Sharma: Conceptualization, Formal analysis, Investigation, Visualization, Validation, Writing - original draft, Writing - review & editing. **Nagarjun Vijay:** Conceptualization, Resources, Writing - original draft, Writing - review & editing, Funding acquisition, Project administration, Supervision.

Conflict of interest statement

The authors declare that they have no conflict of interest.

Acknowledgment

We thank the University Grants Commission for supporting AS with a Ph.D. scholarship. The Department of Biotechnology, Ministry of Science and Technology, India (Grant no. BT/11/IYBA/2018/03) and Science and Engineering Research Board (Grant no. ECR/2017/001430) provided funds used to generate primary sequencing data published in this article and computational resources (i.e., Har Gobind Khorana Computational Biology cluster) used.

Appendix

Supplementary 1. Supplementary figures

Supplementary 2. Supplementary tables

References.

Agarwal, V., Bell, G.W., Nam, J.W., Bartel, D.P., 2015. Predicting effective microRNA target sites in mammalian mRNAs. Elife 4. https://doi.org/10.7554/ELIFE.05005

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local alignment search tool. J. Mol. Biol. 215, 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2

Apopo, S., Liu, H., Jing, L., Du, X., Xie, S., Gong, Y., Xu, R., Li, S., 2015. Identification and profiling of microRNAs associated with white and black plumage pigmentation in the white and black feather bulbs of ducks by RNA sequencing. Anim. Genet. 46, 627–635.

464  https://doi.org/10.1111/AGE.12343

465 Arora, G., Mishra, S.K., Nautiyal, B., Pratap, S.O., Gupta, A., Beura, C.K., Singh, D.P., 2011.

466  Genetics of hyperpigmentation associated with the Fibromelanosis gene (Fm) and analysis

467  of growth and meat quality traits in crosses of native Indian Kadaknath chickens and non-

468  indigenous breeds. Br. Poult. Sci. 52, 675–685.

469  https://doi.org/10.1080/00071668.2011.635637

470 Barrett, R.D.H., Schluter, D., 2008. Adaptation from standing genetic variation. Trends Ecol.

471  Evol. 23, 38–44. https://doi.org/10.1016/J.TREE.2007.09.008

472 Bateson, B.W., -a, F., Punnett, R.C., Contents, M., 1911. The inheritance of the peculiar

473  pigmentation of the silky fowl. J. Genet. 1911 13 1, 185–203.

474  https://doi.org/10.1007/BF02981551

475 Baynash, A.G., Hosoda, K., Giaid, A., Richardson, J.A., Emoto, N., Hammer, R.E., Yanagisawa,

476  M., 1994. Interaction of endothelin-3 with endothelin-B receptor is essential for

477  development of epidermal melanocytes and enteric neurons. Cell 79, 1277–1285.

478  https://doi.org/10.1016/0092-8674(94)90018-3

479 Behrmann, I., Wallner, S., Komyod, W., Heinrich, P.C., Schuierer, M., Buettner, R., Bosserhoff,

480  A.K., 2003. Characterization of Methylthioadenosin Phosphorylase (MTAP) Expression in

481  Malignant Melanoma. Am. J. Pathol. 163, 683–690. https://doi.org/10.1016/S0002-

482  9440(10)63695-4

483 Bellott, D.W., Skaletsky, H., Pyntikova, T., Mardis, E.R., Graves, T., Kremitzki, C., Brown,

484  L.G., Rozen, S., Warren, W.C., Wilson, R.K., Page, D.C., 2010. Convergent evolution of

485  chicken Z and human X chromosomes by expansion and gene acquisition. Nature 466, 612–

486  6. https://doi.org/10.1038/nature09172

487 Bitgood, J.J., 1988. Linear relationship of the loci for barring, dermal melanin inhibitor, and

488  recessive white skin on the chicken Z chromosome. Poult. Sci. 67, 530–533.

489  https://doi.org/10.3382/ps.0670530

490 Bitgood, J.J., 1985. Additional linkage relationships within the Z chromosome of the chicken.

491   Poult. Sci. 64, 2234–2238. https://doi.org/10.3382/PS.0642234

492 Cha, J., Jin, D., Kim, J.H., Kim, S.C., Lim, J.A., Chai, H.H., Jung, S. a., Lee, J.H., Lee, S.H.,
493   2023. Genome-wide association study revealed the genomic regions associated with skin
494   pigmentation in an Ogye x White Leghorn F2 chicken population. Poult. Sci. 102, 102720.
495   https://doi.org/10.1016/J.PSJ.2023.102720

496 Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X.,
497   Ruden, D.M., 2012. A program for annotating and predicting the effects of single nucleotide
498   polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118;
499   iso-2; iso-3. Fly (Austin). 6, 80. https://doi.org/10.4161/FLY.19695

500 Cook, D.E., Andersen, E.C., 2017. VCF-kit: assorted utilities for the variant call format.
501   Bioinformatics 33, 1581–1582. https://doi.org/10.1093/BIOINFORMATICS/BTX011

502 Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E.,
503   Lunter, G., Marth, G.T., Sherry, S.T., McVean, G., Durbin, R., 2011. The variant call
504   format and VCFtools. Bioinformatics 27, 2156–2158.
505   https://doi.org/10.1093/BIOINFORMATICS/BTR330

506 Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A.,
507   Keane, T., McCarthy, S.A., Davies, R.M., 2021. Twelve years of SAMtools and BCFtools.
508   Gigascience 10, 1–4. https://doi.org/10.1093/GIGASCIENCE/GIAB008

509 Dharmayanthi, A.B., Terai, Y., Sulandari, S., Zein, M.S.A., Akiyama, T., Satta, Y., 2017. The
510   origin and evolution of fibromelanosis in domesticated chickens: Genomic comparison of
511   Indonesian Cemani and Chinese Silkie breeds. PLoS One 12, e0173147.
512   https://doi.org/10.1371/JOURNAL.PONE.0173147

513 Dorshorst, B., Molin, A.M., Rubin, C.J., Johansson, A.M., Strömstedt, L., Pham, M.H., Chen,
514   C.F., Hallböök, F., Ashwell, C., Andersson, L., 2011. A complex genomic rearrangement
515   involving the Endothelin 3 locus causes dermal hyperpigmentation in the chicken. PLoS
516   Genet. 7. https://doi.org/10.1371/journal.pgen.1002412

517 Dorshorst, B., Okimoto, R., Ashwell, C., 2010. Genomic regions associated with dermal

hyperpigmentation, polydactyly and other morphological traits in the silkie chicken. J.

Hered. 101, 339–350. https://doi.org/10.1093/jhered/esp120

Dorshorst, B.J., Ashwell, C.M., 2009. Genetic mapping of the sex-linked barring gene in the

chicken. Poult. Sci. 88, 1811–1817. https://doi.org/10.3382/PS.2009-00134

Dunn, L.C., Jull, M.A., 1927. On the inheritance of some characters op the silky fowl. J. Genet.

19, 27–63. https://doi.org/10.1007/BF02983116/METRICS

Ellegren, H., 2011. Sex-chromosome evolution: recent progress and the influence of male and

female heterogamety. Nat. Rev. Genet. 2011 123 12, 157–166.

https://doi.org/10.1038/nrg2948

Evanno, G., Regnaut, S., Goudet, J., 2005. Detecting the number of clusters of individuals using

the software STRUCTURE: A simulation study. Mol. Ecol. 14, 2611–2620.

https://doi.org/10.1111/j.1365-294X.2005.02553.x

Faraco, C.D., Soˆ, S., Vaz, S.A.S., Vé, M., Pa´stor, R.D., Erickson, C.A., 2001.

Hyperpigmentation in the Silkie Fowl Correlates With Abnormal Migration of Fate-

Restricted Melanoblasts and Loss of Environmental Barrier Molecules. Dev. Dyn. 220:212–

225. https://doi.org/10.1002/1097-0177

Garrison, E., Marth, G., 2012. Haplotype-based variant detection from short-read sequencing.

Ge, S.X., Jung, D., Jung, D., Yao, R., 2020. ShinyGO: a graphical gene-set enrichment tool for

animals and plants. Bioinformatics 36, 2628–2629.

https://doi.org/10.1093/BIOINFORMATICS/BTZ931

Graf, J., Voisey, J., Hughes, I., Van Daal, A., 2007. Promoter polymorphisms in the MATP

(SLC45A2) gene are associated with normal human skin color variation. Hum. Mutat. 28,

710–717. https://doi.org/10.1002/HUMU.20504

Heo, S., Cho, S., Dinh, P.T.N., Park, J., Jin, D.H., Cha, J., Kim, Y.K., Koh, Y.J., Lee, S.H., Lee,

J.H., 2023. A genome-wide association study for eumelanin pigmentation in chicken

plumage using a computer vision approach. Anim. Genet. 54, 355–362.

https://doi.org/10.1111/AGE.13303

Ihaka, R., Gentleman, R., 1996. R: A Language for Data Analysis and Graphics. J. Comput.
Graph. Stat. 5, 299–314. https://doi.org/10.1080/10618600.1996.10474713

Kanehisa, M., Goto, S., 2000. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic
Acids Res. 28, 27. https://doi.org/10.1093/NAR/28.1.27

Khanna, A., Larson, D.E., Srivatsan, S.N., Mosior, M., Abbott, T.E., Kiwala, S., Ley, T.J.,
Duncavage, E.J., Walter, M.J., Walker, J.R., Griffith, O.L., Griffith, M., Miller, C.A., 2022.
Bam-readcount - rapid generation of basepair-resolution sequence metrics. ArXiv 7, 3722.
https://doi.org/10.21105/joss.03722

Kopelman, N.M., Mayzel, J., Jakobsson, M., Rosenberg, N.A., Mayrose, I., 2015. Clumpak: A
program for identifying clustering modes and packaging population structure inferences
across K. Mol. Ecol. Resour. 15, 1179–1191. https://doi.org/10.1111/1755-0998.12387

Korneliussen, T.S., Albrechtsen, A., Nielsen, R., 2014. ANGSD: Analysis of Next Generation
Sequencing Data. BMC Bioinformatics 15, 1–13.

Kozomara, A., Griffiths-Jones, S., 2014. miRBase: annotating high confidence microRNAs
using deep sequencing data. Nucleic Acids Res. 42. https://doi.org/10.1093/NAR/GKT1181

Kvaskoff, M., Whiteman, D.C., Zhao, Z.Z., Montgomery, G.W., Martin, N.G., Hayward, N.K.,
Duffy, D.L., 2011. Polymorphisms in Nevus-Associated Genes MTAP, PLA2G6, and IRF4
and the Risk of Invasive Cutaneous Melanoma. Twin Res. Hum. Genet. 14, 422–432.
https://doi.org/10.1375/TWIN.14.5.422

Li, D., Sun, G., Zhang, M., Cao, Y., Zhang, C., Fu, Y., Li, F., Li, G., Jiang, R., Han, R., Li, Z.,
Wang, Y., Tian, Y., Liu, X., Li, W., Kang, X., 2020. Breeding history and candidate genes
responsible for black skin of Xichuan black-bone chicken. BMC Genomics 21, 1–15.
https://doi.org/10.1186/S12864-020-06900-8/FIGURES/8

Li, G., Li, D., Yang, N., Qu, L., Hou, Z., Zheng, J., Xu, G., Chen, S., 2014. A genome-wide
association study identifies novel single nucleotide polymorphisms associated with dermal
shank pigmentation in chickens. Poult. Sci. 93, 2983–2987.
https://doi.org/10.3382/PS.2014-04164

Li, H., Durbin, R., 2009. Fast and accurate short read alignment with Burrows–Wheeler transform. Bioinformatics 25, 1754–1760. https://doi.org/10.1093/BIOINFORMATICS/BTP324

Li, J., Lee, M.O., Chen, J., Davis, B.W., Dorshorst, B.J., Siegel, P.B., Inaba, M., Jiang, T.X., Chuong, C.M., Andersson, L., 2021. Cis-acting mutation affecting GJA5 transcription is underlying the Melanotic within-feather pigmentation pattern in chickens. Proc. Natl. Acad. Sci. U. S. A. 118, e2109363118. https://doi.org/10.1073/PNAS.2109363118/SUPPL_FILE/PNAS.2109363118.SAPP.PDF

Liu, H., Xi, Y., Tang, Q., Qi, J., Zhou, Z., Guo, Z., Fan, W., Hu, J., Xu, Y., Liang, S., Xie, M., Tang, J., Zhang, Y., Hou, S., 2023. Genetic fine-mapping reveals single nucleotide polymorphism mutations in the MC1R regulatory region associated with duck melanism. Mol. Ecol. 32, 3076–3088. https://doi.org/10.1111/MEC.16924

McGeary, S.E., Lin, K.S., Shi, C.Y., Pham, T.M., Bisaria, N., Kelley, G.M., Bartel, D.P., 2019. The biochemical basis of microRNA targeting efficacy. Science (80-.). 366. https://doi.org/10.1126/SCIENCE.AAV1741

Mcgibbon, W.H., 1978. The genetics of green spotting on shanks of Ancona chickens. J. Hered. 69, 97–100. https://doi.org/10.1093/OXFORDJOURNALS.JHERED.A108912

Meisner, J., Albrechtsen, A., 2018. Inferring population structure and admixture proportions in low-depth NGS data. Genetics 210, 719–731. https://doi.org/10.1534/genetics.118.301336

Mione, M., Bosserhoff, A., 2015. MicroRNAs in melanocyte and melanoma biology. Pigment Cell Melanoma Res. 28, 340–354. https://doi.org/10.1111/PCMR.12346

Mukherjee, T.K., Jerome, F.N., Reinhart, B.S., 1969. The Inhibition of Dermal Melanin in the Shanks of the Domestic Fowl. Poult. Sci. 48, 2137–2142. https://doi.org/10.3382/PS.0482137

Quinlan, A.R., Hall, I.M., 2010. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics 26, 841–842. https://doi.org/10.1093/BIOINFORMATICS/BTQ033

599   Rambaut, A., 2014. FigTree v1.4.2., A Graphical Viewer of Phylogenetic Trees. [WWW

600   Document]. http://tree.bio.ed.ac.uk/software/figtree/. URL

601   http://tree.bio.ed.ac.uk/software/figtree/

602   Reedy, M. V., Faraco, C.D., Erickson, C.A., 1998. Specification and migration of melanoblasts

603   at the vagal level and in hyperpigmented silkie chickens. Dev. Dyn. 213, 476–485.

604   https://doi.org/10.1002/(SICI)1097-0177(199812)213:4<476::AID-AJA12>3.0.CO;2-R

605   Ren, S., Lyu, G., Irwin, D.M., Liu, X., Feng, C., Luo, R., Zhang, J., Sun, Y., Shang, S., Zhang,

606   S., Wang, Z., 2021. Pooled Sequencing Analysis of Geese (Anser cygnoides) Reveals

607   Genomic Variations Associated With Feather Color. Front. Genet. 12, 650013.

608   https://doi.org/10.3389/FGENE.2021.650013/BIBTEX

609   Shinde, S.S., Sharma, A., Vijay, N., 2023. Decoding the fibromelanosis locus complex

610   chromosomal rearrangement of black-bone chicken: genetic differentiation, selective

611   sweeps and protein-coding changes in Kadaknath chicken. Front. Genet. 14, 1180658.

612   https://doi.org/10.3389/FGENE.2023.1180658/BIBTEX

613   Shinomiya, A., Kayashima, Y., Kinoshita, K., Mizutani, M., Namikawa, T., Matsuda, Y.,

614   Akiyama, T., 2012. Gene Duplication of endothelin 3 Is Closely Correlated with the

615   Hyperpigmentation of the Internal Organs (Fibromelanosis) in Silky Chickens. Genetics

616   190, 627–638. https://doi.org/10.1534/GENETICS.111.136705

617   Siwek, M., Wragg, D., Sławińska, A., Malek, M., Hanotte, O., Mwacharo, J.M., 2013. Insights

618   into the genetic history of Green-legged Partridgelike fowl: mtDNA and genome-wide SNP

619   analysis. Anim. Genet. 44, 522–532. https://doi.org/10.1111/AGE.12046

620   Skotte, L., Korneliussen, T.S., Albrechtsen, A., 2013. Estimating individual admixture

621   proportions from next generation sequencing data. Genetics 195, 693–702.

622   https://doi.org/10.1534/genetics.113.154138

623   Soh, Y.Q.S., Alföldi, J., Pyntikova, T., Brown, L.G., Graves, T., Minx, P.J., Fulton, R.S.,

624   Kremitzki, C., Koutseva, N., Mueller, J.L., Rozen, S., Hughes, J.F., Owens, E., Womack,

625   J.E., Murphy, W.J., Cao, Q., De Jong, P., Warren, W.C., Wilson, R.K., Skaletsky, H., Page,

626   D.C., 2014. Sequencing the mouse Y chromosome reveals convergent gene acquisition and

amplification on both sex chromosomes. Cell 159, 800–813.

https://doi.org/10.1016/J.CELL.2014.09.052

Sohn, J. Il, Nam, K., Hong, H., Kim, J.M., Lim, D., Lee, K.T., Do, Y.J., Cho, C.Y., Kim, N.,

Chai, H.H., Nam, J.W., 2018. Whole genome and transcriptome maps of the entirely black

native Korean chicken breed Yeonsan Ogye. Gigascience 7, 1–14.

https://doi.org/10.1093/GIGASCIENCE/GIY086

Stephen A. Ainger, Kasturee Jagirdar, Katie J. Lee, H. Peter Soyer, Richard A. Sturm, 2017.

Skin Pigmentation Genetics for the Clinic. Dermatology.

Subramanian, S., Ramasamy, U., Chen, D., 2019. VCF2PopTree: a client-side software to

construct population phylogeny from genome-wide SNPs. PeerJ 7.

https://doi.org/10.7717/PEERJ.8213

Tian, M., Fang, S., Wang, Y., Gu, X., Feng, C., Hao, R., Hu, X., Li, N., 2014a. Inverted

duplication including Endothelin 3 closely related to dermal hyperpigmentation in Silkie

chickens. Front. Agric. Sci. Eng. 1, 121–129. https://doi.org/10.15302/J-FASE-2014026

Tian, M., Hao, R., Fang, S., Wang, Y., Gu, X., Feng, C., Hu, X., Li, N., 2014b. Genomic regions

associated with the sex-linked inhibitor of dermal melanin in Silkie chicken. Front. Agric.

Sci. Eng. 1, 242–249. https://doi.org/10.15302/J-FASE-2014018

Tian, X., Pang, X., Wang, Liangyan, Li, M., Dong, C., Ma, X., Wang, Lei, Song, D., Feng, J.,

Xu, P., Li, X., 2018. Dynamic regulation of mRNA and miRNA associated with the

developmental stages of skin pigmentation in Japanese ornamental carp. Gene 666, 32–43.

https://doi.org/10.1016/J.GENE.2018.04.054

Tixier-Boichard, M., 2002. From phenotype to genotype: major genes in chickens. Worlds.

Poult. Sci. J. 58, 65–75. https://doi.org/10.1079/WPS20020008

Wang, G., Liao, J., Tang, M., Yu, S., 2018. Genetic variation in the MITF promoter affects skin

colour and transcriptional activity in black-boned chickens. Br. Poult. Sci. 59, 21–27.

https://doi.org/10.1080/00071668.2017.1379053

Xu, J., Lin, S., Gao, X., Nie, Q., Luo, Q., Zhang, X., 2017. Mapping of Id locus for dermal shank

melanin in a Chinese indigenous chicken breed. J. Genet. 96, 977–983.

https://doi.org/10.1007/S12041-017-0862-Z

Yan, L., Liqing, Z., Dexiang, Z., Xiquan, Z., Xuemei, L., 2010. Faster evolution of Z-linked

duplicate genes in chicken. J. Genet. Genomics 37, 695–702. https://doi.org/10.1016/S1673-

8527(09)60087-4

Yu, S., Liao, J., Tang, M., Wang, Y., Wei, X., Mao, L., Zeng, C., Wang, G., 2017. A functional

single nucleotide polymorphism in the tyrosinase gene promoter affects skin color and

transcription activity in the black-boned chicken. Poult. Sci. 96, 4061–4067.

https://doi.org/10.3382/PS/PEX217

Yu, S., Wang, G., Liao, J., 2019. Association of a novel SNP in the ASIP gene with skin color in

black-bone chicken. Anim. Genet. 50, 283–286. https://doi.org/10.1111/AGE.12768

Yu, S., Wang, G., Liao, J., Tang, M., Chen, J., 2020. Identification of key microRNAs affecting

melanogenesis of breast muscle in Muchuan black-boned chickens by RNA sequencing. Br.

Poult. Sci. 61, 225–231. https://doi.org/10.1080/00071668.2019.1709619

Yuan, W., Qin, H., Bi, H., Zhao, D., Zhang, Y., Chen, W., 2023. Ssc-mir-221-3p regulates

melanin production in Xiang pigs melanocytes by targeting the TYRP1 gene. BMC

Genomics 24, 1–14. https://doi.org/10.1186/S12864-023-09451-W/FIGURES/6

Zhu, Z., Ma, Y., Li, Y., Cheng, Z., Li, H., Zhang, L., Xu, D., Li, P., 2019. Comparison of

miRNA-101a-3p and miRNA-144a-3p regulation with the key genes of alpaca melanocyte

pigmentation. BMC Mol. Biol. 20, 1–11. https://doi.org/10.1186/S12867-019-0137-

8/FIGURES/8

Figure legends:

Fig. 1: A magnified view of a ~10.5 Mb (72000000-82529921) segment situated on the q-arm

of Chicken chromosome Z: (A) Within this segment, two sets of Z amplicon repeat units

(ZARU) which primarily harbor multiple copies of three genes: *C2orf3*, *ARHGAP33*, and

MRPL19. In the initial part of the first Z amplicon repeat unit (ZARU), the *ADCY10* gene has

multiple copies. In contrast, towards the terminal end of chromosome Z, various copies of

ARHGAP33 are present, depicted in the initial panel. The second panel shows read coverage

within a 10 Kb window of the same region and supports high coverage due to repeat regions. (B)

Genes present in the R1 region have been shown. The direction of the arrows represents the

direction of transcription.

Fig. 2: Phylogenetic tree and population structure analysis of the R1 region: (A)

Phylogenetic tree based on SNPs (Single Nucleotide Polymorphisms) at the R1 region,

showcasing 270 individuals (42 BBC and 228 non-BBC) and visualized using FigTree. **(B)**

Principal Component Analysis (PCA) of the R1 region indicates that PC1 explains 19.8% of the

genetic variance, while PC2 accounts for 9.01%. **(C)** An admixture analysis for individual

ancestry reveals distinct separation of all BBC individuals at K6.

Fig. 3: Signature of selection in the last intron of the *MTAP* gene located in R1: (A)

Comparative analysis of Weighted F_{ST} (KADK vs. WHLH + SILK vs. WHLH/KADK vs. SILK)

between BBC (KADK and SILK) and non-BBC (WHLH). **(B)** F_{ST} comparison specifically

between KADK and SILK. **(C)** F_{ST} comparison between KADK and WHLH. **(D)** FST

comparison between SILK and WHLH. **(E)** Comparison of D_{xy} between KADK and SILK. **(F)**

Comparison of D_{xy} between KADK and WHLH. **(G)** Further F_{ST} comparison between SILK and

WHLH along chromosome Z, using 1Kb non-overlapping windows. A horizontal black dotted

line represents the 90th percentile threshold for both F_{ST} and D_{xy} comparisons. **(H)** The

phastCons conservation score of the GRCg6a genome assembly is displayed in dark blue, with a

red dotted vertical line marking the selected region. **(I)** Identification of 6 SNPs located within a

5kb span in the last intron of the *MTAP* gene. The percentage of bases at each SNP has been

documented for both the BBC (n=42) and non-BBC (n=228) breeds.

Fig. 4: Identification of three SNPs in the promoter region of the *CCDC112* gene: (A) Within

the promoter region of the *CCDC112* gene, three distinctive SNPs, uniquely found in BBC, are

illustrated by orange triangles. **(B)** This section presents the percentage of bases at these three

SNPs for the BBC (n=42) and the non-BBC (n=228) breeds.

Fig. 5: Signature of selection in a 26 Kb long intergenic region located upstream to the

ZARU: (A) Comparative analysis of Weighted F_{ST} (KADK vs. WHLH + SILK vs.

WHLH/KADK vs. SILK) between BBC (KADK and SILK) and non-BBC (WHLH). **(B)** F_{ST} comparison between KADK and SILK. **(C)** F_{ST} comparison between KADK and WHLH. **(D)** F_{ST} comparison between SILK and WHLH. **(E)** Comparison of D_{xy} between KADK and SILK. **(F)** Comparison of D_{xy} between KADK and WHLH. **(G)** Further F_{ST} comparison between SILK and WHLH, along chromosome Z, using 1Kb non-overlapping windows. A horizontal black dotted line represents the 90th percentile threshold for both F_{ST} and D_{xy} comparisons. **(H)** and **(I)** depict π and Watterson theta, respectively. Here, slate blue represents the KADK breed, firebrick represents SILK, and cyan represents WHLH. **(J)** The phastCons conservation score of the GRCg6a genome assembly is displayed in dark blue, with a red dotted vertical line marking the selected region. **(K)** and **(L)** presents the gene enrichment analysis for the target genes of gga-mir-1457 and gga-mir-6552.

Fig. 6: Proposed novel genetic crossing design for identification of *Id* gene: A round-robin cross between a group of five breeds consisting of three BBC breeds (Kadaknath, Silkie, and Yeonsan Ogye) and two non-BBC breeds (White Leghorn and White Plymouth Rock).