The evolution of fitness during range expansions in different dimensions

Kotsar Yurii[1,2] | Hikaru Matsuoka[1] | Gen Tamiya[1,2]

[1] RIKEN Center for Advanced Intelligence Project, Tokyo, Japan

[2] Tohoku University Graduate School of Medicine, Sendai, Japan

Correspondence: Gen Tamiya, RIKEN Center for Advanced Intelligence Project, Tokyo, 103-0027, Japan

Email: gtamiya@med.tohoku.ac.jp

ABSTRACT

We develop a set of programs – the range expansions simulation kit (RESK) – to efficiently
simulate range expansions of populations on a square lattice in 1D, 2D (on a strip, and on a disk)
and 3D (in a cylinder, and in a sphere). In this study, we present the simulation kit to the public
and present some results of using it to simulate a population of diploid individuals with finite
genome regions, each containing infinite sites. Specifically, using the programs, we calculate and
analyse the temporal evolution of population fitness, as well as fitness on the expansion front, in
3D for the first time. We explore the model over different conditions, compare normalisation
methods for fitness, and explore the case of radial (sphere) and axial (cylinder) expansions in 3D,
which might apply in the analysis of the different real-life populations, such as viruses/bacteria
inside a host, marine species in ocean environments, and potentially in future space colonization
planning. In 3D expansions, we find complex spatial fluctuations in deme-average fitness values,
different from those in radial 2D expansions. In axial 3D (cylinder) expansions, we determine that
the highest-valued deme-average fitness lies along the axis of the expansion. We also find the

fluctuation patterns of fitness in 3D cylinder expansions, similar to those previously seen in radial expansions in 2D. In radial 2D (disk) expansions, we find that the fitness of a population undergoing multiple mutations shows a smooth combination of binary segregation pictures against each of those mutations. We confirm the accumulation of deleterious mutations -- a phenomenon known as expansion load -- in all scenarios above. We present the software used in the above to the public as a Julia repository, ready to use with different functions for simulating range expansions with minimal syntax.

1. INTRODUCTION

Most species on Earth exhibit fluctuating habitats, a significant dynamic mode of which is range expansions. Understanding the consequences and dynamics of such changes of habitats is of great importance in evolutionary biology. Their distribution, however, is nontrivial even in a spatially deterministic model of migration, since almost all organisms have multiple segregating genetic loci which accept random mutations (which, in turn, affect the fitness, and thus, the number of individuals). Due to demes on the expansion front having less genetic variation in the early generations after colonisation, while the front is moving, a mutation (neutral, deleterious or beneficial) can fix itself on the front for many generations thereafter, but also spread back into the bulk of the population. This is the principle of *gene surfing* (Klopfstein *et al.*, 2005). Through many generations, the effect of those mutations will accumulate, if the mutations are prevalently either beneficial or deleterious (Peischl *et al.*, 2013). There is evidence to show that 70% to 90% of all mutations are deleterious (Eyre-Walker and Keightley, 2007).

This lowers the value of fitness averaged over the individuals in every deme on the expansion front – simply "*mean front fitness*" hereafter – and exacerbates the difference in fitness between the core population and the expansion front. Nevertheless, if the selection is soft, mean front fitness can never fall to zero, and a population can avoid extinction (Peischl *et al.*, 2015a). In this regime, establishing the means to quantify the change in mean front fitness over the expansion front is an important goal, the results of which have various applications in the field of population ecology, especially in the wake of some range expansions caused by climate change (Bokma, 2010; Parmesan, 2006; Pateman *et al.*, 2012; Yamano *et al.*, 2011). These cases assume a one-dimensional (1D) or two-dimensional (2D) habitat, but this change in fitness could also be modelled for three dimensions (3D), which is the focus of this study. In the 3D case, quantifying the change in mean front fitness is important in elucidating the capabilities of viruses (Bocharov *et al.*, 2016), as well as marine species to spread (Kitchel *et al.*, 2022; Assis *et al.*, 2016), or could be vital information in the planning stages of human space expansion. A particularly important

case is the gradual migration of marine species in depth, in addition to lateral migration –
primarily to deeper latitudes, as a result of climate change (Poloczanska *et al.*, 2016). In these
scenarios, a reliable way to simulate and understand range expansions in three dimensions is
required, especially for the case of an expansion along an axis and a certain radius around it to
allow for slight deviations -- in our terminology, the *3D cylinder* scenario.

So far, an analytical approximation to the change in mean front fitness has been
constructed only for 1D (Peischl *et al.*, 2013). This prior research shows that it's difficult to
construct a theoretical formula for mean front fitness for the case of range expansions with logistic
growth, mutation, selection and genetic drift, and so it is valuable to create a set of simulation
programs with the purpose of generating the dynamics of a range expansion, generation by
generation, and to use the simulated data to quantify the change in mean front fitness. We thus set
the goal of developing a time-effective and robust in terms of starting conditions simulation kit,
that can also work with 2D and 3D expansions. Furthermore, with this study, we set out to provide
an example of our simulation kit with some realistic parameters for human range expansions.

We base this study in part on the research in Peischl *et al.* (2013). Their study includes
a C++ program, which is able to generate fitness values for each deme in a one-dimensional
expansion. In their publication, some simulation results for 1D and 2D have been given. In our
study, based on the aforementioned C++ program, as well as Gilbert *et al.* (2018), we recreate the
functionality for 1D expansions, and independently develop 2D and 3D functionality, in the Julia
programming language. In this study, we consolidate the above into a ready-to-use simulation kit
and present it as well as examples of simulations in different dimensions, while exploring different
steps throughout the algorithm (such as normalisation methods). It is available in the "Data
availability statement".

2. METHODS

2.1 Model: genome regions with infinite sites

Our model is based largely on the model in Peischl *et al.* (2013). For all dimensions, we
start with a habitat of discrete laterally connected lattice of demes, and employ a model which
accepts a population of monoecious diploid individuals of size N, employing the stepping-stone
model (Kimura, 1953), where migration is only present between adjacent demes.

In this model, each diploid individual has a genome of $2L$ independently segregating
regions with no epistasis and no dominance. We set $L = 20$ in this study. These regions are the
basic units during genetic recombination, which occurs once per generation. During genetic

recombination, regions $1 \leq k \leq L$ have a 1/2 chance to swap with their pair in $L+1 \leq k \leq 2L$. Inside

these genome regions, we assume the infinite-site model, so the same segregating region could

have multiple mutations. A mutation has a fixed chance μ of occurring in a gamete, and upon that,

a fixed chance ϕ of being deleterious. We set this value as $\phi = 0.9$, unless stated otherwise.

Deleterious; beneficial mutations change the fitness of a recombining region by $1 - |s|$; $1 + |s|$

respectively. In this study, $|s|$ is a fixed value in each simulation trial, and is called the *selection*

coefficient. Multiplicative fitness is assumed. The fitness values are the main variables in our

model; they are stored for every segregating region for every individual and begin at 1. In order

to model mutations, we sample the number of mutations from a Poisson distribution with mean

μ, determine whether the mutation is beneficial or deleterious, then act on the fitness of the

recombining region picked randomly from a uniform distribution. Since an infinite-site model is

assumed inside the genome regions, the chance of further changing the allele of the same locus is

infinitesimal, thus once the mutation has affected the fitness, the multiplicative factor of its

selective effect always persists in the individual fitness. The evolution of each deme's population

is based on the Beverton-Holt formula for discrete logistic growth:

98
$$N_{t+1} = RN_t \Big/ \left(1 + N_t \frac{R-1}{K} \right)$$

where N_t is the number of individuals in a deme in generation t, $R = 2$ is the proliferation rate, and

K is the constant carrying capacity, set equal to 100, unless specified otherwise.

Furthermore, we sample the actual number of offspring individuals from a Poisson

distribution with mean N_{t+1}. The deme itself does not have a hard limit for the number of

individuals, hence due to Poisson sampling, the number of individuals in a deme can sometimes

exceed K. The fitness of all the segregating regions is multiplied for all individuals within a deme,

and pairs of individuals are randomly selected from the deme. The maximum fitness within a

deme is found, and the individuals from pairs have their fitness divided by the maximum fitness

and compared with a uniform random variable. If both individuals pass, their zygote enters into

the new generation. This is done until the number of offspring sampled above is satisfied. Since

we assume monoecious individuals, selfing is allowed.

Finally, each generation, migration is allowed and equally occurs between all adjacent

demes, with a total migration rate m. In one dimension, migration rate is simply $m/2$ to move to

either of two adjacent demes. In higher dimensions, migration rate is m/l, where l is the number

of laterally adjacent demes. In particular, l is 2 in 1D, 4 in 2D, and 6 in 3D. We then employ an

additional check for the demes at the boundaries of the habitat: we use reflecting borders, in the

sense that if migration has been initiated, the individual shall always move somewhere -- in this

case, away from the boundary of the habitat.

It is difficult, in principle, to calculate a clear proportion of deleterious and beneficial
mutations, since accurate measurement of the effects of single mutations is possible only when
they have fairly large effects on fitness (> 1%) (Eyre-Walker and Keightley, 2007). Nevertheless,
different studies are in agreement that the proportion of deleterious mutations is prevalent and is
between 70% and 90% (Kim *et al.,* 2017). In our model, single mutations can only have constant
deleterious or beneficial effects, which we set for this study as $s = -0.005$ and $s = +0.005$
respectively, and which can be viewed as average values of all possible mutations for
$-1 < s < 0$ and $0 \leqq s < \infty$ respectively, where the weights for $s = -0.005$ and $s = +0.005$
(multiplicative changes in fitness of 1 - 0.005, 1 + 0.005 respectively) need to be decided. We
base the weights on the distribution values in humans (Eyre-Walker and Keightley, 2007): ~90%
for deleterious and ~10% for beneficial mutations, when taken without neutral mutations. The
proportion of deleterious to all mutations can be changed in the program, and our model can be
used for viruses and other populations as well: Sanjuan *et al.* (2010) give ~70% for deleterious
mutations in viruses. There are also lethal mutations, which have to be considered due to their
special logic of potentially erasing an individual from the model. As in the current study we are
focusing on human populations, for which lethal mutations make up less than 0.5% of deleterious
mutations (Gao *et al.,* 2015), we can reasonably ignore them. For viruses, however, more than
25% of mutations could be lethal (Sanjuan *et al.,* 2010). A more accurate approach would be to
consider variable s (e.g. multiple s bands), which we aim to integrate in our simulations in the
future.

2.2   Simulations

We developed multiple methods for our simulations kit to recreate different scenarios of range
expansions, based on the model described above. These methods are named *rangeexp_#d_inf*,
where # is the number of spatial dimensions.

We divide the total number of generations into a burn-in phase and an expansion phase. During
burn-in, the habitat is restricted to an area much smaller than the full space. This is done to allow
the population to reach mutation-selection-drift equilibrium (Peischl *et al.*, 2013). During the
expansion phase, the population is allowed to migrate freely, up until the boundaries of the habitat.
In all simulation trials (1D, 2D and 3D), we picked a size of the habitat that allow the population
to fill the habitat up in several hundreds to thousands of generations. Next, we describe the general
algorithm of the programs (Figure 1).

We start with an array of empty demes, which are themselves arrays of individuals
(Figure S1). We then pick b random demes and fill them to their carrying capacity (given in Table
1) with individuals. Every individual starts out with an array of fitness values of 1 for every

genome region. For all simulations in the current study, we set this parameter b to 5. Every generation, we run a cycle over all demes: we first determine the number of offspring logistically (as described above), then, for every individual in the offspring, we iterate over each individual in the current generation, in a random order, until we find suitable parents for the offspring individual (we call this *draw until pass* below). Specifically, the two candidates for parents must draw samples from a uniform distribution from 0 to the current deme's maximum fitness, that pass the fitness values of their gametes. In one case below (referenced as *first pair*), we also employ a variation wherein, instead of a running a *while* cycle, we run a *for* cycle on the parents: in other words, the fate of offsprings individuals depends on the first random pair of parents to either pass or fail the check; if this check fails, the offspring is automatically less by one individual.

After this step, the two parents' gametes are recombined, and mutations are introduced. The offspring individual is formed, and then allowed to migrate.

The are several parameters that the simulations take as input. We summarise them in Table 1.

The simulations output:

• the **deme-average fitness matrix** per every generation, which stores the deme-average individual fitness for every deme in the habitat,

• the **population matrix** per every generation, which stores deme populations,

• the **deme-average mutations matrices** per every generation, storing the deme-average number of deleterious as well as beneficial mutations.

The focus for this study is to use the output for the deme-average fitness to further calculate its average over the expansion front, and then graph the temporal evolution of the resulting mean front fitness. We do this for different dimensions and configurations, and compare it to the analytical approximation found below in 2.3. More specifically, we find the expansion front as the collection of outermost demes, counting from the borders of the habitat to one side ($x > 0$ half-plane in 2D, and $z > 0$ half-space in 3D) in axial expansions, and all sides in radial expansions. We then find the mean of deme-average fitness values over this expansion front. The population matrix was used to confirm the logistic growth of the population, and to determine the generation at which the whole habitat was filled, i.e. the end of the expansion. The population matrix and mutations matrices are available as output options in our simulation kit.

2.2.1 Normalisation of fitness

It is also common to normalise the mean front fitness. The consideration of a normalisation in time appears when we include a burn-in phase. During this phase, individuals

are locked in a tight environment, and in the absence of expansion mean front fitness grows at a
fast pace. It is then natural to have mean front fitness start at the same value at the onset of the
expansion as the value it would have had if there was no burn-in phase. Since we set our initial
value of fitness in the genome to be 1, we could normalise our mean front fitness such that the
value at the onset generation is 1. To this end, we divide mean front fitness in each generation
after the onset by its value at the onset of the expansion. This approach seems to yield good
correspondence with data (Peischl *et al.*, 2013, 2015b). We call this method "*onset mean*
normalisation" in the results below.

However, this is not the only kind of normalisation applicable. A more literal definition
of normalisation could also be applied, which maps the average values of fitness of each inhabited
deme to the interval [0, 1] at every generation by dividing each deme-average fitness by the
maximum overall inhabited demes. A similar approach is to map the values of fitnesses of each
individual by dividing individual fitnesses by the maximum fitness of all surviving individuals.
We have found that these approaches result in far more rapid decline in normalised mean front
fitness. We included the results of dividing mean front fitness by the maximum deme-average
fitness in the graphs in *Results* as "*maximum normalisation*". Rather thanbeing a method to equate
the scale of fitness values after a certain number of generations (in our case, the burn-in period)
passed with the scale of fitness values obtained without that period, "maximum normalisation" is
a normalisation in space, that gives a good way of quantifying the expansion load. We can also
use this method to confirm that the difference in fitness between the core population and the front
population grows with time in range expansions, even when the "onset mean normalisation"
yields increasing fitness with time.

2.3 Analytical approximation

An analytical approximation to the temporal evolution of mean front fitness in this model was
developed in Peischl *et al.* (2013). In this approximation, migration from a deme is only assumed
possible after the deme has reached the carrying capacity K. This is a good approximation for low
migration rates (m), at least compared to the logarithmic proliferation rate $r = \ln(R) >> m$. Once
at the capacity, $Km/2$ individuals on average migrate to further demes in one general direction
(out of two). This holds especially well in axial expansions. We then only concern ourselves with
those $Km/2$ individuals, since those are assumed to comprise the expansion front. We employ the
Wright-fisher model of evolution of allele frequencies, but split that process into growth and

sampling phases (Wahl and Gerrish, 2001). This allows us to model $T \approx \log(2m^{-1})/r$ intermediate

generations of the growth phase, during which the front population undergoes only reproduction
and selection against the mutation that had just occurred. This is equivalent to a series of
consequent founder effects, and is similar to a model of repeated bottlenecks.

223   2.3.1 Probability of fixation

224   We derive a more general formula for the probability of fixation in this analytical model. In
225   Peischl *et al.* (2013), the mutant frequency x' in the next generation is given by

226
$$x' = \frac{x \exp[(r + s)T]}{x \exp[(r + s)T] + (1 - x) \exp(rT)} = \frac{x \exp(sT)}{x \exp(sT) + 1 - x},$$

227   where x is the mutant frequency in the current generation. We would like to use the difference Δx
228    $= x' - x$, its mean $M[\Delta x]$ and its variance $V[\Delta x]$, to find the probability of fixation of a mutation
229   with effect s on the expansion front. We have

230
$$M_{\Delta x} = x' - x, \quad V_{\Delta x} = \frac{1}{2K\frac{m}{2}} x'(1 - x').$$

231   Meanwhile, the general formula for the gene frequency distribution under irreversible mutation
232   (Kimura, 1964):

$$u(p) = \int_0^p G(x)dx \Big/ \int_0^1 G(x)dx ., \tag{1}$$

235
$$G(x) = \exp\left(-\int \frac{2M_{\Delta x}}{V_{\Delta x}} dx\right).$$

236   In our case:

237
$$-\frac{2M_{\Delta x}}{V_{\Delta x}} = -2Km \cdot \frac{x - x'}{x'(1 - x')} = 2Km(e^{-sT} - 1)[(e^{sT} - 1)x + 1].$$

239   We do not assume small s here, and so this does not simplify further into the formula in Peischl
240   *et al.* (2013). Next, we rearrange and define A and B:

241
$$-\frac{2M_{\Delta x}}{V_{\Delta x}} = 2Km(e^{-sT} - 1)[(e^{sT} - 1)x + 1] =$$

242
$$-\underbrace{\frac{2Km}{e^{sT}}(e^{sT}-1)}_{A}\left[\underbrace{(e^{sT}-1)}_{B}x+1\right].$$

243   Then,

244
$$G(x) = e^{\frac{-AB}{2}x^2 - Ax + C}.$$

245   Then the integrals in Eq. 1 can be seen as Gaussian integrals. Note that $A > 0$, $B > 0$, $AB > 0$, but
246   the coefficient of x^2 in the exponent is a minus, so we are dealing with the error function erf(x).
247   We have:

248
$$\int_0^p e^{-\frac{AB}{2}x^2 - Ax + C}\,dx = \sqrt{\frac{\pi}{2AB}}\, e^{C + \frac{A}{2B}}\big[\text{erf}\big(A^*(Bp+1)\big) - \text{erf}(A^*)\big],$$

249
$$A^* = \sqrt{\frac{A}{2B}} = \sqrt{\frac{Km}{e^{sT}}}.$$

250   Solving the lower integral in a similar manner, Eq. 1 becomes:

$$u(p) = \frac{\text{erf}[A^*(Bp+1)] - \text{erf}(A^*)}{\text{erf}[A^*(B+1)] - \text{erf}(A^*)}. \tag{2}$$

We have tested this function for the fixation probability in a flexible range of parameters
s, m, r, K. Compared to the formula in Peischl et $al.$ (2013), it gives slightly lower values of mean
front fitness over time for small s and other parameters close to values used in this paper, with
relative difference $\delta(\bar{w}) \sim 0.7\%$ after 1000 generations. Unfortunately, this analytical model is
known to underestimate actual mean front fitness, so the difference does not contribute to the
accuracy of approximation of the actual mean front fitness. On the other hand, it differs from the
formula in Peischl et $al.$ (2013) quite drastically when $s \gtrsim 0.03, K \lesssim 10$. Hereafter and in the
results, we employ our derived formula (Eq. 2).

3. RESULTS

3.1 1D

In one dimension, we have carried out 3 trials: one long simulation over 70 replicates, which is
our main trial, designed to measure the long-term behaviour of fitness in range expansions (**1**), a
trial concerning the critical value of the rate of deleterious mutations (**2**), and a trial employing a

different approach to generating the offspring in the model (**3**). All trials are roughly one-
directional, in the sense that individuals start in the leftmost demes of the habitat, and we measure
the front as the (moving) populated outermost demes on the right.

1) We ran 70 simulation replicates, each with a burn-in phase and active phase of 10000 and 2500
generations respectively; as well as 70 replicates without the burn-in phase. Figure 2 shows the
evolution of mean front fitness in time over 70 simulation replicates. Along with it, we have also
run the program from Peischl *et al.* (2013) to simulate 10 and 20 replicates of the expansion with
identical parameters (all of the parameters are given in Table 1). The rise after approximately 1930
generations of free expansion corresponds to the generation in which the population filled up the
whole habitat, thus raising fitness due to efficient selection. For simulations with the burn-in phase
was present, we also display the results of applying different normalisation methods ("onset mean
normalisation" and "maximum normalisation" in Figure 2).

2) We have also performed a trial of the formula for the critical value of the rate of deleterious
mutations, below which an expansion with the parameters used in **1)** (and given in Table 1) would
always have increasing mean fitness at the front (Peischl *et al.*, 2013). We calculated this value to
be equal to $\phi_c \sim 0.57$. Thus we set the simulation parameters to the ones in **1)**, with the exception
of $\phi = 0.45$ and the simulation length: we run a simulation for 5000 burn-in generations and
1250 generations after the onset of the expansion, 10 times. The results confirm that our program
gives an increasing trend of normalised mean front fitness during a range expansion, even with 10
simulation replicates (Figure S2).

3) To test out and compare the "first pair" and the "draw until pass" approaches that we have
described in Chapter 2.1, we ran two sets of simulations under more extreme conditions. The
results are shown in Figure S3. We can see that for mean front fitness with applied "onset mean
normalisation", the trend in both approaches is identical, and the difference in values is within
2%, excluding the rough fluctuations near the onset of the expansion (presumably due to a small
sample of simulations).

3.2 2D strip

294   In two dimensions, we likewise focused on developing simulation programs capable of
295   confirming the results of prior research, concerning the evolution of mean fitness on the expansion
296   front, as well as simulating and visualising the evolution of average fitness at every deme.
297   Specifically, we developed two programs in 2D: expansion on a strip, and expansion on a disk.
298   Below we talk about the main trial we carried out with the former program.

299   We created a program to simulate a range expansion roughly along an axis in two

dimensions. We call this "2D strip" expansion, due to the habitat constituting a long rectangle, and give the parameters in Table 1. Individuals start in the leftmost demes of the habitat, and we measure the front as the (moving) populated outermost demes on the right. We ran 20 simulation replicates, each with a burn-in phase and an active phase of 1000 generations both; and the same number of replicates without the burn-in phase. Figure S4 (top) shows the evolution of mean front fitness in time for these results, averaged over replicates. We have also displayed the non-normalised mean front fitness for reference. Furthermore, we have displayed the results of applying different normalisation methods.

In addition, we have generated animations of a heatmap that shows the evolution of deme-average fitness throughout the whole expansion at every point in the habitat. Specifically, we generated heatmaps for deme-average fitness and normalised deme-average fitness. In this case, normalisation is such that the average fitness $\overline{w_{x,y}}(t)$ at each deme (x, y are space coordinates, t counts generations from the onset of the expansion; negative t is allowed in the presence of a burn-in phase) is divided by $\overline{\overline{W}}(0)$, the average fitness over the whole habitat at the onset of the expansion. These animations are available in the "Data availability statement". A snapshot of the heatmap of deme-average fitness is given in Figure S4 (bottom).

3.3 2D disk

Next, we have simulated a range expansion on a 2D disk, which is bounded by the circle equation at a certain radius -- parameters are given in Table 1. Aside from this space boundary, the individual movement is not constrained. At the start, the founder individuals were put in the centre of the habitat. We have performed one trial, where we likewise found the evolution of mean front fitness in time, averaged over 20 simulation replicates. This result, with different normalisation methods, is shown in Figure 3. Similar to 2D strip, we have generated heatmaps that show the temporal evolution of (normalised) average fitness at every deme in the habitat, at every generation. This heatmap is available as an animation (see "Data availability statement"). A snapshot of the heatmap of deme-average fitness is given in Figure 3 (right).

3.4 3D cylinder

The main goal of this study was to build a simulation tool that can produce data for range expansion in three dimensions, and analyse the evolution of mean front fitness consistently with the results from 1D and 2D. In particular, we developed two methods for 3D: expansion in a cylindrical habitat, and expansion in a sphere. Here, first, we describe the trial we carried out using the former program.

In this trial, we contained a population inside of a cylinder, that is constrained by a circle
equation at the lateral surface and a barrier at both ends of the cylinder. This trial was roughly
one-directional, meaning that the individuals started in the lowermost demes of the z-axis, and we
measured the front as the (moving) populated highest demes in the direction of the z-axis. We ran
10 simulation replicates, each with a burn-in phase and an active phase of 1000 generations both;
and the same number of simulation replicates without the burn-in phase. Figure 4 (left) shows the
evolution of mean front fitness in time, averaged over 10 simulation replicates. As before, here
we also display the non-normalised mean front fitness for reference and the results of applying
different normalisation methods. We have also generated three-dimensional heatmaps of deme-
average fitness, a snapshot of which can be seen in Figure 4 (right). The animations of these
heatmaps are available in the "Data availability statement". A typical slice image of the heatmap
at height $z = 22$ and generation 200 is presented in Figure S5 (left).

3.5 3D sphere

Finally, we developed a simulation program for an expansion bounded by a 3D sphere of a
constant size. At the start, the founder individuals were put in the centre of the habitat. We have
performed one trial, where we likewise found the evolution of mean front fitness front in time,
averaged over 10 simulation replicates. The result, with different normalisation methods, is shown
in Figure 5. Similar to other trials, we have generated heatmaps for deme-average fitness at each
generation. Sample animations of these heatmaps are available in the "Data availability
statement". A snapshot of the heatmap of deme-average fitness is given in Figure 5 (right). A
typical slice image of the heatmap at height $z = 18$ and generation 642 is presented in Figure S5
(right).

4. DISCUSSION

4.1 1D

Even though we have only used 10 and 20 replicates of simulation data generated with the
program provided in previous research, we can see that simulations generated with our program
yield close values and behaviour (relative error of normalised mean front fitness $\leq 1\%$ everywhere
expect the first 170 generations). The discrepancy between the analytical approximation and
simulated values grows visibly during longer expansions.

367   We have confirmed that the onset mean normalisation method yields values of mean
368 front fitness for simulations with a burn-in phase closest in value to mean front fitness for
369 simulations without a burn-in phase. Cf. the normalisation method where all demes are mapped
370 to the interval [0, 1] every generation ("maximum normalisation" in Figure 2). The rapidly
371 decreasing mean front fitness normalised via "maximum normalisation" can be used to measure
372 the growing difference between the fitness in the core population and on the expansion front.

373   In trial **3**, we saw that the population cannot go extinct in range expansions in idealistic
374 settings with soft selection (cf. range shifts, Gilbert *et al.* (2018)). As expected, in severe
375 conditions, the mean front fitness in "first pair" populations was extremely low compared to "draw
376 until pass" populations. However, when applying the "onset mean normalisation" method, these
377 fitness values were almost equal throughout the expansion. Hence, we can say that onset mean
378 normalisation preserves the trend well, and mean front fitness for these two offspring generation
379 approaches differs only in scale of the absolute value.

381 **4.2 2D**

382 Our 2D disk simulations show that, in an expansion where the population is free to move in 4
383 directions (orthogonally), segregation patterns resembling spokes or circular sectors emerge
384 practically from the start of the expansion (Figure 3). This behaviour has been observed
385 extensively in bacterial mixtures with only a single mutant allele (Hallatschek and Nelson, 2010;
386 Korolev *et al.*, 2010). Moreover, the behaviour was recently confirmed in originally monospecific
387 populations partially or fully developing specialist subpopulations during the expansion (Li and
388 Kardar, 2022). Our results show that this binary segregation behaviour (i.e. with zones where the
389 mutant allele frequency is strictly 0 or 1) changes into a picture with smooth transitions between
390 predominantly mutant and predominantly non-mutant zones in expansions of populations with
391 multiple mutation types. This is speculated to be a blend of separate binary segregation pictures
392 against each of those mutations (no matter received before or during the expansion). This was
393 shown in part in Excoffier *et al.*, 2008. The simple model used in this study can be approximated
394 by a model with co-dominant mutations for low *s*, and it should be noted that for dominant or
395 recessive mutations this behaviour might be different. It might also be different when long-range
396 migration is allowed: in their simulations, Paulose and Hallatschek (2020) found that the
397 possibility for long-range dispersal could make the segregating pattern above disappear.

398   On the other hand, in our 2D strip trial shown in Figure S4, large-scale spatial deviations
399 from the monotonous decrease in fitness with distance from the core population were not present
400 in the narrow space we have set up. We've run separate trials to confirm that this is only the case
401 with narrow expansions that approach an axial expansion such as the 1D expansion, and is also

dependent on the simulation parameters such as the selection coefficient and migration rate. Cf.

other simulations with wide rectangles ($0 \leq y \leq 20$) (Peischl *et al.,* 2015b).

4.3 3D

This is the first time that range expansions of this type were simulated in 3D. The deme-average

fitness in **3D cylinder** (Fig. 8, right) was found to exhibit patterns of smooth combination of

binary segregation pictures, similar to the those observed in **2D disk** (Fig. 7, right), but with more

variation in shape, no more resembling spokes or circular sectors. This pertains to the concurrent

variation in the lateral xy plane with the variation along the z -coordinate. The effect from this

joint variation seems greater that the effects from separate respective variations. The combination

of radial and axial variations here provides a more extreme fitness distribution, where any slight

deviation from the central expansion axis has a chance to result in a drastic falloff in fitness. This

suggests that, aside from the monotonous decrease in fitness proportional to the distance from the

origin, colonisers in 3D space need to take into account the dramatic variation in fitness when

straying from the central axis.

On the other hand, the mean front fitness can still be approximated fairly well using the

analytical formula for one-directional 1D expansions. This is also the case for the spherical

expansion (Fig. 9). Moreover, in all of the expansion scenarios in 2D and 3D, mean front fitness

could still be well approximated with the analytical formula for simple one-directional 1D

expansions. We assume this follows from the fact that, for each deme at the front, the probability

to colonise a further deme is approximately equal to the probability to remain in the present

population. However, a thing to note is the sudden drop in mean front fitness just after the onset

of the expansion in simulations with a burn-in phase. This is a flaw of the burn-in phase, and

could be explained by the fact that the population at the time of the onset experiences the lifting

of a migration restriction, and this allows it to migrate and undergo the biggest loss of genetic

diversity per deme, since the population is still comparatively small.

Genetic drift was shown to be a factor in heterozygosity gradients in a population (Liu

et al., 2006), and in future studies, it would be of value to analyse the distribution of heterozygous

loci and calculate the temporal change in expansion-front heterozygosity in different dimensions.

We have already developed the methods for such simulations that use a finite-site model with

dominance – these methods are available in RESK. Upon modifying our programs to a finite

number of loci (1000 and more), the calculations turned out to be computationally intensive in

multiple dimensions, increasing the simulation time severalfold compared to the infinite-sites

finite-region model in 3D. We would like to explore this model, using more powerful hardware

in a future study, as well as investigate the difference between soft and hard selection.

Nonetheless, the infinite-site finite-region model explored in this study proved to be an efficient

tool for studying the general properties of expansion load.

5. CONCLUSION

A program kit was developed for efficiently simulating range expansions in different habitats in

1, 2 and 3 dimensions. Using them, the mean fitness of expansions in three dimensions was

successfully simulated for the first time. Different normalisation methods were investigated.

"Maximum normalisation" was found to be useful to measure the growing difference between the

fitness in the core population and on the expansion front. In radial 2D (disk) expansions, deme-

average fitness of a population that underwent multiple mutations was found to show a smooth

combination of binary segregation pictures against each of those mutations. This pattern was also

observed as a component in the deme-average fitness in 3D expansions. Overall, complex

fluctuations in deme-average fitness, different from those in radial 2D expansions, were found in

3D expansions. Axial 3D (cylinder) expansions were found to have the highest fitness along the

axis of the expansion.

Conflict of interest statement

The authors declare no conflicts of interest.

Author contributions

Conceptualisation and providing the calculation environment: Gen Tamiya.

Simulation software development: Hikaru Matsuoka and Yurii Kotsar.

Planning and creating example simulations used in the study: Yurii Kotsar.

Manuscript: Yurii Kotsar.

Manuscript revision and corrections: Hikaru Matsuoka and Gen Tamiya.

Data availability statement

We have uploaded the program kit, some simulated result data, as well as animations used in the

study to: *https://github.com/HartreeY/RESK*. Thes program kit consists of the core file (*resk.jl*)

with all the simulation methods, including *rangeexp_inf* – the one used in this study; an optional

graphing methods file (*reskplots.jl*); a file for defining default parameters (*consts.jl*); and the

initialisation file that installs the requirements for this program kit (*init.jl*). We have also included

examples in the *study* folder that correspond to trials in this study. In it, we have also uploaded
animations of the heatmaps for results of 2D and 3D simulations to the *animations* folder.

To generate the data for comparison with Peischl *et al.* (2013), we used the following
program linked in their paper: *https://github.com/CMPG/ADMRE/tree/master*.

BIBLIOGRAPHY

Assis, J., Coelho, N.C., Lamy, T., Valero, M., Alberto, F. and Serrão, E.Á. (2016). Deep reefs are
climatic refugia for genetic diversity of marine forests. J. Biogeogr., 43: 833-844.

Beverton R., Holt S. (1957). On the Dynamics of Exploited Fish Populations. Fisheries
Investigation Series 2 (19). Ministry of Agriculture, London, Fisheries and Food.

Bocharov G. (2016). Spatiotemporal dynamics of virus infection spreading in tissues. PLOS ONE
11(12): e0168576.

Bokma, F. (2010). Speciation and patterns of diversity. Syst Biol, 59: 365-367.

Eyre-Walker, A. and Keightley, P. D. (2007). The distribution of fitness effects of new mutations.
Nature Reviews Genetics, 8 (8), 610-618.

Excoffier, L., & Ray, N. (2008). Surfing during population expansions promotes genetic
revolutions and structuration. Trends in ecology & evolution, 23(7), 347–351.

Gao, Z., Waggoner, D., Stephens, M., Ober, C. and Przeworski, M. (2015). An estimate of the
average number of recessive lethal mutations carried by humans. Genetics, 199 (4), 1243-1254.

Gilbert, K. J., Peischl, S. and Excoffier, L. (2018). Mutation load dynamics during
environmentally-driven range shifts. PLOS Genetics, 14 (9), e1007450.

Hallatschek, O. and Nelson, D. R. (2010). Life at the front of an expanding population. Evolution,
64 (1), 193–206.

Kim B., Huber C., Lohmueller K. (2017). Inference of the distribution of selection coefficients
for new nonsynonymous mutations using large samples. Genetics. 206(1), 345–361.

Kimura, M. (1953). Stepping stone model of population, Annual Report of the National Institute
of Genetics, Japan, 3, 62.

Kimura, M. (1964). Discussion models in population genetics. Vol. 1. Cambridge University
Press.

Kitchel, Z. J., Conrad, H. M., Selden, R. L., and Pinsky, M. L. (2022). The role of continental
shelf bathymetry in shaping marine range shifts in the face of climate change. Global Change

Biology, 28, 5185–5199.

Klopfstein, S., Currat, M. and Excoffier, L. (2005). The fate of mutations surfing on the wave of
a range expansion. Molecular Biology and Evolution, 23 (3), 482-490.

Korolev, K. S., Avlund, M., Hallatschek, O. and Nelson, D. R. (2010). Genetic demixing and
evolution in linear stepping stone models. Reviews of Modern Physics, 82 (2), 1691–1718.

Li, L. H. and Kardar, M. (2023). Specialization at an expanding front. Phys. Rev. E 108, L032402.

Liu H., Prugnolle F., Manica A., Balloux F. (2006). A geographically explicit genetic model of
worldwide human settlement history. American Journal of Human Genetics, 79, 230–237.

Parmesan, C. (2006). Ecological and evolutionary responses to recent climate change. Annual
Review of Ecology Evolution and Systematics, 37: 637-669.

Pateman R. M., Hill J. K., Roy D. B., Fox R., Thomas C. D. (2012). Temperature-dependent
alterations in host use drive rapid range expansion in a butterfly. Science, 336: 1028-1030.

Paulose, J., & Hallatschek, O. (2020). The impact of long-range dispersal on gene surfing.
Proceedings of the National Academy of Sciences of the United States of America, 117(14),
7584–7593.

Peischl, S. and Excoffier, L. (2015b). Expansion load: recessive mutations and the role of standing
genetic variation. Mol Ecol, 24: 2084-2094.

Peischl, S., Dupanloup, I., Kirkpatrick, M. and Excoffier, L. (2013). On the accumulation of
deleterious mutations during range expansions. Molecular Ecology, 22 (24), 5972-5982.

Peischl, S., Kirkpatrick, M. and Excoffier, L. (2015a). Expansion load and the evolutionary
dynamics of a species range. The American Naturalist, 185 (4), E81-93.

Poloczanska et al. (2016). Responses of marine organisms to climate change across oceans.
Frontiers in Marine Science, 3, 1–21.

Sanjuan, R., Moya, A. and Elena (2004). The distribution of fitness effects caused by single-
nucleotide substitutions in an RNA virus. Proceedings of the National Academy of Sciences of
the United States of America, 101 (22), 8396-8401.

Wahl, L. M. and Gerrish, P. J. (2001). The probability that beneficial mutations are lost in
populations with periodic bottlenecks. Evolution, 55 (12), 2606-2610.

Yamano H., Sugihara K., Nomura K. (2011). Rapid poleward range expansion of tropical reef
corals in response to rising sea surface temperatures. Geophysical Research Letters, 38.

APPENDIX A. Floating point error

Due to operating memory limitations, we used Float32 arrays to store fitness values. Since the fitness of founders exponentially influences the fitness of next generations, we designed a test for cumulative errors in mean front fitness values.

We performed the test both for 1D and 3D. In 1D, we ran 15 simulation replicates with the default parameters given in Table 1, however only for 1000 burn-in generations and 1000 generations after onset. These were run both by a program which employs Float32, and one which employs Float64. For every simulation, we synchronised the random seeds given to both programs. We calculated the average over 15 replicates of the mean front population and mean front fitness for each generation, for outputs of both of the programs. We then analysed the relative difference between the averages in the two programs. The difference in average population was negligible, with relative error ~0.3%. The difference in average deme fitness was likewise unimportant with the value < 0.3%.

In 3D, we ran 10 simulation replicates for both programs with the default parameters given in Table 1, but setting 100 burn-in generations and 600 generations after onset. Both differences had a maximum at the last generation in the simulation, indicating that this could be a cumulative floating-point error. The maximum of the relative difference in mean front population was ~0.6%, while the maximum of the relative difference in mean front fitness was also ~0.6%.

It goes without saying, that this is a comparison between Float32 and Float64, and not between Float32 and an ideal floating-point datatype. Still, this error component was found not likely to influence the results of the tests described above in a significant way.

APPENDIX B. Parallel processing

For our simulation kit, we have included the possibility to run parallel simulations, to generate multiple replicates in one go. The results in this study were generated in parallel. In Figure 6, we show a simple test, which we have also included in *https://github.com/HartreeY/RESK/programs/examples/distributed_2.ipynb*. The test compares computation times to run 10 replicates of the same simulation on a variable number of parallel processes: from 1 to 8. The test was performed on a machine with 8 CPUs. Figure 6 shows that even with only 2 processes, the performance is boosted almost twofold, and additional processes further decrease the time in a diminishing manner (although there is a reverse effect due to the

small amount of CPUs). In scientific research, having data from a large number of replicates is invaluable, and the built-in option of parallel simulations should add to the efficiency of generating data for studies that rely on simulations.

Parameter	1D (1)	1D (2)	1D (3)	2D strip	2D disk	3D cylinder	3D sphere
Number of replicates	70	10		20		10	
Burn-in generations	10000	5000		1000	1000	1000	1000
Expansion generations	2500	1250		1000	500	1000	150
Space size (burn-in)	$1 \leq X \leq 5$			$1 \leq X \leq 5,$ $1 \leq Y \leq 10$	$0 \leq R \leq 3$	$1 \leq Z \leq 5,$ $0 \leq R \leq 10$	$0 \leq R \leq 3$
Space size	$1 \leq X \leq 500$			$1 \leq X \leq 250,$ $1 \leq Y \leq 10$	$0 \leq R \leq 100$	$1 \leq Z \leq 300,$ $0 \leq R \leq 10$	$0 \leq R \leq 25$
Selection coefficient	0.005		0.03	0.005			
Mutation rate	0.05		0.1	0.05			
Prop. of del. mutations	0.90	0.45		0.90			
Migration rate	0.05						
Carrying capacity	100						
Filled demes at start	5						
Proliferation rate	2						
Number of gene regions	20						

TABLE 1. List of simulation parameters and their values used in this paper.

FIGURE 1. The algorithm of the cycle in each generation of expansion in our program. "dememax" is the maximum fitness in the deme.

FIGURE 2. Temporal evolution of mean front fitness, as described in **1D 1)**. The grey vertical

line signifies the generation when all demes in the space have been filled.

FIGURE 3. Left: temporal evolution of mean front fitness, as described in **2D disk**. The grey vertical line signifies the generation when all demes in the space have been filled. Right: a snapshot of the evolution of deme-average fitness (given by the colour) at every deme in this expansion, at generation 200.

FIGURE 4. Left: temporal evolution of mean front fitness, as described in **3D cylinder**. Right: a snapshot of the evolution of deme-average fitness (given by the colour) at every deme in this expansion, at generation 500.

FIGURE 5. <u>Left</u>: temporal evolution of mean front fitness, as described in **3D sphere**. The grey vertical line signifies the generation when all demes in the space have been filled. <u>Right</u>: a snapshot of the evolution of mean deme fitness (given by the colour) at every deme in this expansion, at generation 130.

FIGURE 6. The results of a simulation time test done with a variable number of processes (from 1 to 10), to simulate 10 replicates of a range expansion in 10x100 deme habitat, for 100 (burn-in) + 200 generations. The method *rangeexp_1d_inf* from RESK was used.