The mutational landscape of *Bacillus subtilis* conditional hypermutators shows how proofreading skews DNA polymerase error rates

Ira Tanneur[1,2], Etienne Dervyn[1], Cyprien Guérin[2], Guillaume Kon Kam King[2], Matthieu Jules[1,‡] and Pierre Nicolas[2,‡]

[1] *Université Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, 78350, Jouy-en-Josas, France*.

[2] *Université Paris-Saclay, INRAE, MaIAGE, 78350, Jouy-en-Josas, France.*

[‡] To whom correspondence should be addressed: matthieu.jules@inrae.fr; pierre.nicolas@inrae.fr.

Abstract

Polymerase errors during DNA replication are a major source of point mutations in genomes. The resulting rate of spontaneous mutation also depends on the counteracting activity of DNA repair mechanisms, with mutator phenotypes appearing constantly and allowing for periods of rapid evolution in nature and in the laboratory. Here, we use the Gram-positive model bacterium *Bacillus subtilis* to disentangle the contributions of DNA polymerase initial nucleotide selectivity, DNA polymerase proofreading, and mismatch repair (MMR) to the mutation rate. To achieve this, we constructed several conditional hypermutators with a proofreading-deficient allele of *polC* and/or a deficient allele of *mutL* and performed mutation accumulation experiments. With their wide range of mutation rates and contrasting mutation profiles, these conditional hypermutators enrich the *B. subtilis* synthetic biology toolbox for directed evolution. Using mathematical models, we investigated how to interpret the apparent probabilities with which errors escape MMR and proofreading, highlighting the difficulties of working with counts that aggregate potentially heterogeneous mutations and with unknowns about the pathways leading to mutations in the wild-type. Aware of these difficulties, the analysis shows that proofreading prevents partial saturation of the MMR in *B. subtilis* and that an inherent drawback of proofreading is to skew the net polymerase error rates by amplifying intrinsic biases in nucleotide selectivity.

Running title: **Proofreading skews polymerase error rates**

INTRODUCTION

Substitutions, insertions and deletions of single base pairs in genomes can have diverse consequences on encoded molecular functions, from no effect to abrupt change, most often in the direction of deterioration (Eyre-Walker and Keightley 2007). As such, point mutations are a constant threat to the integrity of the genetic information, even if they are also essential to adaptive evolution. In the long term, point mutation rates themselves result from an evolutionary process. An hypothesis, supported by the comparison of mutation rates across the tree of life, postulates that they are simply maintained as low as possible; the limit being the "drift-barrier", where the strength of selection is matched by the opposite pressure of random genetic drift and mutation, presumably biased towards creating weak mutators (Lynch et al. 2016). Mutation rate might thus have nothing to do with the benefit of evolvability for long-term survival, the energetic cost of fidelity, or a biophysical limit. In bacteria, this rate is already typically as low as one mutation introduced in the genome per one thousand generations, but mutation rates one order of magnitude below those observed in nature were reported to arise under scenarios of artificial evolution (Deatherage et al. 2018; Dervyn et al. 2023).

However, mutator phenotypes occur everywhere and their contribution to evolution is difficult to estimate (Taddei et al. 1997; Couce et al. 2017). In asexual populations, where mutator alleles and the mutations that they generate remain linked across successive generations, mutators are advantageous when the potential for fitness improvement is high. For example, in pathogenic bacteria, mutators have been associated with complex antibiotic resistance (Dulanto Chiang et al. 2022), rapid evolution within the host during infection (Oliver and Mena 2010), and atypical virulence traits (Rudenko et al. 2020). Mutators also emerge spontaneously during laboratory evolution in response to applied selective pressures (Sniegowski et al. 1997; Swings et al. 2017). To foster adaptation, conditional mutator circuits and systems have been engineered for some organisms as part of the synthetic biology toolbox (Badran and Liu 2015; Sherer and Kuhlman 2020; Molina et al. 2022). Understanding the molecular factors that determine mutation rates is therefore of strong fundamental and applied interest.

The sources of spontaneous mutations in living cells are diverse, including DNA lesions caused by endogenous and exogenous agents, errors introduced by the DNA polymerase during replication, and the activity of error-prone polymerases recruited in response to stress (Maki 2002). The resulting mutation rate depends on the intensity of these sources and of the counteracting activity of DNA repair mechanisms, which work in concert to ensure that the correct genetic material is passed

on to daughter cells. Two essential mechanisms, conserved from prokaryotes to eukaryotes, ensure

accurate repair of both bulky and non-bulky lesions resulting from DNA damage, including those

induced by reactive oxygen species, a major source of DNA errors (Foster et al. 2015). NER

(Nucleotide Excision Repair) repair various drug- and UV-induced lesions (*i.e.* bulky lesions),

while BER (Base Excision Repair) repair lesions caused by various chemical assaults, such as

alkylation, oxidation, deamination, etc. (*i.e.* non-bulky lesions).The accuracy of DNA replication

depends on three critical mechanisms: the initial selectivity of the DNA polymerase, which is

responsible for inserting the correct nucleotide; the proofreading, which removes misincorporated

nucleotides through polymerase-associated exonucleases; and the mismatch repair (MMR), which

adds a second layer of error correction shortly after replication (Ganai and Johansson 2016).

In bacteria, genome replication is carried out by a multiprotein machine classified into the C-family

of DNA polymerase holoenzymes, in which the catalytic polymerase α-subunit exists in two

primary forms, DnaE and PolC (Timinskas et al. 2014). A representative example of DnaE is found

in the extensively studied Gram-negative model bacterium *Escherichia coli*. Conversely, PolC is

predominant in low-GC Gram-positive bacteria, such as *Bacillus subtilis* (Sanjanwala and Ganesan

1991). In this organism, replication elongation involves two essential polymerases, PolC and DnaE

(Dervyn et al. 2001), which have distinct functions: PolC does most of the DNA synthesis, but only

DnaE can elongate from RNA primers on the lagging strand before passing the DNA fragment to

PolC. In the absence of proofreading and MMR, the error rate of the *E. coli* α-subunit replication

machinery has been estimated to be approximately 10^{-6} per base pair per generation both *in vitro*

(Fujii et al. 1999) and *in vivo* (Niccum et al. 2018). Given this error rate and the size of the *E. coli*

genome, about 5 mutations are expected to be introduced per generation.

The exonuclease domain essential for proofreading is encoded as an integral part of the vast

majority of PolC polymerases (Timinskas et al. 2014), including *B. subtilis* PolC. In contrast, DnaE

polymerases do not have their own exonuclease domain. The proofreading activity of the *E. coli*

DNA PolIII holoenzyme containing DnaE is based on an exonuclease domain located in the ε-

subunit. Errors made by polymerases devoid of proofreading can also sometimes be corrected by a

process known as proofreading in *trans,* or extrinsic proofreading, which is well described between

eukaryotic DNA polymerases (Zhou et al. 2021). In *B. subtilis*, data suggest that the PolC

exonuclease is able to correct errors made by the error-prone DnaE polymerase (Bruck et al. 2003;

Paschalis et al. 2017).

The MMR is a universal mechanism that is responsible for correcting errors that occur during DNA

replication and escape proofreading. Upon detection of a replication error, the mismatch-sensing

protein, MutS, recruits MutL. Most prokaryotic and eukaryotic MutL homologs, from humans to

bacteria, possess a highly conserved endonuclease active site that serves to remove mismatches

(Pillon et al. 2010; Bolz et al. 2012). *E. coli* has been the primary model for studying MMR, but its

MutL lacks the endonuclease activity encoded here in a distinct protein, MutH, which specifically

nicks the unmethylated and thus nascent strand (Lenhart et al. 2012). In the absence of MutH and

Dam methylation, the process that guides MutL to the nascent strand remains unclear in most

prokaryotes and all eukaryotes (Kadyrov et al. 2006). In bacteria, MMR increases the fidelity of the

chromosomal DNA replication pathway approximately 100-fold, and MMR is considered to be for

repairing the most frequent replication errors (Lujan et al. 2012). Mutator phenotypes found in

nature are often caused by mutations that inactivate the MMR.

Studying organisms such as *B. subtilis*, with a PolC polymerase and an MMR pathway more widely

conserved across biology can provide important insights into the coordinated functioning of these

systems within living cells (Klocko et al. 2011). Extending previous work characterizing the

mutation profiles of MMR-deficient *B. subtilis* strains (Sung et al. 2015; Schroeder et al. 2016), the

main goal of our study was to jointly assess the respective contributions of initial nucleotide

selectivity, proofreading and MMR to the mutation rates and their interdependencies. For this

purpose, we constructed and analyzed several conditional hypermutators. Analysis of the data in

light of several mathematical models suggests that proofreading is needed to avoid partial saturation

of the MMR, as previously reported in *E. coli* (Schaaper 1988; Niccum et al. 2018), but also that an

inherent effect of proofreading is to skew the net polymerase error rate. With their wide range of

mutation rates and contrasting mutation profiles, the conditional hypermutators also enrich the *B.*

subtilis synthetic biology toolbox for directed evolution.

RESULTS

The mutation rate of *Bacillus subtilis* can be increased up to 6,000 times. We have constructed five mutant strains with expected hypermutator phenotypes from a strain derived from *B. subtilis* 168. The first two strains are constitutively MMR deficient as a result of single deletions of *mutS* and *mutL*. The other three were engineered for conditional inactivation of either or both of MMR or proofreading and were therefore expected to be inducible hypermutators (**Figures S1** and **S2**). In these three strains, the IPTG-inducible promoter P_{hs} controls the expression of mutant alleles selected for their ability to competitively displace their functional counterparts. The first allele, designated *mutL**, has a mutation in the region encoding the ATP hydrolysis active site of MutL which has been reported to have a dominant negative effect (Bolz et al. 2012). The second allele, designated *polC**, encodes a proofreading deficient variant of PolC with a mutation in its exonuclease domain (Sanjanwala and Ganesan 1991). The final strain, which was expected to have the highest mutation rate under full induction, expresses these two deficient alleles in a synthetic operon (*mutL* polC**). The following notations are used for the reference parental strain derived from 168 and the five mutant strains: R^{168}, ΔL, ΔS, L^*, C^*, LC^*.

Fluctuation assays were performed (**Figure S3**) to compare the rate of mutation to rifampicin resistance of these strains. Point estimates and confidence intervals are shown in **Figure 1**, and results are detailed in **Table S1**. In the absence of IPTG, the estimated mutation rate of R^{168} was 9.74×10^{-10} per generation. Constitutive inactivation of the MMR in ΔL and ΔS increased the mutation rate by a factor of approximately 85, with no statistically significant difference between the two strains. This is close to the factor of about 60 previously obtained for a double deletion of *mutL* and *mutS* in the *B. subtilis* PY79 genetic background with a fluctuation assay also based on rifampicin (Schroeder et al. 2016). Mutation rates in the absence of IPTG were slightly higher for the inducible strains (*L**, *C**, *LC**) than for R^{168} (up to 1.09×10^{-8} for *LC**), likely reflecting the low basal activity already described for P_{hs} (Guiziou et al. 2016). From there, the mutation rate of the three inducible strains increased with IPTG concentration, reaching a plateau between 50 and 100 µM. At full induction (100 µM IPTG), the mutation rates ranging from 3.66×10^{-7} for *L** to 5.78×10^{-6} for *LC**. The mutation rate in *L** is comparable to or slightly higher than in ΔL and ΔS, while the mutation rate in *LC** represents an approximately 6,000-fold increase over R^{168}.

Figure 1. Mutation rate to rifampicin resistance measured by fluctuation assays for increasing IPTG concentration. Each color corresponds to a strain; vertical bars represent 95% confidence intervals.

Much higher mutation rates than in the reference strain may induce stress responses, potentially altering the physiology of each mutant differently. However, we did not detect any substantial impact on growth in 96-well microtiter plates (**Figure S4**). We also performed transcriptomics experiments on the R^{168}, $L*$, $C*$ and $LC*$ strains in the presence of 100 µM IPTG. The analyses did not reveal any significantly differentially expressed genes between the strains. However, they did allow us to quantify the expression of the mutant alleles relative to wild-type alleles for *mutL* and *polC*. Upon induction, the mutant allele accounted for 96-98% of the total mRNA pool for the gene in question (**Table S2**), which is consistent with the results of fluctuation assays that give a mutation rate in $L*$ as high as in ΔL and ΔS (*i.e.* complete inactivation of the MMR). We therefore concluded that the mutation profiles of these strains can be attributed to the sole inactivation of the two targeted DNA repair pathways.

Highest mutation rates are counter-selected in mutation-accumulation experiments. Mutation-accumulation experiments give access to the molecular nature of mutations (Lynch et al. 2016), in contrast to fluctuation assays, which only provide a mutation rate aggregated across a range of mutations that confer a screenable phenotype. For each of the six strains (R^{168}, ΔL, ΔS, $L*$, $C*$, $LC*$), four independent mutation-accumulation lines (MA-lines) were propagated by repeated cycles (MA-steps, **Figure S5**) of colony picking, dilution, and plating on LB, in the presence of

17   6

100 µM IPTG when relevant. By randomly selecting a single colony, each MA step creates a

bottleneck in the propagated population, the whose purpose of which is to limit genetic diversity,

thereby maximizing random genetic drift and minimizing natural selection. The interval between

two bottlenecks, one MA-step, was estimated to be 25.6 generations on average. We performed

whole-genome sequencing at the endpoint of each line; we also sequenced intermediate time-points

(up to 4 for LC^*) to detect changes in mutation rates (**Figure S5**). A total of 56 clones isolated after

1 to 37 MA-steps were sequenced. Point mutations - substitutions, insertions (ins), and deletions

(del) - were identified. Mutation rates per base pair (bp) and generation (abbreviated $bp^{-1}.gen^{-1}$)

were estimated for each time interval of each MA-line. Previously collected data from mutation-

accumulation experiments using *B. subtilis* 3610 (R^{3610}) and its corresponding Δ*mutS* mutant strain

(ΔS^{3610}) were also incorporated to increase statistical power (Sung et al. 2015; Sung et al. 2016). The

detailed list of all mutations found is provided in **Table S3**.

In the four independent lines of R^{168}, only one nucleotide substitution was identified after 37 MA-

steps, giving an estimated substitution rate of 7×10^{-11} $bp^{-1}.gen^{-1}$ (**Table 1**). This rate was not

statistically significantly different from a previous report on *B. subtilis* (Sung et al. 2016),

recalculated to 3.4×10^{-10} $bp^{-1}.gen^{-1}$ (**Table 1**).

Table 1. Aggregated number of substitutions, substitution rate, and proportion of
transversions for each strain.

Strain[a]	Lines	Generations [b,c]	Substitutions[c]			Substitution rate ($bp^{-1}.gen^{-1}$) [95% CI]	Proportion of transversions [95% CI]
			Total	ts[d]	tv[e]		
R^{168}	4	3,790	1	1	0	7.0×10^{-11} [0.18-39×10⁻¹⁰]	0.00 [0.00-0.98]
R^{3610}	49	248,920	319	238	81	3.4×10^{-10} [3.0-3.8×10⁻¹⁰]	0.25 [0.21-0.31]
ΔS^{3610}	19	38,000	4,844	4,711	133	3.4×10^{-8} [3.3-3.5×10⁻⁸]	0.03 [0.02-0.03]
ΔL	4	2,151	149	147	2	1.8×10^{-8} [1.5-2.1×10⁻⁸]	0.01 [0.00-0.05]
ΔS	4	2,151	157	155	2	1.9×10^{-8} [1.6-2.2×10⁻⁸]	0.01 [0.00-0.05]
L^*	4	2,151	113	111	2	1.4×10^{-8} [1.1-1.7×10⁻⁸]	0.02 [0.00-0.06]
MMR-[168]	12	6,453	419	413	6	1.7×10^{-8} [1.6-1.9×10⁻⁸]	0.01 [0.01-0.03]
C^*	4	1,895 (256)	395 (19)	348 (18)	47 (1)	5.5×10^{-8} [5.0-6.1×10⁻⁸]	0.12 [0.09-0.16]
LC^*	4	230 (897)	502 (627)	484 (599)	18 (28)	5.5×10^{-7} [5.2-6.3×10⁻⁷]	0.04 [0.02-0.06]

[a] The label MMR-[168] corresponds to the aggregation of data from the three MMR-deficient strains
constructed from R^{168} (ΔL, ΔS, L^*). Data for strains R^{3610} and ΔS^{3610} were obtained from Sung *et al.* (2015)
and mapped to the R^{168} genome sequence.
[b] Conversion between MA-steps and generation based on an estimated average number of generations per
step: 25.6 here, 27.53 in Sung *et al.* (2015).
[c] Number of substitutions in the reference subset of positions well covered by the sequencing data (3,795
kbp). Between parentheses: number of generations or substitutions in time intervals after a detected decrease
in mutation rate (discarded from analysis).

22
[d] Number of transitions.
[e] Number of transversions.

Between 113 and 157 nucleotide substitutions were identified after 21 MA-steps in each of the ΔL,

ΔS and L^* strains, resulting in point estimates of the substitution rates between 1.4×10^{-8} and 1.9×10^{-8}

^{8} bp^{-1}.gen^{-1} (**Table 1**). These rates were not statistically significant from each other which led us to

aggregate the data collected for these three strains under the label MMR-[168] (**Table 1**). Sequencing

an intermediate time-point located at the end of MA-step 11 for the L^* strain did not reveal any

difference in the rates of accumulation between the first and second parts of the evolution (**Figure 2**

and **Figure S6**). This suggests that in MMR-deficient strains, substitutions identified at endpoints of

the MA-lines result from accumulation at a constant rate.

Figure 2. Evolution of substitution rate along mutation accumulation lines. Examples are
shown for one MA line from each strain with an inducible mutation rate. The rate is calculated from
the number of new substitutions identified within each interval. Sequencing intervals are
represented by horizontal brackets and 95% confidence intervals for estimated rates are reported by
vertical bars. Red crosses indicate sequencing intervals with significantly decreased mutation rates
discarded from downstream analyses.

In contrast to L^* MA-lines, both C^* and LC^* MA-lines showed a tendency towards decreasing

substitution rates during their evolution, with differences between the two strains in terms of

frequency, temporality, and magnitude of decrease (**Figure 2** and **Figure S6**). A statistically

significant decrease was detected for a single C^* MA-line (C^*_3) and this decrease was only detected

during the second half of the 21 MA-steps evolution. In contrast, a decrease was detected in all LC^*

MA-lines, as early as during the second MA-step for LC^*_2 (p-value $= 8.4 \times 10^{-3}$), and during MA-

steps 3-6 for the three other LC^* MA-lines. The magnitude of the decrease was a factor

23   8
24

approximately 2.5x for $C*_3$ but reached approximately 50x for $LC*_4$. Therefore, despite

heterogeneity between MA-lines, decreases were globally more frequent, quicker and of larger

magnitude for $LC*$ than for $C*$. Importantly, in $LC*$ MA-lines, 56% of the 1,129 identified

substitutions occurred in intervals affected by a decrease in the substitution rate (**Table 1**). To

determine the mutation rates of the strains, we only considered data from time intervals before any

detected decrease. This led to point estimates of the substitution rates for the $C*$ and $LC*$ strains of

$5.5×10^{-8}$ and $5.5×10^{-7}$ bp^{-1}.gen^{-1}, respectively (**Table 1**).

Statistically significant decreases in indel rate were also observed (**Figure S7**) and correlated with

decreases in substitution rate (**Figure S8**). This is consistent with both proofreading and MMR

contributing to correction of indel errors made by PolC polymerase activity (**Supplementary**

Methods and Results 1.1), the rate of which increases with homopolymer length (**Figure S9**). The

calculated insertions and deletions rates are given in **Table S4**.

Nonsynonymous mutations were found in the inducible synthetic circuits of 4 out of 5 MA-lines

exhibiting a decrease in mutation rate (**Table S5**, **Figure S10, Supplementary Methods and**

Results 1.2). Given the total number of mutations in the $LC*$ lines and the size of the $polC$ gene,

the number of mutations found on the $polC*$ allele is four times higher than expected in the absence

of selection (chi-squared test with simulated p-values, p-value=$4.1×10^{-2}$). This finding echoes

previous studies reporting changes in mutation rates in mutation-accumulation experiments

(Perfeito et al. 2014; Singh et al. 2017). Indeed, recent studies have concluded that positive

selection is possible in mutation-accumulation experiments despite the extreme bottlenecks imposed

on the population (Mahilkar et al. 2022; Wahl and Agashe 2022). Here, the over-representation of

mutations in the genetic elements that confer the strongest hypermutator phenotypes indicates

adaptive evolution through positive selection to reduce the mutation rate. Furthermore, it provides

an *a posteriori* experimental justification for the decision to limit mutation-accumulation

experiments on proofreading-deficient *E. coli* to 3-6 MA-steps to minimize selection (Niccum et al.

2018).

Proofreading repairs transversion errors at least as well as transition errors. The sequence

data do not provide information about the DNA strand on which the error that led to the mutation

originally occurred. To record substitutions, we opted for a framework in which the reference base

is the pyrimidine (C or T) of the Watson-Crick pair at the genomic position where a mutation is

observed. This allows for a prior-free analysis of strand asymmetries in mutation profiles, and

follows the convention used in cancer research (Tate et al. 2019).

In all strains, a slightly higher number of mutations were found on C bases than on T bases (**Figure**

3). All strains also showed a predominance of transitions over transversions, but the strength of this

bias differed between strains (**Figure 3** and **Table 1**). The highest proportion of transversions

among substitutions, found in R[3610] (point estimate 0.25), is about 10 times higher than the lowest,

found in the MMR- strains (ΔS[3610] and MMR-[168] strains, point estimates 0.03 and 0.01,

respectively). The *C** strain showed an intermediate proportion (point estimate 0.12). The small

number of transversions made it difficult to compare those changing the reference pyrimidine (C or

T) to an A and to a G. Nevertheless, the *C** and R[3610] strains may present an excess of C→A over

C→G not seen in other strains. The approximately 10-fold increase in the proportion of transitions

when comparing the MMR-[168] strains to the R[3610] strain is consistent with previously published

results on ΔS[3610] (Sung et al. 2015). Indeed, there is general tendency across microorganisms for

MMR to reduce the transition rate much more than the transversion rate (Lujan et al. 2012; Long et

al. 2018).

Figure 3. Substitution rates for each type of substitution measured by mutation accumulation

experiments. 95% confidence intervals are reported. No C→G and T→G mutations occurred in

MMR-[168], only the upper limit of the 95% CI is shown.

The fold-change in mutation rate was greater when proofreading was inactivated in the presence of

MMR than in its absence (**Figure 3** and **Table 1**), otherwise the fold-change in the mutation rate

when both repair systems are inactivated compared to the wild-type would be the product of the

fold-changes observed when they are inactivated separately (a scenario later referred to as

multiplicative fold-changes). This suggests that the efficiency of correction by MMR decreases

simultaneously with proofreading inactivation (i.e. in C^* compared to R^{3610} or R^{168}). In *E. coli*, the

fold-change in substitution rate upon inactivation of PolIII holoenzyme proofreading was also

reported to be much lower in MMR-deficient cells than in wild-type (Schaaper 1988; Niccum et al.

2018). This was interpreted as a consequence of MMR saturation due to the high number of errors

introduced during DNA replication, a hypothesis further supported by direct assays of MMR

activity and restoration by MMR overexpression (Schaaper 1988; Schaaper and Radman 1989).

Comparing C^* to R^{3610}, inactivation of proofreading significantly reduced the proportion of

transversions among substitutions (from 0.25 to 0.12), which may also result from MMR saturation

in C^*. Below, we will formally explore the MMR saturation hypothesis using a model-based

analysis that integrate information about the chromosomal context of the mutations (adjacent

nucleotides and strand).

In the absence of MMR, inactivation of PolC proofreading resulted in a slight increase in the

proportion of transversions (from 0.01 in MMR-^{168} to 0.04 in LC^*). This difference is not

statistically significant (**Table 1**) but suggests that proofreading corrects errors leading to

transversions with at least as much efficiency as those leading to transitions.

Strand-asymmetry of substitution rate at C:G sites is only visible after proofreading. To

further characterize the two DNA repair systems, we examined for each strain how substitution

rates varied with distance from the origin of replication, strand orientation relative to replication or

transcription, coding or non-coding status, and transcription level. For these analyses, we counted

substitutions in the different chromosomal contexts and calculated the corresponding "local"

substitution rates.

In MMR-^{168} strains, as in ΔS^{3610} and R^{3610} strains, substitution rates were significantly higher when C

is on the leading than on the lagging strand (**Figure 4A**). This agrees with previous results in *B.*

subtilis (Sung et al. 2015), where all mutations were recorded as a change on the "strand templating

the leading strand" (*i.e.* the lagging strand), and which showed higher mutation rates for G bases in

R^{3610} and ΔS^{3610} strains. This bias was not present in the C^* and LC^* strains, which are proofreading

deficient. In addition to the strong asymmetry at C:G sites, we detected a weaker but statistically

significant replication-oriented asymmetry in substitutions at T:A sites: the substitution rate is

higher when T is on the lagging strand, the difference between the two strands being statistically

significant for all our hypermutator strains (**Figure 4A**). In R^{3610}, this bias at T:A sites appears to be

less pronounced and may even be absent. Consistent with the conclusions of Schroeder et al.

(2016), analysis of localization with respect to transcription, which is most often collinear with

replication in *B. subtilis*, did not indicate a contribution of transcription-related processes to these

asymmetries between strands in any of the hypermutator strains (**Supplementary Methods and**

Results 1.3, Figure S11AB).

Figure 4. Substitution rate as a function of replication strand and neighboring nucleotides.
The pyrimidine of the pair determines the strand of a mutation site. Rates are plotted for each
pyrimidine and each genotype. Error bars represent the 95% confidence intervals. **A.** Effect of

orientation with respect to the replication strand. **B.** Effect of orientation both with respect to both the replication and transcription strands. **C.** Effect of the 3'-adjacent nucleotide. **D.** Effect of the 5'-adjacent nucleotide.

In wild-type, the replication-oriented asymmetry of substitution rates at G:C sites is a prominent feature of the mutation profile. Consistent with previous analyses of MMR-deficient strains, including ΔS^{3610}, this bias was also detected in MMR-[168] strains. Our data revealed its absence in *C** and *LC** strains. A similar observation was made in *E. coli* proofreading-deficient strains, leading to the interpretation that proofreading is strand-biased and produces this bias (Niccum et al. 2018), but a concurrent explanation will be discussed below. In contrast, the wild-type had little or no asymmetry at A:T sites, whereas the hypermutator strains have such asymmetry; error correction systems, and in particular the MMR, tend to eliminate this asymmetry.

In parallel, it is intriguing to observe the presence of two trends detected only in wild-type: a higher substitution rate in non-coding regions (**Figure S11C**) and, to a lesser extent, in the half-chromosome near the replication origin (**Figure S11D**). A wild-type specific higher substitution rate in non-coding than coding regions, specific to the wild-type has also been reported in *E. coli* and was interpreted as an indication that "MMR preferentially repairs coding sequences" (Lee et al. 2012; Foster et al. 2018). Alternatively, trends observed exclusively in the wild-type may correspond to substitutions originating from processes not subject to correction by proofreading and MMR, which could be masked by increased substitution rates in hypermutator strains.

Polymerase initial nucleotide selectivity shapes the distribution of mutations even after proofreading. In all strains, the substitution rate was strongly influenced by the nucleotide adjacent to the focal pyrimidine (**Figure 4C** and **Figure 4D**). This observation, previously made in wild-type and MMR-deficient strains (Sung et al. 2015), extends to proofreading-deficient strains. Simultaneously considering the adjacent nucleotides on both sides (**Table S6** and **Figure S12**) and the replication strand requires binning the counts into 64 replication-stranded triplets. To mitigate the dimensionality problem, exemplified by the absence of observed substitutions for some bins, we adopted a Bayesian estimation framework that incorporates the mean and standard deviation of log-transformed rates as strain-specific hyperparameters (**Supplementary Methods and Results 1.4**). Information from the entire distribution is thereby used to establish point estimates and credibility intervals of the substitution rate for a given triplet (**Figure 5A**).

Figure 5. Comparison of stranded-triplet substitution profiles between genotypes. Each stranded triplet corresponds to the pyrimidine of the pair and its 5' and 3' nucleotides, distinguishing pyrimidines on the leading and lagging strands of replication. ΔS, ΔL, and $L*$ (IPTG 100 µM) are aggregated as MMR-[168]. **A.** Heatmap representation of the stranded-triplet substitution profiles ($\log_2$ ratio of the estimated rate with respect to the mean for the genotype) in the different strains. Rates were estimated using a Bayesian method involving a log-normal prior and hyperparameters. Estimates based on the absence of substitutions in this stranded-triplet context are indicated by "0" in the cells of the heatmaps. Triplets are ordered in decreasing order of non-stranded empirical substitution rates. **B.** Beeswarm representations of the distributions of Bayesian estimates of stranded-triplet substitution rates, with the standard deviation of the estimates in log10-scale reported above. **C.** KL divergences from a uniform distribution (same rate for each triplet), derived from robust entropy estimates.

Based on these estimates, we measured a very strong correlation between the stranded-triplet substitution profiles of MMR-[168] and ΔS^{3610} (Pearson correlation r=0.91 for log-transformed rates), confirming that our MMR-[168] background is very close to the previously studied mutant (Sung et al.

2015). There is also a strong correlation between the stranded-triplet substitution profiles of *LC**

and proofreading-proficient MMR-deficient strains (r=0.79 between *LC** and ΔS^{3610}). This is

consistent with the idea that, after proofreading and before correction by MMR, most of the errors

leading to substitutions are due to   misincorporation by the PolC polymerase that escaped

proofreading. Since MMR removes ~98-99% of these errors (**Table 1**), they also represent the

majority of errors corrected by the MMR.

We also found correlations between the wild-type stranded-triplet substitution profile and those of

all hypermutator strains. The correlations are highest with the proofreading-proficient MMR-

deficient strains (r=0.72 with ΔS^{3610}, r=0.71 with MMR-[168]) and remain statistically significant with

the proofreading-deficient strains (r=0.58 with *LC** and r=0.57 with *C**, *p*-values<10^{-6}). The

correlation between the stranded-triplet substitution rates profiles in *LC** and the wild-type, whose

global substitution rate is 1600 times lower (**Table 1**), fits remarkably with the working hypothesis

that PolC misincorporations that escaped proofreading and MMR substantially shape the

substitution profile of the wild-type.

Proofreading squares the biases of the initial polymerase selectivity. As measured by the

standard deviation and the span of their distributions, point estimates of stranded-triplet substitution

rates in MMR-[168] and ΔS^{3610} are approximately 10-fold more dispersed than in other strains,

doubling the dispersion on a logarithmic scale (**Figure 5B**). Because point estimates deduced from

small counts are not precise, we sought an alternative approach that could directly quantify

dispersion. Using a coincidence-counting method for entropy estimation developed to be robust to

sparse sampling (Nemenman et al. 2001), we calculated the Kullback-Leibler (KL) divergence

between the unknown underlying distribution of observed counts and the theoretical distribution

assuming a uniform substitution rate (expected count proportional to the number of occurrences of

the triplet in the chromosome). Estimates of KL divergences (**Figure 5C**) confirmed both the

similar level of dispersion of substitution rates in the wild-type (R^{3610}) and proofreading-deficient

strains, and the comparatively much higher dispersion in the MMR-deficient strains (approximately

2.5-fold higher divergence from uniform). In other words, proofreading disperses substitution rates,

while MMR activity counteracts this dispersion. This observation is consistent with the conclusions

of a study on proofreading of the *E. coli* PolIII holoenzyme (Niccum et al. 2018), and has been

interpreted as suggesting a compensation of proofreading biases by opposite biases of MMR

correction; a concurrent hypothesis to explain the apparent compensation of the biases is presented

in the discussion.

Interestingly, after proofreading (but before MMR correction), the stranded-triplet substitution rates

become more dispersed, but remains very similar in terms of the direction of the biases compared to

before proofreading (**Figure 5A**). This observation is puzzling because it implies that proofreading

amplifies biases that already arise from the sole polymerase activity, rather than simply masking the

initial biases with its own biases of greater amplitude. To better understand the implications of this

observation, we can formulate a minimal model in which the proofreading activity reduces the

initial error probability of the polymerase, denoted $\gamma[i]$, through a two-step process: first, the

detection and removal of a misincorporated nucleotide with probability $d[i]$; second, the

reincorporation of a nucleotide with the error probability $\gamma[i]$ characteristic of the initial polymerase

activity. The error probability after proofreading writes then $e[i]=\gamma[i](1-d[i])/(1-\gamma[i]d[i])$, where the

term $(1-\gamma[i]d[i])$ accounts for the possibility of cycling the two-step process if a new incorporation

error follows the removal. If $d[i]$ is the same for all i, the possibility of cycling, by itself, already

amplifies the initial biases. However, this effect is negligible as long as $\gamma[i]$, which corresponds to

the substitution rate observed in the *LC** strain ($<10^{-5}$ in **Figure 5B**), remains extremely small

compared to 1. Doubling the biases on a logarithmic scale (*i.e.* $e[i]/e[j]=(\gamma[i]/\gamma[j])^2$), as

approximately observed in our data, implies a probability of non-detection and removal in the first

step of proofreading $(1-d[i])$ proportional to $\gamma[i]$, the error probability of the single polymerase

activity. The most common errors made during initial incorporation also the least likely to be

corrected during proofreading.

A theoretical model suggests that MMR can prevent up to 4 mutations per generation before

saturation. The apparent efficiencies of MMR and proofreading are strongly influenced by the

presence or absence of the other system. As evoked above, proofreading reduced the total

substitution rate by a factor 162 in MMR-proficient cells (μ_{R3610}/μ_{C*}, where μ is the mutation rate

under consideration and using the ML estimates from **Table 1**) but only by a factor 32 in MMR-

deficient cells ($\mu_{MMR-168}/\mu_{LC*}$). Similarly, MMR reduced the substitution rate more in the presence

than in the absence of proofreading. According to Bayesian estimates of substitution rates, none of

the triplets or mutation types (ts, tv, ins, del) clearly contradicts this observation (with a posterior

probability of being above the diagonal greater than 5% for all triplets; **Figure 6**). The trend

becomes even more pronounced when ΔS^{3610} is used instead of MMR-[168] to represent the MMR-

deficient proofreading-proficient background (**Figure S12**).

Figure 6. Effects of proofreading and MMR in the presence or absence of the other system. Effects are measured as the log10-ratio of substitution rates. Left plot: effect of proofreading in the presence or absence of MMR. Right plot: effect of MMR in the presence or absence of proofreading. Around each point, the 50% and 95% marginal credibility intervals on horizontal and vertical axes, computed from the quantiles of the posterior distributions, are represented by segments (bold and dark *vs.* thin and light, respectively). The data used for the MMR- substitution profile is MMR-[168].

To thoroughly investigate the compatibility of the data collected in *B. subtilis* with a saturation mechanism, we formulated a mathematical model in which the mutation rate, $\mu_{C*}[i]$, in strain *C** for a given triplet or type of mutation i is determined by the equation:

$$\mu_{C*}[i]=\gamma[i](\theta+(1-\theta).q_{MMR}[i]),$$

where $\gamma[i]$ is the error rate before correction by proofreading or MMR, and θ is a mixture parameter common to all values of i. It corresponds to the proportion of errors made by the proofreading-deficient PolC that occur in a physiological context of MMR saturation (generating mutations distributed as in *LC**). The complementary proportion $(1-\theta)$ is subject to correction by MMR, which reduces the number of errors by a factor qMMR[i]. In this model, whose assumptions are presented along with an associated Bayesian estimation procedure in **Supplementary Methods and Results 1.5**, the mutation rate in *C** can be expressed as a function of the rates in the three other backgrounds by identifying $q_{MMR}[i]$ with $\mu_{wt}[i]/\mu_{MMR-}[i]$ and $\gamma[i]$ with $\mu_{LC*}[i]$.

This parameterization was used to estimate the mixture parameter θ and to check the agreement of the model with the experimental data (**Figure 7**). In practice, the posterior distribution of θ was estimated using either the substitution rates for the 64 replication-oriented triplets (with MMR-[168] or

ΔS^{3610} to represent the MMR-deficient background) or the rates of the 4 types of mutations (ts, tv,

ins, del). These three posterior distributions are very similar (**Figure 7A**), with the posterior mean

for θ varying only from 0.071 to 0.084. The observed counts, aggregated by mutation type (**Figure**

7C) and by triplet (**Figures S14** and **S15**), fall within the prediction intervals of the model,

indicating a good fit to the experimental data. Notably, although the fraction of errors made by the

proofreading-deficient PolC in the context of MMR saturation is small ($\theta<10\%$), these errors

account for the majority of the mutations observed in strain C^* (**Figure 7A**).

Figure 7. Parameter estimation of the MMR-saturation model and evaluation of the fit to

experimental data. A. Posterior distribution of the mixing parameter θ corresponding to the

proportion of polymerase errors in strain C^* arising in the context of saturated MMR. The

corresponding proportion of mutations observed in C^* (i.e. after MMR correction) is shown in the

inset plot. **B.** Relationship between the mixing parameter θ and the number of mutations before

MMR saturation in a simplified model of replication: one replication per generation, the first

mutations are subjected to MMR correction until MMR saturation. **C.** Evaluation of the fit of the

MMR-saturation model to experimentally measured rates of transition, transversion, insertion, and

deletion. Points represent empirically calculated mutation rates, *i.e.* the number of observed

mutations divided by the number of possible sites in the genome and the number of generations.

Colored areas represent the distribution of values for the empirical rates simulated under the

posterior distribution of the model parameters (50% of the density in the dark area, 95% including
also the light area).

In a simple mechanistic model, these values of θ may correspond to the capacity of the MMR to

prevent (with a probability of failure $q_{MMR}[i]$) up to 4 mutations per generation (**Figure 7B**).

Alternatively, θ values can be interpreted as reflecting the fraction of errors made by the

proofreading-deficient polymerase before it travels a certain distance after a first error

(**Supplementary Methods and Results 1.5**).

DISCUSSION

Aggregating mutations in counts: a necessary evil. In principle, our experimental data could also be explained by a model that does not involve MMR saturation. This alternative model recognizes that counting mutations on the genome implies aggregating them by type or context, which are likely to encompass subclasses of mutations arising from errors that are not corrected with the same probability of success. In the extreme scenario where MMR and proofreading correct non-overlapping sets of errors, the increments observed in mutation rates when each system is inactivated would simply add up in cells where both systems are inactivated. This idea has already been mentioned and refuted for mutational signatures in human cancers (Haradhvala et al. 2018). In fact, while changes in mutation rates associated with each repair system are not multiplicative, they are clearly more than additive. This raises interest in a more general model that considers aggregated subclasses of mutations in the counts and arbitrary correction rates. We have algebraically explored this model in its simplest form with only two subclasses (**Supplementary Methods and Results 1.6**). Interestingly, when the probability of error correction differs between subclasses for both MMR and proofreading, the aggregation creates apparent epistasis in the sense that the effect of inactivating one system, as measured by the mutation rate fold-change, depends on the presence or absence of the other system.

With more parameters than data points (5 parameters for 4 mutation rates), even the simplest two-subclass scenario can fit almost any data set, explaining additive to super-multiplicative effects. The model is therefore difficult to falsify. We note, however, that the sign of this epistasis depends on whether the systems have similar or opposite specificities: if they tend to repair the same subclasses of errors the apparent epistasis will be positive (super-multiplicative), otherwise the epistasis will be negative (sub-multiplicative, as in our data). Explaining negative epistasis with this model is therefore at odds with the widely accepted idea that MMR correction is mostly coreplicative and targets the same errors as those corrected by proofreading, *i.e.* Watson-Crick mismatches introduced by DNA polymerase during DNA replication. If subclass aggregation does not contribute significantly to the observed apparent epistasis, which is well explained by MMR saturation, it still complicates the interpretation of the apparent efficiency of repair systems deduced from mutation profiles.

A cautious interpretation of the apparent efficiency of proofreading and MMR. The proofreading-deficient strain C^* should not be considered as fully MMR-proficient due to MMR saturation. Therefore, its mutation rate cannot be used to measure the apparent efficiencies of

proofreading and MMR under wild-type physiological conditions, i.e. as the numerator in a ratio

μ_{C^*}/μ_{LC^*} to estimate the probability that an error escapes proofreading, or as the denominator in a ratio

μ_{wt}/μ_{C^*} to estimate the probability that an error escapes MMR correction. Instead, these repair system

escape probabilities should be estimated from the ratios μ_{MMR-}/μ_{LC^*} and μ_{wt}/μ_{MMR-} shown in **Figure 8**

(where MMR- is either MMR-^{168} or ΔS^{3610}) for each replication-oriented triplet and mutation type

(ts, tv, ins, del). The rates of errors leading to mutation before and after correction by the combined

action of proofreading and MMR (mutation rates in *LC** and wild-type, respectively) are also

shown.

Figure 8. Apparent efficiency of proofreading and MMR across replication-stranded triplets
and mutation types. Estimated proofreading and MMR escape probabilities are shown (middle
plots) along with estimated mutation rates in the absence or presence of both correction systems
(top and bottom plots). Replication-stranded triplets are ordered in decreasing order of non-stranded
empirical substitution rates and then focal pyrimidine on the leading and lagging strands of
replication. Bold and thin vertical bars represent 50% and 95% credibility intervals, respectively.

Proofreading and MMR escape probabilities (μ_{MMR-}/μ_{LC*} and μ_{wt}/μ_{MMR-}) are correlated with triplet

substitution rates (**Figure S16**). In **Figure 8**, the triplets are ordered according to wild-type

substitution rates (as in **Figure 5A**), which highlights general trends: proofreading escape tends to

be higher on triplets where the initial polymerase selectivity is the lowest (**Figure S16**, r=0.32 *p*-

value=0.0097); MMR escape tends to be lower on triplets with high wild-type mutation rate

(**Figure S16**, r=-0.45, *p*-value=0.00018). Note that these two correlations are of opposite sign to the

artifactual correlations that would be generated by noise in the estimates of $\mu_{LC*}[i]$ and $\mu_{wt}[i]$, which

also enter in the ratios $\mu_{MMR-}[i]/\mu_{LC*}[i]$ and $\mu_{wt}[i]/\mu_{MMR-}[i]$.

The positive correlation between proofreading escape and mutation rate in *LC** is consistent with

the observation that proofreading increases the bias of polymerase errors. Statistical uncertainty

about the probability of proofreading escape makes it difficult to draw conclusions for specific

replication-stranded triplets, and few triplets show clear strand asymmetry. It is interesting to note,

however, that this positive link is not seen within pairs of triplets that differ only by the strand. In

particular, GTG and ATG are more affected by substitutions in *LC** when the focal pyrimidine (T)

is on the lagging strand, but no difference in proofreading escape probability between strands is

detected. For these triplets, the initial strand asymmetry may originate from incorporation errors

made by the DnaE polymerase during synthesis of the lagging strand that are efficiently proofread

in-trans by PolC. Conversely, errors affecting GCT and GCG appear to be less corrected by

proofreading when the C-site is on the leading strand, but the substitution rates in *LC** are similar

between strands. In these cases of C-site substitutions on the leading strand, the apparent probability

of proofreading escape most likely overestimates the escape of initial polymerase selectivity errors

because of the likely contribution of deamination damage to the mutation profile of the

proofreading-proficient MMR-deficient strains.

Deamination of cytosine to uracil in the lagging strand template (*i.e.* the leading strand) due to

exposure of single-stranded DNA in the context of the replication fork has long been suspected to

be the reason for the near-universal asymmetry between replication strands in prokaryotes (Frank

and Lobry 1999) and indeed has been shown to be a substantial source of $C \rightarrow T$ transitions in wild-

type (Bhagwat et al. 2016). Strikingly, proofreading-proficient MMR-deficient strains exhibit a

similarly oriented strong strand asymmetry for substitutions at C sites (Lee et al. 2012; Sung et al.

2015). It has been hypothesized that cytosine deamination may also cause this asymmetry (Foster et

al. 2018) which is consistent with other studies reporting a role for the *B. subtilis* MMR in

counteracting the effects of base deamination (López-Olmos et al. 2012; Patlan-Vazquez et al.

2022). In the absence of proofreading, the high rate of polymerase misincorporation on both strands

would mask the deamination-induced asymmetry.

The hypothesis that proofreading efficiency is positively linked to initial polymerase selectivity was

proposed above based on the analysis of substitution rates by triplet context without distinguishing

between transversions and transitions. Transversions account for only 4% of the substitutions in

LC*. The lower apparent proofreading escape probability of errors leading to transversions than to

transitions (**Figure 8**) is thus consistent with the above hypothesis. However, the difference

between the proofreading escape probabilities of the two types of errors, which cannot be precisely

estimated, seems to be smaller than expected. A possible explanation would that the apparent

proofreading escape probability overestimates proofreading escape for initial polymerase selectivity

errors that lead to transversions. This would occur if a substantial fraction of the rare transversions

in proofreading-proficient in MMR-deficient strains did not originate from polymerase errors and

thus was simply not subjected to proofreading.

Regarding MMR escape, which is not the primary focus of this study, its strong negative correlation

with proofreading escape (**Figure S17**, r=-0.63, *p*-value=2.6×10^{-8}) could reflect an evolution of

MMR towards correcting the most common DNA replication errors. Indeed, this would echo the

proposed role of differential MMR efficiency in balancing DNA replication fidelity between the

two strands in eukaryotes (Lujan et al. 2012; Andrianova et al. 2017; Zhou et al. 2021).

However, the negative correlation between the apparent probabilities of MMR and proofreading

escape could also reflect an overestimation of MMR escape for the sites least affected by DNA

replication errors. Indeed, the contribution of different sources of spontaneous mutations in wild-

type remains difficult to characterize (Schroeder et al. 2018), and thus there is a lack of knowledge

about the fraction of the wild-type mutation profile that is not subject to MMR correction. Among

the studies addressing this question, it has been shown that inactivation of oxidative damage repair

pathways increases the mutation rate in *E. coli* under optimal growth conditions without external

stress, suggesting that damage that escapes correction by oxidative damage repair pathways may

also contribute to the wild-type mutation profile (Foster et al. 2015). Since oxidative damage tends

to cause transversions, this hypothesis would be consistent with the higher proportion of

transversions in the wild-type than in any of our hypermutator strains. If a substantial fraction of the

wild-type substitution profile μ_{wt} originates from sources not subjected to MMR correction,

variations not directly related to MMR efficiency may enter the ratio μ_{wt}/μ_{MMR-} used to estimate the

MMR escape probability (**Figure 8**), both via its numerator μ_{wt} (*e.g.*, variations in the amount of

substitutions originating from these other sources) and via its denominator μ_{MMR-} (*e.g.*, variations in

the amount of polymerase errors remaining after proofreading). Indeed, the very strong negative

correlation across triplets between μ_{wt}/μ_{MMR-} and μ_{MMR-} (**Figure S16**, r=-0.94, *p*-value=2.3e^{-31})

suggests that variations in μ_{MMR-} may be largely responsible for variations in the apparent MMR

escape probability.. Combined with the unsurprising and clear positive correlation between the

apparent proofreading escape probability (μ_{MMR-}/μ_{LC*}) and μ_{MMR-} (**Figure S16**, r=0.71, *p*-

value=7.3e^{-11}), the substitutions originating from sources not subjected to MMR correction may thus

contribute to the observed negative correlation between the apparent probabilities of MMR and

proofreading escape.

Conclusion. Saturation of the MMR in the absence of proofreading, greater dispersion of
substitution rates in the presence than in the absence of proofreading, and the existence of strand
biases that become apparent only in the presence of proofreading appear to be traits shared between
B. subtilis and *E. coli*. Given the considerable divergence between these two organisms in terms of
phylogenetic distance and molecular organization of DNA polymerase and MMR, these traits are
likely to be common to many other organisms. The characterization of the overdispersion of
mutation rates in MMR-deficient proofreading-proficient strains compared to other strains led us to
conclude that proofreading skews DNA polymerase error rates. This could be interpreted as an
inherent drawback of the proofreading principle, which relies on the DNA polymerase to detect its
own errors. The role of judge and party leads to the same biases in initial nucleotide selectivity and
error correction efficiency.

This study also examines the consequences of aggregating mutations in counts and, more generally,
recognizes the difficulty of interpreting the apparent efficiencies of repair systems. First, unaware
aggregation of subclasses of mutations resulting from different molecular pathways can create
almost any pattern of apparent interactions in the effects of disabling different repair systems.
Second, interpretation of the effect of disabling one repair system is generally complex because of
uncertainty about the contribution of damage or errors subject to correction by that system to the
mutations observed in its presence. It is thus difficult to interpret the appearance of strand
asymmetry upon proofreading activation, which may be caused by a post-proofreading mutation
process, such as deamination, rather than a proofreading bias. Similarly, the "flattening" of
mutation rates upon activation of the MMR is difficult to interpret, as it could result from a better
efficiency of the MMR in correcting the most common errors, but also from the contribution of
multiple sources to the wild-type mutation profiles. Indeed, a substantial contribution of multiple
sources seems consistent with the drift-barrier hypothesis (Lynch et al. 2016): in the wild-type, the
rates of mutations resulting from different molecular pathways may have been pushed below the
same point where they do not encounter significant counter-selection.

The construction of strains with inducible hypermutator phenotypes addressed the stability issue
that affected the reproducibility of some studies on *E. coli* proofreading-deficient strains discussed
in (Niccum et al. 2018). It was also motivated by the interest in tools for synthetic biology
applications. Regarding the future use of our inducible systems to accelerate evolution in the
laboratory, a proofreading-deficient strain would yield a less skewed mutation spectrum than an
MMR-deficient strain. If necessary, extreme mutation rates can be achieved by disabling both repair

systems simultaneously, but strong counter-selection against mutational load makes such induction

feasible only for a short period of time.

METHODS

Media and bacterial strains. *E. coli* DH5α was used for plasmid construction and transformation

using standard techniques (Sambrook et al. 1989). The *B. subtilis* strains used in this study were

derived from our Master Strain (MS), a prophage-free and *trp*[+] derivative of *B. subtilis* 168 (Dervyn

et al. 2023), denoted here R[168]. Lysogeny broth (LB) was used to grow *E. coli* and *B. subtilis*.

Transformation of *B. subtilis* cells was performed using the protocol of (Konkol et al. 2013). When

relevant, the media were supplemented with the following antibiotics: ampicillin 100 µg.mL[-1] for *E.*

coli and spectinomycin 100 µg.mL[-1] or kanamycin 5 µg.mL[-1] for *B. subtilis*.

Construction of hypermutator strains. The Δ*S* and Δ*L* mutant strains were generated by

transforming the PCR-amplified *kan mutL* and *mutS kan* sequences using the P1-P2 primer pair

together with genomic DNA from Δ*mutS*::*kan* and Δ*mutL*::*kan* mutant strains, respectively. These

strains were obtained from the previously published *B. subtilis* single-gene deletion library (Koo et

al. 2017).

To construct the *L** strain, the first and second halves of the *mutL* gene were PCR-amplified using

primer pairs P5-P8 and P6-P7, respectively (**Figure S1**), where P7 and P8 both carry the desired

point mutation (as indicated in **Table S7**). The two fragments were then assembled by PCR,

resulting in the *mutL*(N34H) allele. The backbone of the pDR111 plasmid (kind gift of D. Rüdner)

containing the isopropyl-β-D-1-thiogalactopyranoside (IPTG) inducible P*hyperspank* promoter (denoted

here P*hs*) and the *spec* gene (conferring resistance to spectinomycin), was PCR-amplified using P3

and P4. The 5' extensions of P5 and P6 then allowed the assembly of the *mutL*(N34H) allele with

the PCR-amplified pDR111 using the HiFi DNA assembly protocol (New England Biolabs, USA).

This resulted in cloning the *mutL*(N34H) allele under the control of P*hs* into a *B. subtilis amyE*-

integrative plasmid (**Figure S1**).

Similarly, to construct the *C** strain, the *polC* allele found in *B. subtilis mut*-1 (Bazill and Gross

1973), characterized by the G430E and S621N mutations (Sanjanwala and Ganesan 1991), was

PCR-amplified using the primer pair P11-P12 and assembled to the PCR-amplified pDR111 (using

P3 and P4) using the HiFi DNA assembly protocol (New England Biolabs, USA). This resulted in

cloning the *polC mut*-1 allele under the control of P*hs* into a *B. subtilis amyE*-integrative plasmid

(**Figure S2**).

For the construction of the *LC** strain, a *mutL** *polC** synthetic operon was generated by

assembling the *mutL*(N34H) allele PCR-amplified from strain *L** using P5 and P11, and the *polC*

mut-1 allele PCR-amplified from *C** using P12 and P10, to the PCR-amplified pDR111 (using P3

and P4) using the HiFi DNA assembly protocol (New England Biolabs, USA). This resulted in the

cloning of the *mutL* polC** synthetic operon under the control of P$_{hs}$ into a *B. subtilis amyE-*

integrative plasmid (**Figure S2**).

Plasmids were transformed into the *B. subtilis amyE* locus by double recombination events. All

strains were verified by sequencing, and transcriptomics experiments were performed to compare

global gene expression. The RNA-seq reads and detailed protocols and results have been deposited

in GEO.

Fluctuation assays. For each strain to be tested, a single colony was grown in 1 mL LB at 37°C for

90 minutes. This preculture was serially diluted in fresh LB to start cultures with a small number of

cells (N_0). Cells were then grown for 7.5 h to reach saturation. When induction with IPTG was

tested, LB medium with the desired concentration of IPTG was prepared from an IPTG stock

concentration of 1 mM just before use. When the culture volume was 1 mL, the cultures were

centrifuged before plating to retain the cells, and 750 µL of supernatant was removed. The

remaining 250 µL were gently vortexed before plating onto LB supplemented with rifampicin (10

µg.mL^{-1}). For each assay, a number of cultures (8 for the R^{168} assay, 4 for the ΔS and ΔL assays, and

3 for all other assays) were not plated on LB medium supplemented with rifampicin, but were

serially diluted and plated on LB agar to determine the final number of cells (N_t). Fluctuation assays

performed on the same day were considered to have the same distribution of final cell numbers. All

other cultures were plated on LB agar with rifampicin to determine the number of Rif-resistant

colony-forming units (CFUs). All plates were incubated at 37°C and scored for CFUs after 24 h of

growth.

The maximum likelihood estimator (MLE) of the number of mutations per assay (m) and the

confidence interval, were calculated under the Luria-Delbrück model taking into account the

variation in the final number of cells (Zheng 2016), using the newton.B0 with default parameters

and confint.B0 functions of the R package "Rsalvador" v1.7 (Zheng 2017). For the calculation of

confidence intervals, the initial guess for the parameter m was taken as the m given by the

"newton.B0" function. The mutation probability was assumed to be constant over the cell cycle, so

that the mutation rate per base per generation is the mutation rate per base per cell division (Foster

2006). The final number of cells, N_t, is the result of N_t - N_0 cell divisions, *i.e.* ~N_t divisions. The rate

of Rif^R emergence was therefore calculated as μ_{Rif}= m/N_t.

Mutation-accumulation experiments and sequencing. One isolated colony was collected each

day (24 h at 37°C), suspended in culture medium + 20% glycerol, and diluted by 2×10^5, a factor

that allows distinguishable colonies, before plating on LB agar (+ 100 µM IPTG for *L**, *C** and

LC* strains) for the next MA-step. Counting of the colonies present on the agar plate provided an

estimate of the number of bacteria initially present in the diluted colony and thus the number of

generations per MA-step. Four parallel MA-lines were propagated per strain (21 consecutive MA-

steps for Δ*S*, Δ*L*, *L** and *C**, 11 for *LC**).

For sequencing at intermediate and end points of the MA-lines, 5 to 50% of the picked colony was

cultured in LB medium to collect cells. DNA was extracted using the GenEluteTM Bacterial

Genomic DNA Kit (Sigma-Aldrich) according to the supplied protocol. DNA samples

corresponding to an intermediate time-point in the four parallel MA-lines for the same strain were

pooled in equimolar proportions. Individual and pooled DNA samples were sequenced (150 bp

paired-end reads) on an Illumina platform (NovaSeq 6000) to an average depth of ~300. Reads are

available in NCBI SRA.

Detection of mutations. The reads were aligned to the reference sequence of the *B. subtilis* 168

genome (GenBank: AL009126.3) using BWA-MEM v0.7.17 (Li and Durbin 2009), after quality

control and trimming using Sickle v1.33 (command "sickle pe" with options "-t sanger -x -q 20 -l

20") (Joshi and Fass 2011). Properly paired reads, selected using "samtools view -f 3" (samtools

v1.14, Li 2011), were locally realigned around indels using ABRA2 v2.24 (Mose et al. 2019). The

number of occurrences of each nucleotide (base read quality ≥35) and indel at each position of the

reference in confidently mapped reads (alignment quality ≥50) was counted using "samtools

mpileup" with the options "-aa -d 5000 -q 50 -Q 35 -x -B". These counts were analyzed in R.

For each position, the effective sequencing depth (DPeff) was calculated as the total number of

informative reads. For the computation of the mutation rates, a reference subset of positions

common to all samples was determined. This reference consisted of positions that were well

covered on both strands (DPeff≥100 and ≥10% of the reads on the less represented strand) in all

samples. Most of the regions with low coverage corresponded to the regions deleted in the

construction of the MS / R[168] strain (Dervyn et al. 2023), which lacks 233.4 kb of chromosome

relative to AL009126.3, and to the multicopy structural RNAs. Over-covered regions were also

eliminated from this reference subset of positions and consisted of: the region of gene *upp* and

downstream (positions 3,788,426 to 3,789,124), which was repeated due to pop-ins and pop-outs at

this locus during the construction of the R[168] strain; the region from position 2,432,478 to 2,433,315,

over-covered in *polC** samples; the regions of the genes *polC* (1,727,133 to 1,731,446) and *mutL*

(1,778,337 to 1,780,539) duplicated by insertion of the mutant alleles *polC** and *mutL**. This

resulted in a reference subset of 3,794,734 positions (out of a total of 4,215,606 bp in AL009126.3),

which served as our reference chromosome for the mutation rate calculations.

The distribution of the proportion of non-reference reads in the different samples was examined

graphically to establish relevant cut-offs for the identification of mutations. A mutation was

identified at the endpoint of an MA-line if a variant accounted for ≥75% of the DPeff at a position,

with ≥10% of the non-reference reads on the less represented strand. If intermediate time-points

were available for this MA-line, the mutation was traced back to the first time-point where it

occurred at frequency ≥5% in the corresponding pooled sequence sample. Due to the detection,

during graphical examination, of contamination from other samples, we lowered the cut-off from

75% to 60% for the identification of mutations in the third MA-line of ΔS and from 5% to 2% for

the analysis of the pool corresponding to the intermediate time-point for *L** strain MA-lines.

Mutations found in all samples or in the four MA-lines of a same strain were interpreted as fixed

before the mutation-accumulation experiment and were discarded for the calculation of mutation

rates.

Detection of mutations in inducible synthetic circuits. The reads were also aligned to the

reference sequences of the inserted regions represented in **Figure S2**. For the mutant alleles *mutL**

and *polC**, reads from the native alleles (*mutL* and *polC*) were also mapped to the insert, and

variant calling *per se* cannot distinguish between a mutation in the native and mutant allele.

However, at positions where bases differed from the reference on these genes in individual samples,

the proportion of these alternative bases was bimodally distributed, with two peaks, located at

approximately at 40% and 60% of the reads **(Figure S10**). Since the characteristic point mutations

of both *polC** and *mutL** accounted for more than 50% of the reads (between 52% and 71% and

between 65% and 75%,, respectively) at their respective positions, we predicted that the mutations

for which the majority of the reads differed from the reference would be on the mutant allele, while

the others would be on the native allele. This prediction is consistent with the chromosomal position

of the *amyE* locus where the mutant alleles are inserted, *i.e.* closer to the origin of replication of the

chromosome than either native allele, and thus expected to be more abundant in the sample due to

ongoing replication. To verify this, we used specific primers to amplify and sequence either the

native or the inserted allele. Of the 5 PCR-verified mutations, all were found on the predicted allele

(**Table S5**).

Chromosome partitioning to assess the effect of transcription and replication on mutation

rate. To assess the effect of transcription, the "gene" features of the GenBank annotation served to

define the dichotomy between "template" and "nontemplate" strand as well as between "coding"

and "noncoding". Since the "noncoding" represents only ~10% of the genome and includes

transcribed untranslated regions (UTRs) we also sought to assess the impact of transcription with

more statistical power and precision than allowed by the GenBank annotation. For this purpose, two

categories of regions of approximately equal size were defined based on the transcribed regions

identified in 269 samples of a wild-type strain representative of a wide variety of growth conditions

(Nicolas et al. 2012). These two categories reflected the amount of transcripts in LB as measured in

9 samples corresponding to growth in liquid LB (triplicate samples for exponential, transition and

stationary phases) and 2 samples corresponding to 16 hours of growth on LB agar (non-onfluent

colonies). The "high" transcription level regions were those that were in the top 30% in at least one

of these 11 samples while the "low" transcription level regions were those that were never in the top

30%. All overlapping regions (*i.e.* both strands were transcribed) were eliminated, as well as all

regions shorter than 100 bp. This resulted in a set of 3,622 non-overlapping transcription-oriented

regions covering 84.9% of the reference genome (43.4% for "high", 41.4% for "low").

To assess the effect of DNA replication asymmetry, the leading and lagging strands were defined

based on the origin of replication (position 1) and its most central terminus (position 2,018,289)

(Wake 1997). To assess the effect of DNA replication timing, the genome was divided into a "first

half" corresponding to the 2 Mbp of the chromosome centered on the origin of replication (position

1) and a "second half".

Mutation rate estimations and comparisons. To include the list of mutations in R^{3610} and ΔS^{3610}

from (Sung et al. 2015) and (Sung et al. 2016) in our analysis, the positions on the *B. subtilis* NCIB

3610 genome (GenBank: CM000488.1) were transferred to the *B. subtilis* 168 genome by mapping

the 41 bp long sequence centered on each mutation site. Retaining only exact and unique matches,

more than 99% of these mutations were transferred (**Table S3**), with perfect collinearity between

the positions of the mutations on both reference genomes.

Maximum-likelihood estimates of mutation rates were obtained as $\mu=m/(T\times G)$, where *m* is the total

number of mutations of a given type in a given genotype and genomic context (nucleotide at focal

position and adjacent nucleotides, orientation with respect to replication, transcription, …), T is the

total number of occurrences of the genomic context in the reference sequence, and G is the number

of generations considered in MA-lines. Confidence intervals for these point estimates were

calculated using the exact method for Poisson counts implemented in R package "epitools" v0.5-

10.1, with *m* as count and T×G as time-person at risk.

To assess whether a factor affects the substitution rates, we used Generalised Linear Models

(GLMs) for Poisson distributed count data with log-link, combined with an Analysis of Variance

(ANOVA) (R package "stats" v3.6.3) to compare the fit of a GLM including and a GLM excluding

the factor of interest. This statistical comparison was performed separately for each genotype.

Markov chain Monte Carlo methods implemented in JAGS (Plummer 2003), accessed via the R

package "rjags", were used for Bayesian estimation via posterior sampling, in particular for the

estimation of mutation rates of replication-stranded triplets and the MMR saturation parameter θ.

Models and algorithm settings are described in detail in **Supplementary Methods and Results 1.4**.

Mathematical modelling of mutation rates

Assumptions and Bayesian estimation procedure for the model with saturation of the MMR are

presented in **Supplementary Methods and Results 1.5**. Algebraic analysis of the general model

with two subclasses of errors and two repair pathways is presented in **Supplementary Methods**

and Results 1.6.

DATA ACCESS

The RNA and DNA sequencing data generated in this study have been submitted to the NCBI Gene Expression Omnibus database under accession number GSE239804 and to the NCBI Sequence Read Archive under BioProject accession number PRJNA995423, respectively.

COMPETING INTEREST STATEMENT

The authors declare no competing interests.

ACKNOWLEDGEMENTS

We express our gratitude to Guillaume Achaz, Elena Bidnenko, Clément Nizak and Paulo Tavares for their advice during the course of this work. We also thank the INRAE MIGALE bioinformatics facility (https://doi.org/10.15454/1.5572390655343293E12) for providing computational resources and Marina Elez for helpful comments on the manuscript.

AUTHOR CONTRIBUTIONS

IT, MJ, and PN conceived the project and designed the experimental plan. ED and IT performed the experiments. IT processed the raw data with the help of CG. GKKK, IT, and PN conducted the statistical and mathematical analyses. IT, MJ, and PN interpreted the results and wrote the manuscript with contributions from all authors.

FUNDING

This work was supported by the French National Research Agency (ANR-18-CE43-0002). The IT PhD fellowship was partially funded by the MathNum division of INRAE.

REFERENCES

Andrianova MA, Bazykin GA, Nikolaev SI, Seplyarskiy VB. 2017. Human mismatch repair system balances mutation rates between strands by removing more mismatches from the lagging strand. *Genome Res* **27**: 1336-1343.

Badran AH, Liu DR. 2015. Development of potent in vivo mutagenesis plasmids with broad mutational spectra. *Nat Commun* **6**: 8425.

Bazill GW, Gross JD. 1973. Mutagenic DNA polymerase in *B. subtilis*. *Nat New Biol* **243**: 241-243.

Bhagwat AS, Hao W, Townes JP, Lee H, Tang H, Foster PL. 2016. Strand-biased cytosine deamination at the replication fork causes cytosine to thymine mutations in *Escherichia coli*. *Proc Natl Acad Sci U S A* **113**: 2176-2181.

Bolz NJ, Lenhart JS, Weindorf SC, Simmons LA. 2012. Residues in the N-terminal domain of MutL required for mismatch repair in *Bacillus subtilis*. *J Bacteriol* **194**: 5361-5367.

Bruck I, Goodman MF, O'Donnell M. 2003. The essential C family DnaE polymerase is error-prone and efficient at lesion bypass. *J Biol Chem* **278**: 44361-44368.

Couce A, Caudwell LV, Feinauer C, Hindre T, Feugeas JP, Weigt M, Lenski RE, Schneider D, Tenaillon O. 2017. Mutator genomes decay, despite sustained fitness gains, in a long-term experiment with bacteria. *Proc Natl Acad Sci U S A* **114**: E9026-E9035.

Deatherage DE, Leon D, Rodriguez AE, Omar SK, Barrick JE. 2018. Directed evolution of *Escherichia coli* with lower-than-natural plasmid mutation rates. *Nucleic Acids Res* **46**: 9236-9250.

Dervyn E, Planson AG, Tanaka K, Chubukov V, Guerin C, Derozier S, Lecointe F, Sauer U, Yoshida KI, Nicolas P et al. 2023. Greedy reduction of *Bacillus subtilis* genome yields emergent phenotypes of high resistance to a DNA damaging agent and low evolvability. *Nucleic Acids Res* doi:10.1093/nar/gkad145.

Dervyn E, Suski C, Daniel R, Bruand C, Chapuis J, Errington J, Janniere L, Ehrlich SD. 2001. Two essential DNA polymerases at the bacterial replication fork. *Science* **294**: 1716-1719.

Dulanto Chiang A, Patil PP, Beka L, Youn JH, Launay A, Bonomo RA, Khil PP, Dekker JP. 2022. Hypermutator strains of *Pseudomonas aeruginosa* reveal novel pathways of resistance to combinations of cephalosporin antibiotics and beta-lactamase inhibitors. *PLoS Biol* **20**: e3001878.

Eyre-Walker A, Keightley PD. 2007. The distribution of fitness effects of new mutations. *Nature Reviews Genetics* **8**: 610-618.

Foster PL. 2006. Methods for determining spontaneous mutation rates. *Methods Enzymol* **409**: 195-213.

Foster PL, Lee H, Popodi E, Townes JP, Tang H. 2015. Determinants of spontaneous mutation in the bacterium *Escherichia coli* as revealed by whole-genome sequencing. *Proc Natl Acad Sci U S A* **112**: E5990-5999.

Foster PL, Niccum BA, Popodi E, Townes JP, Lee H, MohammedIsmail W, Tang H. 2018. Determinants of Base-Pair Substitution Patterns Revealed by Whole-Genome Sequencing of DNA Mismatch Repair Defective *Escherichia coli*. *Genetics* **209**: 1029-1042.

Frank AC, Lobry JR. 1999. Asymmetric substitution patterns: a review of possible underlying mutational or selective mechanisms. *Gene* **238**: 65-77.

Fujii S, Akiyama M, Aoki K, Sugaya Y, Higuchi K, Hiraoka M, Miki Y, Saitoh N, Yoshiyama K, Ihara K et al. 1999. DNA replication errors produced by the replicative apparatus of *Escherichia coli*. *J Mol Biol* **289**: 835-850.

Ganai RA, Johansson E. 2016. DNA Replication-A Matter of Fidelity. *Mol Cell* **62**: 745-755.

Guiziou S, Sauveplane V, Chang HJ, Clerte C, Declerck N, Jules M, Bonnet J. 2016. A part toolbox to tune genetic expression in *Bacillus subtilis*. *Nucleic Acids Res* **44**: 7495-7508.

Haradhvala NJ, Kim J, Maruvka YE, Polak P, Rosebrock D, Livitz D, Hess JM, Leshchiner I, Kamburov A, Mouw KW et al. 2018. Distinct mutational signatures characterize concurrent loss of polymerase proofreading and mismatch repair. *Nat Commun* **9**: 1746.

Joshi NA, Fass JN. 2011. Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files. https://github.com/najoshi/sickle.

Kadyrov FA, Dzantiev L, Constantin N, Modrich P. 2006. Endonucleolytic function of MutLalpha in human mismatch repair. *Cell* **126**: 297-308.

Klocko AD, Schroeder JW, Walsh BW, Lenhart JS, Evans ML, Simmons LA. 2011. Mismatch repair causes the dynamic release of an essential DNA polymerase from the replication fork. *Mol Microbiol* **82**: 648-663.

Konkol MA, Blair KM, Kearns DB. 2013. Plasmid-encoded ComI inhibits competence in the ancestral 3610 strain of *Bacillus subtilis*. *J Bacteriol* **195**: 4085-4093.

Koo BM, Kritikos G, Farelli JD, Todor H, Tong K, Kimsey H, Wapinski I, Galardini M, Cabal A, Peters JM et al. 2017. Construction and Analysis of Two Genome-Scale Deletion Libraries for *Bacillus subtilis*. *Cell Syst* **4**: 291-305.

Lee H, Popodi E, Tang H, Foster PL. 2012. Rate and molecular spectrum of spontaneous mutations in the bacterium *Escherichia coli* as determined by whole-genome sequencing. *Proc Natl Acad Sci U S A* **109**: E2774-2783.

Lenhart JS, Schroeder JW, Walsh BW, Simmons LA. 2012. DNA repair and genome maintenance in *Bacillus subtilis*. *Microbiol Mol Biol Rev* **76**: 530-564.

Li H. 2011. A statistical framework for SP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics* **27**: 2987-2993.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**: 1754-1760.

Long H, Miller SF, Williams E, Lynch M. 2018. Specificity of the DNA Mismatch Repair System (MMR) and Mutagenesis Bias in Bacteria. *Mol Biol Evol* **35**: 2414-2421.

López-Olmos K, Hernández MP, Contreras-Garduño JA, Robleto EA, Setlow P, Yasbin RE, Pedraza-Reyes M. 2012. Roles of endonuclease V, uracil-DNA glycosylase, and mismatch repair in *Bacillus subtilis* DNA base-deamination-induced mutagenesis. *Journal of Bacteriology* **194**: 243-252.

Lujan SA, Williams JS, Pursell ZF, Abdulovic-Cui AA, Clark AB, Nick McElhinny SA, Kunkel TA. 2012. Mismatch repair balances leading and lagging strand DNA replication fidelity. *PLoS Genet* **8**: e1003016.

Lynch M, Ackerman MS, Gout JF, Long H, Sung W, Thomas WK, Foster PL. 2016. Genetic drift, selection and the evolution of the mutation rate. *Nat Rev Genet* **17**: 704-714.

Mahilkar A, Raj N, Kemkar S, Saini S. 2022. Selection in a growing colony biases results of mutation accumulation experiments. *Sci Rep* **12**: 15470.

Maki H. 2002. Origins of spontaneous mutations: specificity and directionality of base-substitution, frameshift, and sequence-substitution mutageneses. *Annu Rev Genet* **36**: 279-303.

Molina RS, Rix G, Mengiste AA, Alvarez B, Seo D, Chen H, Hurtado J, Zhang Q, Donato Garcia-Garcia J, Heins ZJ et al. 2022. In vivo hypermutation and continuous evolution. *Nat Rev Methods Primers* **2**.

Mose LE, Perou CM, Parker JS. 2019. Improved indel detection in DNA and RNA via realignment with ABRA2. *Bioinformatics* **35**: 2966-2973.

Nemenman I, Shafee F, Bialek W. 2001. Entropy and inference, revisited. In *Advances in Neural Information Processing Systems*, Vol 14 (ed. TG Dietterich, et al.). The MIT Press.

Niccum BA, Lee H, MohammedIsmail W, Tang H, Foster PL. 2018. The Spectrum of Replication Errors in the Absence of Error Correction Assayed Across the Whole Genome of *Escherichia coli*. *Genetics* **209**: 1043-1054.

Nicolas P, Mäder U, Dervyn E, Rochat T, Leduc A, Pigeonneau N, Bidnenko E, Marchadier E, Hoebeke M, Aymerich S et al. 2012. Condition-dependent transcriptome reveals high-level regulatory architecture in *Bacillus subtilis*. *Science* **335**: 1103-1106.

Oliver A, Mena A. 2010. Bacterial hypermutation in cystic fibrosis, not only for antibiotic resistance. *Clin Microbiol Infect* **16**: 798-808.

Paschalis V, Le Chatelier E, Green M, Nouri H, Kepes F, Soultanas P, Janniere L. 2017. Interactions of the Bacillus subtilis DnaE polymerase with replisomal proteins modulate its activity and fidelity. *Open Biol* **7**.

Patlan-Vazquez AG, Ayala-Garcia VM, Vallin C, Cortes J, Vasquez-Morales SG, Robleto EA, Nudler E, Pedraza-Reyes M. 2022. Dynamics of Mismatch and Alternative Excision-Dependent Repair in Replicating *Bacillus subtilis* DNA Examined Under Conditions of Neutral Selection. *Front Microbiol* **13**: 866089.

Perfeito L, Sousa A, Bataillon T, Gordo I. 2014. Rates of fitness decline and rebound suggest pervasive epistasis. *Evolution* **68**: 150-162.

Pillon MC, Lorenowicz JJ, Uckelmann M, Klocko AD, Mitchell RR, Chung YS, Modrich P, Walker GC, Simmons LA, Friedhoff P et al. 2010. Structure of the endonuclease domain of MutL: unlicensed to cut. *Mol Cell* **39**: 145-151.

Plummer M. 2003. JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling. In Proceedings of the 3rd International Workshop on Distributed Statistical Computing (DSC 2003), March 20-22, Vienna, Austria. ISSN 1609-395X., volume 124, page 125.

Rudenko O, Engelstadter J, Barnes AC. 2020. Evolutionary epidemiology of *Streptococcus iniae*: Linking mutation rate dynamics with adaptation to novel immunological landscapes. *Infect Genet Evol* **85**: 104435.

Sambrook J, Fritsch EF and Maniatis T. 1989. Molecular Cloning: a Laboratory Manual. Cold Spring Harbor ed. Cold Spring Harbor Laboratory, NY.

Sanjanwala B, Ganesan AT. 1991. Genetic structure and domains of DNA polymerase III of *Bacillus subtilis*. *Mol Gen Genet* **226**: 467-472.

Schaaper RM. 1988. Mechanisms of mutagenesis in the *Escherichia coli* mutator mutD5: role of
DNA mismatch repair. *Proc Natl Acad Sci U S A* **85**: 8126-8130.

Schaaper RM, Radman M. 1989. The extreme mutator effect of *Escherichia coli mutD5* results
from saturation of mismatch repair by excessive DNA replication errors. *EMBO J* **8**: 3511-
3516.

Schroeder JW, Hirst WG, Szewczyk GA, Simmons LA. 2016. The Effect of Local Sequence
Context on Mutational Bias of Genes Encoded on the Leading and Lagging Strands. *Curr
Biol* **26**: 692-697.

Schroeder JW, Yeesin P, Simmons LA, Wang JD. 2018. Sources of spontaneous mutagenesis in
bacteria. *Crit Rev Biochem Mol Biol* **53**: 29-48.

Sherer NA, Kuhlman TE. 2020. Escherichia coli with a Tunable Point Mutation Rate for Evolution
Experiments. *G3 (Bethesda)* **10**: 2671-2681.

Singh T, Hyun M, Siegowski P. 2017. Evolution of mutation rates in hypermutable populations of
Escherichia coli propagated at very small effective population size. *Biol Lett* **13**.

Sniegowski PD, Gerrish PJ, Lenski RE. 1997. Evolution of high mutation rates in experimental
populations of *E. coli*. *Nature* **387**: 703-705.

Sung W, Ackerman MS, Dillon MM, Platt TG, Fuqua C, Cooper VS, Lynch M. 2016. Evolution of
the Insertion-Deletion Mutation Rate Across the Tree of Life. *G3 (Bethesda)* **6**: 2583-2591.

Sung W, Ackerman MS, Gout J-F, Miller SF, Williams E, Foster PL, Lynch M. 2015. Asymmetric
Context-Dependent Mutation Patterns Revealed through Mutation–Accumulation
Experiments. *Molecular Biology and Evolution* **32**: 1672-1683.

Swings T, Weytjens B, Schalck T, Bonte C, Verstraeten N, Michiels J, Marchal K. 2017. Network-
Based Identification of Adaptive Pathways in Evolved Ethanol-Tolerant Bacterial
Populations. *Mol Biol Evol* **34**: 2927-2943.

Taddei F, Radman M, Maynard-Smith J, Toupance B, Gouyon PH, Godelle B. 1997. Role of
mutator alleles in adaptive evolution. *Nature* **387**: 700-702.

Tate JG, Bamford S, Jubb HC, Sondka Z, Beare DM, Bindal N, Boutselakis H, Cole CG, Creatore
C, Dawson E et al. 2019. COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic
Acids Res* **47**: D941-D947.

Timinskas K, Balvociute M, Timinskas A, Venclovas C. 2014. Comprehensive analysis of DNA
polymerase III alpha subunits and their homologs in bacterial genomes. *Nucleic Acids Res*
42: 1393-1413.

Wahl LM, Agashe D. 2022. Selection bias in mutation accumulation. *Evolution* **76**: 528-540.

Wake RG. 1997. Replication fork arrest and termination of chromosome replication in *Bacillus
subtilis*. *FEMS Microbiol Lett* **153**: 247-254.

Zheng Q. 2016. A second look at the final number of cells in a fluctuation experiment. *J Theor Biol*
401: 54-63.

Zheng Q. 2017. rSalvador: An R Package for the Fluctuation Experiment. *G3 (Bethesda)* **7**: 3849-
3856.

Zhou ZX, Lujan SA, Burkholder AB, St Charles J, Dahl J, Farrell CE, Williams JS, Kunkel TA.
2021. How asymmetric DNA replication achieves symmetrical fidelity. *Nat Struct Mol Biol*
28: 1020-1028.