The ribosomal protein L22 binds the *MDM4* pre-mRNA

and promotes exon skipping

to activate p53 upon nucleolar stress

Jennifer Jansen[1], Katherine E. Bohnsack[2], Susanne Böhlken-Fascher[1], Markus T. Bohnsack[2], Matthias Dobbelstein[1,3,*]

[1]Institute of Molecular Oncology, Göttingen Center of Molecular Biosciences (GZMB), University Medical Center Göttingen, Justus-von-Liebig-Weg 11, 37077 Göttingen, Germany

[2]Department of Molecular Biology, University Medical Center Göttingen, Humboldtallee 23, 37073 Göttingen, Germany

[3]Lead contact

*Correspondence: mdobbel@uni-goettingen.de

Running title: Ribosomal protein L22 influences MDM4 splicing

Key words: Ribosomal proteins L22, L22L1, L5, L11, S13; nucleolus; RNA polymerase I; ZMAT3; UBAP2L; splicing; exon skipping; RNA stem-loop

ABSTRACT

The tumor suppressor p53, along with its antagonists MDM2 and MDM4, represents a central integrator of stress signaling. While DNA damage is the most widely explored trigger of a p53 response, stress can also arise from dysbalanced assembly of ribosomes in nucleoli. Deletions of the gene encoding the ribosomal protein L22 (RPL22; eL22) correlate with the presence of full-length *MDM4* mRNA in human cancer, but the mechanistic basis for this phenomenon was hitherto unknown. Here we show that L22, under conditions of ribosomal and nucleolar stress, promotes the skipping of exon 6 within the *MDM4* pre-mRNA. Upon L22 depletion, far more full-length MDM4 is maintained despite treatment with nucleolar stressors, leading to diminished p53 activity and enhanced proliferation. Mechanistically, L22 binds to specific RNA elements within intron 6 of *MDM4* that correspond to a stem-loop consensus, leading to the skipping of exon 6. This intronic RNA overlaps with the region responsible for splice regulation by *ZMAT3*. Targeted deletion of these intronic elements largely abolishes L22-mediated exon skipping and re-enables cell proliferation despite nucleolar stressors such as 5-fluorouracil. L22 also governs alternative splicing of the *RPL22L1* and *UBAP2L* mRNAs. Thus, L22 serves as a signaling intermediate that integrates different layers of gene expression. Defects in ribosome synthesis lead to specific alternative splicing, ultimately triggering p53-mediated transcription and arresting cell proliferation.

INTRODUCTION

p53 is the most frequently mutated tumor suppressor in human cancer (Kandoth et al., 2013; Levine and Oren, 2009; Olivier et al., 2010). When not affected by mutations, the transcription factor p53 is kept in check by its antagonist MDM2, which is itself encoded by a p53-responsive gene (Juven et al., 1993; Leach et al., 1993; Oliner et al., 2016; Wu et al., 1993). MDM2 binds, ubiquitinates and thus destabilizes p53, thereby providing negative feedback (Haupt et al., 1997; Honda et al., 1997; Kubbutat et al., 1997). This system serves to integrate stress signals such as the kinase cascade triggered by DNA damage (Vousden and Prives, 2009). The kinases ATM/ATR and CHK2/CHK1 phosphorylate both p53 and MDM2, thus decreasing MDM2-mediated ubiquitination of p53 and enhancing p53 levels and activity (Li and Kurokawa, 2015; Rizzotto et al., 2021).

Besides DNA damage, nucleolar stress (also known as ribosome assembly stress, triggering the impaired ribosome biogenesis checkpoint) represents an important trigger of p53, but the underlying mechanisms remain to be fully explored. Nucleolar stress occurs when ribosome assembly is impaired, e.g. by alterations in ribosomal RNA (rRNA) synthesis, or by the failure to synthesize or correctly assemble a specific ribosomal component. Under such circumstances, nucleolar proteins, including ribosomal proteins, accumulate in the nucleoplasm. This can induce a p53-response in multiple ways. For example, the 5S ribonucleoprotein complex (5S RNP), consisting of the 5S rRNA and the ribosomal proteins L5 (RPL5; uL18) and L11 (RPL11; uL5), associates with MDM2 and decreases p53 ubiquitination (Boulon et al., 2010; Sloan et al., 2013; Yang et al., 2018).

Besides MDM2, the MDM4 protein is an essential regulator of p53, and the deletion of *MDM4* is embryonic lethal unless accompanied by a deletion of p53 (Parant et al., 2001). MDM4 binds both p53 and MDM2, and it supports MDM2-mediated ubiquitination of p53 (Gu et al., 2002; Shvarts et al., 1996; Tanimura et al., 1999). Thus, MDM4 can also serve as an integrator of stress signals to trigger p53. One of the mechanisms that modulate MDM4 in response to stress consists in the skipping of exon 6 in the *MDM4* pre-mRNA. When alterations in splicing activity lead to elimination of this exon from the pre-mRNA, only a smaller and unstable form of MDM4 is synthesized, whereas inclusion of exon 6 allows the synthesis of full-length, stable and active MDM4. As a consequence, conditions that increase exon 6 skipping enhance p53 activity (Bardot and Toledo, 2017; Dewaele et al., 2016; Wu et al., 2021).

Massive gene expression analyses in tumor tissue revealed a positive correlation between the skipping of exon 6 in *MDM4* and the presence of two intact alleles encoding the ribosomal protein L22 as one of the most striking correlations (Ghandi et al., 2019). This suggests that L22 and MDM4 might affect each other's expression or activity. Interestingly, our earlier research revealed that L22 binds to a specific stem-loop motif in RNA, consisting of a G-C

base pair on top of the stem, and a U at the 3' end of the loop. This motif is found in viral RNAs, but its role in mammalian RNA is less well-defined (Dobbelstein and Shenk, 1995). Research in non-mammalian vertebrates suggested a role for L22 in splicing regulation (Zhang et al., 2017). Moreover, low L22 expression levels correlate with worse outcome of myelodysplastic syndrome in patients and a mouse model (Harris et al., 2023). Taken together, these observations raise the possibility that the *MDM4* mRNA precursor might contain L22 binding motifs, thus allowing L22 to influence MDM4 expression.

Here we show that L22 binds to three cognate elements within intron 6 of the *MDM4* pre-mRNA. L22 promotes skipping of exon 6 and thus prevents the synthesis of stable and active, full-length MDM4, thereby increasing the expression of p53 target genes and diminishing cell proliferation. Thus, L22 provides a link between three different aspects of gene expression, i) its extra-ribosomal pool is increased when ribosome biosynthesis is impaired, ii) it then regulates the splicing of *MDM4* and iii) it ultimately triggers the activation of p53 as a transcription factor of specific mRNAs. This signaling mechanism supports cells in responding to nucleolar stress by activating a key tumor suppressive mechanism.

RESULTS

RNA polymerase I inhibition triggers L22-dependent *MDM4* exon 6 skipping and reduced L22L1 synthesis.

To test the impact of L22 on *MDM4* pre-mRNA splicing, we first induced nucleolar stress in Retinal Pigment Epithelial (RPE) cells, a widely used non-transformed cell line, by treating them with an inhibitor of RNA polymerase I, the enzyme that synthesizes three of the four ribosomal RNAs. As expected, depletion of MDM4 induced the expression of p53 target genes *CDKN1A/p21* and *BBC3/Puma* in these cells (Suppl. Figure 1). In parallel to RNA polymerase I (RNA Pol I) inhibition with the small compound BMH-21 (Peltonen et al., 2014), we efficiently depleted L22 by siRNA transfection (Figures 1A-B). As expected (O'Leary et al., 2013), depletion of L22 triggered a compensatory increase in the mRNA level of the L22 paralogue L22L1 (Figure 1A). RNA Pol I inhibition reduced the level of full-length *MDM4* mRNA while increasing expression of the short isoform in which exon 6 is skipped. Strikingly, however, depletion of L22 dramatically reduced the level of this short isoform and restored expression of full-length *MDM4* mRNA, even when nucleolar stress was induced by RNA Pol I inhibition (Figure 1A). Consistent with this, L22 knockdown also increased the level of MDM4 full-length protein and prolonged its presence despite nucleolar stress (Figure 1B).

Moreover, treatment with BMH-21, but also with a medically relevant inhibitor of rRNA maturation, 5-Fluorouracil (5-FU) (Burger et al., 2010), caused redistribution of L22, which is predominantly present in the nucleolus and cytosol in untreated cells, to enhance its presence within the nucleoplasm (Figures 1C-G). Treatment with these compounds also caused p53 accumulation and augmented the levels of p53 target gene products, p21 and MDM2, while not affecting total L22 protein levels (Figure 1H). Both BMH-21 and 5-FU reduced the levels of MDM4 full-length protein (Figure 1H). We also observed the nucleoplasmic enrichment of L22 upon treatment with BMH-21 or 5-FU in a stably transfected HEK293 Flp-In cell line engineered for the tetracycline-inducible expression of L22 with a C-terminal FLAG tag (Suppl. Figure 2).

We also investigated the impact of L22L1 on expression of MDM4 isoforms in RPE cells and found that its knockdown induced similar splicing changes as depletion of L22, i.e. decreased expression of the short isoform and increased expression of the full-length mRNA (Suppl. Figure 1). Similarly, in HEK293T cells, L22 knockdown reduced the levels of the short *MDM4* isoform and led to increased full-length *MDM4* mRNA and protein levels (Suppl. Figures 3A-B). In this system, however, the effect of L22L1 knockdown on *MDM4* splicing was much less pronounced. To further consolidate the influence of L22 and L22L1 on *MDM4* isoform expression, the paralogous ribosomal proteins were overexpressed in HEK293T cells. Reciprocally to the effects of the knockdowns, this revealed enhanced *MDM4* exon skipping, indicated by reduced expression of the full-length *MDM4* mRNA and increased levels of the

short *MDM4* isoform. (Suppl. Figures 3C-D). Taken together, these results strongly suggest that L22 modulates *MDM4* exon skipping in the context of nucleolar stress.

Depletion of the ribosomal protein S13 triggers L22-dependent nucleolar stress, *MDM4* exon skipping, and p53 activation.

Next, we induced nucleolar stress independent of drugs, i.e. by depletion of a ribosomal protein that belongs to the small ribosomal subunit, S13 (RPS13; uS15), alone or together with L22 depletion (Figures 2A-B). The depletion of S13 alone led to L22 accumulation in the nucleoplasm (Figures 2C-E and Suppl. Figures 4C-E). Lack of S13 alone also reduced the levels of full-length *MDM4* mRNA, increased the levels of the short *MDM4* isoform, and triggered the expression of p53-responsive genes *p21*, *MDM2* and *Puma* (Figure 4A). However, when L22 was depleted along with S13, these effects were substantially diminished (Figure 2A). L22 co-depletion also restored the presence of detectable MDM4 protein upon S13 knockdown, and it diminished the protein levels of the p53-responsive gene products p21 and MDM2 (Figure 2B).

We also induced nucleolar stress by depleting the ribosomal proteins L5 and L11 which are part of the large ribosomal subunit and of the 5S RNP, again combining this with L22 knockdown (Suppl. Figures 4A-B). This again led to nucleoplasmic accumulation of L22 (Figures 2C-E and Suppl. Figures 4C-E). Similar to the previous approach, the reduction of full-length *MDM4* mRNA and protein levels and the increased p53 target gene expression caused by L5 or L11 depletion were L22-dependent (Suppl. Figures 4A-B). Last, S13 or L11 depletion caused a similar L22 re-distribution in the stably transfected HEK293 Flp-In cell line expressing L22-FLAG (Suppl. Figure 5). In conclusion, L22 not only mediates exon-skipping of *MDM4* upon nucleolar stress but also contributes to the subsequent activation of p53.

Intron 6 of the *MDM4* pre-mRNA contains three consensus L22 binding sites and their deletion abolishes the impact of L22 on MDM4 splicing.

L22 has long been known as a ligand of the Epstein Barr Virus Expressed RNAs (EBERs) (Toczyski et al., 1994; Toczyski and Steitz, 1991), and in vitro selection has revealed a specific stem-loop RNA motif that binds L22 (Dobbelstein and Shenk, 1995) (Figure 3A). As L22 induces *MDM4* exon skipping, we suspected a direct interaction of L22 with the pre-mRNA of *MDM4*, perhaps through similar RNA motifs. Consistent with this hypothesis, subjecting the RNA sequence around exon 6 to a folding algorithm, i.e. the RNAfold web server minimum free energy (MFE) function (http://rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi), revealed the presence of three L22-binding consensus motifs within intron 6, in close proximity

to each other and to the adjacent exon 6 (Figures 3B-C). Strikingly, all of the three sequences are conserved in the *MDM4* homologous gene within the murine genome (Suppl. Figure 6A). We therefore tested whether the portion of intron 6 that contains the L22-binding motifs is required for L22-mediated alternative splicing. To this end, we removed this portion of the intron in RPE cells by CRISPR/Cas9 genome editing (Figure 3B and Suppl. Figure 6B). Cells that carry the deletion in a homozygous fashion remained viable. When we subjected these cells to nucleolar stress via S13 depletion, pre-mRNA *MDM4* splicing was not affected, and L22-depletion did not affect the splice pattern of *MDM4* (Figures 3D-G). In line with this, cells lacking the three L22 consensus sequences showed reduced p53 activity upon S13 depletion, as evidenced by diminished expression of *p21* and *Puma* (Suppl. Figures 6C-D). This demonstrates that the region of the *MDM4* pre-mRNA containing the consensus L22 binding sites is strictly required for the impact of L22 on *MDM4* splicing.

L22 also regulates the expression of UBAP2L isoforms.

The previous analysis of exon inclusion vs. genotype in tumor cells not only revealed a correlation between L22-deletions and full-length MDM4 expression, but also between L22-deletions and expression of the gene *UBAP2L* (Ghandi et al., 2019). UBAP2L is a strongly conserved ubiquitin- and RNA-binding protein (Guerber et al., 2022). We therefore hypothesized that L22 might also act as a regulator of *UBAP2L* splicing, and we investigated whether this is the case by combining L22 and RPS13 depletions (Figure 4A). Here, L22-depletion strongly reduced the abundance of a variant that comprises the last *UBAP2L* exon, exon 27, in a more distal version (Figures 4A-B). In contrast, depleting L22 enhanced the inclusion of a more proximal exon 27, corresponding to the isoform that is considered the major one (Figures 4A-B), thereby enhancing the protein levels of this main UBAP2L variant (Figure 4C). This might be explained by the presence of L22 binding sites within the above-mentioned exons (Figure 4B). Similarly, alternative splicing was also induced by L22 in the case of L22L1. Here, the depletion of L22 enhanced the abundance of a functional mRNA encoding L22L1 and reduced a non-coding variant of this mRNA, while RPS13 depletion led to increased levels of L22L1 non-coding RNA (Figures 4A and 4D). We found juxtaposed L22-binding sites in this case as well (Figures 4E-F). In conclusion, L22 regulates the splicing of at least two more genes.

ZMAT3 requires an overlapping portion of intron 6 to induce *MDM4* exon skipping, but does not affect the splicing of *L22L1* and *UBAP2L*.

In previous studies, the protein ZMAT3 was characterized as a splice factor that also triggers *MDM4* exon skipping (Bieging-Rolett et al., 2020). We therefore asked whether ZMAT3 might

act on similar RNA sequence elements as L22 to modulate splicing. To investigate this, we combined L22 and ZMAT3 depletions (Figures 5A-C). Indeed, the depletion of ZMAT3 enhanced the abundance of full-length *MDM4* and reduced the amount of short isoform-mRNA. Moreover, the partial deletion of intron 6 (including the consensus L22 binding sites) abolished the response of *MDM4*-splicing to ZMAT3, at first glance suggesting a similar splice-regulatory mode for ZMAT3 and L22 (Figures 5A-B). However, the simultaneous depletion of ZMAT3 and L22 diminished the short *MDM4* isoform even more strongly than the single knockdowns, arguing that they may not solely be part of the same splice-regulatory mechanism (Figures 5A-B). Moreover, ZMAT3 depletion did not affect the ratio of *L22L1* and *UBAP2L* isoforms, again suggesting that the activity of L22 and ZMAT3 is not the same (Figure 5A). We propose that ZMAT3 and L22 act through similar portions of the *MDM4* intron 6, but that they still use different and at least partially independent mechanisms for doing so.

L22-binding motifs 1 and 2 are especially important for interaction with the *MDM4* pre-mRNA and its splicing.

To more precisely define how L22 influences *MDM4* pre-mRNA splicing, we performed reporter assays. The reporter expresses the enhanced green fluorescent protein (eGFP) upon exon skipping. We cloned *MDM4* exon 6, along with the flanking intronic regions, into this reporter (Figure 6A). As expected, siRNA-mediated depletion of L22 led to enhanced exon inclusion and thus diminished eGFP levels (Figures 6B-C and Suppl. Figure 7A). Next, we mutated the three L22 binding motifs of intron 6 within this reporter construct, individually or in combination, or deleted all three motifs. Upon removal of the first and/or second L22 consensus motif, exon skipping became less dependent on, or even entirely independent of, L22 (Figures 6B-C). Hence, these two RNA motifs are of particular importance for pre-mRNA splicing regulation by L22.

Next, we addressed whether L22 directly binds to its consensus motifs in intron 6 of the *MDM4* pre-mRNA to promote exon skipping. HEK293T cells were transfected with the wildtype *MDM4* reporter plasmid (containing *MDM4* intron 6 and flanking regions) or the construct with mutations of the three L22 motifs, along with an expression construct for flag-tagged L22, and performed RNA immunoprecipitation (RIP) (Figure 6D and Suppl. Figures 7B-C). *MDM4* RNA corresponding to the region directly downstream of the L22 consensus sequences in intron 6 was co-precipitated with L22 (Figures 6D), and this precipitation was reduced by two thirds if the exogenous *MDM4* RNA contained mutated L22 motifs (Figure 6D). Thus, L22 binds to its consensus motifs in the *MDM4* pre-mRNA to regulate *MDM4* isoform expression.

Removing L22-binding sites within intron 6 of the *MDM4* pre-mRNA suppresses p53 activation and enables cell proliferation during nucleolar stress.

Finally, we sought to clarify how L22-mediated splicing regulation of the *MDM4* pre-mRNA influences p53 activity and cell proliferation, especially when cells are confronted with a clinically relevant compound inducing nucleolar stress, i.e. 5-fluorouracil (5-FU) (Burger et al., 2010). We treated RPE cells, with or without the critical region within MDM4 intron 6 containing the L22 binding motifs, with 5-FU, followed by quantification of cell proliferation (Figures 7A-D), and monitoring of p53-responsive gene expression (Figure 7E). As expected, 5-FU induced a p53-response, and it compromised cell proliferation during the following days. However, when the L22-binding sites within *MDM4* intron 6 had been removed, the cells not only displayed reduced expression of p53-responsive genes, but they also proliferated to a significantly higher extent than the parental cells (Figures 7A-E). As expected, 5-FU induced *MDM4* exon 6 skipping and reduced full-length *MDM4* mRNA expression only if the L22 consensus sequences in *MDM4* intron 6 were present (Figures 7E-G). In contrast, the reduction of L22L1 expression upon 5-FU treatment was similar in both cell lines (Figure 7E). In conclusion, L22 serves to integrate nucleolar stress signaling into growth arrest, reaching from compromised rRNA synthesis or processing, through alternative *MDM4* splicing and p53-mediated gene expression, to a pronounced block in cell proliferation (Figure 7H).

DISCUSSION

Our results identify L22 as a signaling transmitter that triggers a p53 response upon impaired ribosome synthesis. When biogenesis of the translation machinery is perturbed, L22 accumulates in the nucleoplasm and associates with a cognate binding motif within a subset of pre-mRNAs, including that of MDM4. Splicing of the MDM4 pre-mRNA is adjusted towards the skipping of exon 6, leading to increased expression of a truncated and poorly functional isoform of MDM4. This promotes p53 to activate the transcription of its target genes.

Interestingly, a link between RPL22 and alternative splicing of the MDM4 pre-mRNA was recently reported in the context of colon cancer of the microsatellite instability subtype (Weinstein et al., 2023), implying that our finding in RPE and HEK293T cells are broadly applicable. These complementary studies both also reveal a role for L22 in modulating the splicing of the L22L1 pre-mRNA. Importantly, our work provides additional mechanistic details on the critical nucleotides within the L22-binding motifs in the MDM4 pre-mRNA and the role of ZMAT3, as well as identifying the *UBAP2L* pre-mRNA as an additional target of L22-modulated alternative splicing.

Nucleolar stress is a widely encountered cell condition. It is not only triggered by dysbalanced ribosomal protein synthesis, e.g. in aneuploid cells, but also occurs in response to DNA damage, as the DNA encoding rRNA responds to damage by perturbed transcription and disintegration of nucleoli (Kruhlak et al., 2007; Rubbi and Milner, 2003). Accordingly, a number of chemotherapeutics can trigger nucleolar stress (Burger et al., 2010; Hein et al., 2013). 5-Fluorouracil (5-FU) does so through incorporation into rRNA (Ghoshal and Jacob, 1997; Sun et al., 2007). Oxaliplatin differs from cisplatin by its ability to modify rRNA, whereas cisplatin primarily acts on DNA (Bruno et al., 2017; Sutton and DeRose, 2021). These agents are thus plausible triggers of L22-mediated MDM4 exon skipping.

Alongside L22, the p53-inducible factor ZMAT3, the arginine methyl transferases PRMT5 and PRMT1, and the splicing factor PRPF19 can modulate the splicing of MDM4 (Bezzi et al., 2013; Bieging-Rolett et al., 2020; Jackson-Weaver et al., 2020; Yano et al., 2021). This emphasizes the role of MDM4 exon skipping in integrating the responses to multiple stresses, all leading to p53 activation. Our data suggest, however, L22 is the first factor that transmits nucleolar stress signals to MDM4.

Besides MDM4, the MDM2 protein is also a target of stress signaling, and this is already known to include nucleolar stress. Not only the 5S RNP, including the ribosomal proteins L5 and L11, binds to MDM2 and prevents p53 ubiquitination (Donati et al., 2013; Onofrillo et al., 2017); other nucleolar proteins, such as nucleophosmin and perhaps even p14[ARF], move from nucleoli to nucleoplasm upon nucleolar stress, where they bind and antagonize MDM2 (Kurki et al., 2004; Llanos et al., 2001; Sherr, 2006; Yang et al., 2016). Besides the 5S RNP, additional

ribosomal proteins were found in association with MDM2, including L22 (Cao et al., 2017; Liu et al., 2016). Thus, L22 may activate p53 in two ways, by triggering MDM4 exon skipping and by directly binding MDM2. However, impaired activation of p53 in the absence of L22-binding RNA motifs within MDM4 intron 6 emphasize the importance of the interaction between L22 and MDM4 pre-mRNA.

Strikingly, in a genome-wide search for associations between exon expression levels and gene mutations in cancers, the link between MDM4 exon 6 inclusion and deletions in the L22-coding gene was one of the top hits (Ghandi et al., 2019). Our results provide a mechanistic explanation for this; when L22 levels are lowered by deletion of at least one allele, this likely prevents L22-induced exon skipping of MDM4 and thus increases the inclusion of exon 6. Even more remarkably, gene losses of L22 are particularly found in endometrial carcinomas of the microsatellite instability (MSI) subtype (Novetsky et al., 2013). This subtype often retains wildtype p53 (Schultheis et al., 2016). In this situation, it is conceivable that elevating full-length MDM4 through L22-deletions will diminish p53 activity and thus allow enhanced tumor cell proliferation. L22 may thus be considered a haploinsufficient tumor suppressor, based on the tumor-promoting traits induced by deleting one of its alleles. L22 is not only unique in this regard, but it is also the only dispensable ribosomal protein, due to the existence of the paralogue L22L1 (O'Leary et al., 2013). This gene duplication event may have facilitated the evolution of tumor-suppressive functions for L22.

Beyond MDM4, our study highlights two additional pre-mRNA targets for L22, i.e. the *L22L1* and *UBAP2L* pre-mRNAs. A broader spectrum of splicing regulation is consistent with a previously reported role of both L22 and L22L1 in zebrafish embryos; there the ribosomal proteins affect the splicing of Smad2 and Smad1 in opposite directions (Zhang et al., 2013; Zhang et al., 2017). The existence of two distinct paralogue genes encoding L22/L22L1 is a unique feature of genes that encode ribosomal proteins in mammalian cells. Indeed, the targeted deletion of L22 does not preclude the survival of mice due to back-up by L22L1 (O'Leary et al., 2013). To maintain the total levels of L22/L22L1, the negative control of L22L1 expression by L22 appears to be required. The ability of L22 to bind the *L22L1* pre-mRNA and suppress its expression was reported previously (O'Leary et al., 2013), but the precise impact of L22 on the L22L1 splicing pattern is first described here (Figure 4).

The other newly identified L22 splicing target, UBAP2L (aka NICE4), is essential for RNA polymerase II degradation when the enzyme hits a DNA lesion (Herlihy et al., 2022). Moreover, it is a core component of stress granules that modulate mRNA processing and storage under various stress conditions (Asano-Inami et al., 2023; Cirillo et al., 2020). Interestingly, UBAP2L is an RNA-binding protein that regulates the expression of genes that determine the global level of translation, such as EIF4G1 (Luo et al., 2020). It is thus tempting to speculate that L22-

mediated suppression of UBAP2L might serve to adapt global translational activity to the availability of ribosomes.

L22 was first discovered as a target of viral RNAs, especially the EBERs, i.e. highly abundant transcripts of RNA polymerase III, encoded by Epstein-Barr virus (Toczyski et al., 1994). It is still unknown which evolutionary advantage led viruses to express L22-interacting RNAs, but one possibility is that this might allow them to modulate cellular pre-mRNA splicing, to modulate the p53-regulatory system.

According to our results, L22 connects three layers of gene expression: biogenesis of the translational machinery, mRNA maturation via pre-mRNA splicing, and RNA synthesis by transcription. Dysfunctional production of the ribosomal subunits triggers an extra-ribosomal splicing-regulatory function of L22, and subsequently, L22-driven alternative splicing of MDM4 determines the activity of p53, the most prominent transcriptional regulator in the context of cancer biology.

ACKNOWLEDGEMENTS

This work was supported by the Deutsche Forschungsgemeinschaft (DFG) via SFB1565, project number 469281184 (P12 to K.E.B., P06 to M.T.B. and P08 to M.D.). J.J. was supported by a doctoral stipend provided by the Studienstiftung des deutschen Volkes.

MATERIAL AND METHODS

<u>Antibodies</u>

Antibody	Source	Identifier
Alexa Fluor 488 goat anti-rabbit	Thermo Fisher Scientific	Cat# A-11034; RRID: AB_2576217
Donkey anti-mouse IgG, HRP conjugated	Jackson ImmunoResearch Labs	Cat# 715-036-150; RRID: AB_2340773
Donkey anti-rabbit IgG, HRP conjugated	Jackson ImmunoResearch Labs	Cat# 711-036-152; RRID: AB_2340590
DYKDDDDK Tag (IF)	Cell Signaling	Cat# 14793; RRID: AB_2572291
DYKDDDDK Tag (WB)	GenScript	Cat# A00187; RRID: AB_1720813
FLAG M2-Peroxidase (HRP) (WB)	Sigma-Aldrich	Cat# A8592; RRID: AB_439702
GAPDH (WB)	Abcam	Cat# ab8245; RRID: AB_2107448
MDM2 IF2 (WB)	Millipore	Cat# OP46-100UG; RRID: AB_564803
MDM4 (WB)	Millipore	Cat# 04-1555; RRID: AB_10562652
p21 (WB)	Cell Signaling Technology	Cat# 2947; RRID: AB_823586
p53 DO-1 (WB)	Santa Cruz Biotechnology	Cat# sc-126; RRID: AB_628082
RPL11 (WB)	Cell Signaling Technology	Cat# 18163; RRID: AB_2798794
RPL22 (IF)	Proteintech	Cat# 25002-1-AP; RRID: AB_2879841
RPL22 (WB)	Santa Cruz Biotechnology	Cat# sc-136413; RRID: AB_10658965
RPL5 (WB)	Cell Signaling Technology	Cat# 51345; RRID: AB_2799391
RPS13 (WB)	Santa Cruz Biotechnology	Cat# sc-398690
UBAP2L (WB)	Cell Signaling Technology	Cat# 40199

ZMAT3 (WB)	Proteintech	Cat# 10504-1-AP; RRID: AB_2217579
β-actin (WB)	Abcam	Cat# ab6276; RRID: AB_2223210

Chemicals, peptides, and recombinant proteins

Reagent	Source	Identifier
5-Fluorouracil	Selleckchem	Cat# S1209
6x DNA Gel Loading Dye	Life Technologies	Cat# R0611
Anti-FLAG M2 Affinity Gel	Sigma-Aldrich	Cat# A2220
Aprotinin	Life Technologies	Cat# A2132
Blasticidin S	Thermo Fisher Scientific	Cat# R21001
BMH-21	Absource Diagnostics/Selleckchem	Cat# S7718
cOmplete, EDTA-free Protease Inhibitor Cocktail	Sigma-Aldrich	Cat# 5056489001
cOmplete, Mini, EDTA-free Protease Inhibitor Cocktail	Sigma-Aldrich	Cat# 11836170001
DAPI	Sigma-Aldrich	Cat# D9542
DMSO	Geyer	Cat# A3672
dNTP Set, 100 mM Solutions	Thermo Fisher Scientific	Cat# R0182
Fluorescence Mounting Medium	Agilent	Cat# S302380-2
GlycoBlue Coprecipitant	Thermo Fisher Scientific	Cat# AM9516
Hygromycin B	Thermo Fisher Scientific	Cat# 10687010
Leupeptin hemisulfate	AppliChem	Cat# A2183
Nuclease-Free Water	Life Technologies	Cat# AM9939
OneTaq® Quick-Load® 2X Master Mix	New England Biolabs	Cat# M0486L
Pefabloc	Carl Roth	Cat# A154
Pepstatin A	VWR	Cat# A2205
Protein G Sepharose	Thermo Fisher Scientific	Cat# 115544935
Proteinase K	Thermo Fisher Scientific	Cat# 25530049
RiboLock RNase Inhibitor	Thermo Fisher Scientific	Cat# EO0381
RNase Inhibitor, Human Placenta	New England Biolabs	Cat# M0307

RNAsIn #4 DG3275	Provided by Prof. Dirk Görlich, Göttingen	N/A
SERVA DNA Stain Clear G	SERVA	Cat# 39804.01
SYBR Green	Thermo Fisher Scientific	Cat# S7567
Taq Polymerase	Primetech	Cat# 1800
TRIzol™ reagent	Life Technologies	Cat# 15596018
Yeast RNA (10 mg/mL)	Thermo Fisher Scientific	Cat# AM7118

Critical commercial assays

Assay	Source	Identifier
Ambion RNase III	Thermo Fisher Scientific	Cat# AM2290
DNase I, RNase-free	Thermo Fisher Scientific	Cat# EN0521
Immobilon Western HRP Substrate	Geyer/Millipore	Cat# WBKLS0500
Lipofectamine 3000 transfection reagent	Thermo Fisher Scientific	Cat# L3000015
MinElute PCR Purification Kit	Qiagen	Cat# 28006X4
M-MuLV Reverse Transcriptase	New England Biolabs	Cat# M0253
Pierce BCA Protein Assay Kit	Thermo Fisher Scientific	Cat# 23227
QuikChange XL Site-Directed Mutagenesis Kit	Agilent	Cat# 200516
SuperSignal West Femto Maximum Sensitivity Substrate	Thermo Fisher Scientific	Cat# 34095

Experimental models: Cell lines

Cell line	Source	Identifier
Human: H1299	ATCC	Cat# CRL-5803; RRID: CVCL_0060
Human: HEK293 Flp-In T-REx	ThermoFisher Scientific	Cat# R78007
Human: HEK293 Flp-In L22-FLAG	This study	N/A

Human: HEK293T	ATCC	Cat# CRL-3216; RRID: CVCL_0063
Human: RPE	ATCC	Cat# CRL-4000; RRID: CVCL_4388
Human: RPE Cas9	Provided by Prof. Manuel Kaulich, Frankfurt	N/A
Human: RPE MDM4 Δ intron 6	This study	N/A

Oligonucleotides

RNA Oligonucleotide	Source	Identifier
Alt-R CRISPR-Cas9 tracrRNA	Integrated DNA Technologies	Order# 3319514
crRNA MDM4_sli6 all_1 TTTATAGTCATCTTGGTACG	This study/Integrated DNA Technologies	Order# 3657380
crRNA Mdm4I61 CAACAAAAGAAGTCACCGTG	This study/Integrated DNA Technologies	Order# 3319514
siRNA ctrl #1 (ssc1)	Thermo Fisher Scientific	Cat# 4390844
siRNA ctrl #2 (Luciferase #1)	Thermo Fisher Scientific	Cat# s237070, 4399665
siRNA ctrl #3 (Luciferase #2)	Thermo Fisher Scientific	Cat# s237071, 4399665
siRNA L22 #1	Thermo Fisher Scientific	Cat# s12190
siRNA L22 #2	Thermo Fisher Scientific	Cat# s12191
siRNA L22L1 #1	Thermo Fisher Scientific	Cat# s47290
siRNA MDM2 #1	Thermo Fisher Scientific	Cat# cs
siRNA MDM4 #1	Thermo Fisher Scientific	Cat# s8631
siRNA RPL11 #1	Thermo Fisher Scientific	Cat# s12168
siRNA RPL5 #1	Thermo Fisher Scientific	Cat# s12151

siRNA RPS13 #1	Thermo Fisher Scientific	Cat# s12304
siRNA RPS13 #2	Thermo Fisher Scientific	Cat# s12305
siRNA ZMAT3 #2	Thermo Fisher Scientific	Cat# s34677

DNA Oligonucleotide (Primer)	Sequence
36B4 forward (V. Manzini)	5'-GATTGGCTACCCAACTGTTG-3'
36B4 reverse (V. Manzini)	5'-CAGGGGCAGCAGCCACAAA-3'
FMv2 3' eGFP forward	5'-ACTCGCAGACCATTACCAGC-3'
FMv2 3' eGFP reverse	5'-CTTTGCTCAGCGCAGATTGG-3'
FMv2-MDM4_A1 (A forward)	5'-CATAGGTACAGACTAGGGGCCAGTCTG TACAACAAAAGAAGTCACC-3'
FMv2-MDM4_A2 (A reverse)	5'-GGTGACTTCTTTTGTTGTACAGACTGGC CCCTAGTCTGTACCTATG-3'
FMv2-MDM4_B1 (B forward)	5'-GGCAGAACAGACTAGGTTCAGTCTGTC CCTGGTCTGTGAAC-3'
FMv2-MDM4_B2 (B reverse)	5'-GTTCACAGACCAGGGACAGACTGAACC TAGTCTGTTCTGCC-3'
FMv2-MDM4_C1 (C forward)	5'-AAGAAAGGTTCAGACTCTGGAAGTCTGT TATGGGCAGAACAGAGAAG-3'
FMv2-MDM4_C2 (C reverse)	5'-CTTCTCTGTTCTGCCCATAACAGACTTC CAGAGTCTGAACCTTTCTT-3'
FMv2-MDM4_KO1 forward	5'-GGTAAAATCACCACACACGTGGCGAGGG TTGG-3'
FMv2-MDM4_KO1 reverse	5'-CCAACCCTCGCCACGTGTGTGGTGATTT TACC-3'
L22 forward	5'-AGCAAGAGCAAGATCACCGT-3'
L22 reverse	5'-AACTACGCGCAACCAGTCA-3'
L22L1 common forward	5'-TGGACCTTACTCATCCAGTAGA-3'
L22L1 common reverse	5'-GCAACCACTCGAAGCCAATC-3'
L22L1 forward	5'-GGAATGTTGTTCACATTGAACGCT-3'
L22L1 reverse	5'-CGAAGCCAATCACGAAGATTGTT-3'
L22L1 mRNA forward	5'-TTTGAGCAATTTCTACGGGAGAAGG-3'
L22L1 mRNA reverse	5'-TGCAACCACTCGAAGCCAA-3'

L22L1 ncRNA forward	5'-TTGGTACCTTCCCTAGTGAATATCT-3'
L22L1 ncRNA reverse	5'-GAAGCCAATCACGAAGATTGTT-3'
MDM2 forward (V. Manzini)	5'-TCAGGATTCAGTTTCAGATCAG-3'
MDM2 reverse (V. Manzini)	5'-CATTTCCAATAGTCAGCTAAGG-3'
MDM4 common forward	5'-TGTCACTTTAGCCACTGCTACT-3'
MDM4 common reverse	5'-AGTGGAACTTTCCTCTGCACTT-3'
MDM4 intron 6 3' of L22 motifs forward	5'-TGATCTTCAGTTCTTCCAGTTGC-3'
MDM4 intron 6 3' of L22 motifs reverse	5'-TGAACTTAACTGCAACACGCC-3'
MDM4_sli6all_1 forward	5'-TCAGACTCTCGCTCTCGCAC-3'
MDM4_sli6all_1 reverse	5'-AAGGCAACACCTCTCCTCAAC-3'
MDM4_sli6all_2 forward	5'-TCTCGCACAGGATCACAGTATG-3'
MDM4_sli6all_2 reverse	5'-CCTCAACCCACTGTGAGCAAA-3'
MDM4-FL forward (K. Wohlberedt)	5'-CTCAGACTCTCGCTCTCGCA-3'
MDM4-FL reverse (K. Wohlberedt)	5'-CTCAAATCCAAGGTCCAGCCT-3'
MDM4-S forward	5'-CTACTGGGACGTCAGAGCTTC-3'
MDM4-S reverse	5'-CCTCTGCACTTTGCTGTAGTA-3'
Nascent p21 forward (K. Henningsen)	5'-AGACCAGCATGACAGGTGCG-3'
Nascent p21 reverse (K. Henningsen)	5'-GCCTGGCATAATGAACATTCCCA-3'
Oligo-dT	5'-TTTTTTTTTTTTTTTTTTTTTTTTTTTV-3' with wobble N 3' modification
p21 forward (V. Manzini)	5'-TAGGCGGTTGAATGAGAGG-3'
p21 reverse (V. Manzini)	5'-AAGTGGGGAGGAGGAAGTAG-3'
p53 forward (V. Manzini)	5'-ATGGAGGAGCCGCAGTCAGATC-3'
p53 reverse (V. Manzini)	5'-GGGAGCAGCCTCTGGCATTCTG-3'
Puma forward (V. Manzini)	5'-GACGACCTCAACGCACAGTA-3'
Puma reverse (V. Manzini)	5'-CTAATTGGGCTCCATCTCG-3'
Random nonamer	5'-NNNNNNNNN-3'
RPL11 forward	5'-GATCCTTTGGCATCCGGAGAA-3'
RPL11 reverse	5'-CAGGCCGTAGATACCAATGCT-3'
RPL5 forward	5'-CAGCGTATGCACACGAACTG-3'
RPL5 reverse	5'-AACCTATTGAGAAGCCTGCGG-3'
RPS13 forward	5'-AGGGACTTGCTCCTGATCTTCC-3'
RPS13 reverse	5'-ACCAGGGCAGAGGCTGTAGAT-3'
UBAP2L-201 forward	5'-CAGCAGCCGCATTCTCAGAT-3'
UBAP2L-201 reverse	5'-CCAAGCTGGTCATCGACGAAA-3'
UBAP2L-207 forward	5'-TGCCTTCAGCCCTAGGAAGT-3'

UBAP2L-207 reverse	5'-AGCTGTAGCTGTTGTAGGCAG-3'
ZMAT3 forward	5'-ATCCTCAGAGCTGGGTCAAC-3'
ZMAT3 reverse	5'-TGGCCACTTGGAGTAACACA-3'
L22_fw_Kozak_BamHI	5'-ATATATGGATCCGCCACCATGGCTCCTGTGAAAAAGCTTGTGG-3'
L22_rv_NheI_nostop	5'-ATATATGCTAGCATCCTCGTCTTCCTCCTCTTCTTCG-3'

Recombinant DNA

Plasmid	Source	Identifier
FMv2 (control)	Fukushima et al., 2020; this study; Thermo Fisher Scientific (GeneArt Gene Synthesis)	Project ID 2023AAAUVC
FMv2-MDM4 (reporter)	This study; Thermo Fisher Scientific (GeneArt Gene Synthesis)	Project ID 2023AAA6HC
FMv2-MDM4_del_1,2,3	This study (generated from FMv2-MDM4 using FMv2-MDM4_KO1 primers)	N/A
FMv2-MDM4_mut_1	This study (generated from FMv2-MDM4 using FMv2-MDM4_A1+2 primers)	N/A
FMv2-MDM4_mut_1,2,3	This study (generated from FMv2-MDM4_mut_1+3 using FMv2-MDM4_B1+2 primers)	N/A
FMv2-MDM4_mut_1+2	This study (generated from FMv2-MDM4_mut_2 using FMv2-MDM4_A1+2 primers)	N/A
FMv2-MDM4_mut_1+3	This study (generated from FMv2-MDM4_mut_1 using FMv2-MDM4_C1+2 primers)	N/A
FMv2-MDM4_mut_2	This study (generated from FMv2-MDM4 using FMv2-MDM4_B1+2 primers)	N/A
FMv2-MDM4_mut_2+3	This study (generated from FMv2-MDM4_mut_3 using FMv2-MDM4_B1+2 primers)	N/A
FMv2-MDM4_mut_3	This study (generated from FMv2-MDM4 using FMv2-MDM4_C1+2 primers)	N/A
pCMV6Entry (empty control)	Origene/VWR	Cat# PS100001

pCMV6Entry-L22	Origene	Cat# RC208910
pCMV6Entry-L22L1	Origene	Cat# RC211790
pOG44	ThermoFisher Scientific	Cat# V600520
pcDNA5-L22-FLAG	This study (generated from Bohnsack lab plasmid pMB187 using primers RPL22_fw_Kozak_BamHI and RPL22_rv_NheI_nostop)	N/A

Software and algorithms

Software and algorithms	Source	Identifier
Fiji v. 2.0.0	NIH	N/A
Image Lab v. 5.2.1	Bio Rad	N/A
Prism v. 9.3.1	GraphPad	N/A
RNAfold web server	Institute for Theoretical Chemistry, University of Vienna	http://rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi

Cell culture

The human non-small cell lung cancer cell line H1299 and the human embryonic kidney cell line containing the SV40 T-antigen HEK293T were maintained in Dulbecco´s Modified Eagle´s Medium (DMEM, 31600091, Thermo Fisher Scientific) supplemented with 10% Fetal Bovine Serum (FBS) (ACSM0190, Anprotec), 2 mM glutamine (25030123, Thermo Fisher Scientific), 50 U/mL penicillin, 50 µg/mL streptomycin (15140122, GIBCO), 10 µg/mL ciprofloxacin (PZN03277618, Fresenius Kabi), and 2 µg/mL tetracycline (87128, Sigma-Aldrich). To generate a stably transfected cell line for the inducible expression of L22 with a C-terminal FLAG tag, HEK293 Flp-In T-Rex cells were reverse transfected with an appropriate pcDNA5-based plasmid and pOG44, which expresses the Flp recombinase, using X-tremeGENE HP DNA transfection reagent (6366546001 Sigma-Aldrich/Roche) according to the manufacturer's instructions. Cells with appropriate genomic integrations in the Flp-In locus were selected using 100 µg/mL hygromycin B (10687010, Thermo Fisher Scientific) and 10 µg/mL blasticidin S (R21001, Thermo Fisher Scientific) for approximately 2 weeks. Colonies were resuspended, pooled and cells were maintained in DMEM GlutaMAX (61965059, Gibco) supplemented with 10% FBS, penicillin, streptomycin, and 100 µg/mL hygromycin B (10687010, Thermo Fisher Scientific). Additionally, 10 µg/mL blasticidin S (R21001, Thermo Fisher Scientific) was added to the culture medium once a week. The selection markers hygromycin B and blasticidin S were removed from culturing medium 24 h prior to experiments. Expression of L22 was induced by adding 1 µg/mL tetracycline for 24 h. The non-tumorigenic human retinal pigment

epithelial cell line RPE and its derivatives RPE p53$^{-/-}$, RPE Cas9 and RPE MDM4 Δ intron 6 were cultured in DMEM GlutaMAX supplemented with 10% FBS, penicillin, streptomycin, ciprofloxacin, and 50 µg/mL hygromycin B for hTERT activation. Hygromycin B was removed from culturing medium 24 h prior to experiments. All cell lines were cultured at 37 °C with 5% CO_2 and routinely tested for mycoplasma contamination.

Cells were treated with BMH-21 (S7718, Absource Diagnostics/Selleckchem), 5-fluorouracil (S1209, Selleckchem), and DMSO (A3672, Geyer).

Transfections of siRNAs were performed using Lipofectamine 3000 (L3000015, Thermo Fisher Scientific). Cells were reverse transfected with final siRNA concentration of 12.5 nM siRNA. Medium was refreshed after 24 h, and cells were harvested after 48 h or 72 h.

Cells were forward transfected with plasmids in amounts of 2.5 µg per well of 6-well plates Medium was refreshed after 6 h except for HEK293T cells to avoid cell detachment, and cells were harvested after 42 h or 48 h.

CRISPR/Cas9-mediated generation of knockout cells

To delete all three L22 consensus sequences in MDM4 intron 6, crRNAs binding upstream (Mdm4I61) and downstream (MDM4_sli6all_1) of these sequences were designed using the web tool CRISPOR.org (Concordet and Haeussler, 2018). For crRNA:tracrRNA 50 µM duplex formation, equal volumes of 100 µM crRNA and 100 µM tracrRNA were mixed, incubated at 95 °C for 5 min and cooled to room temperature.

crRNA:tracrRNA duplexes were transfected into RPE Cas9 cells by reverse transfection using Lipofectamine 3000 transfection reagent (L3000015, Thermo Fisher Scientific) with a final crRNA:tracrRNA concentration of 25 nM (12.5 nM upstream crRNA:tracrRNA, 12.5 nM downstream cr:tracrRNA). Medium was refreshed after 24h.

At 48h post transfection, cells were trypsinized and 50, five or two cells per well were seeded on 96-well plates. After around three weeks, cells from different wells were passaged further to larger plates. RPE MDM4 Δ intron 6 cells, a cell line generated from the 96-well plate with 50 cells per well, were chosen for subsequent experiments since they showed a complete knockout in genotyping and sequencing.

DNA isolation, genotyping, and sequencing

Genotyping and sequencing were performed to identify RPE cell lines generated by CRISPR/Cas9 with deletion of all three L22 consensus sequences in MDM4 intron 6. For DNA isolation, a 2 mL aliquot of cell suspension was centrifuged briefly (3000 rpm, 5 min, 4 °C) and

pellets were washed once with PBS. Each pellet was resuspended in 50 µL 0.1% SDS shearing buffer (1% Triton X-100, 0.15 M NaCl, 1 mM EDTA, 0.5 mM EGTA, 20 mM HEPES, 0.1% SDS), followed by the addition of 50 µL 10 mM Tris, 100 µL 2x Weinmann buffer pH 8.0 (100 mM Tris-HCl pH 8.8, 20 mM EDTA, 2% SDS) and 2 µL 20 mg/mL proteinase K (25530049, Thermo Fisher Scientific). Samples were incubated overnight at 65 °C and 800 rpm, 200 µL 10 mM Tris was added, and DNA was isolated using the MinElute PCR Purification Kit (28006X4, Qiagen).

For genotyping, MDM4_sli6all_1 primers binding upstream and downstream of the crRNA binding sites were used. One genotyping PCR sample contained 10 µL OneTaq® Quick-Load® 2X Master Mix (M0486L, New England Biolabs), 1 µL 10 µM forward primer, 1 µL 10 µM reverse primer, 3 µL ddH$_2$O, and 5 µL ddH$_2$O. The PCR program was a preheating step (2 min, 95 °C), followed by 36 cycles of 95 °C for 30 sec, 56 °C for 1 min and 72 °C for 1.5 min, with a final step at 72 °C for 5 min. Genotyping PCR products were separated by agarose gel electrophoresis using SERVA DNA Stain Clear G (39804.01, SERVA) for DNA detection under a UV illuminator.

DNA samples showing the PCR product sizes expected for successful knockout in genotyping were sequenced for further confirmation. For this, genotyping PCR products were purified using MinElute PCR Purification Kit (28006X4, Qiagen), DNA concentration was measured by spectrophotometry, and samples were sequenced with MDM4_sli6all_2 forward and reverse primers (Eurofins Genomics).

RNA isolation, reverse transcription, and real-time quantitative PCR

Cells were washed with PBS and harvested in TRIzol™ reagent containing phenol (15596018, Life Technologies). For total RNA extraction, 200 µL chloroform per 1 mL TRIzol™ was added. Samples were centrifuged for phase separation, and the upper aqueous phase was used for precipitating the RNA by adding 500 µL isopropanol – and optionally 2 µL GlycoBlue Coprecipitant (AM9516, Thermo Fisher Scientific) – and centrifuging. The RNA pellet was washed once with 75% ethanol, dried, and resuspended in nuclease-free H$_2$O (AM9939, Life Technologies). RNA concentration was measured by spectrophotometry.

For reverse transcription, 1 µg total RNA was incubated with 6.25 µM oligo-dT (Metabion) and 1.875 µM random nonamer primers (Metabion) as well as 0.625 mM dNTPs (R0182, Thermo Fisher Scientific) at 70 °C for 5 min for primer annealing. Next, 10 U of RNase Inhibitor, Human Placenta (M0307, New England Biolabs) and 25 U of M-MuLV Reverse Transcriptase with corresponding buffer (M0253, New England Biolabs) was added and the reaction was incubated at 42 °C for 1 h, followed by 5 min incubation at 95 °C for inactivating the reverse

transcriptase. A control reaction without reverse transcriptase (no reverse transcriptase control, NRT) was included. Finally, the synthesized cDNA was diluted by adding 130 µL ddH₂O per 20 µL cDNA reaction mixture and used as a template for real-time quantitative PCR (RT-qPCR).

For quantification, RT-qPCR was performed in technical duplicates. One well of a 96-well plate (721980010, Sarstedt) contained one reaction consisting of 1 µL 10 µM forward primer, 1 µL 10 µM reverse primer, 3 µL cDNA, 6 µL ddH₂O, and 14 µL of a homemade RT-qPCR reaction mixture containing 518 mM trehalose (stock in 10 mM Tris pH 8.0), 130 mM Tris-HCl pH 8.8, 35 mM ammonium sulphate, 0.017% Tween-20, 0.43% Triton X-100, 5.18 mM MgCl₂, 0.43x concentrated SYBR Green (S7567, Life Technologies), 0.35 mM dNTPs (R0182, Thermo Fisher Scientific), and 35 U/mL Taq polymerase (1800, Primetech). The program used for the thermocycler was a preheating step (2 min, 95 °C), followed by 45 cycles of 95 °C for 15 sec and 60 °C for 1 min. Subsequently, a melting curve was generated, followed by a final incubation at 95 °C for 30 sec.

Gene expression levels were normalized to the mRNA levels of the housekeeping gene 36B4, and the analysis was conducted using the ΔΔCt method.

For semi-quantitative detection of MDM4 and L22L1 splice variants using common primers leading to products of different sizes for the respective splice variants, RT-qPCR was performed as usual. Next, 6x DNA Gel Loading Dye (R0611, Life Technologies) was added to RT-qPCR products and they were separated and visualized by agarose gel electrophoresis using SERVA DNA Stain Clear G (39804.01, SERVA) for DNA detection under a UV illuminator. Band quantification was performed using ImageJ. Each band was defined as a region of interest (ROI) and a background ROI was defined as well. Band intensities were determined and resulting "mean" values of sample minus background were used to determine splice variant ratios.

Immunoblot analysis

Cells were washed with PBS and harvested in lysis buffer, i.e. 14 mM Tris-HCl pH 7.5, 100 mM NaCl, 9 mM EDTA, 1% Triton X-100, 1% deoxycholate, 0.1% SDS, 2.7 M urea, supplemented with protease inhibitors (aprotinin, leupeptin hemisulfate, aminoethyl-benzene-sulfonyl fluoride/Pefabloc, pepstatin A). The samples were briefly sonicated and pushed through a syringe to disrupt chromatin. Protein concentration was measured using a BCA protein assay kit (23227, Thermo Fisher Scientific). Equal protein amounts of each sample were boiled in Laemmli buffer at 95 °C for 5 min and separated by SDS-PAGE. Proteins were transferred to a nitrocellulose membrane. Membranes were blocked with 5% (w/v) non-fat milk

(Roth) in TBS containing 0.1% Tween-20 (VWR) for 1h and incubated with the primary antibodies at 4 °C overnight. Next, membranes were incubated with horse radish peroxidase (HRP)-conjugated secondary antibodies (donkey anti-mouse or anti-rabbit IgG, Jackson ImmunoResearch Labs), followed by protein detection using either Immobilon Western HRP Substrate (Geyer/Millipore) or SuperSignal West Femto Maximum Sensitivity Substrate (Thermo Fisher Scientific).

RNA immunoprecipitation (RIP)

For RIP to precipitate FLAG-tagged L22 and its bound RNAs, cells grown on 6 wells of 6-well plates per sample were washed once with PBS, and RNA and protein were crosslinked with 0.75% paraformaldehyde (PFA) in PBS for 10 min. Fixation was quenched by adding 0.125 M glycine for 5 min. Next, cells were washed once with PBS and harvested on ice in RIP high salt buffer (500 mM NaCl, 50 mM Tris, 5 mM EDTA, 0.5% NP-40), supplemented with protease inhibitors (11836170001, Sigma-Aldrich) and, in case of samples used not only for protein but also for bound RNA precipitation, with 0.1 U/µL RNase inhibitor (RNAsIn #4 DG3275, kindly provided by D. Görlich). All following steps were performed on ice. Samples were pushed through a syringe and sonicated to disrupt chromatin, followed by centrifugation for 10 min at 16100 rcf and 4 °C to remove cell debris, proceeding with the supernatant. The cell lysate was pre-cleared with Protein G Sepharose (115544935, Thermo Fisher Scientific) for 2h at 4 °C, and 5% of each sample was preserved as input. This was followed by an overnight incubation at 4 °C with 30 µL (15 µL beads) per sample of Anti-FLAG M2 Affinity Gel (A2220, Sigma-Aldrich). Additionally, 20 µg/mL yeast RNA (AM7118, Thermo Fisher Scientific) was added for this precipitation step as a competitor RNA. Beads were washed three times with RIP high salt buffer and three times with RIP low salt buffer (5 mM NaCl, 50 mM Tris, 5 mM EDTA, 0.5% NP-40), supplemented with protease inhibitors and RNase inhibitor as described above. Samples only used for protein precipitation were washed once more with RIP high salt buffer. Beads were then resuspended in Laemmli buffer and boiled at 95 °C for 5 min prior to SDS-PAGE and immunoblot analysis using FLAG antibody covalently conjugated to horseradish peroxidase for detection of precipitated L22 (A8592, Sigma-Aldrich).

For samples also used for RNA precipitation, a digestion with RNase III (AM2290, Thermo Fisher Scientific) was performed to generate smaller fragments from precipitated RNAs. For this, beads were incubated with 0.05 U/µL RNase III and corresponding buffer at 37 °C for 30 min, followed by another washing step with RIP high salt buffer. The RNA-protein complexes of immunoprecipitation (IP) and input samples were then decrosslinked by adding 0.4 µg/µL proteinase K (25530049, Thermo Fisher Scientific) in RIP high salt buffer for 1h at 60 °C. 500 µL TRIzol™ reagent was added per sample to input and beads, and RNA was isolated as

usual. Each RNA pellet was resuspended in 11 µL nuclease-free H_2O, and DNase treatment was performed by incubating 8.8 µL RNA with 2 U DNase I, RNase-free with corresponding buffer (EN0521, Thermo Fisher Scientific) and 20 U RiboLock RNase Inhibitor (EO0381, Thermo Fisher Scientific) at 37 °C for 30 min. Next, 4.5 mM EDTA was added and samples were incubated at 65 °C for 10 min for DNase inactivation. 10 µL from 22 µL DNase-treated RNA was used as a template for reverse transcription as described above, except that RNA was incubated with 6.25 µM random nonamer primers (Metabion). A control reaction without reverse transcriptase (no reverse transcriptase control, NRT) was included for each sample. Finally, the synthesized cDNA was used as a template for RT-qPCR. The quantification of the RT-qPCR signal was performed relative to the input.

Immunofluorescence

Cells were grown in 8-well chambers on PCA slides with removable frame (946140802, Sarstedt). They were fixed in 3.7% PFA in PBS for 30 min, rinsed with PBS, and permeabilized with 0.5% Triton X-100 in PBS for 30 min. After 2x 5 min washing with PBS, blocking solution (10% FBS 0.1% Tween-20 in PBS) was added to the cells for 10 min, followed by incubation with primary antibody produced in rabbit, diluted in blocking solution, at 4 °C overnight. Cells were washed 2x 5 min with PBS, and secondary antibody (Alexa Fluor 488 goat anti-rabbit; A-11034, Thermo Fisher Scientific) (1:500) and 4',6-diamidino-2-phenylindole (DAPI; D9542, Sigma-Aldrich) (1:3000) diluted in blocking solution were added for 1h at room temperature. Two more washing steps with PBS were performed, gaskets were removed and slides were mounted onto coverslips using Fluorescence Mounting Medium (S302380-2, Agilent). Images were acquired using a Zeiss Axio Scope.A1 fluorescence microscope with 40x and 100x magnifications and further analyzed with ImageJ. For quantification of nuclear L22, the DAPI channel was used to create binary images (segmentation) to define the nuclei as regions of interest (ROIs). These ROIs were then used to quantify the L22 signal within these regions. Mean fluorescence intensity values were visualized.

MDM4 splicing reporter plasmid generation

FMv2 (control plasmid) was generated by inserting the following sequences into the pcDNA3.1/Hygro(+) vector containing the CMV promoter using the restriction enzymes *Nhe*I and *Xho*I: the mCherry coding region, the 5′ end of the eGFP gene, the 5′ end of DMD intron 18, a Multiple Cloning Site (MCS), the 3′ end of DMD intron 19, and the 3′ end of the eGFP gene. FMv2-MDM4 (MDM4 splicing reporter plasmid) was generated by inserting the following sequences into the MCS of FMv2 using the restriction enzymes *Hind*III and *Bam*HI: the 3′ end

(34 bp) of MDM4 intron 5, MDM4 exon 6 (68 bp), and the 5′ end (398 bp) of MDM4 intron 6 containing all three L22 consensus sequences. For both plasmids, the GeneArt Gene Synthesis service (Thermo Fisher Scientific) was used. Different FMv2-MDM4 versions containing mutations or deletions of single or combined L22 consensus sequences in MDM4 intron 6 were generated by QuikChange site-directed mutagenesis (200516, Agilent).

MDM4 splicing reporter assay

The principle of this assay is as follows: upon overexpression of either FMv2 or FMv2-MDM4 or its mutated versions in cells, mCherry is constitutively expressed and serves as a transfection marker. eGFP, the sequence of which is split into two parts in all plasmids, is also constitutively expressed upon FMv2 overexpression, but only if the MDM4 exon 6 is skipped in case of FMv2-MDM4 (or variant) overexpression, thus serving as a splicing marker.

For the assay, cells were grown on 24-well plates (3524, Corning Costar/Geyer), transfected with control (ctrl#1) or L22 (L22#1) siRNAs for 72h, and with control plasmid (FMv2) (Fukushima et al., 2020), MDM4 reporter plasmid (FMv2-MDM4) or different FMv2-MDM4 versions for 48h. Using the ZEISS Celldiscoverer 7 live cell fluorescence microscope with 5x magnification, red fluorescence (mCherry, exposure time 100 ms, intensity 30%) and green fluorescence (eGFP, exposure time 20 ms, intensity 30%) were detected, and five to six images per well were taken.

Images were further analyzed with ImageJ. The mCherry channel was used to create binary images (segmentation) to define strongly transfected cells (i.e., cells with high mCherry intensity above a threshold) as regions of interest (ROIs). These ROIs were then used to quantify the eGFP signal within these regions. The average of all resulting "mean" values was calculated and the L22#1/ctrl#1 ratio of these values was determined.

Cell confluence measurements

Cells were plated on 24-well plates (3524, Corning Costar/Geyer) and continuously treated with the indicated compounds starting on the day after seeding, with a refreshment of treatment every other day. On each day of treatment, cell confluence was measured and analyzed using the Celigo Adherent Cell Cytometer.

<u>Statistical analysis</u>

Statistical analysis was performed using GraphPad Prism software. Unpaired Student 's t test was used for statistical analysis of RT-qPCR, agarose gel band quantification, reporter experiments, and confluence measurements. Mann-Whitney U test was used for statistical analysis of quantification of single nuclear intensities of L22 immunofluorescence signal. *, P≤0.05; **, P≤0.01; ***, P≤0.001; ****, P≤0.0001; ns, not significant.

REFERENCES

Asano-Inami, E., A. Yokoi, M. Sugiyama, T. Hyodo, T. Hamaguchi, and H. Kajiyama. 2023. The association of UBAP2L and G3BP1 mediated by small nucleolar RNA is essential for stress granule formation. *Communications biology*. 6:415.

Bardot, B., and F. Toledo. 2017. Targeting MDM4 Splicing in Cancers. *Genes*. 8.

Bezzi, M., S.X. Teo, J. Muller, W.C. Mok, S.K. Sahu, L.A. Vardy, Z.Q. Bonday, and E. Guccione. 2013. Regulation of constitutive and alternative splicing by PRMT5 reveals a role for Mdm4 pre-mRNA in sensing defects in the spliceosomal machinery. *Genes & development*. 27:1903-1916.

Bieging-Rolett, K.T., A.M. Kaiser, D.W. Morgens, A.M. Boutelle, J.A. Seoane, E.L. Van Nostrand, C. Zhu, S.L. Houlihan, S.S. Mello, B.A. Yee, J. McClendon, S.E. Pierce, I.P. Winters, M. Wang, A.J. Connolly, S.W. Lowe, C. Curtis, G.W. Yeo, M.M. Winslow, M.C. Bassik, and L.D. Attardi. 2020. Zmat3 Is a Key Splicing Regulator in the p53 Tumor Suppression Program. *Molecular cell*. 80:452-469.e459.

Boulon, S., B.J. Westman, S. Hutten, F.M. Boisvert, and A.I. Lamond. 2010. The nucleolus under stress. *Molecular cell*. 40:216-227.

Bruno, P.M., Y. Liu, G.Y. Park, J. Murai, C.E. Koch, T.J. Eisen, J.R. Pritchard, Y. Pommier, S.J. Lippard, and M.T. Hemann. 2017. A subset of platinum-containing chemotherapeutic agents kills cells by inducing ribosome biogenesis stress. *Nature medicine*. 23:461-471.

Burger, K., B. Mühl, T. Harasim, M. Rohrmoser, A. Malamoussi, M. Orban, M. Kellner, A. Gruber-Eber, E. Kremmer, M. Hölzel, and D. Eick. 2010. Chemotherapeutic drugs inhibit ribosome biogenesis at various levels. *The Journal of biological chemistry*. 285:12416-12425.

Cao, B., Z. Fang, P. Liao, X. Zhou, J. Xiong, S. Zeng, and H. Lu. 2017. Cancer-mutated ribosome protein L22 (RPL22/eL22) suppresses cancer cell survival by blocking p53-MDM2 circuit. *Oncotarget*. 8:90651-90661.

Cirillo, L., A. Cieren, S. Barbieri, A. Khong, F. Schwager, R. Parker, and M. Gotta. 2020. UBAP2L Forms Distinct Cores that Act in Nucleating Stress Granules Upstream of G3BP1. *Current biology : CB*. 30:698-707.e696.

Concordet, J.P., and M. Haeussler. 2018. CRISPOR: intuitive guide selection for CRISPR/Cas9 genome editing experiments and screens. *Nucleic acids research*. 46:W242-w245.

Dewaele, M., T. Tabaglio, K. Willekens, M. Bezzi, S.X. Teo, D.H. Low, C.M. Koh, F. Rambow, M. Fiers, A. Rogiers, E. Radaelli, M. Al-Haddawi, S.Y. Tan, E. Hermans, F. Amant, H. Yan, M. Lakshmanan, R.C. Koumar, S.T. Lim, F.A. Derheimer, R.M. Campbell, Z. Bonday, V. Tergaonkar, M. Shackleton, C. Blattner, J.C. Marine, and E. Guccione. 2016. Antisense oligonucleotide-mediated MDM4 exon 6 skipping impairs tumor growth. *The Journal of clinical investigation*. 126:68-84.

Dobbelstein, M., and T. Shenk. 1995. In vitro selection of RNA ligands for the ribosomal L22 protein associated with Epstein-Barr virus-expressed RNA by using randomized and cDNA-derived RNA libraries. *Journal of virology*. 69:8027-8034.

Donati, G., S. Peddigari, C.A. Mercer, and G. Thomas. 2013. 5S ribosomal RNA is an essential component of a nascent ribosomal precursor complex that regulates the Hdm2-p53 checkpoint. *Cell reports*. 4:87-98.

Fukushima, S., M. Farea, K. Maeta, A.Q.M. Rani, K. Fujioka, H. Nishio, and M. Matsuo. 2020. Dual Fluorescence Splicing Reporter Minigene Identifies an Antisense Oligonucleotide to Skip Exon v8 of the CD44 Gene. *International journal of molecular sciences*. 21.

Ghandi, M., F.W. Huang, J. Jané-Valbuena, G.V. Kryukov, C.C. Lo, E.R. McDonald, 3rd, J. Barretina, E.T. Gelfand, C.M. Bielski, H. Li, K. Hu, A.Y. Andreev-Drakhlin, J. Kim, J.M. Hess, B.J. Haas, F. Aguet, B.A. Weir, M.V. Rothberg, B.R. Paolella, M.S. Lawrence, R. Akbani, Y. Lu, H.L. Tiv, P.C. Gokhale, A. de Weck, A.A. Mansour, C. Oh, J. Shih, K. Hadi, Y. Rosen, J. Bistline, K. Venkatesan, A. Reddy, D. Sonkin, M. Liu, J. Lehar, J.M. Korn, D.A. Porter, M.D. Jones, J. Golji, G. Caponigro, J.E. Taylor, C.M. Dunning, A.L. Creech, A.C. Warren, J.M. McFarland, M. Zamanighomi, A. Kauffmann, N. Stransky, M. Imielinski, Y.E. Maruvka, A.D. Cherniack, A. Tsherniak, F. Vazquez, J.D. Jaffe, A.A. Lane, D.M. Weinstock, C.M. Johannessen, M.P. Morrissey, F. Stegmeier, R. Schlegel, W.C. Hahn, G. Getz, G.B. Mills, J.S. Boehm, T.R. Golub, L.A. Garraway, and W.R. Sellers. 2019. Next-generation characterization of the Cancer Cell Line Encyclopedia. *Nature*. 569:503-508.

Ghoshal, K., and S.T. Jacob. 1997. An alternative molecular mechanism of action of 5-fluorouracil, a potent anticancer drug. *Biochemical pharmacology*. 53:1569-1575.

Gu, J., H. Kawai, L. Nie, H. Kitao, D. Wiederschain, A.G. Jochemsen, J. Parant, G. Lozano, and Z.M. Yuan. 2002. Mutual dependence of MDM2 and MDMX in their functional inactivation of p53. *The Journal of biological chemistry*. 277:19251-19254.

Guerber, L., E. Pangou, and I. Sumara. 2022. Ubiquitin Binding Protein 2-Like (UBAP2L): is it so NICE After All? *Frontiers in cell and developmental biology*. 10:931115.

Harris, B., D.K. Singh, M. Verma, S.P. Fahl, M. Rhodes, S.R. Sprinkle, M. Wang, Y. Zhang, J. Perrigoue, R. Kessel, S. Peri, J. West, O. Giricz, J. Boultwood, A. Pellagatti, K.H. Ramesh, C. Montagna, K. Pradhan, J.W. Tyner, B.K. Kennedy, M. Holinstat, U. Steidl, S. Sykes, A. Verma, and D.L. Wiest. 2023. Ribosomal protein control of hematopoietic stem cell transformation through direct, non-canonical regulation of metabolism. *bioRxiv : the preprint server for biology*.

Haupt, Y., R. Maya, A. Kazaz, and M. Oren. 1997. Mdm2 promotes the rapid degradation of p53. *Nature*. 387:296-299.

Hein, N., K.M. Hannan, A.J. George, E. Sanij, and R.D. Hannan. 2013. The nucleolus: an emerging target for cancer therapy. *Trends in molecular medicine*. 19:643-654.

Herlihy, A.E., S. Boeing, J.C. Weems, J. Walker, A.B. Dirac-Svejstrup, M.H. Lehner, R.C. Conaway, J.W. Conaway, and J.Q. Svejstrup. 2022. UBAP2/UBAP2L regulate UV-induced ubiquitylation of RNA polymerase II and are the human orthologues of yeast Def1. *DNA repair*. 115:103343.

Honda, R., H. Tanaka, and H. Yasuda. 1997. Oncoprotein MDM2 is a ubiquitin ligase E3 for tumor suppressor p53. *FEBS letters*. 420:25-27.

Jackson-Weaver, O., N. Ungvijanpunya, Y. Yuan, J. Qian, Y. Gou, J. Wu, H. Shen, Y. Chen, M. Li, S. Richard, Y. Chai, H.M. Sucov, and J. Xu. 2020. PRMT1-p53 Pathway Controls Epicardial EMT and Invasion. *Cell reports*. 31:107739.

Juven, T., Y. Barak, A. Zauberman, D.L. George, and M. Oren. 1993. Wild type p53 can mediate sequence-specific transactivation of an internal promoter within the mdm2 gene. *Oncogene*. 8:3411-3416.

Kandoth, C., M.D. McLellan, F. Vandin, K. Ye, B. Niu, C. Lu, M. Xie, Q. Zhang, J.F. McMichael, M.A. Wyczalkowski, M.D.M. Leiserson, C.A. Miller, J.S. Welch, M.J. Walter, M.C. Wendl, T.J. Ley, R.K. Wilson, B.J. Raphael, and L. Ding. 2013. Mutational landscape and significance across 12 major cancer types. *Nature*. 502:333-339.

Kruhlak, M., E.E. Crouch, M. Orlov, C. Montaño, S.A. Gorski, A. Nussenzweig, T. Misteli, R.D. Phair, and R. Casellas. 2007. The ATM repair pathway inhibits RNA polymerase I transcription in response to chromosome breaks. *Nature*. 447:730-734.

Kubbutat, M.H., S.N. Jones, and K.H. Vousden. 1997. Regulation of p53 stability by Mdm2. *Nature*. 387:299-303.

Kurki, S., K. Peltonen, L. Latonen, T.M. Kiviharju, P.M. Ojala, D. Meek, and M. Laiho. 2004. Nucleolar protein NPM interacts with HDM2 and protects tumor suppressor protein p53 from HDM2-mediated degradation. *Cancer cell*. 5:465-475.

Leach, F.S., T. Tokino, P. Meltzer, M. Burrell, J.D. Oliner, S. Smith, D.E. Hill, D. Sidransky, K.W. Kinzler, and B. Vogelstein. 1993. p53 Mutation and MDM2 amplification in human soft tissue sarcomas. *Cancer research*. 53:2231-2234.

Levine, A.J., and M. Oren. 2009. The first 30 years of p53: growing ever more complex. *Nature reviews. Cancer*. 9:749-758.

Li, J., and M. Kurokawa. 2015. Regulation of MDM2 Stability After DNA Damage. *Journal of cellular physiology*. 230:2318-2327.

Liu, Y., C. Deisenroth, and Y. Zhang. 2016. RP-MDM2-p53 Pathway: Linking Ribosomal Biogenesis and Tumor Surveillance. *Trends in cancer*. 2:191-204.

Llanos, S., P.A. Clark, J. Rowe, and G. Peters. 2001. Stabilization of p53 by p14ARF without relocation of MDM2 to the nucleolus. *Nature cell biology*. 3:445-452.

Luo, E.C., J.L. Nathanson, F.E. Tan, J.L. Schwartz, J.C. Schmok, A. Shankar, S. Markmiller, B.A. Yee, S. Sathe, G.A. Pratt, D.B. Scaletta, Y. Ha, D.E. Hill, S. Aigner, and G.W. Yeo. 2020. Large-scale tethered function assays identify factors that regulate mRNA stability and translation. *Nature structural & molecular biology*. 27:989-1000.

Novetsky, A.P., I. Zighelboim, D.M. Thompson, Jr., M.A. Powell, D.G. Mutch, and P.J. Goodfellow. 2013. Frequent mutations in the RPL22 gene and its clinical and functional implications. *Gynecologic oncology*. 128:470-474.

O'Leary, M.N., K.H. Schreiber, Y. Zhang, A.C. Duc, S. Rao, J.S. Hale, E.C. Academia, S.R. Shah, J.F. Morton, C.A. Holstein, D.B. Martin, M. Kaeberlein, W.C. Ladiges, P.J. Fink, V.L. Mackay, D.L. Wiest, and B.K. Kennedy. 2013. The ribosomal protein Rpl22 controls ribosome composition by directly repressing expression of its own paralog, Rpl22l1. *PLoS genetics*. 9:e1003708.

Oliner, J.D., A.Y. Saiki, and S. Caenepeel. 2016. The Role of MDM2 Amplification and Overexpression in Tumorigenesis. *Cold Spring Harbor perspectives in medicine*. 6.

Olivier, M., M. Hollstein, and P. Hainaut. 2010. TP53 mutations in human cancers: origins, consequences, and clinical use. *Cold Spring Harbor perspectives in biology*. 2:a001008.

Onofrillo, C., A. Galbiati, L. Montanaro, and M. Derenzini. 2017. The pre-existing population of 5S rRNA effects p53 stabilization during ribosome biogenesis inhibition. *Oncotarget*. 8:4257-4267.

Parant, J., A. Chavez-Reyes, N.A. Little, W. Yan, V. Reinke, A.G. Jochemsen, and G. Lozano. 2001. Rescue of embryonic lethality in Mdm4-null mice by loss of Trp53 suggests a nonoverlapping pathway with MDM2 to regulate p53. *Nature genetics*. 29:92-95.

Peltonen, K., L. Colis, H. Liu, R. Trivedi, M.S. Moubarek, H.M. Moore, B. Bai, M.A. Rudek, C.J. Bieberich, and M. Laiho. 2014. A targeting modality for destruction of RNA polymerase I that possesses anticancer activity. *Cancer cell*. 25:77-90.

Rizzotto, D., L. Englmaier, and A. Villunger. 2021. At a Crossroads to Cancer: How p53-Induced Cell Fate Decisions Secure Genome Integrity. *International journal of molecular sciences*. 22.

Rubbi, C.P., and J. Milner. 2003. Disruption of the nucleolus mediates stabilization of p53 in response to DNA damage and other stresses. *The EMBO journal*. 22:6068-6077.

Schultheis, A.M., L.G. Martelotto, M.R. De Filippo, S. Piscuglio, C.K. Ng, Y.R. Hussein, J.S. Reis-Filho, R.A. Soslow, and B. Weigelt. 2016. TP53 Mutational Spectrum in Endometrioid and Serous Endometrial Cancers. *International journal of gynecological pathology : official journal of the International Society of Gynecological Pathologists*. 35:289-300.

Sherr, C.J. 2006. Divorcing ARF and p53: an unsettled case. *Nature reviews. Cancer*. 6:663-673.

Shvarts, A., W.T. Steegenga, N. Riteco, T. van Laar, P. Dekker, M. Bazuine, R.C. van Ham, W. van der Houven van Oordt, G. Hateboer, A.J. van der Eb, and A.G. Jochemsen. 1996. MDMX: a novel p53-binding protein with some functional properties of MDM2. *The EMBO journal*. 15:5349-5357.

Sloan, K.E., M.T. Bohnsack, and N.J. Watkins. 2013. The 5S RNP couples p53 homeostasis to ribosome biogenesis and nucleolar stress. *Cell reports*. 5:237-247.

Sun, X.X., M.S. Dai, and H. Lu. 2007. 5-fluorouracil activation of p53 involves an MDM2-ribosomal protein interaction. *The Journal of biological chemistry*. 282:8052-8059.

Sutton, E.C., and V.J. DeRose. 2021. Early nucleolar responses differentiate mechanisms of cell death induced by oxaliplatin and cisplatin. *The Journal of biological chemistry*. 296:100633.

Tanimura, S., S. Ohtsuka, K. Mitsui, K. Shirouzu, A. Yoshimura, and M. Ohtsubo. 1999. MDM2 interacts with MDMX through their RING finger domains. *FEBS letters*. 447:5-9.

Toczyski, D.P., A.G. Matera, D.C. Ward, and J.A. Steitz. 1994. The Epstein-Barr virus (EBV) small RNA EBER1 binds and relocalizes ribosomal protein L22 in EBV-infected human B lymphocytes. *Proceedings of the National Academy of Sciences of the United States of America*. 91:3463-3467.

Toczyski, D.P., and J.A. Steitz. 1991. EAP, a highly conserved cellular protein associated with Epstein-Barr virus small RNAs (EBERs). *The EMBO journal*. 10:459-466.

Vousden, K.H., and C. Prives. 2009. Blinded by the Light: The Growing Complexity of p53. *Cell*. 137:413-431.

Weinstein, H.N.W., K. Hu, L. Fish, Y.A. Chen, P. Allegakoen, K.S.F. Hui, J.H. Pham, M.B. Baco, H. Song, A.O. Giacomelli, F. Vazquez, M. Ghandi, H. Goodarzi, and F.W. Huang. 2023. RPL22 is a tumor suppressor in MSI-high cancers and a key splicing regulator of MDM4. *bioRxiv : the preprint server for biology*.

Wu, J., G. Lu, and X. Wang. 2021. MDM4 alternative splicing and implication in MDM4 targeted cancer therapies. *American journal of cancer research*. 11:5864-5880.

Wu, X., J.H. Bayle, D. Olson, and A.J. Levine. 1993. The p53-mdm-2 autoregulatory feedback loop. *Genes & development*. 7:1126-1132.

Yang, K., M. Wang, Y. Zhao, X. Sun, Y. Yang, X. Li, A. Zhou, H. Chu, H. Zhou, J. Xu, M. Wu, J. Yang, and J. Yi. 2016. A redox mechanism underlying nucleolar stress sensing by nucleophosmin. *Nature communications*. 7:13599.

Yang, K., J. Yang, and J. Yi. 2018. Nucleolar Stress: hallmarks, sensing mechanism and diseases. *Cell stress*. 2:125-140.

Yano, K., R.U. Takahashi, B. Shiotani, J. Abe, T. Shidooka, Y. Sudo, Y. Yamamoto, S. Kan, H. Sakagami, and H. Tahara. 2021. PRPF19 regulates p53-dependent cellular senescence by modulating alternative splicing of MDM4 mRNA. *The Journal of biological chemistry*. 297:100882.

Zhang, Y., A.C. Duc, S. Rao, X.L. Sun, A.N. Bilbee, M. Rhodes, Q. Li, D.J. Kappes, J. Rhodes, and D.L. Wiest. 2013. Control of hematopoietic stem cell emergence by antagonistic functions of ribosomal protein paralogs. *Developmental cell*. 24:411-425.

Zhang, Y., M.N. O'Leary, S. Peri, M. Wang, J. Zha, S. Melov, D.J. Kappes, Q. Feng, J. Rhodes, P.S. Amieux, D.R. Morris, B.K. Kennedy, and D.L. Wiest. 2017. Ribosomal Proteins Rpl22 and Rpl22l1 Control Morphogenesis by Regulating Pre-mRNA Splicing. *Cell reports*. 18:545-556.

LEGENDS TO FIGURES

Figure 1: Regulation of *MDM4* and *L22L1* pre-mRNA splicing via L22 upon inhibition of RNA polymerase I.

A.-B. RPE cells were transfected with siRNA to deplete L22 for 48 h. Treatment with 1 µM BMH-21 was performed for the indicated periods to inhibit RNA polymerase I and cause nucleolar stress. Ctrl., no BMH-21 treatment. **A.** RT-qPCR analyses using primers against the indicated sequences were performed. Expression levels were normalized to the mRNA levels of 36B4 (housekeeping gene). The average of five biological replicates is depicted. The statistical significance was assessed by an unpaired t test. *, P≤0.05; **, P≤0.01; ***, P≤0.001; ****, P≤0.0001; ns, not significant. Error bars, standard deviation of the mean. **B.** Western blot analysis of the indicated proteins was performed, with GAPDH as a sample control. **C.-H.** RPE cells were treated with 10 or 20 µM 5-FU or 1 µM BMH-21 for 24 h or 48 h to induce nucleolar stress. **C.-G.** Immunofluorescence was performed to detect L22. **C.-D.** Representative images (100x objective, bar: 40 µm) of the nuclear (DAPI, blue) and L22 (green) staining upon 24 h (**C**) and 48 h treatment (**D**). **E.-F.** Quantification of nuclear intensities of L22 signal in single cells upon 24 h (**E**) and 48 h (**F**) treatment. Red lines represent the mean fluorescence intensity values. A minimum of 180 cells were quantified for each condition. The statistical significance was assessed by a Mann-Whitney U test. **G.** Quantification of cytoplasmic versus nucleoplasmic intensity of the L22 signal. At least 190 cells were analyzed in each case. The average of three biological replicates is depicted. The statistical significance was assessed by an unpaired t test. **H.** Western blot analysis monitoring the levels of the indicated proteins upon induction of nucleolar stress by treatment with 10 or 20 µM 5-FU or 1 µM BMH-21 for 24 h. *, background band.

Figure 2: p53 activity upon nucleolar stress, dependent on L22.

A.-B. RPE cells were transfected with siRNA to deplete L22 and/or RPS13 (the latter to induce nucleolar stress), for 48h. **A.** RT-qPCR analyses of the indicated mRNAs were performed. Expression levels were normalized to the mRNA levels of 36B4. **B.** Western blot analysis of the proteins indicated was performed, using β-actin as a sample control. **C.-E.** RPE cells were transfected with siRNA to deplete L22 and/or RPS13 or RPL11 for 48h. Immunofluorescence staining of L22 was performed. **C.** Representative images (100x objective, bar: 40 µm) of the nuclear (DAPI, blue) and L22 (green) staining. **D.** Quantification of single nuclear intensities of L22 signal. Red lines represent the mean fluorescence intensity values. A minimum of 130

cells were quantified for each condition. **E.** Quantification of cytoplasmic versus nucleoplasmic intensity of L22 signal. At least 150 cells were analyzed for each condition.

Figure 3: Abolished splice regulation by L22 upon deletion of L22-binding consensus sequences from *MDM4* intron 6.

A. RNA motif that mediates the interaction with L22: a stem-loop structure with a GC base pair at the top of the stem (G on the 5' side, C on the 3' side) and U at the 3' end of the loop. The conserved bases G, C and U are marked in bold. Adapted from Dobbelstein and Shenk, 1995. **B.** Sequence of exon 6 (marked in yellow) and part of intron 6 of the *MDM4* gene. Sequences in intron 6 predicted to form L22-binding stem loops are marked in green, and the characteristic bases G, C and U (T) are indicated in bold. Red scissor symbols show cleavage sites for CRISPR/Cas9-mediated deletion of all three L22-binding stem loop sequences in MDM4 intron 6. **C.** RNA folding prediction of the last 46 bp of exon 6 and the first 268 bp of intron 6 of the MDM4 gene using the RNAfold web server (http://rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi) minimum free energy (MFE) function. Sequences predicted to form L22-binding stem loops are marked in green as in **B**. **D.-G.** RPE Cas9 or RPE MDM4 Δ intron 6 cells were transfected with siRNA to deplete L22 or RPS13 for 48h. **D.** RT-qPCR analyses using primers against the indicated sequences were performed. Expression levels were normalized to the mRNA levels of 36B4. **E.-F.** RT-qPCR analysis using primers leading to a longer product for MDM4-FL and a shorter product for MDM4-S was performed. **E.** RT-qPCR products were visualized by agarose gel electrophoresis and staining with SERVA DNA Stain Clear G. **F.** Based on band quantification, MDM4-S/MDM4-FL ratios were calculated. The average of three biological replicates is depicted. The statistical significance was assessed by an unpaired t test. **G.** Western blot analysis was performed to indicate MDM4 protein levels largely corresponding to the mRNA.

Figure 4: Splice regulation of *L22L1* and *UBAP2L* by L22.

A. RPE p53$^{-/-}$ cells were transfected with siRNA to deplete L22 or RPS13 as indicated for 48h. RT-qPCR analyses using primers against the indicated sequences were performed. Expression levels were normalized to the mRNA levels of 36B4. **B.** End of the sequence of the UBAP2L gene. The last exon, exon 27, is marked in yellow for the main *UBAP2L* variant UBAP2L-207, and in orange for the *UBAP2L* variant UBAP2L-201. Within both exons, sequences predicted to form L22-binding stem loops are marked in green, and the characteristic bases G, C and U (T) are indicated in bold. **C.** RPE p53$^{-/-}$ cells were transfected with siRNA to deplete L22 or RPS13 for 48h. Western blot analysis revealed L22-dependency

of UBAP2L protein levels. **D.** RT-qPCR analysis of the same samples as in **A** was performed, using primers leading to a longer product for L22L1 ncRNA and a shorter product for L22L1 mRNA. RT-qPCR products were visualized by agarose gel electrophoresis. **E.** Sequence of exons 2 and 3 (marked in yellow) and intron 2 of the L22L1 gene. Sequences in intron 2 predicted to form L22-binding stem loops are marked in green as in **B**. The splice acceptor site of intron 2 and an alternative splice acceptor site leading to the formation of the L22L1 non-coding RNA (ncRNA) variant are indicated in bold and orange. **F.** RNA folding prediction of the last 10 bp of exon 2, intron 2, and the first 10 bp of exon 3 of the L22L1 gene using the RNAfold web server (http://rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi) minimum free energy (MFE) function. L22-binding motifs are indicated according to the scheme used in **B** and **E**.

Figure 5: Overlapping intronic sequences required for splice regulation of MDM4 by L22 and ZMAT3.

RPE Cas9 or RPE MDM4 Δ intron 6 cells were transfected with siRNA to deplete L22 and/or ZMAT3 for 48h. **A.** RT-qPCR analyses using primers against the indicated sequences were performed. Expression levels were normalized to the mRNA levels of 36B4. **B.** RT-qPCR analysis using primers leading to a longer product for MDM4-FL and a shorter product for MDM4-S was performed. RT-qPCR products were visualized by agarose gel electrophoresis. **C.** Western blot analysis of the proteins indicated was performed, using GAPDH as a sample control.

Figure 6: L22 binding to consensus sequences in intron 6 of *MDM4* pre-mRNA.

A. A splicing reporter assay based on Fukushima et al., 2020, was used to investigate the sequence requirements needed for L22 to bind MDM4 pre-mRNA and regulate its splicing. To generate the control plasmid FMv2, the following sequences were inserted into the pcDNA3.1/Hygro(+) vector containing the CMV promoter by using the restriction enzymes *Nhe*I and *Xho*I: the mCherry coding sequence, the 5′ end of the eGFP gene (5′-eGFP), the 5′ end of DMD intron 18 (5′ DMD intron 18), a Multiple Cloning Site (MCS), the 3′ end of DMD intron 19 (3′ DMD intron 19), and the 3′ end of the eGFP gene (3′-eGFP). To generate the MDM4 splicing reporter plasmid FMv2-MDM4, the following sequences were inserted into the MCS of FMv2 by using the restriction enzymes *Hind*III and *Bam*HI: the 3′ end (34 bp) of MDM4 intron 5 (3′ MDM4 intron 5), MDM4 exon 6 (68 bp), and the 5′ end (398 bp) of MDM4 intron 6 (5′

MDM4 intron 6) containing all three previously identified L22 consensus sequences. Upon transfection of either FMv2 or FMv2-MDM4, mCherry is constitutively expressed and serves as a transfection marker. eGFP is constitutively expressed upon FMv2 overexpression, but only if *MDM4* exon 6 is skipped in case of FMv2-MDM4 overexpression, therefore serving as a splicing marker. **B.-C.** FMv2-MDM4 versions with either deletion of all three L22 consensus sequences in MDM4 intron 6 (FMv2-MDM4_del_1,2,3) or with mutations of the three conserved nucleotides of the single or combined L22 consensus sequences as indicated (FMv2-MDM4_mut_[…]) were generated by site-directed mutagenesis. H1299 cells were transfected with siRNA to deplete L22 for 72h, and with FMv2, FMv2-MDM4 and the different mutated versions of FMv2-MDM4, at 24 hrs after the siRNA transfection, followed by incubation for 48h. Red fluorescence (mCherry) and green fluorescence (eGFP) were detected. **B.** The eGFP fluorescence intensity among cells with high mCherry intensity (>20 cells per sample) was quantified, and the change of the eGFP/mCherry ratio with L22 compared to control depletion was visualized. The average of three biological replicates is depicted. The statistical significance was assessed by an unpaired t test. **C.** Images (5x objective, bar: 800 µm) of the eGFP (green) and mCherry (red) staining. **D.** HEK293T cells were transfected with pCMV6Entry or pCMV6Entry-L22 and with either FMv2, FMv2-MDM4 or FMv2-MDM4_mut_1,2,3 for 42h, followed by RNA immunoprecipitation (RIP) using FLAG beads to precipitate tagged L22. RT-qPCR analyses using primers against the indicated sequences were performed for the immunoprecipitation (IP) and 5% input samples shown, and IP signal was normalized to input signal.

Figure 7: Resistance towards 5-fluorouracil upon abrogation of MDM4 splicing regulation by L22.

A.-D. RPE Cas9 (dark) or RPE MDM4 Δ intron 6 (light) cells were treated with 5-fluorouracil (5-FU) or control-treated (ctrl) as indicated for 20 days. Every other day, the treatment was refreshed and cell confluence was measured. **A.-C.** For each condition, results of three biological replicates (with three technical replicates each) are depicted. **D.** Statistical analysis was performed by comparing the area under the curve (AUC) values, which were calculated from the confluence graphs (A-C), of both cell lines using an unpaired t test. **E.-G.** RPE Cas9 or RPE MDM4 Δ intron 6 cells were treated with 5-FU as in A-D. After 10 days of treatment, cells were harvested for RNA isolation. **E.** RT-qPCR analyses to quantify mRNA levels corresponding to the indicated genes were performed, normalized to 36B4. **F.-G.** RT-qPCR analysis using primers leading to a longer product for MDM4-FL and a shorter product for MDM4-S was performed. **F.** RT-qPCR products were visualized by agarose gel

electrophoresis. **G.** Based on band quantification, the MDM4-S/MDM4-FL ratio was calculated. **H.** L22, by promoting alternative MDM4 splicing and thus p53-mediated gene expression, translates nucleolar stress into growth arrest.

LEGENDS TO SUPPLEMENTARY FIGURES

Supplementary Figure 1: Gene expression upon depletion of L22 and L22L1, as well as MDM4 and MDM2.

RPE cells were transfected with siRNAs to deplete L22, L22L1, MDM2, or MDM4 for 48 h. RT-qPCR analyses to detect the indicated target mRNAs were performed. Expression levels were normalized relative to the mRNA levels of 36B4 (housekeeping gene) in the control sample. The average of three biological replicates is depicted. Error bars, standard deviation of the mean. The statistical significance was assessed by an unpaired t test. *, P≤0.05; **, P≤0.01; ***, P≤0.001; ****, P≤0.0001; ns, not significant.

Supplementary Figure 2: L22-FLAG accumulation in the nucleoplasm upon treatment with BMH-21 or 5-FU.

A.-B. Stably transfected HEK293 Flp-In cells for the inducible expression of L22-FLAG were treated with 10 or 20 µM 5-FU or 1 µM BMH-21, for 24 h or 48 h, to induce nucleolar stress. Tetracycline (1 µg/mL) was added for all conditions, including the control, during the entire treatment duration (24 h or 48 h), to induce the expression of FLAG-tagged L22. Immunofluorescence staining of L22 using a FLAG antibody was performed. Representative images (100 objective, bar: 40 µm) of the nuclear (DAPI, blue) and L22 (green) staining upon 24 h (**A**) and 48 h treatment (**B**).

Supplementary Figure 3: *MDM4* exon 6 inclusion upon depletion or overexpression of L22 or L22L1 in HEK293T cells.

A.-B. HEK293T cells were transfected with siRNA to deplete L22 or L22L1 for 48 h. **C.-D.** HEK293T cells were transfected with plasmids to overexpress L22 or L22L1 for 48 h. **A., C.** RT-qPCR analyses to detect the indicated target mRNAs were performed, with normalization to the 36B4 mRNA level in the control sample. **B., D.** Western blot analyses of the proteins indicated were performed, using GAPDH (**B)** and β-actin (**D**) as loading controls.

Supplementary Figure 4: p53 activity as a function of L22 upon nucleolar stress.

A.-B. RPE cells were transfected with siRNA to deplete L22 and/or RPL5 or RPL11 for 48h (the latter two to induce nucleolar stress). **A.** RT-qPCR analyses with normalization to 36B4. **B.** Western blot analyses. Note that the depletion of L5 also diminishes the levels of L11 and vice versa, presumably due to mutual stabilization of the two proteins within the 5S RNP. **C.-E.** RPE cells were transfected with siRNA to deplete L22 and/or RPS13 or RPL11 for 72h. Immunofluorescence staining of L22 was performed. **C.** Images (100x objective, bar: 40 μm) of the nuclear (DAPI, blue) and L22 (green) staining. **D.** Quantification of single nuclear intensities of the L22 signal. Red lines represent the mean fluorescence intensity values. A minimum of 90 cells were quantified for each condition. **E.** Quantification of cytoplasmic versus nucleoplasmic intensity of L22 signal. More than 100 cells were analyzed for each condition.

Supplementary Figure 5: L22 localization to the nucleoplasm upon depletion of RPS13 or RPL11.

A.-B. HEK293 eL22(C) cells were transfected with siRNA to deplete L22. To cause nucleolar stress, RPS13 or RPL11 were depleted in parallel, for 48h or 72h. 24h after seeding and reverse siRNA transfection, i.e. for a total duration of 24h respectively 48h, tetracycline (1 μg/mL) was added for all conditions including the control, to induce the expression of FLAG-tagged L22. Immunofluorescence staining of L22 using a FLAG antibody was performed. Representative images (100x objective, bar: 40 μm) of the nuclear (DAPI, blue) and L22 (green) staining upon 48h (**A**) and 72h knockdown (**B**).

Supplementary Figure 6: Conserved L22-binding sequences in *MDM4* intron 6 sustaining p53 activity.

A. Sequence of exon 6 (marked in yellow) and part of intron 6 of the human and murine *MDM4* genes. Sequences in intron 6 predicted to form L22-binding stem loops are marked in green, and the characteristic bases G, C and U (T) are indicated in bold. Nucleotide differences in the murine compared to the human gene are marked in grey. **B.** Genotyping PCR products of RPE Cas9 and RPE MDM4 Δ intron 6 cells were sequenced using a reverse sequencing primer binding at the 3′ end of the reverse genotyping primer. The following sequences are marked as also indicated in the figure: 3′ end of *MDM4* exon 6 (yellow frames), upstream (us) and downstream (ds) crRNA binding sites (RPE Cas9) respectively parts thereof (RPE MDM4 Δ

intron 6) (red frames), protospacer-adjacent motif (PAM) sequences (black frames), sequences predicted to form L22-binding stem loops (green frames). In addition, the binding site of the upstream (forward) genotyping primer respectively its 3′ end is marked with a dark blue line. **C.-D.** RPE Cas9 or RPE MDM4 Δ intron 6 cells were transfected with siRNA to deplete L22 or RPS13 for 48h. **C.** RT-qPCR analyses with normalization to 36B4. **D.** Western blot analysis.

Supplementary Figure 7: Confirmation of L22 depletion, overexpression and immunoprecipitation.

A. H1299 cells were transfected with siRNA to deplete L22 for 72h, and with FMv2, FMv2-MDM4 and the different mutated versions of FMv2-MDM4, at 24 hrs after the siRNA transfection, followed by incubation for 48h. Western blot analysis was performed to confirm L22 depletion, using GAPDH as a loading control. **B.-C.** HEK293T cells were transfected with pCMV6Entry or pCMV6Entry-L22 and with either FMv2, FMv2-MDM4 or FMv2-MDM4_mut_1,2,3 for 42h, followed by RNA immunoprecipitation (RIP) using FLAG beads to precipitate tagged L22. Overexpression (**B**) and precipitation (**C**) of L22 were confirmed by Western blot analysis.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

FMv2

FMv2-MDM4

5′-eGFP   5′ DMD intron 18   5′ MDM4 intron 6   3′ DMD intron 19   3′-eGFP

3′ MDM4 intron 5   Multiple Cloning Site (MCS)

MDM4 exon 6

B

eGFP/mCherry

reporter plasmid: FMv2, FMv2-MDM4, del_1,2,3, mut_1, mut_2, mut_3, mut_1+2, mut_1+3, mut_2+3, mut_1,2,3

FMv2-MDM4 with deletions or mutations of L22 motifs in intron 6

C

D

MDM4 intron 6 downstream of L22 motifs

FMv2 3′ eGFP

Plasmid: FMv2-MDM4, FMv2-MDM4_mut_1,2,3, FMv2-MDM4, FMv2-MDM4_mut_1,2,3

pCMV6Entry   pCMV6Entry-L22

Plasmid: pCMV6Entry, pCMV6Entry-L22

FMv2

Figure 6

Figure 7

Relative mRNA levels

MDM4-FL

Puma

MDM4-S

p53

p21

L22L1

MDM2

L22

siRNA: ctrl#1, L22#1, L22L1#1, MDM2#1, MDM4#1

siRNA: ctrl#1, L22#1, L22L1#1, MDM2#1, MDM4#1

Supplementary Figure 1

A

24h

	Ctrl	10 µM 5-FU	20 µM 5-FU	1 µM BMH-21

DAPI

FLAG (L22)

B

48h

	Ctrl	10 µM 5-FU	20 µM 5-FU	1 µM BMH-21

DAPI

FLAG (L22)

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

A

48h

siRNA: ctrl#1   L22#1   RPS13#1   L22#1 +   RPL11#1   L22#1 +
                                      RPS13#1   RPL11#1

DAPI

FLAG
(L22)

B

72h

siRNA: ctrl#1   L22#1   RPS13#1   L22#1 +   RPL11#1   L22#1 +
                                      RPS13#1   RPL11#1

DAPI

FLAG
(L22)

<u>**Supplementary Figure 5**</u>

A

B

C

<u>**Supplementary Figure 7**</u>