Identification of SLC25A46 interaction interfaces with

mitochondrial membrane fusogens Mfn2 and Opa1

Sivakumar Boopathy[1,2], Camila Makhlouta Lugo[1*], Bridget E. Luce[1], Julie McDonald[1#],

Pusparanee Hakim[1], Jackeline Ponce[3], Beatrix M. Ueberheide[3,4], Luke H. Chao[1,2,5]

[1]Department of Molecular Biology, Massachusetts General Hospital, Boston MA 02114,

USA

[2]Department of Genetics, Harvard Medical School, Boston MA 02115, USA

3 Proteomics Resource Center, Division of Advanced Research Technologies, New

York University Langone Health Center, New York NY 10016, USA

[4]Department of Biochemistry and Molecular Pharmacology, New York University

Langone Health Center, New York NY 10016, USA

[5]Corresponding author

*Present address: Program in Neurobiology, University of Chicago, Chicago IL 60637,

USA

#Present address: Department of Biology, Massachusetts Institute of Technology,

Cambridge MA 02139, USA

Running Title

SLC25A46 interaction interfaces with Mfn2 and Opa1

Keywords

1. Mitochondria

2. Membrane fusion

3. GTPase

4. Structural model

5. Protein cross-linking

6. Mass spectrometry

7. Protein-protein interaction

8. Mitochondrial solute carrier

Abstract

Mitochondrial fusion requires the sequential merger of four bilayers to two. The outer-membrane solute carrier protein SLC25A46 interacts with both the outer and inner-membrane dynamin family GTPases Mfn1/2 and Opa1. While SLC25A46 levels are known affect mitochondrial morphology, how SLC25A46 interacts with Mfn1/2 and Opa1 to regulate membrane fusion is not understood. In this study, we use crosslinking mass-spectrometry and AlphaFold 2 modeling to identify interfaces mediating a SLC25A46-Opa1-Mfn1/2 complex. We reveal that the bundle signaling element of Opa1 interacts with SLC25A46, and the helical repeat 1 region of Mfn2 interacts with the SLC25A46 N-terminus. We validate these newly identified interaction interfaces and show that they play a role in mitochondrial network maintenance.

Introduction

Mitochondrial fission and fusion are membrane remodeling processes required for maintenance of a healthy mitochondrial reticulum (1, 2), which regulates the number, and distribution of mitochondrial DNA and respiratory chain proteins (1-5). A distinct feature of mitochondrial fusion is the sequential fusion of two pairs of bilayers. Key questions regarding the spatiotemporal control of these events remain. Specifically, how are the activities of the outer and inner membrane fusion GTPases, Mfn1/2 and Opa1 respectively, coordinated and regulated?

SLC25A46 interacts with both fusion GTPases and has emerged as an inter-membrane bridging protein which plays roles in regulating mitochondrial membrane dynamics (6-9). Mutations in SLC25A46 result in optic atrophy spectrum disorder, Leigh syndrome, and pontocerebellar hypoplasia (10-24). SLC25A46 is a member of the solute carrier family 25 of mitochondrial transporters, which transport substrates such as nucleotides, vitamins, amino acids, fatty acids across the inner mitochondrial membrane (IMM) (25). SLC25A46 resides in the outer mitochondrial membrane (OMM), lacks the signature sequence motifs of the SLC25 family (25), does not have an identified substrate and contains a distinct N-terminal extension.

In SLC25A46 knockout and knockdown models of human neurological disease, mitochondria are often enlarged and aggregated in nervous tissues, exhibiting detached, vesiculated cristae (21, 24, 26-29). Respiratory chain complex activity is impaired in these models (26, 27), and ATP levels are reduced (27, 28). In human cell lines, knockdown of SLC25A46 induces a tubular mitochondrial network whereas overexpression results in mitochondrial fragmentation (6, 9, 24), suggesting a role in downregulating fusion. The phenotypes of SLC25A46 knockout varied depending on the cell lines tested, displaying mitochondrial fragmentation in human fibroblasts and HeLa cells (8), and hyper-filamentous network in other cell lines (30, 31). Knockout of the yeast orthologoue Ugo1 (~22% sequence identity) also induced fragmentation (32-34). These observations argue for a role of SLC25A46 in regulating mitochondrial fusion where SLC25A46 levels at the OMM are critical. In fibroblasts and iPSC-derived cerebral neurons, SLC25A46 is focally present at tips and branches of mitochondrial tubules and at sites of fusion and fission (8). Interestingly the SLC25A46 puncta were

proximal to Opa1 in human fibroblasts (8), suggesting spatially-defined interactions with
SLC25A46 play a role in fusion regulation.

Understanding the molecular details of the SLC25A46-Mfn2-Opa1 interactions
can provide constraints on how SLC25A46 organizes sites for mitochondrial membrane
remodeling and inform mechanisms for coordinated OMM/IMM fusion. In yeast, Ugo1
physically interacts with Fzo1 and Mgm1, the orthologous fusion GTPases (34-37).
Previous studies indicate Opa1 and Mfn2 interacts with SLC25A46 (6-9), however the
specific interface mediating these interactions was undefined. In this study, we used
crosslinking-mass spectrometry and AlphaFold 2 modeling to identify the protein-protein
interaction surfaces of SLC25A46-Opa1 and SLC25A46-Mfn2. We validated this newly
identified SLC25A46-Opa1 interface by mutagenesis, and demonstrate a role for this
interface in mitochondrial network maintenance.

Results

SLC25A46 interacts with Opa1 via its bundle signaling element.

Previous studies used crosslinking, immunoprecipitation and mass spectrometry
to identify an interaction network of Opa1, Mfn1/2 and SLC25A46 (6-9). We over-
expressed human SLC25A46 with Opa1 in *Pichia pastoris* under the control of a strong
promoter (alcohol oxidase) (**Figure 1A**), and found that detergent (GDN)-solubilized
SLC25A46 in *P. pastoris* lysate interacts with column-immobilized Opa1 from lysate in
the absence of crosslinkers (**Figure 1B**). Likewise, column-immobilized SLC25A46 was
also found to interact with Opa1 (**Supporting Figure 1A**).

To identify the specific interface responsible for the SLC25A46-Opa1 interaction,
we crosslinked the purified complex for analysis by mass spectrometry. Using
bis(sulfosuccinimidyl)suberate (BS3) we identified a crosslink between K241 of
SLC25A46 and K568 of Opa1 bundle signaling element (BSE) helix 2. Intralinks
consistent with folded Opa1 secondary structure elements were identified (**Supporting**
Figure 1B). A second crosslinker Bis-N-succinimidyl-(pentaethylene glycol)
(BS(PEG)5) with a larger spacer length also crosslinked K241(SLC25A46)-K568(Opa1),
and identified a second crosslink between K256 of SLC25A46 and the K960 of Opa1
BSE helix 3 (**Figure 1C**). We modeled this interaction using Alphafold 2 (38, 39), which

consistently revealed a high-confidence and robust interface mediated by the Opa1
BSE, with a topology consistent with the proteins' sub-organellar localization (6, 21, 24,
40, 41) (**Figure 1D, Supporting Figure 1C**).

Validation of SLC25A46:Opa1-BSE interaction

Examination of the SLC25A46-Opa1 interface in the AlphaFold 2 model revealed
two salt bridges comprising residues R257 (SLC25A46) and E561 (Opa1); R347
(SLC25A46) and D565 (Opa1). Adjacent to the salt bridges, a cluster of hydrophobic
residues, I229, I230, I349, Y357, V359, L360 in SLC25A46 and A569, F572 and T576 in
Opa1 BSE helix 2 appear to further stabilize the interaction (**Supporting Figure 2A**).
We tested this newly identified SLC25A46-Opa1 interface by mutating the salt bridges
and hydrophobic site residues in SLC25A46, and evaluating if the interaction with Opa1
was maintained (**Figure 2A**). While recombinantly expressed and purified WT
SLC25A46 bound column-immobilized Opa1 (consistent with observations in lysates),
both the I349D mutation and the R257A/R347A double mutation diminished the
interaction (**Figure 2B and Supporting Figure 2B**).

The BSE is essential for transducing GTP-hydrolysis stimulated activities of
dynamin related protein (42, 43). In dynamin, the BSE frequently moves closer to the
GTPase upon GTP hydrolysis (44-46). To determine if the nucleotide-state of Opa1
affects its interaction with SLC25A46, we incubated the lysates with GTPγS, GMPPCP
and GDP prior to affinity capture on StrepTactin resin. We found that the presence of
nucleotide or nucleotide analogues did not affect co-elution, indicating nucleotide-
coupled BSE rearrangement does not affect Opa1-SLC25A46 interaction (**Supporting
Figure 2C**).

We also generated a minimal construct consisting of the GTPase domain and the
BSE alone (Opa1(MGD)), to test that the "paddle" and stalk self-assembly regions are
dispensable for the Opa1-SLC25A46 interaction (**Figure 3A**). SLC25A46 bound
Opa1(MGD) as shown by the co-elution of the two proteins, indicating an Opa1(MGD)
construct is sufficient for binding (**Figure 3B**). We found the same two crosslinks
observed for Opa1-SLC25A46 between Opa1(MGD) and SLC25A46. In addition, we

observed a third crosslink between K387 of SLC25A46 and K568 of Opa1 (**Figure 3C**
and 3D).

The cytosolic unstructured domain of SLC25A46 interacts with the HR1 of Mfn2.

We next investigated if SLC25A46 also interacts with the outer membrane fusion
protein Mfn2. We co-expressed the two proteins in *P. pastoris* (**Supporting Figure 3A**)
and subjected the GDN-solubilized lysates to column-binding experiments. SLC25A46
co-elutes upon elution of column-immobilized Mfn2 (**Supporting Figure 3B**), and Mfn2
co-elutes with resin bound SLC25A46 (**Supporting Figure 3C**). To test the SLC25A46-
Mfn2 interaction was nucleotide-state dependent, we treated lysates with GTPγS,
GMPPCP or GDP during detergent extraction. The presence of these nucleotides did
not enhance or diminish interaction of Mfn2 with SLC25A46 (**Supporting Figure 3H**).

To obtain insights into the SLC25A46-Mfn2 interaction, we performed
crosslinking and mass-spectrometry of the purified complex. Two crosslinks were
identified – between K60 residue of SLC25A46 and residues K416 and K420 of Mfn2
(**Supporting Figure 3D**). K60 of SLC25A46 is part of the first ~83 residues that are
predicted to be disordered in the cytosol (**Supporting Figure 3F**). Residues K416 and
K420 are part of the cytosol facing heptad repeat 1 (HR1) of Mfn2 (**Supporting Figure
3F**). The AlphaFold 2 predicted interface was low confidence for SLC25A46-Mfn2
complex, particularly for the N-terminal region of SLC25A46.

Since crosslinks were observed at K60 in SLC25A46, we tested the role of these
N-terminal residues in binding Mfn2. We deleted the first 83 amino acids of SLC25A46
termed (Δ2-83)SLC25A46, and co-expressed this protein with Mfn2 in *P. pastoris*.
Interestingly, (Δ2-83)SLC25A46 still co-eluted with Mfn2 in the affinity capture
experiments (**Supporting Figure 3G**).

Disease-causing SLC25A46 mutants do not affect interaction with Opa1 or Mfn2.

A cluster of SLC25A46 disease mutations face the IMS (**Figure 4A**). While many
mutations were reported to destabilize the protein, certain disease-causing mutations
(T142I and R340C) cause a hyper-fused mitochondrial phenotype (6, 8, 9, 15, 24). We
co-expressed these SLC25A46 mutants with Opa1, and found these disease mutants

still bound Opa1 (**Figure 4B**). We note that in these experiments we were unable to
achieve similar expression levels of the various proteins. We also saw no effect of
patient-derived mutations on the Mfn2-SLC25A46 interaction (**Supporting Figure 3G**).
170

Mutations disrupting SLC25A46-Opa1 interaction alter mitochondrial morphology.

Previous studies showed that the pathogenic mutants of SLC25A46 that are
unstable and express poorly in cells (6, 8, 9, 15, 21) induce a hyper-filamentous
phenotype similar to that observed with siRNA/shRNA knockdown of SLC25A46 (6, 9,
24). To revisit mitochondrial phenotypes upon SLC25A46 loss, we imaged
mitochondrial networks in the human colorectal carcinoma cell line HCT116. Loss of
SLC25A46 did not affect mitochondrial morphology in HCT116 cells, but upon serum
starvation, a fragmented mitochondrial network was observed. Reintroduction of WT
SLC25A46 rescued this phenotype (**Figure 5, Supporting Video 1**). This phenotype is
consistent with previous observations in human fibroblasts and HeLa cells (8). We
found that expression of R340C or R257Q SLC25A46 mutants induced interconnected
mitochondria, also in agreement with previous studies (8, 24) (**Figure 5B, Supporting
Video 1**). The mitochondrial morphology in G249D-expressing cells was similar to WT
(**Figure 5, Supporting Video 1**). Expression levels of R340C and G249D mutants
never reached WT-like levels (**Figure 5A**), likely due to protein instability (6, 8, 9, 15,
21).

We next tested the effects of the newly identified Opa1-SLC25A46 interactions
on mitochondrial morphology. Expression of I349D SLC25A46 (disrupts a hydrophobic
interface between the Opa1 BSE) produced shorter mitochondrial tubules compared to
WT, whereas R257A/R347A mutant (disrupt SLC25A46-Opa1 salt bridge interactions)
resulted in more fragmented mitochondria (**Figure 5B, Supporting Video 1**). These
results suggest that the identified SLC24A45-Opa1 interaction interface is important for
maintaining a tubular mitochondrial morphology. Expression of the (Δ2-83)SLC25A46
resulted in an increase in short mitochondrial tubules (**Figure 5B, Supporting Video 1**).
In these experiments, we did not observe changes in levels of Mfn2 or Opa1 due to
SLC25A46 knockout or expression of varying levels of mutant SLC25A46 (**Figure 5A**).

We could not evaluate the expression level of (Δ2-83)SLC25A46 due to the inability of
commercial antibodies to detect this mutant (**Figure 5A**).

Discussion

Mitochondria undergo morphological changes tightly coupled to cellular
physiology (4). While the identity of the GTPases responsible for sequential
mitochondrial fusion have long been identified (47), and mechanisms for individual
fusion steps have been described (48-51), questions remain regarding how outer and
inner-membrane activities are coordinated and regulated. Emerging evidence points to
SLC25A46 as a regulator of fusion activity, with protein levels influencing the fusion-
fission balance (6, 8, 24). Consistent with this role, SLC25A46 is observed to localize to
the fusion sites (8).

In this study, we investigate the physical interaction between SLC25A46 and the
membrane fusogens Opa1 and Mfn1/2. Combining crosslinking mass-spectrometry, and
AlphaFold 2 modeling, we identify the BSE region in Opa1 interacting with the IMS-face
of SLC25A46. We observe that this interaction is nucleotide state independent. We also
describe interactions between SLC25A46 N-terminal region and the HR1 region of
Mfn2. The crosslinked residues K416 and K420 are conserved in Mfn1, suggesting the
interaction we see for Mfn2 is also possible for Mfn1.

Our pull-down experiments showed SLC25A46-Mfn2 interaction was maintained
despite deletion of the N-terminal region of SLC25A46. In yeast, residues 1-294 of Ugo1
interacts with Fzo1 in a region comprising the transmembrane segments, residues N-
terminal to the first transmembrane helix and the C-terminal HR2 (34). From these
observations, we speculate that the region downstream of residue 83 in SLC25A46 also
interacts with Mfn2 and may involve transmembrane-transmembrane interactions.

Reports from several studies indicate that mitochondrial morphology is sensitive
to SLC25A46 levels. Schuettpelz *et al.* reported mitochondrial fragmentation in human
fibroblasts and HeLa cells upon SLC25A46 KO (8). In SLC25A46 KO HCT116 cells, we
observed mitochondrial fragmentation upon serum starvation. In addition, we observed
a ~2-fold slower growth of this cell line, consistent observations in SLC25A46 deleted
fibroblasts deletion (8). Overexpression of SLC25A46 in all tested cell lines caused

mitochondrial fragmentation (6, 24, 31). The overexpression findings suggest that at
high levels, SLC25A46 can inhibit the activity of the protein fusogens. One possible
mode of inhibitory action may be SLC25A46 binding would be sufficient to result in
steric competition for Opa1 self-assembly, which relies on the BSE and stalk regions to
mediate formation of helical filaments (52-54). Consistent with this prediction, levels of
Opa1 oligomeric complexes were found to increase upon loss of SLC25A46 (8).

In our study, SLC25A46 mutants targeting Opa1 interaction did not promote a
hyperfused phenotype, but instead disrupted the mitochondrial network. This suggests
that a SLC25A46-Opa1 interaction promotes fusion necessary for a tubular network.
Although we could not estimate the relative levels of (Δ2-83)SLC25A46, this N-terminal
deletion reduced the filamentous network, while increasing the proportion of short
tubules, also suggesting a fusion defect. Taken together, these observations provide
evidence for SLC25A46 playing an activating role in fusion when expressed at
physiological levels.

Our results provide a basis for future functional analyses of the SLC25A46
interaction with the fusion machinery. SLC25A46 protein-protein interactions also may
regulate cristae morphology, which is dramatically affected upon loss of SLC25A46 (6,
8, 24, 26, 27, 29). Proteomics studies have also implicated SLC25A46 endoplasmic
reticulum interaction partners in maintaining ER-mitochondria contacts facilitating lipid
flux (6-8). Future mechanistic structure-activity studies are necessary to explore these
other roles of SLC25A46 in cristae regulation and lipid metabolism.

Experimental procedures

Plasmids

Genes for SLC25A46, Mfn2, Opa1(253-960), Opa1(MGD) constructs were
synthesized (GenScript) and cloned into pPICZ A for *P. pastoris* expression. C-terminus
of SLC25A46 was fused to a $(G_4S)_3$ linker followed by TEV cleavage site and deca-
histidine. N-terminus of Opa1(253-960), Opa1(263-960), Opa1(MGD) was tagged with
Twin-Strep tag followed by HRV 3C cleavage site and $(G_4S)_3$ linker. C-terminus of Mfn2
was fused to G_3S linker, HRV 3C cleavage site and Twin-Strep tag. For co-expression,
SLC25A46 and either Mfn2 or Opa1 with tags with its own AOX1 promoter and
terminator were cloned into the same plasmid. Strep-Opa1(253-960) and Strep-
Opa1(MGD) were also cloned into pET-28a(+) vector for expression in *E. coli*.

Generation of *Pichia* strains

Expression strains were generated from *Pichia pastoris* strain SMD1163 (*his4*
pep4 prb1) (kind gift from Dr. Tom Rapaport (Harvard Medical School) according to
established protocols (55). Plasmids were transformed into SMD1163 by electroporation
and grown on YPDS (1% yeast extract, 2% peptone, 2% dextrose, 1 M sorbitol) zeocin
(InvivoGen, ant-zn-1p) plates. 7-14 resulting colonies were grown in BMGY (1% yeast
extract, 2% peptone, 100 mM potassium phosphate pH 6.0, 1.34% yeast nitrogen base,
4×10^{-5}% biotin, 1% glycerol) and induced in BMMY (replaced glycerol with 0.5%
methanol in BMGY). Cultures were screened for protein expression by Western using
anti-His, anti-Strep, anti-SLC25A46, anti-Opa1 or anti-Mfn2 antibodies.

Binding experiments using lysates

P. pastoris (1 L) expressing SLC25A46, Opa1, Mfn2 alone or in combination
were harvested and cryo-milled using SPEX SamplePrep 6875D Freezer/Mill. For
StrepTactin column-binding and elution, milled cells (125 mg) was resuspended in 3 ml
of 25 mM HEPES, 150 mM NaCl, pH 8.0 buffer containing 1% GDN and protease
inhibitors, tumbled for 1-1:30 h at 4 °C. Suspension was centrifuged at 20,000 for 30
min at 4 °C. Supernatant (100 µl) was incubated with 50 µl of StrepTactin XT 4Flow

high-capacity resin in a spin column (Abcam, ab286861) at 4°C for 15 min followed by

centrifugation at 500 xg for 30 s. Resin was washed 3x with 100 µl buffer containing

0.1% GDN. To elute, the resin was incubated with 100 µl buffer with 50 mM biotin for 15

min and centrifuged. Same procedure for Ni-NTA binding experiments except that the

binding and elution buffer contained 80 mM and 500 mM imidazole, respectively.

AlphaFold 2 modeling

Protein sequences for human SLC25A46, Opa1 and Mfn2 were run through in a

AlphaFold2 Google ColabFold – v 1.5.3 workbook (38, 39, 56). Five independent runs

generated the top three ranked relaxed models for SLC25A46-Opa1 and SLC25A46-

Mfn2 models respectively.

Data availability

Mass spectrometry data available on request.

Supporting information

This article contains supporting information.

Acknowledgements

We thank Dr. Yuan Gao and Dr. Tom Rapoport for providing *Pichia pastoris* strain SMD1163. We thank all the members of the Chao lab for critical feedback.

Funding and additional information

Mass spectrometric experiments supported by a shared instrumentation grant from the NIH, 1S10OD010582-01A1 towards purchase of Orbitrap Eclipse. This work was supported by funding from the National Institutes of Health (R35GM142553 to L.H.C.)

References

References

1.   Wai, T., andLanger, T. (2016) Mitochondrial Dynamics and Metabolic Regulation
Trends Endocrinol Metab **27**, 105-117 10.1016/j.tem.2015.12.001

2.   Yapa, N. M. B., Lisnyak, V., Reljic, B., andRyan, M. T. (2021) Mitochondrial
dynamics in health and disease FEBS Lett **595**, 1184-1204 10.1002/1873-
3468.14077

3.   Giacomello, M., Pyakurel, A., Glytsou, C., andScorrano, L. (2020) The cell
biology of mitochondrial membrane dynamics Nat Rev Mol Cell Biol **21**, 204-224
10.1038/s41580-020-0210-7

4.   Chan, D. C. (2020) Mitochondrial Dynamics and Its Involvement in Disease Annu
Rev Pathol **15**, 235-259 10.1146/annurev-pathmechdis-012419-032711

5.   Meyer, J. N., Leuthner, T. C., andLuz, A. L. (2017) Mitochondrial fusion, fission,
and mitochondrial toxicity Toxicology **391**, 42-53 10.1016/j.tox.2017.07.019

6.   Janer, A., Prudent, J., Paupe, V., Fahiminiya, S., Majewski, J., Sgarioto, N. *et al.*
(2016) SLC25A46 is required for mitochondrial lipid homeostasis and cristae
maintenance and is responsible for Leigh syndrome EMBO Mol Med **8**, 1019-
1038 10.15252/emmm.201506159

7.   Perivolidi, V. I., Violitzi, F., Ioannidou, E., Rinotas, V., Stamatakis, G., Samiotaki,
M. *et al.* (2022) Proteomic Identification of the SLC25A46 Interactome in
Transgenic Mice Expressing SLC25A46-FLAG J Proteome Res **21**, 375-394
10.1021/acs.jproteome.1c00728

8.   Schuettpelz, J., Janer, A., Antonicka, H., andShoubridge, E. A. (2023) The role of
the mitochondrial outer membrane protein SLC25A46 in mitochondrial fission
and fusion Life Sci Alliance **6**, 10.26508/lsa.202301914

9.   Steffen, J., Vashisht, A. A., Wan, J., Jen, J. C., Claypool, S. M., Wohlschlegel, J.
A. *et al.* (2017) Rapid degradation of mutant SLC25A46 by the ubiquitin-
proteasome system results in MFN1/2-mediated hyperfusion of mitochondria Mol
Biol Cell **28**, 600-612 10.1091/mbc.E16-07-0545

10. Kodal, L. S., Hammer-Hansen, S., Holm-Yildiz, S., Gronskov, K., Karstensen, H.
G., andDysgaard, T. (2022) Case report: A novel variant in SLC25A46 causing
sensorimotor polyneuropathy and optic atrophy Front Neurol **13**, 1066040
10.3389/fneur.2022.1066040

11. Yamada, M., Suzuki, H., Adachi, H., Noguchi, A., Miya, F., Takahashi, T. *et al.*
(2022) Diagnosis of SLC25A46-related pontocerebellar hypoplasia in two siblings
with fulminant neonatal course: role of postmortem CT and whole genomic
analysis: a case report BMC Neurol **22**, 20 10.1186/s12883-021-02540-x

12. Li, Q., Madden, J. A., Lin, J., Shi, J., Rosen, S. M., Schmitz-Abe, K. *et al.* (2021)
Reanalysis of Exome Data Identifies Novel SLC25A46 Variants Associated with
Leigh Syndrome J Pers Med **11**, 10.3390/jpm11121277

13. Raju, S., Medarametla, S., andBoraiah, N. (2021) Dystonia and Hereditary Motor
Sensory Neuropathy 6B Due to SLC25A46 Gene Mutations Mov Disord Clin
Pract **8**, 480-482 10.1002/mdc3.13139

14. Bitetto, G., Malaguti, M. C., Ceravolo, R., Monfrini, E., Straniero, L., Morini, A. *et*
al. (2020) SLC25A46 mutations in patients with Parkinson's Disease and optic
atrophy Parkinsonism Relat Disord **74**, 1-5 10.1016/j.parkreldis.2020.03.018

15. Abrams, A. J., Fontanesi, F., Tan, N. B. L., Buglo, E., Campeanu, I. J., Rebelo,
A. P. *et al.* (2018) Insights into the genotype-phenotype correlation and molecular
function of SLC25A46 Hum Mutat **39**, 1995-2007 10.1002/humu.23639
16. Ivanov, I., Atkinson, D., Litvinenko, I., Angelova, L., Andonova, S., Mumdjiev, H.
et al. (2018) Pontocerebellar hypoplasia type 1 for the neuropediatrician:
Genotype-phenotype correlations and diagnostic guidelines based on new cases
and overview of the literature Eur J Paediatr Neurol **22**, 674-681
10.1016/j.ejpn.2018.03.011
17. Braunisch, M. C., Gallwitz, H., Abicht, A., Diebold, I., Holinski-Feder, E., Van
Maldergem, L. *et al.* (2018) Extension of the phenotype of biallelic loss-of-
function mutations in SLC25A46 to the severe form of pontocerebellar hypoplasia
type I Clin Genet **93**, 255-265 10.1111/cge.13084
18. van Dijk, T., Rudnik-Schoneborn, S., Senderek, J., Hajmousa, G., Mei, H., Dusl,
M. *et al.* (2017) Pontocerebellar hypoplasia with spinal muscular atrophy (PCH1):
identification of SLC25A46 mutations in the original Dutch PCH1 family Brain
140, e46 10.1093/brain/awx147
19. Hammer, M. B., Ding, J., Mochel, F., Eleuch-Fayache, G., Charles, P., Coutelier,
M. *et al.* (2017) SLC25A46 Mutations Associated with Autosomal Recessive
Cerebellar Ataxia in North African Families Neurodegener Dis **17**, 208-212
10.1159/000464445
20. Sulaiman, R. A., Patel, N., Alsharif, H., Arold, S. T., andAlkuraya, F. S. (2017) A
novel mutation in SLC25A46 causes optic atrophy and progressive limb
spasticity, with no cerebellar atrophy or axonal neuropathy Clin Genet **92**, 230-
231 10.1111/cge.12963
21. Wan, J., Steffen, J., Yourshaw, M., Mamsa, H., Andersen, E., Rudnik-
Schoneborn, S. *et al.* (2016) Loss of function of SLC25A46 causes lethal
congenital pontocerebellar hypoplasia Brain **139**, 2877-2890
10.1093/brain/aww212
22. Charlesworth, G., Balint, B., Mencacci, N. E., Carr, L., Wood, N. W., andBhatia,
K. P. (2016) SLC25A46 mutations underlie progressive myoclonic ataxia with
optic atrophy and neuropathy Mov Disord **31**, 1249-1251 10.1002/mds.26716
23. Nguyen, M., Boesten, I., Hellebrekers, D. M., Mulder-den Hartog, N. M., de Coo,
I. F., Smeets, H. J. *et al.* (2017) Novel pathogenic SLC25A46 splice-site mutation
causes an optic atrophy spectrum disorder Clin Genet **91**, 121-125
10.1111/cge.12774
24. Abrams, A. J., Hufnagel, R. B., Rebelo, A., Zanna, C., Patel, N., Gonzalez, M. A.
et al. (2015) Mutations in SLC25A46, encoding a UGO1-like protein, cause an
optic atrophy spectrum disorder Nat Genet **47**, 926-932 10.1038/ng.3354
25. Kunji, E. R. S., King, M. S., Ruprecht, J. J., andThangaratnarajah, C. (2020) The
SLC25 Carrier Family: Important Transport Proteins in Mitochondrial Physiology
and Pathology Physiology (Bethesda) **35**, 302-327 10.1152/physiol.00009.2020
26. Duchesne, A., Vaiman, A., Castille, J., Beauvallet, C., Gaignard, P., Floriot, S. *et*
al. (2017) Bovine and murine models highlight novel roles for SLC25A46 in
mitochondrial dynamics and metabolism, with implications for human and animal
health PLoS Genet **13**, e1006597 10.1371/journal.pgen.1006597

27.   Li, Z., Peng, Y., Hufnagel, R. B., Hu, Y. C., Zhao, C., Queme, L. F. *et al.* (2017)
396   Loss of SLC25A46 causes neurodegeneration by affecting mitochondrial
397   dynamics and energy production in mice Hum Mol Genet **26**, 3776-3791
398   10.1093/hmg/ddx262
399   28.   Suda, K., Ueoka, I., Azuma, Y., Muraoka, Y., Yoshida, H., andYamaguchi, M.
400   (2018) Novel Drosophila model for mitochondrial diseases by targeting of a
401   solute carrier protein SLC25A46 Brain Res **1689**, 30-44
402   10.1016/j.brainres.2018.03.028
403   29.   Terzenidou, M. E., Segklia, A., Kano, T., Papastefanaki, F., Karakostas, A.,
404   Charalambous, M. *et al.* (2017) Novel insights into SLC25A46-related
405   pathologies in a genetic mouse model PLoS Genet **13**, e1006656
406   10.1371/journal.pgen.1006656
407   30.   Qiu, K., Zou, W., Fang, H., Hao, M., Mehta, K., Tian, Z. *et al.* (2022) Light-
408   activated mitochondrial fission through optogenetic control of mitochondria-
409   lysosome contacts Nat Commun **13**, 4303 10.1038/s41467-022-31970-5
410   31.   Santo-Domingo, J., Lassueur, S., Galindo, A. N., Alvarez-Illera, P., Romero-
411   Sanz, S., Caldero-Escudero, E. *et al.* (2023) SLC25A46 promotes mitochondrial
412   fission and mediates resistance to lipotoxic stress in INS-1E insulin-secreting
413   cells J Cell Sci **136**, 10.1242/jcs.260049
414   32.   Coonrod, E. M., Karren, M. A., andShaw, J. M. (2007) Ugo1p is a multipass
415   transmembrane protein with a single carrier domain required for mitochondrial
416   fusion Traffic **8**, 500-511 10.1111/j.1600-0854.2007.00550.x
417   33.   Sesaki, H., andJensen, R. E. (2001) UGO1 encodes an outer membrane protein
418   required for mitochondrial fusion J Cell Biol **152**, 1123-1134
419   10.1083/jcb.152.6.1123
420   34.   Sesaki, H., andJensen, R. E. (2004) Ugo1p links the Fzo1p and Mgm1p
421   GTPases for mitochondrial fusion J Biol Chem **279**, 28298-28303
422   10.1074/jbc.M401363200
423   35.   Hoppins, S., Horner, J., Song, C., McCaffery, J. M., andNunnari, J. (2009)
424   Mitochondrial outer and inner membrane fusion requires a modified carrier
425   protein J Cell Biol **184**, 569-581 10.1083/jcb.200809099
426   36.   Sesaki, H., Southard, S. M., Yaffe, M. P., andJensen, R. E. (2003) Mgm1p, a
427   dynamin-related GTPase, is essential for fusion of the mitochondrial outer
428   membrane Mol Biol Cell **14**, 2342-2356 10.1091/mbc.e02-12-0788
429   37.   Wong, E. D., Wagner, J. A., Scott, S. V., Okreglak, V., Holewinske, T. J.,
430   Cassidy-Stone, A. *et al.* (2003) The intramitochondrial dynamin-related GTPase,
431   Mgm1p, is a component of a protein complex that mediates mitochondrial fusion
432   J Cell Biol **160**, 303-311 10.1083/jcb.200209015
433   38.   Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O. *et al.*
434   (2021) Highly accurate protein structure prediction with AlphaFold Nature **596**,
435   583-589 10.1038/s41586-021-03819-2
436   39.   Mirdita, M., Schutze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., andSteinegger,
437   M. (2022) ColabFold: making protein folding accessible to all Nat Methods **19**,
438   679-682 10.1038/s41592-022-01488-1

40. Ishihara, N., Fujita, Y., Oka, T., andMihara, K. (2006) Regulation of mitochondrial
morphology through proteolytic cleavage of OPA1 EMBO J **25**, 2966-2977
10.1038/sj.emboj.7601184
41. Olichon, A., Emorine, L. J., Descoins, E., Pelloquin, L., Brichese, L., Gas, N. *et*
al. (2002) The human dynamin-related protein OPA1 is anchored to the
mitochondrial inner membrane facing the inter-membrane space FEBS Lett **523**,
171-176 10.1016/s0014-5793(02)02985-x
42. Chappie, J. S., Acharya, S., Liu, Y. W., Leonard, M., Pucadyil, T. J., andSchmid,
S. L. (2009) An intramolecular signaling element that modulates dynamin
function in vitro and in vivo Mol Biol Cell **20**, 3561-3571 10.1091/mbc.e09-04-
0318
43. Sever, S., Muhlberg, A. B., andSchmid, S. L. (1999) Impairment of dynamin's
GAP domain stimulates receptor-mediated endocytosis Nature **398**, 481-486
10.1038/19024
44. Anand, R., Eschenburg, S., andReubold, T. F. (2016) Crystal structure of the
GTPase domain and the bundle signalling element of dynamin in the GDP state
Biochem Biophys Res Commun **469**, 76-80 10.1016/j.bbrc.2015.11.074
45. Chappie, J. S., Acharya, S., Leonard, M., Schmid, S. L., andDyda, F. (2010) G
domain dimerization controls dynamin's assembly-stimulated GTPase activity
Nature **465**, 435-440 10.1038/nature09032
46. Kong, L., Sochacki, K. A., Wang, H., Fang, S., Canagarajah, B., Kehr, A. D. *et al.*
(2018) Cryo-EM of the dynamin polymer assembled on lipid membrane Nature
560, 258-262 10.1038/s41586-018-0378-6
47. Gao, S., andHu, J. (2021) Mitochondrial Fusion: The Machineries In and Out
Trends Cell Biol **31**, 62-74 10.1016/j.tcb.2020.09.008
48. Anand, R., Wai, T., Baker, M. J., Kladt, N., Schauss, A. C., Rugarli, E. *et al.*
(2014) The i-AAA protease YME1L and OMA1 cleave OPA1 to balance
mitochondrial fusion and fission J Cell Biol **204**, 919-929 10.1083/jcb.201308006
49. Ban, T., Ishihara, T., Kohno, H., Saita, S., Ichimura, A., Maenaka, K. *et al.* (2017)
Molecular basis of selective mitochondrial fusion by heterotypic action between
OPA1 and cardiolipin Nat Cell Biol **19**, 856-863 10.1038/ncb3560
50. Ge, Y., Shi, X., Boopathy, S., McDonald, J., Smith, A. W., andChao, L. H. (2020)
Two forms of Opa1 cooperate to complete fusion of the mitochondrial inner-
membrane Elife **9**, 10.7554/eLife.50973
51. Mishra, P., Carelli, V., Manfredi, G., andChan, D. C. (2014) Proteolytic cleavage
of Opa1 stimulates mitochondrial inner membrane fusion and couples fusion to
oxidative phosphorylation Cell Metab **19**, 630-641 10.1016/j.cmet.2014.03.011
52. Faelber, K., Dietrich, L., Noel, J. K., Wollweber, F., Pfitzner, A. K., Muhleip, A. *et*
al. (2019) Structure and assembly of the mitochondrial membrane remodelling
GTPase Mgm1 Nature **571**, 429-433 10.1038/s41586-019-1372-3
53. Nyenhuis, S. B., Wu, X., Strub, M. P., Yim, Y. I., Stanton, A. E., Baena, V. *et al.*
(2023) OPA1 helical structures give perspective to mitochondrial dysfunction
Nature **620**, 1109-1116 10.1038/s41586-023-06462-1
54. von der Malsburg, A., Sapp, G. M., Zuccaro, K. E., von Appen, A., Moss, F. R.,
3rd, Kalia, R. *et al.* (2023) Structural mechanism of mitochondrial membrane

remodelling by human OPA1 Nature **620**, 1101-1108 10.1038/s41586-023-
06441-6
55.	Hartmann, L., Kugler, V., andWagner, R. (2016) Expression of Eukaryotic
Membrane Proteins in Pichia pastoris Methods Mol Biol **1432**, 143-162
10.1007/978-1-4939-3637-3_10
56.
https://colab.research.google.com/github/deepmind/alphafold/blob/main/n
otebooks/AlphaFold.ipynb
57.	Wang, R., Mishra, P., Garbis, S. D., Moradian, A., Sweredoski, M. J., andChan,
D. C. (2021) Identification of new OPA1 cleavage site reveals that short isoforms
regulate mitochondrial fusion Mol Biol Cell **32**, 157-168 10.1091/mbc.E20-09-
0605
58.	Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch,
T. *et al.* (2012) Fiji: an open-source platform for biological-image analysis Nat
Methods **9**, 676-682 10.1038/nmeth.2019
499

Abbreviations and nomenclature

OMM – outer mitochondrial membrane

IMM – inner mitochondrial membrane

SLC – solute carrier family

SLC25A46 – solute carrier family 25 member A46

Opa1 – optic atrophy 1

Mfn1/2 – mitofusin 1/2

iPSC – induced pluoripotent stem cells

BSE – bundle signaling element

HR1/2 – heptad repeat 1/2

ER – endoplasmic reticulum

MGD – minimal GTPase domain

DDM – n-Dodecyl-β-D-Maltopyranoside

GDN – glyco-diosgenin

PA – phosphatidic acid

TM – transmembrane segment

MTS – mitochondrial targeting sequence

IMS – inter-membrane space

Figure legends

Figure 1. SLC25A46 interacts with the bundle signaling element (BSE) of Opa1.

A. Domain diagrams of SLC25A46 and Opa1. s-Opa1 (amino acids 253-960) used in this study is designated below full length Opa1 sequence.

B. GDN extracts prepared from *P. pastoris* expressing SLC25A46-His, Strep-Opa1 or co-expressing SLC25A46-His and Strep-Opa1 subjected to StrepTactin column binding and elution.

C. Crosslinks identified in BSE helices 2 and 3 of Opa1. Circular map of SLC25A46 and Opa1 with identified lysine-lysine crosslinks (black arcs). Crosslinker used to generate the crosslink is indicated.

D. AlphaFold 2 model of the SLC25A46-Opa1 interaction. N-terminal region of SLC25A46 comprising residues 1-83 (disordered) not displayed, residues 1-263 of Opa1 not modeled. Box region displaying identified crosslinks and the calculated C□-C□ distances based on the model.

Figure 2. SLC25A46 mutations in the binding interface diminish binding to Opa1.

A. Mutations designed to disrupt the interaction are mapped on the AlphaFold 2 model of the SLC25A46-Opa1 complex. Box enlarged and depicted in open book format.

B. Purified SLC25A46-His WT, I349D mutant or R257A/R347A double mutant were incubated with 2 µM of Strep-Opa1 and subjected to StrepTactin binding and elution experiments. Input and elution analyzed by Western blotting.

Figure 3. The BSE of Opa1 in sufficient to bind SLC25A46.

A. Domain diagrams of SLC25A46 and Opa1. Minimal GTPase domain of Opa1 designated Opa1(MGD) is composed of the GTPase domain and three BSE helices, shown as line diagram below full length Opa1 sequence.

B. Purified SLC25A46-His WT was incubated with 2 µM of Strep-Opa1 or Strep-Opa1(MGD) and subjected to StrepTactin binding and elution. Input and elution analyzed by Western blotting. Control reaction: 10 µM of SLC25A46-His alone.

C. Circular map of SLC25A46 and Opa1(MGD) with identified lysine-lysine crosslinks (black arcs) shown. Crosslinker used is indicated. Crosslinks were identified in BSE helices 2 and 3 of Opa1(MGD) as in Opa1.

D. A model of the SLC25A46-Opa1(MGD) interaction. The N-terminal region of SLC25A46 (residues 1-83) is disordered and not displayed. Boxed region enlarged on right displaying the identified crosslinks and the calculated C□-C□ distances based on the model.

Figure 4. Disease-causing SLC25A46 mutants retain binding to Opa1 or Mfn2.

A. Identified SLC25A46 disease mutations mapped on AlphaFold 2 model of SLC25A46-Opa1 complex. Box regions enlarged on the right to display the mutations.

B. GDN extracts from *P. pastoris* co-expressing Strep-Opa1 and SLC25A46-His WT or mutants subjected to StrepTactin (middle) or Ni-NTA (bottom) column binding and elution.

Figure 5. Mutations disrupting Opa1 interaction alters mitochondrial morphology.

A. Western blot of expression levels for Opa1, Mfn2, SLC25A46 or MTCO1 in HCT116 WT, SLC25A46-KO or SLC25A46-KO cells expressing SLC25A46 variants. (Δ2-

580 83)SLC25A46 not detected as commercially available antibodies detect the N-terminal
581 epitope of SLC25A46. MTCO1 is used as loading control.
582
583 B. Mitochondrial morphology in HCT116 WT, SLC25A46-KO or SLC25A46-KO cells
584 expressing SLC25A46 variants. Mitochondrial morphology for 100 cells were scored for
585 each condition and evaluated (see Materials and Methods).

Figure 1

Figure 2

Figure 3

594　**Figure 4**

Figure 5

 Supporting Information

Supporting experimental procedures

Binding experiments using lysates – additional details

For nucleotide treatments, StrepTactin binding experiments were performed as

described but the buffer used was 25 mM HEPES, 100 mM NaCl, 50 mM KCl, 10 mM

MgCl$_2$.6H$_2$O, pH 8.0, 1% GDN (extraction) or 0.1% GDN (binding and elution) and

containing 2 mM of GTPγS (Sigma-Aldrich, 10220647001), GMPPCP (Sigma-Aldrich,

M3509-25MG) or GDP (Sigma-Aldrich, G7127-25MG).

To assess binding of SLC25A46-His disease mutants to Opa1 or Mfn2, *P.*

pastoris strains co-expressing the proteins were cultured in 5-10 ml BMMY. Equal

number of cells (estimated by OD600) were harvested from each culture. The cells were

resuspended in 0.5 ml of 25 mM HEPES, 150 mM NaCl, pH 8.0 containing protease

inhibitors. An equal volume of 0.5 mm glass beads was added and lysed using a Retsch

TissueLyzer. To the recovered lysate, GDN was added from a stock solution of 10% to

a final concentration of 1% and incubated in the cold room for 1.5 h tumbling gently. The

lysate was then centrifuged at 21,100 xg for 30 min at 4 °C. 150-300 ul of the lysate was

applied to either Ni-NTA (Bio-Rad, 780-0800) or StrepTactin XT 4Flow resin pre-

equilibrated with the above buffer containing 10 mM imidazole (for Ni-NTA) or 1 mM

EDTA (StrepTactin) and 0.1% GDN. Resins were washed 3x and eluted with buffer

containing 500 mM imidazole (for Ni-NTA) or 50 mM biotin (StrepTactin) and 0.1%

GDN. The samples were subjected to western blotting using anti-Strep (Mfn2 or Opa1)

and anti-His (SLC25A46) antibodies.

Protein expression

For protein expression in *P. pastoris*, a single desired clone was inoculated in 5-

10 ml BMGY and then scaled up to 500 ml BMGY media and grown overnight at 30 °C.

The culture was harvested and resuspended in 6 litres BMMY at an OD600 of 1.0 for

protein induction. The induced cultures were then grown at 20 °C and harvested 24 h

later. Cell pellets were frozen in liquid nitrogen and stored at –80 °C.

Opa1 and Opa1(MGD) were also purified from *E. coli.* For expression in *E. coli*,

transformed BL21-CodonPlus (DE3)-RIL cells (Agilent, 230245) were grown in LB at 37

°C to an OD600 of 0.7 and induced with 0.2 mM IPTG (Gold Biotechnology, I2481C200)

and grown overnight at 16 °C. Cultures were harvested and stored at –80 °C until use.

636   Purification of SLC25A46

Harvested *P. pastoris* cells expressing WT, I349D mutant or R257A/R347A

double mutant were milled under cryogenic conditions using SPEX SamplePrep 6875D

Freezer/Mill. All purification steps were done at 4 °C. The milled cells were resuspended

in 50 mM HEPES, 250 mM NaCl, 50 mM imidazole, pH 7.8 containing protease

inhibitors (Sigma-Aldrich, 5056489001) and benzonase nuclease (Sigma-Aldrich,

E1014-25KU). The suspension was centrifuged at 235,000 xg for 0:45 h to collect the

insoluble pellet. The pellet was then solubilized in the same buffer containing 2% n-

Dodecyl-β-D-Maltopyranoside (DDM) (Anatrace, D310) and stirred for 1 h. The extract

was centrifuged at 100,000 xg for 1 h and the supernatant was loaded on Ni-NTA

Agarose resin (Qiagen, 30230) pre-equilibrated with 25 mM HEPES, 150 mM NaCl, 50

mM imidazole, pH 7.8, 0.05-0.1% glyco-diosgenin (GDN) (Anatrace, GDN101). The

resin was washed with this buffer before elution using buffer containing 500 mM

imidazole. The elution fractions were concentrated using 50,000 MWCO concentrators

and dialyzed using 20,000 MWCO cassettes (Thermo Fisher Scientific, A52976) against

25 mM HEPES, 150 mM NaCl, pH 8.0, 0.05% GDN. The dialyzed sample was collected

and concentrated (50,000 MWCO) further. Protein concentration was estimated by

absorbance at 280 nm.

655   Purification of Opa1 and Opa1(MGD)

All steps were done at 4 °C. Harvested *E. coli* cells expressing Opa1(253-960) or

Opa1(MGD) were resuspended in 50 mM HEPES, 250 mM NaCl, 1 mM EDTA, pH 8.0,

1 mg/ml lysozyme, protease inhibitors and benzonase nuclease. The cell suspension

was lysed by sonication and centrifuged at 30,000 xg for 1 h. The supernatant was

flowed over StrepTactin XT 4Flow high-capacity resin (IBA Lifesciences, 2-5030-010)

packed in a gravity column and equilibrated with 25 mM HEPES, 150 mM NaCl, pH 8.0.

The same buffer was used to wash the resin and the bound protein was eluted using
buffer containing 50 mM biotin (IBA Lifesciences, 2-1016-005). The fractions were
concentrated (50,000 MWCO for Opa1, 20,000 MWCO for Opa1(MGD)) and dialyzed
(20,000 MWCO) against 25 mM HEPES, 150 mM NaCl, pH 8.0. The concentrations
were estimated using absorbance at 280 nm. The proteins were aliquoted and frozen in
liquid nitrogen and stored at -80 °C.

Generation of SLC25A46-Opa1 and SLC25A46-Mfn2 complexes

All purification steps were performed at 4 °C or on ice. To purify SLC25A46-Opa1
complex, *P. pastoris* co-expressing SLC25A46-His and Strep-Opa1 was harvested and
cryo-milled as before. The milled cells were resuspended in 25 mM HEPES, 150 mM
NaCl, 25 mM imidazole, pH 8.0 containing protease inhibitors and benzonase nuclease.
GDN (2%) and 18:1 PA (Avanti Polar Lipids, 840875) (0.1 mg/ml solubilized in GDN)
were added and stirred for 2 h. The suspension was centrifuged at 100,000 xg for 1 h.
The supernatant was loaded onto a 1 ml HisTrap HP column and washed with 25 mM
HEPES, 150 mM NaCl, 25 mM imidazole, pH 8.0, 0.025 mg/ml 18:1 PA and 0.1% GDN
and eluted with the same buffer containing 500 mM imidazole. The elution was diluted
with StrepTactin binding buffer, 25 mM HEPES, 150 mM NaCl, 1 mM EDTA, pH 8.0,
0.025 mg/ml PA, 0.1% GDN to bring the imidazole concentration below 100 mM and
added to 1 ml StrepTactin XT 4Flow resin, washed with the above binding buffer and
eluted in buffer containing 50 mM biotin. The elution fractions containing SLC25A46 and
Opa1 were combined and concentrated (50,000 MWCO) to ~5.5 mg/ml.

To purify SLC25A46-Mfn2 complex, *P. pastoris* co-expressing SLC25A46-His
and Mfn2-Strep was harvested, cryo-milled and resuspended in 25 mM HEPES, 150
mM NaCl, 25 mM imidazole, pH 8.0 with protease inhibitors and benzonase nuclease
and centrifuged at 235,000 xg for 45 min to collect the membrane fractions. The
membrane fractions were resuspended in the same buffer containing 2% GDN and 0.1
mg/ml PA and stirred for 2 h. The extract was centrifuged and processed using Ni-NTA
and StrepTactin columns as described above. The elution fractions from StrepTactin
column were concentrated (50,000 MWCO) to ~2.4 mg/ml.

Binding experiments using purified proteins

Binding reactions were prepared in 25 mM HEPES, 150 mM NaCl, pH 8.0, 0.1% GDN (binding buffer). Purified Opa1 at 2 µM concentration was incubated with varying concentrations of SLC25A46 at 4 °C for 3 h. The reaction mixtures (100 µl) were added to 50 µl StrepTactin XT 4Flow high-capacity resin packed in spin column and incubated with gentle agitation for 15 min. The spin columns were centrifuged at 500 xg for 30 s to collect the flow-through. The columns were washed thrice with 100 µl of binding buffer. The bound proteins were eluted by incubating with 100 µl of buffer containing 50 mM biotin for 15 min followed by centrifugation at 500 xg for 30 s. Samples analyzed by Western blotting.

Western blotting

Following SDS PAGE, proteins were transferred to PVDF membrane (Bio-Rad, 1704156) using Bio-Rad Trans-Blot Turbo Transfer system. The membranes were incubated with the primary antibodies as indicated and were probed with the appropriate secondary antibodies conjugated to horseradish peroxidase. The blots were developed using Western Lightning Plus Chemiluminescent Substrate (Revvity, NEL105001EA) and imaged using Amersham Imager 680. Densitometry was performed using ImageJ. Antibodies: anti-His (Thermo Fisher Scientific, MA1-21315, 1:1000), anti-Strep (IBA Lifesciences, 2-1509-001, ~1:33000), anti-SLC25A46 (Abcam, ab237760, 1:2500), anti-Opa1 (Novus Biologicals, NBP2-59770, 1:1000), anti-Mfn2 (Thermo Fisher Scientific, 702768, 1:1000), anti-MTCO1 (Abcam, ab14705, 1:2000), anti-mouse, HRP conjugated (Sigma-Aldrich, GENA931-1ML, 1:5000), anti-rabbit, HRP conjugated (Sigma-Aldrich, GENA934-1ML, 1:5000).

Crosslinking mass spectrometry

For crosslinking, BS3 (Thermo Fisher Scientific, A39266) was added to the concentrated SLC25A46-Opa1(263-960) and SLC25A46-Mfn2 complexes at a final concentration of 0.75 mM. Reactions were incubated on ice for 1 h and quenched by adding Tris-HCl, pH 7.8 at a final concentration of ~45 mM) and incubating 15 min on

ice. The samples were separated on a 7.5% polyacrylamide gel and bands excised for mass spectrometry.

While we identified several monolinks and intralinks in Opa1 with BS3 crosslinker, we were unable to identify any crosslinks with SLC25A46 as the sequencing depth for SLC25A46 in this sample was poor. To overcome this, we performed crosslinking experiments with excess purified proteins. We used BS(PEG)$_5$ (Thermo Fisher Scientific, 21581) (25 mM stock in 10% DMSO) in addition. The crosslinker (BS3 or BS(PEG)5) was added at a final concentration of 2.5 mM to 20 µM SLC25A46 and 30 µM Opa1(253-960) or Opa1(MGD) and the sample separated on a 12% Bis-Tris 1.0 mm, Mini Protein Gels. The subsequent steps were same as above. The excised gel bands were destained with 1:1 v/v methanol and 100 mM ammonium bicarbonate. Gel pieces were partially dehydrated with acetonitrile then further dehydrated by SpeedVac. Proteins were reduced at 57 °C for 1 h by adding 100 µl of 20 mM aqueous DTT solution to the dehydrated gel piece. The solution was removed and replaced with 100 µl of a 50 mM iodoacetamide solution and the gel pieces incubated in the dark for 45 min. the solution was removed, and the gel pieces again partially dehydrated with an acetonitrile rinse and further dried in a SpeedVac. 250 ng of sequencing grade modified trypsin (Promega) was added to each sample followed by 300 µl of 100 mM ammonium bicarbonate to cover the gel pieces. Digestion was allowed to proceeded overnight on a shaker at room temperature. To stop the digestion and assist with peptide extraction a solution of 5% formic acid in acetonitrile 1:2 (v:v) was added and incubated for 15 min with agitation. The solution was transferred into a new tube and the procedure repeated two more times. The combined solutions were dried down in a SpeedVac concentrator to remove the acetonitrile. The samples resuspended in 0.1% acetic acid and loaded onto equilibrated Ultra-Micro SpinColumns™ (Harvard Apparatus) using a microcentrifuge. The spin columns were washed three times with 0.1% trifluoroacetic acid and the last wash with 0.5% acetic acid. Peptides were eluted with 40% acetonitrile in 0.5% acetic acid, followed by 80% acetonitrile in 0.5% acetic acid. The solutions combined and dried down using a SpeedVac concentrator. The samples were reconstituted in 0.5% acetic acid and stored at -80 °C until analysis.

Approximately 1 µg of each sample was analyzed individually by LC/MS/MS. Samples were loaded onto an Acclaim PepMap trap column (2 cm x 75 µm) in line with an EASY-Spray analytical column (50 cm x 75 µm ID PepMap C18, 2 µm bead size) using the autosampler of an EASY-nLC 1200 HPLC (Thermo Scientific). Solvent A was 2% acetonitrile in 0.5% acetic acid and solvent B was 80% acetonitrile in 0.5% acetic acid. The gradient used was the following: 5 min held at 5% solvent B, in 60 min to 35% solvent B, in 10 min to 45% solvent B, and in 10 min to 100% solvent B. Peptides were gradient eluted directly into a Thermo Scientific Orbitrap Eclipse Mass Spectrometer.

Mfn2 and SLC25A46 Crosslink analysis:

High resolution full spectra were acquired with a resolution of 240,000, an AGC target of 1e6, with a maximum ion time of 50 ms, and a scan range of 400 to 1500 m/z. All precursors with charge states between 2-7 were selected for fragmentation with a dynamic exclusion of 20 sec. All MS/MS HCD spectra were collected in the ion trap using the rapid scan mode, an AGC target of 2e4, maximum inject time of 18 ms, one microscan, 0.7 m/z isolation window, and normalized collision energy (NCE) of 27. The instrument was set to run at top speed with a cycle time of 3 sec. Samples were run a second time to specifically only fragment precursor ions that are likely crosslinked (charge state 4-10) using either HCD or ETD fragmentation. The gradient used was the following: 5 min held at 5% solvent B, in 120 min to 35% solvent B, in 10 min to 45% solvent B, and in 10 min to 100% solvent B. High resolution full spectra were acquired with a resolution of 60,000, an AGC target of 4e5, with a maximum ion time of 50 ms, and a scan range of 400 to 1500 m/z. Charge states 4-6 were analyzed in the Orbitrap using HCD with an NCE of 27, resolution of 30K, AGC target of 2e5, maximum inject time of 200 msec and an isolation window of 2. Charge states 7-10 were analyzed in the Orbitrap using ETD with a charge state dependent reaction time, resolution of 30K, AGC target of 2e5, maximum inject time of 54 msec and an isolation window of 2.

Opa1 and SLC25A46 Crosslink analysis:

High resolution full spectra were acquired with a resolution of 120,000, an AGC target of 4e6, with a maximum ion time of 50 ms, and a scan range of 400 to 1500 m/z. All precursors with charge states between 2-5 were selected for fragmentation with a dynamic exclusion of 30 sec. All MS/MS HCD spectra were collected in the Orbitrap

785 with a resolution of 30K, an AGC target of 2e5, maximum inject time of 200 ms, one
786 microscan, 2 m/z isolation window, and normalized collision energy (NCE) of 27.
787 MS/MS spectra were searched in Byos by Protein Metrics against the human
788 SLC25A46, Opa1 and Mfn2 sequences to identify crosslinked peptides.

Cell lines and culture conditions

HCT116 WT and SLC25A46 KO cells were kind gifts from CeMM (Research
Center for Molecular Medicine, Medical University of Vienna, Austria). The cells
maintained in RPMI 1640 with 10% fetal bovine serum (FBS) and 1%
penicillin/streptomycin at 37 °C and 5% CO_2. Cell genotype confirmed by Western
blotting. Human SLC25A46 sequence from plasmid pDONR221_SLC25A46 (a gift from
RESOLUTE Consortium & Giulio Superti-Furga (Addgene plasmid #131971;
http://n2t.net/addgene:131971; RRID:Addgene_131971) was cloned into pCDNA3.1(+)
plasmid to generate the construct pAH9. Site-directed mutagenesis (New England
Biolabs) was used to generate expression plasmids bearing the mutations G249D
(pAH10), R257Q (pAH11), R340C (pAH12), I349D (pAH13), Δ1-83 (pAH14) and
R257A/R347A (pAH36) in the SLC25A46 coding sequence of pAH9. To transiently
express SLC25A46 and its mutant derivatives in HCT116 SLC25A46 KO cells, cells
seeded onto 35 mm glass-bottom dishes (MatTek Life Sciences) coated with poly-D-
lysine (0.1 mg/ml) were transfected with 2.5 μg expression plasmid mixed with
Lipofectamine 3000 in a total volume of 250 μl Opti-MEM (Thermo Fisher Scientific).
Cells were grown for 36-48 h prior to imaging.

For Western blotting, mitochondria were isolated using Qiagen Qproteome
Mitochondria Isolation Kit (Qiagen, 37612). Protease inhibitors were added to isolated
mitochondria and solubilized using 0.5% DDM. Protein content was estimated using
Bradford assay (Bio-Rad, 5000205). ~1 μg of total protein was loaded on.

Fluorescence imaging and quantification

HCT116 cells were seeded onto glass-bottom dishes and grown to ~80%
confluency. Cells were washed with PBS and fixed in 4% paraformaldehyde (PFA) in
PBS for 20 min. Following three PBS washes, cells were permeabilized in 0.1% Triton

X-100 diluted in PBS for 5 min. Cells were incubated in 2% normal goat serum and
0.05% Triton X-100 in PBS for 1 h, then incubated in the same buffer with 1:300 Tom20
rabbit antibody (Invitrogen, MAS-34964) for 1 h at room temperature or overnight at
4°C. After 3 washes with PBS, cells were incubated with 1:1000 goat anti-rabbit Alexa
Fluor® 488 AffiniPure antibody (Jackson ImmunoResearch, 111-545-144) for 1 h at
room temperature. Cells were washed in PBS and coated with UltraCruz® Aqueous
Mounting Medium with DAPI (Santa Cruz) prior to imaging. Videos were captured using
A1R HD25 point scanning confocal with GaAsP and PMT detectors, equipped with an
Apo TIRF 60x/1.49 NA objective lens and Ti2 Z-drive. 100 cells per transfected
condition were manually scored into four morphological classifications (57):
"Fragmented" refers to cells that contain spherical mitochondrial fragments with two or
less short tubules present. "Short tubular" refers to cells containing a mixture of
fragmented and short tubular mitochondria. "Long tubular" refers to cells with elongated
mitochondria, but not fused into a mitochondrial mesh. "Interconnected" refers to cells
with a highly interconnected network of mitochondrial filaments, with few isolated
mitochondria.

Live cell confocal microscopy

Cells were stained with 1:1000 PKmito Deep Red (Spirochrome) at 37°C for 30
min. Following three 1X PBS washes, cells were placed in Live Cell Imaging Solution
(Invitrogen). Acquisition was performed using Nikon A1R HD25 confocal with GaAsP
and PMT detectors and an Apo 60x/1.49 NA objective. Temperature, humidity, and CO_2
were controlled within an Oko-Lab chamber equipped with Live Cell^{TM} and Solent
Scientific controllers. Image acquisition done using NIS-Elements (Nikon Instruments
Inc.) and analysis performed using Fiji (58). Time-lapse videos of mitochondria taken at
1 frame per 10 sec for 5 min.

Supporting Figures

Supporting Figure 1. SLC25A46 interacts with Opa1.

A. Strep-Opa1 co-elutes upon elution of bound SLC25A46-His. GDN extracts prepared from *P. pastoris* expressing SLC25A46-His, Strep-Opa1 or co-expressing SLC25A46-His and Strep-Opa1 were subjected to Ni-NTA column binding and elution.

B. The intralinks (black arcs) and monolinks (red circles) identified from crosslinking-mass spectrometry experiments are mapped on the domain diagrams of SLC25A46 and Opa1.

C. AlphaFold2 model for SLC25A46-Opa1 complex colored by pLDDT (red = high confidence, blue = low confidence).

Supporting Figure 2. Electrostatic and hydrophobic interactions are predicted to stabilize SLC25A46-Opa1 interaction.

A. Two different views of the SLC25A46-Opa1 interface predicted by AlphaFold 2 is enlarged on the right. Top: Predicted salt-bridge interactions between R257 of SLC25A46 and E561 of Opa1, and R347 of SLC25A46 and D565 of Opa1. Bottom: Adjacent to the salt bridges is a cluster of hydrophobic residues comprising of I229, I230, I349, Y357, V359, L360 of SLC25A46 and T576, F572, A569 of Opa1.

B. Intensity of the anti-His (SLC25A46) bands in the elution in Figure 2 were normalized against the corresponding bands from the input and plotted as a function of SLC25A46 concentration used in the reaction mixtures.

C. Presence of the nucleotides did not affect binding of SLC25A46 to Opa1. GDN extracts prepared from *P. pastoris* co-expressing SLC25A46-His and Strep-Opa1 were

incubated with 2 mM of the indicated guanosine nucleotide prior to StrepTactin binding

and elution.

Supporting Figure 3. SLC25A46 interacts with Mfn2.

A. Domain diagrams of SLC25A46 and Mfn2 are depicted.

B. SLC25A46-His co-elutes upon elution of bound Mfn2-Strep. GDN extracts prepared

from *P. pastoris* expressing Mfn2-Strep or co-expressing SLC25A46-His and Mfn2-

Strep were subjected to StrepTactin column binding and elution.

C. Mfn2-Strep co-elutes upon elution of bound SLC25A46-His. GDN extracts prepared

from *P. pastoris* expressing Mfn2-Strep or co-expressing SLC25A46-His and Mfn2-

Strep were subjected to Ni-NTA column binding and elution.

D. Circular map of SLC25A46 and Mfn2 with the identified lysine-lysine crosslinks (black

arcs). Crosslinker used is indicated. Crosslinks were identified between the cytosol-

exposed N-terminal K60 of SLC25A46 and the HR1 lysines K416 and K420 of Mfn2.

E. The intralinks (black arcs) and monolinks (red circles) identified from crosslinking-

mass spectrometry experiments are mapped on the domain diagram of Mfn2.

F. Model of the SLC25A46-Mfn2 interaction. The identified crosslinks are depicted as

dotted lines in the AlphaFold 2 generated models of SLC25A46 and Mfn2. The N-

terminal region of SLC25A46 comprising residues 1-83 is predicted to be disordered

and is displayed as a hand-drawn cartoon. The transmembrane helices of Mfn2 were

poorly modeled and are shown as cylinders.

G. GDN extracts prepared from *P. pastoris* co-expressing Mfn2-Strep and either

SLC25A46-His WT, (Δ2-83)SLC25A46-His or disease mutants were subjected to

StrepTactin (middle) or Ni-NTA (bottom) column binding and elution. (Δ2-83)SLC25A46

still interacts with Mfn2 as seen by co-elution of both proteins. Disease mutations did

not abrogate binding of SLC25A46 to Opa1 or Mfn2.

H. GDN extracts prepared from *P. pastoris* co-expressing SLC25A46-His and Mfn2-

Strep were incubated with 2 mM of the indicated guanosine nucleotide prior to

StrepTactin binding and elution. Presence of the nucleotides did not affect binding of

SLC25A46 to Mfn2.

Supporting Video 1.

Live-cell confocal microscopy of PKmito Deep Red-stained mitochondria in indicated

HCT116 cell lines. Movies were taken at 15 sec per frame for 5 min. Playback at 4

frames per sec (37.5x real-time). Scale bar = 10 μm.

 Supporting Figure 1

Supporting Figure 2

Supporting Figure 3