Identification of SLC25A46 interaction interfaces with

mitochondrial membrane fusogens Opa1 and Mfn2

Sivakumar Boopathy[1,2], Bridget E. Luce[1], Camila Makhlouta Lugo[1*], Pusparanee

Hakim[1], Julie McDonald[1#], Ha Lin Kim[1], Jackeline Ponce[3], Beatrix M. Ueberheide[3,4],

Luke H. Chao[1,2,5]

[1]Department of Molecular Biology, Massachusetts General Hospital, Boston MA 02114,

USA

[2]Department of Genetics, Harvard Medical School, Boston MA 02115, USA

[3]Proteomics Resource Center, Division of Advanced Research Technologies, New York

University Langone Health Center, New York NY 10016, USA

[4]Department of Biochemistry and Molecular Pharmacology, New York University

Langone Health Center, New York NY 10016, USA

[5]Corresponding author

*Present address: Program in Neurobiology, University of Chicago, Chicago IL 60637,

USA

#Present address: Department of Biology, Massachusetts Institute of Technology,

Cambridge MA 02139, USA

Running Title

SLC25A46 interaction interfaces with Opa1 and Mfn2

Keywords

1. Mitochondria
2. Membrane fusion
3. GTPase
4. Structural model
5. Protein cross-linking
6. Mass spectrometry
7. Protein-protein interaction
8. Mitochondrial solute carrier

Abstract

Mitochondrial fusion requires the sequential merger of four bilayers to two. The outer-membrane solute carrier protein SLC25A46 interacts with both the outer and inner-membrane dynamin family GTPases Mfn1/2 and Opa1. While SLC25A46 levels are known to affect mitochondrial morphology, how SLC25A46 interacts with Mfn1/2 and Opa1 to regulate membrane fusion is not understood. In this study, we use crosslinking mass-spectrometry and AlphaFold 2 modeling to identify interfaces mediating a SLC25A46 interactions with Opa1 and Mfn2. We reveal that the bundle signaling element of Opa1 interacts with SLC25A46, and present evidence of a Mfn2 interaction involving the SLC25A46 cytosolic face. We validate these newly identified interaction interfaces and show that they play a role in mitochondrial network maintenance.

Introduction

Mitochondrial fission and fusion are membrane remodeling processes required for maintenance of a healthy mitochondrial reticulum (1, 2), which regulates the number and distribution of mitochondrial DNA and respiratory chain proteins (1-5). A distinct feature of mitochondrial fusion is the sequential fusion of two pairs of bilayers. Key questions regarding the spatiotemporal control of these events remain. Specifically, how are the activities of the outer and inner membrane fusion GTPases, Mfn1/2 and Opa1 respectively, coordinated and regulated?

SLC25A46 interacts with both fusion GTPases and has emerged as an inter-membrane bridging protein, playing roles in regulating mitochondrial membrane dynamics (6-9). SLC25A46 is a member of the solute carrier family 25 of mitochondrial transporters, which transport substrates such as nucleotides, vitamins, amino acids, fatty acids across the inner mitochondrial membrane (IMM) (10). Residing in the outer mitochondrial membrane (OMM), SLC25A46 lacks the signature sequence motifs of the SLC25 family (10), does not have an identified substrate and contains a distinct N-terminal extension. Mutations in SLC25A46 result in optic atrophy spectrum disorder, Leigh syndrome and pontocerebellar hypoplasia (11-25).

In SLC25A46 knockout and knockdown models of human neurological disease, mitochondria are often enlarged and aggregated in nervous tissues, exhibiting detached, vesiculated cristae (22, 25-29). Respiratory chain complex activity is impaired in these models (26, 27), and ATP levels are reduced (27, 28). In human cell lines, knockdown of SLC25A46 induces a tubular mitochondrial network whereas overexpression results in mitochondrial fragmentation (6, 9, 25), suggesting a role in downregulating fusion. The phenotypes of SLC25A46 knockout varied depending on the cell lines tested, displaying mitochondrial fragmentation in human fibroblasts and HeLa cells (8), and hyper-filamentous network in other cell lines (30, 31). Knockout of the yeast orthologue Ugo1 (~22% sequence identity) also induced fragmentation (32-34). These observations argue for a role of SLC25A46 in regulating mitochondrial fusion where SLC25A46 levels at the OMM are critical. In fibroblasts and iPSC-derived cerebral neurons, SLC25A46 is focally present at tips and branches of mitochondrial tubules and at sites of fusion and fission (8). Interestingly the SLC25A46 puncta were

proximal to Opa1 in human fibroblasts (8), suggesting spatially-defined interactions with
SLC25A46 play a role in fusion regulation.

Understanding the molecular details of the SLC25A46-Mfn2-Opa1 interactions
can provide constraints on how SLC25A46 organizes sites for mitochondrial membrane
remodeling and inform mechanisms for coordinated OMM/IMM fusion. In yeast, Ugo1
physically interacts with Fzo1 and Mgm1, the orthologous fusion GTPases (34-37).
Previous studies indicate Opa1 and Mfn2 interacts with SLC25A46 (6-9), however the
specific interfaces mediating these interactions was undefined. In this study, we used
crosslinking-mass spectrometry and AlphaFold 2 modeling to identify the protein-protein
interaction surfaces of SLC25A46-Opa1 and SLC25A46-Mfn2. We validate this newly
identified SLC25A46-Opa1 interface by mutagenesis and demonstrate a role for this
interface in mitochondrial network maintenance.

Results

SLC25A46 interacts with Opa1 via its bundle signaling element.

Previous studies used crosslinking, immunoprecipitation and mass spectrometry
to identify an interaction network of Opa1, Mfn1/2 and SLC25A46 (6-9). We over-
expressed human SLC25A46 with Opa1 in *Pichia pastoris* under the control of a strong
promoter (alcohol oxidase) (**Figure 1A**) and found that detergent (GDN)-solubilized
SLC25A46 in *P. pastoris* lysate interacts with column-immobilized Opa1 from lysate in
the absence of crosslinkers (**Figure 1B**). Likewise, column-immobilized SLC25A46 was
also found to interact with Opa1 (**Supporting Figure 1A**).

To identify the specific interface responsible for the SLC25A46-Opa1 interaction,
we crosslinked the purified complex for analysis by mass spectrometry. Using
bis(sulfosuccinimidyl)suberate (BS3) we identified a crosslink between K241 of
SLC25A46 and K568 of Opa1 bundle signaling element (BSE) helix 2. Intralinks
consistent with folded Opa1 secondary structure elements were identified (**Supporting
Figure 1B**).   A second crosslinker Bis-N-succinimidyl-(pentaethylene glycol)
(BS(PEG)5) with a larger spacer length also crosslinked K241(SLC25A46)-K568(Opa1),
and identified a second crosslink between K256 of SLC25A46 and the K960 of Opa1
BSE helix 3 (**Figure 1C**). We modeled this interaction using Alphafold 2 (38, 39), which

consistently revealed a high-confidence and robust interface mediated by the Opa1

BSE, with a topology consistent with the proteins' sub-organellar localization (6, 22, 25,

40, 41) (**Figure 1D, Supporting Figure 1C**).

Validation of SLC25A46:Opa1-BSE interaction

Examination of the SLC25A46-Opa1 interface in the AlphaFold 2 model revealed

two salt bridges comprising residues R257 (SLC25A46) and E561 (Opa1); R347

(SLC25A46) and D565 (Opa1). Adjacent to the salt bridges, a cluster of hydrophobic

residues, I229, I230, I349, Y357, V359 and L360 in SLC25A46 and A569, F572 and

T576 in Opa1 BSE helix 2 appear to further stabilize the interaction (**Figure 2A**). We

tested this newly identified SLC25A46-Opa1 interface by mutating the salt bridges and

hydrophobic site residues in SLC25A46 and evaluating if the interaction with Opa1 was

maintained. While recombinantly expressed and purified WT SLC25A46 bound column-

immobilized Opa1 (consistent with observations in lysates), both the I349D mutation

and the R257A/R347A double mutation diminished the interaction (**Figure 2B and

Supporting Figure 2A**).

The BSE is essential for transducing GTP-hydrolysis stimulated activities of

dynamin related protein (42, 43). In dynamin, the BSE frequently moves closer to the

GTPase upon GTP hydrolysis (44-46). To determine if the nucleotide-state of Opa1

affects its interaction with SLC25A46, we incubated the lysates with GTPγS, GMPPCP

and GDP prior to affinity capture on StrepTactin resin. We found that the presence of

nucleotide or nucleotide analogues did not affect co-elution, indicating nucleotide-

coupled BSE rearrangement does not affect Opa1-SLC25A46 interaction (**Supporting

Figure 2B**).

We also generated a minimal construct consisting of the GTPase domain and the

BSE alone (Opa1(MGD)), to test that the "paddle" and stalk self-assembly regions are

dispensable for the Opa1-SLC25A46 interaction (**Figure 1A**). SLC25A46 co-elutes with

Opa1(MGD), indicating an Opa1(MGD) construct is sufficient for binding (**Supporting

Figure 3A**). We found the same two crosslinks observed for Opa1-SLC25A46 between

Opa1(MGD) and SLC25A46. In addition, we observed a third crosslink between K387 of

SLC25A46 and K568 of Opa1 in the same vicinity, strengthening our findings

(**Supporting Figure 3B and 3C**).

Cytosolic facing SLC25A46 interactions with the Mfn2 ectodomain.

We next investigated if SLC25A46 also interacts with the outer membrane fusion

protein Mfn2. We co-expressed the two proteins in *P. pastoris* (**Figure 3A**) and

subjected the GDN-solubilized lysates to column-binding experiments. SLC25A46 co-

elutes upon elution of column-immobilized full-length Mfn2 (**Figure 3B**), and Mfn2 co-

elutes with resin bound SLC25A46 (**Supporting Figure 4A**). To test the SLC25A46-

Mfn2 interaction was nucleotide-state dependent, we treated lysates with GTPγS,

GMPPCP or GDP during detergent extraction. The presence of these nucleotides did

not enhance or diminish interaction of Mfn2 with SLC25A46 (**Supporting Figure 4B**).

To gain insight into the nature of the SLC25A46-Mfn2 interaction, we performed

crosslinking and mass-spectrometry on the purified complex. In addition to

intramolecular crosslinks consistent with folded Mfn2 (**Supporting Figure 4C**), two

inter-links were identified – between K60 residue of SLC25A46 and residues K416 and

K420 of Mfn2 (**Figure 3C**). K60 of SLC25A46 is part of the first ~83 residues that are

predicted to be disordered in the cytosol (**Figure 3D**). Residues K416 and K420 are part

of the cytosol facing heptad repeat 1 (HR1) of Mfn2 (**Figure 3D**). In contrast to the

Opa1-SLC25A46 models, AlphaFold 2 predicted interface was low confidence for the

SLC25A46-Mfn2 complex, particularly for the N-terminal region of SLC25A46, and the

Mfn2 transmembrane segments (**Supporting Figure 4D**).

Since crosslinks were observed at K60 in SLC25A46, we tested the role of these

N-terminal residues in binding Mfn2. We deleted the first 82 amino acids of SLC25A46

termed (Δ2-83)SLC25A46, and co-expressed this protein with Mfn2 in *P. pastoris*.

Interestingly, (Δ2-83)SLC25A46 still co-eluted with Mfn2 in the affinity capture

experiments (**Figure 3E**), suggesting interactions outside this N-terminal segment

mediating Mfn2 interactions. Consistent with this idea, MBP-fusion of the SLC25A46 N-

terminus (SLC25A46(1-88)-MBP) was insufficient to capture Mfn2 (**Figure 3F**). To

exclude the possibility that the transmembrane segments of Mfn2 interact with

SLC25A46, we co-expressed SLC25A46 with Mfn2 lacking the transmembrane helices

(Mfn2(ΔTM)). We observed co-elution of SLC25A46 with Mfn2(ΔTM) (**Figure 3G**),

supporting an interaction between the Mfn2 ectodomain and cytosol-exposed regions of

SLC25A46. Expression of a minimal Mfn2(MGD) construct, containing only the helical

bundle (HB1) and GTPase domain (47), still bound SLC25A46, suggesting this region of

the ectodomain interacts with the SLC25A46 cytosolic face (**Figure 3H**).

Mutations disrupting SLC25A46-Opa1 interaction alter mitochondrial morphology.

A cluster of disease mutations reported in SLC25A46 face the IMS (**Figure 4A**)

(16). While many mutations were reported to destabilize the protein, certain disease-

causing mutations (T142I and R340C) cause a hyperfused mitochondrial phenotype (6,

8, 9, 16, 25), similar to that observed with siRNA/shRNA knockdown of SLC25A46 (6, 9,

25).

To investigate cellular mitochondrial phenotypes upon SLC25A46 perturbations,

we imaged mitochondrial networks in the human colorectal carcinoma cell line HCT116.

Under steady-state conditions, WT and SLC25A46 KO HCT116 cells demonstrate

similar mitochondrial network morphology. Previous studies show that nutrient

starvation triggered stress can induce mitochondrial hyperfusion (48, 49). To test if

mitochondria in HCT116 cells undergo similar stress-induced change, cells were grown

in serum-free medium. Under such conditions, we observed significant mitochondrial

fragmentation in SLC25A46 KO cells but not in WT cells. Reintroduction of WT

SLC25A46 rescued this phenotype (**Figure 4B, Supporting Video 1**), consistent with

previous observations in human fibroblasts and HeLa cells (8). We found that

expression of R257Q or R340C SLC25A46 mutants induced interconnected

mitochondrial networks, also in agreement with previous studies (8, 25) (**Figure 4B,**

Supporting Video 1). The mitochondrial morphology in G249D-expressing cells was

similar to WT (**Figure 4B, Supporting Video 1**). Expression levels of R340C and

G249D mutants never reached WT-like levels (**Figure 4C**), likely due to protein

instability (6, 8, 9, 16, 22).

We next tested the effects of the newly identified Opa1-SLC25A46 interactions

on mitochondrial morphology. Expression of I349D SLC25A46 (disrupts a hydrophobic

interface between the Opa1 BSE) produced shorter mitochondrial tubules compared to

WT, whereas R257A/R347A mutant (disrupt SLC25A46-Opa1 salt bridge interactions) resulted in more fragmented mitochondria (**Figure 4B, Supporting Video 1**). These results suggest that the identified SLC24A45-Opa1 interaction interface is important for maintaining a tubular mitochondrial morphology. Expression of the (Δ2-83)SLC25A46 resulted in an increase in short mitochondrial tubules (**Figure 4B, Supporting Video 1**). In these experiments, we did not observe changes in levels of Mfn2 or Opa1 due to SLC25A46 knockout or expression of varying levels of mutant SLC25A46 (**Figure 4C**). The expression level of (Δ2-83)SLC25A46 could not be evaluated due to the inability of commercial antibodies to detect this mutant (**Figure 4C**).

We also investigated SLC25A46 disease mutants expressed in yeast. Under these conditions, many of SLC25A46 mutants still bound Opa1 (**Supporting Figure 5B**). We also saw little effect of patient-derived mutations on the Mfn2-SLC25A46 interaction (**Supporting Figure 5B**). We note that in these binding experiments we were unable to achieve similar expression levels of the respective proteins, and binding may only describe one aspect a complex phenotype only observable under cellular conditions.

Discussion

Mitochondria undergo morphological changes tightly coupled to cellular physiology (4). While the identity of the GTPases responsible for sequential mitochondrial fusion have long been identified (50), and mechanisms for individual fusion steps have been described (51-54), questions remain regarding how outer and inner-membrane activities are coordinated and regulated. Emerging evidence points to SLC25A46 as a regulator of fusion activity, with protein levels influencing the fusion-fission balance (6, 8, 25). Consistent with this role, SLC25A46 is observed to localize to the fusion sites (8). In this study, we investigate the physical interaction between SLC25A46 and the membrane fusogens Opa1 and Mfn1/2. Combining crosslinking mass-spectrometry, and AlphaFold 2 modeling, we identify and validate the BSE region in Opa1 interaction with the IMS-face of SLC25A46. We also present evidence for an interaction between SLC25A46 cytosolic face and the Mfn2 ectodomain. The Mfn2 region we identify includes helical bundle 1 (HB1) and GTPase domains, but not the

transmembrane segments. This is consistent with yeast studies, where residues 1-294
of Ugo1 interacts with Fzo1 in a region comprising residues N-terminal to the first
transmembrane helix and the C-terminal helical repeat 2 (34).

Reports from several studies indicate that mitochondrial morphology is sensitive
to SLC25A46 levels. Schuettpelz *et al.* reported mitochondrial fragmentation in human
fibroblasts and HeLa cells upon SLC25A46 KO (8, 55). In SLC25A46 KO HCT116 cells,
we observed mitochondrial fragmentation upon serum starvation. In addition, we
observed a ~2-fold slower growth of this cell line, consistent with observations in
SLC25A46 deleted fibroblasts (8). In previous studies, overexpression of SLC25A46
increased mitochondrial fragmentation, suggesting that at high levels, SLC25A46 may
downregulate the activities of the protein fusogens. (6, 25, 31). One possible mode of
inhibitory action may be SLC25A46 binding resulting in steric competition for Opa1 self-
assembly, which relies on the BSE and stalk regions to mediate formation of helical
filaments (56-58). Consistent with this prediction, levels of Opa1 oligomeric complexes
were found to increase upon loss of SLC25A46 (8).

In our study, SLC25A46 mutants targeting Opa1 interaction did not promote a
hyperfused phenotype, but instead disrupted the mitochondrial network. This suggests
that a SLC25A46-Opa1 interaction promotes fusion necessary for a tubular network.
Although we could not estimate the relative levels of (Δ2-83)SLC25A46, this N-terminal
deletion reduced the filamentous network, while increasing the proportion of short
tubules, also suggesting a fusion defect. Since the SLC25A46 N-terminal region alone
is insufficient to bind with Mfn2, we speculate that this region may play a stabilizing role
facilitating interactions of additional cytosolic-facing regions with the ectodomain of
Mfn2. Taken together, these observations provide evidence for SLC25A46 interactions
playing an activating role in fusion when expressed at physiological levels.

Our results provide a basis for future functional analyses of SLC25A46 regulation
of the fusion machinery. SLC25A46 protein-protein interactions also may regulate
cristae morphology, which is dramatically affected upon loss of SLC25A46 (6, 8, 25-27,
29). Proteomics studies have also implicated SLC25A46 endoplasmic reticulum
interaction partners in maintaining ER-mitochondria contacts facilitating lipid flux (6-8).

Future mechanistic structure-activity studies are necessary to explore these other roles

of SLC25A46 in cristae regulation and lipid metabolism.

Experimental procedures

Plasmids

      Genes for SLC25A46, Mfn2, Opa1(253-960), Opa1(MGD) constructs were synthesized (GenScript) and cloned into pPICZ A for *P. pastoris* expression. C-terminus of SLC25A46 was fused to a $(G_4S)_3$ linker followed by TEV cleavage site and deca-histidine. N-terminus of Opa1(253-960), Opa1(263-960), Opa1(MGD) was tagged with Twin-Strep tag followed by HRV 3C cleavage site and $(G_4S)_3$ linker. The C-terminus of Mfn2 was fused to G_3S linker, HRV 3C cleavage site and Twin-Strep tag. For co-expression, SLC25A46 and either Mfn2 or Opa1 with tags with its own AOX1 promoter and terminator were cloned into the same plasmid. (SLC25A46(1-88)-MBP) was generated from SLC25A46(1-88) fused to MBP followed by a GGGS linker TEV site and His10 tag. Mfn2(ΔTM) was generated from Mfn2(2-544)-GGS linker-Mfn2(651-757). Mfn2(MGD) was generated from Mfn2(2-400:706-757). Strep-Opa1(253-960) and Strep-Opa1(MGD) were also cloned into pET-28a(+) vector for expression in *E. coli*.

Generation of *Pichia* strains

      Expression strains were generated from *Pichia pastoris* strain SMD1163 (*his4 pep4 prb1*) (kind gift from Dr. Tom Rapaport (Harvard Medical School) according to established protocols (59). Plasmids were transformed into SMD1163 by electroporation and grown on YPDS (1% yeast extract, 2% peptone, 2% dextrose, 1 M sorbitol) zeocin (InvivoGen, ant-zn-1p) plates. 7-14 resulting colonies were grown in BMGY (1% yeast extract, 2% peptone, 100 mM potassium phosphate pH 6.0, 1.34% yeast nitrogen base, 4×10^{-5}% biotin, 1% glycerol) and induced in BMMY (replaced glycerol with 0.5% methanol in BMGY). Cultures were screened for protein expression by Western using anti-His, anti-Strep, anti-SLC25A46, anti-Opa1 or anti-Mfn2 antibodies.

Binding experiments using lysates

      P. pastoris (1 L) expressing SLC25A46, Opa1, Mfn2 alone or in combination were harvested and cryo-milled using SPEX SamplePrep 6875D Freezer/Mill. For StrepTactin column-binding and elution, milled cells (125 mg) were resuspended in 3 ml

of 25 mM HEPES, 150 mM NaCl, pH 8.0 buffer containing 1% GDN and protease

inhibitors, tumbled for 1-1:30 h at 4 °C. Milled cell input was increased to 375 mg (3x

original input) when needed for increased signal. Suspension was centrifuged at 20,000

for 30 min at 4 °C. Supernatant (100 µl) was incubated with 50 µl of StrepTactin XT

4Flow high-capacity resin in a spin column (Abcam, ab286861) at 4°C for 15 min

followed by centrifugation at 500 xg for 30 s. Resin was washed 3x with 100 µl buffer

containing 0.1% GDN. To elute, the resin was incubated with 100 µl buffer with 50 mM

biotin for 15 min and centrifuged. Same procedure for Ni-NTA binding experiments

except that the binding and elution buffer contained 80 mM and 500 mM imidazole,

respectively.

AlphaFold 2 modeling

Protein sequences for human SLC25A46, Opa1 and Mfn2 were run through in a

AlphaFold2 Google ColabFold – v 1.5.3 workbook (38, 39, 60). Five independent runs

generated the top three ranked relaxed models for SLC25A46-Opa1 and SLC25A46-

Mfn2 models respectively.

Data availability

Mass spectrometry data available on request.

Supporting information

This article contains supporting information.

Acknowledgements

We thank Dr. Yuan Gao and Dr. Tom Rapoport for providing *Pichia pastoris* strain SMD1163. We thank all the members of the Chao lab for critical feedback.

Funding and additional information

Mass spectrometric experiments supported by a shared instrumentation grant from the NIH, 1S10OD010582-01A1 towards purchase of Orbitrap Eclipse. This work was supported by funding from the National Institutes of Health (R35GM142553 to L.H.C.).

References

1. Wai, T., andLanger, T. (2016) Mitochondrial Dynamics and Metabolic Regulation Trends Endocrinol Metab **27**, 105-117 10.1016/j.tem.2015.12.001

2. Yapa, N. M. B., Lisnyak, V., Reljic, B., andRyan, M. T. (2021) Mitochondrial dynamics in health and disease FEBS Lett **595**, 1184-1204 10.1002/1873-3468.14077

3. Giacomello, M., Pyakurel, A., Glytsou, C., andScorrano, L. (2020) The cell biology of mitochondrial membrane dynamics Nat Rev Mol Cell Biol **21**, 204-224 10.1038/s41580-020-0210-7

4. Chan, D. C. (2020) Mitochondrial Dynamics and Its Involvement in Disease Annu Rev Pathol **15**, 235-259 10.1146/annurev-pathmechdis-012419-032711

5. Meyer, J. N., Leuthner, T. C., andLuz, A. L. (2017) Mitochondrial fusion, fission, and mitochondrial toxicity Toxicology **391**, 42-53 10.1016/j.tox.2017.07.019

6. Janer, A., Prudent, J., Paupe, V., Fahiminiya, S., Majewski, J., Sgarioto, N. *et al.* (2016) SLC25A46 is required for mitochondrial lipid homeostasis and cristae maintenance and is responsible for Leigh syndrome EMBO Mol Med **8**, 1019-1038 10.15252/emmm.201506159

7. Perivolidi, V. I., Violitzi, F., Ioannidou, E., Rinotas, V., Stamatakis, G., Samiotaki, M. *et al.* (2022) Proteomic Identification of the SLC25A46 Interactome in Transgenic Mice Expressing SLC25A46-FLAG J Proteome Res **21**, 375-394 10.1021/acs.jproteome.1c00728

8. Schuettpelz, J., Janer, A., Antonicka, H., andShoubridge, E. A. (2023) The role of the mitochondrial outer membrane protein SLC25A46 in mitochondrial fission and fusion Life Sci Alliance **6**, 10.26508/lsa.202301914

9. Steffen, J., Vashisht, A. A., Wan, J., Jen, J. C., Claypool, S. M., Wohlschlegel, J. A. *et al.* (2017) Rapid degradation of mutant SLC25A46 by the ubiquitin-proteasome system results in MFN1/2-mediated hyperfusion of mitochondria Mol Biol Cell **28**, 600-612 10.1091/mbc.E16-07-0545

10. Kunji, E. R. S., King, M. S., Ruprecht, J. J., andThangaratnarajah, C. (2020) The SLC25 Carrier Family: Important Transport Proteins in Mitochondrial Physiology and Pathology Physiology (Bethesda) **35**, 302-327 10.1152/physiol.00009.2020

11. Kodal, L. S., Hammer-Hansen, S., Holm-Yildiz, S., Gronskov, K., Karstensen, H. G., andDysgaard, T. (2022) Case report: A novel variant in SLC25A46 causing sensorimotor polyneuropathy and optic atrophy Front Neurol **13**, 1066040 10.3389/fneur.2022.1066040

12. Yamada, M., Suzuki, H., Adachi, H., Noguchi, A., Miya, F., Takahashi, T. *et al.* (2022) Diagnosis of SLC25A46-related pontocerebellar hypoplasia in two siblings with fulminant neonatal course: role of postmortem CT and whole genomic analysis: a case report BMC Neurol **22**, 20 10.1186/s12883-021-02540-x

13. Li, Q., Madden, J. A., Lin, J., Shi, J., Rosen, S. M., Schmitz-Abe, K. *et al.* (2021) Reanalysis of Exome Data Identifies Novel SLC25A46 Variants Associated with Leigh Syndrome J Pers Med **11**, 10.3390/jpm11121277

14. Raju, S., Medarametla, S., andBoraiah, N. (2021) Dystonia and Hereditary Motor Sensory Neuropathy 6B Due to SLC25A46 Gene Mutations Mov Disord Clin Pract **8**, 480-482 10.1002/mdc3.13139

15. Bitetto, G., Malaguti, M. C., Ceravolo, R., Monfrini, E., Straniero, L., Morini, A. *et*
al. (2020) SLC25A46 mutations in patients with Parkinson's Disease and optic
atrophy Parkinsonism Relat Disord **74**, 1-5 10.1016/j.parkreldis.2020.03.018
16. Abrams, A. J., Fontanesi, F., Tan, N. B. L., Buglo, E., Campeanu, I. J., Rebelo,
A. P. *et al.* (2018) Insights into the genotype-phenotype correlation and molecular
function of SLC25A46 Hum Mutat **39**, 1995-2007 10.1002/humu.23639
17. Ivanov, I., Atkinson, D., Litvinenko, I., Angelova, L., Andonova, S., Mumdjiev, H.
et al. (2018) Pontocerebellar hypoplasia type 1 for the neuropediatrician:
Genotype-phenotype correlations and diagnostic guidelines based on new cases
and overview of the literature Eur J Paediatr Neurol **22**, 674-681
10.1016/j.ejpn.2018.03.011
18. Braunisch, M. C., Gallwitz, H., Abicht, A., Diebold, I., Holinski-Feder, E., Van
Maldergem, L. *et al.* (2018) Extension of the phenotype of biallelic loss-of-
function mutations in SLC25A46 to the severe form of pontocerebellar hypoplasia
type I Clin Genet **93**, 255-265 10.1111/cge.13084
19. van Dijk, T., Rudnik-Schoneborn, S., Senderek, J., Hajmousa, G., Mei, H., Dusl,
M. *et al.* (2017) Pontocerebellar hypoplasia with spinal muscular atrophy (PCH1):
identification of SLC25A46 mutations in the original Dutch PCH1 family Brain
140, e46 10.1093/brain/awx147
20. Hammer, M. B., Ding, J., Mochel, F., Eleuch-Fayache, G., Charles, P., Coutelier,
M. *et al.* (2017) SLC25A46 Mutations Associated with Autosomal Recessive
Cerebellar Ataxia in North African Families Neurodegener Dis **17**, 208-212
10.1159/000464445
21. Sulaiman, R. A., Patel, N., Alsharif, H., Arold, S. T., andAlkuraya, F. S. (2017) A
novel mutation in SLC25A46 causes optic atrophy and progressive limb
spasticity, with no cerebellar atrophy or axonal neuropathy Clin Genet **92**, 230-
231 10.1111/cge.12963
22. Wan, J., Steffen, J., Yourshaw, M., Mamsa, H., Andersen, E., Rudnik-
Schoneborn, S. *et al.* (2016) Loss of function of SLC25A46 causes lethal
congenital pontocerebellar hypoplasia Brain **139**, 2877-2890
10.1093/brain/aww212
23. Charlesworth, G., Balint, B., Mencacci, N. E., Carr, L., Wood, N. W., andBhatia,
K. P. (2016) SLC25A46 mutations underlie progressive myoclonic ataxia with
optic atrophy and neuropathy Mov Disord **31**, 1249-1251 10.1002/mds.26716
24. Nguyen, M., Boesten, I., Hellebrekers, D. M., Mulder-den Hartog, N. M., de Coo,
I. F., Smeets, H. J. *et al.* (2017) Novel pathogenic SLC25A46 splice-site mutation
causes an optic atrophy spectrum disorder Clin Genet **91**, 121-125
10.1111/cge.12774
25. Abrams, A. J., Hufnagel, R. B., Rebelo, A., Zanna, C., Patel, N., Gonzalez, M. A.
et al. (2015) Mutations in SLC25A46, encoding a UGO1-like protein, cause an
optic atrophy spectrum disorder Nat Genet **47**, 926-932 10.1038/ng.3354
26. Duchesne, A., Vaiman, A., Castille, J., Beauvallet, C., Gaignard, P., Floriot, S. *et*
al. (2017) Bovine and murine models highlight novel roles for SLC25A46 in
mitochondrial dynamics and metabolism, with implications for human and animal
health PLoS Genet **13**, e1006597 10.1371/journal.pgen.1006597

27. Li, Z., Peng, Y., Hufnagel, R. B., Hu, Y. C., Zhao, C., Queme, L. F. *et al.* (2017)
Loss of SLC25A46 causes neurodegeneration by affecting mitochondrial
dynamics and energy production in mice Hum Mol Genet **26**, 3776-3791
10.1093/hmg/ddx262

28. Suda, K., Ueoka, I., Azuma, Y., Muraoka, Y., Yoshida, H., andYamaguchi, M.
(2018) Novel Drosophila model for mitochondrial diseases by targeting of a
solute carrier protein SLC25A46 Brain Res **1689**, 30-44
10.1016/j.brainres.2018.03.028

29. Terzenidou, M. E., Segklia, A., Kano, T., Papastefanaki, F., Karakostas, A.,
Charalambous, M. *et al.* (2017) Novel insights into SLC25A46-related
pathologies in a genetic mouse model PLoS Genet **13**, e1006656
10.1371/journal.pgen.1006656

30. Qiu, K., Zou, W., Fang, H., Hao, M., Mehta, K., Tian, Z. *et al.* (2022) Light-
activated mitochondrial fission through optogenetic control of mitochondria-
lysosome contacts Nat Commun **13**, 4303 10.1038/s41467-022-31970-5

31. Santo-Domingo, J., Lassueur, S., Galindo, A. N., Alvarez-Illera, P., Romero-
Sanz, S., Caldero-Escudero, E. *et al.* (2023) SLC25A46 promotes mitochondrial
fission and mediates resistance to lipotoxic stress in INS-1E insulin-secreting
cells J Cell Sci **136**, 10.1242/jcs.260049

32. Coonrod, E. M., Karren, M. A., andShaw, J. M. (2007) Ugo1p is a multipass
transmembrane protein with a single carrier domain required for mitochondrial
fusion Traffic **8**, 500-511 10.1111/j.1600-0854.2007.00550.x

33. Sesaki, H., andJensen, R. E. (2001) UGO1 encodes an outer membrane protein
required for mitochondrial fusion J Cell Biol **152**, 1123-1134
10.1083/jcb.152.6.1123

34. Sesaki, H., andJensen, R. E. (2004) Ugo1p links the Fzo1p and Mgm1p
GTPases for mitochondrial fusion J Biol Chem **279**, 28298-28303
10.1074/jbc.M401363200

35. Hoppins, S., Horner, J., Song, C., McCaffery, J. M., andNunnari, J. (2009)
Mitochondrial outer and inner membrane fusion requires a modified carrier
protein J Cell Biol **184**, 569-581 10.1083/jcb.200809099

36. Sesaki, H., Southard, S. M., Yaffe, M. P., andJensen, R. E. (2003) Mgm1p, a
dynamin-related GTPase, is essential for fusion of the mitochondrial outer
membrane Mol Biol Cell **14**, 2342-2356 10.1091/mbc.e02-12-0788

37. Wong, E. D., Wagner, J. A., Scott, S. V., Okreglak, V., Holewinske, T. J.,
Cassidy-Stone, A. *et al.* (2003) The intramitochondrial dynamin-related GTPase,
Mgm1p, is a component of a protein complex that mediates mitochondrial fusion
J Cell Biol **160**, 303-311 10.1083/jcb.200209015

38. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O. *et al.*
(2021) Highly accurate protein structure prediction with AlphaFold Nature **596**,
583-589 10.1038/s41586-021-03819-2

39. Mirdita, M., Schutze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., andSteinegger,
M. (2022) ColabFold: making protein folding accessible to all Nat Methods **19**,
679-682 10.1038/s41592-022-01488-1

40.   Ishihara, N., Fujita, Y., Oka, T., andMihara, K. (2006) Regulation of mitochondrial
459   morphology through proteolytic cleavage of OPA1 EMBO J **25**, 2966-2977
460   10.1038/sj.emboj.7601184

461   41.   Olichon, A., Emorine, L. J., Descoins, E., Pelloquin, L., Brichese, L., Gas, N. *et*
462   *al.* (2002) The human dynamin-related protein OPA1 is anchored to the
463   mitochondrial inner membrane facing the inter-membrane space FEBS Lett **523**,
464   171-176 10.1016/s0014-5793(02)02985-x

465   42.   Chappie, J. S., Acharya, S., Liu, Y. W., Leonard, M., Pucadyil, T. J., andSchmid,
466   S. L. (2009) An intramolecular signaling element that modulates dynamin
467   function in vitro and in vivo Mol Biol Cell **20**, 3561-3571 10.1091/mbc.e09-04-
468   0318

469   43.   Sever, S., Muhlberg, A. B., andSchmid, S. L. (1999) Impairment of dynamin's
470   GAP domain stimulates receptor-mediated endocytosis Nature **398**, 481-486
471   10.1038/19024

472   44.   Anand, R., Eschenburg, S., andReubold, T. F. (2016) Crystal structure of the
473   GTPase domain and the bundle signalling element of dynamin in the GDP state
474   Biochem Biophys Res Commun **469**, 76-80 10.1016/j.bbrc.2015.11.074

475   45.   Chappie, J. S., Acharya, S., Leonard, M., Schmid, S. L., andDyda, F. (2010) G
476   domain dimerization controls dynamin's assembly-stimulated GTPase activity
477   Nature **465**, 435-440 10.1038/nature09032

478   46.   Kong, L., Sochacki, K. A., Wang, H., Fang, S., Canagarajah, B., Kehr, A. D. *et al.*
479   (2018) Cryo-EM of the dynamin polymer assembled on lipid membrane Nature
480   **560**, 258-262 10.1038/s41586-018-0378-6

481   47.   Li, Y. J., Cao, Y. L., Feng, J. X., Qi, Y., Meng, S., Yang, J. F. *et al.* (2019)
482   Structural insights of human mitofusin-2 into mitochondrial fusion and CMT2A
483   onset Nat Commun **10**, 4914 10.1038/s41467-019-12912-0

484   48.   Labbe, K., Mookerjee, S., Le Vasseur, M., Gibbs, E., Lerner, C., andNunnari, J.
485   (2021) The modified mitochondrial outer membrane carrier MTCH2 links
486   mitochondrial fusion to lipogenesis J Cell Biol **220**, 10.1083/jcb.202103122

487   49.   Tondera, D., Grandemange, S., Jourdain, A., Karbowski, M., Mattenberger, Y.,
488   Herzig, S. *et al.* (2009) SLP-2 is required for stress-induced mitochondrial
489   hyperfusion EMBO J **28**, 1589-1600 10.1038/emboj.2009.89

490   50.   Gao, S., andHu, J. (2021) Mitochondrial Fusion: The Machineries In and Out
491   Trends Cell Biol **31**, 62-74 10.1016/j.tcb.2020.09.008

492   51.   Anand, R., Wai, T., Baker, M. J., Kladt, N., Schauss, A. C., Rugarli, E. *et al.*
493   (2014) The i-AAA protease YME1L and OMA1 cleave OPA1 to balance
494   mitochondrial fusion and fission J Cell Biol **204**, 919-929 10.1083/jcb.201308006

495   52.   Ban, T., Ishihara, T., Kohno, H., Saita, S., Ichimura, A., Maenaka, K. *et al.* (2017)
496   Molecular basis of selective mitochondrial fusion by heterotypic action between
497   OPA1 and cardiolipin Nat Cell Biol **19**, 856-863 10.1038/ncb3560

498   53.   Ge, Y., Shi, X., Boopathy, S., McDonald, J., Smith, A. W., andChao, L. H. (2020)
499   Two forms of Opa1 cooperate to complete fusion of the mitochondrial inner-
500   membrane Elife **9**, 10.7554/eLife.50973

501   54.   Mishra, P., Carelli, V., Manfredi, G., andChan, D. C. (2014) Proteolytic cleavage
502   of Opa1 stimulates mitochondrial inner membrane fusion and couples fusion to
503   oxidative phosphorylation Cell Metab **19**, 630-641 10.1016/j.cmet.2014.03.011

55. Schuettpelz, J., Watt, K., Antonicka, H., Janer, A., Larsson, O., andShoubridge,
E. A. (2024) SLC25A46 is in contact with lysosomes and plays a role in
mitochondrial cholesterol homeostasis bioRxiv 2024.2004.2001.587618
10.1101/2024.04.01.587618
56. Faelber, K., Dietrich, L., Noel, J. K., Wollweber, F., Pfitzner, A. K., Muhleip, A. *et*
al. (2019) Structure and assembly of the mitochondrial membrane remodelling
GTPase Mgm1 Nature **571**, 429-433 10.1038/s41586-019-1372-3
57. Nyenhuis, S. B., Wu, X., Strub, M. P., Yim, Y. I., Stanton, A. E., Baena, V. *et al.*
(2023) OPA1 helical structures give perspective to mitochondrial dysfunction
Nature **620**, 1109-1116 10.1038/s41586-023-06462-1
58. von der Malsburg, A., Sapp, G. M., Zuccaro, K. E., von Appen, A., Moss, F. R.,
3rd, Kalia, R. *et al.* (2023) Structural mechanism of mitochondrial membrane
remodelling by human OPA1 Nature **620**, 1101-1108 10.1038/s41586-023-
06441-6
59. Hartmann, L., Kugler, V., andWagner, R. (2016) Expression of Eukaryotic
Membrane Proteins in Pichia pastoris Methods Mol Biol **1432**, 143-162
10.1007/978-1-4939-3637-3_10
60.
https://colab.research.google.com/github/deepmind/alphafold/blob/main/n
otebooks/AlphaFold.ipynb
524

Abbreviations and nomenclature

OMM – outer mitochondrial membrane

IMM – inner mitochondrial membrane

SLC – solute carrier family

SLC25A46 – solute carrier family 25 member A46

Opa1 – optic atrophy 1

Mfn1/2 – mitofusin 1/2

iPSC – induced pluoripotent stem cells

BSE – bundle signaling element

HR1/2 – heptad repeat 1/2

ER – endoplasmic reticulum

MGD – minimal GTPase domain

DDM – n-Dodecyl-β-D-Maltopyranoside

GDN – glyco-diosgenin

PA – phosphatidic acid

TM – transmembrane segment

MTS – mitochondrial targeting sequence

IMS – inter-membrane space

Figure legends

Figure 1. SLC25A46 interacts with the bundle signaling element (BSE) of Opa1.

A. Domain diagrams of SLC25A46 and Opa1. s-Opa1 (amino acids 253-960) and

Opa1(MGD) (minimal GTPase domain comprising the GTPase domain and three BSE

helices) are shown as line diagrams below full length Opa1 sequence.

B. GDN extracts prepared from *P. pastoris* expressing SLC25A46-His, Strep-Opa1 or

co-expressing SLC25A46-His and Strep-Opa1 subjected to StrepTactin column binding

and elution.

C. Crosslinks identified in BSE helices 2 and 3 of Opa1. Circular map of SLC25A46 and

Opa1 with identified lysine-lysine crosslinks (black arcs). Crosslinker used to generate

the crosslink is indicated.

D. AlphaFold 2 model of the SLC25A46-Opa1 interaction. N-terminal region of

SLC25A46 comprising residues 1-83 (disordered) not displayed, residues 1-263 of

Opa1 not modeled. Box region displaying identified crosslinks and the calculated C□-

C□ distances based on the model.

Figure 2. SLC25A46 mutations in the binding interface diminish binding to Opa1.

A. SLC25A46-Opa1 interface observed in AlphaFold 2 model. Top view: Adjacent to the

salt bridges is a cluster of hydrophobic residues comprising of I229, I230, I349, Y357,

V359, L360 of SLC25A46 and T576, F572, A569 of Opa1. Bottom view: Predicted salt-

bridge interactions between R257 of SLC25A46 and E561 of Opa1, and R347 of

SLC25A46 and D565 of Opa1. Interface residues highlighted. Mutated residues

indicated in red.

B. Purified SLC25A46-His WT, I349D mutant or R257A/R347A double mutant were

incubated with 2 µM of Strep-Opa1 and subjected to StrepTactin binding and elution

experiments. Input and elution analyzed by Western blotting.

Figure 3. SLC25A46 interactions with Mfn2.

A. Domain diagrams of full length and truncated constructs of SLC25A46 and Mfn2 are

depicted.

B. SLC25A46-His co-elutes upon elution of bound Mfn2-Strep. GDN extracts prepared

from *P. pastoris* expressing Mfn2-Strep or co-expressing SLC25A46-His and Mfn2-

Strep were subjected to StrepTactin column binding and elution.

C. Circular map of SLC25A46 and Mfn2 with the identified lysine-lysine crosslinks (black

arcs). Crosslinker used is indicated. Crosslinks were identified between the cytosol-

exposed N-terminal K60 of SLC25A46 and the HR1 lysines K416 and K420 of Mfn2.

D. Model of the SLC25A46-Mfn2 interaction. The identified crosslinks are depicted as

dotted lines in the AlphaFold 2 generated models of SLC25A46 and Mfn2. Helical

bundle 1 (HB1) indicated. The N-terminal region of SLC25A46 comprising residues 1-83

is predicted to be disordered and is displayed as a hand-drawn cartoon. The

transmembrane helices of Mfn2 were poorly modeled and are shown as transparent

cylinders, to indicate uncertainty.

E. (Δ2-83)SLC25A46-His co-elutes upon elution of bound Mfn2-Strep. GDN extracts

prepared from *P. pastoris* expressing Mfn2-Strep and co-expressing SLC25A46-His or

(Δ2-83)SLC25A46-His were subjected to StrepTactin column binding and elution.

F. Residues 1-88 of SLC25A46 fused to MBP and His tag at the C-terminus

(SLC25A46(1-88)-MBP-His) does not co-elute with immobilized Mfn2-Strep.

G. A Mfn2 lacking the two transmembrane segments (Mfn2(ΔTM)-Strep) co-elutes with
SLC25A46-His.

H. A minimal Mfn2 construct comprising helical bundle 1 (HB1) and the GTPase domain
co-elutes with SLC25A46-His.

Figure 4. Mutations disrupting Opa1 interaction alter mitochondrial morphology.

A. Identified SLC25A46 disease mutations mapped on AlphaFold 2 model of
SLC25A46-Opa1 complex. Box regions enlarged on the right to display the mutations.

B. Graph bar representing the relative proportion of scored mitochondrial morphologies
in HCT116 WT, SLC25A46-KO or SLC25A46-KO cells expressing SLC25A46 variants.
Mitochondrial morphology for 300 cells were scored for each condition and evaluated
(see Materials and Methods). Significance of difference is tested relative to wild type
using unpaired t test with Welch's correction. Blue ****: $p<0.0001$ when compared to
scored WT cells showing fragmented mitochondria. Green *: $p<0.05$ when compared to
scored WT cells showing interconnected mitochondrial network.

C. Western blot of expression levels for Opa1, Mfn2, SLC25A46 or MTCO1 in HCT116
WT, SLC25A46-KO or SLC25A46-KO cells expressing SLC25A46 variants. (Δ2-
83)SLC25A46 not detected as commercially available antibodies detect the N-terminal
epitope of SLC25A46. MTCO1 is used as loading control.

Figure 1

632
633

634 **Figure 2**

635
636

637　**Figure 3**

638
639

640 **Figure 4**

A

B

Percentage (%)

Legend:
- Interconnected
- Long tubular
- Short tubular
- Fragmented

****

X-axis: WT, SLC25A46-KO, WT, G249D, R257Q, R340C, I349D, R257A,R347A, Δ(2-83)

SLC25A46-KO +

C

SLC25A46-KO +

WT | SLC25A46-KO | WT | G249D | R257Q | R340C | I349D | R257A/R347A | Δ(2-83)

Anti-Opa1

Anti-Mfn2

Anti-SLC25A46

Anti-MTCO1